

In this response, bold parts in quotes are direct extracts of referee comments, blue italic parts are changes made in the article and black normal texts are answers/explanations about each comment made by the referees and the reviewer. The authors thank both referees and the reviewer for their pertinent comments.

Referee 1 comments:

5 The authors thank the referee for his positive general evaluation of the paper, pointing out its originality and interest. Their detailed comments have allowed its further improvement. The paper has been carefully reread and very long sentences split.

Referee 1 general comments:

10 **1. “Page 6, line 2: I don’t understand the aim of the sentence "Four volatility bins are added for anthropogenic and biogenic SOA ranging from 1 to 1000 $\mu\text{g}\cdot\text{m}^{-3}$ ". You say on page 5, that you have 9 volatility bins between 0.01 and 10⁶, corresponding to the figure SI2. On the Figure 2, you represent the 4 volatility bins between 1 and 1000. Which bins do you use in the model? Please make the manuscript clearer.”**

15 There is a distinction in this work and also other previous articles concerning VBS between POA volatility bins and ASOA/BSOA volatility bins because of their different nature and also different sources. For ASOA/BSOA species four volatility bins ranging between 1 to 1000 $\mu\text{g}\cdot\text{m}^{-3}$ are employed, while for POA 9 bins ranging between 0.01 and 10⁶ are used. In the article, this distinction has been shown in figure 2 for BSOA/ASOA species, while same type of figure for POA is shown in the supplementary materials (Figure SI-1). To show this fact better a text was added to the legend of figure 2:

Note that this schematic represents BSOA/ASOA where four bins are used, for SVOC/IVOC (where 9 bins are used) a schematic is presented in SI-1.

2. “Page 10, line 34: I don’t understand what you mean by "for regimes".”

25 The sentence was intended to present meteorological regimes, meaning periods with different meteorological conditions. The wording is changed to “synoptical meteorological conditions” in the revised manuscript.

30 *While the general comparison between the hourly meteorological fields used as input for CHIMERE simulations and observations is in general already satisfying, the correlation becomes higher and the bias lower when daily averages representative for different meteorological conditions are compared instead of hourly values. (Page 11, lines 24-26)*

3. “Page 12, line 18-20: Could you explain a little more the cloud effect affecting aerosols concentrations.”

The following sentences have been added for the cloud scavenging effect seen in the observations:

35 *Cloud scavenging processes are already taken into account in the model. However, because of the unique geographical characteristics mentioned before for this site, the meteorological inputs did not simulate these fog events and therefore cloud scavenging was not activated in the simulation. Since these decreases concern only a small percentage of the observations, they do not have a major effect on the outcome of these comparisons. While this effect is very visible for sulfates, it is less pronounced for other particulate species such as black carbon and OA. (Page 13, lines 10-15)*

40 **4. “Page 18, line 10-12: The concentrations of allPOA around Corsica does not reach 1.5 $\mu\text{g}\cdot\text{m}^{-3}$ in figure 12.g. Same for figure 13.e and the 30% of allPOA to the OA. Could you correct the text?”**

Yes, we agree with the first referee. It has been corrected in the revised version of the article.

Referee 1 figure related comments:

1. **“Figure 4: You mention a, b and c, but they are not reported on the figure.”**

Yes, it has been corrected in the revised version of the article.

2. **“Figure 5.b1: The blue ribbon around the date 20130726 seems weird. As I understood your method, it should be placed around the simulation results (blue line), but on these part of the graph it seems to be around the black line corresponding to the observation. Could you explain why?”**

This error is rectified in the revised version. The maximum of the ribbon being higher than the limit of the plot, the coding program unfortunately removed the whole upper ribbon for this date.

Referee 1 technical comments:

1. **Page 5, line 26: Please add "SI1" after "in the supplementary information".**

Added in the revised version.

2. **Page 6, line 33: as in Shrivastava et al. 2013 -> as in Shrivastava et al. (2013)**

Modified in the revised version.

3. **Page 6, line 36: life time -> lifetime**

Modified in the revised version.

4. **Page 10, line 14: Please add "Orographic Representativeness Error" for the clarity of the text.**

Modified in the revised version.

5. **Page 12, line 30: You say the correlation is 0.26, but in the table it is 0.36 for Ersar and 0.47 for Es Pinar. Could you correct, or explain?**

Modified in the revised version. It was just a confusion between orographic representativeness of 26 % and a correlation of 0.36.

6. **Page 14, line 22: Could you precise what US stands for?**

US stands for United States, it has been changed to *the USA* in the revised version.

Referee 2 comments:

Referee 2 general comments:

We wish to thank referee 2 for his evaluation of the paper, which has allowed us to improve the paper. This response integrates our earlier short response to referee 2, which replied to the major issues raised. We hope that our reply can assure the referee that our paper presents indeed original and novel results.

1. **“In addition, the authors do not give appropriate credit of the work of Chrit et al. (2017) who simulated this same data, using a different approach. The authors indicate that the concentrations and properties are well simulated – so this calls into question the need for the current paper. What does this new study tell us that the work of Chrit et al. (2017) did not?”**

Although the two papers are based on results from the same measurements campaign, there are fundamental differences between them. Apart from obvious differences like different models with different inputs (emissions, meteorological fields ...), the approach taken in the two articles regarding the simulation of organic aerosols is clearly different. In our article, we use different VBS schemes, taking into account formation of non-volatile SOA and fragmentation processes in one of them. In Chrit et al. (2017), they use a surrogate based approach where SOA are divided into three types, hydrophilic,

hydrophobic or both (Couvidat et Sartelet 2015). In addition, ELVOCs are also taken into account. To our point of view, it is important to compare all these different schemes to detailed measurement data and not limit this comparison to one particular scheme.

In several aspects, our paper goes significantly beyond the Chrit et al. (2017) paper, which was published earlier.

(1) In our work, we use measurements from a second site at Mallorca in addition to data from Cap Corsica for the same period, which is not the case for the work of Chrit et al. (2017). The two sites have different characteristics as it is seen in the results of ^{14}C measurements and PMF (Positive Matrix Factorization) method. The one at Mallorca is more strongly influenced by anthropogenic emissions. Since the goal of this article is to evaluate the performance of different SOA simulation schemes in the western Mediterranean area, it is more representative to take into account campaign data from these two sites. The use of previously untapped data from a second site at Mallorca not only enriches the comparison, but also opens the research and analysis work to different mixtures of anthropogenic and biogenic emissions, allowing to evaluate and discuss further the capabilities of the model.

(2) Our work provides, in addition to that of OA, a detailed comparison of meteorological inputs, SOA precursors (isoprene and monoterpenes) and intermediate compounds (MVK+MACR) for both sites while Chrit et al. (2017) only focuses on OA.

(3) For the Cap Corsica station, because of its unique geographical characteristics not well represented by CTMs, an orographical representativeness error study is performed in our work. This is actually cited in Chrit et al. (2017) to indicate an error for organic aerosol (Chrit et al. 2017, Page 7, section 4, subsection 2). The novelty of this approach is discussed in more detail below.

(4) A source apportionment study for the western Mediterranean basin at two different altitudes (model layers between 0-50m and 300-450m above ground) is also included in our study which is not the case in Chrit et al. (2017). Our shows the impact of biogenic and anthropogenic sources for the formation of OA not only on the surface, but also for a higher altitude.

As a side note, the two articles were written at almost the same time period. While Chrit et al. (2017) was accepted for publication on 23 October 2017, this article was submitted for discussion on 26 August 2017. The Chrit et al. (2017) paper being now published, we cited it at several places:

For example, Chrit et al. (2017) modelled SOA formation in the western Mediterranean area during the ChArMEx summer campaigns, with a surrogate scheme that also contains ELVOCs (Extremely Low Volatile Organic Compounds). The simulated concentrations and properties (oxidation and affinity with water) of organic aerosols agree well to the observations performed at Ersa (Corsica), after they had included the formation of extremely low volatility organic aerosols and organic nitrate from monoterpenes oxidation in the model. (Page 3, lines 26-28)

Chrit et al. (2017) used a two-step surrogate scheme for the simulation of the Ersa site measurements. They found that their modified SOA simulation scheme corresponds well with the data, with a correlation of 0.67 and a Mean Fractional Bias (MFB) of -0.15 for hourly values of the period between June to August 2013. For the period of July to August 2013 for hourly values, we find 0.52 correlation and an MFB of -0.03 for the modified VBS scheme, which shows that both these schemes work reasonably well for the simulated area. (Page 19, line 20-25)

2. “As a result of these facts, I do not find this manuscript very novel.”

Here, we wish to show that our work is indeed novel and original in several important aspects. While we appreciate the comments made by referee 2, we strongly disagree with this statement for the reasons specified in detail below.

5 The Cap Corsica site has unique geographical characteristics which raises issues when comparing
simulations to measurements. Indeed, it is located on the northerly edge of a crest and surrounded from
the west, north and east by slopes falling rapidly to the sea. Its altitude is at 530 m, while in CHIMERE
simulations, even at a 1 km horizontal resolution, it is represented at 360 m height. This height difference
10 might induce differences between simulated and observed concentrations, which need to be assessed.
This is even truer in a marine environment with low boundary layer heights. Therefore, a novel approach
was developed in this work to calculate an orographic representativeness error. This error expresses the
uncertainty in the simulated concentration due to the fact that the orography is not perfectly represented
15 in the model. It is calculated based on the comparison between simulated concentrations at different
altitudes (100-500 m) over a narrow domain, with the measurement site at the center. This method allows
attributing representativeness errors to different species, which turn out to be large for primary
compounds especially emitted in the marine boundary layer like NO_x and BC, and smaller for (mainly)
secondary compounds as O₃ and OA. To the best of our knowledge, this approach has not been used
anywhere else for the estimation of errors produced by orographic representativeness of a site (given
20 that earlier work mostly focusses on problems of horizontal representativeness). This method can be
applied to other sites with the same characteristics (high altitude remote sites with strong change of
altitude over small distances). The results obtained from this approach are also cited in Chrit et al 2017
(Page 7, section 4, sub-section 2). We would argue that this development, while not the primary goal of
the paper, is clearly novel.

20 For the eastern Mediterranean area, relatively ample literature is available both for aerosol chemical
composition (Bardouki et al., 2003 ; Koçak et al., 2007b ; Koulouri et al., 2008 ; Hildebrandt et al.,
2010) and also for OA comparisons to simulations (e.g Fountoukis et al., 2014). However, for the
western part of the basin, less studies exist for the occidental basin with exception of those done for
coastal cities in the area (for example Querol et al., 2009 ; Pey et al., 2013). Furthermore these data have
25 not been used to the best of our knowledge, specifically for CTM model evaluation. Thus, detailed
model-observation comparisons with different aspects of OA (total concentrations, oxidation states,
modern/fossil fractions from ¹⁴C) including comparisons of precursors, intermediary compounds,
meteorological conditions and other gaseous/particulate species for two different sites, as presented in
our paper, have not been performed before for the western Mediterranean area. Comparisons of OA for
30 the Cap Corse area was done by Chrit et al. (2017) with a different scheme and a different model;
however, the comparisons for the Mallorca station are not used in any other articles, and are therefore
clearly new. Also, articles comparing different SOA simulation schemes have not yet focused on this
part of the basin, this basin introduces interesting challenges in this area with its particular characteristics
and different emission sources including continental, coastal, and marine anthropogenic and biogenic
35 emissions.

Source apportionment studies with simulations for the western Mediterranean area is also a subject that
has not been much explored before. Of course, observational based source apportionment analyses have
been presented for different sites of the Mediterranean basin (Koçak et al., 2007 ; Querol et al., 2009 ;
Minguillón et al., 2011 ; Pey et al., 2013) and also for the European area there are studies that discuss
40 the simulated effects of biogenic emissions (Sartelet et al., 2012) on OA. Our study is original and novel,
because it uses a model tied to observations of source apportionment; allowing us to extrapolate results
from two sites to a larger area. This part of our work, answers one of the principal goals of the ChArMEx
campaign directly, that is exploring the source apportionment of OA over the western Mediterranean
area. We think that this part of our paper makes an important contribution to the ACP /AMT special
45 ChArMEx section.

3. “However, no mention of cloud processing in the aerosol modules is mentioned. Why not link the updated VBS with a cloud processing module?”

Cloud scavenging processes are already taken into account in the model. However, because of the unique
geographical characteristics mentioned before for this site, these fog events were not simulated by the

meteorological inputs. Since these decreases concern only a small percentage of the observations, they do not have a major effect on the outcome of these comparisons. While this effect is very visible for sulfates, it is less pronounced for other particulate species such as black carbon and OA.

5 **4. “However, these types of evaluations were done during development of those models.”**

Actually, these schemes were not tested and compared to measurements for the Mediterranean area. They were tested in the USA (Robinson et al., 2007 ; Lane et al., 2008), different parts of Europe (Petetin et al., 2014 ; Zhang et al., 2013 ; Fountoukis et al., 2014) and in South America (Shrivastava et al., 2011 ; Hodzic and Jimenez, 2011 ; Shrivastava et al., 2013 ; Shrivastava et al., 2015). Only in Chrit et al (2017) a scheme for the simulation of organic aerosols is tested for the western Mediterranean, but the evaluated scheme is different from the four schemes evaluated in our work (mainly VBS derived schemes in our work versus a one-step surrogate scheme in Chrit et al. 2017).

Referee 2 specific comments:

15 **1. “Page 4, Line 13. I did not find Table 1 particularly useful. It could be incorporated into the Figure 1 caption. “**

Table 1 has been removed in the revised version of the article, and incorporated in figure 1.

2. “Page 4, Line 33. What is a shipping snap sector?”

20 Shipping SNAP (Selective Nomenclature for Air Pollution) sector is the sector in anthropogenic emissions representing shipping emissions. Some anthropogenic emission inventories use SNAP sectors while others use NFR09 sectors. Both are explained in the references given for MACC-III and Edgar-HTAP emission inventories. In the revised paper, “snap” is changed to SNAP (Selective Nomenclature for Air Pollution).

3. “Page 5, Line 10. Has MELCHIOR been updated since its inception in 2003?”

25 As a response to this comment, the following text is added to the article:

This mechanism has around 120 reactions to describe the whole gas-phase chemistry. The reaction rates used in MELCHIOR are constantly updated (last update in 2015), however, the reaction scheme itself has not been updated since 2003. Some reactions have been added to it by Bessagnet et al. (2009) regarding the oxidation of organic aerosol precursors, but they don't affect gas-phase chemistry. Also, MELCHIOR has been compared to SAPRC-07A, a more recent scheme (Carter, 2010), and the results show acceptable differences between the two schemes, for example, when compared to EEA (European Economic Area) ozone measurements, both produce a correlation coefficient of 0.71. These comparisons are presented in Menut et al. (2013) and Mailler et al. (2017). (Page 5, lines 13-19)

35 **4. “Page 5, Line 13. What is the distribution of particle sizes in the model? That is, the authors provide the range but not any information about how the bins are spaced.”**

The following short explanation was added for this comment:

This module distributes aerosols in a number of size bins, here 10 bins ranging from 40nm to 40µm, in a logarithmic sectional distribution, each bin spanning over a size range of a factor of two (40-20 µm, 20-10 µm, ...). (Page 5, lines 21-23)

40 **5. Page 5, Line 24. Pun and Seigneur were not the source of the experimental data. This citation should be to the original manuscripts.**

References are added for experimental data.

The SOA simulation scheme in CHIMERE (Bessagnet et al., 2008) consists of a single-step oxidation process, where VOC lumped species are directly transformed into SVOC (Semi-volatile Organic Compounds) with yields that are taken from experimental data (Odum et al., 1997 ; Griffin et al., 1999 ; Pun and Seigneur, 2007). (Page 5, line 33-35)

- 5 **6. Page 6, Line 27. Please provide more detail on how this ratio of NO reaction rates is used to determine the low versus high NOx yields. What values of this ratio correspond to low versus high NOx? What are the bounds of this ratio?**

The following explanations are added to the text: (Page 7, lines 1-8)

10 *For this purpose, a parameter (α) is added to the scheme, which calculates the ratio of the reaction rate of RO₂ radicals with NO (v_{NO} ;high-NO_x regime) with respect to the sum of reaction rates of the reactions with HO₂ (v_{HO_2}), and RO₂ (v_{RO_2} ;low-NO_x regime).The parameter α is expressed as follows:*

$$\alpha = \frac{v_{NO}}{v_{NO}+v_{HO_2}+v_{RO_2}} \quad \text{Eq. 1}$$

15 *This α value represents the part of RO₂ radicals reacting with NO (which leads to apply “high NO_x yields”). It is calculated for each grid cell by using the instantaneous NO, HO₂ and RO₂ concentrations in the model. Then, the following equation is used to calculate an adjusted SOA yield using this α value (Carlton et al., 2009).*

$$Y = \alpha \times Y_{highNO_x} + (\alpha - 1) \times Y_{lowNO_x} \quad \text{Eq. 2}$$

- 20 **7. Section 3 Page 7, Line 19. Was only total NOx measured and compared? Were there any issues due to conversion of other NOy species into NO?**

For Ersa, due to its photolytic converter, the CRANOX II instrument measures NO and NO₂ without interferences. However, for Es Pinar interferences from NO_y on NO are possible. Figure 5 was modified to include NO_y time series as well as NO_x to address this issue for Es Pinar. The following text is added to the article as a response:

25 *At Ersa, NO_x (nitrogen oxides) were measured by CraNOx analyzer using ozone chemiluminescence with a resolution of 5 minutes. The photolytic converter in the analyzer allows the conversion of direct measurements of NO₂ into NO in a selective way, thus avoiding interferences with other NO_y species. At Es Pinar, an API Teledyne T200 with molybdenum converter was used; therefore, the measurements are not specific to NO₂ and interferences of NO_y are possible for these measurements. (Page 8, lines 1-5)*

Also, table 1 has been updated for the NOx measurement instrument in Es Pinar. In figure 5, NO_y time series are added to the comparisons of simulated and observed NO_x to take into account the possible interference of NO_y in the measurements. Statistics for the comparison of simulated NO_y to NO_x measurements are added to table 3.

- 35 **8. Page 8, Lines 2 and 5. I do not understand why a constant value of HOA was assumed for each PMF. This needs to be clarified as model results are later compared to these values.**

40 A constant value of HOA was not considered for none of the station. The a-value mentioned in the article refers to the extent to which the output HOA factor is allowed to vary from the input HOA reference mass spectra (i.e. 10% for Ersa and 5% for Cap Es Pinar). The a value is a scalar ranging from 0 to 1 (i.e. 0 to 100%). If the a value is set to 1 then we are in the case of a pure unconstrained PMF approach and if the a value is set to 0 then we are in the case of a Chemical Mass Balance approach (CMB). The use of the a-value approach still accounts for variability of the profile. This method is fully described in Canonaco et al., (2013) and Sturtz, (2014). In such remote environments such as the two
45 stations we are using, the HOA factor could not be extracted from the OA mass spectral matrix with a

classical unconstrained PMF. In order to assess the impact of primary fossil fuel combustions, the HOA factor must be constrained using a reference HOA mass spectra. Note that only the HOA factor was constrained, all the other factors being fully free.

5 *For the Ersa site, HOA (Hydrocarbon-like Organic Aerosol) profile was constrained with a reference HOA factor using an a value of 0.1. The a value refers to the extent to which the output HOA factor is allowed to vary from the input HOA reference mass spectra (i.e. 10% in this case, Canonaco et al., 2013). In such remote environment, the HOA factor could not be extracted from the OA mass spectral matrix with a classic unconstrained PMF approach. Two other factors were extracted, without any constraints, including SVOOA (Semi-Volatile Oxygenated Organic Aerosol) and LVOOA (Low-Volatile Oxygenated Organic Aerosol). For the Es Pinar site, HOA has been constrained using an a-value of 0.05. Three additional factors were retrieved, including an SVOOA (Semi-Volatile Oxygenated Organic Aerosol) and 2 LVOOA (Low Volatile Oxygenated Organic Aerosol) factors. (Page 8, lines 23-31)*

15 **9. Section 4 Page 9, Line 6. The wording here should be changed to indicate that it is the simulated height of the cell in which the site was located was underpredicted – not that the simulated height of the site was underpredicted (since the authors are not actually simulating the height of the site).**

Yes, this is true, the phrasing is changed in the article to the following:

20 *For the 10-km domain (D10), we noticed that there was an inconsistency between simulated and real altitude of the cell where the Ersa site is located. (Page 9, lines 29-31)*

10. Page 9, Line 27. Higher than a fixed value? What value? What was its basis?

As mentioned in the article, more detailed explanation about the method of the calculation of ORE (Orographic Representativeness Error) is presented in SI3. The threshold value for correlation between fitted and simulated values mentioned in the article is 0.50, but the method was tested with 0.60 and 0.70 correlations as well, but no significant changes were seen in the resulting ORE for any of the species. The threshold value was chosen somewhat arbitrarily, for assuring satisfying correlation.

11. Page 9, Line 39. The authors state that secondary species appear less influenced by the orographic uncertainty. However, the impact on MVK + MACR is as large as that for primary species. The authors need to be careful with their word choice.

This is true, the phrasing of the paragraph should have been more precise. It is changed to the following:

35 *A general conclusion is that secondary pollutants with higher atmospheric lifetimes appear to be well represented from a geographic point of view. On the contrary, model-observation comparisons for more reactive primary and secondary pollutants with short lifetimes (primary such as NO_x and reactive secondary such as MVK+MACR) should be performed with caution keeping in mind the fact that the simulated altitude is not representative of the orography for this specific station. (Page 10, lines 27-31)*

12. Page 10, Line 35. I do not believe that this last sentence is necessary.

Sentence removed.

13. Page 11, Line 7. The caption from Table 4 needs to refer to Figure 5.

Figure number corrected.

14. Page 11, Line 32. Does this requirement of a regional look at the aerosol call into question the use of two sites?

No, it does not. Since the regional look mentioned here refers to precursors of SOA, meaning isoprene and mono-terpenes (as also mentioned in the article). These two are unstable components with short lifetimes, thus, it is necessary to have a regional look to be able to compare them to simulations more

accurately. However, OA has a much longer lifetime, making it much more stable, therefore, two stations can be enough for an accurate observation-simulation comparison.

5 **15. Page 12, Line 19. Here is evidence for cloud processing (again, it is presented for BC later). However, no mention of cloud processing in the aerosol modules is mentioned. Why not link the updated VBS with a cloud processing module? That would certainly enhance the novelty of the work.**

The following has been added to the text:

10 *Cloud scavenging processes are already taken into account in the model. However, because of the unique geographical characteristics mentioned before for this site, the meteorological inputs did not simulate these fog events and therefore cloud scavenging was not activated in the simulation. Since these decreases concern only a small percentage of the observations, they do not have a major effect on the outcome of these comparisons. While this effect is very visible for sulfates, it is less pronounced for other particulate species such as black carbon and OA. (Page 13, lines 10-15)*

15 **16. Page 13, Line 3. I do not find this summary paragraph necessary.**

The summary paragraph has been removed.

17. Page 13, Line 27. Figure 7 caption has an error in the word standard.

The word “standard” has been modified in the figure 7 caption.

18. Page 13, Lines 30 and 31. Please provide standard deviations on the averages.

20 Standard deviation values were added for table 5.

19. Page 14, Line 12. What is meant by ‘respected’?

The phrase was removed.

20. Page 14, Line 34. I do not see much use for Table 7. This text can be included in the main body of the manuscript.

25 Table 7 has been removed and its contents incorporated in the manuscript.

30 *ASOA is considered to be in the fossil fraction and BSOA in the non-fossil fraction. For carbonaceous aerosol, residential/domestic uses are considered as non-fossil as they are mostly related to wood burning (Sasser et al., 2012). Therefore, they are attributed to the non-fossil bin (3.6% for BC and 12.3% for OC, Sasser et al., 2012). The non-fossil contribution of ASOA and POA due to biofuel usage is ignored here, as it is minor (<5%).(Page 15, lines 12-16)*

21. Page 15 and beyond, The colors in Figures 8, 9, and 11 need to be differentiated to a greater extent if they are included in any future submission.

35 White spaces and numbers were added make these images clearer for the reader. But the color palette was not changed, since the plot separates schemes by their order, and not by their color. The only colors that are of importance in these figures are green in its different shades for non-fossil OA and brown in its different shades for fossil OA.

22. Page 17, Line 15. I find this summary paragraph unnecessary.

The summary paragraph is removed.

40 **Reviewer comments:**

Thank you

1. p3 , line 38 at at

Fixed in the final version

2. p5, line 16 don't -> do not

Fixed in the final version

3. Please avoid acronyms inside figure captions, give the definition.

Image legends were revised and explanations are added for all acronyms.

- 5 **4. I recommend to do something very simple such as adding one paragraph to introduce the focus on western Mediterranean, the use of the Orographic Representativeness Error and the consideration of factors other than OA for model evaluation.**

The following phrases were added to the introduction:

10 *This enables us to perform model-observation comparisons with unprecedented detail over this region, including not only the OA concentration, but also its origin (I4C analyses), its oxidation state (PMF method results). In addition, a comparison for meteorological parameters and gaseous/particulate species which could affect the production of OA or could help analyze the robustness of the used schemes has been performed. Moreover, because of the orographic complexity of one of the sites (Ersa, Cap Corse) explored in this work, a novel method is*
15 *designed to calculate the orographic representativeness error of different species. To the best of our knowledge, this is the first time that the concentrations of precursors, intermediary products and OA concentrations and properties have been simulated for different OA simulation schemes and compared for each scheme to multiple series of measurements in different stations for the western Mediterranean area.*

- 20 Also, a phrase was added to the abstract to explain about ORE.

Because of the complex orography of the Ersa site, an original method for calculating an orographic representativeness error (ORE) has been developed.

References:

- 25 Bardouki, H., Liakakou, H., Economou, C., Sciare, J., Smolík, J., Ždímal, V., Eleftheriadis, K., Lazaridis, M., Dye, C. and Mihalopoulos, N.: Chemical composition of size-resolved atmospheric aerosols in the eastern Mediterranean during summer and winter, *Atmos. Environ.*, 37(2), 195–208, doi:10.1016/S1352-2310(02)00859-2, 2003.
- 30 Chrit, M., Sartelet, K., Sciare, J., Pey, J., Marchand, N., Couvidat, F., Sellegri, K. and Beekmann, M.: Modelling organic aerosol concentrations and properties during ChArMEx summer campaigns of 2012 and 2013 in the western Mediterranean region, *Atmos. Chem. Phys. Discuss.*, 1–33, doi:10.5194/acp-2017-312, 2017.
- 35 Fountoukis, C., Megaritis, A. G., Skyllakou, K., Charalampidis, P. E., Pilinis, C., Denier Van Der Gon, H. A. C., Crippa, M., Canonaco, F., Mohr, C., Prévôt, A. S. H., Allan, J. D. and Poulain, L.: Organic aerosol concentration and composition over Europe: insights from comparison of regional model predictions with aerosol mass spectrometer factor analysis, *Atmos. Chem. Phys.*, 14, 9061–9076, doi:10.5194/acp-14-9061-2014, 2014.
- 40 Hildebrandt, L., Engelhart, G. J., Mohr, C., Kostenidou, E., Lanz, V. A., Bougiatioti, A., Decarlo, P. F., Prevot, A. S. H., Baltensperger, U., Mihalopoulos, N., Donahue, N. M. and Pandis, S. N.: Aged organic aerosol in the Eastern Mediterranean: the Finokalia Aerosol Measurement Experiment – 2008, *Atmos. Chem. Phys. Atmos. Chem. Phys.*, 10, 4167–4186, doi:10.5194/acp-10-4167-2010, 2010.

- Hodzic, A. and Jimenez, J. L.: Modeling anthropogenically controlled secondary organic aerosols in a megacity: a simplified framework for global and climate models, *Geosci. Model Dev.*, 4(4), 901–917, doi:10.5194/gmd-4-901-2011, 2011.
- 5 Koçak, M., Mihalopoulos, N. and Kubilay, N.: Chemical composition of the fine and coarse fraction of aerosols in the northeastern Mediterranean, *Atmos. Environ.*, 41(34), 7351–7368, doi:10.1016/j.atmosenv.2007.05.011, 2007a.
- Koçak, M., Mihalopoulos, N. and Kubilay, N.: Contributions of natural sources to high PM10 and PM2.5 events in the eastern Mediterranean, *Atmos. Environ.*, 41(18), 3806–3818, doi:10.1016/j.atmosenv.2007.01.009, 2007b.
- 10 Koulouri, E., Saarikoski, S., Theodosi, C., Markaki, Z., Gerasopoulos, E., Kouvarakis, G., Makela, T., Hillamo, R. and Mihalopoulos, N.: Chemical composition and sources of fine and coarse aerosol particles in the Eastern Mediterranean, *Atmos. Environ.*, 42(26), 6542–6550, doi:10.1016/j.atmosenv.2008.04.010, 2008.
- Lane, T. E., Donahue, N. M. and Pandis, S. N.: Simulating secondary organic aerosol formation using the volatility basis-set approach in a chemical transport model, *Atmos. Environ.*, 42(32), 7439–7451, doi:10.1016/j.atmosenv.2008.06.026, 2008.
- 15 Minguillón, M. C., Perron, N., Querol, X., Szidat, S., Fahrni, S. M., Alastuey, A., Jimenez, J. L., Mohr, C., Ortega, A. M., Day, D. A., Lanz, V. A., Wacker, L., Reche, C., Cusack, M., Amato, F., Kiss, G., Hoffer, A., Decesari, S., Moretti, F., Hillamo, R., Teinilä, K., Seco, R., Peñuelas, J., Metzger, A., Schallhart, S., Müller, M., Hansel, A., Burkhardt, J. F., Baltensperger, U. and Prévôt, A. S. H.: Fossil versus contemporary sources of fine elemental and organic carbonaceous particulate matter during the DAURE campaign in Northeast Spain, *Atmos. Chem. Phys.*, 11(23), 12067–12084, doi:10.5194/acp-11-12067-2011, 2011.
- Petetin, H., Beekmann, M., Sciare, J., Bressi, M., Rosso, A., Sanchez, O. and Gherzi, V.: A novel model evaluation approach focusing on local and advected contributions to urban PM2.5 levels - Application to Paris, France, *Geosci. Model Dev.*, 7(4), 1483–1505, doi:10.5194/gmd-7-1483-2014, 2014.
- 25 Pey, J., Pérez, N., Cortés, J., Alastuey, A. and Querol, X.: Chemical fingerprint and impact of shipping emissions over a western Mediterranean metropolis: Primary and aged contributions, *Sci. Total Environ.*, 463–464, 497–507, doi:10.1016/j.scitotenv.2013.06.061, 2013.
- 30 Querol, X., Alastuey, a., Pey, J., Cusack, M., Pérez, N., Mihalopoulos, N., Theodosi, C., Gerasopoulos, E., Kubilay, N. and Koçak, M.: Variability in regional background aerosols within the Mediterranean, *Atmos. Chem. Phys. Discuss.*, 9(2), 10153–10192, doi:10.5194/acpd-9-10153-2009, 2009.
- Robinson, A. L., Donahue, N. M., Shrivastava, M. K., Weitkamp, E. A., Sage, A. M., Grieshop, A. P., Lane, T. E., Pierce, J. R. and Pandis, S. N.: Rethinking Organic Aerosols: Semivolatile Emissions and Photochemical Aging, *Science (80-.)*, 315(5816), 1259–1262, doi:10.1126/science.1133061, 2007.
- 35 Sartelet, K. N., Couvidat, F., Seigneur, C. and Roustan, Y.: Impact of biogenic emissions on air quality over Europe and North America, *Atmos. Environ.*, 53, 131–141, doi:10.1016/J.ATMOSENV.2011.10.046, 2012.
- Shrivastava, M., Fast, J., Easter, R., Gustafson, W. I., Zaveri, R. A., Jimenez, J. L., Saide, P. and Hodzic, A.: Modeling organic aerosols in a megacity: Comparison of simple and complex representations of the volatility basis set approach, *Atmos. Chem. Phys.*, 11(13), 6639–6662, doi:10.5194/acp-11-6639-2011, 2011.
- 40 Shrivastava, M., Zelenyuk, A., Imre, D., Easter, R., Beranek, J., Zaveri, R. A. and Fast, J.: Implications of low volatility SOA and gas-phase fragmentation reactions on SOA loadings and their spatial and

temporal evolution in the atmosphere, *J. Geophys. Res. Atmos.*, 118(8), 3328–3342, doi:10.1002/jgrd.50160, 2013.

Shrivastava, M., Easter, R. C., Liu, X., Zelenyuk, A., Singh, B., Zhang, K., Ma, P., Chand, D., Ghan, S., Jimenez, J. L., Zhang, Q., Fast, J., Rasch, P. J. and Tiitta, P.: Global transformation and fate of SOA: Implications of low-volatility SOA and gas-phase fragmentation reactions, *J. Geophys. Res. Atmos.*, 1–27, doi:10.1002/2014JD022563. Received, 2015.

Zhang, Q. J., Beekmann, M., Drewnick, F., Freutel, F., Schneider, J., Crippa, M., Prevot, A. S. H., Baltensperger, U., Poulain, L., Wiedensohler, A., Sciare, J., Gros, V., Borbon, A., Colomb, A., Michoud, V., Doussin, J. F., Denier Van Der Gon, H. A. C., Haefelin, M., Dupont, J. C., Siour, G., Petetin, H., Bessagnet, B., Pandis, S. N., Hodzic, A., Sanchez, O., Honoré, C. and Perrussel, O.: Formation of organic aerosol in the Paris region during the MEGAPOLI summer campaign: Evaluation of the volatility-basis-set approach within the CHIMERE model, *Atmos. Chem. Phys.*, 13(11), 5767–5790, doi:10.5194/acp-13-5767-2013, 2013.

15

20

25

30

35

Simulation of fine organic aerosols in the western Mediterranean area during the ChArMEx 2013 summer campaign

Arineh Cholakian^{1,2}, Matthias Beekmann¹, Augustin Colette², Isabelle Coll¹, Guillaume Siour¹, Jean Sciare^{3,5}, Nicolas Marchand⁴, Florian Couvidat², Jorge Pey^{4*}, Valerie Gros³, Stéphane Sauvage^{6**}, Vincent Michoud^{96**}, Karine Sellegri⁷, Aurélie Colomb⁷, Karine Sartelet⁸, Helen Langley DeWitt⁴, Miriam Elser^{910***}, André S. H. Prévot⁹¹⁰, Sonke Szidat¹⁰⁺, and François Dulac³

1 Laboratoire Inter-Universitaire des Systèmes Atmosphériques (LISA), UMR CNRS 7583, Université Paris Est Créteil et Université Paris Diderot, Institut Pierre Simon Laplace, Créteil, France (arineh.cholakian@lisa.u-pec.fr)

2 Institut National de l'Environnement Industriel et des Risques, Parc Technologique ALATA, Verneuil en Halatte, France

3 Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, Gif-sur-Yvette, France

4 Aix-Marseille Université, CNRS, LCE FRE 3416, Marseille, 13331, France

5 The Cyprus Institute, Energy, Environment and Water Research Center, Nicosia, Cyprus

6 IMT Lille Douai, Univ. Lille, Département Sciences de l'Atmosphère et Génie de l'Environnement, F-59000 Lille, France

7 LAMP, Campus universitaire des Cezeaux, 4 Avenue Blaise Pascal, 63178 Aubiere, France

8 CEREA, Joint Laboratory École des Ponts ParisTech – EDF R & D, Université Paris-Est, 77455 Marne la Vallée, France

~~9 Mines Douai, Département Sciences de l'Atmosphère et Génie de l'Environnement (SAGE), 59508 Douai, France~~

~~910 Paul Scherrer Institute, 5232 Villigen-PSI, Switzerland~~

~~10+ University of Bern, Freiestrasse 3, CH-3012 Bern, Switzerland~~

* Now at the Spanish Geological Survey, IGME, 50006 Zaragoza, Spain

** Now at ~~Mines Douai, Département Sciences de l'Atmosphère et Génie de l'Environnement (SAGE), 59508 Douai, France~~ laboratoire Inter-Universitaire des Systèmes Atmosphériques (LISA), UMR CNRS 7583, Université Paris Est

*** Laboratory for Advanced Analytical Technologies, Empa, Dübendorf, 8600, Switzerland

Abstract. The simulation of fine organic aerosols with CTMs (Chemistry Transport Models) in the western Mediterranean basin has not been studied until recently. The ChArMEx (the Chemistry-Aerosol Mediterranean Experiment) SOP 1b2 (Special Observation Period 1b2) intensive field campaign in summer of 2013 gathered a large and comprehensive dataset of observations allowing the study of different aspects of the Mediterranean atmosphere including the formation of organic aerosols (OA) in 3D models. In this study, we used the CHIMERE CTM to perform simulations for the duration of the SAFMED (Secondary Aerosol Formation in the MEDiterranean) period (July to August 2013) of this campaign. In particular, we evaluated four schemes for the simulation of OA, including the CHIMERE standard scheme, the VBS (Volatility Basis Set) standard scheme with two parameterizations including aging of biogenic secondary OA, and a modified version of the VBS scheme which includes fragmentation and formation of non-volatile OA. The results ~~from~~ these four schemes are compared to observations at two stations in the western Mediterranean basin, located ~~on Ersa, in~~ Cap Corse (Corsica Isl, France) and at Cap Es Pinar (Mallorca Isl, Spain). These observations include OA mass concentration, PMF (positive matrix factorization) results of different OA fractions, and ¹⁴C observations showing the fossil or non-fossil origins of carbonaceous particles. Because of the complex orography of the Ersa site, an original method for calculating an orographic representativeness error (ORE) has been developed. It is concluded that the modified VBS scheme is close to observations in all three aspects mentioned above; the standard VBS scheme without BSOA (Biogenic Secondary Organic Aerosol) aging also has a satisfactory performance in simulating the mass

concentration of OA, but not for the source origin analysis comparisons. In addition, the OA sources over the western Mediterranean basin are explored. OA shows a major biogenic origin, especially at several hundred meters height from the surface; however over the Gulf of Genoa near the surface, the anthropogenic origin is of similar importance. A general assessment of other species was performed to evaluate the robustness of the simulations for this particular domain before evaluating OA simulation schemes. It is also shown that the Cap Corse site presents important orographic complexity which makes comparison between model simulations and observations difficult. A method was designed to estimate an orographic representativeness error for ~~a list of species~~ measured at Ers and yields an uncertainty of between 50-85% for primary pollutants, and around 2-10% for secondary species, ~~for these species model to observations comparisons are only little impacted by orography.~~

10 **1 Introduction**

The Mediterranean basin is subject to multiple emission sources; anthropogenic emissions that are transported from adjacent continents or are produced within the basin, local or continental biogenic and natural emissions among which the dust emissions from northern Africa can be considered as an important source (Pey et al., 2013 ; Vincent et al., 2016). All these different sources, the geographic particularities of the region favoring accumulation of pollutants (Gangoiti et al., 2001), the prevailing meteorological conditions favorable ~~to~~ intense photochemistry and thus secondary aerosol formation, make the Mediterranean area a region experiencing a heavy burden of aerosols (Monks et al., 2009 ; Nabat et al., 2012). In the densely populated coastal areas, this aerosol burden constitutes a serious sanitary problem considering the harmful effects of fine aerosols on human health (Martinelli et al., 2013). In addition, studies have shown that the Mediterranean area could be highly sensitive to future climate change effects (Giorgi, 2006 ; Lionello and Giorgi, 2007). This could affect aerosol formation processes, but in turn the aerosol load also affects regional climate (Nabat et al., 2013). These interactions, the high aerosol burden in the area, and its health impact related to the high population density situated around the basin makes this region particularly important to study.

The aforementioned primary emissions can be in form of gaseous species, as particulate matter, or as semi-volatile species distributed between both phases (Robinson et al., 2007a). In the atmosphere, they can subsequently undergo complex chemical processes lowering their volatility which leads to the formation secondary particles. These processes are not entirely elucidated especially for the formation of SOA (Secondary Organic Aerosols, for example (Kroll and Seinfeld, 2008), starting from initially emitted biogenic or anthropogenic VOC (Volatile Organic Compounds) and SVOC (semi-volatile organic compounds).

The chemical composition of aerosols has been studied in detail in the eastern Mediterranean area (Lelieveld et al., 2002 ; Bardouki et al., 2003 ; Sciare et al., 2005 ; Koçak et al., 2007 ; Koulouri et al., 2008 ; Sciare et al., 2008) and to a less extent in the western part (Sellegrì et al., 2001 ; Querol et al., 2009 ; Minguillón et al., 2011 ; Ripoll et al., 2014 ; Menut et al., 2015 ; Arndt et al., 2017). Little focus has been given on the formation of organic aerosol (OA) over the western Mediterranean even if OA can play a significant role in both local and global climate (Kanakidou et al., 2005) and can affect health (Pöschl, 2005 ; Mauderly and Chow, 2008). Its contribution has been calculated in studies to be nearly 30% in PM₁ for the eastern Mediterranean area during the FAME 2008 campaign (i.e. Hildebrandt et al., 2010). It is also important to know the contribution of different sources (biogenic, anthropogenic) to the total concentration of OA in the western part of the basin. Such studies have been performed for the eastern Mediterranean basin (e.g. Hildebrandt et al., 2010), while for the western part of the basin, they

have been in general restricted to coastal cities such as Marseilles and Barcelona (El Haddad et al., 2011 ; Mohr et al., 2012 ; Ripoll et al., 2014).

The ChArMEx (the Chemistry Aerosol Mediterranean Experiment; <http://charmex.lsce.ipsl.fr>) project was organized in this context, with a focus over the western Mediterranean basin during the period of 2012-2014, in order to better assess the sources, formation, transformation and mechanisms of transportation of gases and aerosols. During this project, detailed measurements were acquired not only for the chemical composition of aerosols, but also for a large number of gaseous species from both ground-based and airborne platforms. The project ChArMEx is divided into different sub-projects, each with a different goal; among those, the SAFMED (Secondary Aerosol Formation in the MEDiterranean) project aimed at understanding and characterizing the concentrations and properties of OA in the western Mediterranean (for example Nicolas, 2013 ; Di Biagio et al., 2015 ; Chrit et al., 2017 ; Arndt et al., 2017 ; Freney et al., ~~submitted~~, 2017 ; Pey et al., ~~2017~~ in prep). To reach these goals, two intense ground-based and airborne campaigns were organized during July-August of 2013 and also summer of 2014. The focus of the present study is on the SAFMED campaign in summer of 2013, since detailed measurements on the formation of OA and precursors were obtained during this period namely at Ersu (Corsica) and Es Pinar (Mallorca). Other ChArMEx sub-projects and campaigns included the TRAQA (Transport et Qualité de l'Air) campaign in summer 2012 set-up to study the transport and impact of continental air on atmospheric pollution over the basin (Sič et al., 2016), the ADRIMED (Aerosol Direct Radiative Impact in the Mediterranean, June –July 2013) campaign aiming at understanding and assessing the radiative impact of various aerosol sources (Mallet et al., 2016).

Modelling of aerosol processes and properties is a difficult task. Aside from the lack of knowledge of aerosol formation processes, the difficulty lies in the fact that organic aerosols present an amalgam of thousands of different species that cannot all be represented in a 3D chemistry transport model due to limits in computational resources. Therefore a small number of lumped species with characteristics that are thought to be representative of all the species in each group are used instead. There are many different approaches that can be used in creating representative groups for organic aerosols (e.g. which characteristics to use to group the species, which species to lump together, physical processes that should be presented for their simulation...). It is therefore necessary to test these different simulation schemes for organic aerosols in different regions and compare the results to experimental data to check for their robustness. For example, Chrit et al. (2017) modelled SOA formation in the western Mediterranean area during the ChArMEx summer campaigns, with a surrogate scheme that also contains ELVOCs (Extremely Low Volatile Organic Compounds). –The simulated concentrations and properties (oxidation and affinity with water) of organic aerosols agree well to the observations performed at Ersu (Corsica), after they had included the formation of extremely low volatility organic aerosols and organic nitrate from monoterpene oxidation in the model. ~~In this work, different configurations of the Volatility Basis Set (Donahue et al., 2006 ; Robinson et al., 2007 ; Shrivastava et al., 2013) and the base parameterization of the CHIMERE 3D model (Bessagnet et al., 2008 ; Menut et al., 2013) are used for this purpose. Although Chrit et al. (2017) modelled SOA formation in the western Mediterranean during the ChArMEx summer campaigns, they used a surrogate approach rather than a VBS approach in their air quality modeling platform. The simulated concentrations and properties (oxidation and affinity with water) of organic aerosols agree well to the observations performed at Ersu (Corsica), after they had modified their model to include the formation of extremely low volatility organic aerosols and organic nitrate from monoterpene oxidation.~~

The present study focuses on the comparison of different OA formation schemes implemented in the CHIMERE chemistry-transport model for simulation of OA over the western Mediterranean area, ~~In this work, Different configurations of the Volatility Basis Set (Donahue et al., 2006 ; Robinson et al., 2007 ; Shrivastava et al., 2013) and the base parameterization of the CHIMERE 3D model (Bessagnet et al., 2008 ; Menuet et al., 2013) are used~~ for this purpose. Our work takes advantage of the ~~with an~~ extensive experimental data pool obtained during the SAFMED campaign. This enables us to perform model-observation comparisons with unprecedented detail over this region, including not only the OA concentration, but also its origin (¹⁴C analyses), its oxidation state (PMF method results). In addition, a comparison for meteorological parameters and gaseous/particulate species which could affect the production of OA or could help analyze the robustness of the used schemes has been performed. Moreover, because of the orographic complexity of one of the sites (Ersa, Cap Corse) explored in this work, a novel method is designed to calculate the orographic representativeness error of different species. To the best of our knowledge, this is the first time that the concentrations of precursors, intermediary products and OA concentrations and properties have been simulated for different OA simulation schemes and compared for each scheme to multiple series of measurements in different stations for the western Mediterranean area. For OA schemes, the paper aims at ~~comparisons aim~~ at assessing the robustness of each scheme with regard to different criteria as mass, fossil and modern fraction ~~and~~ -, volatility. Section 2 describes the model and the inputs used for the simulations. Also, the evaluated schemes are explained in this section in more detail. The experimental data used for simulation-observation comparisons are discussed in section 3. An overall validation of the model is presented in section 4, together with comparisons of different gaseous and particulate species and meteorological parameters to observations. In section 5, comparison of implemented schemes to measurements regarding concentration, oxidation state and origin of organic aerosol are presented. In section 6, the contribution of different sources to the organic aerosol present in the whole basin is explored, before the conclusion in section 7.

2 Model setup

The CHIMERE model (Menuet et al., 2013; <http://www.lmd.polytechnique.fr/chimere>) is an offline regional CTM (Chemistry-Transport Model) which has been tested rigorously for Europe and France (Zhang et al., 2013 ; Petetin et al., 2014; Colette et al., 2015; Menuet et al., 2015 ; Rea et al., 2015). It is also widely used in both research and forecasting activities in France, Europe and other countries (Hodzic and Jimenez, 2011). In this work, a slightly modified version of the CHIMERE 2014b configuration is used to perform the simulations. The modifications concern an updated description of the changes in aerosol size distribution due to condensation and evaporation processes (Mailler et al., 2016). Four domains are used in the simulations; a coarse domain covering whole Europe and Northern Africa with a 30km resolution, and three nested domains inside the coarse domain with resolutions of 10km, 3km and 1km (figure 1). The 10km resolved domain covers the western Mediterranean area and the two smaller domains are centered on the Cap Corse, where the main field observations in SAFMED were performed. Such highly resolved domains are necessary to resolve the complex orography of Cap Corse ground-based measurements site which will be discussed in section 4, while for the flatter Es Pinar site, a 10km resolution is sufficient. The simulations for each domain contain 15 vertical levels starting from 50 m to about 12 km above sea level (asl) with an average vertical resolution of 400 m within the continental boundary layer (CBL) and 1 km above ~~The simulations for each domain contain 15 vertical levels starting from 50m to about 12km asl (vertical resolution of an average of 400m within the continental boundary layer (CBL) and 1km above CBL).~~ The

CHIMERE model needs as mandatory inputs a set of gridded data: meteorological data, emission data for both biogenic and anthropogenic sources, landuse parameters, boundary/limit conditions and other optional inputs such as dust and fire emissions. Given these inputs, the model produces the concentrations and deposition fluxes for major gaseous and particulate species and also intermediate compounds. -The simulations presented in this article cover the period of 1 month from July 10th to August 9th.

Meteorological inputs are calculated with the WRF (Weather Research Forecast) regional model (Wang et al., 2015) forced by NCEP (National Centers of Environmental Predictions) reanalysis meteorological data (<http://www.ncep.noaa.gov>) with a base resolution of 1°. Slightly larger versions of each domain with the same horizontal resolutions (Figure 1) are used for the meteorological simulations. The WRF configuration used for this study consists of the Single Moment-5 class microphysics scheme (Hong et al., 2004), the RRTMG radiation scheme (Mlawer et al., 1997), the Monin-Obukhov surface layer scheme (Janjic, 2003), and the NOAA Land Surface Model scheme for land surface physics (Chen et al., 2001). Sea surface temperature update, surface grid nudging (Liu et al., 2012, Bowden et al., 2012), ocean physics and topographic wind options are activated. Also, the feedback option is activated, meaning that the simulation of the nested domains can influence the parent domain. Some observation-simulation comparisons are presented for meteorological parameters in section 4.

Anthropogenic emissions for all but shipping ~~SNAP (Selective Nomenclature for Air Pollution)~~ SNAP (Selective Nomenclature for Air Pollution) sectors come from the HTAP-V2 (Hemispheric Transport of Air Pollution, (http://edgar.jrc.ec.europa.eu/htap_v2/index.php) inventory. The shipping sector in this inventory was judged to overestimate ship traffic around the Cap Corse area, especially on the shipping lines between Marseilles and Corsica Island, due to overweighing ferries with respect to cargos (Van der Gon, personal communication). This could, which can be explained by the fact that the boat traffic description is based on voluntary information. Therefore HTAP-V2 shipping emissions were replaced by those of the MACC-III inventory (Kuenen et al., 2014). The base resolution of HTAP inventory is 10km × 10km and that of MACC-III inventory is 7km × 7km. For both inventories the emissions for the year 2010 were used, since this year was the latest common year in the two inventories.

Biogenic emissions are calculated using MEGAN (Model of Emissions of Gases and Aerosols from Nature, Guenther et al., 2006) including isoprene, limonene, α -pinene, β -pinene, ocimene and other monoterpenes with a base horizontal resolution of 0.008° × 0.008°. The land-use data comes from Globcover (Arino et al., 2008) with a base resolution of 300m × 300m. Initial and boundary conditions of chemical species are taken from the climatological simulations of LMDz-INCA3 (Hauglustaine et al., 2014) for gaseous species and GOCART (Chin et al., 2002) for particulate matter.

The chemical mechanism used for the baseline gas-phase chemistry is the MELCHIOR2 scheme (Derognat et al., 2003). This mechanism has around 120 reactions to describe the whole gas-phase chemistry. The reaction rates used in MELCHIOR are constantly updated (last update in 2015), however, the reaction scheme itself has not been updated since 2003. Some reactions have been added to it by Bessagnet et al. (2009) regarding the oxidation of organic aerosol precursors, but they do not affect gas-phase chemistry. Also, MELCHIOR has been compared to SAPRC-07A, a more recent scheme (Carter, 2010), and the results show acceptable differences between the two schemes, for example, when compared to EEA (European Economic Area) ozone measurements, both produce a correlation coefficient of 0.71. These comparisons are presented in Menut et al. (2013) and Mailler et al. (2017).

The CHIMERE aerosol module is responsible for the simulation of physical and chemical processes that influence the size distribution and chemical speciation of aerosols (Bessagnet et al., 2008). This module distributes aerosols in a number of size bins, here 10 bins ranging from 40nm to 40µm, ~~in a logarithmic sectional distribution, each bin spanning over a size range of a factor of two (40-20 µm, 20-10 µm, ...)~~, in a logarithmic sectional distribution, each bin spanning over a size range of a factor of two (40-20 µm, 20-10 µm, ...). The module also addresses coagulation, nucleation, condensation, as well as dry and wet deposition processes. The basic chemical speciation includes Elemental Carbon (EC), sulfate, nitrate, ammonium, SOA, dust, salt and PPM (primary particulate matter other than ones mentioned above).

2.2 Organic aerosol simulation

10 The SOA particles are divided, depending on their precursors, into two groups: ASOA (anthropogenic SOA), and BSOA (biogenic SOA). Four schemes were tested to simulate their formation, more detail on each scheme is presented below.

2.2.1 CHIMERE standard scheme

15 The ~~scheme for simulation of~~ SOA simulation scheme in CHIMERE (Bessagnet et al., 2008) consists of a single-step oxidation process, where VOC lumped species are directly transformed into SVOC (Semi-volatile Organic Compounds) with yields that are taken from experimental data (Odum et al., 1997 ; Griffin et al., 1999 ; Pun and Seigneur, 2007). ~~These SVOC species are and~~ then ~~are~~ distributed into gaseous and particulate phases (figure 2-a) following the partitioning theory of Pankow (Pankow, 1987). The precursors for this scheme are presented in the supplementary information (~~(SII)~~-(S11)). A number of 11 semi-volatile surrogate compounds are formed from these
20 precursors, which include six hydrophilic species, three hydrophobic species and two surrogates for isoprene oxidation. The sum of all 11 species results in the concentration of simulated SOA in this scheme.

2.2.2 VBS scheme

As an alternative to single step schemes like the one in CHIMERE, the VBS (volatility basis set) approach was developed. In these types of schemes, SVOC are divided into volatility bins regardless of their chemical
25 characteristics, but only depending on their saturation concentration. Therefore it becomes possible to add aging processes in the simulation of OA by adding reactions that shift species from one volatility bin to another (Donahue et al., 2006). This scheme was implemented and tested in CHIMERE for Mexico city (Hodzic and Jimenez, 2011) and the Paris region (Zhang et al., 2013). The volatility profile used for this scheme consists of nine volatility bins with saturation concentrations in the range of 0.01 to 10⁶ µg.m⁻³ (convertible to saturation pressure using
30 atmospheric standard conditions), across which the emissions of SVOC and IVOC (Intermediate Volatility Organic Compounds) are distributed, following a specific aggregation table (Robinson et al., 2007). Four volatility bins are ~~used~~ for anthropogenic and biogenic SOA ranging from 1 to 1000 µg.m⁻³. SOA yields are taken from the literature (Lane et al., 2008 ; Murphy and Pandis, 2009) using low-NO_x condition (VOC/NO_x > 10ppbC.ppb⁻¹). The SVOC species can age, by decreasing their volatility by one bin independent of their origin
35 with a given constant rate. SVOC species are either (e.g directly emitted or formed from anthropogenic or biogenic VOC precursors) ~~with a given constant rate~~. Fragmentation processes and the production of non-volatile SOAs are ignored in this scheme. In the basic VBS scheme, the BSOA aging processes are usually ignored since they tend

to result in a significant overestimation in biogenic SOA (Lane et al., 2008). Although physically present, their kinetic constants for this aging process are considered the same as anthropogenic compounds and seem to be overestimated. However, in Zhang et al. (2013), including BSOA aging was necessary to explain the observed experimental data. Therefore, in this work, the VBS scheme is evaluated both with and without including the BSOA aging processes. Figure 2-b shows a simplified illustration for anthropogenic and biogenic SOA, while the partition for SVOC is presented in supplementary information (SI2). For all bins, regardless of their origin, the partitioning between gaseous and the particulate phases is performed following Raoult's law and depends on total organic aerosol concentration. Under normal atmospheric conditions, SVOC with the volatility range of 0.01 to $10^3 \mu\text{g.m}^{-3}$ can form aerosols. In total, the sum of 24 species in the model (with 10 size distribution bins each, i.e. 240 species in total) makes up the total concentration of SOA simulated by this scheme.

2.2.3 Modified VBS scheme

The basic VBS scheme does not include fragmentation processes, corresponding to the break-up of oxidized OA compounds in the atmosphere into smaller and thus more volatile molecules (Shrivastava et al., 2011). It also does not include the formation of non-volatile SOA, where SOA can become non-volatile after formation (Shrivastava et al., 2015). In this work, these two processes were added to the VBS scheme presented above and tested for the Mediterranean basin. The volatility bins for the VBS model were not changed (ranges presented in the previous section). SOA yields were kept as in the standard VBS scheme, however, instead of using the low- NO_x or the high- NO_x regimes, an interpolation between the yields of these two regimes was added to the model. For this purpose, a parameter is added to the scheme, which calculates the ratio of the reaction rate of RO_2 radicals with NO (high- NO_x regime) with respect to the sum of reaction rates of the reactions with HO_2 , and RO_2 (low- NO_x regime). For this purpose, a parameter (α) is added to the scheme, which calculates the ratio of the reaction rate of RO_2 radicals with NO (ν_{NO} ; high- NO_x regime) with respect to the sum of reaction rates of the reactions with HO_2 (ν_{HO_2}), and RO_2 (ν_{RO_2} ; low- NO_x regime). The parameter α is expressed as follows:

$$\alpha = \frac{\nu_{\text{NO}}}{\nu_{\text{NO}} + \nu_{\text{HO}_2} + \nu_{\text{RO}_2}} \quad \text{Eq. 1}$$

This α value represents the part of RO_2 radicals reacting with NO (which leads to apply "high NO_x yields"). It is calculated for each grid cell by using the instantaneous NO , HO_2 and RO_2 concentrations in the model. Then, the following equation is used to calculate an adjusted SOA yield using this α value (Carlton et al., 2009).

$$Y = \alpha \times Y_{\text{highNO}_x} + (1 - \alpha) \times Y_{\text{lowNO}_x} \quad \text{Eq. 2}$$

The fragmentation processes for the SVOC start after the third generation of oxidation, because fragmentation is favored with respect to functionalization for more oxidized compounds. Therefore, three series of species in different volatility bins were added to present each generation, similar to the approach set up in Shrivastava et al. (2013). ~~F~~However for biogenic VOC, fragmentation processes come into effect starting from the first generation, as in Shrivastava et al. (2013), because the intermediate species are considered as more oxidized. A fragmentation rate of 75% (with 25% left for functionalization) is used in this work for each oxidation step following Shrivastava et al. (2015). The formation of non-volatile SOA is performed by moving a part of each aerosol bin to non-volatile bins with a reaction constant ~~of~~ corresponding to a lifetime of one hour, similar to Shrivastava et al. (2015). Figure 2-c shows a scheme of the modified VBS for the VOC. In total, 40 species (with 10 size distribution bins each, i.e. 400 species in total) are added together to calculate the total concentration of SOA simulated by this scheme. The

resulting model has a total of 740 species in the output files (including gas-phase chemistry), which makes this scheme the most time consuming among the tested schemes. In section 5 of this paper, the results from the three schemes introduced above will be compared to observations.

3 Experimental dataset

5 During the SAFMED subproject, measurements were made at two major sites, the Ersa, Cap Corse station and the Cap Es Pinar, Mallorca station. The geographical characteristics and the measurements performed at each site are presented in the following section.

3.1 ChArMEx measurements

10 The Ersa supersite (42°58'04.1", 9°22'49.1") is located on the northern edge of the Corsica Island, in a rural environment, at an altitude of 530_m ~~above sea level (asl)~~. The station is located on a crest that dominates the northern part of the cape. It has a direct view of the sea on the three western, northern and eastern sides. Measurements carried out in this station are reported in table [12](#). More details about the instrumental setup in Ersa can be found in Michoud et al. (2017), Arndt et al. (2017) and Pey et al. (in prep).

15 The Es Pinar supersite (39°53'04.6", 3°11'40.9") is located on the northeastern part of Mallorca. The monitoring station was placed in the "Es Pinar" military facilities belonging to the Spanish Ministry of Defense. The environment is a non-urbanized area surrounded by pine forested slopes, being one of the most insulated zones in the Mallorca Isle, in between the Alcudia and Pollença bays, but still can be influenced by local anthropogenic emissions. The site is located in an altitude of 20 m.asl. The location of the station and a list of available measurements are presented in figure 3 and table [12](#) respectively.

20 ~~At Ersa~~~~For both sites~~, NO_x (nitrogen oxides) were measured by CraNO_x analyzer using ozone chemiluminescence with a resolution of 5 minutes. ~~The photolytic converter in the analyzer allows the conversion of direct measurements of NO₂ into NO in a selective way, thus avoiding interferences with other NO_y species. At Es Pinar, an API Teledyne T200 with molybdenum converter was used; therefore, the measurements are not specific to NO₂ and interferences of NO_y are possible for these measurements.~~~~A photolytic converter was incorporated in the~~
25 ~~analyzer to convert direct measurements of NO to NO₂.~~ Volatile Organic compounds were monitored at both supersites by a PTR-ToF-MS (KoreTM 2nd generation at Ersa, and IonikonTM PTR-ToF-MS 8000 at Es Pinar). A detailed procedure of VOCs quantification is provided in Michoud et al (2017) for Ersa and Pey et al. (in prep) for Es Pinar. Briefly, both instruments were calibrated daily using gas standard calibration bottles and blanks performed by means of a catalytic converter (stainless steel tubing filled with Pt wool held at 350°C).

30 A Quadrupole Aerosol Chemical Speciation Monitor (Q-ACSM, Aerodyne Research Inc, Ng et al., 2011) was used for the measurements of the chemical composition of non-refractory submicron aerosol at Ersa ~~with a time resolution of 30 minutes~~. This instrument has the same general structure of [an](#) AMS (Aerosol Mass Spectrometer), with the difference that it was developed specifically for long term monitoring. A High-Resolution Time-of-Flight AMS (HR-ToF-AMS, Aerodyne Research Inc, Decarlo et al., 2006) operated under standard conditions (i.e.
35 temperature of the vaporizer set at 600 °C, electronic ionization (EI) at 70 eV) was deployed with a temporal resolution of 8 minutes to determine the bulk chemical composition of the non-refractory fraction of the aerosol ~~and~~ for the Es Pinar site. AMS data were processed and analyzed using HR-ToF-AMS Analysis software

SQUIRREL (SeQUential Igor data RetRiEvaL) v.1.52L and PIKA (Peak Integration by Key Analysis) v.1.11L for IGOR Pro software package (Wavemetrics, Inc., Portland, OR, USA). Q-ACSM and AMS source apportionment results discussed in this work are detailed in Michoud et al. (2017) and Pey et al. (in prep) respectively. For both sites, source contributions were obtained from PMF analysis (Paatero and Tapper, 1994) of Q-ACSM and AMS OA mass spectra. PMF was solved using the multi-linear engine algorithm (ME-2, Paatero, 1997) ~~algorithm~~, using the Source Finder toolkit (SoFi, Canonaco et al., 2013). For the Ersa site, HOA (Hydrocarbon-like Organic Aerosol) profile was constrained with a reference HOA factor using an a value of 0.1. The a value refers to the extent to which the output HOA factor is allowed to vary from the input HOA reference mass spectra (i.e. 10% in this case, Canonaco et al., 2013). In such remote environment, the HOA factor could not be extracted from the OA mass spectral matrix with a classic unconstrained PMF approach. Two other factors were extracted, without any constraints, including SVOOA (Semi-Volatile Oxygenated Organic Aerosol) and LVOOA (Low-Volatile Oxygenated Organic Aerosol). For the Es Pinar site, HOA has been constrained using an a-value of 0.05. Three additional factors were retrieved, including an SVOOA (Semi-Volatile Oxygenated Organic Aerosol) and 2 LVOOA (Low Volatile Oxygenated Organic Aerosol) factors. In addition to HOA, 3 other factors have been extracted from the PMF analysis: SVOOA (Semi-Volatile Oxygenated Organic Aerosol) and 2 LVOOA (Low Volatile Oxygenated Organic Aerosol) factors. Differences between these 2 LVOOA factors are mainly linked to air masses origins and to the probable influence of marine emissions. ~~For the marine one of the~~ LVOOA factor at both sites, ~~with~~ correlation coefficient (R^2) of 0.43 and 0.47 with the main fragment derived from methane sulfonic acid (MSA, fragment CH_3SO_2^+) for Es Pinar and Ersa, respectively. For ~~the~~ sake of clarity and for ~~the~~ purpose of intercomparison ~~purpose~~ with the model outcomes, we merge the 2 LVOOA factors into one LVOOA for Cap Es Pinar site to be compared to the Ersa site results. Online aerosol chemical characterization was complemented by an Aethalometer (AE33, MAGEE, Drinovec et al., 2015) at Es Pinar and a Multiangle Absorption Photometer (MAAP5012, Thermo) for the quantification of BC at Ersa. PM_{10} total mass measurements were taken from TEOM-FDMS (Tapered Element Oscillating Microbalance-Filter Dynamic Measurement System) ~~for at~~ Es Pinar and ~~BETA~~ corrected by a factor obtained after comparison to gravimetric ~~measurements for~~ Es Pinar. Finally, daily PM_{10} aerosol samples were collected onto 150 mm quartz fiber filters (Tissuquartz, Pall) at both sites. Eighteen and eight samples were selected for Ersa and Es Pinar, respectively, for a subsequent analysis of radiocarbon performed on both OC and EC fraction following the method developed in Zhang et al. (2012).

3.2 Other measurements

For the validation of meteorological parameters, along with the meteorological surface measurements performed at ChArMEx stations, ~~other datasets were also used~~. ~~R~~adio-sounding data for three stations in the western Mediterranean basin were used for simulation-observation comparisons for meteorological parameters. The radio-soundings are performed by Météo-France at the two stations of Ajaccio, France (41°55'5", 8°47'38") and Nîmes, France (43°51'22", 4°24'22") and by AEMet at Palma, Spain (39°36'21", 2°42'24"). Each day two balloons at about 00 and 12 UTC are available for each station, a total of 96 balloons are included in the comparisons. Ajaccio and Palma are coastal stations, but Nîmes is farther from the coast compared to the other two stations. Each day two balloons were launched at about 00 and 12 UTC at each station, and a total of 96 balloons are included in the comparisons for an altitude between surface to 10 km.

4 Model validation

The CHIMERE model has been previously validated for different parts of the world (Hodzic & Jimenez, 2011 ; Solazzo et al., 2012 ; Borrego et al., 2013 ; Berezin et al., 2013 ; Petetin et al., 2014 ; Rea et al., 2015 ; ; Kononov et al., 2015 ; Mallet et al., 2016). The data set presented in section 3 is used for model validation. First, a representativeness error within simulations is calculated for a list of pollutants, which is necessary to distinguish between uncertainties due to limitations in model resolution and due to other reasons. Then a validation for the meteorological parameters is presented, before comparison of simulation results to gaseous and aerosol measurements.

4.1 Orographic representativeness of Cap Corse simulations

As explained before, during the ChArMEx campaign, an important number of observations were made at Ersa, Cap Corse. In order to use this data set for model evaluation, potential discrepancies due to a crude representation of the complex orography of Cap Corse need to be minimized and quantified since the measurements were performed on the crest line.

For the 10-km domain (D10), we noticed that there was an inconsistency between simulated and real altitude of the cell where the Ersa site is located; altitude being simulated at 360 m asl lower-below than the real altitude of measurements (530 m asl). Therefore, 1-km horizontally-resolved simulations were performed for the inner domain. However, even for the 1-km simulations the simulated altitude remains too low (365 m asl). This error occurs because the altitude of each cell in CHIMERE is calculated using the average of altitudes of points inside the cell, therefore if the altitude of the ground surface inside a cell happens to vary greatly, the average would be lower than the higher points seen in the cell (which corresponds in our case to the Ersa site located on the crest). In addition, the average of the marine boundary layer height is typically around 500 m (Stull, 1988), therefore a discrepancy in the simulated altitude could cause significant errors in the simulations. These two reasons ~~facts~~, make it important to explore the representativeness of the simulations regarding this station.

This led us to perform an orographic representativeness test on the 1-km domain (D1) at the Ersa site. A matrix of neighboring cells around the grid cell covering the Ersa station (up to 5km distance) was taken (figure 4-a), and species concentrations were plotted against the variation of the altitude of these different cells. The highest altitude reached by one of the cells is about 450 m. Then, the concentration on the exact altitude (530 m) was extrapolated using a non-linear regression between the altitude and the concentration of the selected cells with several different equations for each time step. In total, 9 non-linear equations were tested, among ~~from~~ which only five were finally used for the calculation of the representativeness error. For the other four equations, due to convergence problems occurred and for the other four equations no stable solution could be found for some of the hourly time-steps (see supplementary material, SI3 for the details). Regressions were performed separately for each of the 720 hourly time-steps of the one month simulations. An example of this regression for organic aerosols for one time step and one of the equations is shown in figure 4-b (equation n.1 from SI3). The results were filtered using two criteria (convergence of regression for each time-step and a correlation coefficient between fitted and simulated points of higher than a fixed value ~~threshold~~) depending on statistical values of the regressions (see supplementary material, SI3, for details) and only regressions conforming to these criteria were retained. If at least two converging regressions were not retained for a given time step, the results for that time step were not further used. Figure 4-c shows the compiled results for all equations and all simulation times in one time series for total OA concentration.

Note that model output was generated with an hourly time step. Using these results, for a list of different species, an orographic representativeness error (ORE) was calculated using the average of the difference of the upper and lower confidence intervals for all equations. As an example, carbon monoxide, which is a well-mixed and a more stable component in the atmosphere, presents the lowest error among the tested species (2%). Ozone also presents one of the lowest errors (4%) and nitrogen oxides one of the highest (75%). Organic aerosol, of particular interest for this study, shows a moderate error of 10%. In terms of meteorological parameters, relative humidity appears more affected (relative ORE of 18%) than temperature (the relative ORE is calculated on values of T in °C). A summary of results of this test is shown in table 23.

~~A general conclusion is that secondary pollutants with higher atmospheric lifetimes appear to be well represented from a geographic point of view. On the contrary, model-observation comparisons for more reactive primary and secondary pollutants with short lifetimes (primary such as NO_x and reactive secondary such as MVK+MACR) should be performed with caution keeping in mind the fact that the simulated altitude is not representative of the orography for this specific station. A general conclusion is that secondary pollutants appear to be well represented from a geographic point of view, however model-observation comparisons for specially more variable primary pollutants should be performed with caution and keeping in mind the fact that the simulated altitude is not representative of the orography for this specific station.~~ This is due to the fact that ~~for~~ short-lived primary species, ~~have not yet had the chance to vertically mix, if~~ emission sources happen to be nearby (which is the case here). ~~Vertical layering of these concentrations results then in significant sensitivity to the simulated altitude of a site. ; then the air masses in the area have not had yet the chance to properly mix.~~ On the contrary, secondary species, which ~~partly~~ have partly been transported from the continental boundary layer, are believed to be better mixed vertically. ~~Therefore, for secondary~~ For those species, differences in the simulated versus observed altitude lead to relatively smaller errors.

The question remains on which domain should be used for model-measurement comparisons remains. As seen above, D10, despite having a sufficiently fine resolution for most continental areas, is not capable of representing the complex orography of the Cap Corse, therefore the D1 simulation results have been used for comparisons, except for meteorological parameters, where all possible domains (and resolutions) are compared. The Es Pinar station does not have the same intense altitudinal gradient ~~that is~~ seen at Ersa, therefore the aforementioned test was not performed for this station and the D10 simulations are used for comparisons for Es Pinar.

In the following section, ~~for each model-observation comparison at Ersa, the~~ confidence intervals representing ~~which stands for~~ the Orographic Representativeness Error (ORE) ~~value that was~~ derived in this section are considered for the model-observation comparisons ~~is added~~.

4.2 Meteorology evaluation

Meteorological output of the mesoscale WRF model at different resolutions has been used as input to the CHIMERE CTM. The meteorological data used by CHIMERE ~~were~~ compared to various meteorological observations such as radio-soundings and surface observations at the measurement sites. Detailed results of these comparisons are given in the supplementary material (SI4). Here, a short overview of the results and the implications for the model ability to simulate transport to the measurement sites is given.

Comparisons for temperature, a basic variable to control the quality of meteorological simulations show a good correlation for radio-sounding comparisons (typically from 0.60 to 0.85 for hourly values) and a low bias (typically

from -1.16 to -0.39°C) for the three sites of Palma, Nîmes and Ajaccio. Wind speed shows a good correlation at higher altitudes, and also near the surface for Nîmes and Palma stations for radio-sounding stations, while for the Ajaccio station the sea/land breezes are probably not well represented in the model. For Es Pinar, the coastal feature of the site is difficult to take into account in a 10 km horizontally resolved simulation and leads to larger errors. Ground based meteorological measurements were also compared at both sites (SI4). ~~whereas at~~ At Ers, the correlation in finer domains is better than that of D10 for wind speed (typically around 0.66 versus around 0.60). For the E-OBS network (surface datasets provided by European Climate Assessment & Dataset (ECAD) project for monitoring and analyzing climate extremes, Haylock et al., 2008; Hofstra et al., 2009) comparisons (SI4) also show a good correlation and a low bias for temperature (correlation of 0.79 with a bias of -0.54°C for mean temperature observed for 71 stations in D10 domain), while the daily minima seems to be underestimated (bias of -3°C ~~for daily minima~~ observed for 71 stations).

While the general comparison between the hourly meteorological fields used as input for CHIMERE simulations and observations is in general already satisfying, ~~when daily averages representative for different meteorological conditions are compared instead of hourly values,~~ the correlation becomes higher and the bias lower when daily averages representative for different meteorological conditions are compared instead of hourly values. ~~Differences in wind speed and wind direction may lead to errors in representing the short term variability of gaseous and aerosol species.~~

4.3 Gaseous species

Among all the gaseous species available in the observations, four were chosen in this study: nitrogen oxides (NO_x), isoprene (C₅H₈), monoterpenes and the sum of methacrolein and methyl vinyl ketone (here after called MACR+MVK). These four species were chosen since isoprene and monoterpenes are the principal precursors for biogenic SOA, MACR+MVK are formed during isoprene oxidation and NO_x is a good tracer for local pollution. The comparisons for the Ers and Es Pinar stations are shown in figure 5, and statistics of the comparison are shown in table 34. For Ers, the orographic representativeness errors derived in section 4.1 are also shown. In all comparisons, the results for the simulations with the modified VBS scheme are used, but the choice of the organic aerosol scheme only slightly affects the simulation of gaseous species (mainly via heterogeneous reactions on aerosol surfaces included within CHIMERE).

~~For nitrogen oxides (figure 5-a1),~~ Results show that there is a good correspondence between the averages of simulated and observed nitrogen oxides at Ers (figure 5-a1). The low correlation for nitrogen oxides at Ers might be partly explained by the high representativeness error (75%) for this component. This is because the altitude in the simulations is lower and therefore the emission sources are closer in the model than they are in reality. At Es Pinar (figure 5-a2), since the measurements are not specific to NO₂, the NO_y time series are added to the figure as well. As a consequence, if the model had no error, the NO_x observations would be expected to lie between NO_x and NO_y simulations. This is the case because NO_x observations are on the average 40% higher than the NO_x simulations and 9% lower than the NO_y simulations. both observed and simulated NO_x concentrations becomes higher than at Ers, indicating the more locally polluted nature of this site, but the model underestimates observed NO_x by about 40%.

For isoprene (figure 5-b1 and 5-b2), a good correlation (0.76, 0.71) between simulations and observations appears at both sites. However there is an important overestimation (by a factor of 2.5) in the simulations for the Ers site,

which could also be linked to the high orographic representativeness error (85%), and also to the fact that local emissions sources ~~may~~ not ~~be~~ correctly taken into account in the MEGAN emission model. On the contrary, at Es Pinar isoprene is underestimated by about 25%. The sum of MACR+MVK (figure 5-c1 and 5-c2) is overestimated by about a factor of two at Ersa, following the pattern of overestimation of isoprene at this site, while the bias is small at Es Pinar. Monoterpenes (figure 5-d1 and 5-d2) show an underestimation by about 70% at both sites, observations being about 5 times larger at Ersa than at Es Pinar. Again, this could be related to the orographic representativeness error at Ersa and to ~~non~~-accounted-for local vegetation at both sites.

Daily correlations of 0.35, 0.87, 0.85 and 0.58 (instead of 0.37, 0.76, 0.62 and 0.35 hourly values) are seen at Ersa for nitrogen oxides, isoprene, MACR+MVK and monoterpenes, respectively. These values change to 0.16, 0.51, 0.72 and 0.10 for daily comparisons (instead of 0.12, 0.69, 0.41 and 0.14 when correlating hourly values) at Es Pinar. ~~While showing the same improvement in correlation as for meteorological parameters for Ersa, meaning that chemical regime changes are better simulated than short term variations, the results do not change as much for Es Pinar. Improvements in correlation for daily averages could be related to those in meteorological parameters, at least for Ersa. They show the difficulty to correctly simulate short term (hourly) variations both in meteorological parameters and chemical species.~~

~~A drawback of the comparisons for isoprene, and monoterpenes, contributing respectively to 40 and 60% of SOA simulated with the modified VBS scheme in the western Mediterranean region, is the measurements representativeness restricted to local scales. A more regional evaluation of these precursor species would have been necessary, since SOA simulated and observed at Ersa and El Pinar was partly formed far away from these sites.~~

~~Isoprene and monoterpenes are the most important precursors of biogenic organic aerosols. Sensitivity tests using the modified VBS scheme showed that there is a 40 to 60% repartition for the SOA formed from isoprene and monoterpenes, respectively, in the western Mediterranean region. A drawback of the comparisons for these two species is the fact that the measurements present a very local point of view, while a more regional look would have been more pertinent for these species, since a significant amount of SOA is transported to both sites mainly from continental areas.~~

As mentioned before (section 2), isoprene and terpene emissions in our work are generated by the MEGAN model (Guenther et al., 2006). Zare et al. (2012) evaluated MEGAN model derived isoprene emissions coupled to the hemispheric DEHM chemistry-transport model against measurements. On average over 2006, they found a simulated isoprene overestimation ~~at~~for 4 European sites (between a factor 2 and 10), good agreement (within +/- 30%) ~~at~~for 2 sites and an underestimation ~~at~~for 2 sites (also between a factor 2 and 10). However, none of the sites was located close to the Mediterranean Sea. Curci et al. (2010) performed an inverse modelling study to correct European summer (May to September) 2005 MEGAN isoprene emissions from formaldehyde vertical column OMI measurements (given that isoprene is a major formaldehyde precursor). For western Mediterranean countries area, they found an isoprene emissions underestimation using MEGAN of 40 % over Spain, a tendency for an underestimation, but with regional differences over Italy, and only small differences over France. This comparison globally lends confidence to MEGAN derived isoprene emissions. Unfortunately, to our best knowledge, no comparable studies exist in order to validate monoterpene emissions. For the Rome area, Italy, Fares et al. (2013) concluded for a variety of typical Mediterranean tree and vegetation species mix, that MEGAN correctly simulated ~~(within 10% error)~~ mean observed monoterpene fluxes over the last two weeks of August 2007 ~~within 10% error~~

for a variety of Mediterranean tree and vegetation species mix, as long as a canopy model was included (as in our study).

4.4 Particulate species

Figure 6 shows the comparison between observations and simulations for particulate sulfate and black carbon in PM₁. These two species are chosen as two important fine aerosol components, before the comparison of organic aerosol in chapter 5. The left panel shows the comparison for Ersa and the right one for Es Pinar. Statistical information for these species is given in table 45. There is an overestimation for sulfate particles (figures 6-b1 and 6-b2) by about 45%, well beyond the representativeness error for this species (15%). Besides, the short and sharp decreases in the measurements of sulfates correspond to low clouds passing at the level of the station which are not simulated by the model. Cloud scavenging processes are already taken into account in the model. However, because of the unique geographical characteristics mentioned before for this site, the meteorological inputs did not simulate these fog events and therefore cloud scavenging was not activated in the simulation. Since these decreases concern only a small percentage of the observations, they do not have a major effect on the outcome of these comparisons. While this effect is very visible for sulfates, it is less pronounced for other particulate species such as black carbon and OA. A large sulfate peak simulated in the morning of 29 July is not present in the observations; it originates in the model from an air mass arriving from Marseille which is both a busy harbor and an important industrial area with large SO₂ emissions. This ~~very specific transport event is not observed, which~~ is probably due to small errors in the wind fields. In addition, there are two periods of overestimation of this species in the simulations; the period of 18 to 20 July and the period of 29 July to the night of first of August. During the second period, the ACSM PM₁ observations show concentrations of close to zero, which are consistent with PM₁₀ PILS-IC sulfate measurements. For this period elevated southerly winds are observed in the Corsica area, and the absence of strong SO₂ sources in this sector might explain the lower concentrations that are seen not only for sulfate but also for black carbon. The constant overestimation of sulfates during this period may suggest an overestimation of boundary conditions for these species.

A moderate correlation (0.326) between the model and the observations is seen for BC (figures 6-a1 and 6-a2), but the representativeness error is important for this species (26%), since one of its primary origins is emission of ships passing nearby the coasts of the Cap Corse. The local emissions due to shipping activities at Ersa and anthropogenic activities at Es Pinar are visible in the observations as frequent narrow (in time) peaks. These activities are also visible in the simulations at Ersa. However, at Es Pinar the model doesn't succeed in simulating them, as already observed to an even larger extent for NO_x. The slight overestimation of BC (5%) in the model for the Ersa site could be explained by the orographic representativeness error. The same cloud effect seen for the sulfate particles at Ersa is visible to a lesser extent for BC for the Ersa site as well. Correlation between daily observations and the simulations was also calculated, ~~R² is which shows~~ 0.51 and 0.56 for the Ersa station (instead of 0.36 and 0.42 for hourly values) and 0.81 and 0.65 for the Es Pinar station (instead of 0.47 and 0.52 for hourly values) for sulfate and BC respectively. Similar to meteorological parameters, the model can reproduce the daily concentration changes for these two species better than the hourly changes.

~~As a conclusion of this chapter, the comparison between the meteorological fields used as input for CHIMERE simulations and observations at surface sites and from radio soundings is satisfying, and shows low bias. However, differences in wind speed and wind direction may suggest that the variability in advection to the measurement~~

sites, in particular at short timescales (less than a day), is not always properly simulated. This may lead to errors in representing the short term variability of gaseous species. Among gaseous species, the biases found for the short lived compounds could mainly be explained by strong orographic representativeness error, which were evaluated in the range 60-85% in this section. The local character of the comparisons for biogenic VOCs was pointed out, as well as the fact that biogenic OA is formed at a regional scale from these precursors rather than a local scale. Available regional comparisons show acceptable comparisons for BVOC observed and simulated concentrations (Curci et al., 2010 ; Zare et al., 2012 ; Fares et al., 2013). Therefore, a systematic misrepresentation of BVOC in the model seem to be unlikely. For aerosol species, low bias was seen for BC, while an overestimation in simulations is observed for sulfates.

5 Organic aerosol simulation

A description of each of the 4 schemes for the simulation of organic aerosols tested within the CHIMERE model for the Mediterranean area has been given in section 2. These four schemes are the CHIMERE standard scheme, the VBS scheme with BSOA aging (noted as Standard VBS_ba), the VBS scheme without BSOA aging (noted as Standard VBS_nba) and the modified VBS (noted as modified VBS) scheme which includes fragmentation and formation of non-volatile organic aerosol. For each scheme, four domains were used: one coarse domain, and three others nested inside the coarse one with increasing resolutions. As before, the simulations from the finest domain are used for the comparisons. The domain with the finest resolution (1km) was used for the representativeness tests. For each scheme, meteorological and boundary conditions are the same, hence only the simulation of organic aerosols and subsequently the aggregation of anthropogenic PM emissions differ between simulations.

5.1 Comparison of PM₁ total organic aerosols concentration

Figure 7 shows the time series of comparison of organic aerosols for these four schemes with the measurements at Ersa and Es Pinar. The circles beside each plot show the average concentration for different time series, in addition table 56 shows the statistical parameters corresponding to these time series. The observed OA concentration at the Ersa site measured by the ACSM for the PM₁ fraction has an average of 3.71 $\mu\text{g}\cdot\text{m}^{-3}$ and that of the Es Pinar site measured by the AMS for the PM₁ fraction a somewhat lower average of 2.88 $\mu\text{g}\cdot\text{m}^{-3}$. The difference between the two sites may be attributed to the fact that Cap Corse is closer to both local and transported (continental) biogenic sources than Es Pinar. The VBS scheme with BSOA aging greatly overestimates the organic aerosol with a (bias of more than a factor of three). As mentioned before, the aging of biogenic aerosols in the VBS scheme usually results in an overestimation in organic aerosols (Lane et al., 2008). With the BSOA aging option turned off, the standard VBS scheme comes much closer to the average of total OA concentration measured at both sites (relative biases of below 50%). The CHIMERE standard scheme also overestimates the mass concentration of organic aerosols, but in a lesser extent compared to the VBS scheme with BSOA aging (bias of a bit less than a factor of two). The modified VBS scheme corresponds much better to observations (negative biases about 20%). The model results agree for both stations in this regard. At Ersa, biases of +244%, +91%, +34% and -18% respectively are observed for the VBS standard scheme with BSOA aging, the CHIMERE scheme, the standard VBS scheme without BSOA aging and the VBS modified scheme respectively. The corresponding numbers are of +218%, +98%, +43% and -23% for Es Pinar.

The daily correlation for the modified VBS scheme is 0.63 and 0.51 for Ersa and Es Pinar respectively (instead of 0.50 and 0.29 for hourly values). This shows, again, that the model better represents long-scale day-to-day changes better than hourly variations. While the concentrations of both the modified VBS scheme and the standard VBS scheme without BSOA aging correspond well with the experimental data, other aspects such as origins of the formed organic aerosol and its oxidation state have to be considered before reaching any conclusion about the robustness of these schemes. ~~It is also noticeable that the large-scale tendencies are in most cases simulated in each scheme, therefore different regimes are well predicted by the model. However, smaller changes in the observations (on shorter time scales, a day or less) are not reproduced in any of the schemes. This might be because of wind direction simulation, which, as explained before is difficult to represent in coastal areas.~~

The overestimation of secondary organic aerosol with the CHIMERE standard scheme is a new feature, not apparent in previous studies (Bessagnet et al., 2008 ; Hodzic and Jimenez, 2011 ; Petetin et al., 2014). A previous comparison of CHIMERE with organic aerosol simulations using the same scheme and coupled to MEGAN biogenic emissions, resulted in a good comparison or underestimation with OC observations over Europe from the Carbosol project (Gelencsér et al., 2007) for summer 2003, when biogenic SOA was dominant. However, this comparison included only one site close to the western Mediterranean basin (Montelibretti, Italy). The overestimation of BSOA in the VBS scheme when the BSOA aging is activated was documented for the USA in several occasions (Robinson et al., 2007b ; Lane et al., 2008). Ultimately, it cannot be excluded that the OA overestimation with both schemes at Ersa and Es Pinar is also due to BSOA-BCOV overestimation. However, the available material does not support this hypothesis: (i) OA overestimation is observed at two independent, distant sites, (ii) local monoterpene underestimation at both these sites, (iii) no evidence for MEGAN monoterpene overestimation is available in literature and MEGAN isoprene overestimation over the western Mediterranean area was ruled out by satellite observations (Curci et al., 2010).

5.2 Total carbonaceous particles origins based on ¹⁴C measurements

The results of ¹⁴C measurements in carbonaceous aerosol filter samples for the PM₁ fraction at the Ersa and Es Pinar sites were used in order to better discriminate between the modern and mostly biogenic versus fossil and anthropogenic origin of organic aerosols, and to compare it to simulations with different organic aerosol schemes tested. It must be noted that in the simulations, species are not separated automatically into fossil/ non-fossil parts, therefore these fractions need to be calculated as a post-treatment of simulations, affecting each relevant particulate species to both fractions. ASOA is considered to be in the fossil fraction and BSOA in the non-fossil fraction. For carbonaceous aerosol, residential/domestic uses are considered as non-fossil as they are mostly related to wood burning (Sasser et al., 2012). Therefore, they are attributed to the non-fossil bin (3.6% for BC and 12.3% for OC, Sasser et al., 2012). The non-fossil contribution of ASOA and POA due to biofuel usage is ignored here, as it is minor (<5%). No major biomass burning events were seen in the period of this study, but minor contributions of this source cannot be excluded. It should also be mentioned that the ¹⁴C measurements show the mass of carbonaceous particles in each filter, therefore have a unit of μgC.m⁻³, while the simulations show the total organic aerosols in μg.m³. In the comparisons that follow, it is pertinent to use an OM/OC conversion factor to be able to compare the ¹⁴C measurements to simulations. For this purpose, an average OM/OC factor of 2 is used for the secondary aerosol fraction (both for the LV-OOA and SV-OOA factors), while a factor of 1.3 is used for HOA according to Aiken et al.(2008). However, the choice of the OM/OC factor has a small effect on the outcome of

general comparisons since HOA values are marginal compared to other factors and we are only interested in relative contribution of fossil/non-fossil factors ~~the comparisons are all shown in percentage of fossil/non fossil distribution.~~

Figure 8-a for Ersa and 8-b for Es Pinar show the average of all filters for each scheme compared to the observations, while figure 9-a and 9-b show the relative distribution of fossil/non fossil sources at both sites. Among these total averages, the distribution at Ersa is 81% \pm 1.5% non-fossil and 19% \pm 1.5% fossil, and 67% \pm 3% versus 33% \pm 3% at Es Pinar. Apparently, biogenic contributions to OA are dominant at both sites, but larger for the Ersa site.

While the comparison of averages for all schemes with Ersa measurements shows that the modified VBS scheme is the closest in fossil/non-fossil partitioning (19%/81% \pm 1%), the CHIMERE standard scheme is also performing well when looking at the percentage of distribution for each source (20%/80% \pm 1%). The VBS scheme with BSOA aging shows an underestimation in the percentage of fossil carbons (16%/84% \pm 0.7%), which can be due to the overestimation of biogenic aging of secondary organic aerosols in this scheme; on the contrary, the VBS scheme without BSOA aging shows an important overestimation of the fossil percentage (34%/66% \pm 2%).

A distribution of 33%/67% \pm 3% of the fossil-non-fossil fraction is observed at Es Pinar. As for the Ersa site, the modified VBS scheme is the closest to the observations (32%/68% \pm 2.5%). The CHIMERE standard scheme shows an underestimation of the fossil contribution with a distribution of 28%/72% \pm 3%. The standard VBS scheme with BSOA aging underestimates the fossil contribution with a distribution of 21%/79% \pm 1.5%, while, the standard VBS scheme without BSOA aging largely overestimates the fossil contribution (42%/58% \pm 2%). These results show that the two sites differ greatly when taking into account nearby sources and geographical conditions; the Ersa site is an elevated rural station at the interface between the marine boundary layer and the residual boundary layer, and the Es Pinar site is a sea-side station closer to anthropogenic sources. These differences should normally be represented in the percentage of fossil carbon concentrations both in observations and in simulations. Although this difference is noticeable in observations, and also in the modified VBS scheme, it is less emphasized in the CHIMERE standard scheme and in the VBS standard scheme with either parameterization of aging.

A more detailed look to individual filters for the modified VBS scheme for the Ersa and Es Pinar sites is presented in figure 9-a and figure 9-b respectively. The tendencies from one day to another are in most cases not reproduced correctly for both sites. Therefore while the average mass repartition of modern and fossil sources is well-represented by this scheme, the day-to-day variability is not fully consistent with the measurements. This inconsistency is also seen in the other tested schemes.

5.3 Volatility and oxidation state comparison with PMF results

The PMF (Positive Matrix Factorization) results of the ACSM/AMS (Aerosol Mass Spectrometer) measurements give us the chance to learn more about the oxidation state of the organic aerosols (Michoud et al., 2017 for Ersa and Pey et al. in prep for Es Pinar). The PMF analysis allows to divide PM₁ organic aerosol measurements into different groups with distinctive mass spectra corresponding to distinctive oxidation state (Lanz et al., 2010). The most common retrieved factors of such an analysis are HOA (Hydrogen-like Organic Aerosol), SV-OOA (Semi-Volatility Oxidized Organic Aerosol) and LV-OOA (Low-Volatility Oxidized Organic Aerosol). However, it does not give direct information about the volatility ~~range-distribution~~ of each group. This makes the PMF output

difficult to compare with our model results, which give the volatility distribution of organic aerosol, but not its oxidative state. Here, we first compare volatility distributions obtained with the four aerosol schemes, and then try to attribute the simulated aerosol to the three factors HOA, SV-OOA and LV-OOA, in order to compare it to the observed distributions. The three schemes based on the VBS scheme already distribute aerosols in volatility bins (Robinson et al., 2007a). The distribution to LV-OOA and SV-OOA for these three schemes was done mainly by taking into account the saturation concentration, with the threshold chosen as a saturation concentration $C^* \geq 1 \mu\text{g}\cdot\text{m}^{-3}$ for SV-OOA and $C^* \leq 0.1 \mu\text{g}\cdot\text{m}^{-3}$ for LV-OOA (Donahue et al., 2012). Primary organic aerosols were considered to be in the HOA factor regardless of their saturation concentration. For the CHIMERE standard scheme, each surrogate species is associated with a saturation vapor pressure, which was used to calculate the saturation concentration for each component at ambient temperature.

The observations show that at Ersa, the LV-OOA factor dominates the PMF results (88%), with 10% SV-OOA and a minor (only 2%) contribution of HOA. For the Es Pinar site the contribution of LV-OOA drops to (75%) and the contribution of ~~HOA and SV-OOA~~ SV-OOA and HOA becomes somewhat larger (214% and 424% respectively). As mentioned before, the Es Pinar site is more influenced by local anthropogenic sources. Therefore, more local OA emissions are expected which corresponds to the increase in the HOA percentage seen in the observations. These emissions are oxidized locally to form SVOCs that fall in the SV-OOA group explaining the rise in the percentage of this group.

Figure 10-a and figure 10-b show the relative distribution of all organic aerosols in seven volatility bins for all tested schemes for Ersa and Es Pinar sites, respectively, and for all tested schemes. For each scheme, the average for the total period of the simulations was used to calculate the percentage in each bin. The bins shown are in the range of $10^{-3} - 10^3 \mu\text{g}\cdot\text{m}^{-3}$, all the aerosols with a volatility higher or lower than the extremes are put in the last high or low bin respectively.

These figures show that the percentage of aerosols in each volatility bin for the two different sites is relatively similar. The CHIMERE standard scheme distributes most of the aerosol in the three bins in 1-100 $\mu\text{g}\cdot\text{m}^{-3}$ volatility ranges which falls into the SV-OOA group obtained by PMF. This is due to the initial distribution of volatilities of surrogate SVOC species used that best fit chamber measurements, and the fact that there is no aging mechanism in this scheme to make the aerosols less volatile. The 100 $\mu\text{g}\cdot\text{m}^{-3}$ volatility bin corresponds to SVOCs produced from isoprene and monoterpene oxidation and presents the highest percentage for this scheme. The standard scheme also produces 16% of organic aerosols in the LV-OOA range (volatilities corresponding to a C^* between 0.001 and 0.1 $\mu\text{g}\cdot\text{m}^{-3}$). The standard VBS scheme with BSOA aging has only small fraction of aerosol in the LV-OOA range (from SVOC emission aging) since, by construction of the scheme, the most aged BSOA and ASOA aerosols fall in the 1 $\mu\text{g}\cdot\text{m}^{-3}$ volatility bin, which is actually the bin with the highest percentage of OA. The standard VBS scheme without BSOA aging presents relatively similar results to the scheme with biogenic aging, with as expected larger percentages for higher volatilities in the absence of aging. It also show a larger LV-OOA fraction, probably because the lower total OA concentration favors the contribution of lower volatility SVOC's to the aerosol phase. On the whole, the two schemes yield much too low LV-OOA fractions as compared to observations. The modified VBS scheme has a more realistic distribution into the three oxidation groups. In this scheme, the highest percentage of OA falls in the $10^{-3} \mu\text{g}\cdot\text{m}^{-3}$ volatility bin which is in the LV-OOA range. The rest of the aerosols are distributed almost equally in volatility bins between 10^{-2} and $10^2 \mu\text{g}\cdot\text{m}^{-3}$ with a slight decrease in percentage in the higher volatility bins. The percentage in higher volatility bins at Ersa are slightly lower than the

ones at Es Pinar which could be explained by the stronger local sources at the Es Pinar site with stronger primary SVOC emissions.

Figure 11-a and figure 11-b show the calculated HOA, LV-OOA and SV-OOA groups in simulations compared to the ones in observations at Ersa and at Es Pinar respectively. ~~At For the Ersa, site~~ the results of the modified VBS scheme are consistent with the observations with a slight underestimation of HOAs which is more visible at Es Pinar. The standard CHIMERE scheme leads to higher, overestimated HOA values, as primary OA emissions are considered non-volatile. Only the modified VBS scheme succeeds ~~to reproduce~~ in reproducing the major contribution of LV-OOAs at both sites. However, while staying close to observations at Ersa, it seems to overestimate the formation of LV-OOAs at Es Pinar. The standard VBS schemes with or without BSOA aging, greatly underestimate the formation of LV-OOAs, and as a counterpart overestimate the formation of SV-OOAs. Therefore, among the tested schemes, only the modified VBS scheme allows representing the distribution of the different PMF factors in a satisfying way.

~~A general conclusion for this section is that among the tested schemes, the modified VBS scheme seems to be closer to observations in all three criteria chosen for this study (mass, modern vs. fossil origin, volatility/oxidation state). It succeeds to reproduce the mass average of the aerosols observed at both sites, represents the fossil/non-fossil distribution of OA both when looking at the percentages and simulated concentrations, and it comes closest to observations when comparing the oxidation state.~~

6 Budget of organic aerosols

In section 5, we have highlighted the best performance of the modified VBS scheme for organic aerosol simulation amongst the tested schemes by comparison ~~with to~~ observational data at two different sites in the western Mediterranean basin. In the present section, we use this scheme in order to simulate the organic aerosol distribution and its anthropogenic and biogenic origins over the western Mediterranean basin during the SAFMED campaign. Figure 12 shows a series of figures where the left column always corresponds to simulations near the surface, and the right column shows the same concentration at an altitude of between 300 to 450m (for marine grid cells, called for simplicity boundary layer BL). Each row shows a different component, first row corresponding to OA concentrations, second row to biogenic OA concentrations, third row to anthropogenic OA concentrations and last row presents the sum of POA and all its subsequent oxidation products. The ~~bigger larger~~ part of each figure corresponds to D10 simulations, while the part inside the black rectangle shows the D3 simulations, both showing the average of the organic aerosols on the whole simulation period (1 month from July 10th to August 9th). Figure 13 shows the same type of figures for the percentage of contribution of biogenic OAs, anthropogenic OAs and the sum of POAs (that is POA and POA oxidation products). The small differences at the interface of the two domains is because CHIMERE model is a one-way chemistry-transport model: the simulations for the parent domain influence the simulations for the nested domain, however, the inverse is not applied; therefore any concentrations observed in the nested domain will not change the concentrations seen in the parent domain.

Examining figures corresponding to OA concentrations (Figures 12-a and 12-b), at the surface level there is a region of high concentration of OA in the Gulf of Genoa (between Genoa and Corsica) reaching nearly 4 $\mu\text{g}\cdot\text{m}^{-3}$. The high concentration in this area is less pronounced in the BL (Figure 12-b). The concentration of biogenic OAs (Figures 12-c and 12-d) is high over the basin as well as over Europe, and less important over North Africa. The percentage of contribution of this type of aerosol to the overall OA concentration is shown in figure 13-a and 13-

b. As expected from looking at the total concentration of biogenic OAs, their contribution stays on average around 70% over the basin, but is lower in the Gulf of Genoa (about 60% at surface, nearly 70% at altitude). In this area the secondary anthropogenic organic aerosols show a bit higher contributions compared to their contribution to the rest of the domain (around 14% instead of 12%), also, the contribution of primary organic aerosols (and its subsequent oxidations) is also quite high in this region (around 20%). The areas between the Corsica and Marseilles and also the northern coasts of Africa are main shipping routes in the western Mediterranean basin, with high amounts of shipping related emissions. They affect primary organic aerosol POA and its semi-volatile oxidation products as shown in figure figures 12-g and 12-h, with concentrations as high as 4.50.9 $\mu\text{g}\cdot\text{m}^{-3}$ around Corsica and near the African coasts. This corresponds to a contribution of around 2230% over Corsica and 252% over the coasts of Africa (Figures 13-e and 13-f). These values may actually be upper limits sinceas shipping primary SVOC and IVOC emissions were treated as those from other activity sectors (see section 2.2.3), and as recent chamber study data suggest lower SOA yields from shipping emissions (Pieber et al., 2016) than used here. The influence of shipping emissions is not visible in the BL, because vertical mixing within the marine boundary layer is weak, thus in the 300 to 400m layer the contribution of biogenic OA becomes dominant for the whole basin. Still, in the Gulf of Genoa, some effects of anthropogenic influences are visible in the BL, which could be linked to emissions originating from industrial sites and the harbor of Genoa area, which are apparently better mixed vertically over the continental convective boundary layer (in the coastal region with strong orography). These anthropogenic emissions are visible in figures corresponding to anthropogenic OA, formed from anthropogenic VOC's and especially aromatic compounds (Figures 12-e and 12-f for absolute concentrations and figures 13-c and 13-d for relative percentages). While the average concentration of anthropogenic OAs stay relatively low over the western Mediterranean basin (an average of $0.30 \mu\text{g}\cdot\text{m}^{-3}$ near the surface), they become more pronounced around the Gulf of Genoa and the eastern part of the domain both near the surface and in the BL. A contribution of about 12-15% both near the surface and in the BL is seen for this component above the Gulf of Genoa and over the highly industrialized and densely populated Po Valley.

25 **7 Conclusion/discussion**

Three schemes for the simulation of organic aerosol were implemented and tested along with the standard scheme in the CHIMERE chemistry-transport model. The simulations from each of the four schemes were compared to detailed experimental data obtained from two different stations in the western Mediterranean area during the ChArMEx campaign in summer 2013. The simulations were performed over one month in the summer of 2013 on 4 nested domains with increasing resolutions, the largest one covering Europe and northern Africa with a 30-km horizontal resolution, to the smallest one focused on the Cap Corse area with a 1km horizontal resolution.

For the comparisons of OA simulated with different schemes to observations, we explored three different aspects: mass concentration, distribution with respect to volatility and oxidative state of OA classes derived by PMF, and ^{14}C measurements discriminating fossil or non-fossil origin. Results show that the modified VBS scheme (i.e. including the fragmentation and formation of non-volatile organic aerosols), better corresponds to the observations at both sites, and this for all three aspects. The modified VBS scheme succeeds at simulating the average concentration of OA for the one-month campaign period with low bias (about -20% at both sites), even if the hourly variability is not perfectly displayed (as for other aerosol components). Comparisons for OA precursors (isoprene and terpenes) and isoprene oxidation products (sum of methyl vinyl ketone and methacroleine) were

performed and showed significant differences, which do however not necessarily affect the model ability to form BSOA, because the BVOC measurements are representative for a local scale, while BSOA formation occurs on a larger regional scale. Chrit et al. (2017) used a two-step surrogate scheme for the simulation of the Ersa site measurements. They found that their modified SOA simulation scheme corresponds well with the data, with a correlation of 0.67 and a Mean Fractional Bias (MFB) of -0.15 for hourly values of the period between June to August 2013. For the period of July to August 2013 for hourly values, we find 0.52 correlation and an MFB of -0.03 for the modified VBS scheme, which shows that both these schemes work reasonably well for the simulated area.

Furthermore, the fossil/non-fossil distribution of OA was explored in our different schemes. The modified VBS scheme corresponds better to available data in this regard as well. It is also the only scheme among the four tested that represents the distribution of the different PMF factors in a satisfying way attributing the major OA fraction to LV-OOA (but slightly overestimating this part). The differences between the sites, especially the more local anthropogenic character of the Es Pinar site (larger ¹⁴C fossil fuel origin, larger SV-OOA fraction, higher NO_x concentration which is a good tracer for anthropogenic pollution) and the lower OA concentration at this site are qualitatively simulated well with this scheme. While the standard VBS scheme without BSOA aging is as close to mass and origin comparisons as the modified VBS scheme, by construction, it does not include the formation of LV-OOA, resulting in an important overestimation of the SV-OOA at both sites.

A closer look at OA sources over the western Mediterranean basin simulated with the modified VBS scheme, selected because of its good results (at least for the summer of 2013 period and for this given region), shows that the OA with biogenic origins is dominant in the whole basin. In areas between Corsica and Marseilles, the ~~gulf~~ Gulf of Genoa and also the northern coast of Africa, the contribution of biogenic organic aerosols is less than for other parts. This fact points to the influence of shipping emissions for the areas between Marseilles and Corsica and also the northern coast of Africa, which can be seen in the contribution of POA and their oxidation products to the formation of OA over the basin, even if this part may be overestimated in the current simulations. For the gulf of Genoa, slightly higher contribution of anthropogenic organic aerosols was observed compared to other parts of the domain. However, at a higher altitude, the contribution of the biogenic sources becomes dominant in the whole basin, with a significant drop in the contribution of POA over the basin and leaving only a small trace of anthropogenic contribution in the gulf of Genoa. This contribution is attributed to ASOA rather than POA for the industrial area in the northern Italy, which is a persistent source of ASOA both near the surface and at higher altitudes.

It would be useful to compare the precursor components for the formation of OA such as isoprene and monoterpenes to multiple stations rather than only two, since the measurements for these species tend to have a more local profile-representativeness rather than a regional one, therefore multiple stations spread in the domain would give the needed regional aspect of the comparisons. Longer periods of simulations with comparisons to observations are also necessary, since processes leading to the formation of OA can change in other seasons and especially in winter when the biogenic contribution is much lower. Since airborne measurements for OA and biogenic gas-phase precursors were also performed during the summer of 2014 in the SAFMED+ campaign over different forested areas, it would be useful to continue the simulations for this year to compare the results to in-situ measurements and airborne measurements at the same time.

Acknowledgements. This research has received funding from the French National Research Agency (ANR) projects SAF-MED (grant ANR-15 12-BS06-0013). This work is part of the ChArMEx project supported by ADEME, CEA, CNRS-INSU and Météo-France through the multidisciplinary programme MISTRALS (Mediterranean Integrated Studies at Regional And Local Scales). The station at Ersa was partly supported by the
5 CORSiCA project funded by the Collectivité Territoriale de Corse through the Fonds Européen de Développement Régional of the European Operational Program 2007-2013 and the Contrat de Plan Etat-Région. The EEA is acknowledged for air quality data for several stations in Europe which was used for observation-simulation comparisons. The NCEP is acknowledged for the meteorological input data used in the WRF meteorological model. E-OBS datasets is acknowledged. The thesis work of Arineh Cholakian is supported by ADEME, INERIS
10 (with the support of the French Ministry in charge of Ecology), and via the ANR SAF-MED project. [Eric Hamonou and the ChArMEx team](#) are acknowledged for [their](#) great help in organizing the measurement campaign at Ersa, as well as the MISTRALS management and accounting team. Jorge Pey is currently granted with a Ramón y Cajal research contract (RYC-2013-14159) from the Spanish Ministry of Economy, Industry and Competitiveness.

15

This work was performed using HPC
resources from GENCI-CCRT
(Grant 2017- t2015017232).

References

- 20 Aiken, A. C., DeCarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty, K. S., Ulbrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y., Zhang, Q., Trimborn, A., Northway, M., Ziemann, P. J., Canagaratna, M. R., Onasch, T. B., Alfarra, M. R., Prevot, A. S. H., Dommen, J., Duplissy, J., Metzger, A., Baltensperger, U., and Jimenez, J. L.: O/C and OM/OC Ratios of Primary, Secondary, and Ambient Organic Aerosols with High-Resolution Time-of-Flight Aerosol Mass Spectrometry, *Environ. Sci. Technol.*, 42, 4478–4485,
25 doi:10.1021/es703009q, 2008.
- Arino, O., Bicheron, P., Achard, F., Latham, J., Witt, R., and Weber, J.: Globcover: The most detailed portrait of Earth, *Eur. Sp. Agency Bull.*, 36, 24–31, 2008.
- Arndt, J., Sciare, J., Mallet, M., Roberts, G. C., Marchand, N., Sartelet, K., Sellegri, K., Dulac, F., Healy, R. M., and Wenger, J. C.: Sources and mixing state of summertime background aerosol in the north-western
30 Mediterranean basin, *Atmos. Chem. Phys.*, 17, 6975–7001, doi:10.5194/acp-17-6975-2017, 2017.
- Bardouki, H., Liakakou, H., Economou, C., Sciare, J., Smolík, J., Ždímal, V., Eleftheriadis, K., Lazaridis, M., Dye, C., and Mihalopoulos, N.: Chemical composition of size-resolved atmospheric aerosols in the eastern Mediterranean during summer and winter, *Atmos. Environ.*, 37, 195–208, doi:10.1016/S1352-2310(02)00859-2, 2003.
- 35 Berezin, E. V., Konovalov, I. B., Ciais, P., Richter, A., Tao, S., Janssens-Maenhout, G., Beekmann, M., and Schulze, E.-D.: Multiannual changes of CO₂ emissions in China: indirect estimates derived from satellite measurements of tropospheric NO₂ columns, *Atmos. Chem. Phys.*, 13, 9415–9438, doi:10.5194/acp-13-9415-2013, 2013.
- Bessagnet, B., Menut, L., Curd, G., Hodzic, A., Guillaume, B., Liousse, C., Moukhtar, S., Pun, B., Seigneur, C., and Schulz, M.: Regional modeling of carbonaceous aerosols over Europe-focus on secondary organic
40 aerosols, *J. Atmos. Chem.*, 61, 175–202, doi:10.1007/s10874-009-9129-2, 2008.
- Di Biagio, C., Doppler, L., Gaimoz, C., Grand, N., Ancellet, G., Raut, J.-C., Beekmann, M., Borbon, A., Sartelet, K., Attié, J.-L., Ravetta, F. and Formenti, P.: Continental pollution in the western Mediterranean basin: vertical profiles of aerosol and trace gases measured over the sea during TRAQA 2012 and SAFMED 2013, *Atmos. Chem. Phys.*, 15, 9611–9630, doi:10.5194/acp-15-9611-2015, 2015.
- 45 Borrego, C., Souto, J. A., Dios, M., Monteiro, A., Ferreira, J., Rodriguez, A., Saavedra, S., Casares, J. J., and Miranda, A. I.: The role of transboundary air pollution over galicia and north portugal area, *Env. Sci Pollut Res*, 20, doi:10.1007/s11356-012-1201-9, 2013.
- Bowden, J. H., Otte, T. L., Nolte, C. G., Otte, M. J., Bowden, J. H., Otte, T. L., Nolte, C. G., and Otte, M. J.:

- Examining interior grid nudging techniques using two-way nesting in the wrf model for regional climate modeling, *J. Clim.*, 25, 2805–2823, doi:10.1175/JCLI-D-11-00167.1, 2012.
- Canonaco, F., Crippa, M., Slowik, J. G., Baltensperger, U., and Prévôt, A. S. H.: SoFi, an IGOR-based interface for the efficient use of the generalized multilinear engine (ME-2) for source apportionment: ME-2 application to aerosol mass spectrometer data, *Atmos. Meas. Tech.*, 6, 3649–3661, doi:10.5194/amt-6-3649-2013, 2013.
- Carlton, A. G., Wiedinmyer, C., and Kroll, J. H.: A review of Secondary Organic Aerosol (SOA) formation from isoprene, *Atmos. Chem. Phys.*, 9, 4987–5005, doi:10.5194/acp-9-4987-2009, 2009.
- Chen, F., Dudhia, J., Chen, F., and Dudhia, J.: Coupling an Advanced Land Surface–Hydrology Model with the Penn State–NCAR MM5 Modeling System. Part I: Model Implementation and Sensitivity, *Mon. Weather Rev.*, 129, 569–585, doi:10.1175/1520-0493(2001)129<0569:CAALSH>2.0.CO;2, 2001.
- Chin, M., Ginoux, P., Kinne, S., Torres, O., Holben, B. N., Duncan, B. N., Martin, R. V., Logan, J. A., Higurashi, A., and Nakajima, T.: Tropospheric Aerosol Optical Thickness from the GOCART Model and Comparisons with Satellite and Sun Photometer Measurements, *J. Atmos. Sci.*, 59, 461–483, doi:10.1175/1520-0469(2002)059<0461:TAOTFT>2.0.CO;2, 2002.
- Chrit, M., Sartelet, K., Sciare, J., Pey, J., Marchand, N., Couvidat, F., Sellegri, K., and Beekmann, M.: Modelling organic aerosol concentrations and properties during ChArMEx summer campaigns of 2012 and 2013 in the western Mediterranean region, *Atmos. Chem. Phys.*, 17, 12503–12531, doi:10.5194/acp-17-12509-2017, 2017.
- Colette, A., Andersson, C., Baklanov, A., Bessagnet, B., Brandt, J., Christensen, J. H., Doherty, R., Engardt, M., Geels, C., Giannakopoulos, C., Hedegaard, G. B., Katragkou, E., Langner, J., Lei, H., Manders, A., Melas, D., Meleux, F., Rouïl, L., Sofiev, M., Soares, J., Stevenson, D. S., Tombrou-Tzella, M., Varotsos, K. V and Young, P.: Is the ozone climate penalty robust in Europe?, *Environ. Res. Lett.*, 10, 84015, doi:10.1088/1748-9326/10/8/084015, 2015.
- Curci, G., Palmer, P. I., Kurosu, T. P., Chance, K. and Visconti, G.: Estimating European volatile organic compound emissions using satellite observations of formaldehyde from the Ozone Monitoring Instrument, *Atmos. Chem. Phys.*, 10, 11501–11517, doi:10.5194/acp-10-11501-2010, 2010.
- Decarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R. and Jimenez, J. L.: Field-Deployable, High-Resolution, Time-of-Flight Aerosol Mass Spectrometer, *Anal. Chem.*, 78, 8281–8289, doi:10.1021/ac061249n, 2006.
- Derognat, C.: Effect of biogenic volatile organic compound emissions on tropospheric chemistry during the Atmospheric Pollution Over the Paris Area (ESQUIF) campaign in the Ile-de-France region, *J. Geophys. Res.*, 108, doi:10.1029/2001JD001421, 2003.
- Donahue, N. M., Robinson, A. L., Stanier, C. O. and Pandis, S. N.: Coupled partitioning, dilution, and chemical aging of semivolatile organics, *Environ. Sci. Technol.*, 40, 2635–2643, doi:10.1021/es052297c, 2006.
- Donahue, N. M., Kroll, J. H., Pandis, S. N. and Robinson, A. L.: A two-dimensional volatility basis set-Part 2: Diagnostics of organic-aerosol evolution, *Atmos. Chem. Phys.*, 12, 615–634, doi:10.5194/acp-12-615-2012, 2012.
- Drinovec, L., Močnik, G., Zotter, P., Prévôt, A. S. H., Ruckstuhl, C., Coz, E., Rupakheti, M., Sciare, J., Müller, T., Wiedensohler, A. and Hansen, A. D. A.: The “dual-spot” Aethalometer: an improved measurement of aerosol black carbon with real-time loading compensation, *Atmos. Meas. Tech.*, 8, 1965–1979, doi:10.5194/amt-8-1965-2015, 2015.
- Fares, S., Schnitzhofer, R., Jiang, X., Guenther, A., Hansel, A. and Loreto, F.: Observations of diurnal to weekly variations of monoterpene-dominated fluxes of volatile organic compounds from mediterranean forests: implications for regional modeling., *Environ. Sci. Technol.*, 47, 11073–82, doi:10.1021/es4022156, 2013.
- Freney, E., Sellegri, K., Chrit, M., Adachi, K., Brito, J., Waked, A., Borbon, A., Colomb, A., Dupuy, R., Pichon, J.-M., Bouvier, L., Delon, C., Jambert, C., Durand, P., Bourianne, T., Gaimoz, C., Triquet, S., Féron, A., Beekmann, M., Dulac, F., and Sartelet, K.: Aerosol composition and the contribution of SOA formation over Mediterranean forests, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2017-482>, in review, 2017.
- Gangoiti, G., Millán, M. M., Salvador, R., and Mantilla, E.: Long-range transport and re-circulation of pollutants in the western Mediterranean during the project Regional Cycles of Air Pollution in the West-Central Mediterranean Area, *Atmos. Environ.*, 35, 6267–6276, doi:10.1016/S1352-2310(01)00440-X, 2001.
- Gelencsér, A., May, B., Simpson, D., Sánchez-Ochoa, A., Kasper-Giebl, A., Puxbaum, H., Caseiro, A., Pio, C. and Legrand, M.: Source apportionment of PM_{2.5} organic aerosol over Europe: Primary/secondary, natural/anthropogenic, and fossil/biogenic origin, *J. Geophys. Res.*, 112, D23S04, doi:10.1029/2006JD008094, 2007.
- Giorgi, F.: Climate change hot-spots, *Geophys. Res. Lett.*, 33, 1–4, doi:10.1029/2006GL025734, 2006.
- Griffin, R. J., Cocker, D. R., Flagan, R. C. and Seinfeld, J. H.: Organic aerosol formation from the oxidation of biogenic hydrocarbons, *J. Geophys. Res. Atmos.*, 104, 3555–3567, doi:10.1029/1998JD100049, 1999.
- Guenther, A., Karl, T., Harley, P., C.Wiedinmyer, Palmer, P. I. and Geron3, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, doi:10.1016/j.cognition.2008.05.007, 2006.

- El Haddad, I., Marchand, N., Wortham, H., Piot, C., Besombes, J.-L., Cozic, J., Chauvel, C., Armengaud, A., Robin, D. and Jaffrezo, J.-L.: Primary sources of PM_{2.5} organic aerosol in an industrial Mediterranean city, Marseille, *Atmos. Chem. Phys.*, 11, 2039–2058, doi:10.5194/acp-11-2039-2011, 2011.
- 5 Hauglustaine, D. A., Balkanski, Y. and Schulz, M.: A global model simulation of present and future nitrate aerosols and their direct radiative forcing of climate, *Atmos. Chem. Phys.*, 14, 11031–11063, doi:10.5194/acp-14-11031-2014, 2014.
- Haylock, M. R., Hofstra, N., Klein Tank, A. M. G., Klok, E. J., Jones, P. D. and New, M.: A European daily high-resolution gridded data set of surface temperature and precipitation for 1950–2006, *J. Geophys. Res.*, 113, D20119, doi:10.1029/2008JD010201, 2008.
- 10 Hildebrandt, L., Kostenidou, E., Mihalopoulos, N., Worsnop, D. R., Donahue, N. M. and Pandis, S. N.: Formation of highly oxygenated organic aerosol in the atmosphere: Insights from the Finokalia Aerosol Measurement Experiments, *Geophys. Res. Lett.*, 37, 6–10, doi:10.1029/2010GL045193, 2010.
- Hodzic, A. and Jimenez, J. L.: Modeling anthropogenically controlled secondary organic aerosols in a megacity: a simplified framework for global and climate models, *Geosci. Model Dev.*, 4, 901–917, doi:10.5194/gmd-4-901-2011, 2011.
- 15 Hofstra, N., Haylock, M., New, M. and Jones, P. D.: Testing E-OBS European high-resolution gridded data set of daily precipitation and surface temperature, *J. Geophys. Res.*, 114, D21101, doi:10.1029/2009JD011799, 2009.
- Hong, S.-Y., Dudhia, J., Chen, S.-H., Hong, S.-Y., Dudhia, J. and Chen, S.-H.: A Revised Approach to Ice Microphysical Processes for the Bulk Parameterization of Clouds and Precipitation, *Mon. Weather Rev.*, 132(1), 103–120, doi:10.1175/1520-0493(2004)132<0103:ARATIM>2.0.CO;2, 2004.
- 20 Janjic, Z. I.: A nonhydrostatic model based on a new approach, *Meteorol. Atmos. Phys.*, 82, 271–285, doi:10.1007/s00703-001-0587-6, 2003.
- Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G. and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, doi:10.5194/acp-5-1053-2005, 2005.
- 25 Koçak, M., Mihalopoulos, N. and Kubilay, N.: Chemical composition of the fine and coarse fraction of aerosols in the northeastern Mediterranean, *Atmos. Environ.*, 41, 7351–7368, doi:10.1016/j.atmosenv.2007.05.011, 2007.
- Konovalov, I. B., Beekmann, M., Berezin, E. V., Petetin, H., Mielonen, T., Kuznetsova, I. N. and Andreae, M. O.: The role of semi-volatile organic compounds in the mesoscale evolution of biomass burning aerosol: a modeling case study of the 2010 mega-fire event in Russia, *Atmos. Chem. Phys.*, 15, 13269–13297, doi:10.5194/acp-15-13269-2015, 2015.
- 35 Koulouri, E., Saarikoski, S., Theodosi, C., Markaki, Z., Gerasopoulos, E., Kouvarakis, G., Makela, T., Hillamo, R. and Mihalopoulos, N.: Chemical composition and sources of fine and coarse aerosol particles in the Eastern Mediterranean, *Atmos. Environ.*, 42, 6542–6550, doi:10.1016/j.atmosenv.2008.04.010, 2008.
- Kroll, J. H. and Seinfeld, J. H.: Chemistry of secondary organic aerosol: Formation and evolution of low-volatility organics in the atmosphere, *Atmos. Environ.*, 42(16), 3593–3624, doi:10.1016/j.atmosenv.2008.01.003, 2008.
- 40 Kuenen, J. J. P., Visschedijk, A. J. H., Jozwicka, M. and Denier Van Der Gon, H. A. C.: TNO-MACC-II emission inventory; A multi-year (2003–2009) consistent high-resolution European emission inventory for air quality modelling, *Atmos. Chem. Phys.*, 14, 10963–10976, doi:10.5194/acp-14-10963-2014, 2014.
- Lane, T. E., Donahue, N. M. and Pandis, S. N.: Simulating secondary organic aerosol formation using the volatility basis-set approach in a chemical transport model, *Atmos. Environ.*, 42, 7439–7451, doi:10.1016/j.atmosenv.2008.06.026, 2008.
- 45 Lanz, V. A., Prévôt, A. S. H., Alfarra, M. R., Weimer, S., Mohr, C., DeCarlo, P. F., Gianini, M. F. D., Hueglin, C., Schneider, J., Favez, O., D’Anna, B., George, C. and Baltensperger, U.: Characterization of aerosol chemical composition with aerosol mass spectrometry in Central Europe: an overview, *Atmos. Chem. Phys.*, 10, 10453–10471, doi:10.5194/acp-10-10453-2010, 2010.
- 50 Lelieveld, J., Lelieveld, J., Berresheim, H., Borrmann, S., Crutzen, P. J., Dentener, F. J., Fischer, H., Feichter, J., Flatau, P. J. and Heland, J.: Global Air Pollution Crossroads over the Mediterranean.pdf, , 794, doi:10.1126/science.1075457, 2002.
- Lionello, P. and Giorgi, F.: Winter precipitation and cyclones in the Mediterranean region: future climate scenarios in a regional simulation, *Adv. Geosci.*, 12, 153–158, doi:10.5194/adgeo-12-153-2007, 2007.
- 55 Liu, P., Tsimpidi, A. P., Hu, Y., Stone, B., Russell, A. G. and Nenes, A.: Differences between downscaling with spectral and grid nudging using WRF, *Atmos. Chem. Phys.*, 12, 3601–3610, doi:10.5194/acp-12-3601-2012, 2012.
- Mailler, S., Menut, L., Khvorostyanov, D., Valari, M., Couvidat, F., Siour, G., Turquety, S., Briant, R., Tuccella, P., Bessagnet, B., Colette, A., Létinois, L., Markakis, K., and Meleux, F.: CHIMERE-2017: from urban to hemispheric chemistry-transport modeling, *Geosci. Model Dev.*, 10, 2397–2423, doi:10.5194/gmd-10-2397-
- 60

2017, 2017.

- Mallet, M., Dulac, F., Formenti, P., Nabat, P., Sciare, J., Roberts, G., Pelon, J., Ancellet, G., Tanré, D., Parol, F., Denjean, C., Brogniez, G., Di Sarra, A., Alados-Arboledas, L., Arndt, J., Auriol, F., Blarel, L., Bourrienne, T., Chazette, P., Chevaillier, S., Claey's, M., D'Anna, B., Derimian, Y., Desboeufs, K., Di Iorio, T., Doussin, J. F., Durand, P., Féron, A., Freney, E., Gaimoz, C., Goloub, P., Gómez-Amo, J. L., Granados-Muñoz, M. J., Grand, N., Hamonou, E., Jankowiak, I., Jeannot, M., Léon, J. F., Maillé, M., Mailler, S., Meloni, D., Menut, L., Momboisse, G., Nicolas, J., Podvin, T., Pont, V., Rea, G., Renard, J. B., Roblou, L., Schepanski, K., Schwarzenboeck, A., Sellegri, K., Sicard, M., Solmon, F., Somot, S., Torres, B., Totems, J., Triquet, S., Verdier, N., Verwaerde, C., Waquet, F., Wenger, J. and Zapf, P.: Overview of the Chemistry-Aerosol Mediterranean Experiment/Aerosol Direct Radiative Forcing on the Mediterranean Climate (ChArMEx/ADRIMED) summer 2013 campaign, *Atmos. Chem. Phys.*, 16, 455–504, doi:10.5194/acp-16-455-2016, 2016.
- Martinelli, N., Olivieri, O., Girelli, D., Wilkins, E. T., Logan, W. P., Brunekreef, B., Holgate, S. T., Franchini, M., Mannucci, P. M., Franchini, M., Mannucci, P. M., Brunekreef, B., Forsberg, B., Venkataraman, C., Kao, A. S., Dockery, D. W., Pope, C. A., Xu, X., Spengler, J. D., Ware, J. H., Fay, M. E., al., et, Pope, C. A., Thun, M. J., Namboodiri, M. M., Dockery, D. W., Evans, J. S., Speizer, F. E., al., et, Brook, R. D., Rajagopalan, S., Pope, C. A., Brook, J. R., Bhatnagar, A., Diez-Roux, A. V., al., et, Brook, R. D., Franklin, B., Cascio, W., Hong, Y., Howard, G., Lipsett, M., al., et, Kloog, I., Coull, B. A., Zanobetti, A., Koutrakis, P., Schwartz, J. D., Bruzzi, P., Green, S. B., Byar, D. P., Brinton, L. A., Schairer, C., Nawrot, T. S., Perez, L., Künzli, N., Munters, E., Nemery, B., Zanobetti, A., Schwartz, J., Dominici, F., Peng, R. D., Bell, M. L., Pham, L., McDermott, A., Zeger, S. L., al., et, Puett, R. C., Schwartz, J., Hart, J. E., Yanosky, J. D., Speizer, F. E., Suh, H., al., et, Peters, A., Klot, S. von, Heier, M., Trentinaglia, I., Hörmann, A., Wichmann, H. E., al., et, Pope, C. A., Muhlestein, J. B., May, H. T., Renlund, D. G., Anderson, J. L., Horne, B. D., Pope, C. A., Burnett, R. T., Thurston, G. D., Thun, M. J., Calle, E. E., Krewski, D., al., et, Atkinson, R. W., Carey, I. M., Kent, A. J., Staa, T. P. van, Anderson, H. R., et al.: Air particulate matter and cardiovascular disease: a narrative review., *Eur. J. Intern. Med.*, 24, 295–302, doi:10.1016/j.ejim.2013.04.001, 2013.
- Mauderly, J. L. and Chow, J. C.: Health Effects of Organic Aerosols, *Inhal. Toxicol.*, 20, 257–288, doi:10.1080/08958370701866008, 2008.
- Menut, L., Bessagnet, B., Khvorostyanov, D., Beekmann, M., Blond, N., Colette, A., Coll, I., Curci, G., Foret, G., Hodzic, A., Mailler, S., Meleux, F., Monge, J.-L., Pison, I., Siour, G., Turquety, S., Valari, M., Vautard, R. and Vivanco, M. G.: CHIMERE 2013: a model for regional atmospheric composition modelling, *Geosci. Model Dev.*, 6, 981–1028, doi:10.5194/gmd-6-981-2013, 2013.
- Menut, L., Rea, G., Mailler, S., Khvorostyanov, D. and Turquety, S.: Aerosol forecast over the Mediterranean area during July 2013 (ADRIMED/CHARMEX), *Atmos. Chem. Phys.*, 15, 7897–7911, doi:10.5194/acp-15-7897-2015, 2015.
- Michoud, V., Sciare, J., Sauvage, S., Dusanter, S., Léonardis, T., Gros, V., Kalogridis, C., Zannoni, N., Féron, A., Petit, J.-E., Crenn, V., Baisnée, D., Sarda-Estève, R., Bonnaire, N., Marchand, N., DeWitt, H. L., Pey, J., Colomb, A., Gheusi, F., Szidat, S., Stavroulas, I., Borbon, A., and Locoge, N.: Organic carbon at a remote site of the western Mediterranean Basin: sources and chemistry during the ChArMEx SOP2 field experiment, *Atmos. Chem. Phys.*, 17, 8837–8865, doi:10.5194/acp-17-8837-2017, 2017.
- Minguillón, M. C., Perron, N., Querol, X., Szidat, S., Fahrni, S. M., Alastuey, A., Jimenez, J. L., Mohr, C., Ortega, A. M., Day, D. A., Lanz, V. A., Wacker, L., Reche, C., Cusack, M., Amato, F., Kiss, G., Hoffer, A., Decesari, S., Moretti, F., Hillamo, R., Teinilä, K., Seco, R., Peñuelas, J., Metzger, A., Schallhart, S., Müller, M., Hansel, A., Burkhardt, J. F., Baltensperger, U. and Prévôt, A. S. H.: Fossil versus contemporary sources of fine elemental and organic carbonaceous particulate matter during the DAURE campaign in Northeast Spain, *Atmos. Chem. Phys.*, 11, 12067–12084, doi:10.5194/acp-11-12067-2011, 2011.
- Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J. and Clough, S. A.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, *J. Geophys. Res. Atmos.*, 102, 16663–16682, doi:10.1029/97JD00237, 1997.
- Mohr, C., DeCarlo, P. F., Heringa, M. F., Chirico, R., Slowik, J. G., Richter, R., Reche, C., Alastuey, A., Querol, X., Seco, R., Peñuelas, J., Jiménez, J. L., Crippa, M., Zimmermann, R., Baltensperger, U. and Prévôt, A. S. H.: Identification and quantification of organic aerosol from cooking and other sources in Barcelona using aerosol mass spectrometer data, *Atmos. Chem. Phys.*, 12, 1649–1665, doi:10.5194/acp-12-1649-2012, 2012.
- Monks, P. S., Granier, C., Fuzzi, S., Stohl, A., Williams, M. L., Akimoto, H., Amann, M., Baklanov, A., Baltensperger, U., Bey, I., Blake, N., Blake, R. S., Carslaw, K., Cooper, O. R., Dentener, F., Fowler, D., Fragkou, E., Frost, G. J., Generoso, S., Ginoux, P., Grewe, V., Guenther, A., Hansson, H. C., Henne, S., Hjorth, J., Hofzumahaus, A., Huntrieser, H., Isaksen, I. S. A., Jenkin, M. E., Kaiser, J., Kanakidou, M., Klimont, Z., Kulmala, M., Laj, P., Lawrence, M. G., Lee, J. D., Liousse, C., Maione, M., McFiggans, G., Metzger, A., Mieville, A., Moussiopoulos, N., Orlando, J. J., O'Dowd, C. D., Palmer, P. I., Parrish, D. D., Petzold, A., Platt, U., Pöschl, U., Prévôt, A. S. H., Reeves, C. E., Reimann, S., Rudich, Y., Sellegri, K., Steinbrecher, R., Simpson,

- D., ten Brink, H., Theloke, J., van der Werf, G. R., Vautard, R., Vestreng, V., Vlachokostas, C. and von Glasow, R.: Atmospheric composition change – global and regional air quality, *Atmos. Environ.*, 43, 5268–5350, doi:10.1016/j.atmosenv.2009.08.021, 2009.
- 5 Murphy, B. N. and Pandis, S. N.: Simulating the Formation of Semivolatile Primary and Secondary Organic Aerosol in a Regional Chemical Transport Model, *Environ. Sci. Technol.*, 43, 4722–4728, doi:10.1021/es803168a, 2009.
- Nabat, P., Solmon, F., Mallet, M., Kok, J. F., and Somot, S.: Dust emission size distribution impact on aerosol budget and radiative forcing over the Mediterranean region: a regional climate model approach, *Atmos. Chem. Phys.*, 12, 10545–10567, doi:10.5194/acp-12-10545-2012, 2012.
- 10 Nabat, P., Somot, S., Mallet, M., Chiapello, I., Morcrette, J. J., Solmon, F., Szopa, S., Dulac, F., Collins, W., Ghan, S., Horowitz, L. W., Lamarque, J. F., Lee, Y. H., Naik, V., Nagashima, T., Shindell, D. and Skeie, R.: A 4-D climatology (1979–2009) of the monthly tropospheric aerosol optical depth distribution over the Mediterranean region from a comparative evaluation and blending of remote sensing and model products, *Atmos. Meas. Tech.*, 6, 1287–1314, doi:10.5194/amt-6-1287-2013, 2013.
- 15 Ng, N. L., Herndon, S. C., Trimborn, A., Canagaratna, M. R., Croteau, P. L., Onasch, T. B., Sueper, D., Worsnop, D. R., Zhang, Q., Sun, Y. L. and Jayne, J. T.: An Aerosol Chemical Speciation Monitor (ACSM) for Routine Monitoring of the Composition and Mass Concentrations of Ambient Aerosol, *Aerosol Sci. Technol.*, 45, 780–794, doi:10.1080/02786826.2011.560211, 2011.
- Odum, J. R., Jungkamp, T. P. W., Griffin, R. J., Forstner, H. J. L., R. C. Flagan, A. and Seinfeld*, J. H.: Aromatics, Reformulated Gasoline, and Atmospheric Organic Aerosol Formation, , doi:10.1021/ES960535L, 1997.
- 20 Paatero, P.: Least squares formulation of robust non-negative factor analysis, *Chemom. Intell. Lab. Syst.*, 37, 23–35, doi:10.1016/S0169-7439(96)00044-5, 1997.
- Paatero, P. and Tapper, U.: Positive matrix factorization: A non-negative factor model with optimal utilization of error estimates of data values, *Environmetrics*, 5, 111–126, doi:10.1002/env.3170050203, 1994.
- 25 Pankow, J. F.: Review and comparative analysis of the theories on partitioning between the gas and aerosol particulate phases in the atmosphere, *Atmos. Environ.*, 21, 2275–2283, doi:10.1016/0004-6981(87)90363-5, 1987.
- Petetin, H., Beekmann, M., Sciare, J., Bressi, M., Rosso, A., Sanchez, O. and Gherzi, V.: A novel model evaluation approach focusing on local and advected contributions to urban PM_{2.5} levels - Application to Paris, France, *Geosci. Model Dev.*, 7, 1483–1505, doi:10.5194/gmd-7-1483-2014, 2014.
- 30 Pey, J., Querol, X., Alastuey, A., Forastiere, F. and Stafoggia, M.: African dust outbreaks over the Mediterranean Basin during 2001–2011: PM₁₀ concentrations, phenomenology and trends, and its relation with synoptic and mesoscale meteorology, *Atmos. Chem. Phys.*, 13, 1395–1410, doi:10.5194/acp-13-1395-2013, 2013.
- Pieber, S. M., Zhao, Y., Orasche, J., Stengel, B., Czech, H., Corbin, J. C., Haddad, I. El, Klein, F., Kilic, D., Slowik, J. G., Donahue, N., Robinson, A., Zimmermann, R., Baltensperger, U. and Prévôt, A. S. H.: Characterization of organic low-, semi-, intermediate-and volatile organic compounds from four-stroke ship engine emissions: implications for atmospheric processing, *EAC Abstr.*, 2016.
- Pöschl, U.: Atmospheric Aerosols: Composition, Transformation, Climate and Health Effects, *Angew. Chem. Int. Ed.*, 44, 7520–7540, doi:10.1002/anie.200501122, 2005.
- 40 Pun, B. K. and Seigneur, C.: Investigative modeling of new pathways for secondary organic aerosol formation, *Atmos. Chem. Phys.*, 7, 2199–2216, doi:10.5194/acp-7-2199-2007, 2007.
- Querol, X., Alastuey, A., Pey, J., Cusack, M., Pérez, N., Mihalopoulos, N., Theodosi, C., Gerasopoulos, E., Kubilay, N., and Koçak, M.: Variability in regional background aerosols within the Mediterranean, *Atmos. Chem. Phys.*, 9, 4575–4591, doi:10.5194/acp-9-4575-2009, 2009.
- 45 Rea, G., Turquety, S., Menut, L., Briant, R., Mailler, S. and Siour, G.: Source contributions to 2012 summertime aerosols in the Euro-Mediterranean region, *Atmos. Chem. Phys.*, 15, 8013–8036, doi:10.5194/acp-15-8013-2015, 2015.
- Ripoll, A., Pey, J., Minguillón, M. C., Pérez, N., Pandolfi, M., Querol, X. and Alastuey, A.: Three years of aerosol mass, black carbon and particle number concentrations at Montsec (southern Pyrenees, 1570 m a.s.l.), *Atmos. Chem. Phys.*, 14, 4279–4295, doi:10.5194/acp-14-4279-2014, 2014.
- 50 Robinson, A. L., Donahue, N. M., Shrivastava, M. K., Weitkamp, E. A., Sage, A. M., Grieshop, A. P., Lane, T. E., Pierce, J. R. and Pandis, S. N.: Rethinking Organic Aerosols: Semivolatile Emissions and Photochemical Aging, *Science (80-.)*, 315, 1259–1262, doi:10.1126/science.1133061, 2007a.
- Robinson, A. L., Donahue, N. M., Shrivastava, M. K., Weitkamp, E. A., Sage, A. M., Grieshop, A. P., Lane, T. E., Pierce, J. R. and Pandis, S. N.: Rethinking Organic Aerosols: Semivolatile Emissions and Photochemical Aging, *Science (80-.)*, 315, 1259–1262, doi:10.1126/science.1133061, 2007b.
- 55 Sasser, E., Hemby, J., Adler, K., Anenberg, S., Bailey, C., Brockman, L., Chappell, L., DeAngelo, B., Damberg, R., Dawson, J., Frank, N., Geller, M., Hagler, G., Hemming, B., Jantarasami, L., Luben, T., Mitchell, J. and Moss, J.: Report to Congress on Black Carbon, EPA, (March), 2012.
- 60 Sciare, J., Oikonomou, K., Cachier, H., Mihalopoulos, N., Andreae, M. O., Maenhaut, W. and Sarda - Es eve, R.:

Aerosol mass closure and reconstruction of the light scattering coefficient over the Eastern Mediterranean Sea during the MINOS campaign, *Atmos. Chem. Phys.*, 5, 2253–2265 [online] Available from: www.atmos-chem-phys.org/acp/5/2253/ (Accessed 24 March 2017), 2005.

- 5 Sciare, J., Oikonomou, K., Favez, O., Liakakou, E., Markaki, Z., Cachier, H. and Mihalopoulos, N.: Long-term measurements of carbonaceous aerosols in the Eastern Mediterranean: evidence of long-range transport of biomass burning, *Atmos. Chem. Phys.*, 8, 5551–5563 [online] Available from: www.atmos-chem-phys.net/8/5551/2008/ (Accessed 24 March 2017), 2008.
- 10 Sellegri, K., Gourdeau, J., Putaud, J.-P. and Despiiau, S.: Chemical composition of marine aerosol in a Mediterranean coastal zone during the FETCH experiment, *J. Geophys. Res. Atmos.*, 106, 12023–12037, doi:10.1029/2000JD900629, 2001.
- Shrivastava, M., Fast, J., Easter, R., Gustafson, W. I., Zaveri, R. A., Jimenez, J. L., Saide, P. and Hodzic, A.: Modeling organic aerosols in a megacity: Comparison of simple and complex representations of the volatility basis set approach, *Atmos. Chem. Phys.*, 11, 6639–6662, doi:10.5194/acp-11-6639-2011, 2011.
- 15 Shrivastava, M., Zelenyuk, A., Imre, D., Easter, R., Beranek, J., Zaveri, R. A. and Fast, J.: Implications of low volatility SOA and gas-phase fragmentation reactions on SOA loadings and their spatial and temporal evolution in the atmosphere, *J. Geophys. Res. Atmos.*, 118, 3328–3342, doi:10.1002/jgrd.50160, 2013.
- Shrivastava, M., Easter, R. C., Liu, X., Zelenyuk, A., Singh, B., Zhang, K., Ma, P., Chand, D., Ghan, S., Jimenez, J. L., Zhang, Q., Fast, J., Rasch, P. J. and Tiitta, P.: Global transformation and fate of SOA: Implications of low-volatility SOA and gas-phase fragmentation reactions, *J. Geophys. Res. Atmos.*, 1–27, doi:10.1002/2014JD022563. Received, 2015.
- 20 Sič, B., El Amraoui, L., Piacentini, A., Marécal, V., Emili, E., Cariolle, D., Prather, M. and Attié, J.-L.: Aerosol data assimilation in the chemical transport model MOCAGE during the TRAQA/ChArMEx campaign: aerosol optical depth, *Atmos. Meas. Tech.*, 9, 5535–5554, doi:10.5194/amt-9-5535-2016, 2016.
- 25 Solazzo, E., Bianconi, R., Pirovano, G., Matthias, V., Vautard, R., Moran, M. D., Wyat Appel, K., Bessagnet, B., Brandt, J., Christensen, J. H., Chemel, C., Coll, I., Ferreira, J., Forkel, R., Francis, X. V., Grell, G., Grossi, P., Hansen, A. B., Miranda, A. I., Nopmongkol, U., Prank, M., Sartelet, K. N., Schaap, M., Silver, J. D., Sokhi, R. S., Vira, J., Werhahn, J., Wolke, R., Yarwood, G., Zhang, J. and Galmarini, S.: Operational model evaluation for particulate matter in Europe and North America in the context of AQMEII, *Atmos. Environ.*, 53, 75–92, doi:10.1016/j.atmosenv.2012.02.045, 2012.
- 30 Stull, R. B.: Mean Boundary Layer Characteristics, in *An Introduction to Boundary Layer Meteorology*, pp. 1–27, Springer Netherlands, Dordrecht., 1988.
- Vincent, J., Laurent, B., Losno, R., Bon Nguyen, E., Roullet, P., Sauvage, S., Chevaillier, S., Coddeville, P., Ouboulmane, N., di Sarra, A. G., Tovar-Sánchez, A., Sferlazzo, D., Massanet, A., Triquet, S., Morales Baquero, R., Fournier, M., Coursier, C., Desboeufs, K., Dulac, F. and Bergametti, G.: Variability of mineral dust deposition in the western Mediterranean basin and south-east of France, *Atmos. Chem. Phys.*, 16, 8749–8766, doi:10.5194/acp-16-8749-2016, 2016.
- 35 Zare, A., Christensen, J. H., Irannejad, P. and Brandt, J.: Evaluation of two isoprene emission models for use in a long-range air pollution model, *Atmos. Chem. Phys.*, 12, 7399–7412, doi:10.5194/acp-12-7399-2012, 2012.
- 40 Zhang, Q. J., Beekmann, M., Drewnick, F., Freutel, F., Schneider, J., Crippa, M., Prevot, A. S. H., Baltensperger, U., Poulain, L., Wiedensohler, A., Sciare, J., Gros, V., Borbon, A., Colomb, A., Michoud, V., Doussin, J. F., Denier Van Der Gon, H. A. C., Haeffelin, M., Dupont, J. C., Siour, G., Petetin, H., Bessagnet, B., Pandis, S. N., Hodzic, A., Sanchez, O., Honor??, C. and Perrussel, O.: Formation of organic aerosol in the Paris region during the MEGAPOLI summer campaign: Evaluation of the volatility-basis-set approach within the CHIMERE model, *Atmos. Chem. Phys.*, 13, 5767–5790, doi:10.5194/acp-13-5767-2013, 2013.
- 45 Zhang, Q. J., Beekmann, M., Drewnick, F., Freutel, F., Schneider, J., Crippa, M., Prevot, A. S. H., Baltensperger, U., Poulain, L., Wiedensohler, A., Sciare, J., Gros, V., Borbon, A., Colomb, A., Michoud, V., Doussin, J. F., Denier Van Der Gon, H. A. C., Haeffelin, M., Dupont, J. C., Siour, G., Petetin, H., Bessagnet, B., Pandis, S. N., Hodzic, A., Sanchez, O., Honor??, C. and Perrussel, O.: Formation of organic aerosol in the Paris region during the MEGAPOLI summer campaign: Evaluation of the volatility basis set approach within the CHIMERE model, *Atmos. Chem. Phys.*, 13, 5767–5790, doi:10.5194/acp-13-5767-2013, 2013.
- 50

Tables

10

1: Ersa, France (42°58'04.1", 9°22'49.1")

Gases: NO_x (CRANOX), VOCs (PTR-ToF-MS-KORE)

Aerosols: PM₁₀ total mass (TEOM-FDMS), On-line (non refractive) PM₁

Chemistry : OA, SO₄²⁻, NH₄⁺, NO₃⁻ (ACSM), ¹⁴C (PM₁, daily filters),
BC (PM_{2.5}, MAAP)

Meteorology: temperature, wind speed, wind direction, relative humidity

2: Es Pinar, Spain (39°53'04.6", 3°11'40.9")

Gases: NO_x (~~API Teledyne T200~~CRANOX), VOCs (PTR-ToF-MS, Ionikon)

Aerosols: PM₁₀ total mass (BETA corrected by factor obtained after

comparison with gravimetric), PM₁ on-line Chemistry: OA, SO₄²⁻,

NH₃⁺, NO₃⁻ (HR-ToF-AMS), ¹⁴C (PM₁, daily filters),

BC (PM_{2.5}, Aethalometer AE33)

Meteorology: temperature, wind speed, wind direction, relative humidity

3 : Palma, Spain (39° 36' 20.88", 2° 42' 23.7594"), 4 : Nîmes, France (43° 51' 21.6", 4°

24' 21.5994"), 5 : Ajaccio, France (41° 55' 5.3256", 8° 47' 38.0538")

Radio-soundings : T, RH, wind speed, wind direction; at 00 and 12 UTC each day

Table 21. Gas/aerosol and meteorological measurements used for this study.

15

20

25

Pollutant	ORE (%)	Pollutant	ORE (%)	Pollutant	ORE (%)	Parameter	ORE (%)
O ₃	4	C ₅ H ₈	85	Mono-terpenes	59	Temperature	0.5
OA	10	BC	26	SO ₂	62	Relative humidity	18
SO ₄ ²⁻	15	NO _x	75	Aromatic species	49		
PM ₁₀	9	CO	2	MACR+MVK	60		

Table 23. Calculated relative orographic representativeness error (ORE) for a list of species and meteorological parameters. MACR+MVK presents the sum of methyl vinyl ketone and methacrolein.

10

15

20

25

30

	<u>Ersa</u>					<u>Es Pinar</u>			
	<u>ORE</u>	<u>R</u>	<u>RMSE</u>	<u>Bias</u>	<u>Mean_obs</u>	<u>R</u>	<u>RMSE</u>	<u>Bias</u>	<u>Mean_obs</u>
	<u>%</u>	<u>ppb</u>							
<u>NO_x</u>	<u>75</u>	<u>0.37</u>	<u>0.68</u>	<u>0.16</u>	<u>0.62</u>	<u>0.12</u> <u>(0.11)</u>	<u>2.76</u> <u>(2.56)</u>	<u>-1.51</u> <u>(0.35)</u>	<u>3.53</u> <u>(3.53)</u>
<u>C₅H₈</u>	<u>85</u>	<u>0.76</u>	<u>0.44</u>	<u>0.24</u>	<u>0.19</u>	<u>0.69</u>	<u>0.10</u>	<u>-0.04</u>	<u>0.17</u>
<u>MACR+MVK</u>	<u>60</u>	<u>0.62</u>	<u>0.22</u>	<u>0.09</u>	<u>0.09</u>	<u>0.41</u>	<u>0.09</u>	<u>0.005</u>	<u>0.10</u>
<u>Mono terpenes</u>	<u>59</u>	<u>0.35</u>	<u>0.69</u>	<u>-0.38</u>	<u>0.52</u>	<u>0.14</u>	<u>0.11</u>	<u>-0.06</u>	<u>0.09</u>

10 **Table 34.** Statistical data for time series shown in figure 56, Mean_obs shows the average of observations. Values in parenthesis for Es Pinar NO_x statistics show the comparison of NO_y simulations to NO_x measurements.

5

10

		Ersa				Es Pinar			
	ORE	R	RMSE	Bias	Mean_obs	R	RMSE	Bias	Mean_obs
	%	$\mu\text{g.m}^{-3}$							
BC	26	0.36	0.16	0.02	0.39	0.47	0.22	-0.13	0.39
SO ₄ ²⁻	15	0.42	1.72	1.21	1.90	0.52	1.93	0.91	2.70

Table 45. Statistical data for time series shown in figure 6, Mean_obs column shows the average of observations.

15

20

25

30

35

5

10

		CHIMERE standard		Standard VBS ba		Standard VBS nba		Modified VBS	
		Ersa	Es Pinar	Ersa	Es Pinar	Ersa	Es Pinar	Ersa	Es Pinar
R		0.46	0.16	0.50	0.11	0.45	0.22	0.50	0.29
RMSE	$\mu\text{g}\cdot\text{m}^{-3}$	3.96	2.65	9.63	7.29	2.17	1.89	1.59	1.49
Bias		3.39	2.49	9.05	7.17	1.26	1.25	-0.68	-0.65
Mean_sim		7.10	5.36	12.76	10.17	4.97	4.13	3.02	2.22
Mean_obs		3.71	2.88	3.71	2.88	3.71	2.88	3.71	2.88
SD_sim		2.21	1.64	3.78	3.62	1.73	1.34	0.95	0.79
SD_obs		1.66	1.16	1.66	1.16	1.66	1.16	1.66	1.16

Table 56. Statistical data for hourly time series shown in figure 7. R is the correlation coefficient between model and measurements, RMSE the root mean square error, mean_sim and mean_obs show the average and SD_sim and SD_obs show the standard deviation of simulations and observations respectively.

15

20

25


30

Figures

10

15

20



25

Domain name	Cell numbers	Resolution
D30	71×64	30km×30km
D10	110×110	10km×10km
D3	110×110	3km×3km
D1	80×80	1km×1km

Figure 1. The four domains used in simulations (, in order of size: D30, D10, D3, D1). The resolution for each domain is given in the table below the image.




Figure 2. Organic aerosol simulation schemes - a: CHIMERE standard scheme (Bessagnet et al., 2008): from a parent volatile organic compounds (VOC), different semi-volatile VOC (SVOC) compounds (only one represented) are formed in a single step by oxidation; they are in equilibrium between gas and aerosol phase (pSVOC); b: VBS standard scheme (Robinson et al., 2007b): from a parent VOC, SVOC with regularly spaced volatility ranges are formed and are in equilibrium with the aerosol phase. Aging of SVOC by functionalization is included by passing species to classes with lower volatility; c: modified VBS scheme (Shrivastava et al., 2015): here SVOC aging also includes fragmentation, leading to transfer of species to classes with higher volatility. In addition, semi-volatile aerosol can be irreversibly transformed into non-volatile one (yellow-filled circle). For each bin saturation concentration is shown in $\mu\text{g}\cdot\text{m}^{-3}$. Note that this schematic represents BSOA/ASOA (Biogenic/anthropogenic secondary organic aerosol) where four bins are used, for SVOC/IVOC (Semi/Intermediate volatility organic compound, where 9 bins are used) a schematic is presented in SI-1.



Figure 3. Experimental data sites, including in-situ surface stations (1: Ersaa, France; 2: Cap Es Pinar, Spain), and meteorological sounding stations (3: Palma, Spain; 4: Nîmes, France; 5: Ajaccio, France). More details about exact location and measurements performed in table 12. 1- Ersaa, France, 2- Cap Es Pinar, Spain, 3- Palma, Spain, 4- Nîmes, France, 5- Ajaccio, France

5

10

15

20




Figure 4. Orographic representativeness error- a: neighbouring cells used in the orographic representativeness test; b: an example of non-linear regressions performed on one time-step for **organic aerosol (OA)**, 1 point corresponds to 1 grid cell); c: **results from all hourly simulations for OA** results for all hourly simulation times for **OA**. In b and c, the purple ribbon shows the confidence interval of the regression results. In c, the blue line shows the simulations at the nominal Ersa site grid cell.




Figure 5. Time series showing the comparison of simulated (in blue) and measured (in black) gaseous species during the ChArMEx/SAFMED campaign period. a: nitrogen oxides (time series in red for Es Pinar represents NO_y concentrations); b: isoprene; c: MACR+MVK (methyl vinyl ketone+methacrolein), d: monoterpenes. Statistical data for these comparisons are given in table 23. Beside each time table two points are shown presenting the average for simulations (in blue circle) and observations (in black square). Ribbon around Ersa simulations presents orographic representativeness error.




Figure 6. As for figure 5, but for particulate matter. a: BC (**black carbon**); b: sulfate particles. Statistical data for these comparisons [areis](#) given in table [45](#).

5

10

15

20

25




Figure 7. Compared time series of total PM₁ organic aerosol (OA) concentration at Ersa (top) and Es Pinar (bottom) – black: observations; green: modified VBS; brown: standard VBS without BSOA aging; blue: CHIMERE standard; red: standard VBS with BSOA aging. For each site the average of simulations is shown with circles in corresponding colors and the observations with black squares. Shaded area for Ersa presents the orographic representativeness error. Statistical data is shown in table [56](#).




Figure 8. Comparison of simulations to ^{14}C measurements - a: comparison of filters to four schemes for Erska; b: comparison of filters to four schemes for Es Pinar. For both figures from outer ring to inner ring: observations, modified VBS, standard VBS with BSOA aging, standard VBS without BSOA (biogenic secondary organic aerosol) aging and CHIMERE standard scheme.

5




Figure 9. Comparison of simulations to ^{14}C measurements - a: Erska individual filters; b: Es Pinar individual filters. For each filter, measurements are on left and modified VBS simulations are on right. Dark/light brown shows fossil and dark/light green shows non-fossil sources.





Figure 10. Distribution of simulated organic aerosol (OA) in volatility bins - a: Ersa volatility bins; b: Es Pinar volatility bins.



5

Figure 11. Comparison to PMF results - a: comparison of observed PMF factors to that derived from four schemes for Ersa; b: for Es Pinar. For both figures from outer ring to inner ring: observations, modified VBS, standard VBS with BSOA aging, standard VBS without BSOA (biogenic secondary organic aerosol) aging and CHIMERE standard scheme. LVOOA refers to low-volatility oxidized organic aerosol, SVOOA to semi-volatile oxidized organic aerosol and HOA to Hydrocarbon-like Organic Aerosol.




Figure 12. Average organic aerosol (OA) concentrations over the western Mediterranean basin simulated from 10 July to 9 August 2013. Simulations were performed with the modified VBS scheme. Left column is near the surface, and right column for an altitude between 300-450 m asl. From top to bottom: a-b: total OA concentration ($\mu\text{g m}^{-3}$); c-d: BSOA (Biogenic secondary OA) concentration ($\mu\text{g m}^{-3}$); e-f: total ASOA (Anthropogenic secondary OA concentration) concentration ($\mu\text{g m}^{-3}$); g-h: sum of POA (Primary OA) and its subsequent oxidations products ($\mu\text{g m}^{-3}$). Results are from the D3 simulation (3-km horizontal resolution) within the framed area, from the D10 simulation outside (10-km resolution).




Figure 13. As figure 12 but for relative contributions of BSOA/ASOA/POA to total OA (%). From top to bottom: a-b: biogenic secondary organic aerosol (BSOA); c-d: anthropogenic secondary organic aerosol (ASOA); e-f: POA and its subsequent oxidation products.