

Supplement of

Agricultural ammonia emissions in China: reconciling bottom-up and top-down estimates

Lin Zhang, et al.

5 Correspondence to: Lin Zhang (zhanglg@pku.edu.cn)

Table S1. Dates of fertilizer application for main crops in China¹

	Crops	First	Second	Third	Fourth	Fifth	Deviation ²
Number of day after the planting date	Early rice	10	20	30	45	80	32
	Late rice	10	20	30	45	80	32
	Spring wheat	0	20	45	\	\	29
	Winter wheat	0	30	168	\	\	27
	Spring maize	0	15	25	40		36
	Summer maize	0	15	25	40		36
	Cotton	0	14	54	100	110	24
	Sweet potato	0	10	60	80	\	17
	Potato	12	29	37	\	\	23
	Rapeseed	0	30	144	151	\	30
	Soybean	0	16	71	\	\	32
	Peanut	0	20	\	\	\	22
Calendar day of year	Tobacco	90	99	109	132	\	9
	Apple	305	64	115	152	274	9
	Banana	244	35	181	\	\	9
	Grape	274	91	158	188	\	9
	Citrus	319	74	110	140	213	9
	Pear	60	105	135	316	\	9

¹Timing of fertilizer application (up to five times) based on Liao (1993) and Zhang and Zhang (2012).

²Mean deviation (number of days) from the fertilizer application dates for each crop over China used in this study (σ in Eq. (4) in the main text).

15 **Table S2.** Fertilizer application rates for main crops in China¹

Crops	First	Second	Third	Fourth	Fifth	Total
Early rice	0.8	0.8	55.2	55.2	27.6	139.6
Late rice	0.8	0.8	41.3	70.5	16.5	129.9
Spring wheat	48.9	36.4	36.4	\	\	121.7
Winter wheat	48.9	36.4	36.4	\	\	121.7
Spring maize	71	13	22	22	\	128
Summer maize	71	13	22	22	\	128
Cotton	61	9.3	16.2	16.2	29	131.7
Sweet potato	62.5	5.2	6.9	6.9	\	81.5
Potato	65.5	77	65.5	\	\	208
Rapeseed	105	64	20	25	\	214
Soybean	16.2	13.4	16.2	\	\	45.8
Peanut	70	21	\	\	\	91
Tobacco	1.3	18	30	42	\	91.3
Apple	150	30	60	30	30	300
Banana	144	108	108	\	\	360
Grape	124	37.2	62	24.8	\	248
Citrus	162	162	107	18	25	474
Pear	108	81	53	28	\	270

¹Fertilizer application rates in unit of kg N ha⁻¹ are based on Liao (1993) and Zhang and Zhang (2012). Timing of fertilizer application (up to five times) is shown in Table S1.

Table S3. Emission factor index for synthetic fertilizer application¹

Factor	Value
Crop type	
Upland crops	-0.045
Flooded crops	0
Fertilizer type	
Ammonium sulfate	0.429
Urea	0.666
Ammonium nitrate	-0.35
Other straight N	-0.507
Compound NPK	0.014
Ammonium bicarbonate ²	0.928
Application mode	
Broadcast	-1.305
Injection	-1.895
Soil pH ³	$0.067 \times \text{pH}^2 - 0.69 \times \text{pH} + 0.68$
Soil CEC	
CEC ≤ 16	0.088
$16 < \text{CEC} \leq 24$	0.012
$24 < \text{CEC} \leq 32$	0.163
CEC > 32	0

20 ¹Values used for calculating NH₃ emission factors from fertilizer application (Eq. (5) in the main text) based on Bouwman et al. (2002) with modifications noted in the table.

²The value for ammonium bicarbonate is corrected according to observations from Zhu et al. (1989).

25 ³For soil pH, a second-order function is applied to fit the segmental values given in Bouwman et al. (2002).

Table S4. A summary of reported emission factors (EF) for NH₃ from fertilizer application in China

Study	Region	Method ¹	Fertilizer type ²	pH	CEC ³	Crops	Application mode ⁴	EF ⁵
Su et al., 2006	Beijing	WTS	U	8	<16	Bare soil	b/f	25.4
Su et al., 2006	Beijing	WTS	U	8	<16	Bare soil	b/f	23
Su et al., 2006	Beijing	WTS	U	8	<16	Bare soil	b/f	25.8
Su et al., 2006	Beijing	WTS	CAN	8	<16	Bare soil	b/f	2.9
Su et al., 2006	Beijing	WTS	ASN	8	<16	Bare soil	b/f	19.8
Su et al., 2006	Beijing	WTS	U	8	<16	Bare soil	b/f	17.5
Su et al., 2007	Beijing	WTS	U	8	<16	Summer Maize	b	33.1
Su et al., 2007	Beijing	WTS	U	8	<16	Summer Maize	b	23.7
Wang et al., 2012	Jiangxi	EIV	U	5.24	16-24	Late Rice	i	18.9
Wang et al., 2012	Jiangxi	EIV	U	5.24	16-24	Late Rice	b	9.2
Wu et al., 2015	Yunnan	VM	U	6.34	<16	Snow Peas	b	26.02
Wu et al., 2015	Yunnan	VM	U	6.34	<16	Snow Peas	b	29.8
Wu et al., 2015	Yunnan	VM	U	6.34	<16	Snow Peas	b	29.1
Wu et al., 2015	Yunnan	VM	U	6.34	<16	Zucchini	b	15.42
Wu et al., 2015	Yunnan	VM	U	6.34	<16	Zucchini	b	14.56
Wu et al., 2015	Yunnan	VM	U	6.34	<16	Zucchini	b	14.08
Wu et al., 2015	Yunnan	VM	U	6.34	<16	Sweet Maize	b	11
Wu et al., 2015	Yunnan	VM	U	6.34	<16	Sweet Maize	b	12.35
Wu et al., 2015	Yunnan	VM	U	6.34	<16	Sweet Maize	b	11.62
Cai et al., 2002	Henan	MT	U	8.6	<16	Maize	b/f	18
Cai et al., 2002	Henan	MT	U	8.6	<16	Maize	b	44
Cai et al., 2002	Henan	MT	U	8.6	<16	Maize	b	48
Yang et al., 2010	Henan	VM	U	8.2	<16	Winter Wheat	i	15.96
Zhu et al., 1989	Henan	MT	U	8.8	<16	Rice	i	30.1
Zhu et al., 1989	Henan	MT	ABC	8.8	<16	Rice	i	39.1
Zhao et al., 1988	Tianjin	¹⁵ N	AS	8	16-24	Bare Soil	b	18.32
Zhao et al., 1988	Tianjin	¹⁵ N	AS	8	16-24	Bare Soil	b	22.07
Zhao et al., 1988	Tianjin	¹⁵ N	AS	8	16-24	Bare Soil	b	15
Zhao et al., 1988	Tianjin	¹⁵ N	AS	8	16-24	Bare Soil	b	32
Zhao et al., 1988	Tianjin	¹⁵ N	ABC	8	16-24	Bare Soil	b	50.92
Zhao, et al. 1988	Tianjin	¹⁵ N	AS	8	16-24	Bare Soil	b	17.69
Dong et al., 2014	Liaoning	VM	U	6.13	16-24	Bare Soil	b	26.94
Wu et al., 2009	Hunan	CC	U	4.9	<16	Early Rice	i	2.2

Wu et al., 2009	Hunan	CC	U	4.9	<16	Late Rice	i	5
Wu et al., 2009	Hunan	CC	U	4.9	<16	Late Rice	b	11.3
Xi et al., 1987	Shanghai	CC	U	7.9	<16	Cabbage	i	11.12
Li et al., 1993	Shannxi	Ha	AS	7.9	<16	Bare Soil	b	34.6
Tian et al., 2007	Jiangsu	EIV	U	7.3	16-24	Rice	i	8.4
Tian et al., 2007	Jiangsu	EIV	U	7.3	16-24	Rice	b	27.4
Yu et al., 2013	Jiangsu	VM	U	6.99	16-24	Rice		19.4
Zhu et al., 1985	Jiangsu	EIV	AS	6.6	16-24	Early Rice	i	1.5
Zhu et al., 1985	Jiangsu	EIV	U	6.6	16-24	Early Rice	i	4.5
Zhu et al., 1985	Jiangsu	EIV	AS	8	16-24	Early Rice	i	10.2
Zhu et al., 1985	Jiangsu	EIV	U	8	16-24	Early Rice	i	11.3
Zhu et al., 1985	Jiangsu	EIV	ABC	5.7	<16	Late Rice	i	15.4
Zhu et al., 1985	Jiangsu	EIV	U	5.7	<16	Late Rice	i	16
Zhu et al., 1985	Jiangsu	EIV	AS	5.7	<16	Late Rice	i	10.2
Fan et al., 2005	Jiangsu	MT	U	7.15	16-24	Wheat	b	22.7
Fan et al., 2005	Jiangsu	MT	U	7.15	16-24	Wheat	b	26.2
Cai et al., 1985	Jiangsu	MT	U	5.3	16-24	Rice	b	8.8
Cai et al., 1985	Jiangsu	MT	ABC	5.3	16-24	Rice	b	19.5
Xia et al., 2010	Hubei	CAEM	U	6.3	<16	Rice	i	13.79

¹For the methods, WTS denotes wind tunnel system, EIV denotes enclosed intermittent vent method, VM denotes venting method, ¹⁵N denotes ¹⁵N stable tracer isotope, MT denotes micrometeorological technique, CC denotes closed air-flow chamber, Ha denotes Hargroue's method, and CAEM denotes continuous air flow enclosure method.

²U denotes Urea, ABC denotes ammonium bicarbonate, AS denotes ammonium sulfate, AN denotes ammonium nitrate, CAN denotes calcium ammonium nitrate, and ASN denotes ammonium sulfate-nitrate.

³Soil cation exchange capacity (CEC) in unit of cmol(+)/kg.

⁴b denotes broadcast, i denotes cultivate after broadcasting, and b/f denotes flood after broadcasting.

⁵EF in unit of %.

40

45

References

- Bouwman, A. F., Boumans, L. J. M., and Batjes, N. H.: Estimation of global NH₃ volatilization loss from synthetic fertilizers and animal manure applied to arable lands and grasslands, *Global Biogeochem. Cy.*, 16, 2, 1024, doi:10.1029/2000GB001389, 2002.
- 50 Cai, G. X., Zhu, Z. L., Zhu, Z. W., Trevitt, A. C. F., Freney, J. R., and Simpsom, J. R.: The research of nitrogen losses from ammonium bicarbonate and urea in rice paddy field *Soils*, 17, 225-256, doi:10.13758/j.cnki.tr.1985.05.001, 1985 (in Chinese).
- Cai, G. X., Chen, D. L., Ding, H., Pacholski, A., Fan, X. H., and Zhu, Z. L.: Nitrogen losses from fertilizers applied to maize, wheat and rice in the North China Plain, *Nutr. Cycl. Agroecosys.*, 63, 187-195, doi:10.1023/a:1021198724250, 2002.
- 55 Dong, Y. H., Zhang, Y. G., Sun, S. L., Jin, D., and Wang, Y. G.: Soil ammonia volatilization under different urea combined fertilization treatments, *Chinese Journal of Ecology*, 33, 2943-2949, 2014 (in Chinese with English abstract).
- Fan, X. H., Song, Y. S., Lin, D. X., Yang, L. Z., and Zhou, J. M.: Ammonia volatilization losses from urea applied to wheat on a paddy soil in Taihu Region, China, *Pedosphere*, 15, 60 59-65, 2005.
- Li, S. X., and Liu, C. Y.: Ammonia volatilization from calcareous soil I. effects of soil properties on N loss by volatilization, *Agricultural Research in the Arid Areas*, 11 Suppl., 125-129, 1993 (in Chinese with English abstract).
- 65 Liao, J.: Fertilizer application and analysis, Shanghai science and Technology Press, Shanghai, 1993 (in Chinese).
- Su, F., Ding, X. Q., Gao, Z. L., Huang, B. X., Chen, X. P., and Zhang, F. S.: Ammonia volatilization from nitrogen fertilization of winter wheat-summer maize rotation system in the North China Plain, *China Environmental Science*, 27, 409-413, 2007 (in Chinese with 70 English abstract).
- Su, F., Huang, B. X., Ding, X. Q., Gao, Z. L., Chen, X. P., Zhang, F. S., Martin, K., and Volker, R.: Ammonia volatilization of different nitrogen fertilizer types, *Soils*, 38, 682-686, 2006 (in Chinese with English abstract).
- Tian, Y. H., He, F. Y., Yin, B., and Zhu, Z. L.: Ammonia volatilization from paddy field in 75 the Taihu Lake Region as affected by N and P combination in fertilization, *Acta*

Pedologica Sinica, 44, 893-900, 2007 (in Chinese with English abstract).

Wang, C., Zhou, W., LI, Z. Z., Liu, X. M., Sun, G., Xia, W. J., Wang, X. B., and Liu, G. R.:
Effects of different nitrogen application rates on ammonia volatilization from paddy fields
under double-harvest rice system, *Plant Nutrition and Fertilizer Science*, 18, 349-358,
80 2012 (in Chinese with English abstract).

Wu, P. P., Liu, J. J., Yang, X. X., Shang, Q. Y., Zhou, Y., Xie, X. L., Sheng, Q. R., and Guo,
S. W.: Effects of different fertilization systems on ammonia volatilization from double-rice
cropping field in red soil region, *Chinese Journal of Rice Science*, 23, 85-93, 2009 (in
Chinese with English abstract).

85 Wu, T. C., Lan, Z. Q., Hu, Z. Y., Wang, H. H., and Zhu, X. Q.: Effect of different nitrogen
fertilizer amount on soil ammonia volatilization and crop yield at vegetables field in
Chaihe Basin of Dianchi Lake, *Modern Agricultural Sciences and Technology*, 205-208,
2015 (in Chinese with English abstract).

Xi, Z. B., Shi, X. Z., Huang, W. Y., Yao, Z., Humphereys, E., Freney, J. R., and Simpson, J.
90 R.: Measuring urea loss from ammonia volatilization by a micrometeorological method,
Acta Agriculturae Shanghai, 3, 47-56, 1987(in Chinese with English abstract).

Xia, W. J., Zhou, W., Liang, G. Q., Wang, X. B., Sun, J. W., Li, S. L., Hu, C., and Chen, Y.
F.: Effect of optimized nitrogen application on ammonia volatilization from paddy field
under wheat-rice rotation system, *Plant Nutrition and Fertilizer Science*, 16, 6-13, 2010 (in
95 Chinese with English abstract).

Yang, S. L., Zhu, A. N., Zhang, J. B., Chen, X. M., and Zhu, Q. G.: Ammonia volatilization
loss and its affecting factors under different amounts and ways of N application in field,
Arid Zone Research, 27, 415-421, 2010 (in Chinese with English abstract).

Yu, Y. L., Xue, L. H., and Yang, L. Z.: Ammonia volatilization from paddy fields under
100 different nitrogen schemes in Tai Lake Region, *Journal of Agro-Environment Science*, 32,
1682-1689, doi:10.11654/jaes.2013.08.028, 2013 (in Chinese with English abstract).

Zhang, W. F. and Zhang, F. S.: Report on the development of fertilizer in China, China
Agriculture University Press, 2012b (in Chinese).

Zhao, Z. D., Zhang, J. S., Ren, S. R., and Zhou, Y. M.: The Nitrogen balance of crop field soil
105 in north area of China, *Acta Pedologica Sinica*, 3, 67-72, 1988 (in Chinese with English

abstract).

Zhu, Z. L., Cai, G. X., Xu, Y. H., and Zhang, S. L.: Ammonia volatilization and its significance to the losses of fertilizer nitrogen applied to paddy soil, *Acta Pedologica Sinica*, 22, 321-328, 1985 (in Chinese with English abstract).

- 110 Zhu, Z. L., Simpson, J. R., Zhang, S. L., Cai, G. X., Chen, D. L., Freney, J. R., and Jackson, A. V.: Investigations on nitrogen losses from fertilizers applied to flooded calcareous paddy soil, *Acta Pedologica Sinica*, 26, 337-343, 1989 (in Chinese with English abstract).