

1 Assessment and economic valuation of air pollution impacts on human health over Europe
2 and the United States as calculated by a multi-model ensemble in the framework of
3 AQMEII3

4 Ulas Im^{1*}, Jørgen Brandt¹, Camilla Geels¹, Kaj Mantzius Hansen¹, Jesper Heile Christensen¹,
5 Mikael Skou Andersen¹, Efisio Solazzo², Ioannis Kioutsioukis³, Ummugulsum Alyuz⁴,
6 Alessandra Balzarini⁵, Rocio Baro⁶, Roberto Bellasio⁷, Roberto Bianconi⁷, Johannes Bieser⁸,
7 Augustin Colette⁹, Gabriele Curci^{10,11}, Aidan Farrow¹², Johannes Flemming¹³, Andrea
8 Fraser¹⁴, Pedro Jimenez-Guerrero⁶, Nutthida Kitwiroon¹⁵, Ciao-Kai Liang¹⁶, Uarporn
9 Nopmongcol¹⁷, Guido Pirovano⁵, Luca Pozzoli^{4,2}, Marje Prank^{18,19}, Rebecca Rose¹⁴, Ranjeet
10 Sokhi¹², Paolo Tuccella^{10,11}, Alper Unal⁴, Marta Garcia Vivanco^{9,20}, Jason West¹⁶, Greg
11 Yarwood¹⁷, Christian Hogrefe²¹, Stefano Galmarini²

12 ¹ Aarhus University, Department of Environmental Science, Frederiksborgvej 399, DK-4000,
13 Roskilde, Denmark.

14 ² European Commission, Joint Research Centre (JRC), Ispra (VA), Italy.

15 ³ University of Patras, Department of Physics, University Campus 26504 Rio, Patras, Greece

16 ⁴ Eurasia Institute of Earth Sciences, Istanbul Technical University, Istanbul, Turkey

17 ⁵ Ricerca sul Sistema Energetico (RSE SpA), Milan, Italy

18 ⁶ University of Murcia, Department of Physics, Physics of the Earth, Campus de Espinardo, Ed.
19 CIOyN, 30100 Murcia, Spain

20 ⁷ Enviroware srl, Concorezzo, MB, Italy

21 ⁸ Institute of Coastal Research, Chemistry Transport Modelling Group, Helmholtz-Zentrum
22 Geesthacht, Germany

23 ⁹ INERIS, Institut National de l'Environnement Industriel et des Risques, Parc Alata, 60550 Verneuil-
24 en-Halatte, France

25 ¹⁰ Dept. Physical and Chemical Sciences, University of L'Aquila, L'Aquila, Italy

26 ¹¹ Center of Excellence CETEMPS, University of L'Aquila, L'Aquila, Italy

27 ¹² Centre for Atmospheric and Instrumentation Research (CAIR), University of Hertfordshire,
28 Hatfield, UK

29 ¹³ European Centre for Medium Range Weather Forecast (ECMWF), Reading, UK

30 ¹⁴ Ricardo Energy & Environment, Gemini Building, Fermi Avenue, Harwell, Oxon, OX11 0QR, UK

31 ¹⁵ Environmental Research Group, Kings' College London, London, UK

32 ¹⁶ Department of Environmental Sciences and Engineering, University of North Carolina at Chapel
33 Hill, Chapel Hill, North Carolina, USA

34 ¹⁷ Ramboll Environ, 773 San Marin Drive, Suite 2115, Novato, CA 94998, USA

35 ¹⁸ Finnish Meteorological Institute, Atmospheric Composition Research Unit, Helsinki, Finland

36 ¹⁹ Cornell University, Department of Earth and Atmospheric Sciences, Ithaca, USA

37 ²⁰ CIEMAT. Avda. Complutense 40., 28040 Madrid, Spain

38 ²¹ Computational Exposure Division, National Exposure Research Laboratory, Office of Research and
39 Development, United States Environmental Protection Agency, Research Triangle Park, NC, USA.

40

41 *Correspondence to:* Ulas Im (ulas@envs.au.dk)

42 **Abstract**

43 The impact of air pollution on human health and the associated external costs in Europe and
44 the United States (U.S.) for the year 2010 is modelled by a multi-model ensemble of regional
45 models in the frame of the third phase of the Air Quality Modelling Evaluation International
46 Initiative (AQMEII3). The modelled surface concentrations of O₃, CO, SO₂ and PM_{2.5} are
47 used as input to the Economic Valuation of Air Pollution (EVA) system to calculate the

48 resulting health impacts and the associated external costs from each individual model. Along
49 with a base case simulation, additional runs were performed introducing 20% anthropogenic
50 emission reductions both globally and regionally in Europe, North America and East Asia, as
51 defined by the second phase of the Task Force on Hemispheric Transport of Air Pollution
52 (TF-HTAP2).

53 Health impacts estimated by using concentration inputs from different chemistry and
54 transport models (CTMs) to the EVA system can vary up to a factor of three in Europe
55 (twelve models) and the United States (three models). In Europe, the multi-model mean total
56 number of premature deaths (acute + chronic) is calculated to be 414 000 while in the U.S., it
57 is estimated to be 160 000, in agreement with previous global and regional studies. The
58 economic valuation of these health impacts are calculated to be 300 and 145 billion Euros in
59 Europe and the U.S., respectively. A subset of models that produce the smallest error
60 compared to the surface observations at each time step against an all-models mean ensemble
61 results in increase of health impacts by up to 30% in Europe, while in the U.S., the optimal
62 ensemble mean led to a decrease in the calculated health impacts by ~11%.

63 A total of 54 000 and 27 500 premature deaths can be avoided by a 20% reduction of global
64 anthropogenic emissions in Europe and the U.S., respectively. A 20% reduction of North
65 American anthropogenic emissions avoids a total premature death of ~1 000 in Europe and
66 25 000 total premature deaths in the U.S. A 20% decrease of anthropogenic emissions within
67 the European source region avoids a total premature death of 47 000 in Europe. Reducing the
68 East Asian anthropogenic emissions by 20% avoids ~2000 total premature deaths in the U.S.
69 These results show that the domestic anthropogenic emissions make the largest impacts on
70 premature death on a continental scale, while foreign sources make a minor contributing to
71 adverse impacts of air pollution.

72 **1. Introduction**

73 According to the World Health Organization (WHO), air pollution is now the world's largest
74 single environmental health risk (WHO, 2014). Around 7 million people died prematurely in
75 2012 as a result of air pollution exposure from both outdoor and indoor emission sources
76 (WHO, 2014). WHO estimates 3.7 million premature deaths in 2012 from exposure to
77 outdoor air pollution from urban and rural sources worldwide. According to the Global
78 Burden of Disease (GBD) study, exposure to ambient particulate matter pollution remains
79 among the ten leading risk factors. Air pollution is a transboundary phenomenon with global,
80 regional, national and local sources, leading to large differences in the geographical
81 distribution of human exposure. Short-term exposure to ozone (O₃) is associated with
82 respiratory morbidity and mortality (e.g. Bell et al., 2004), while long-term exposure to O₃
83 has been associated with premature respiratory mortality (Jerrett et al., 2009). Short-term
84 exposure to particulate matter (PM_{2.5}) has been associated with increases in daily mortality
85 rates from respiratory and cardiovascular causes (e.g. Pope and Dockery, 2006), while long-
86 term exposure to PM_{2.5} can have detrimental chronic health effects, including premature
87 mortality due to cardiopulmonary diseases and lung cancer (Burnett et al., 2014). The Global

88 Burden of Disease Study 2015 estimated 254 000 O₃-related and 4.2 million anthropogenic
89 PM_{2.5}-related premature deaths per year (Cohen et al., 2017).

90 Changes in emissions from one region can impact air quality over others, affecting also air
91 pollution-related health impacts due to intercontinental transport (Anenberg et al., 2014;
92 Zhang et al., 2017). In the framework of the Task Force on Hemispheric Transport of Air
93 Pollution (TF-HTAP), Anenberg et al. (2009) found that reduction of foreign ozone precursor
94 emissions can contribute to more than 50% of the deaths avoided by simultaneously reducing
95 both domestic and foreign precursor emissions. Similarly, they found that reducing emissions
96 in North America (NA) and Europe (EU) has largest impacts on ozone-related premature
97 deaths in downwind regions than within (Anenberg et al., 2009). This result agrees with
98 Duncan et al. (2008), which showed for the first time that emission reductions in NA and EU
99 have greater impacts on ozone mortality outside the source region than within. Anenberg et
100 al. (2014) estimates that 93–97 % of PM_{2.5}-related avoided deaths from reducing emissions
101 occurs within the source region while 3–7 % occur outside the source region from
102 concentrations transported between continents. In spite of the shorter lifetime of PM_{2.5}
103 compared to O₃, it was found to cause more deaths from intercontinental transport (Anenberg
104 et al., 2009; 2014). In the frame of the second phase of the Task Force on Hemispheric
105 Transport of Air Pollution (TF-HTAP2; Galmarini et al., 2017), an ensemble of global
106 chemical transport model simulations calculated that 20% emission reductions from one
107 region generally lead to more avoided deaths within the source region than outside (Liang et
108 al., 2017).

109 Recently, Lelieveld et al. (2015) used a global chemistry model and calculated that outdoor
110 air pollution led to 3.3 million premature deaths globally in 2010. They calculated that in
111 Europe and North America, 381 000 and 68 000 premature deaths occurred, respectively.
112 They have also calculated that these numbers are likely to roughly double in the year 2050
113 assuming a business-as-usual scenario. Silva et al. (2016), using the ACCMIP model
114 ensemble, calculated that the global mortality burden of ozone is estimated to markedly
115 increase from 382 000 deaths in 2000 to between 1.09 and 2.36 million in 2100. They also
116 calculated that the global mortality burden of PM_{2.5} is estimated to decrease from 1.70
117 million deaths in 2000 to between 0.95 and 1.55 million deaths in 2100. Silva et al. (2013)
118 estimated that in 2000, 470 000 premature respiratory deaths are associated globally and
119 annually with anthropogenic ozone, and 2.1 million deaths with anthropogenic PM_{2.5}-related
120 cardiopulmonary diseases (93%) and lung cancer (7%). These studies employed global
121 chemistry and transport models with coarse spatial resolution ($\geq 0.5^\circ \times 0.5^\circ$), therefore health
122 benefits from reducing local emissions were not able to be adequately captured. Higher
123 resolutions are necessary to calculate more robust estimates of health benefits from local vs.
124 non-local sources (Fenech et al., 2017). In addition, these studies calculated number of
125 premature deaths due to air pollution, however none of them addresses morbidity such as
126 number of lung cancer or asthma cases, or restricted activity days. Finally, these studies did
127 not include economic costs either. On the other hand, there are a number of regional studies
128 that calculate health impacts on finer spatial resolutions, and address morbidity. However,
129 they are mostly based on single air pollution models or do not evaluate the health benefits

130 from local vs. non-local emissions. Therefore, a comprehensive study employing multi model
131 ensemble of high spatial resolution and focusing on both mortality and morbidity from local
132 vs. non-local sources lacks in the literature.

133 In Europe, recent results show that outdoor air pollution due to O₃, CO, SO₂ and PM_{2.5} causes
134 a total number of 570 000 premature deaths in the year 2011 (Brandt et al., 2013a; 2013b).
135 The external (or indirect) costs to society related to health impacts from air pollution are
136 tremendous. OECD (2014) estimates that outdoor air pollution is costing its member
137 countries USD 1.57 trillion in 2010. Among the OECD member countries, the economic
138 valuation of air pollution in the U.S. was calculated to be ~500 billion USD and ~660 USD in
139 Europe. In the whole of Europe, the total external costs have been estimated to approx. 800
140 billion Euros in year 2011 (Brandt et al., 2013a). These societal costs have great influence on
141 the general level of welfare and especially on the distribution of welfare both within the
142 countries as air pollution levels are vastly heterogeneous both at regional and local scales and
143 between the countries as air pollution and the related health impacts are subject to long-range
144 transport. Geels et al. (2015), using two regional chemistry and transport models, estimated a
145 premature mortality of 455 000 and 320 000 in Europe (EU28 countries) for the year 2000,
146 respectively, due to O₃, CO, SO₂ and PM_{2.5}. They also estimated that climate change alone
147 leads to a small increase (15%) in the total number of O₃-related acute premature deaths in
148 Europe towards the 2080s and relatively small changes (<5%) for PM_{2.5}-related mortality.
149 They found that the combined effect of climate change and emission reductions will reduce
150 the premature mortality due to air pollution, in agreement with the results from Schucht et al.
151 (2015).

152 The U.S. Environmental Protection Agency estimated that in 2010 there were ~160 000
153 premature deaths in the U.S. due to air pollution (U.S. EPA, 2011). Fann et al. (2012)
154 calculated 130,000 - 350,000 premature deaths associated with O₃ and PM_{2.5} from the
155 anthropogenic sources in the U.S. for the year 2005. Caizzo et al. (2013) estimated 200 000
156 cases of premature death in the U.S. due to air pollution from combustion sources for the year
157 2005.

158 The health impacts of air pollution and their economic valuation are estimated based on
159 observed and/or modelled air pollutant concentrations. Observations have spatial limitations
160 particularly when assessments are needed for large regions. The impacts of air pollution on
161 health can be estimated using models, where the level of complexity can vary depending on
162 the geographical scale (global, continental, country or city), concentration input
163 (observations, model calculations, emissions) and the pollutants of interest that can vary from
164 only few (PM_{2.5} or O₃) to a whole set of all regulated pollutants. The health impact models
165 normally used may differ in the geographical coverage, spatial resolutions of the air pollution
166 model applied, complexity of described processes, the exposure-response functions (ERFs),
167 population distributions and the baseline indices (see Anenberg et al., 2015 for a review).

168 Air pollution related health impacts and associated costs can be calculated using Chemical
169 Transport Model (CTM) or with standardized source-receptor relationships characterizing the
170 dependence of ambient concentrations on emissions. (e.g. EcoSense model: ExternE, 2005,

171 TM5-FASST: Van Dingenen et al., 2014). Source-receptor relationships have the advantage
172 of reducing the computing time significantly and have therefore been extensively used in
173 systems like GAINS (Amann et al., 2011). On the other hand, full CTM simulations have the
174 advantage of better accounting for non-linear chemistry-transport processes in the
175 atmosphere.

176 CTMs are useful tools to calculate the concentrations of health-related pollutants taking into
177 account non-linearities in the chemistry and the complex interactions between meteorology
178 and chemistry. However, the CTMs include different chemical and aerosol schemes that
179 introduce differences in the representation of the atmosphere as well as differences in the
180 emissions and boundary conditions they use (Im et al., 2015a,b). These different approaches
181 are present also in the health impact estimates that use CTM results as basis for their
182 calculations. Multi-model (MM) ensembles can be useful to the extent that allow us to take
183 into consideration several model results at the same time, define the relative weight of the
184 various members in determining the mean behavior, and produce also an uncertainty
185 estimated based on the diversity of the results (Potemski and Galmarini, 2010; Riccio et al.,
186 2013; Solazzo et al., 2013).

187 The third phase of the Air Quality Modelling Evaluation International Initiative (AQMEII3)
188 project brought together fourteen European and North American modelling groups to
189 simulate the air pollution levels over the two continental areas for the year 2010 (Galmarini et
190 al., 2017). Within AQMEII3, the simulated surface concentrations of health related air
191 pollutants from each modelling group serves as input to the Economic Valuation of Air
192 Pollution (EVA) model (Brandt et al., 2013a; 2013b). This is the first study in our knowledge
193 that uses a common approach across the two continents regarding the economic valuation of
194 health impacts of air pollution, as also pointed in Andersen (2017). The EVA model is used
195 to calculate the impacts of health-related pollutants on human health over the two continents
196 as well as the associated external costs. EVA model has also been tested and validated for the
197 first time outside Europe. We adopt a multi-model ensemble (MM) approach, in which the
198 outputs of the modelling systems are statistically combined assuming equal contribution from
199 each model and used as input for the EVA model. In addition, the human health impacts (and
200 the associated costs) of reducing anthropogenic emissions, globally and regionally have been
201 calculated, allowing to quantify the trans-boundary benefits of emission reduction strategies.
202 Finally, following the conclusions of Solazzo and Galmarini (2015), the health impacts have
203 been calculated using an optimal ensemble of models, determined by error minimization .
204 This approach can assess the health impacts with reduced model bias, which we can then
205 compare with the classically derived estimates based on model averaging.

206 **2. Material and Methods**

207 **2.1. AQMEII**

208 *2.1.1. Participating Models*

209 In the framework of the AQMEII3 project, fourteen groups participated to simulate the air
210 pollution levels in Europe and North America for the year 2010. In the present study, we use

211 results from the thirteen groups that provided all health-related species (Table 1). As seen in
212 Table 1, six groups have operated the CMAQ model. The main differences among the CMAQ
213 runs reside in the number of vertical levels and horizontal spacing (Table 1) and in the
214 estimation of biogenic emissions. UK1, DE1, and US3 calculated biogenic emissions using the
215 BEIS (Biogenic Emission Inventory System version 3) model, while TR1, UK1, and UK2
216 calculated biogenic emissions through the MEGAN model (Guenther et al., 2012). Moreover,
217 DE1 does not include the dust module, while the other CMAQ instances use the inline
218 calculation (Appel et al., 2013) and TR1 uses the dust calculation previously calculated for
219 AQMEII Phase 2. Finally, all runs were carried out using CMAQ version 5.0.2 except for TR1,
220 which is based on the 4.7.1 version. The gas-phase mechanisms and the aerosol models are
221 used by each group is also presented in Table 1. More details of the model system are provided
222 in the supplementary material. The differences in the meteorological drivers and aerosol
223 modules can lead to substantial differences in modelled concentrations (Im et al., 2015b).

224 2.1.2. Emission and Boundary Conditions

225 The base-case emission inventories that are used in AQMEII for Europe and North America
226 are extensively described in Pouliot et al. (2015). For Europe, the 2009 inventory of TNO-
227 MACC anthropogenic emissions was used. In regions not covered by the emission inventory,
228 such as North Africa, five modelling systems have complemented the standard inventory with
229 the HTAPv2.2 datasets (Janssens-Maenhout et al., 2015). For the North American domain,
230 the 2008 National Emission Inventory was used as the basis for the 2010 emissions,
231 providing the inputs and datasets for processing with the SMOKE emissions processing
232 system (Mason et al., 2012). For both continents the regional scale emission inventories were
233 embedded in the global scale inventory (Janssens-Maenhout et al., 2015) used by the global-
234 scale HTAP2 modelling community so that to guarantee coherence and harmonization of the
235 information used by the regional scale modelling community. The annual totals for European
236 and North American emissions in the HTAP inventory are the same as the MACC and
237 SMOKE emissions. However, there are differences in the temporal distribution, chemical
238 speciation and the vertical distribution used in the models. The C-IFS model (Flemming et
239 al., 2015 and 2017) provided chemical boundary conditions. The C-IFS model has been
240 extensively evaluated in Flemming et al. (2015 and 2017), and in particular for North
241 America (Hogrefe et al., 2017; Huang et al., 2017). Galmarini et al. (2017) provides more
242 details on the setup of the AQMEII3 and HTAP2 projects.

243 2.1.3. Model Evaluation

244 The models' performance on simulating the surface concentrations of the health-related
245 pollutants were evaluated using Pearson's Correlation (r), normalized mean bias (NMB),
246 normalized mean gross error ($NMGE$) and root mean square error ($RMSE$) to compare the
247 modelled and observed hourly pollutant concentrations over surface measurement stations in
248 the simulation domains. The hourly modelled vs. observed pairs are averaged and compared
249 on a monthly basis. The modelled hourly concentrations were first filtered based on
250 observation availability before the averaging has been performed. The observational data
251 used in this study are the same as the dataset used in second phase of AQMEII (Im et al.,

252 2015a, b). Surface observations are provided in the Ensemble system
253 (<http://ensemble2.jrc.ec.europa.eu/public/>) that is hosted at the Joint Research Centre (JRC).
254 Observational data were originally derived from the surface air quality monitoring networks
255 operating in EU and NA. In EU, surface data were provided by the European Monitoring and
256 Evaluation Programme (EMEP, 2003; <http://www.emep.int/>) and the European Air Quality
257 Database (AirBase; <http://acm.eionet.europa.eu/databases/airbase/>). In NA observational data
258 were obtained from the NAtChem (Canadian National Atmospheric Chemistry) database and
259 from the Analysis Facility operated by Environment Canada (<http://www.ec.gc.ca/natchem/>).

260 The model evaluation has been conducted for 491 European and 626 North American stations
261 for O₃, 541 European stations and 37 North American stations for CO, 500 European station
262 and 277 North American stations for SO₂, and 568 European stations and 156 North
263 American stations for PM_{2.5}.

264 *2.1.4. Emissions Perturbations*

265 In addition to the base case simulations in AQMEII3, a number of emission perturbation
266 scenarios have been simulated (Table 1). The perturbation scenarios feature a reduction of
267 20% in the global anthropogenic emissions (GLO) as well as the HTAP2-defined regions of
268 Europe (EUR), North America (NAM) and East Asia (EAS), as explained in detail in
269 Galmarini et al. (2017) and Im et al. (2017). To prepare these scenarios, both the regional
270 models and the global C-IFS model that provides the boundary conditions to the participating
271 regional models have been operated with the reduced emissions. The global perturbation
272 scenario (GLO) reduces the global anthropogenic emissions by 20%, introducing a change in
273 the boundary conditions as well as a 20% decrease in the anthropogenic emissions used by
274 the regional models. The North American perturbation scenario (NAM) reduces the
275 anthropogenic emissions in North America by 20%, introducing a change in the boundary
276 conditions while anthropogenic emissions remain unchanged for Europe, showing the impact
277 of long-range transport while for North America, while the scenarios introduces a 20%
278 reduction of anthropogenic emissions in the HTAP-defined North American region. The
279 European perturbation scenario (EUR) reduces the anthropogenic emissions in the HTAP-
280 defined Europe domain by 20%, introducing a change in the anthropogenic emissions while
281 boundary conditions remain unchanged in the regional models, showing the contribution
282 from the domestic anthropogenic emissions only. Finally, the East Asian perturbation
283 scenario (EAS) reduces the anthropogenic emissions in East Asia by 20%, introducing a
284 change in the boundary conditions while anthropogenic emissions remain unchanged in the
285 regional models, showing the impact of long-range transport from East Asia on the NA
286 concentrations.

287 **2.2. Health Impact Assessment**

288 All modeling groups interpolate their model outputs on a common 0.25°×0.25° resolution
289 AQMEII grid predefined for Europe (30°W - 60°E, 25°N - 70°N) and North America
290 (130°W - 59.5°W, 23.5°N - 58.5°N). All the analyses performed in the present study use the
291 pollutant concentrations on these final grids. Health impacts are first calculated for each

292 individual model and then the ensemble mean, median and standard deviation are calculated
293 for each health impact. In order to be able to estimate an uncertainty in the health impacts
294 calculations, none of the models were removed from the ensemble.

295 Along with the individual health impact estimates from each model, a multi-model mean
296 dataset (MM_m , in which all the modelling systems are averaged assuming equally weighted
297 contributions) has been created for each grid cell and time step, hence creating a new model
298 set of results that have the same spatial and temporal resolution of the ensemble-contributing
299 members. In addition to this simple MM_m , an optimal MM ensemble (MM_{opt}) has been
300 generated. MM_{opt} is created following the criteria extensively discussed and tested in the
301 previous phases of the AQMEII activity (Riccio et al., 2012; Kioutsioukis et al., 2016;
302 Solazzo and Galmarini, 2016), where it was shown that there are several ways to combine the
303 ensemble members to obtain a superior model, mostly depending on the feature we wish to
304 promote (or penalize). For instance, generating an optimal ensemble that maximizes the
305 accuracy would require a minimization of the mean error or of the bias, while maximizing the
306 associativity (variability) would require maximize the correlation coefficient (standard
307 deviation). In this study, the sub-set of models whose mean minimize the mean squared error
308 (MSE) is selected as optimal (MM_{opt}). MM_m and MM_{opt} have therefore the same spatial
309 resolution with the individual models. The MSE is chosen for continuity with previous
310 AQMEII-related works. The MSE is chosen in the light of its property of being composed by
311 bias, variance and covariance types of error, thus lumping together measures of accuracy
312 (bias), variability (variance) and associativity (covariance) (Solazzo and Galmarini, 2016).
313 The minimum MSE has been calculated at the monitoring stations, where observational data
314 are available and then extended to the entire continental areas. This approximation might
315 affect remote regions away from the measurements. However, considering that for the main
316 pollutants (O_3 and $PM_{2.5}$) the network of measurements is quite dense around densely
317 populated areas (where the inputs of the MM ensemble are used for assessing the impact of
318 air pollutants on the health of the population), errors due to inaccurate model selection in
319 remote regions might be regarded as negligible (Solazzo and Galmarini, 2015). It should be
320 noted that the selection of the optimal combinations of models is affected by the model's bias
321 that might stem from processes that are common to all members of the ensemble (e.g.
322 emissions). Therefore, such a common bias does not cancel out when combining the models,
323 possibly creating a biased ensemble. Current work is being devoted to identify the optimal
324 combinations of models from which the offsetting bias is removed (Solazzo et al., 2017b).

325 **2.2.1. EVA System**

326 The EVA system (Brandt et al., 2013a, b) is based on the impact-pathway chain (e.g.
327 Friedrich and Bickel, 2001), consisting of the emissions, transport and chemical
328 transformation of air pollutants, population exposure, health impacts and the associated
329 external costs. The EVA system requires hourly gridded concentration input from a regional-
330 scale CTM as well as gridded population data, exposure-response functions (ERFs) for health
331 impacts, and economic valuations of the impacts from air pollution. A detailed description of
332 the integrated EVA model system along with the ERFs and the economic valuations used are
333 given in Brandt et al. (2013a).

334 The gridded population density data over Europe and the U.S. used in this study are presented
335 in Fig. 1. The population data over Europe are provided on a 1km spatial resolution from
336 Eurostat for the year 2011 (<http://www.efgs.info>). The U.S. population data has been
337 provided from the U.S. Census Bureau for the year 2010. The total populations used in this
338 study are roughly 532 and 307 million in Europe and the U.S., respectively. As the health
339 outcomes are age-dependent, the total population data has been broken down to a set of age
340 intervals being babies (under 9 months), children (under 15), adult (above 15), above 30, and
341 above 65. The fractions of population in these intervals for Europe is derived from the
342 EUROSTAT 2000 database, where the number of persons of each age at each grid cell was
343 aggregated into the above clusters (Brandt et al., 2011), while for the U.S. they are derived
344 from the U.S. Census Bureau for the year 2010 at 5-year intervals.

345 The EVA system can be used to assess the number of various health outcomes including
346 different morbidity outcomes as well as short-term (acute) and long-term (chronic) mortality,
347 related to exposure of O₃, CO and SO₂ (short-term) and PM_{2.5} (long-term). Furthermore,
348 impact on infant mortality in response to exposure of PM_{2.5} is calculated. The health impacts
349 are calculated using an ERF of the following form:

$$350 \quad R = \alpha \times \delta_c \times P$$

351 where R is the response (in cases, days, or episodes), c denotes the pollutant concentration, P
352 denotes the affected share of the population, and α an empirically determined constant for the
353 particular health outcome. EVA uses ERFs that are modelled as a linear function, which is a
354 reasonable approximation as showed in several studies (e.g. Pope et al., 2000; the joint World
355 Health Organization/UNECE Task Force on Health (EU, 2004; Watkiss et al., 2005)). The
356 concentration metrics used in each ERF is shown in Table 2. The sensitivity of EVA to the
357 different pollutant concentrations are further evaluated in the supplementary material and
358 depicted in Fig. S1. EVA calculates and uses the annual mean concentrations of CO, SO₂ and
359 PM_{2.5}, while for O₃, it uses the SOMO35 metric that is defined as the yearly sum of the daily
360 maximum of 8-hour running average over 35 ppb, following WHO (2013) and EEA (2017).

361 The morbidity outcomes include chronic bronchitis, restricted activity days, congestive heart
362 failure, lung cancer, respiratory and cerebrovascular hospital admissions, asthmatic children
363 (<15 years) and adults (>15 years), which includes bronchodilator use, cough, and lower
364 respiratory symptoms. The exposure-response functions are broadly in line with estimates
365 derived with detailed analysis in EU funded research (Rabl, Spadaro and Holland, 2014;
366 EEA, 2013) To figure out the total number of premature deaths from the years of life lost due
367 to PM_{2.5}, they have been converted into lost lives according to a lifetable method (explained
368 in detail in Andersen, 2017) but using the factor of 10.6, as reported by (Watkiss et al., 2005).
369 To these deaths are added the acute deaths due to O₃ and SO₂. The ERFs used, along with
370 their references, in both continents as well as the economic valuations for each health
371 outcome in Europe and the U.S., respectively, are presented in Table 2. Baseline incidence
372 rates are not assumed to be dissimilar, which is a coarse approach for morbidity. The baseline
373 rates are from Statistics Denmark
374 (<http://www.statistikbanken.dk/statbank5a/default.asp?w=1280>) and lifetables are based on

375 Denmark, which is close to the US and Eurozone average (Andersen, 2017). For a description
376 of the morbidity ERFs, see Andersen et al. (2004 and 2008). The economic valuations are
377 provided by Brandt et al. (2013a); see also EEA (2013).

378 ERF for all-cause chronic mortality due to PM_{2.5} were based on the findings of Pope et al.
379 (2002), which is the most extensive study available, following conclusions from the scientific
380 review of the Clean Air For Europe (CAFÉ) programme (Hurley et al., 2005; Krupnick et al.,
381 2005). The results from Pope et al. (2002) are further supported by Krewski et al. (2009), and
382 more recently by the latest HRAPIE project report (WHO, 2013a). Therefore, as
383 recommended by WHO (2013a), EVA uses the ERFs based on the meta-analysis of 13 cohort
384 studies as described in Hoek et al. (2013). In EVA, the number of lost life years for a Danish
385 population cohort with normal age distribution, when applying the ERF of Pope et al. (2002)
386 for all-cause mortality (relative risk, RR= 1.062 (1.040-1.083) on 95% confidence interval),
387 and the latency period indicated, sums to 1138 yr of life lost (YOLL) per 100 000 individuals
388 for an annual PM_{2.5} increase of 10 µg m⁻³ (Andersen, 2008)..EVA uses a counterfactual
389 PM_{2.5} concentration of 0 µgm⁻³ following the EEA methodology, meaning that the impacts
390 have been estimated for the full range of modelled concentrations from 0 µgm⁻³ upwards.
391 Applying a low counterfactual concentration can underestimate health impacts at low
392 concentrations if the relationship is linear or close to linear (Anenberg et al., 2016). However,
393 it is important to note that uncertainty in the health impact results may increase at low
394 concentrations due to sparse epidemiological data. Assuming linearity at very low
395 concentrations may distort the true health impacts of air pollution in relatively clean
396 atmospheres (Anenberg et al., 2016).

397 It has been shown that O₃ concentrations above the level of 35 ppb involve an acute mortality
398 increase, presumably for weaker and elderly individuals. EVA applies the ERFs selected in
399 CAFE for post-natal death (age group 1–12 months) and acute death related to O₃ (Hurley et
400 al., 2005). WHO (2013a) also recommends the use of the daily maximum of 8-hour mean O₃
401 concentrations for the calculation of the acute mortality due to O₃. There are also studies
402 showing that SO₂ is associated with acute mortality, and EVA adopts the ERF identified in
403 the APHENA study – Air Pollution and Health: A European Approach (Katsouyanni et al.,
404 1997).

405 Chronic exposure to PM_{2.5} is also associated with morbidity, such as lung cancer. EVA
406 employs the specific ERF (RR = 1.08 per 10 µg m⁻³ PM_{2.5} increase) for lung cancer indicated
407 in Pope et al. (2002). Bronchitis has been shown to increase with chronic exposure to PM_{2.5}
408 and we apply an ERF (RR = 1.007) for new cases of bronchitis based on the AHSMOG study
409 (involving non-smoking Seventh-Day Adventists; Abbey et al., 1999), which is the same
410 epidemiological study as in CAFE (Abbey, 1995; Hurley et al., 2005). The ExternE crude
411 incidence rate was chosen as a background rate (ExternE, 1999), which is in agreement with
412 a Norwegian study, rather than the pan-European estimates used in CAFE (Eagan et al.,
413 2002). Restricted activity days (RADs) comprise two types of responses to exposure: so-
414 called minor restricted activity days as well as work-loss days (Ostro, 1987). This distinction
415 enables accounting for the different costs associated with days of reduced well-being and
416 actual sick days. It is assumed that 40% of RADs are work-loss days based on Ostro (1987).

417 The background rate and incidence are derived from ExternE (1999). Hospital admissions are
418 deducted to avoid any double counting. Hospital admissions and health effects for asthmatics
419 (here corresponding to the three responses bronchodilator use, cough and lower respiratory
420 symptoms) are also based on ExternE (1999).

421 Table 2 lists the specific valuation estimates applied in the modelling of the economic
422 valuation of mortality and morbidity effects. A principal value of EUR 1.5 million was
423 applied for preventing an acute death, following expert panel advice (EC 2001). For the
424 valuation of a life year, the results from a survey relating specifically to air pollution risk
425 reductions were applied (Alberini et al., 2006), implying a value of EUR 57.500 per year of
426 life lost (YOLL). With the more conservative metric of estimating lost life years, rather than
427 'full' statistical lives, there is no adjustment for age. This is due to the fact that government
428 agencies in Europe, including the European Commission, apply a methodology for costing of
429 air pollution that is based on accounting for lost life years, rather than for entire statistical
430 lives as is customary in USA. While the average traffic victim, for instance, is mid-aged and
431 likely to lose about 35-40 years of life expectancy, pollution victims are believed to suffer
432 significantly smaller losses of years (EAHEAP, 1999:64; Friedrich and Bickel, 2001). To
433 avoid overstating the benefits of air pollution control, these are treated as proportional to the
434 number of life years lost. Most of the excess mortality is due to chronic exposure to air
435 pollution over many years and the life year metric is based on the number of lost life years in
436 a statistical cohort. Following the guidelines of the Organisation for Economic Co-operation
437 and Development (OECD, 2006), the predicted acute deaths, mainly from O₃, are valued
438 here with the adjusted value for preventing a fatality (VSL, Value of a Statistical Life). The
439 life tables are obtained from European data and are applied to the U.S. as the average life
440 expectancy in the U.S. is similar to that in Europe, and close to the OECD average (OECD,
441 2016). The willingness to pay for reductions in risk obviously differs across income levels.
442 However, in the case of air pollution costs, adjustment according to per capita income
443 differences among different states is not regarded as appropriate, because long-range
444 transport implies that emissions from one state will affect numerous other states and their
445 citizens. The valuations are thus adjusted with regional purchasing power parities (PPP) of
446 EU27 and USA.

447
448 Cost-benefit analysis in the U.S. related to air pollution proceeds from a standard approach,
449 where abatement measures preventing premature mortality are considered according to the
450 number of statistical fatalities avoided, which are appreciated according to the value of VSL
451 (presently USD 7.4 million). In contrast, and following recommendations from the UK
452 working group on Economic Appraisal of the Health Effects of Air Pollution (EAHEAP,
453 1999), focus in EU has been on the possible changes in average life expectancy resulting
454 from air pollution. In EU, the specific number of life years lost as a result of changes in air
455 pollution exposures are estimated based on lifetable methodology, and monetized with Value-
456 Of-Life-Year (VOLY) unit estimates (Holland et al. 1999; Leksell and Rabl 2001). The
457 theoretical basis is a life-time consumption model according to which the preferences for risk
458 reduction will reflect expected utility of consumption for remaining life years (Hammit
459 2007; OECD 2006:204). The much lower VSL values customary in Europe (presently €2.2

460 million) add decisively to the differences, as VOLY is deducted from this value. By using a
461 common valuation framework according the EU approach we allow for direct comparisons of
462 the monetary results. It follows from OECD recommendations (2012) to correct with PPP
463 when doing such benefit transfer. The unit values have been indexed to 2013 prices as
464 indicated in Table 2.

465

466 3. Results

467 3.1. Model Evaluation

468 Observed and simulated hourly surface O₃, CO, SO₂ and daily PM_{2.5}, which are species used
469 in the EVA model to calculate the health impacts, over Europe and North America for the
470 entire 2010 were compared in order to evaluate each model's performance. The statistical
471 parameters to evaluate the models and their equations are provided in the supplementary
472 material. For a more thorough evaluation of models and species, see Solazzo et al. (2017a).
473 The results of this comparison are presented in Table S1 for EU and NA, along with the
474 multi-model mean and median values. The monthly time series plots of observed and
475 simulated health-related pollutants are also presented in Figs. 2 and 3. The monthly means are
476 calculated using the hourly pairs of observed and modelled concentrations at each station.
477 The results show that over Europe, the temporal variability of all gaseous pollutants is well
478 captured by all models with correlation coefficients (*r*) higher than 0.70 in general. The
479 normalized mean biases (*NMB*) in simulated O₃ levels are generally below 10% with few
480 exceptions up to -35%. CO levels are underestimated by up to 45%, while the majority of the
481 models underestimated SO₂ levels by up to 68%, while some models overestimated SO₂ by
482 up to 49%. PM_{2.5} levels are underestimated by 19% to 63%. Over Europe, the median of the
483 ensemble performs better than the mean in terms of model bias (*NMB*) for O₃ (by 52%),
484 while for CO, SO₂ and PM_{2.5}, the mean performs slightly better than the median (Table S1).

485 We have further evaluated the models' performance on simulating the annual mean pollutant
486 levels over individual measurements stations and plotted the geographical distribution of the
487 bias. Fig. 4 presents the multi model mean geographical distribution of bias over Europe,
488 while Fig. S2-S5 for O₃, CO, SO₂ and PM_{2.5}, respectively. O₃ levels over central to western
489 Europe are overestimated by up to ~10 µgm⁻³, while over eastern Europe, O₃ levels are
490 underestimated by up to ~10 µgm⁻³ (Fig. 4a) Over southern Europe, overestimations are
491 larger (10-20 µgm⁻³). The geographical pattern is similar among the models with slight
492 differences ($\pm 10 \mu\text{gm}^{-3}$) in the bias (Fig. S2). CO levels are underestimated over all stations
493 by up to 600 µgm⁻³ except for few stations where CO levels are overestimated by up to 100
494 µgm⁻³ (Fig. 4b). All models underestimated CO levels over the majority of the stations (Fig.
495 S3). SO₂ levels are slightly overestimated over central and southern Europe (Fig. 4c). There
496 are also underestimation over few stations with no specific geographical pattern. Similar to
497 CO, all models underestimated SO₂ levels over the majority of the stations (Fig. S4). Finally,
498 PM_{2.5} levels are underestimated by up to 10 µgm⁻³ over most of Europe (Fig. 4d), with larger
499 underestimations over the eastern Europe up to 30 µgm⁻³.

500 Over North America, the hourly O₃ variation is well captured by all models (Table S1), with
501 DK1 having slightly lower *r* coefficient compared to the other models and largest *NMB* (Fig.
502 3a). The hourly variation of CO and SO₂ levels are simulated with relatively lower *r* values
503 (Figs. 3b, c), with SO₂ levels having the highest underestimations. The PM_{2.5} levels are
504 underestimated by ~15% except for the DE1 model, having a large underestimation of 63%
505 (Table S1). As DE1 and US3 use the same SMOKE emissions and CTM, the large difference
506 in PM_{2.5} concentrations can be partly due to the differences in horizontal and vertical
507 resolutions in the model setups, as can also be seen in the differences in the CO
508 concentrations. There are also differences in the aerosol modules and components that each
509 model simulates. For example, DE1 uses an older version of the secondary organic aerosol
510 (SOA) module, producing ~3 μgm⁻³ less SOA, which can explain ~20% of the bias over
511 North America. Over the North American domain, the median outscore the mean for O₃ (by
512 35%), CO (by 52%) and PM_{2.5} (by 29%) while for SO₂, the median produces 26% higher
513 *NMB* compared to the mean. DK1 model simulates a much higher bias for O₃ and SO₂
514 compared to other models in the North American domain, while DE1 has the largest bias for
515 CO and PM_{2.5}.

516 O₃ levels are generally overestimated by the MM mean over the eastern U.S. by up to 15 ppb,
517 while over the western U.S. there are also overestimations by up to 10 ppb (Fig. 5a). As seen
518 in Fig. S6, all three models have very similar performance over the U.S., with DK1
519 simulating a slightly lower underestimation and a higher overestimation compared to DE1
520 and US3. DE1 and DK1 have very similar spatial pattern in terms of CO bias, in particular
521 over the eastern coast of the U.S. (Fig. S7). CO levels are underestimated by ~100 ppb over
522 majority of the stations, especially over the eastern U.S., while there are much larger
523 underestimation over the western U.S. by up to 1000 ppb (Fig. 5b). SO₂ levels are
524 underestimated by up to 5 ppb over the majority of the stations in the U.S., with few
525 overestimations of up to 5 ppb (Fig. 5c). DE1 and DK1 have very similar spatial distribution
526 of bias, while US3 has slightly more overestimations (Fig. S8) Finally, PM_{2.5} levels are
527 underestimated over majority of the stations by up to 6 μgm⁻³, with few overestimations by 2-
528 4 μgm⁻³ (Fig. 5d). DE1 has the largest underestimations compared to DK1 and US3 (Fig. S9).

529 Table S1 shows that the ensemble median performs slightly better than the ensemble mean
530 for all pollutants over both continents regarding the bias and error, while the difference on *r*
531 is rather small. Over the European stations, the median has improved results over the mean
532 by up to 14% for *r* and up to 9% for the *RMSE*. The improvements in *r* over the U.S. are
533 much smaller compared to Europe (up to ~4%), while the *RMSE* is improved by up to 27%,
534 except for SO₂ where the median has 14% higher *RMSE* than the mean.

535 3.2. Health outcomes and their economic valuation in Europe

536 The different health outcomes calculated by each model in Europe as well as their multi
537 model mean and median are presented in Table S2. Table 3 presents the mean of the
538 individual model estimates as *MM_{mi}*. Standard deviations calculated from the individual
539 model estimates are presented along with the *MM_{mi}* in the text. The health impact estimates
540 vary significantly between different models. The different estimates obtained are found to

541 vary up to a factor of three. Among the different health outcomes, the individual models
542 simulated the number of congestive heart failure (CHF) cases to be between 19 000 to 41 000
543 (mean of all individual models, MM_{mi} , $31\ 000 \pm 6\ 500$). The number of lung cancer cases due
544 to air pollution are calculated to be between 30 000 to 78 000 (mean of all individual models,
545 MM_{mi} , $55\ 000 \pm 14\ 000$). Finally, the total (acute + chronic) number of premature death due
546 to air pollution is calculated to be 230 000 to 570 000 (mean of all individual models, MM_{mi} ,
547 $414\ 000 \pm 100\ 000$). The health impacts calculated as the median of individual models differ
548 slightly ($\sim \pm 1\%$) from those calculated as the mean of individual models (Table S2) due to the
549 slight differences in the model bias (NMB) and error ($NMGE$ and $RMSE$) between the mean
550 and the median performance statistics of the models.

551 In addition to averaging the health estimates from individual models (MM_{mi}), we have also
552 produced a multi-model mean concentration data (MM_m) by taking the average of
553 concentrations of each species calculated by all models at each grid cell and hour, and fed it
554 to the EVA model. We have calculated the number of premature death cases in Europe as
555 410 000 (Table 3) using MM_m . Difference between the health impacts calculated using MM_m
556 data from the mean of all individual model (MM_{mi}) estimates is smaller than 1%. The number
557 of premature death cases in Europe as calculated as the average of all models in the multi
558 model ensemble, MM_{mi} , due to exposure to O_3 is $12\ 000 \pm 6\ 500$, while the cases due to
559 exposure to $PM_{2.5}$ is calculated to be $390\ 000 \pm 100\ 000$ [180 000 – 550 000]. The O_3 -related
560 mortality well agrees with Liang et al. (2017) that used the multi-model mean of the HTAP2
561 global model ensemble, which calculated an O_3 -related mortality of 12 800 [600 - 28 100].
562 The multi-model mean (MM_{mi}) $PM_{2.5}$ -related mortality in the present study is much higher
563 than that from the HTAP2 study (195 500 [4 400 – 454 800]). The results also agree with the
564 most recent EEA findings (EEA, 2015), which calculated a total premature death of 419 000
565 die to O_3 and $PM_{2.5}$ in the EU-28 countries. There is also agreement with Geels et al. (2015)
566 that calculated 388 000 premature death cases in Europe for the year 2000. This difference
567 can be attributed to the number of mortality cases as calculated by the individual models,
568 where the HTAP2 ensemble calculates a much lower minimum while the higher ends from
569 the two ensembles well agree.

570 The differences between the health outcomes calculated by the HTAP2 and AQMEII
571 ensembles arise firstly from the differences in the concentrations fields due to the differences
572 in models, in particular spatial resolutions as well as the gas and aerosols treatments in
573 different models, but also the differences in calculating the health impacts from these
574 concentrations fields. EVA calculates the acute premature death due to O_3 by using the
575 SOMO35 metric. On the other hand, in HTAP2 O_3 -related premature death is calculated by
576 using the 6-month seasonal average of daily 1-h maximum O_3 concentrations. Both groups
577 use the annual mean $PM_{2.5}$ to calculate the $PM_{2.5}$ -related premature death. In addition to O_3
578 and $PM_{2.5}$, EVA also takes into account the health impacts from CO and SO_2 , which is
579 missing in the HTAP2 calculations.

580 Among all models, DE1 model calculated the lowest health impacts for most health
581 outcomes, which can be attributed to the largest underestimation of $PM_{2.5}$ levels ($NMB=-$
582 63%: Table S2) due to lower spatial resolution of the model that dilutes the pollution in the

583 urban areas, where most of the population lives. The number of premature deaths calculated
584 by this study is in agreement with previous studies for Europe using the EVA system (Brandt
585 et al., 2013a; Geels et al., 2015). Recently, EEA (2015) estimated that air pollution is
586 responsible for more than 430 000 premature deaths in Europe, which is in good agreement
587 with the present study.

588 Fig. 6a. presents the geographical distribution of the number of premature death in Europe in
589 2010. The figure shows that the numbers of cases are strongly correlated to the population
590 density (Fig. 1a), with the largest numbers seen in the Benelux and the Po Valley regions that
591 are characterized as the pollution hot spots in Europe as well as in megacities such as
592 London, Paris, Berlin and Athens.

593 The economic valuation of the air pollution-associated health impacts calculated by the
594 different models along with their mean and median are presented in Table 4. A total cost of
595 196 to 451 billion Euros (MM mean cost of 300 ± 70 billion Euros) was estimated over
596 Europe (EU28). Results show that 5% [1% - 11%] of the total costs is due to exposure to O₃,
597 while 89% [80% - 96%] is due to exposure to PM_{2.5}. Brandt et al. (2013a) calculated a total
598 external cost of 678 billion Euros for the year 2011 for Europe, larger than the estimates of
599 this study, which can be explained by the differences in the simulation year and the emissions
600 used in the models as well as the countries included in the two studies (the previous study
601 includes e.g. Russia).

602 3.3. Health outcomes and their economic valuation in the U.S.

603 The different health outcomes calculated by each model for the U.S. as well as their mean
604 and median are presented in Table S2. The variability among the models (~3) is similar to
605 that in Europe. The number of congestive heart failure cases in the U.S. as calculated as the
606 average of all models in the ensemble (MM_{mi}) is calculated to be 13 000 [7 000 – 18 000],
607 while the lung cancer cases due to air pollution are calculated to be 22 000 [9 000 – 31 000].
608 Finally, the number of premature deaths due to air pollution is calculated to be $165\,000 \pm$
609 $75\,000$, where $25\,000 \pm 6\,000$ cases are calculated due to exposure to O₃ and $140\,000 \pm 72$
610 000 cases due to exposure to PM_{2.5}. The MM_m dataset leads to a number of premature death
611 of 149 000 that is 6% smaller than the average estimate from individual models (MM_{mi}). Due
612 to the large reduction of *NMB* by the median compared to the mean of individual models
613 (Table S1), the multi-model health impacts calculated as the median of health impacts from
614 individual models are ~13% higher than the health impacts calculated from the MM_{mi} . The
615 O₃- and PM_{2.5} mortality cases as calculated by the AQMEII and HTAP2 model ensembles
616 reasonably agree. Liang et al. (2017) calculated an O₃-related mortality of 14 700 [900 –
617 30 400] and a PM_{2.5}-related mortality of 78 600 [4 500 – 162 600]. These results are in very
618 good agreement with the U.S. EPA (2011) estimates of number of premature death cases of
619 160 000 in year 2010 and with Caizzo et al. (2013), who calculated 200 000 premature death
620 cases from combustion sources in the U.S. Among all models, DE1 model calculated the
621 lowest health impacts for most health outcomes, which can be attributed to the largest
622 underestimation of PM_{2.5} levels ($NMB=-63\%$: Table S2).

623 The premature death cases in North America are mostly concentrated over the New York
624 area, as well as in hot spots over Chicago, Detroit, Houston Los Angeles and San Francisco
625 (Fig. 6b). The figure shows that the number of cases is following the pattern of the population
626 density. The economic valuation of the air pollution-associated health impacts calculated by
627 the different models in the U.S. are shown in Table 4. As seen in the table, a total cost of
628 ~145 billion Euros is calculated. Results show that ~22% of the total costs is due to exposure
629 to O₃ while ~78% is due to exposure to PM_{2.5}. The major health impacts in terms of their
630 external costs are slightly different in North America compared to Europe.

631 3.4. Health impacts and their economic valuation through optimal reduced ensemble subset

632 The effect of pollution concentrations (EVA input) on health impacts (EVA output) is
633 investigated in order to estimate the contribution of each air pollutant in the EVA system to
634 health impacts over different concentration levels. The technical details are provided in the
635 supplement.

636 Results show that for the particular input (gridded air pollutant concentrations from
637 individual model)-output (each health outcome) configuration, the PM_{2.5} drives the variability
638 of the different health impact and that at least 81% of the variation of the health impacts are
639 explained by sole variations in the pollutants (i.e. without interactions: Table S3). Table S1
640 also shows that the most important contribution to the health impacts is from PM_{2.5}, followed
641 by CO and O₃ (with much smaller influence though). The impact of perturbing PM_{2.5} by a
642 fixed fraction of its standard deviation on the health impact is roughly double compared to
643 CO and O₃.

644 We have run the EVA system over an all-models mean (MM_m) dataset and an optimal
645 reduced ensemble dataset (MM_{opt}) calculated for each of the pollutants in the two domains in
646 order to see how and whether an optimal reduced ensemble changes the assessment of the
647 health impacts compared to an all- models ensemble mean. Table 5 shows some sensible
648 error reduction, although the temporal and spatial averages mask the effective improvement
649 in accuracy from MM_m to MM_{opt} . In Europe, the optimal reduced ensemble decreases the
650 RMSE by up to 24%, while in NA, the error reduction is much larger (4% to up to 147%). On
651 a seasonal basis, MM_{opt} reduces RMSE in PM_{2.5} over Europe by 23% in winter while smaller
652 decreases are achieved in other seasons (~10%). Regarding O₃, improvement is 16%-22%,
653 with the largest improvement in spring. In NA, the improvement in winter RMSE in PM_{2.5} is
654 smallest (~2%) while larger improvements are achieved in other seasons (~7% - ~9%). For
655 O₃, the largest RMSE reduction in NA is achieved for the summer period by 14%.

656 The analysis of the aggregated health indices data for Europe (Table S1) shows that EVA
657 indices rely principally on the PM_{2.5} levels and then the CO and O₃ values. Therefore, the
658 relative improvement of the indices with the optimal ensemble should be proportional to the
659 relative improvement in PM_{2.5}, CO and O₃. The proportionality rate for each pollutant is
660 given in Table S3, assuming all pollutants are varied (from MM_m to MM_{opt}) away from their
661 mean by the same fraction of their variance. As seen in the Table 3, from MM_m to MM_{opt} , the
662 health indices increase by up to 30% in Europe. This increase is due to a 27% increase in the

663 domain mean PM_{2.5} levels when the optimal reduced ensemble is used, as well a slight
664 increase in O₃ by ~1%. The number of premature deaths in Europe increase from 410 000 to
665 524 000 (28%), resulting in a much higher estimate compared to previous mortality studies.
666 On the contrary, in the U.S., the mean PM_{2.5} and O₃ levels decrease from 2.94 µg m⁻³ to 2.62
667 µgm⁻³ (~11%) and 18.7 ppb to 18.4 ppb (~2%), respectively. In response, the health indices
668 decrease by ~11% (Table 3). The number of premature death cases in NA decrease from
669 149 000 to 133 000.

670 3.5. Impact of anthropogenic emissions on the health impacts and their economic valuation

671 The impacts of emission perturbations on the different health outcomes over Europe and the
672 U.S. as calculated by the individual models are presented in Tables S4-S6. Table 6 shows the
673 impacts of the different emission perturbations on the premature death cases in Europe and
674 the U.S as calculated by a subset of models that simulated the base case and all three
675 perturbation scenarios (*MM_c*). Results show that in Europe, the 20% reduction in the global
676 anthropogenic emissions leads to ~17% domain-mean reduction in all the health outcomes,
677 with a geographical variability as seen in Fig. 6c. The figure shows that the larger changes in
678 mortality is calculated in the central and northern parts of Europe (15-20% decreases), while
679 the changes are smaller in the Mediterranean region (5-10%), highlighting the non-linearity
680 of the response to emission reductions. However, it should be noted that global models or
681 coarse-resolution regional models (as in this study) cannot capture the urban features and
682 pollution levels and thus, non-linearities should be addressed further using fine spatial
683 resolutions or urban models. The models vary slightly simulating the response to the 20%
684 reduction in global emissions, estimating decreases of ~11% to 20%. The number of
685 premature deaths decreased on average by ~50 000, ranging from -39 000 (DK1) to -103 000
686 (IT1). This number is in good agreement with the ~45 000 premature death calculated by the
687 HTAP2 global models (Liang et al., 2017). The *MM_c* ensemble calculated a 15% and 17%
688 decrease in the O₃- and PM_{2.5}-related premature death cases, respectively, in response to the
689 GLO scenario. This decrease in the global anthropogenic emissions leads to an estimated
690 decrease of 56 ± 18 billion Euros in associated costs in Europe (Table 6).

691 As seen in Table 8, a 20% reduction of anthropogenic emissions in the EUR region, as
692 defined in HTAP2, avoids 47 000 premature death, while a 20% reduction of the
693 anthropogenic emissions in the NAM region leads to a much smaller decrease of premature
694 deaths in Europe (~1 000). These improvements in the number of premature deaths are in
695 agreement with a recent HTAP2 global study that calculated reductions of ~34 000 and
696 ~1 000 for the EUR and NAM scenarios, respectively (Liang et al., 2017) and with Anenberg
697 et al. (2009 and 2014), which totals to a sum of avoided premature deaths being ~39 000 and
698 1 800 as calculated by the MM mean. Both the global and regional models agree that the
699 largest impacts of reducing emissions with respect to premature deaths come from emission
700 within the source region, while foreign sources contribute much less to improvements in
701 avoiding adverse impacts of air pollution. The decreases in health impacts in EUR and NAM
702 scenarios corresponds to decreases in the associated costs by -47 ± 16 billion Euros and -1.4
703 ± 0.4 billion Euros, respectively. This is consistent with results in Brandt et al. (2012), where
704 a contribution of ~1% to PM_{2.5} concentrations in Europe is originating from the NAM region.

705 The 20% reduction in global anthropogenic emissions leads to 18% reduction in the health
706 outcomes (Table 8) in the U.S., with a geographical variability in the response. Fig. 6d shows
707 that the largest decreases in mortality is calculated for the western coast of the U.S. (~20%)
708 and slightly lower response in the central and eastern parts of the U.S. (15-20%). The number
709 of premature death cases, as calculated by the mean of all individual models decreases from
710 $\sim 160\,000 \pm 70\,000$ to $\sim 130\,000 \pm 60\,000$, avoiding 24 ± 10 billion Euros (Table 6) in
711 external costs, also in agreement with the ensemble of HTAP2 global models ($\sim 23\,000$) The
712 O₃-related premature death cases decreased by 42% while the PM_{2.5}-related cases decreased
713 by 18%.

714 A 20% reduction of the North American emissions avoids $\sim 25\,000 \pm 12\,000$ premature
715 deaths (-16%), suggesting that $\sim 80\%$ of avoided premature deaths are achieved by reductions
716 within the source region while 20% ($\sim 5\,000$ premature deaths) is from foreign sources. This
717 number is also in good agreement with Liang et al. (2017) that estimated a reduction of
718 premature deaths of $\sim 20\,000$ due to O₃ and PM_{2.5} in the United States due to an emission
719 reduction of 20% within the region itself, using the ensemble mean of the HTAP2 global
720 models. These results are much larger than the number of avoided premature death of
721 $\sim 11\,000$ as calculated by the sum of Anenberg et al. (2009 and 2104). The corresponding
722 benefit is calculated to be 21 ± 9 billion Euros in the NAM scenario. According to results
723 from the EAS scenario, among these 5 000 avoided cases that are attributed to the foreign
724 emission sources, $1\,900 \pm 2\,000$ premature deaths can be avoided by a 20% reduction of the
725 East Asian emissions, avoiding 2.5 ± 3 billion Euros. Our number of avoided premature
726 deaths due to the EAS scenario is much higher than 580 avoided premature deaths calculated
727 by Liang et al. (2017) and 380 avoided cases as calculated by Anenberg et al. (2009 and
728 2014).

729 **Conclusions**

730 The impact of air pollution on human health and their economic valuation for the society
731 across Europe and the United States is modelled by a multi-model ensemble of regional
732 models from the AQMEII3 project. All regional models used boundary conditions from the
733 C-IFS model, and emissions from either the MACC inventory in Europe or the EPA
734 inventory for the North America, or the global inventory from HTAP. Sensitivity analysis on
735 the dependence of models on different sets of boundary conditions has not been conducted so
736 far but large deviations from the current results in terms of health impacts are not expected.
737 The modelled surface concentrations by each individual model are used as input to the EVA
738 system to calculate the resulting health impacts and the associated external costs from O₃,
739 CO, SO₂ and PM_{2.5}. Along with a base case simulation for the year 2010, some groups
740 performed additional simulations, introducing 20% emission reductions both globally and
741 regionally in Europe, North America and East Asia.

742 The base case simulation of each model is evaluated with available surface observations in
743 Europe and North America. Results show large variability among models, especially for
744 PM_{2.5}, where models underestimate by $\sim 20\%$ - $\sim 60\%$, introducing a large uncertainty in the
745 health impact estimates as PM_{2.5} is the main driver for health impacts. The differences in the

746 models are largely due to differences in the spatial and vertical resolutions, meteorological
747 inputs, inclusion of natural emissions, dust in particular, as well as missing or underestimated
748 SOA mass, which is critical for the PM_{2.5} mass. As shown in the supplementary material, the
749 CTMs diverge a lot on the representation of particles and their size distribution, SOA
750 formation, as well as the inclusion of natural sources. As the anthropogenic emissions are
751 harmonized in the models, they represent a minor uncertainty in terms of model-to-model
752 variation. However, differences in the treatment of the temporal, vertical and chemical
753 distributions of the particulate and volatile organic species have an influence in the model
754 calculations and therefore lead to model-to-model variations.

755 The variability of health impacts among the models can be up to a factor of three in Europe
756 (twelve models) and the U.S. (three models), among the different health impacts. The multi-
757 model mean total number of premature death is calculated to be 414 000 in Europe and
758 160 000 in the U.S., where PM_{2.5} contributes by more than 90%. These numbers agree well
759 with previous global and regional studies for premature deaths due to air pollution. In order to
760 reduce the uncertainty coming from each model, an optimal ensemble set is produced, that is,
761 the subset of models that produce the smallest error compared to the surface observations at
762 each time step. The optimum ensemble results in an increase of health impacts by up to 30%
763 in Europe and a decrease by ~11% in the United States. These differences clearly
764 demonstrate the importance of the use of optimal-reduced multi-model ensembles over
765 traditional all model-mean ensembles, both in terms of scientific results, but also in policy
766 applications.

767 Finally, the role of domestic versus foreign emission sources on the related health impacts is
768 investigated using the emission perturbation scenarios. A global reduction of anthropogenic
769 emissions by 20% decreases the health impacts by 17%, while the reduction of foreign
770 emissions decreases the health impacts by less than 1%. The decrease of emissions within the
771 source region decreases the health impacts by 16%. These results show that the largest
772 impacts of reducing emissions with respect to the premature death come from emissions
773 within the source region, while foreign sources contributing to much less improvements in
774 avoiding adverse impacts of air pollution.

775 **Outlook**

776 Currently health assessments of airborne particles are carried out under the assumption that
777 all fine fraction particles affect health to a similar degree independent of origin, age and
778 chemical composition of the particles. A 2013 report from WHO concludes that the
779 cardiovascular effects of ambient PM_{2.5} are greatly influenced, if not dominated, by their
780 transition metal contents (WHO, 2013b). It is known that trace metals and traffic markers are
781 highly associated with daily mortality (Lippmann, 2014). Even low concentrations of trace
782 metals can be influential on health related responses.

783 Regarding ambient concentrations of PM and the exposure-response functions (ERFs), there
784 is a rich set of studies providing information on total PM mass. However, only few studies
785 focus on individual particulate species, mainly black carbon and carbonaceous particles. In

786 addition to PM, studies on human populations have not been able to isolate potential effects
787 of NO₂, because of its complex link to PM and O₃. The WHO REVIHAAP review from 2013
788 concludes that health assessments based on PM_{2.5} ERFs will be most inclusive (WHO,
789 2013b). In addition, the ERFs are based on urban background measurements, introducing
790 uncertainties regarding non-urban areas or high pollution areas as e.g. street canyons. Current
791 state-of-the-art health impact estimates, in particular on regional to global scales, assume a
792 correlation with exposure to outdoor air pollution, while in reality, exposure is dynamic and
793 depends on the behavior of the individual. In addition, differences in age groups, gender,
794 ethnicity and behavior should be considered in the future studies. There are also uncertainties
795 originating from the representations of the aerosols in the atmospheric models used in the
796 calculation of pollutant concentrations as well as the emissions. Further developments in the
797 aerosol modules, such as the representation of organic aerosols and windblown and
798 suspended dust, are need in order to achieve mass closure of PM to get robust estimates of
799 health impacts. In addition, new findings show that O₃ has also chronic health impacts in
800 addition to its acute impacts (WHO, 2013a; Turner et al., 2016).

801 Due to above reasons, there is a large knowledge gap regarding the health impacts of
802 particles. There are a number of ongoing projects trying to identify the health impacts from
803 individual particle components and produce individual ERFs for these components.
804 NordicWelfAir project (<http://projects.au.dk/nordicwelfair/>) aims to investigate the potential
805 causal impact of individual chemical air pollutants as well as mixtures of air pollutants on
806 health outcomes. In pursuing this aim, the project uses the unique Nordic population-based
807 registers allowing linkage between historical residential address, air pollutants over decades
808 and later health outcomes. By linking the exposure to health outcomes, new exposure-
809 response relationships can be determined of health effects for different population groups
810 (e.g. age, education, ethnicity, gender, lifestyle, and working life vs. retirement conditions)
811 related to air pollution for the individual chemical air pollutants. In addition, the high
812 resolution simulations conducted will enable us to have a better understanding of non-
813 linearities between the emissions, health impacts, and their economic valuation.

814 **ACKNOWLEDGEMENTS**

815 We gratefully acknowledge the contribution of various groups to the third air Quality Model
816 Evaluation international Initiative (AQMEII) activity. Joint Research Center Ispra/Institute
817 for Environment and Sustainability provided its ENSEMBLE system for model output
818 harmonization and analyses and evaluation. Although this work has been reviewed and
819 approved for publication by the US Environmental Protection Agency, it does not necessarily
820 reflect the views and policies of the agency. Aarhus University gratefully acknowledges the
821 NordicWelfAir project funded by the NordForsk's Nordic Programme on Health and Welfare
822 (grant agreement no. 75007), the REEEM project funded by the H2020-LCE Research and
823 Innovation Action (grant agreement no.: 691739), and the Danish Centre for Environment
824 and Energy (AU-DCE). University of L'Aquila thanks the EuroMediterranean Center for
825 Climate Research (CMCC) for providing the computational resources. RSE contribution to
826 this work has been financed by the research fund for the Italian Electrical System under the
827 contract agreement between RSE S.p.A. and the Ministry of Economic Development –

828 General Directorate for Nuclear Energy, Renewable Energy and Energy Efficiency in
829 compliance with the decree of 8 March 2006.

830

831 REFERENCES

832 Abbey, D.E., Lebowitz, M.D., Mills, P.K., Petersen, F.F., Beeson, W.L. Burchette, R.J, 1995.
833 Long-term ambient concentrations of particulates and oxidants and development of chronic
834 disease in a cohort of non-smoking California residents. *Inhalation Toxicology* 7, 19-34.

835 Abbey, D.E., Nishino, N., Mcdonnell, W.F., Burchette, R.J., Knutsen, S.F., Lawrence Beeson,
836 W., Yang, J.X., 1999. Long-term inhalable particles and other air pollutants related to
837 mortality in nonsmokers. *Am. J. Respir. Crit. Care Med.*, 159, 373–382.

838 Alberini, A., Hunt, A. and Markandya, A., 2006, Willingness to pay to reduce mortality risks:
839 Evidence from a three-country contingent valuation study. *Environmental and Resource*
840 *Economics*, 33, 251–264.

841 Amann, M., Bertok, I., Borcken-Kleefeld, J., Cofala, J., Heyes, C., Höglund-Isaksson, L.,
842 Klimont, Z., Nguyen, B., Posch, M., Rafaj, P., Sandler, R., 2011. Cost-effective control of air
843 quality and greenhouse gases in Europe: modeling and policy applications. *Environmental*
844 *Modelling & Software*, 26 (12), 1489–1501.

845 Andersen, M.S., 2017. Co-benefits of climate mitigation: Counting statistical lives or life-
846 years? *Ecological Indicators*, 79, 11-18.

847 Andersen, M.S., Frohn, L.M., Jensen, S.S., Nielsen, J.S., Sørensen, P.B., Hertel, O., Brandt,
848 J., Christensen, J.H., 2014. Sundhedseffekter af luftforurening – beregningspriser, Faglig
849 rapport fra DMU, nr. 507
850 (http://www.dmu.dk/1_viden/2_Publikationer/3_fagrapporter/rapporter/FR507.PDF).

851 Andersen, M.S., Frohn, L.M., Nielsen, J.S., Nielsen, M., Jensen, S.S., Christensen, J.H.,
852 Brandt, J., 2008. A Non-linear Eulerian Approach for Assessment of Health-cost
853 Externalities of Air Pollution. Proceedings of the European Association of Environmental and
854 Resource Economists 16th Annual Conference, Gothenburg, Sweden, 25–28 June 2008, 23
855 pp.

856 Anderson, H.R., Ponce de Leon, A., Bland, J.M., Bower, J.S., Strachan, D.P., 1996. Air
857 Pollution and daily mortality in London: 1987-92. *British Medical Journal*, 312, 665-669.

858 Anenberg, S. C., A. Belova, J. Brandt, N. Fann, S. Greco, S. Guttikunda, M.-E. Heroux, F.
859 Hurley, M. Krzyzanowski, S. Medina, B. Miller, K. Pandey, J. Roos, R. Van Dingenen, 2015.
860 Survey of ambient air pollution health risk assessment tools. *Risk Analysis*. DOI:
861 10.1111/risa.12540.

862 Anenberg, S.C., West, J.J., Yu, H., Chin, M., Schulz, M., Bergmann, D., Bey, I., Bian, H.,
863 Diehl, T., Fiore, A., Hess, P., Marmer, E., Montanaro, V., Park, R., Shindell, D., Takemura,

864 T., Dentener, F., 2014. Impacts of intercontinental transport of anthropogenic fine particulate
865 matter on human mortality. *Air Quality, Atmosphere & Health*, 7, (3), 369-379,
866 doi:10.1007/s11869-014-0248-9.

867 Anenberg, S.C., West, J.J., Fiore, A.M., Jaffe, D.A., Prather, M.J., Bregmann, D., Cuvelier,
868 K., Dentener, F.J., Duncan, B.N., Gauss, M., Hess, P., Jonson, J.E., Lupu, A., MacKenzie,
869 I.A., Marmor, E., Park, R.J., Sanderson, M.G., Schultz, M., Shindell, D.T., Szopa, S.,
870 Vivanco, M.G., Wild, O., Zeng, G., 2009. Intercontinental impacts of ozone air pollution on
871 human mortality. *Environ Science and Technology*, 43,6482–6487.

872 Bell, M.L., McDermott, A., Zeger, S.L., Samet, J.M., Dominici, F., 2004. Ozone and short-
873 term mortality in 95 US urban communities, 1987–2000. *Journal of American Medical*
874 *Association*, 292, 2372–2378.

875 Brandt, J., J. D. Silver, L. M. Frohn, C. Geels, A. Gross, A. B. Hansen, K. M. Hansen, G. B.
876 Hedegaard, C. A. Skjøth, H. Villadsen, A. Zare, and J. H. Christensen, 2012. An integrated
877 model study for Europe and North America using the Danish Eulerian Hemispheric Model
878 with focus on intercontinental transport. *Atmospheric Environment*, Volume 53, June 2012,
879 pp. 156-176, doi:10.1016/j.atmosenv.2012.01.011

880 Brandt, J., Silver, J. D., Christensen, J. H., Andersen, M. S., Bønløkke, J. H., Sigsgaard, T.,
881 Geels, C., Gross, A., Hansen, A. B., Hansen, K. M., Hedegaard, G. B., Kaas, E., and Frohn,
882 L. M.: Contribution from the ten major emission sectors in Europe and Denmark to the
883 health-cost externalities of air pollution using the EVA model system – an integrated
884 modelling approach. *Atmospheric Chemistry and Physics*, 13, 7725–7746, doi:10.5194/acp-
885 13-7725-2013, 2013a.

886 Brandt, J., Silver, J. D., Christensen, J. H., Andersen, M. S., Bønløkke, J. H., Sigsgaard, T.,
887 Geels, C., Gross, A., Hansen, A. B., Hansen, K. M., Hedegaard, G. B., Kaas, E., and Frohn,
888 L. M.: Assessment of past, present and future health-cost externalities of air pollution in
889 Europe and the contribution from international ship traffic using the EVA model system.
890 *Atmospheric Chemistry and Physics*, 13, 7747-7764, doi:10.5194/acp-13-7747-2013, 2013b.

891 Brandt, J., Silver, J.D., Frohn, L.M., Christensen, J.H., Andersen, M.S., Bønløkke, J.H.,
892 Sigsgaard, T., Geels, C., Gross, A., Hansen, A.B., Hansen, K.M., Hedegaard, G.B., Kaas, E.,
893 2011. Assessment of Health-Cost Externalities of Air Pollution at the National Level using
894 the EVA Model System, CEEH Scientific Report No 3, pp. 96.
895 (www.ceeh.dk/CEEH_Reports/Report_3)

896 Burnett, R. T., Arden Pope, C., Ezzati, M., Olives, C., Lim, S. S., Mehta, S., Shin, H. H.,
897 Singh, G., Hubbell, B., Brauer, M., Ross Anderson, H., Smith, K. R., Balme, J. R., Bruce, N.
898 G., Kan, H., Laden, F., Prüss-Ustün, A., Turner, M. C., Gapstur, S. M., Diver, W. R., Cohen,
899 A., 2014. An integrated risk function for estimating the global burden of disease attributable
900 to ambient fine particulate matter exposure. *Environmental Health Perspectives*, 122, 397–
901 403, doi:10.1289/ehp.1307049.

902 Caiazzo, F., Ashok, A., Waitz, I., Yim, S.H.L., Barrett, S.R.H., 2013. Air pollution and early
903 deaths in the United States. Part I: Quantifying the impact of major sectors in 2005.
904 *Atmospheric Environment*, 79,198–208, doi:10.1016/j.atmosenv.2013.05.081.

905 Cohen, A.J., Brauer, M., Burnett, R., Anderson, H.R., Frostad, J., Estep, K., Balakrishnan, K.,
906 Brunekreef, B., Dandona, L., Dandona, R., Feigin, V., Freedman, G., Hubbell, B., Jobling,
907 A., Kan, H., Knibbs, L., Liu, Y., Martin, R., Morawska, L., Pope III, C.A., Shin, H., Straif,
908 K., Shaddick, G., Thomas, M., van Dingenen, R., van Donkelaar, A., Vos, T., Murray, C.J.L.,
909 Forouzanfar, M.H., 2017. Estimates and 25-year trends of the global burden of disease
910 attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases
911 Study 2015. *The Lancet*, 389, 10082, 1907–1918. [http://dx.doi.org/10.1016/S0140-
912 6736\(17\)30505-6](http://dx.doi.org/10.1016/S0140-6736(17)30505-6).

913 Dab, W., Medina, S., Quénel, P., le Moullec, Y., le Tertre, A., Thelot, B., Monteil, C.,
914 Lameloise, P., Pirard, P., Momas, I., Ferry, R., Festy, B., 1996. Short Term Respiratory
915 Health Effects of Ambient Air Pollution: Results of the APHEA Project in Paris. *Journal of
916 Epidemiology and Community Health*, 50 (suppl 1), S42-S46.

917 Duncan, B.N., West, J.J., Yoshida, Y., Fiore, A.M., Ziemke, J.R., 2008. The influence of
918 European pollution on the air quality in the Near East and northern Africa. *Atmospheric
919 Chemistry and Physics*, 8, 2267–2283.

920 Dusseldorp, A., Kruize, H., Brunekreef, B., Hofschreuder, P., de Meer, G., Oudvorst, A.B.
921 van, 1995. Associations of PM10 and Airborne Iron with Respiratory Health of Adults Living
922 Near a Steel Factory. *American Journal of Critical Care Medicine*, 152, 1932-1939.

923 Eagan, T.M.L., Bakke, P.S., Eide, G.E., Gulsvik, A., 2002. Incidence of asthma and
924 respiratory symptoms by sex, age and smoking in a community study. *Eur. Respir. J.*, 19,
925 599–605.

926 European Commission (EC) 2001, Recommended interim values for the value of preventing
927 a fatality in DGEEnvironment Cost Benefit analysis, Bruxelles:
928 (http://ec.europa.eu/environment/enveco/others/pdf/recommended_interim_values.pdf),
929 accessed 28 Sept 2016.

930 EEA, 2015. Air quality in Europe, Technical report 5/2015, Copenhagen: European
931 Environment Agency.

932 EEA, 2013. Road user charges for heavy goods vehicles: Tables with external costs of air
933 pollution, Technical report 1/2013, Copenhagen: European Environment Agency.

934 EU 2004: Modelling and assessment of the health impact of particulate matter and ozone.
935 Economic commission for Europe, Executive body for the convention on long-range
936 transboundary air pollution, Working group on effects, twenty-third session, Geneva, 1–3
937 September 2004.

938 ExternE, 2005. Externalities of Energy Methodology 2005 update, European Commission,
939 Directorate-General for Research Sustainable Energy Systems. Brussels, www.externe.info.

940 ExternE: ExternE – Externalities of Energy: Vol. 7 Methodology 1998 update, European
941 Commission, Brussels, www.externe.info (last access: 2 March 2013), 1999.

942 Fann, N., Lamson, A.D., Anenberg, S.C., Wesson, K., Risley, D., Hubbell, B., 2012.
943 Estimating the national public health Burden associated with exposure to ambient PM_{2.5} and
944 ozone. *Risk Anal.* 32, 81-95.

945 Fenech, S., Doherty, R. M., Heaviside, C., Vardoulakis, S., Macintyre, H. L., and O'Connor,
946 F. M.: The influence of model spatial resolution on simulated ozone and fine particulate
947 matter: implications for health impact assessments, *Atmos. Chem. Phys. Discuss.*,
948 <https://doi.org/10.5194/acp-2017-1074>, in review, 2017.

949 Flemming, J., Benedetti, A., Inness, A., Engelen, R. J., Jones, L., Huijnen, V., Remy, S.,
950 Parrington, M., Suttie, M., Bozzo, A., Peuch, V.-H., Akritidis, D., and Katragkou, E.: The
951 CAMS interim Reanalysis of Carbon Monoxide, Ozone and Aerosol for 2003–2015, *Atmos.*
952 *Chem. Phys.*, 17, 1945-1983, <https://doi.org/10.5194/acp-17-1945-2017>, 2017.

953 Flemming, J., Huijnen, V., Arteta, J., Bechtold, P., Beljaars, A., Blechschmidt, A.-M.,
954 Diamantakis, M., Engelen, R. J., Gaudel, A., Inness, A., Jones, L., Josse, B., Katragkou, E.,
955 Marecal, V., Peuch, V.-H., Richter, A., Schultz, M. G., Stein, O., and Tsikerdekis, A., 2015.
956 Tropospheric chemistry in the Integrated Forecasting System of ECMWF. *Geoscientific*
957 *Model Development*, 8, 975-1003, doi:10.5194/gmd-8-975-2015.

958 Friedrich, R. and Bickel, P., 2001, *Environmental External Costs of Transport*, München:
959 Springer.

960 Galmarini, S., Koffi, B., Solazzo, E., Keating, T., Hogrefe, C., Schulz, M., Benedictow, A.,
961 Griesfeller, J. J., Janssens-Maenhout, G., Carmichael, G., Fu, J., and Dentener, F., 2017.
962 Technical note: Coordination and harmonization of the multi-scale, multi-model activities
963 HTAP2, AQMEII3, and MICS-Asia3: simulations, emission inventories, boundary
964 conditions, and model output formats. *Atmospheric Chemistry and Physics*, 17, 1543-1555,
965 doi:10.5194/acp-17-1543-2017.

966 Geels, C., C. Andersson, O. Hänninen, A. S. Lansø, P. Schwarze and J. Brandt, 2015. Future
967 Premature Mortality due to Air Pollution in Europe – Sensitivity to Changes in Climate,
968 Anthropogenic Emissions, Population and Building stock. *International Journal of*
969 *Environmental Research Public Health*, 12, 2837-2869.

970 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C., 2006.
971 Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of
972 Gases and Aerosols from Nature). *Atmos. Chem. Phys.*, 6, 3181–3210, doi:10.5194/acp-6-
973 3181-2006.

974 Hogrefe, C., Liu, P., Pouliot, G., Mathur, R., Roselle, S., Flemming, J., Lin, M., and Park, R.
975 J.: Impacts of Different Characterizations of Large-Scale Background on Simulated
976 Regional-Scale Ozone Over the Continental United States, *Atmos. Chem. Phys. Discuss.*,
977 <https://doi.org/10.5194/acp-2017-676>, in review, 2017.

978 Huang, M., Carmichael, G. R., Pierce, R. B., Jo, D. S., Park, R. J., Flemming, J., Emmons, L.
979 K., Bowman, K. W., Henze, D. K., Davila, Y., Sudo, K., Jonson, J. E., Tronstad Lund, M.,
980 Janssens-Maenhout, G., Dentener, F. J., Keating, T. J., Oetjen, H., and Payne, V. H.: Impact
981 of intercontinental pollution transport on North American ozone air pollution: an HTAP
982 phase 2 multi-model study, *Atmos. Chem. Phys.*, 17, 5721-5750, <https://doi.org/10.5194/acp-17-5721-2017>, 2017.

984 Hurley, F., Hunt, A., Cowie, H., Holland, Miller, B., Pye, S., Watkiss, P., 2005. Development
985 of Methodology for the CBA of the Clean Air For Europe (CAFE) Programme, Volume 2:
986 Health Impact Assessment, Report for European Commission DG Environment.

987 Im, U., Geels, C., Hansen, K.M., Christensen, J.H., Brandt, J., Solazzo, E., Alyuz, U.,
988 Balzarini, A., Baro, R., Bellasio, R., Bianconi, R., Bieser, J., Colette, A., Curci, G., Farrow,
989 A., Flemming, J., Fraser, A., Jimenez-Guerrero, P., Kitwiroon, N., Pirovano, G., Pozzoli, L.,
990 Prank, M., Rose, R., Sokhi, R., Tuccella, P., Unal, A., Vivanco, M.G., Yardwood, G.,
991 Hogrefe, C., Galmarini, S., 2017. Impacts of emission perturbations on multi-model
992 simulations of major air pollutants over Europe and North America in frame of AQMEII3, In
993 Preparation for Atmospheric Chemistry and Physics.

994 Im, U., Bianconi, R., Solazzo, E., Kioutsioukis, I., Badia, A., Balzarini, A., Baro, R., Bellasio,
995 R., Brunner, D., Chemel, C., Curci, G., Denier van der Gon, H., Flemming, J., Forkel,
996 R., Giordano, L., Jimenez-Guerrero, P., Hirtl, M., Hodzic, A., Honzak, L., Jorba, O., Knote,
997 C., Makar, P. A., Manders-Groot, A., Neal, L., Pérez, J. L., Pirovano, G., Pouliot, G., San
998 Jose, R., Savage, N., Schroder, W., Sokhi, R. S., Syrakov, D., Torian, A., Tuccella, P., Wang,
999 K., Werhahn, J., Wolke, R., Zabkar, R., Zhang, Y., Zhang, J., Hogrefe, C., and Galmarini, S.:
1000 Evaluation of operational online coupled regional air quality models over Europe and North
1001 America in the context of AQMEII phase 2, Part II: particulate matter, *Atmos. Environ.*, 115,
1002 421–441, 2015a.

1003 Im, U., Bianconi, R., Solazzo, E., Kioutsioukis, I., Badia, A., Balzarini, A., Baro, R., Bellasio,
1004 R., Brunner, D., Chemel, C., Curci, G., Flemming, J., Forkel, R., Giordano, L., Jimenez-
1005 Guerrero, P., Hirtl, M., Hodzic, A., Honzak, L., Jorba, O., Knote, C., Kuenen, J. J. P., Makar,
1006 P. A., Manders-Groot, A., Neal, L., Pérez, J. L., Pirovano, G., Pouliot, G., San Jose, R.,
1007 Savage, N., Schroder, W., Sokhi, R. S., Syrakov, D., Torian, A., Tuccella, P., Werhahn,
1008 J., Wolke, R., Yahya, K., Zabkar, R., Zhang, Y., Zhang, J., Hogrefe, C., and Galmarini, S.:
1009 Evaluation of operational online-coupled regional air quality models over Europe and
1010 North America in the context of AQMEII phase 2, Part I: ozone, *Atmos. Environ.*, 115, 404–
1011 420, 2015b.

1012 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Dentener, F., Muntean, M., Pouliot, G.,
1013 Keating, T., Zhang, Q., Kurokawa, J., Wankmüller, R., Denier van der Gon, H., Kuenen, J. J.

1014 P., Klimont, Z., Frost, G., Darras, S., Koffi, B., and Li, M., 2015. HTAP_v2.2: a mosaic of
1015 regional and global emission grid maps for 2008 and 2010 to study hemispheric transport of
1016 air pollution. *Atmospheric Chemistry and Physics*, 15, 11411–11432, doi:10.5194/acp-15-
1017 11411-2015.

1018 Jerrett, M., Burnett, R.T., Pope, C.A., Ito, K., Thurston, G., Krewski, D., Shi, Y.L., Calle, E.,
1019 Thun, M., 2009. Long-term ozone exposure and mortality. *New England Journal of*
1020 *Medicine*, 360, 1085–95

1021 Katsouyanni, K., Touloumi, G., Spix, C., Schwartz, J., Balducci, F., Medina, S., Rossi, G.,
1022 Wojtyniak, B., Sunyer, J., Bacharova, L., Schouten, J. P., Ponka, A., Anderson, H. R., 1997.
1023 Short-term effects of ambient sulphur dioxide and particulate matter on mortality in 12
1024 European cities: results from time series data from the APHEA project. *Air Pollution and*
1025 *Health: a European Approach. British Med. J.*, 314, 1658–1663.

1026 Kioutsioukis, I., Im, U., Solazzo, E., Bianconi, R., Badia, A., Balzarini, A., Baró, R.,
1027 Bellasio, R., Brunner, D., Chemel, C., Curci, G., van der Gon, H. D., Flemming, J., Forkel,
1028 R., Giordano, L., Jiménez-Guerrero, P., Hirtl, M., Jorba, O., Manders-Groot, A., Neal, L.,
1029 Pérez, J. L., Pirovano, G., San Jose, R., Savage, N., Schroder, W., Sokhi, R. S., Syrakov, D.,
1030 Tuccella, P., Werhahn, J., Wolke, R., Hogrefe, C., Galmarini, S., 2016. Insights into the
1031 deterministic skill of air quality ensembles from the analysis of AQMEII data. *Atmospheric*
1032 *Chemistry and Physics*, 16, 15629–15652, doi:10.5194/acp-16-15629-2016.

1033 Krewski, D., Jerrett, M., Burnett, R.T., Ma, R., Hughes, E., Shi, Y., Turner, M.C., Arden
1034 Pope III, C., Thurston, G., Calle, E.E., Thun, M.J., 2009. Extended Follow-Up and Spatial
1035 Analysis of the American Cancer Society Study Linking Particulate Air Pollution and
1036 Mortality. *Health Effects Institute Research Report*, 140, 1–154.

1037 Krupnick, A., Ostro, B., Bull, K., 2005. Peer review of the methodology of cost-benefit
1038 analysis of the clean air for Europe programme [http://www.cafe-cba.org/reports-on-](http://www.cafe-cba.org/reports-on-developing-the-cba-framework/)
1039 [developing-the-cba-framework/](http://www.cafe-cba.org/reports-on-developing-the-cba-framework/).

1040 Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., Pozzer, A., 2015. The contribution of
1041 outdoor air pollution sources to premature mortality on a global scale. *Nature*, 25, 367-371.

1042 Liang, C., Silva, R.A., West, J.J., Emmons, L., Jonson, J.E., Bian, H., Pan, X., Chin, M.,
1043 Henze, D., Lund, M.T., Sudo, K., Sekiya, T., Takemura, T., Flemming, J., Park, R., Lin, M.,
1044 Pierce, R.B., Lenzen, A., Kucsera, T., Folberth, G., 2017. Multi-model estimates of
1045 premature human mortality due to intercontinental transport of air pollution. *Atmospheric*
1046 *Chemistry and Physics*, In preparation.

1047 Lippmann, P., 2014. Toxicological and epidemiological studies of cardiovascular effects of
1048 ambient air fine particulate matter (PM_{2.5}) and its chemical components: Coherence and
1049 public health implications. *Critical Reviews in Toxicology*, 44(4), 299-347.

1050 Mason, R., Zubrow, A., Eyth, A., 2007. Technical Support Document (TSD) Preparation of
1051 Emissions Inventories for the Version 5.0, 2007 Emissions Modeling Platform, available at:

1052 <https://www.epa.gov/air-emissions-modeling/2007-version-50-technical-support-document>,
1053 last access: 24 May 2017.

1054 OECD (2014). *The Cost of Air Pollution: Health Impacts of Road Transport*. OECD
1055 Publishing, Paris. <http://dx.doi.org/10.1787/9789264210448-en>

1056 OECD, 2006, *Cost-benefit analysis and the environment: recent developments*, Paris,
1057 Organisation for Economic Co-operation and Development.

1058 Ostro, B.D., 1987. Air Pollution and Morbidity Revisited: A Specification Test. *Journal of*
1059 *Environmental Economics and management*, 14, 87-98.

1060 Pope, C.A., Dockery, D.W., 2006. Health effects of fine particulate air pollution: lines that
1061 connect. *Journal of the Air and Waste Management Association*, 56, 709-742.

1062 Pope, C.A., Burnett, R.T., Thun, M.J., Calle, E.E., Krewski, D., Ito, K., Thurston, G.D.,
1063 2002. Lung cancer, cardiopulmonary mortality and long-term exposure to fine particulate air
1064 pollution. *Journal of American Medical Association*, 287 (9), 1132-1141.

1065 Pope, C.A., 2000. Particulate matter-mortality exposure-response relations and threshold.
1066 *Am. J. Epidemiol.*, 152, 407–412.

1067 Pope, C.A. Thun, M.J., Namboodiri, M.M., Dockery, D.W., Evans, J.S., Speizer, F.E., Heath
1068 Jr, C.W., 1995. Particulate air pollution as a predictor of mortality in a prospective study of
1069 US adults. *American Journal of Respiratory and Critical Care Medicine*, 151, 669-674.

1070 Pope, C.A., Dockery, D.W., 1992. Acute Health Effects of PM10 Pollution on Symptomatic
1071 and Asymptomatic Children. *The American Review of Respiratory Disease*, 145, 1123-1126.

1072 Potempski, S., Galmarini, S., 2009. Est modus in rebus: analytical properties of multi-model
1073 ensembles. *Atmos. Chem. Phys.*, 9, 9471-9489, doi:10.5194/acp-9-9471-2009.

1074 Pouliot, G., Denier van der Gon, H. A. C., Kuenen, J., Zhang, J., Moran, M. D., Makar, P. A.,
1075 2015. Analysis of the emission inventories and model-ready emission datasets of Europe and
1076 North America for phase 2 of the AQMEII project. *Atmospheric Environment*, 115, 345–360.

1077 Rabl, A., Spadaro, J.V., Holland, M., 2014. *How Much Is Clean Air Worth? Calculating the*
1078 *Benefits of Pollution Control*. Cambridge University Press, ISBN: 9781107337831.

1079 Riccio, A., Ciaramella, A., Giunta, G., Galmarini, S., Solazzo, E., Potempski, S., 2012. On
1080 the systematic reduction of data complexity in multimodel atmospheric dispersion ensemble
1081 modeling. *Journal of Geophysical Research*, 117, D05314.
1082 dx.doi.org/10.1029/2011JD016503.

1083 Roemer, W., Hoek, G., Brunekreef, B., 1993. Effect of Ambient Winter Air Pollution on
1084 Respiratory Health of Children with Chronic Respiratory Symptoms. *The American Review*
1085 *of Respiratory disease*, 147, 118-124.

1086 Schucht, S., Colette, A., Rao, S., Holland, M., Schopp, W., Kolp, P., Klimont, Z., Bessagnet,
1087 B., Szopa, S., Vautard, P., Brignon, J.-M., Rouil, L., 2015. Moving towards ambitious
1088 climate policies: Monetised health benefits from improved air quality could offset mitigation
1089 costs in Europe. *Environmental Science & Policy*, 50, 252-269.

1090 Schwartz, J., Morris, R., 1995. Air Pollution and Hospital Admissions for Cardiovascular
1091 Disease in Detroit, Michigan. *American Journal of Epidemiology*, 142 (1). 23-35.

1092 Silva, R. A., West, J. J., Lamarque, J.-F., Shindell, D. T., Collins, W. J., Dalsoren, S.,
1093 Faluvegi, G., Folberth, G., Horowitz, L. W., Nagashima, T., Naik, V., Rumbold, S. T., Sudo,
1094 K., Takemura, T., Bergmann, D., Cameron-Smith, P., Cionni, I., Doherty, R. M., Eyring, V.,
1095 Josse, B., MacKenzie, I. A., Plummer, D., Righi, M., Stevenson, D. S., Strode, S., Szopa, S.,
1096 Zengast, G., 2016. The effect of future ambient air pollution on human premature mortality to
1097 2100 using output from the ACCMIP model ensemble. *Atmospheric Chemistry and Physics*,
1098 16, 9847-9862, doi:10.5194/acp-16-9847-2016.

1099 Silva, R. A., West, J. J., Zhang, Y., Anenberg, S. C., Lamarque, J.-F., Shindell, D. T.,
1100 Collins, W. J., Dalsoren, S., Faluvegi, G., Folberth, G., Horowitz, L. W., Nagashima, T., Naik,
1101 V., Rumbold, S., Skeie, R., Sudo, K., Takemura, T., Bergmann, D., Cameron-Smith, P.,
1102 Cionni, I., Doherty, R. M., Eyring, V., Josse, B., MacKenzie, I. A., Plummer, D., Righi, M.,
1103 Stevenson, D. S., Strode, S., Szopa, S., Zeng, G., 2013. Global premature mortality due to
1104 anthropogenic outdoor air pollution and the contribution of past climate change.
1105 *Environmental Research Letters*, 8, 034005, doi:10.1088/1748-9326/8/3/034005.

1106 Solazzo, E., Van Dingenen, R., Riccio, A., Galmarini, S., 2017b. The role of multi-model
1107 ensembles in assessing the air quality impact on crop yields and mortality. In preparation for
1108 submission to ACP.

1109 Solazzo, E., Bianconi, R., Hogrefe, C., Curci, G., Tuccella, P., Alyuz, U., Balzarini, A., Baró,
1110 R., Bellasio, R., Bieser, J., Brandt, J., Christensen, J. H., Colette, A., Francis, X., Fraser, A.,
1111 Vivanco, M. G., Jiménez-Guerrero, P., Im, U., Manders, A., Nopmongkol, U., Kitwiroon, N.,
1112 Pirovano, G., Pozzoli, L., Prank, M., Sokhi, R. S., Unal, A., Yarwood, G., Galmarini, S.,
1113 2017a. Evaluation and error apportionment of an ensemble of atmospheric chemistry
1114 transport modeling systems: multivariable temporal and spatial breakdown. *Atmospheric
1115 Chemistry and Physics*, 17, 3001-3054, doi:10.5194/acp-17-3001-2017.

1116 Solazzo, E., Galmarini, S., 2016. Error Apportionment for atmospheric chemistry transport
1117 models – a new approach to model evaluation. *Atmospheric Chemistry and Physics* 16, 6263-
1118 6283.

1119 Solazzo, E., Galmarini, S., 2015. A science-based use of ensembles of opportunities for
1120 assessment and scenario studies. *Atmospheric Chemistry and Physics*, 15, 2535-2544,
1121 doi:10.5194/acp-15-2535-2015.

1122 Solazzo, E., Riccio, A., Kioutsioukis, I., Galmarini, S., 2013. Pauci ex tanto numero: reduce
1123 redundancy in multi-model ensemble. *Atmospheric Chemistry and Physics*, 13, 8315–8333.

- 1124 Touloumi, G., Samoli, E., Katsuyanni, K., 1996. Daily mortality and "winter type" air
1125 pollution in Athens, Greece - a time series analysis within the APHEA project. *Journal of*
1126 *Epidemiology and Community Health*, 50 (suppl 1), S47 - S51
- 1127 Turner, M. 2016. Long-Term Ozone Exposure and Mortality in a Large Prospective Study.
1128 *Am. J. Respir. Crit. Care Med.* 193:1134–1142; doi: 10.1164/rccm.201508-1633OC.
- 1129 U.S. EPA, 2011. The Benefits and Costs of the Clean Air Act: 1990 to 2020. Final Report of
1130 U.S. Environmental Protection Agency Office of Air and Radiation, pp. 5-10.
- 1131 Van Dingenen, R., Leitao, J., Dentener, F., 2014. A multi-metric global source-receptor
1132 model for integrated impact assessment of climate and air quality policy scenarios. *European*
1133 *Geophysical Union General Assembly 2014*.
- 1134 Watkiss P., Pye S., Holland M., 2005. Cafe CBA: Baseline Analysis 2000 to 2020. Service
1135 Contract for Carrying out Cost-Benefit Analysis of Air Quality Related Issues, in Particular
1136 in the Clean Air for Europe (Cafe) Programme. 2005.. Available online:
1137 [http://ec.europa.eu/environment/archives/cafe/activities/pdf/cba_baseline_results2000_2020.](http://ec.europa.eu/environment/archives/cafe/activities/pdf/cba_baseline_results2000_2020.pdf)
1138 [pdf](http://ec.europa.eu/environment/archives/cafe/activities/pdf/cba_baseline_results2000_2020.pdf), accessed on 24 May 2017.
- 1139 Woodruff, T.J., Grillo, J., Schoendorf, K.C., 1997. The relationship between selected causes
1140 of postneonatal infant mortality and particulate air pollution in the United States.
1141 *Environmental Health Perspectives*, 105, 608-612.
- 1142 Wordley, I., Walters, S., Ayres J.G., 1997. Short term variations in hospital admissions and
1143 mortality and particulate air pollution. *Journal of Occupational Environmental Medicine*, 54,
1144 108-116.
- 1145 World Health Organization (WHO) 2014.
1146 <http://www.who.int/mediacentre/news/releases/2014/air-pollution/en/>
- 1147 World Health Organization (WHO) 2013a. Health risks of air pollution in Europe – HRAPIE:
1148 Recommendations of concentration-response functions for cost-benefit analysis of particulate
1149 matter, ozone and nitrogen dioxide.
- 1150 World Health Organization (WHO), 2013b. Review of evidence on health aspects of air
1151 pollution (REVIHAAP). WHO Technical Report.
- 1152 Zhang, Q., Jiang, X., Tong, D., Davis, S.J., Zhao, H., Geng, G., Feng, T., Zheng, B., Lu, Z.,
1153 Streets, D.G., Ni, R., Brauer, M., van Donkelaar, A., Martin, R.V., Huo, H., Liu, Z., Pan, D.,
1154 Kan, H., Yan, Y., Lin, J., He, K., Guan, D., 2017. Transboundary health impacts of
1155 transported global air pollution and international trade. *Nature*, 543, 705-709,
1156 doi:10.1038/nature21712.

Table 1. Key features (meteorological/chemistry and transport models, emissions, horizontal and vertical grids) of the regional models participating to the AQMEII3 health impact study and the perturbation scenarios they performed.

Group Code	Model	Emissions	Horizontal Resolution	Vertical Resolution	Gas Phase	Aerosol Model	Europe				North America			
							BASE	GLO	NAM	EUR	BASE	GLO	EAS	NAM
DE1	COSMO-CLM/CMAQ	HTAP	24 km × 24 km	30 layers, 50 hPa	CB5-TUCL	3 modes	×	×	×	×	×	×	×	×
DK1	WRF/DEHM	HTAP	50 km × 50 km	29 layers, 100 hPa	Brandt et al. (2012)	2 modes	×	×	×	×	×	×	×	×
ES1	WRF/CHEM	MACC	23 km × 23 km	33 layers, 50 hPa	RADM2	3 modes, MADE/SORGAM	×		×					
FI1	ECMWF/SILAM	MACC	0.25° × 0.25°	12 layers, 13 km	CB4	1-5 bins, VBS	×	×	×	×				
FRES1	ECMWF/CHIMERE	HTAP	0.25° × 0.25°	9 layers, 50 hPa	MELCHIOR2	8 bins	×	×	×	×				
IT1	WRF/CHEM	MACC	23 km × 23 km	33 layers, 50 hPa	RACM-ESRL	3 modes, MADE/VBS	×	×		×				
IT2	WRF/CAMx	MACC	23 km × 23 km	14 layers, 8 km	CB5	3 modes	×	×						
NL1	LOTOS/EUROS	MACC	0.50° × 0.25°	4 layers, 3.5 km	CB4	2 modes, VBS	×							
TR1	WRF/CMAQ	MACC	30 km × 30 km	24 layers, 10hPa	CB5	3 modes	×	×	×					
UK1	WRF/CMAQ	MACC	15 km × 15 km	23 layers, 100 hPa	CB5-TUCL	3 modes	×	×	×	×				
UK2	WRF/CMAQ	HTAP	30 km × 30 km	23 layers, 100 hPa	CB5-TUCL	3 modes	×	×						
UK3	WRF/CMAQ	MACC	18 km × 18 km	35 layers, 16 km	CB5	3 modes	×	×	×					
US3	WRF/CMAQ	SMOKE	12 km × 12 km	35 layers, 50 hPa	CB5-TUCL	3 modes					×	×	×	×

Table 2. Exposure-response functions, the concentrations metrics, and economic valuations used in the EVA model.

Health effects (compounds)	Exposure-response coefficient	Valuation, € ₀₁₃
	(α)	(EU27 & NA)
Morbidity		
Chronic Bronchitis ¹ , CB (PM)	8.2E-5 cases/ μgm^{-3} (adults)	38,578 per case
Restricted activity days ² , RAD (PM)	=8.4E-4 days/ μgm^{-3} (adults)	98 per day
	-3.46E-5 days/ μgm^{-3} (adults)	
	-2.47E-4 days/ μgm^{-3} (adults>65)	
	-8.42E-5 days/ μgm^{-3} (adults)	
Congestive heart failure ³ , CHF (PM)	3.09E-5 cases/ μgm^{-3}	10,998 per case
Congestive heart failure ³ , CHF (CO)	5.64E-7 cases/ μgm^{-3}	
Lung cancer ⁴ , LC (PM)	1.26E-5 cases/ μgm^{-3}	16,022 per case
Hospital admissions		
Respiratory ⁵ , RHA (PM)	3.46E-6 cases/ μgm^{-3}	5,315 per case
Respiratory ⁵ , RHA (SO ₂)	2.04E-6 cases/ μgm^{-3}	
Cerebrovascular ⁶ , CHA (PM)	8.42E-6 cases/ μgm^{-3}	6,734 per case
Asthma children (7.6 % < 16 years)		
Bronchodilator use ⁷ , BUC (PM)	1.29E-1 cases/ μgm^{-3}	16 per case
Cough ⁸ – COUC (PM)	4.46E-1 days/ μgm^{-3}	30 per day
Lower respiratory symptoms ⁷ , LRSA (PM)	1.72E-1 days/ μgm^{-3}	9 per day
Asthma adults (5.9 % > 15 years)		
Bronchodilator use ⁹ , BUA (PM)	2.72E-1 cases/ μgm^{-3}	16 per case
Cough ⁹ , COUA (PM)	2.8E-1 days/ μgm^{-3}	30 per day
Lower respiratory symptoms ⁹ , LRSA (PM)	1.01E-1 days/ μgm^{-3}	9 per day
Mortality		
Acute mortality ^{10,11} (SO ₂)	7.85E-6 cases/ μgm^{-3}	1,532,099 per case
Acute mortality ^{10,11} (O ₃)	3.27E-6*SOMO35 cases/ μgm^{-3}	
Chronic mortality ^{4,12} , YOLL (PM)	1.138E-3 YOLL/ μgm^{-3} (>30 years)	57,510 per YOLL
Infant mortality ¹³ , IM (PM)	6.68E-6 cases/ μgm^{-3} (> 9 months)	2,298,148 per case

¹ Abbey et al. (1995), ² Ostro (1987), ³ Schwartz and Morris (1995), ⁴ Pope et al. (2002), ⁵ Dab et al. (1996), ⁶ Wordley et al. (1997), ⁷ Roemer et al. (1993), ⁸ Pope and Dockerey (1992), ⁹ Dusseldorp et al. (1995), ¹⁰ Anderson (1996), ¹¹ Touloumi (1996), ¹² Pope et al. (1995), ¹³ Woodruff et al. (1997).

Table 3. Health impacts calculated by the mean of individual model estimates (denoted as MM_{mi}) and the standard deviation, multi-model mean ensemble without error reduction (MM_m) and the optimal ensemble (MM_{Opt}) in Europe and the U.S. See Table 2 for the definitions of health impacts. PD stands for premature death. All health impacts are in units of number of cases $\times 1000$, except for Infant Mortality (IM), which reports directly the number of cases.

	EU			NA		
	MM_{mi}	MM_m	MM_{Opt}	MM_{mi}	MM_m	MM_{Opt}
CB	360±89	360	468	142±74	142	125
RAD	368 266±90 670	368245	478073	145 337±75 250	145337	127921
RHA	23±5	23	28	10±4	8	7
CHA	46±11	46	60	19±10	19	16
CHF	31±6	31	38	13±6	9	8
LC	55±14	55	72	22±11	22	19
BDUC	10 766±2 650	10766	13976	4 566±2 383	4566	4019
BDA	70 492±17 400	70489	91511	27 819±14 400	27819	24485
COUC	37 198±9 160	37196	48289	15 776±8 230	15776	13886
COUA	72 566±17 900	72562	94203	28 637±14 830	28637	25206
LRSC	14 355±3 530	14354	18635	6 088±3 180	6088	5359
LRSA	26 175±6 400	26174	33980	10 330±5 350	10330	9092
AYOLL	26±13	23	20	25±7	9	9
YOLL	4 111±1 010	4111	5337	1 481±762	1481	1304
PD	414±98	410	524	165±76	149	133
IM*	403±99	403	524	143±75	143.3667	126.1

Table 4. External costs (in million Euros) related to the health impacts of air pollution as calculated by the individual models over Europe and the United States.

Models	CO	SO ₂	O ₃	PM _{2.5}	TOTAL
Europe					
DE1	70	19 000	22 000	155 000	196 000
DK1	80	13 000	24 000	237 000	274 000
ES1	70	8 000	6 000	339 000	353 000
FI1	90	18 000	5 000	335 000	358 000
FRES1	90	15 000	13 000	305 000	333 000
IT1	80	17 000	21 000	413 000	451 000
IT2	70	11 000	6 000	253 000	270 000
NL1	70	12 000	18 000	215 000	245 000
TR1	110	30 000	35 000	376 000	441 000
UK1	80	28 000	25 000	280 000	333 000
UK2	80	34 000	27 000	340 000	401 000
UK3	80	47 000	25 000	279 000	351 000
MEAN	81	21 000	19 000	294 000	334 000
MEDIAN	80	17 500	21 500	292 500	342 000
The United States					
DE1	30	9 000	21 000	46 000	76 000
DK1	55	11 000	39 000	123 000	172 000
US3	60	14 000	22 000	155 000	191 000
MEAN	50	11 500	27 000	108 000	146 000
MEDIAN	55	11 000	22 000	123 000	172 000

Table 5. Annual average RMSE of the multi-model ensemble mean (MM_m) and of the optimal reduced ensemble mean (MM_{opt}) for the health impact-related species. Units are ppb for the gaseous species and $\mu\text{g m}^{-3}$ for $\text{PM}_{2.5}$.

	O_3		CO		SO_2		$\text{PM}_{2.5}$	
	MM_m	MM_{opt}	MM_m	MM_{opt}	MM_m	MM_{opt}	MM_m	MM_{opt}
Europe								
Winter	10.3	8.6	502.4	490.3	6.3	5.6	22.5	20.7
Spring	12.4	9.6	247.1	239.5	4.6	3.1	9.9	7.8
Summer	13.4	10.7	197.4	188.0	3.9	2.3	8.2	5.7
Autumn	10.7	8.8	314.5	305.5	4.6	3.1	11.0	8.7
Annual	11.7	9.4	315.3	305.8	4.8	3.5	12.9	10.7
North America								
Winter	10.9	10.4	356.7	328.1	5.7	5.5	8.3	8.1
Spring	12.0	11.4	288.7	270.2	5.4	5.1	7.2	6.6
Summer	15.1	13.0	258.3	238.7	5.4	5.0	9.7	8.8
Autumn	12.8	11.6	330.6	307.6	5.8	5.3	7.8	7.2
Annual	12.7	11.6	308.6	286.1	5.6	5.2	8.2	7.7

Table 6. Impact of the emission reduction scenarios on avoided premature death (Δ PD) and corresponding change in external cost as calculated by the multi-model mean over Europe and the United States.

Source	Receptor			
	Europe		The United States	
	Δ PD	Δ Total Cost (billion €)	Δ PD	Δ Total Cost (billion €)
GLO	-54 000 \pm 18 000	-56 \pm 18	-27 500 \pm 14 000	-24 \pm 10
NAM	-940 \pm 1100	-1.4 \pm 0.4	-25 000 \pm 12 000	-21 \pm 9
EUR	-47 000 \pm 24 000	-47 \pm 16	-	-
EAS	-	-	-1 900 \pm 2 200	-2.5 \pm 3




Fig.1. Population density (population per grid box) over a) the United States and b) Europe.




Fig. 2. Observed and simulated (base case) monthly a) O₃, b) CO, c) SO₂ and d) PM_{2.5} concentrations over Europe.




Fig. 3. Observed and simulated (base case) monthly a) O₃, b) CO, c) SO₂ and d) PM_{2.5} concentrations over the U.S.




Fig. 4. Spatial distribution of annual MM mean bias ($\mu\text{g}/\text{m}^3$) for a) O₃, b) CO, c) SO₂ and d) PM_{2.5} over Europe.




Fig. 5. Spatial distribution of annual MM mean bias (ppb for gases and $\mu\text{g}/\text{m}^3$ for $\text{PM}_{2.5}$) for a) O_3 , b) CO , c) SO_2 and d) $\text{PM}_{2.5}$ over North America.




Fig. 6. Spatial distribution of the number of total premature death (PD: units in number of cases) in a) the United States and b) Europe and the relative change (%) in the number of premature death (PD) in response to the GLO scenario in c) the United States and d) Europe in 2010 as calculated by the multi-model mean ensemble.