

## Response to Co-Editor and Reviewers

Comparing airborne and satellite retrievals of cloud optical thickness and particle effective radius using a spectral radiance ratio technique: Two case studies for multilayer cirrus and deep convective clouds

acp-2017-758 : Krisna et al.

Email to: [trismono\\_candra.krisna@uni-leipzig.de](mailto:trismono_candra.krisna@uni-leipzig.de)

We thank the reviewer and editor for the time and efforts reading our manuscript and providing valuable comments. We have made revisions according to your comments below. The changes in the manuscript are written in *italic* inside quotation marks (""). The diff file indicating changes in the manuscript is enclosed in the end of this document.

Comments to the Author:

I got one formal review and one informal review.

Both reviewers recommend acceptance, one asked for minor revisions:

The second reviewer wrote:

The authors have done a remarkable work addressing my questions and concerns. There are only a few lingering (minor) issues:

- Please, revise the writing style of the manuscript. There are several places in which the sentences lack style, i.e. they are grammatically correct but need to be reworded (e.g. first two sentences of section 4.6). I do acknowledge that the latest manuscript is in better shape, though.

### **Response:**

We have changed the manuscript according to the language style along with the grammar checking. Regarding the specific comment given by the reviewer, we revised the first two sentences of Section 4.6.

*"Time series of  $\tau$  and reff retrieved from SMART and MODIS radiance measurements, along with the MODIS cloud product, are compared for the two cloud cases. The MODIS cloud product, namely MYD06\_L2, provides three different reff (so-called reff\_L,1640, reff\_L,2130, and reff\_L,3700), which are retrieved using respective near-infrared wavelengths centered at  $\lambda = 1640$  nm, 2130 nm, and 3700 nm (Platnick et al., 2017)."*

- Regarding the parameterization to derive the new MODIS band 6 radiances. It says that the validity of the parameterization was tested with observations, but how are you going to test it if the actual MODIS band 6 is not functional. Did you use synthetic measurements to test the parameterization?

**Response:**

The reviewer is correct that the actual MODIS band 6 is noisy or nonfunctional. For more precise, 15 of 20 detectors are in trouble. The striping due to this problem is shown by Fig. 1a below. To check the parameterization, we are not only using synthetic measurement. If we do so, of course it will result a perfect agreement (Fig. 1b). The validity of band 6 retrieval is examined using remaining pixels of MODIS band 6 (along the flight legs and neighbouring pixels). The analysis of a linear regression results in a slope of 0.95 (5% bias) and a correlation coefficient of 0.94 (Fig. 1c).




Fig. 1. Actual MODIS band 6 (a). Comparison of actual and retrieved band 6 using synthetic measurements for different prescribed cloud properties (b). Comparison of actual and retrieved MODIS radiance band 6.

- Page 16, line 3: Rather than citing Zhang and King, I would recommend to mention Liang et al. (2009) because, to the best of my knowledge, Liang et al. is the first study that applies the heterogeneity index to MODIS reflectances. In addition, inhomogeneities are also producing a spatial heterogeneity effect, which is unrelated to the 3D radiative effects.

Liang, L., L. Di Girolamo, and S. Platnick (2009), View-angle consistency in reflectance, optical thickness and spherical albedo of marine water-clouds over the northeastern

**Response:**


We thank for the reference suggested by the reviewer. In the revised manuscript, we added Liang et al. (2009) in the following form:

*“Liang et al. (2009), Zhang and Platnick (2011), and King et al. (2013) estimated the influence of 3-D radiative effects using the cloud heterogeneity index ( $\sigma_{sub}$ )...”*

King et al. (2013) investigated specifically the influence of cloud heterogeneity (both sub-pixel and surrounding-pixel) on the retrieval of  $r_{eff}$  which highlighted the potentially large influence of 3-D effects when comparing  $r_{eff}$  from different instruments. They found that the differences between retrieved and in situ values are getting higher with increasing cloud heterogeneity.

- Figure 9: This is a nice figure. My only criticism is on the use of over/underestimation of tau and  $r_{eff}$ , as it is a little bit confusing. It sounds better for instance: “prescribed tau\_liq is overestimated (underestimated).”

We have changed the figure as follows:



- Table 4 is interesting. It is confusing the use of the eta letter. Just use  $r_e$  plus/minus sigma. I am not aware of many studies that show such a nice comparison between the three different MODIS  $r_{eff}$  and in-situ observations. Kokhanovsky et al. (2013) attempted to retrieve the cloud vertical structure of stratiform clouds with three MODIS channels, and then compare their results with in-situ observations. However, their attempt was applied to warm clouds only.

A. A. Kokhanovsky, D. Painemal and V. V. Rozanov, "The Intercomparison of Satellite-Derived and In Situ Profiles of Droplet Effective Radii in Marine Stratocumulus Clouds," in IEEE Geoscience and Remote Sensing Letters, vol. 10, no. 5, pp. 1147-1151, Sept. 2013. doi: 10.1109/LGRS.2012.2233710

## Response :

The reviewer is correct that Kokhanovsky et al. (2013) aimed to compare remote sensing and in situ reff of liquid water clouds. However, what they did specifically is to retrieve the profile of reff by assuming a linear increase of reff. A similar work has been done by Wang et al. (2009), but for ice clouds (has been cited in our paper). Nevertheless, we think that such comparisons are rather different with our work because we did not retrieve the full profile.

Instead, we have cited Painemal and Zuidema (2011) and King et al. (2013), who performed rather similar studies with us but for liquid water clouds. Painemal and Zuidema (2011) compared MODIS and in situ reff at cloud top (averaged between geometrical depth of 25 m). King et al. (2013) re-implemented the approach by Painemal and Zuidema (2011), and additionally calculated the weighting reff for comparing with MODIS reff. Both references have been cited in our paper. We also encountered similar results with them, that the retrieved and in situ reff differ of up to 20%.

We have revised Table 4 as follows:

**Table 4.** The mean  $\pm$  standard deviation of  $r_{\text{eff}}$  from in situ (CCP), retrievals (SMART and MODIS), and MODIS cloud product (MYD06\_L2) for different near-infrared wavelengths between 1240 nm - 3700 nm. The wavelengths have been sorted in order that the degree of absorption by cloud particles increases to the right. The last line is the the weighting-altitude  $z_w^*$ .

$r_{\text{eff}} \pm \sigma$	1240 nm	1700 nm	1640 nm	2130 nm	1550 nm	1500 nm	3700 nm
CCP ( $\mu\text{m}$ )	$19.0 \pm 9.8$	$18.3 \pm 9.6$	$18.0 \pm 9.5$	$17.5 \pm 9.4$	$17.0 \pm 9.3$	$16.7 \pm 9.3$	$7.0 \pm 5.0$
SMART ( $\mu\text{m}$ )	$22.7 \pm 8.8$	$16.5 \pm 6.7$	$15.6 \pm 3.9$	-	$13.9 \pm 3.7$	$15.7 \pm 2.1$	-
MODIS ( $\mu\text{m}$ )	$22.4 \pm 8.6$	-	$15.0 \pm 1.9$	$14.8 \pm 4.9$	-	-	$7.2 \pm 5.1$
MYD06_L2 ( $\mu\text{m}$ )	-	-	-	$6.2 \pm 1.2$	-	-	$4.8 \pm 3.7$
$z_w^*$ (km)	11.39	11.42	11.44	11.46	11.48	11.49	11.89

Based on the comments by the two (experienced) reviewers, this paper is accepted subject to the above recommended minor revisions.

Non-public comments to the Author:

One of the reviewers recommended confidentially (in my words) careful revision by the prime author in close cooperation with his senior co-authors.

# Comparing airborne and satellite retrievals of cloud optical thickness and particle effective radius using a spectral radiance ratio technique: Two case studies for **multilayer**-cirrus and deep convective clouds

Trismono C. Krisna<sup>1</sup>, Manfred Wendisch<sup>1</sup>, André Ehrlich<sup>1</sup>, Evelyn Jäkel<sup>1</sup>, Frank Werner<sup>1,\*</sup>, Ralf Weigel<sup>3,4</sup>, Stephan Borrmann<sup>3,4</sup>, Christoph Mahnke<sup>3</sup>, Ulrich Pöschl<sup>4</sup>, Meinrat O. Andreae<sup>4,6</sup>, Christiane Voigt<sup>2,3</sup>, and Luiz A. T. Machado<sup>5</sup>

<sup>1</sup>Leipziger Institut für Meteorologie (LIM), Universität Leipzig, Leipzig, Germany

<sup>2</sup>Institut für Physik der Atmosphäre, Deutsches Zentrum für Luft und Raumfahrt (DLR), Oberpfaffenhofen, Germany

<sup>3</sup>Institut für Physik der Atmosphäre, Johannes Gutenberg-Universität Mainz, Mainz, Germany

<sup>4</sup>Biogeochemistry, Multiphase Chemistry, and Particle Chemistry Departments, Max Planck Institute for Chemistry (MPIC), Mainz, Germany

<sup>5</sup>Center of Weather Forecast and Climates Studies, National Institute for Space Research, Sao Jose Dos Campos, Brazil

<sup>6</sup>Scripps Institution of Oceanography, University of California San Diego, La Jolla, California, USA

\* now at : Joint Center for Earth Systems Technology, University of Maryland, Baltimore, MD, USA

*Correspondence to:* Trismono Candra Krisna  
(trismono\_candra.krisna@uni-leipzig.de)

**Abstract.** Solar radiation reflected by cirrus and deep convective clouds (DCCs) was measured by the Spectral Modular Airborne Radiation Measurement System (SMART) installed on the German HALO (High Altitude and Long Range Research Aircraft) during the ML-CIRRUS and the ACRIDICON-CHUVA campaigns. On particular flights, HALO performed measurements closely collocated with overpasses of the Moderate Resolution Imaging Spectroradiometer (MODIS) on board of the Aqua satellite. A cirrus cloud located above liquid water clouds and a DCC topped by an anvil cirrus are analyzed in this paper. Based on the nadir spectral upward radiance measured above the two clouds, the optical thickness  $\tau$  and the particle effective radius  $r_{\text{eff}}$  of the cirrus and DCC are retrieved using a radiance ratio algorithm technique, which considers the cloud thermodynamic phase, the vertical profile of cloud microphysical properties, the presence of multilayer clouds, and the heterogeneity of the surface albedo. For the cirrus case, the comparison of  $\tau$  and  $r_{\text{eff}}$  retrieved on the basis of SMART and MODIS measurements yields a normalized mean absolute deviation of up to 1.2% for  $\tau$  and 2.1% for  $r_{\text{eff}}$ . For the DCC case, deviations of up to 3.5% for  $\tau$  and 6.5% for  $r_{\text{eff}}$  are obtained. The larger deviations in the DCC case are mainly attributed to the fast cloud evolution and significant three-dimensional (3-D) radiative effects. Measurements of spectral upward radiance at near-infrared wavelengths are employed to investigate the vertical profile of  $r_{\text{eff}}$  in the cirrus. The retrieved values of  $r_{\text{eff}}$  are compared with corresponding in situ measurements using a vertical weighting method. Compared to the MODIS observations, measurements of SMART provide more information on the vertical distribution of particle sizes close to the cloud top, which allow reconstructing the profile of  $r_{\text{eff}}$  at close to the cloud top. The comparison between retrieved and in situ values  $r_{\text{eff}}$  yields a normalized mean absolute deviation which ranges between 1.5 – 10.3% and a robust correlation coefficient of 0.82.

## 1 Introduction

Clouds constitute an important component of the global climate system. Covering about 75% of the Earth, their high albedo strongly affects ~~to~~ the Earth's energy budget (Wylie et al., 2005; Kim and Ramanathan, 2008; Stubenrauch et al., 2013). In particular, optically thin cirrus is expected to ~~contribute to a warming of~~ warm the atmosphere below the cloud, while thick  
5 cirrus may ~~cool it~~ rather cool (e.g., Liou, 1986; Wendisch et al., 2005, 2007; Voigt et al., 2017). Cirrus clouds reflect solar radiation and reduce the loss of radiative energy to space due to absorption of terrestrial radiation and re-emission at a lower temperature (greenhouse effect). ~~They pose large challenges in predicting future climate changes (Heymsfield et al., 2017) because they are not adequately represented in general circulation models in spite of~~ Given the fact that their regional coverage can be as high as about 50% in the tropics and 30% over Europe, cirrus clouds pose large challenges when predicting future  
10 climate changes (Heymsfield et al., 2017) because they are not adequately represented in general circulation models.

On the other hand, deep convective clouds (DCCs) alter the radiative energy distribution in the atmosphere by reflection of solar and absorption or emission of terrestrial radiation, as well as by changes of liquid and ice water and hydrometeor profiles (Jensen and Del Genio, 2003; Sherwood et al., 2004; Sohn et al., 2015). Their life cycle is determined by complex microphysical processes including different cloud particle growth/shrinking mechanisms, changes of the thermodynamic phase,  
15 and the development of precipitation. DCCs are optically thick and often associated with heavy precipitation, strong turbulence, considerable vertical motion, lightning, hail formation and icing (Mecikalski et al., 2007; Lane and Sharman, 2014).

Two important properties which determine the cloud radiative impact are the cloud optical thickness  $\tau$  and particle effective radius  $r_{\text{eff}}$  (King et al., 2013). ~~They will decide if a cloud has cooling or warming effect~~ These parameters determine the cooling and warming effects of clouds (Slingo, 1990; Shupe and Intrieri, 2004). Several passive remote sensing techniques have been  
20 developed to retrieve  $\tau$  and  $r_{\text{eff}}$  using spectral upward (cloud-reflected) solar or emitted thermal-infrared radiance measured by airborne and satellite sensors, where the most common technique relies on the bi-spectral methods (e.g., Nakajima and King, 1990; King et al., 1997; Stephens and Kummerow, 2007; Platnick et al., 2017). ~~A~~ Furthermore, a radiance ratio method was introduced by Werner et al. (2013), ~~who showed that the use of radiance ratios is~~ which capable to reduce the retrieval uncertainties. ~~The cloud properties are retrieved by inversion of radiative transfer model simulations, which is often realized~~  
25 ~~by pre-calculated lookup tables.~~

Airborne remote sensing of cirrus and ~~DCCs~~ DCC properties gives a snapshot of the cloud field only, whereas satellite remote sensing (e.g., MODIS) may provide statistical data on a global scale and record long time series to determine temporal changes of cloud properties (Rosenfeld and Lensky, 1998; Lindsey et al., 2006; Berendes et al., 2008). Post-launch validation activities of satellite measurements are crucial to ~~verifying~~ verify the quality of satellite products. It is essential to address all  
30 components of the measurement system, i.e., sensors, algorithms, along with the originally measured radiances and derived data products, and continue validation activities throughout the satellite lifetime (Larar et al., 2010). Radiance measurements above highly reflecting surfaces such as salt lake, desert, snow/ice (Wan, 2014) and clouds (Mu et al., 2017) are usually evaluated in order to monitor the long term stability of the satellite sensors. An estimated uncertainty of about 1 – 5% in case of MODIS reflective solar bands (RSBs) was reported by Xiong et al. (2003). This measurement error propagates into the retrieval

results. Additional uncertainties may arise from inappropriate assumptions of ~~the~~ surface albedo and ~~the~~ ice crystal habit in case of ice or mixed-phase clouds. According to Rolland and Liou (2001), Fricke et al. (2014), and Ehrlich et al. (2017), an inaccurate assumption of ~~the~~ surface albedo can lead to uncertainties of up to 83% for  $\tau$  and 62% for  $r_{\text{eff}}$ . Eichler et al. (2009) demonstrated, that uncertainties of up to 70% for  $\tau$  and 20% for  $r_{\text{eff}}$  are obtained when an inappropriate ice crystal habit is assumed in cirrus retrievals. ~~Further~~Furthermore, the influence of three-dimensional (3-D) radiative effects ~~that can due to the horizontal heterogeneity that~~ enhance the retrieval uncertainties has been demonstrated by ~~King et al. (2013), and therefore should be considered when analyzing comparisons of cloud properties retrieved from different instruments~~Liang et al. (2009), Zhang and Platnick (2011), and King et al. (2013).

Among others, Platnick (2000) and van Diedenhoven et al. (2016) emphasized ~~the fact~~ that  $r_{\text{eff}}$  retrieved from reflected solar radiation measurements depends on the vertical penetration of reflected photons into the cloud. At a wavelength with higher absorption by cloud particles, the probability of photons being scattered back out of the cloud without being absorbed decreases. Therefore, retrievals ~~of  $r_{\text{eff}}$  using different near-infrared wavelengths with different absorption by cloud particles~~ will result in  $r_{\text{eff}}$  ~~related to~~ from different cloud altitudes. This ~~approach~~ technique commonly assumes in-cloud vertical homogeneity, where the ~~result is a single retrieved  $r_{\text{eff}}$  represents a~~ bulk value of  $r_{\text{eff}}$  ~~representing~~ the entire cloud layer. Thus, the retrieved  $r_{\text{eff}}$  ~~does not represent a particle size at a single layer as measured by~~ is hardly comparable with in situ observations, where the particle effective radius is sampled at a specific cloud altitude  $r_{\text{eff}}(z)$ . ~~These different approaches need to be kept in mind when comparing remote sensing and in situ measurements, otherwise a systematic discrepancy might be misinterpreted.~~ A direct comparison at a certain cloud altitude is problematic because it is unclear for what level the retrieved  $r_{\text{eff}}$  corresponds to the in situ  $r_{\text{eff}}$ . This needs to be kept in mind when comparing remote sensing and in situ measurements, otherwise a systematic discrepancy might be misinterpreted.

Studies ~~for~~ of liquid water clouds by Painemal and Zuidema (2011) and King et al. (2013), who compared ~~the~~  $r_{\text{eff}}$  retrieved from MODIS observations with the mean value of  $r_{\text{eff}}$  measured by cloud probes near the cloud top, revealed absolute deviations of up to 20%. King et al. (2013) argued that there is no apparent link between the variation of the  $r_{\text{eff}}$  retrieved using different near-infrared wavelengths of MODIS and the vertical structure of  $r_{\text{eff}}$  measured by in situ methods, ~~while~~ while Painemal and Zuidema (2011) identified four ~~reasons, such as~~ potential reasons for this deviation: the variability of droplet size distributions, ~~forming the formation~~ of precipitation, water vapor absorption above the cloud, and viewing geometry dependent biases, ~~as potential contributors to the deviation.~~ For cirrus clouds, Wang et al. (2009) and Zhang et al. (2010) demonstrated that the differences between retrievals and in situ measurements are also influenced by the ~~simplification in the retrieval algorithm which assumes~~ simplifying assumption of in-cloud vertical homogeneity.

Standard satellite ~~retrieval methods~~ retrievals such as that applied by MODIS commonly assume a priori, that there is only one single homogeneous cloud layer with a specific thermodynamic phase, either liquid water or ice (Platnick et al., 2017). However, studies by Hahn et al. (1984) and Warren et al. (1985) analyzing ground-based observations reported, that the coexistence of multilayer clouds (e.g., cirrus above liquid water clouds) is found in about 50% of the data cases. Chang and Li (2005) and Sourdeval et al. (2015) have demonstrated, that omitting the low liquid water cloud in the retrieval algorithm ~~will introduce~~ introduces significant uncertainties in the retrieved cirrus properties.



In order to assess the ~~aspects~~ issues discussed above, collocated airborne and satellite remote sensing measurements accompanied by in situ observations are necessary. The similar observation geometry of airborne and satellite radiation sensors allows a direct comparison of upward radiance data and a stringent validation of methodologies and retrieval algorithms. The validity of the retrieval results can be explored by comparison with collocated in situ measurements. This has been realized in this paper

5 for two ~~different~~ typical cloud cases, a cirrus above low liquid water clouds and a DCC topped by an anvil cirrus. Measurements of spectral solar radiation using the Spectral Modular Airborne Radiation Measurement System (SMART) installed on board of HALO High Altitude and Long Range Research Aircraft (HALO) during the Mid-Latitude Cirrus (ML-CIRRUS) campaign (Voigt et al., 2017) and the Aerosol, Cloud, Precipitation, and Radiation Interaction and Dynamic of Convective Clouds System - Cloud Processes of the Main Precipitation Systems in Brazil: A Contribution to Cloud Resolving Modelling and to the Global Precipitation Measurement (ACRIDICON-CHUVA) ~~campaigns~~ campaign (Wendisch et al., 2017) are

10 analyzed. For the purpose of airborne-satellite validation, designated flights above clouds were carried out during the ML-CIRRUS and ACRIDICON-CHUVA (Wendisch et al., 2016; Voigt et al., 2017) campaigns, which were closely collocated with overpasses of the A-Train satellites (Savtchenko et al., 2008). HALO with its long endurance of up to 8 hours and high ceiling of up to 15 km altitude is optimally suited to fly above cirrus and DCCs. ~~High cirrus and DCCs are an appropriate target to perform airborne-satellite comparison because in high altitudes, measurements of upward radiance (cloud-reflected) are only marginally affected by atmospheric interferences due to scattering and absorption by gas molecules and aerosol particles.~~

15 The two airborne campaigns, involved instrumentations, and satellite observations are introduced in Section 2. In Section 3, the comparison techniques, data filters, and results of upward radiance comparison are presented. The radiance ratio algorithm and uncertainty estimation, impact of underlying liquid water cloud on the cirrus retrieval, forward simulation of vertically inhomogeneous cloud, vertical weighting function, and results of  $\tau$  and  $r_{\text{eff}}$  comparison are discussed in Section 4. In Section 5, the methods and results of the comparison between in situ and retrieved effective radius are presented. Finally, the conclusions are given in Section 6.

20

## 2 Observations

### 2.1 Airborne campaigns

25 ~~Data from two airborne campaigns with HALO are used in this study.~~ Between 21 March 2014 and 15 April 2014, the ML-CIRRUS campaign performed 16 research flights over Europe and the Atlantic ocean to study nucleation, life-cycle, and climate impact of natural cirrus and aircraft-induced contrail cirrus (Voigt et al., 2017; Schumann et al., 2017). Between 1 September 2014 and 4 October 2014, the ACRIDICON-CHUVA campaign performed 14 research flights combined with satellite and ground-based observations over the Brazilian Amazon rainforest to quantify aerosol-cloud-precipitation interactions and the thermodynamic, dynamic, and radiative effects of tropical deep convective clouds (DCCs) over the Amazon rainforest (Wendisch et al., 2016). One common objective of ML-CIRRUS and ACRIDICON-CHUVA was to compare airborne and satellite measurements and products. Therefore, closely collocated measurements with overpasses of the A-Train satellites were performed. One flight from the ML-CIRRUS flight number 15 (ML-15, 13 April 2014) and another one from the ACRIDICON-




5 CHUVA flight number 18 (AC-18, 28 September 2014) were selected for detailed analyses. The flight path of ML-15 is shown in Fig. 1a. During the MODIS overpass at 13:55:00 UTC, HALO flew west of Portugal over the North Atlantic. In this area, a wide field of cirrus was located above low liquid water clouds (stratocumulus). Fig. 1b shows the flight trajectory of AC-18, when HALO flew in the north-west of Brazil over Amazonian rainforest during MODIS overpass at 17:55:00 UTC, ~~where~~. [In this location](#), a DCC topped by an anvil cirrus was observed.

## 10 2.2 Airborne instrumentation

A comprehensive overview of commonly applied airborne instrumentation is given by (Wendisch and Brenguier, 2013). During the ML-CIRRUS and ACRIDICON-CHUVA campaign, a set of remote sensing and in situ instruments were operated on board of HALO (Wendisch et al., 2016; Voigt et al., 2017). SMART measured spectral upward radiance  $I_{s,\lambda}^{\uparrow}$ , as well as spectral upward  $F_{s,\lambda}^{\uparrow}$  and downward irradiance  $F_{s,\lambda}^{\downarrow}$ . The index "s" refers to measurements by SMART, while  $\lambda$  indicates spectral quantities in units of  $\text{nm}^{-1}$ . The irradiance data can be used to determine the spectral surface albedo (Wendisch et al., 2001; Wendisch and Mayer, 2003; Wendisch et al., 2004). An active stabilization system keeps the optical inlets in a horizontal position during aircraft movements of up to  $\pm 6^\circ$  from the horizontal plane (Wendisch et al., 2001). The spectral resolution defined by the full width at half maximum (FWHM) is 2 - 3 nm for the VNIR spectrometer and 8 - 10 nm for the SWIR spectrometer (Werner et al., 2013).

SMART has two separate types of spectrometers, which measure in the solar spectrum. The Visible to Near Infrared (VNIR) spectrometer covers wavelengths from 300 - 1000 nm and the Shortwave-Infrared (SWIR) spectrometer covers from 1000 - 2200 nm. The combination of both spectrometers covers approximately 97% of the entire solar spectrum (Bierwirth, 2008). However, due to the decreasing sensitivity of the spectrometers at small and large wavelengths, the reasonable wavelength range was restricted to 400 - 1800 nm.



**Figure 1.** (a) is the HALO flight path of ML-15 on 13 April 2014 while (b) is for AC-18 on 28 September 2014 overlaid with MODIS true color image. The yellow cross indicates the flight section that is selected for the analysis.

In this study, only the radiance data are analyzed. The radiance optical inlet has a field of view (FOV) of  $2^\circ$  looking at nadir (Wolf et al., 2017). The nadir radiance measured by SMART is comparable to measurements of MODIS reflective solar bands (RSBs) in the band number 1 - 19, and 26 ranging between 410 - 2130  $\mu\text{m}$  (Xiong and Barnes, 2006). SMART is calibrated radiometrically before, during, and after each campaign using certified calibration standards traceable to ~~NIST~~ (the National Institute of Standards and Technology (NIST)) and by the secondary calibration using a travelling standard. The measurement uncertainty of  $I_{s,\lambda}^\uparrow$  is determined by the accuracy of the spectral calibration, spectrometer noise and dark current, as well as radiometric and transfer calibrations (Eichler et al., 2009; Brückner et al., 2014; Wolf et al., 2017). The main uncertainty results from the Signal-to-Noise-Ratio (SNR) and the calibration standard, while spectral and transfer calibration errors are negligible (Wolf et al., 2017). The resulting total uncertainty is about 4% for the VNIR and 10% for the SWIR.

The Cloud Combination Probe (CCP) incorporates two separate instruments, the Cloud Droplet Probe (CDP) and the greyscale Cloud Imaging Probe (CIPgs) (Weigel et al., 2016). The CCP overall covers a diameter range from 2  $\mu\text{m}$  to 960  $\mu\text{m}$ , including large aerosol particles, liquid cloud droplets and small frozen hydrometeors (Klingebiel et al., 2015). The CDP part detects the forward-scattered laser light when cloud particles cross the CDP laser beam (Lance et al., 2010). Thus, the CDP provides an improved replacement for the Forward Scattering Spectrometer Probe (FSSP) (Dye and Baumgardner, 1984; Baumgardner et al., 1985). Molleker et al. (2014) showed, that the CCP exhibits a nominal limit for cloud particle diameters from 3  $\mu\text{m}$  up to 50  $\mu\text{m}$ . The CIPgs records two-dimensional shadow images of cloud particles in a size range from 15  $\mu\text{m}$  up to 960  $\mu\text{m}$  with an optical resolution of 15  $\mu\text{m}$  (Klingebiel et al., 2015; Weigel et al., 2016). Special algorithms are used to process and analyze the captured images in order to estimate particle number concentrations, particle size distributions, and to differentiate particle shapes (Korolev, 2007).

The CCP measurements are employed to derive the  $r_{\text{eff}}$  for the comparison with the retrieval products from SMART and MODIS. The  $r_{\text{eff}}$  from the CCP is derived from the geometrical properties and number of detected particles. Many definitions of  $r_{\text{eff}}$  exist as summarized in McFarquhar and Heymsfield (1998). In general,  $r_{\text{eff}}$  as a measure for the cloud radiative properties is defined as the ratio of the third to the second moment of a cloud particle size distribution, implying spheres of equivalent cross-sectional area for any cloud particle shape (McFarquhar and Heymsfield, 1996; Frey et al., 2011). The accuracy of the cloud particle sizing is conservatively estimated to be about 10% for spherical particles (Molleker et al., 2014). The sizing uncertainty increases as a function of particles shape complexity (i.e., when dendrites or particles with elevated aspect ratio were predominating). The size bin limits of the CCP cloud particle data are adapted to reduce ambiguities due to the Mie curve, particularly for cloud particles with small sizes less than 5  $\mu\text{m}$ . The instrument sample volume is calculated as a product of the probe air speed (measurement condition) and the instrument specific effective detection area. All concentration data are corrected concerning the air compression upstream of the underwing cloud probe at the high flight speeds (Weigel et al., 2016). The robust performance of the specific CCP instrument used in this study was demonstrated by Frey et al. (2011) for tropical convective outflow, by Molleker et al. (2014) for polar stratospheric clouds, Klingebiel et al. (2015) for low level mixed-phase clouds in the Arctic, as well as by Braga et al. (2017) and Cecchini et al. (2017) for tropical convective clouds.

Water vapor was measured by the Water Vapor Analyzer (WARAN), which is a tunable diode laser hygrometer based on the absorption of a laser beam by gaseous water molecules at  $\lambda = 1370 \text{ nm}$  (Voigt et al., 2014; Kaufmann et al., 2014). The

WARAN is installed on the forward-facing HALO trace gas inlet (HALO-TGI). The instrument measures total water, i.e., gas phase plus enhanced ice water content (IWC), in the range between 50 - 40,000 ppm with an accuracy of about  $\pm 50$  ppm or 5% of reading. Detailed descriptions about the measurement strategy and uncertainties in the data processing are discussed in Afchine et al. (2017). IWC is derived from the difference between the amount of total enhanced water ( $\text{H}_2\text{O}_{\text{tot}}$ ) and the amount of gas phase water ( $\text{H}_2\text{O}_{\text{gas}}$ ) (Kaufmann et al., 2016). Due to the enhancement factor (Voigt et al., 2006) at the HALO-TGI, which is about 20 - 35, the minimum detectable IWC is in the range between 1 - 2000 ppm ( $1 - 2000 \times 10^{-2} \text{ mg m}^{-3}$ ). In this study, the IWC is used to obtain the profile of cloud optical thickness  $\tau(z)$ .

## 2.3 Satellite observations

Satellite data used in this study stem from the Level 1B Moderate Resolution Imaging Spectroradiometer (MODIS) - Aqua collection 6. Detailed instrument specifications and features of MODIS have been described by Platnick et al. (2003), Xiong and Barnes (2006), and others. The data contain calibrated and geolocated radiances and reflectances for 36 discrete spectral bands distributed between 0.41  $\mu\text{m}$  and 14.2  $\mu\text{m}$ , including 20 reflective solar bands (RSBs) and 16 thermal emissive bands (TEBs) (Platnick et al., 2003; Xiong and Barnes, 2006), with a nadir horizontal resolutions of about 1 km. The radiances are generated from MODIS Level 1A scans of raw radiance and in the process converted to geophysical units. The solar reflectance values are based on a solar diffuser panel for reflectance calibration up through the RSBs and an accompanying diffuser stability monitor for assessing the stability of the diffuser of up to 1  $\mu\text{m}$  (Platnick et al., 2003). The spectral response is determined by an interference filter overlying a detector array imaging a 10-km along track scene for each scan (40, 20, and 10 elements arrays for the 250 m, 500 m, and 1 km bands, respectively). Onboard instruments used for in-orbit radiometric calibration were discussed by Xiong et al. (2003) and Sun et al. (2007).

## 3 Comparison of upward radiance

### 3.1 Spectral and spatial resolution adjustment

SMART and MODIS have different spectral resolutions. MODIS measures in broad spectral bands, while SMART measures in much narrower spectral with FWHM between 2 - 10 nm. To allow the comparison, the spectral upward radiance measured by SMART  $I_{s,\lambda}^\uparrow$  must be convoluted with the MODIS relative spectral response  $R(\lambda)$ . The convoluted radiance of SMART  $I_{S,\lambda}^\uparrow$  is calculated by:

$$I_{S,\lambda}^\uparrow = \frac{\int_{\lambda_1}^{\lambda_2} I_{s,\lambda}^\uparrow \cdot R(\lambda) d\lambda}{\int_{\lambda_1}^{\lambda_2} R(\lambda) d\lambda} \quad (1)$$

In this study, upward radiances centered at the MODIS band 1 ( $\lambda = 645$  nm), band 5 ( $\lambda = 1240$  nm), and band 6 ( $\lambda = 1640$  nm) will be primarily used to retrieve  $\tau$  and  $r_{\text{eff}}$ . It is known that 15 of the 20 detectors in the MODIS-Aqua band 6 are either nonfunctional or noisy. However, according to Wang et al. (2006), the MODIS radiance band 6  $I_{M,B6}$  can be retrieved using band 7  $I_{M,B7}$  ( $\lambda = 2130$  nm). This technique was originally developed and tested on the basis of snow surfaces assuming that

the spectral characteristics of the snow reflectivity between MODIS band 6 and 7 do not change significantly for different snow types. Assuming that ice clouds and snow have similar optical properties, the same approach can be applied. Similar to Wang et al. (2006), a parameterization of  $I_{M,B6}$  is developed on the basis of radiative transfer simulations of upward radiance performed for cirrus with different  $\tau$  and  $r_{\text{eff}}$ . A polynomial fit is applied to quantify the relation between  $I_{M,B6}$  and  $I_{M,B7}$  which result the parameterization:

$$I_{M,B6} = -81.033 \cdot I_{M,B7}^2 + 3.257 \cdot I_{M,B7} + 0.002 \quad (2)$$

The validity of the parameterization is tested using the remaining detectors of MODIS band 6 for observations above cirrus (not shown here). The linear regression between original and retrieved  $I_{M,B6}$  showed shows differences below 5% (slope of 0.95 and zero bias) with a correlation coefficient of 0.94.

MODIS data used in this study are delivered at a horizontal resolution of 1 km at nadir, whereas the spatial resolution of SMART varies depending on the flight altitude and temporal resolution. At a flight altitude of 10 km, SMART has a swath of approximately 349 m at the Earth surface. During the two campaigns, the temporal resolution of SMART was between 0.2 - 0.5 s, depending on the measurement conditions. This has to be considered in the data analysis. In order to decrease biases resulting from comparisons of individual measurements, SMART data are averaged over 1 s resolution using a binning method.

### 3.2 Data filter


Only clouds with a top altitude higher than 8 km are selected for this study. The higher proximity to TOA reduces the influence of scattering and absorption by atmospheric molecules and aerosol particles above cloud. Consequently, no correction for the influence of the atmospheric layer above HALO is needed. To assure a similar viewing zenith angle of SMART and MODIS, only nadir observations in the center of MODIS swath were selected for the comparison. Werner et al. (2013) discussed that off-nadir measurements of less than  $5^\circ$  may lead to a bias in the retrieved  $\tau$  and  $r_{\text{eff}}$  of up to 1% and 5%, respectively. To minimize this bias, SMART measurements with roll and pitch angles larger than  $3^\circ$  are discarded and only straight flight legs with altitude changes of less than 50 m are analyzed.

**Table 1.** Flight descriptions and atmospheric conditions during cloud measurements. Horizontal wind speed  $v$  and solar zenith angle  $\theta_0$  are averaged during the selected time series.

Flight	Date	Cloud Type	Appearance	$z_t$ (km)	Time - UTC (HH:MM:SS)	$v$ ( $\text{m s}^{-1}$ )	$\theta_0$ ( $^\circ$ )
ML-15	04/13/2014	Cirrus above liquid cloud	Homogeneous	12	13:56:20 - 13:57:35	21	37
AC-18	09/28/2014	Anvil topped DCC	Inhomogeneous	8	17:56:00 - 17:57:30	9	26


The nadir point of MODIS moves much faster than the aircraft. Therefore, it is impossible that SMART and MODIS always measure exactly above each other along the joint flight track. To analyze the effects caused by time shifts between SMART and

MODIS measurements, data from the ML-CIRRUS and ACRIDICON-CHUVA are divided into groups within and without a threshold  $|\Delta t|$  of 500 s for the cirrus and 300 s for the DCC. Scatter plots of SMART and MODIS radiance at  $\lambda = 645$  nm are shown in Fig. 2a for the cirrus and Fig. 2b for the DCC. For the cirrus (Fig. 2a),  $I_{S,645}^{\uparrow}$  and  $I_{M,645}^{\uparrow}$  are in a better agreement for  $|\Delta t| < 500$  s with a correlation coefficient  $R^2 = 0.96$ , while for  $|\Delta t| > 500$  s deviations are larger with  $R^2 = 0.58$ . The large scatter for  $|\Delta t| > 500$  s is mainly attributed to the fast horizontal wind speed during ~~cirrus~~ the measurements which was 21  $\text{ms}^{-1}$  on average. Additionally, the wind direction is also a key factor causing a significant cloud drift for the larger time delay. For the DCC (Fig. 2b), the scatter is significantly larger compared to the cirrus for the given threshold of  $|\Delta t| < 300$  s and even worse for the threshold of  $|\Delta t| > 300$  s with  $R^2 = 0.79$  and  $-0.09$ , respectively. In this case, the horizontal wind speed is smaller with an average of 9  $\text{ms}^{-1}$ , but the fast cloud evolution is the major issue. Luo et al. (2014) and Schumacher et al. (2015) reported, that tropical DCCs located at altitude between 6 - 8 km typically have an updraft velocity about 2 - 4  $\text{ms}^{-1}$ . According to this analysis, the comparison are restricted to  $|\Delta t| < 500$  s for the cirrus case, while for the DCC case the threshold is tightened to  $|\Delta t| < 300$  s.



**Figure 2.** Scatter plots of upward radiance at  $\lambda = 645$  nm measured by SMART ( $I_{S,645}^{\uparrow}$ ) and MODIS ( $I_{M,645}^{\uparrow}$ ) within a threshold of 500 s for the cirrus (a) and 300 s the DCC (b). Blue circles and red triangles represents data within and without the predetermined threshold.

After the filtering, two suitable cases are left which fulfill most requirements of the analysis. The first case, a cirrus cloud located above low liquid water clouds (stratocumulus) is selected from ML-15 between 13:56:20 - 13:57:35 UTC as shown in Fig. 3a. The cloud top altitude  $z_t$  of the cirrus was about 12 km while HALO flew at about 12.3 km altitude. The second case, a DCC topped by an anvil cirrus is selected from AC-18 between 17:56:00 - 17:57:30 UTC as presented in Fig. 3b. The  $z_t$  of the selected DCC was about 8 km while HALO flew at 8.3 km altitude. Flight descriptions and atmospheric conditions during cloud measurements are summarized in Table 1. The selected time periods extend to 75 s for the cirrus and 90 s for the DCC case. For HALO flying at constant altitude, those correspond to horizontal distances of about 15 km and 18 km, respectively. For the MODIS data, the cloud mask algorithm by Ackerman et al. (1998) is employed to discriminate clear and cloudy pixels for the MODIS data. Cloud edges are associated with sharp changes of  $I_{\lambda}^{\uparrow}$  and higher 3-D radiative effects.



**Figure 3.** MODIS radiance band 1 ( $\lambda = 645$  nm) for the cirrus case (a) and the DCC case (b) superimposed with the selected flight legs of HALO during cloud measurements (red line). The flight direction is from point A to B.

Fisher (2014) discussed variations in cloud height and surface orology to find an offset distance assigned to an uncertainty of  $\pm 40$  m. Therefore, the first and the last pixel of MODIS cloudy pixels are discarded in the data analysis.


### 3.3 Result of upward radiance comparison

- 5 Upward radiances measured by SMART and MODIS are compared for the two selected cloud cases. Fig. 4 shows time series of upward radiance measured by SMART  $I_{S,\lambda}^\uparrow$  and MODIS  $I_{M,\lambda}^\uparrow$  centered at  $\lambda = 645$  nm (a), 1240 nm (b), and 1640 nm (c) for the cirrus case, while Fig. 5 shows the same for the DCC case. Those three wavelengths will be primarily utilized to retrieve the cloud properties in this study. The scatter plots of the respective measurements are shown in Fig. 6. ~~The time series~~ Time series of upward radiances in Fig. 4 and Fig. 5 illustrate, that the cirrus is more homogeneous along the flight legs compared to the
- 10 DCC. For the DCC, the cloud anvil is observed between 17:56:00 - 17:56:20 UTC. Later,  $I_{645}^\uparrow$  increases sharply corresponding to the DCC core and decreases again towards the cloud edge. Fig. 6 shows **obviously**, that the scatters are larger for the DCC case which are **mainly** caused by the remaining effects of the cloud evolution. For the cirrus case, the scatters are significantly smaller because high cirrus typically change less rapidly.


- Fig. 7 shows the comparison of mean spectral upward radiance measured by SMART and MODIS for the cirrus (a) and
- 15 DCC case (b). The solid line represents spectral radiance measured by SMART  $I_{S,\lambda}^\uparrow$ , while  $I_{S,\lambda}^\uparrow$  is the convoluted radiance of SMART using Eq. 1, and  $I_{M,\lambda}^\uparrow$  is the radiance measured by MODIS. The values of mean  $\pm$  standard deviation  $\eta$  at each spectral wavelength are summarized in Table 2. To quantify the agreement, the normalized mean absolute deviation  $\zeta$  is calculated by:

$$\zeta = \frac{1}{n} \sum_{i=1}^n \left| \frac{x_i - \bar{x}}{\bar{x}} \right| \quad (3)$$






**Figure 4.** Time series of  $I_{\lambda}^{\uparrow}$  centered at  $\lambda = 645$  nm (a), 1240 nm (b), and 1640 nm (c) measured by SMART (black) and MODIS (red) for the cirrus case. Shaded areas are measurement uncertainties. Gaps on the time series indicate when the shutter of SMART closed for dark current measurements.




**Figure 5.** Same as Fig. 4 but for the DCC case.





**Figure 6.** Scatter plots of upward radiance centered at  $\lambda = 645$  nm (a), 1240 nm (b), and 1640 nm (c) measured by SMART  $I_{S,\lambda}^\uparrow$  and MODIS  $I_{M,\lambda}^\uparrow$  for the cirrus (red triangles) and the DCC (blue dots) case. The dashed line represents the one-to-one line.



**Figure 7.** Comparison of mean  $I_\lambda^\uparrow$  measured by SMART and MODIS for the cirrus case (a) and the DCC case (b) at  $\lambda$  between 400 - 1800 nm. Error bars represent measurement uncertainties. Wavelengths centered at  $\lambda = 645$  nm, 1240 nm, and 1640 nm are indicated by dashed lines while grey band correspond to the interval of MODIS relative spectral response  $R(\lambda)$  for the respective wavelengths.

where  $n$  is the number of observed values,  $x_i$  are the individual values, and  $\bar{x}$  is the mean value of the radiances measured by SMART and MODIS along the selected time series. For the cirrus case,  $\zeta_{645}$  is found to be 0.04%, while  $\zeta_{1240}$  and  $\zeta_{1640}$  are 7.68% and 1.36%, respectively. For the DCC case,  $\zeta_{645}$  yields a value of 4.25%, while  $\zeta_{1240}$  and  $\zeta_{1640}$  are 6.72% and 5.61%, respectively. The good agreement between SMART  $I_{S,1640}^\uparrow$  and MODIS  $I_{M,1640}^\uparrow$  again justifies the application of the retrieval of MODIS band 6 using the parameterization given in Eq. 2. Overall, all the values of  $\zeta$  in Table 2 lie within

**Table 2.** Comparison of SMART  $I_{S,\lambda}^\dagger$  and MODIS  $I_{M,\lambda}^\dagger$  for the cirrus (ci) and DCC case.  $\eta$  is the mean  $\pm$  standard deviation with a subscript of "S" for SMART and "M" for MODIS.  $\zeta$  is the normalized mean absolute deviation between SMART and MODIS measurements.

$\lambda$ (nm)	$\eta_{S,ci}$	$\eta_{M,ci}$	$\zeta_{ci}$ (%)	$\eta_{S,DCC}$	$\eta_{M,DCC}$	$\zeta_{DCC}$ (%)
421	0.231 $\pm$ 0.014	0.234 $\pm$ 0.011	0.81	0.295 $\pm$ 0.122	0.251 $\pm$ 0.013	8.06
469	0.266 $\pm$ 0.018	0.265 $\pm$ 0.014	0.20	0.335 $\pm$ 0.149	0.351 $\pm$ 0.050	2.34
555	0.229 $\pm$ 0.018	0.224 $\pm$ 0.013	1.19	0.290 $\pm$ 0.135	0.303 $\pm$ 0.047	2.12
645	0.193 $\pm$ 0.016	0.193 $\pm$ 0.012	0.04	0.241 $\pm$ 0.117	0.263 $\pm$ 0.042	4.25
858	0.125 $\pm$ 0.011	0.128 $\pm$ 0.008	1.29	0.162 $\pm$ 0.069	0.167 $\pm$ 0.018	1.47
905	0.096 $\pm$ 0.008	0.104 $\pm$ 0.007	4.36	0.124 $\pm$ 0.059	0.129 $\pm$ 0.016	1.96
936	0.048 $\pm$ 0.005	0.056 $\pm$ 0.005	7.49	0.069 $\pm$ 0.043	0.080 $\pm$ 0.018	7.95
940	0.062 $\pm$ 0.006	0.071 $\pm$ 0.005	7.18	0.084 $\pm$ 0.047	0.099 $\pm$ 0.018	8.26
1240	0.052 $\pm$ 0.004	0.061 $\pm$ 0.004	7.68	0.057 $\pm$ 0.029	0.065 $\pm$ 0.009	6.72
1375	0.005 $\pm$ 0.001	0.005 $\pm$ 0.001	3.24	0.004 $\pm$ 0.004	0.004 $\pm$ 0.003	6.17
1640	0.024 $\pm$ 0.002	0.025 $\pm$ 0.001	1.36	0.016 $\pm$ 0.010	0.018 $\pm$ 0.001	5.61

the measurement uncertainties. The ~~radiance measurements of SMART and MODIS agree better~~ comparison yields a better agreement for the cirrus ~~case~~ than for the DCC case. The larger deviations in case of  $\eta$  of DCC are not only influenced by the cloud evolution, but also due to larger 3-D radiative effects. ~~Zhang et al. (2011) and King et al. (2013) estimate~~ Liang et al. (2009), Zhang and Platnick (2011), and King et al. (2013) estimated the influence of 3-D radiative effects using the cloud heterogeneity index  $\sigma_{sub}$ , ~~which is calculated as the ratio of~~. The  $\sigma_{sub}$  is defined as a ratio between the standard deviation and the mean value of MODIS radiance band ~~2~~,  ~~$\sigma_{sub} = 2$  ( $\lambda = 0.1$  is obtained for 858 nm and 250 m spatial resolution)~~. For the cirrus case ~~while the DCC case shows higher inhomogeneities~~, the  $\sigma_{sub}$  results in a value of about 0.1. Higher heterogeneities are found for the DCC case with  $\sigma_{sub} =$  of about 0.4. These values suggests This shows, that 3-D radiative effects are ~~obviously~~ potentially larger for the DCC case, and therefore have to be considered when interpreting the retrieval results from different instruments.

## 4 Retrieval of cloud optical thickness $\tau$ and particle effective radius $r_{eff}$

### 4.1 Radiance ratio retrieval and uncertainty estimation

A radiance ratio technique adapted from Werner et al. (2013) is applied to retrieve  $\tau$  and  $r_{eff}$  of the cirrus and the DCC based on the nadir upward radiance measured by SMART and MODIS. In case radiance ratios are applied, the uncertainties are reduced because the uncertainties of the radiation source identically influence all measured radiances, and therefore do not contribute to the uncertainty of the ratio. In the radiance ratio algorithm, the upward radiance at the MODIS bands centered at  $\lambda_0 = 645$

nm (band 1),  $\lambda_1 = 1240$  nm (band 5), and  $\lambda_2 = 1640$  nm (band 6) are employed to calculate the following radiance ratios,

$$\mathfrak{R}_{1240} = I_{\lambda_1}^\uparrow / I_{\lambda_0}^\uparrow \text{ and } \mathfrak{R}_{1640} = I_{\lambda_2}^\uparrow / I_{\lambda_0}^\uparrow.$$

In the retrieval algorithm, a decision tree is applied to select the retrieval mode. The retrieval can be performed either in the liquid water or ice mode. To decide which mode is used, a cloud phase index  $I_p$  is determined by the spectral slope method according to Jäkel et al. (2013). In this study,  $I_p$  is defined from the spectral slope of SMART radiance measurements at  $\lambda = 1550$  nm and 1700 nm, where the value is typically larger than zero for ice clouds. A threshold of 0.2 is used to discriminate between ice and liquid water clouds. For the cirrus case, time series of  $I_p$  calculated from the SMART observations yield values larger than 0.4 indicating ice clouds. This indicates, that for the cirrus case the underlying liquid water clouds did not significantly influence  $I_p$ . Additionally, the high values of  $I_p$  show that  $I_p$  is mostly sensitive to the thermodynamic phase of the top cloud layer (cirrus), while the underlying liquid water clouds below the cirrus have a limited influence on the radiances within the wavelength range analyzed for the  $I_p$ . For the DCC case,  $I_p$  varies between 0.2 - 0.4 along the time series with a mean value of 0.25. Based on the high  $I_p$  values, the retrievals in both analyzed cloud cases are performed by assuming ice clouds.

Forward simulations of upward radiance have been performed by 1-D radiative transfer simulations using the radiative transfer package LibRadtran 2.0 (Mayer, 2005; Emde et al., 2016), the discrete ordinate radiative transfer solver (DISORT) version 2 (Stamnes et al., 2000), and assuming vertically homogeneous clouds. The atmospheric profiles of gases and constituents are adapted from the standard profile (Anderson et al., 1986) "mid-latitude" for ML-CIRRUS and "tropical" for ACRIDICON-CHUVA, and are adjusted to radio soundings close to the measurement area. Extraterrestrial spectral irradiance is taken from Gueymard (2004). The standard aerosol particle profile for "spring/summer condition" of "maritime aerosol type" is applied (Shettle, 1989). For the cirrus case, the spectral surface albedo  $\rho$  of ocean implemented in the forward simulations was measured by SMART. For the DCC case, which is above Amazonian rainforest, no corresponding SMART albedo measurements at low altitude covering exactly the same flight path are available. In this area, the heterogeneity of the surface albedo is very high because forested and deforested areas are located close to each other. This implies, that a representative assumption of a homogeneous surface for the whole flight legs is not appropriate. Therefore,  $\rho$  derived from the MODIS BRDF/Albedo product (Strahler et al., 1999) is used to include the horizontal variability of the surface albedo of tropical rainforest.


In the forward simulations, the optical properties of liquid water droplet are derived from Mie calculation (Wiscombe, 1980). The assumption of ice crystal habit considers ice crystal shapes measured by the in situ probes during the two campaigns (Voigt et al., 2017; Järvinen et al., 2016). For the cirrus, representative ice crystal properties of a general habit mixture based on severely roughened aggregates (so-called GHM) by Baum et al. (2014) is applied, while for the DCC ice properties of plate with a high surface roughness (Yang et al., 2013) are assumed. These particle habits differ from the MODIS collection 6 retrievals which use severely-roughened compact aggregates of solid columns (so-called aggregated columns) by Yang et al. (2013). A sensitivity study infers that the retrievals assuming GHM and plate generally will result in a larger  $\tau$  and smaller  $r_{\text{eff}}$  (not shown here), which is in agreement with findings by van Diedenhoven et al. (2014) and Holz et al. (2016). The radiance is simulated for both, the actual flight altitude of HALO for the SMART measurements and the top of atmosphere (TOA) for MODIS observations. Due to the high flight altitude, no significant differences are ~~observed~~obtained.

Due to the multilayer cloud situation in the ~~For the~~ cirrus case, a liquid water cloud layer is considered in the forward simulations due to the multilayer cloud situation. The properties of the liquid water cloud are estimated by comparing the entire spectral signature of the radiance measured by SMART and the simulations assuming different combinations of cloud properties. For the average of the selected time series, a simulation (not shown here) with a liquid water cloud located between 1.5 and 2 km,  $\tau = 8$ , and  $r_{\text{eff}} = 10 \mu\text{m}$  shows the best agreements with the measurements particularly in the water vapor absorption bands (e.g.,  $\lambda = 940 \text{ nm}$  and  $1135 \text{ nm}$ ) and the  $\text{O}_2$  A-band ( $\lambda = 760 \text{ nm}$ ), which are sensitive to such multilayer cloud conditions (Rozanov and Kokhanovsky, 2004; Wind et al., 2010). The radiance lookup tables for the cirrus case are shown in Fig. 8a and 8b, whereas Fig. 8c and 8d are for the DCC case. The upward radiance at a non-absorbing wavelength  $I_{645}^\uparrow$  is combined with  $\mathfrak{R}_{1240}$  (combination 1 - C1) and with  $\mathfrak{R}_{1640}$  (combination 2 - C2).  $I_{645}^\uparrow$  is most sensitive to  $\tau$ , while ratios  $\mathfrak{R}_{1240}$  and  $\mathfrak{R}_{1640}$  are most sensitive to  $r_{\text{eff}}$ . For the cirrus case, the lookup tables cover  $\tau$  between 1 - 5 with steps of 1 and  $r_{\text{eff}}$  between 5 - 60  $\mu\text{m}$  with steps of 3  $\mu\text{m}$ . For the DCC case, the lookup tables cover  $\tau$  between 6 - 40 with steps of 1 for  $\tau$  between 6 - 22 and steps of 2 for  $\tau$  between 24 - 40, while  $r_{\text{eff}}$  ranges between 5 - 90  $\mu\text{m}$  with steps of 3  $\mu\text{m}$  for  $r_{\text{eff}}$  between 5 - 56  $\mu\text{m}$  and steps of 4  $\mu\text{m}$  for  $r_{\text{eff}}$  between 60 - 90  $\mu\text{m}$ .

The measurements of SMART (black crosses) and MODIS (blue circles) are included for both scenes in Fig. 8. For the C1 which is based on  $I_{1240}^\uparrow$ , the MODIS data does not match the lookup table solution space. The results in Section 3.3 show clearly, that  $I_{M,1240}^\uparrow$  are higher than  $I_{S,1240}^\uparrow$  by about 15%. Using the original  $I_{M,1240}^\uparrow$  for the cirrus case, all the retrievals of  $r_{\text{eff}}$  fail because the measurements lie far outside the lookup table solution space (see Fig. 8a), while for the DCC case the retrieval failure is smaller (see Fig. 8c). Enhancing retrieval failure in the cirrus case is due to the larger  $\theta_0$ . At a larger  $\theta_0$ , the upward radiance becomes more insensitive to the changes of  $r_{\text{eff}}$  and consequently the lookup tables are denser. To gain meaningful retrieved cloud properties, a correction of  $I_{M,1240}^\uparrow$  is applied. Following Lyapustin et al. (2014), a correction factor  $g$  is calculated by the slope of linear regression between  $I_{M,1240}^\uparrow$  and  $I_{S,1240}^\uparrow$ , which results in  $g = 0.88$  for the cirrus case and  $g = 0.90$  for the DCC case. The corrected  $I_{M,1240}^\uparrow$  (red circles) are added in Fig. 8 ~~, which and~~ now match the lookup table solution space. Therefore, all following radiance ratio retrievals for the two selected cloud cases use these corrected  $I_{M,1240}^\uparrow$ . In the radiance ratio method, measurement uncertainties of 4% for  $I_{645}^\uparrow$  and 6% for  $\mathfrak{R}_{1240}$  and  $\mathfrak{R}_{1640}$  are considered. The retrieval uncertainties are estimated by considering the measurement uncertainties expressed by its double standard deviation  $2\sigma$ . The retrieval is performed by varying each measurements separately by adding and subtracting  $2\sigma$  which resulted in four solutions. The median of the four solutions is used as the retrieval result of  $\tau$  and  $r_{\text{eff}}$ , while the standard deviation is used to represent the retrieval uncertainties,  $\Delta\tau$  for  $\tau$  and  $\Delta r_{\text{eff}}$  for  $r_{\text{eff}}$ . Note that the retrievals of  $r_{\text{eff}}$  using C1 will result in larger uncertainties than by using C2 due to smaller absorption by cloud particles at  $\lambda = 1240 \text{ nm}$ . As a the result, the lookup tables of  $r_{\text{eff}}$  for C1 are more narrow. At a given 6% measurement uncertainty of  $\mathfrak{R}_{1240}$ , ~~the retrieval a retrieval of~~ a retrieval of  $r_{\text{eff}}$  can result in uncertainties up to 50%.

## 4.2 Impact of underlying liquid layer clouds on the cirrus retrieval

For the cirrus case, the properties of the low liquid water cloud are assumed to be constant along the flight legs. This assumption might not hold in reality and affect the retrieved cirrus properties. Therefore, the sensitivity of the cirrus retrieval on the




**Figure 8.** Radiance lookup tables for the cirrus case (a,b) and DCC case (c,d). (a) and (c) are using C1 ( $I_{645}^{\uparrow}$  and  $\mathcal{R}_{1240}$ ), while (b) and (d) are using C2 ( $I_{645}^{\uparrow}$  and  $\mathcal{R}_{1640}$ ). For the cirrus case, the simulations are performed with  $\theta_0 = 37^\circ$  and assuming GHM (Baum et al., 2014), while for the DCC case  $\theta_0 = 26^\circ$  and the ice habit of plate (Yang et al., 2013) are applied. Radiance measurements of SMART and MODIS are illustrated by symbols.

assumed properties of the low liquid water cloud is quantified using radiative transfer simulations. Spectral upward radiances are simulated for different combinations of liquid water cloud and cirrus properties. The liquid water cloud is varied for  $\tau_{liq} = 6 - 10$  (steps of 1) and  $r_{eff,liq} = 6 - 14 \mu\text{m}$  (steps of 2  $\mu\text{m}$ ), while the cirrus is changed for  $\tau_{ci} = 2 - 8$  (steps of 1  $\mu\text{m}$ ) and  $r_{eff,ci} = 10 - 40 \mu\text{m}$  (steps of 5). These simulated upward radiances are used as synthetic measurements and analyzed with the retrieval algorithm using C2 ( $I_{645}^{\uparrow}$  and  $\mathcal{R}_{1640}$ ), which assumes a liquid water cloud with  $\tau_{liq} = 8$  and  $r_{eff,liq} = 10 \mu\text{m}$ . The comparison of synthetically retrieved and original  $\tau_{ci}$  and  $r_{eff,ci}$  is shown in Fig. 9. The annotation of "overestimation" (below one-to-one line) and "underestimation" (above one-to-one line) corresponds to when the retrieval is run with an overestimation and underestimation of the properties of the low liquid water cloud. The retrieved  $\tau_{ci}$  are analyzed in Fig. 9a for different  $\tau_{liq}$ , while  $r_{eff,ci}$  and  $r_{eff,liq}$  are fixed to 20  $\mu\text{m}$  and 10  $\mu\text{m}$ , respectively. Similarly, the retrieved  $r_{eff,ci}$  are analyzed in Fig. 9b for different  $r_{eff,liq}$  but for a fixed combination of  $\tau_{ci} = 3$  and  $\tau_{liq} = 8$ . In general, the simulations results show that an overestimation of  $\tau_{liq}$  leads to an underestimation of



$\tau_{ci}$  because in this case, the liquid water cloud contributes more strongly to the reflected radiation than in reality. Therefore, a smaller  $\tau_{ci}$  is required to match the measurement, and vice versa. For the range of  $\tau_{ci}$  analyzed here, the retrieved  $\tau_{ci}$  is found to be over- or underestimated by 1.3 when in reality  $\tau_{liq}$  is 6 or 10, while the retrieval assumes  $\tau_{liq} = 8$ . These biases of  $\tau_{ci}$  show, that  $\tau_{liq}$  needs to be estimated accurately because a wrong assumption of  $\tau_{liq}$  almost directly propagates in the uncertainties of  $\tau_{ci}$ .

A similar behavior is found for the retrieval of  $r_{eff,ci}$ , where an overestimation of  $r_{eff,liq}$  leads to an underestimation of  $r_{eff,ci}$ , and vice versa. Assuming larger liquid droplets than in reality implies that these droplets contribute more strongly to the measured absorption at  $\lambda = 1640$  nm, and therefore the ice crystals only contribute less (smaller  $r_{eff,ci}$ ). Fig. 9b illustrates, that the impact of  $r_{eff,liq}$  is strongest when small liquid droplets ( $r_{eff,liq} \leq 8 \mu\text{m}$ ) are present. For larger liquid droplets ( $r_{eff,liq} > 10 \mu\text{m}$ ), the impact is reduced. The maximum uncertainties of  $r_{eff,ci}$  found for the range of  $r_{eff,ci}$  and  $r_{eff,liq}$  considered here are about  $8 \mu\text{m}$  for the underestimation of  $r_{eff,liq}$ , which show a tendency of higher uncertainties for higher  $r_{eff,ci}$ . The retrieval of  $r_{eff,ci}$  is less affected by  $r_{eff,liq}$ , when the cirrus layer is sufficiently thick ( $\tau_{ci} > 5$ ) since then the cirrus layer will dominate the reflected radiation in the absorption bands.



**Figure 9.** Comparison of synthetically retrieved  $\tau_{ci}$  (a) and  $r_{eff,ci}$  (b). Calculations in (a) are performed by changing  $\tau_{liq}$  while the original value is 8 and  $r_{eff,ci} = 20 \mu\text{m}$  and  $r_{eff,liq} = 10 \mu\text{m}$  are fixed. In (b),  $r_{eff,liq}$  is changed while the original value is  $10 \mu\text{m}$  and  $\tau_{ci} = 3$  and  $\tau_{liq} = 8$  are fixed.

### 4.3 Forward simulation of vertically inhomogeneous clouds

It is known from measurements, that the cloud particle sizes can significantly vary with altitudes. For nonprecipitating ice clouds, the ice crystal size typically decreases as a function of altitude (Heymsfield et al., 2017, e.g.,). However, to simplify the retrieval algorithm vertically homogeneous clouds are commonly assumed in the forward radiative transfer simulations. To quantify the effects of such simplifications, simulations with vertically inhomogeneous ice clouds are performed. Analytical

profiles of effective radius as a function of geometrical height are developed using a modified parameterization that was originally proposed by Platnick (2000):

$$r_{\text{eff}}(z, h) = a_0 - \left( a_1 - a_2 \cdot \frac{z}{h} \right)^{1/k} \quad (4)$$

- 10 where the altitude  $z$  ranges from 0 at the cloud base to  $h$  at the cloud top. The parameters  $a_0 = r_{\text{eff,t}} + r_{\text{eff,b}}$ ,  $a_1 = r_{\text{eff,t}}^k$ , and  $a_2 = r_{\text{eff,t}}^k - r_{\text{eff,b}}^k$  are determined from prescribed boundary condition of the cloud top effective radius  $r_{\text{eff,t}}$  and the cloud base effective radius  $r_{\text{eff,b}}$ . To represent a typical vertical structure of ice clouds,  $k = 3$  is chosen. The profiles of effective radius are coupled with the profiles of ice water content, which typically decrease as a function of altitude in ice clouds.

**Table 3.** Total optical thickness  $\tau_c$ , effective radius at the cloud top  $r_{\text{eff,t}}$  and cloud base  $r_{\text{eff,b}}$ , ice water content (IWC) from the cloud base (CB) to the cloud top (CT), with  $z_b$  and  $z_t$  are the boundary altitude of geometrical height  $z$  the cloud base and thickness  $h$  cloud top, respectively. Retrieved ~~effective radius~~  $r_{\text{eff,ret}}$  is compared to the weighting-estimate  $r_{\text{eff,w}}^*$  for two near-infrared wavelengths at  $\lambda = 1240$  nm and 1640 nm.

Cloud	Specification							Validation			
	$\tau_c$	$r_{\text{eff,b}}$	$r_{\text{eff,t}}$	$k$	IWC	$z_b$	$z_t$	$r_{\text{eff,w}}^*$ ( $\mu\text{m}$ )		$r_{\text{eff,ret}}$ ( $\mu\text{m}$ )	
		( $\mu\text{m}$ )	( $\mu\text{m}$ )		( $\text{g m}^{-3}$ )	(km)	(km)	1240 nm	1640 nm	1240 nm	1640 nm
A	3	40	10	3	0.1 - 0.04	10	12	18.3	17.7	17.9	17.3
B	15	50	20	3	0.2 - 0.1	6	8	26.6	24.1	26.1	24.0

- Fig. 10a and 10b show the profile of effective radius for a representative cirrus (cloud A) and a DCC composed of ice particles only (cloud B). The cloud profiles are divided into 20 layers for the implementation in the radiative transfer simulation, where each layer has a homogeneous thin layer of  $\Delta\tau = 0.15$  for cloud A and 0.75 for cloud B. The parameters used to set up both clouds A and B are summarized in Table 3. Forward radiative transfer simulations are performed to calculate spectral upward radiance above the cloud using an adding-superposition technique from the cloud top (CT) to the cloud base (CB).

#### 4.4 Vertical weighting function

- 20 The vertical photon transport depends on the absorption characteristics at the considered wavelengths. With increasing absorption the probability of a photon being scattered back out of the cloud without being absorbed decreases. Thus, utilizing different near-infrared wavelengths with different absorption characteristics in the retrieval will result  $r_{\text{eff}}$  from different altitudes in the cloud (King et al., 2013). To quantify this effect, the vertical weighting function  $w_m$  is investigated. The  $w_m$  describes the contribution of each cloud layer to the absorption considering multiple scattering (Platnick, 2000). Therefore, it can be used to
- 5 characterize the cloud level where the retrieved  $r_{\text{eff}}$  is most representative. For nadir observation,  $w_m$  as a function of optical



thickness  $\tau$  is expressed by:

$$w_m(\lambda, \tau, \tau_c, \mu_0, r_{\text{eff}}) = \left| \frac{dI(\lambda, \tau, \mu_0, r_{\text{eff}})}{d\tau} \right| \cdot \frac{1}{\int_0^{\tau_c} \left| \frac{dI(\lambda, \tau, \mu_0, r_{\text{eff}})}{d\tau} \right| d\tau} \quad (5)$$

$I$  is the radiance above the cloud and  $\tau_c$  is the total cloud optical thickness. Platnick (2000) showed that  $w_m$  can be used to estimate the retrieved value of effective radius  $r_{\text{eff,w}}^*$  (so-called weighting-estimate) from a given profile of  $r_{\text{eff}}(\tau)$  by:

$$r_{\text{eff,w}}^*(\lambda, \tau_c, \mu_0, r_{\text{eff}}) = \int_0^{\tau_c} w_m(\lambda, \tau, \tau_c, \mu_0, r_{\text{eff}}) r_{\text{eff}}(\tau) d\tau \quad (6)$$

$w_m$  calculated for cloud A and B are shown in Fig. 10c and 10d, respectively. For cloud A with  $\tau_c = 3$ , it is found that  $w_m$  for  $\lambda = 1240$  nm and 1640 nm are almost homogeneously distributed along the entire profile. Each cloud layer has nearly equal contribution to the absorption, and therefore to the retrieved  $r_{\text{eff}}$ . Whereas for cloud B with  $\tau_c = 15$ , the upper cloud layers contribute most to the absorption.  $w_m$  profiles for cloud A and B show, that for  $\lambda = 1640$  nm the maximum is found closer to the cloud top, while for  $\lambda = 1240$  nm it is located in a deeper layer. This illustrates, that a retrieval of  $r_{\text{eff}}$  using  $\lambda = 1640$  nm will result in an  $r_{\text{eff}}$  that represents particle sizes located at a higher altitude compared to  $\lambda = 1240$  nm. For the two idealized cloud cases (cloud A and B), this would in general lead to  $r_{\text{eff},1640} < r_{\text{eff},1240}$ . Additionally, the results show clearly that each cloud layer has a contribution to the absorption. Therefore, it should be noted that  $r_{\text{eff}}$  retrieved by this remote sensing technique

5 does not represent a particle size at a single cloud layer only.


Fig. 11a shows the spectral  $w_m$  calculated for cloud A at  $\lambda$  between 1000 - 2000 nm, while Fig. 11b is the single scattering albedo  $\tilde{\omega}_0$  of GHM with  $r_{\text{eff}}$  of 10  $\mu\text{m}$  and 15  $\mu\text{m}$ . The  $\tilde{\omega}_0$  strongly depends on  $r_{\text{eff}}$  and describes the degree of absorption by cloud particles at each individual wavelength. The  $\tilde{\omega}_0$  is smaller for larger particles, and therefore the absorption is higher. The spectral  $w_m$  at each individual cloud layer clearly shows a wavelength dependence. For a wavelength with smaller  $\tilde{\omega}_0$

10 (high absorption by cloud particles), the maximum of  $w_m$  is located closer to the cloud top. In contrast, for a wavelength with  $\tilde{\omega}_0 \approx 1$  (small absorption by cloud particles), the  $w_m$  in the lower layers significantly increases and the maximum  $w_m$  is reduced correspondingly. The spectral  $w_m$  also shows that spectral measurements in the near-infrared wavelengths offers more information on the particle sizes located in different cloud altitudes.

It is found, that  $w_m$  is a function of the cloud profile itself. Assuming a vertically homogeneous profile in the forward simulation

15 will result in different  $w_m$  compared to assuming a realistic profile. This may lead to discrepancies in the  $r_{\text{eff}}$  retrieved using both assumptions. With the help of  $w_m$ , possible impacts are investigated by comparing the weighting-estimate  $r_{\text{eff,w}}^*$  and the retrieved  $r_{\text{eff,ret}}$  using  $\lambda = 1240$  nm and 1640 nm. Radiances above cloud A and B calculated for the entire cloud layer  $I_{\lambda, \tau_c}^\uparrow$ , as described in Section 4.3, serve as synthetic measurements for the radiance ratio retrieval. Both combinations, C1 (1240 nm) and C2 (1640 nm), are employed. The resulting  $r_{\text{eff,w}}^*$  and  $r_{\text{eff,ret}}$  are summarized in Table 3. The absolute deviation between  $r_{\text{eff,ret},1240}$  and  $r_{\text{eff,w},1240}^*$  is 0.4  $\mu\text{m}$  for cloud A and 0.5  $\mu\text{m}$  for cloud B. Between  $r_{\text{eff,ret},1640}$  and  $r_{\text{eff,w},1640}^*$ , the

5 absolute deviation is 0.4  $\mu\text{m}$  for cloud A and 0.1  $\mu\text{m}$  for cloud B. The  $r_{\text{eff}}$  retrieved by using measurements at  $\lambda = 1640$  nm is consistently smaller than  $\lambda = 1240$  nm, which agree with a condition where the particle size decreases towards the cloud top.




**Figure 10.** (a) is an analytic effective radius profile of a cirrus (cloud A) while (b) is for a DCC composed of ice particles only (cloud B). Detailed specifications of the two analytic profiles are summarized in Table 3. (c) is  $w_m$  calculated for cloud A while (d) is for cloud B.

The comparisons between  $r_{\text{eff},w}^*$  and  $r_{\text{eff,ret}}$  for cloud A and B yield a systematic deviation. It is found, that retrievals using a vertically homogeneous assumption result in a slight underestimation of  $r_{\text{eff,ret}}$  compared to  $r_{\text{eff},w}^*$  which assumes a realistic cloud profile with decreasing particle size towards the cloud top. For the two realistic profiles (cloud A and B), larger particles with higher absorption are located in the lower layers. Consequently,  $w_m$  in the lower cloud layers becomes higher, while  $w_m$  in the upper cloud layers is slightly smaller compared to a vertically homogeneous cloud profile (not shown here). The Nevertheless, the impact of vertical profile assumption will decrease-reduce for retrievals using wavelengths with higher absorption by cloud particles such as  $\lambda = 1640$  nm.

#### 4.5 Impact of underlying liquid water cloud on the vertical weighting function


The changes of the vertical weighting function  $w_m$  due to the presence of liquid water clouds below cloud A and B are investigated. Therefore, the calculations of  $w_m$  for cloud A and B presented in Section 4.4 are repeated by adding a liquid water cloud layer. For cloud A, the liquid water cloud is located between 1.5 - 2 km with  $\tau = 8$  and  $r_{\text{eff}} = 10 \mu\text{m}$ , which represent a cirrus above a low liquid water cloud. For cloud B, the liquid water cloud is located between 5 and 6 km with  $\tau = 15$  and  $r_{\text{eff}} = 15 \mu\text{m}$ , which represents a DCC topped by an anvil cirrus, where the lower core of DCC is assumed to be a liquid



**Figure 11.** (a) Spectral vertical weighting function calculated for cloud A. (b) Single scattering albedo  $\tilde{\omega}_0$  of GHM (Baum et al., 2014) with  $r_{\text{eff}} = 10 \mu\text{m}$  (dashed line) and  $15 \mu\text{m}$  (solid line).

water cloud. For simplification, the profiles of liquid water cloud are assumed to be vertically homogeneous. For comparison,  $w_m$  are calculated and normalized for the ice cloud only. Fig. 12a and 12b show  $w_m$  at  $\lambda = 1240 \text{ nm}$  (black) and  $1640 \text{ nm}$  (red) calculated for cloud A and cloud B in a condition with (solid line) and without (dashed line) the presence of the liquid water cloud. Additionally, the single scattering albedo  $\tilde{\omega}_0$  of GHM (blue) and liquid droplets (red) with  $r_{\text{eff}}$  of  $10 \mu\text{m}$  (dashed line) and  $15 \mu\text{m}$  (solid line) is displayed in Fig. 12c.

According to Platnick (2000), it is expected that the low liquid water cloud changes  $w_m$  similar to a bright surface, where the maximum weighting at cloud top will be reduced and shifted to a lower altitude due to the enhanced reflection of transmitted radiation back to the cloud base eventually reaching the sensor above cloud top. Consequently, this will result in a larger retrieved  $r_{\text{eff}}$  for clouds with decreasing particle size towards the cloud top. The results in Fig. 12a and 12b show, that this indeed holds for the  $w_m$  at  $\lambda = 1240 \text{ nm}$  where scattering by cloud particles dominates. For cloud A and B, the maximum of  $w_m$  is shifted to lower altitudes due to multiple reflections of radiation between the surface (liquid water cloud) and cloud base (ice cloud). The  $w_m$  at  $\lambda = 1640 \text{ nm}$  changes differently when adding a liquid water cloud below the ice cloud. The changes of  $w_m$  for cloud A are significantly larger compared to cloud B. This behavior results from the stronger absorption by the ice particles at  $\lambda = 1640 \text{ nm}$ . For optically thick cloud B with  $\tau_c = 15$ , the ice cloud does not transmit sufficient radiation to have a strong interaction with the low level cloud which leads to a similar  $w_m$ . In contrast,  $w_m$  at cloud top is modified for optically thin cloud A with  $\tau_c = 3$  due to the underlying liquid water cloud. Here the different particle phase and size of the liquid water cloud layer lead to a reduction of the upward radiance  $I_\lambda^\uparrow$  when an ice cloud layer is added to the simulations. Given that small liquid droplets have a higher  $\tilde{\omega}_0$  at  $\lambda = 1640 \text{ nm}$ , the liquid water cloud alone reflects more strongly than together with the ice



**Figure 12.** (a) is  $w_m$  at  $\lambda = 1240$  nm and  $1640$  nm calculated for cloud A, while (b) is for cloud B. Solid line and dashed line describe  $w_m$  calculated with and without the presence of underlying liquid water cloud, respectively. (c) Single scattering albedo  $\tilde{\omega}_0$  of GHM and liquid water droplets with  $r_{\text{eff}}$  of  $10 \mu\text{m}$  and  $15 \mu\text{m}$ .

cloud which adds large ice crystals characterized by smaller  $\tilde{\omega}_0$  reducing the total  $I_\lambda^\uparrow$ . Decreasing  $I_\lambda^\uparrow$  strongly contributes to the  $w_m$  close to the cloud top, while at about  $\tau = 1$  the minimum of  $w_m$  is observed where  $I_\lambda^\uparrow$  changes only slightly. Below  $\tau = 1$  (lower altitudes), the impact of the liquid water cloud vanishes and scattering by the ice particles increases  $I_\lambda^\uparrow$  again corresponding to higher  $w_m$  towards cloud base. In general, a similar behavior is imprinted in the  $w_m$  of cloud B but not relevant for the entire  $w_m$  due to the higher  $\tau_c$  of the ice cloud. This also demonstrates, that for optically thick clouds such as the DCC case investigated in this study, a retrieval assuming only ice cloud can be applied to retrieve  $r_{\text{eff}}$  of the upper most cloud layer, even if liquid water clouds are present below the ice cloud layer.

#### 4.6 Comparison of optical thickness and effective radius retrieved by SMART and MODIS

Time series of  $\tau$  and  $r_{\text{eff}}$  retrieved from SMART and MODIS radiance measurements, and from along with the MODIS cloud product, are compared for the two cloud cases. The MODIS cloud product collection-6, namely MYD06\_L2, provides three different  $r_{\text{eff}}$  (so-called  $r_{\text{eff,L,1640}}$ ,  $r_{\text{eff,L,2130}}$ , and  $r_{\text{eff,L,3700}}$ ) retrieved using three, which are retrieved using respective near-infrared wavelengths centered at  $\lambda = 1640$  nm,  $2130$  nm, and  $3700$  nm (Platnick et al., 2017). However, the information of  $r_{\text{eff,L,1640}}$  is very limited due to problems of the detectors, and therefore it cannot be used in this comparison. Due to the similar


ice crystal absorption at  $\lambda = 1640$  nm and 2130 nm, both wavelengths have an almost identical  $w_m$  (Wang et al., 2009; Zhang et al., 2010). For typical cloud profiles analyzed in Section 4.4, the differences of  $r_{\text{eff}}$  retrieved using  $\lambda = 1640$  nm and 2130 nm are less than 1  $\mu\text{m}$ . Therefore,  $r_{\text{eff,L,2130}}$  can be compared with SMART and MODIS  $r_{\text{eff}}$  retrieved using C2 (1640 nm). For observations over land, the MODIS algorithm combines the reflectivity at  $\lambda = 645$  nm and 2130 nm (combination 3 - C3). While over ocean, it combines the reflectivity at  $\lambda = 858$  nm and 2130 nm (combination 4 - C4).

~~Time series of cirrus  $\tau$  (a) and  $r_{\text{eff}}$  (b) retrieved from SMART (black) and MODIS (red) using combination 1 (C1). The dark shaded area describes retrieval uncertainties.  $\eta_S$  (SMART) and  $\eta_M$  (MODIS) represent the mean  $\pm$  standard deviation along time series. (c) and (d) are the respective properties retrieved using combination 2 (C2). Cloud properties derived from the MODIS cloud product (MYD06\_L2),  $\tau_L$  and  $r_{\text{eff,L,2130}}$ , are shown in blue (only in panel c and d) with the corresponding  $\eta_L$ .~~

Time series of cirrus optical thickness and effective radius retrieved using C1,  $\tau_{\text{ci,C1}}$  and  $r_{\text{eff,ci,C1}}$ , are presented in Fig. 13a and 13b, respectively. The  $\eta$  describes the mean  $\pm$  standard deviation of the corresponding cloud properties along the selected time series with the subscript of "S" for SMART and "M" for MODIS. To quantify the agreement of the retrieved cirrus properties based on SMART and MODIS, the normalized mean absolute deviation  $\zeta$  is calculated. A  $\zeta_{\tau_{\text{ci,C1}}}$  of 1.2% and a  $\zeta_{r_{\text{eff,ci,C1}}}$  of 0.7% are obtained. Fig. 13c and 13d show time series of cirrus optical thickness and effective radius retrieved using C2,  $\tau_{\text{ci,C2}}$  and  $r_{\text{eff,ci,C2}}$ , respectively. A  $\zeta_{\tau_{\text{ci,C2}}}$  of 0.5% and a  $\zeta_{r_{\text{eff,ci,C2}}}$  of 2.1% are obtained. The analysis shows, that deviations between SMART and MODIS in the retrieved cloud properties are only slightly enhanced by the non-linearity in the retrieval algorithm. Additionally, cloud properties derived from the MODIS cloud product (blue) are also shown in Fig. 13c and 13d, where  $\eta$  with the subscript of "L" describes the respective mean  $\pm$  standard deviation along the selected time series.

Cirrus properties retrieved using combinations C1 and C2 are compared to the MODIS cloud product (combination C4). Along the selected time series, all combinations show that  $\tau_{\text{ci}}$  is homogeneous indicated by the small standard deviation  $\sigma_{\tau_{\text{ci}}} < 1$ . However, it is found that  $\tau_{\text{ci,L,C4}}$  derived from the MODIS cloud product significantly overestimates  $\tau_{\text{ci,C2}}$  (see Fig. 13c). The absolute deviation between the mean value  $\bar{\tau}_{\text{ci,L,C4}}$  and  $\bar{\tau}_{\text{ci,C2}}$  is found up to 4.7 (160% relative difference). For the MODIS cloud product, the retrieval is always performed with the assumption of a single cloud layer even if a multilayer condition is detected (Platnick et al., 2017). Omitting the low liquid water cloud consequently results in a significant overestimation of the retrieved  $\tau_{\text{ci}}$ . Including a low liquid water cloud in the radiance ratio retrieval as applied to SMART and MODIS, more realistic  $\tau_{\text{ci}}$  are obtained. Furthermore, small differences between  $\tau_{\text{ci,C1}}$  and  $\tau_{\text{ci,C2}}$  are found. For a cirrus cloud where the particle size decreases towards the cloud top, it is expected that  $r_{\text{eff,C1}} > r_{\text{eff,C2}}$ . Due to the remaining coupling between  $\tau$  and  $r_{\text{eff}}$  (non-orthogonal radiance lookup tables), these differences propagate into the retrieved  $\tau$ , and lead to  $\tau_{\text{C1}} > \tau_{\text{ci,C2}}$ .

The results from all approaches show that the mean  $\bar{r}_{\text{eff,ci,C1}} > \bar{r}_{\text{eff,ci,C2}} > \bar{r}_{\text{eff,ci,C4}}$ . It should be noted, that due to omitting the underlying liquid water cloud  $\bar{r}_{\text{eff,ci,C4}}$  underestimates the actual value. The difference between  $r_{\text{eff,C1}}$  and  $r_{\text{eff,C2}}$  results from the different  $w_m$  as discussed in Section 4.4, which makes  $r_{\text{eff,C1}} > r_{\text{eff,C2}}$  for a cirrus with decreasing particle size towards the cloud top. Additionally, the results show that the standard deviation  $\sigma_{r_{\text{eff,ci,C1}}} > \sigma_{r_{\text{eff,ci,C2}}} > \sigma_{r_{\text{eff,ci,C4}}}$ . This indicates, that the horizontal variability of ice crystals is higher in lower cloud layers, while close to the cloud top the ice crystals are distributed more homogeneously along the flight legs. Smaller ice particles with low sedimenting velocity remain at the higher altitudes, while larger ice particles with faster sedimenting velocity drop into the cloud layers below. This sedimentation is horizontally




**Figure 13.** Time series of cirrus  $\tau$  (a) and  $r_{\text{eff}}$  (b) retrieved from SMART (black) and MODIS (red) using combination 1 (C1). The dark shaded area describes retrieval uncertainties.  $\eta_S$  (SMART) and  $\eta_M$  (MODIS) represent the mean  $\pm$  standard deviation along time series. (c) and (d) are the respective properties retrieved using combination 2 (C2). Cloud properties derived from the MODIS cloud product (MYD06\_L2),  $\tau_L$  and  $r_{\text{eff},L,2130}$ , are shown in blue (only in panel c and d) with the corresponding  $\eta_L$ .

inhomogeneous due to the variability of the vertical wind velocity and leads to a size sorting and the observed horizontal variability of the particle sizes. The analysis shows, that the uncertainty  $\Delta r_{\text{eff},\text{ci},\text{C1}} > \Delta r_{\text{eff},\text{ci},\text{C2}}$ . This confirms, that retrievals of  $r_{\text{eff}}$  using a wavelength with a smaller absorption by cloud particles will result in a larger uncertainty. Additionally, it is found that increasing  $\tau$  and  $r_{\text{eff}}$  has a positive correlation with increasing  $\Delta\tau$  and  $\Delta r_{\text{eff}}$ , which is due to decreasing sensitivity in the radiance lookup tables for larger  $\tau$  and  $r_{\text{eff}}$ .

Same as Fig. 13 but for the DCC case.

Time series of DCC optical thickness and effective radius retrieved using C1,  $\tau_{\text{dcc},\text{C1}}$  and  $r_{\text{eff},\text{dcc},\text{C1}}$ , are shown in Fig. 14a and 14b, respectively. A  $\zeta_{\tau_{\text{dcc},\text{C1}}}$  of 1.1% and a  $\zeta_{r_{\text{eff},\text{dcc},\text{C1}}}$  of 6.5% are obtained between SMART and MODIS retrievals. Compared to the cirrus case, the larger horizontal variability indicates a strong evolution of microphysical properties in the deeper layer of DCC. Fig. 14c and Fig. 14d show time series of DCC optical thickness and effective radius retrieved using C2,  $\tau_{\text{dcc},\text{C2}}$  and  $r_{\text{eff},\text{dcc},\text{C2}}$ . A  $\zeta_{\tau_{\text{dcc},\text{C2}}}$  of 3.5% and a  $\zeta_{r_{\text{eff},\text{dcc},\text{C2}}}$  of 4.1% are obtained in this case. In addition to the fast cloud





**Figure 14.** Same as Fig. 13 but for the DCC case.

evolution, larger 3-D radiative effects are likely influencing the observations, which can enhance the deviations of retrieved  
 25 cloud properties. The cloud properties derived from the MODIS cloud product (blue) are also presented in Fig. 14c and 14d. In  
 this case (over land), the MODIS cloud product algorithm uses C3. The high values of standard deviation  $\sigma_{\tau_{dcc}}$  from approach  
 C1, C2, and C3, which are up to 10.3, indicate that  $\tau_{dcc}$  is heterogeneous except in the anvil region. The DCC anvil is observed  
 between 17:56:00 - 17:56:20 UTC, which is characterized by relatively smaller  $\tau$  between 8 - 15. Later,  $\tau_{dcc}$  increases sharply  
 corresponding to the DCC core and decreases again towards the cloud edge. The mean value  $\bar{r}_{eff,dcc,C1} > \bar{r}_{eff,dcc,C2}$  indicates  
 30 decreasing particle size towards the cloud top. It is found, that  $\bar{r}_{eff,dcc,C3}$  is larger than  $\bar{r}_{eff,dcc,C2}$  corresponding to the different  
 assumptions of the ice crystal habit of plate (SMART and MODIS retrievals) and aggregated columns (MODIS cloud product).  
 Given that  $\sigma_{r_{eff,dcc,C1}} > \sigma_{r_{eff,dcc,C2}}$  and  $\sigma_{r_{eff,dcc,C2}} < \sigma_{r_{eff,dcc,C3}}$ , this illustrates that the particle sizes are more homogeneous in  
 the level of  $r_{eff,dcc,C2}$  compared to the level of  $r_{eff,dcc,C1}$  and  $r_{eff,dcc,C3}$ .



## 5 Comparison of retrieval results with in situ measurements for the cirrus case

5 The retrieved and in situ  $r_{\text{eff}}$  are compared for the cirrus case. Here, the terminology of  $r_{\text{eff}}(z)$  is used to describe the particle effective radius sampled at a specific vertical layer  $z$ , while the retrieved  $r_{\text{eff}}$  represents a bulk property of the entire cloud as discussed in Section 4.4. ~~The~~ CCP provides  $r_{\text{eff}}(z)$  at 1 Hz temporal resolution. ~~These~~ ~~Further~~ ~~the~~ data are averaged to derive  $r_{\text{eff}}(z)$  with a vertical resolution of 65 m. Fig. 15a shows, that ~~the~~ CCP detected a cirrus between 10.7 and 12 km with the mean values (solid line) ranging between 3 - 30  $\mu\text{m}$ . The grey area illustrates the estimated uncertainties of the in situ data. The

10 smallest particles with  $r_{\text{eff}} = 3.1 \mu\text{m}$  are found at the cloud base  $z_b = 10.7 \text{ km}$  and grow rapidly up to 30.2  $\mu\text{m}$  at  $z = 10.8 \text{ km}$ . Later,  $r_{\text{eff}}$  decreases reaching a value of 8.4  $\mu\text{m}$  at the cloud top  $z_b = 11.97 \text{ km}$ .

To compare retrieved and in situ  $r_{\text{eff}}$ , the vertical weighting function  $w_m$  has to be considered. A direct comparison between  $r_{\text{eff}}$  and  $r_{\text{eff}}(z)$  at a single layer is inappropriate because both are defined differently. Note that the  $w_m$  in this study is calculated in terms of  $\tau$  increasing from ~~the~~ cloud top towards ~~the~~ cloud base. Therefore, the conversion of geometrical altitude and optical

15 thickness  $\tau(z)$  ~~has needs~~ to be specified ~~and considered in the analysis~~. For this purpose, IWC( $z$ ) measured by WARAN and  $r_{\text{eff}}(z)$  derived from CCP are converted into a profile of the extinction coefficient  $\beta(z)$  following the scheme introduced by ~~introduced by~~ Fu and Liou (1993) and Wang et al. (2009):

$$\beta_e(z) \approx \text{IWC}(z) \cdot \left[ a + \frac{b}{r_{\text{eff}}(z)} \right] \quad (7)$$

where  $a = -6.656 \times 10^{-3}$ ,  $b = 3.686$ .  $\beta_e(z)$  is in the unit of  $\text{m}^{-1}$ , IWC( $z$ ) in  $\text{g m}^{-3}$ , and  $r_{\text{eff}}(z)$  in  $\mu\text{m}$ . Further, the extinction

20 profile is used to calculate  $\tau(z)$  by integrating  $\beta_e(z)$  from ~~the~~ cloud top to the altitude level  $z$ :

$$\tau(z) = \int_z^{z_t} \beta_e(z) dz \quad (8)$$

Using  $\tau(z)$ ,  $r_{\text{eff}}(z)$  can be converted into  $r_{\text{eff}}(\tau)$ . To calculate the  $w_m$ , the cloud is divided into 20 ~~inhomogeneous~~ layers, where each cloud layer is assigned to a  $r_{\text{eff}}(\tau)$ . Finally, the  $r_{\text{eff}}(\tau)$  is convoluted with the  $w_m$  to calculate the in situ weighting-estimate  $r_{\text{eff},w}^*$  given by Eq. 6 to allow a comparison with the retrieved  $r_{\text{eff}}$ . Similarly, the weighting-altitude  $z_w^*$  which characterizes the

5 altitude corresponding to the  $r_{\text{eff},w}^*$  and the retrieved  $r_{\text{eff}}$  can be calculated by:

$$z_w^*(\lambda, \tau_c, \mu_0, r_{\text{eff}}) = \int_0^{\tau_c} w_m(\lambda, \tau, \tau_c, \mu_0, r_{\text{eff}}) z(\tau) d\tau \quad (9)$$

Due to different absorption characteristics, it is expected that  $z_w^*$  will ~~differ vary~~ for different near-infrared wavelengths. The stronger the absorption by cloud particles in the wavelength, the higher the  $z_w^*$  (closer to ~~the~~ cloud top).

The comparison of  $r_{\text{eff},w}^*$  and the mean value of retrieved  $r_{\text{eff}}$  is presented in Fig. 15b by symbols. Horizontal error bars represent the standard deviation of ~~partiele sizes~~  $r_{\text{eff}}$ . Vertical error bars indicate the estimated uncertainty of the ~~weighting-altitude~~  $z_w^*$  ~~with a value~~ of 40 m, ~~which represents~~. ~~This value is defined as~~ the standard deviation of  $z_w^*$  by varying ice crystal habits in the forward simulations. Additionally, the  $r_{\text{eff}}$  retrieved using SMART radiances measurements at  $\lambda = 1500 \text{ nm}$ ,  $1550 \text{ nm}$ , and  $1700 \text{ nm}$ , and also MODIS radiances centered at  $\lambda = 2130 \text{ nm}$ , and  $3700 \text{ nm}$  (band 20) are applied in this comparison.

**Table 4.** The mean  $\pm$  standard deviation of  $r_{\text{eff}}$  from in situ (CCP), retrievals (SMART and MODIS), and MODIS cloud product (MYD06 L2) for different near-infrared wavelengths between 1240 nm - 3700 nm. The wavelengths have been sorted in order that the degree of absorption by cloud particles increases to the right. The last line is the the weighting-altitude  $z_w^*$ .

$\bar{r}_{\text{eff}} \pm \sigma$	1240 nm	1700 nm	1640 nm	2130 nm	1550 nm	1500 nm	3700 nm
<u>CCP (<math>\mu\text{m}</math>)</u>	<u><math>19.0 \pm 9.8</math></u>	<u><math>18.3 \pm 9.6</math></u>	<u><math>18.0 \pm 9.5</math></u>	<u><math>17.5 \pm 9.4</math></u>	<u><math>17.0 \pm 9.3</math></u>	<u><math>16.7 \pm 9.3</math></u>	<u><math>7.0 \pm 5.0</math></u>
<u>SMART (<math>\mu\text{m}</math>)</u>	<u><math>22.7 \pm 8.8</math></u>	<u><math>16.5 \pm 6.7</math></u>	<u><math>15.6 \pm 3.9</math></u>	<u>~</u>	<u><math>13.9 \pm 3.7</math></u>	<u><math>15.7 \pm 2.1</math></u>	<u>~</u>
<u>MODIS (<math>\mu\text{m}</math>)</u>	<u><math>22.4 \pm 8.6</math></u>	<u>~</u>	<u><math>15.0 \pm 1.9</math></u>	<u><math>14.8 \pm 4.9</math></u>	<u>~</u>	<u>~</u>	<u><math>7.2 \pm 5.1</math></u>
<u>MYD06 L2 (<math>\mu\text{m}</math>)</u>	<u>~</u>	<u>~</u>	<u>~</u>	<u><math>6.2 \pm 1.2</math></u>	<u>~</u>	<u>~</u>	<u><math>4.8 \pm 3.7</math></u>
<u><math>z_w^*</math> (km)</u>	<u>11.39</u>	<u>11.42</u>	<u>11.44</u>	<u>11.46</u>	<u>11.48</u>	<u>11.49</u>	<u>11.89</u>


The retrieval and the calculation of  $w_m$  for  $\lambda = 3700$  nm are performed by considering both solar and thermal radiation. Using these additional wavelengths allows to enhance the vertical resolution of retrieved  $r_{\text{eff}}$ . Fig. 15b shows, that in situ  $r_{\text{eff,w}}^*$  and retrieved  $r_{\text{eff}}$  agree within the standard deviation for all altitudes and reproduce the decrease of particle size towards the cloud top. However, it is obvious that although retrievals of  $r_{\text{eff}}$  using multi near-infrared wavelengths result in particle sizes from different cloud altitudes, this **passive**-retrieval technique only provides information of particle size in the cloud top layers. This is because the retrieved  $r_{\text{eff}}$  represents a vertically weighted value, where ~~cloud the~~ the cloud top layers are weighted at most.

15  ~~$\eta_{\text{CCP}}(\text{m}) 19.0 \pm 9.8 18.3 \pm 9.6 18.0 \pm 9.5 17.5 \pm 9.4 17.0 \pm 9.3 16.7 \pm 9.3 7.0 \pm 5.0$~~   
 ~~$\eta_{\text{SMART}}(\text{m}) 22.7 \pm 8.8 16.5 \pm 6.7 15.6 \pm 3.9 13.9 \pm 3.7 15.7 \pm 2.1$~~   
 ~~$\eta_{\text{MODIS}}(\text{m}) 22.4 \pm 8.6 15.0 \pm 1.9 14.8 \pm 4.9 7.2 \pm 5.1$~~   
 ~~$\eta_{\text{MYD06}}(\text{m}) 6.2 \pm 1.2 4.8 \pm 3.7$~~   
 ~~$z_w^*(\text{km}) 11.39 11.42 11.44 11.46 11.48 11.49 11.89$~~

20 Table 4 summarizes the mean  $\pm$  standard deviation  $\eta$  of in situ  $r_{\text{eff,w}}^*$  and retrieved  $r_{\text{eff}}$  from SMART and MODIS, and  $z_w^*$  for near-infrared wavelengths between 1240 nm - 3700 nm. Additionally, MODIS cloud products (MYD06 L2),  $r_{\text{eff,L},2130}$  and  $r_{\text{eff,L},3700}$  are included in the table for the comparison. To quantify the agreement between in situ  $r_{\text{eff,w}}^*$  and retrieved  $r_{\text{eff}}$ , the normalized mean absolute deviation  $\zeta$  is calculated. The deviations of in situ  $r_{\text{eff,w}}^*$  and SMART  $r_{\text{eff}}$  range between  $\zeta = 3.2\%$  ( $\lambda = 1500$  nm) and  $\zeta = 10.3\%$  ( $\lambda = 1550$  nm). Between  $r_{\text{eff,w}}^*$  and MODIS  $r_{\text{eff}}$ , the  $\zeta$  results in a value between 1.5% for  $\lambda = 3700$  nm and 9.1% for  $\lambda = 1640$  nm. Overall, the values of  $\zeta$  are in the range between 1.5 – 10.3% and agree within the horizontal standard deviation, as shown in Fig. 15b.

(a) ~~Profile of effective radius  $r_{\text{eff}}(z)$  from the in situ measurements (CCP – solid line). The grey area represents the uncertainties of in situ data.~~ (b) ~~Comparison of the in situ  $r_{\text{eff,w}}^*$  and the mean value of  $r_{\text{eff}}$  retrieved by SMART and MODIS for  $\lambda$  between 1240 nm – 3700 nm. Horizontal error bars represent the standard deviation of effective radius, while vertical error bars are the uncertainty of the weighting-altitude.~~ (c) ~~Scatter plots between the in situ  $r_{\text{eff,w}}^*$  and the mean value of retrieved  $r_{\text{eff}}$ . The dashed line is the one-to-one line. The labels at each data point describe the wavelengths used to derive the  $r_{\text{eff}}$ .~~

30



**Figure 15.** The mean  $\pm$  standard deviation  $\eta$  (a) Profile of cirrus effective radius determined by  $r_{\text{eff}}(z)$  derived from in situ weighting estimate (CCP (solid line) and retrievals with the corresponding uncertainties (grey area). (b) Comparison of the in situ  $r_{\text{eff},w}^*$  and the mean value of  $r_{\text{eff}}$  retrieved from SMART, MODIS, and MYD06\_L2) MODIS using near-infrared wavelengths  $\lambda$  between 1240 nm - 3700 nm. The wavelengths have been sorted in order that horizontal error bars represent the degree standard deviation of absorption by cloud particles increases to  $r_{\text{eff}}$ , while vertical error bars are the right uncertainty of  $z_w^*$ . (c) Scatter plots between the in situ  $r_{\text{eff},w}^*$  and the mean value of retrieved  $r_{\text{eff}}$ . The dashed line is the weighting altitude one-to-one line. The labels at each data point describe the wavelengths used to retrieve the  $r_{\text{eff}}$ .

The  $r_{\text{eff}}$  derived from the MODIS cloud product are obviously affected by the low liquid water cloud, which is not included in the algorithm of MODIS operational retrieval. Therefore, a  $\zeta$  of 47.5% and 19.3% is obtained for  $r_{\text{eff},L,2130}$  and  $r_{\text{eff},L,3700}$ , respectively. The deviation for  $r_{\text{eff},L,2130}$  is smaller because the absorption by the ice crystals at  $\lambda = 3700$  nm is very strong. Consequently, the first top layers will dominate the absorption and significantly reduce the effect of the underlying liquid water cloud. Fig. 15c shows scatter plots of in situ  $r_{\text{eff},w}^*$  and  $r_{\text{eff}}$  retrieved from SMART (black triangles) and MODIS (red dots), while the dashed line represents the one-to-one line. There is a robust agreement between in situ  $r_{\text{eff},w}^*$  and retrieved  $r_{\text{eff}}$  with

5 a correlation coefficient  $R^2$  of 0.82. The variability of particle size distributions, the uncertainties of deriving  $r_{\text{eff}}$  from the in situ measurements, the presence of liquid water cloud below cirrus, and the uncertainties caused by the choice of ice crystal shapes for the retrievals are considered as the main contributor to address the discrepancies between in situ  $r_{\text{eff,w}}^*$  and retrieved  $r_{\text{eff}}$ .

## 6 Conclusions

10 Accurate solar radiation measurements are necessary to retrieve high-quality cloud products such as the optical thickness  $\tau$  and particle effective radius  $r_{\text{eff}}$ . Small measurement uncertainties propagate through the retrieval processes. Additional retrieval uncertainties ~~may arise from~~, e.g., arise from the assumption of the surface albedo, ice crystal ~~shapes, and habit~~, cloud vertical profile, ~~as well as and~~ multilayer cloud scenes. Such situations make remote sensing of cloud properties complex and challenging. Collocated airborne and satellite measurements incorporated with in situ observation is one option to assess

15 the uncertainties. Two selected cloud cases, a cirrus above low liquid water clouds and a DCC topped by an anvil cirrus measured during the ML-CIRRUS and ACRIDICON-CHUVA campaigns, are investigated in this study.

Spectral upward radiance measured above the clouds by SMART and MODIS are compared ~~for the cirrus and DCC case~~. Comparisons of spectral upward radiance at  ~~$\lambda$~~  wavelengths between 400 - 1800 nm yield a normalized mean absolute deviation between 0.2 - 7.7% for the cirrus ~~case~~ and 1.5 - 8.3% for the DCC case. The deviation is larger for the DCC case due to

20 the fast cloud evolution, which changes the cloud properties during the time delay between SMART and MODIS observation. A radiance ratio retrieval is applied to ~~simultaneously~~ retrieve  $\tau$  and  $r_{\text{eff}}$  simultaneously. Two combinations, C1 ( $I_{645}^\uparrow$  and  $\mathfrak{R}_1$ ) and C2 ( $I_{645}^\uparrow$  and  $\mathfrak{R}_2$ ) are used in the retrieval algorithm, where  $\mathfrak{R}_1 = I_{1240}^\uparrow/I_{645}^\uparrow$  and  $\mathfrak{R}_2 = I_{1240}^\uparrow/I_{645}^\uparrow$ . By applying the ratios, the measurement uncertainties due to the radiometric calibration of the sensor are reduced. ~~In this way~~ Therefore, the uncertainties of radiance ratio retrieval are smaller compared to the ~~common-usual~~ bi-spectral ~~method~~ technique. Using different

25 near-infrared wavelengths with different absorption by cloud particles in the retrieval algorithm provides  $r_{\text{eff}}$  from different cloud altitudes. ~~A~~ The vertical weighting function shows, that a retrieval using C1 (1240 nm) results in ~~an~~  $r_{\text{eff}}$  from a lower cloud layer, while using C2 (1640 nm) results in ~~an~~  $r_{\text{eff}}$  from a layer ~~closer to~~ which is closer to the cloud top. To some degree, retrievals using these two combinations ~~can~~ give a snapshot of the vertical variation of particle sizes in the cloud. ~~However, it should be noted that retrievals using low absorption wavelength (e.g.,  $\lambda = 1240$  nm) will result in higher retrieval uncertainties.~~

30

The vertical weighting function is used to analyze the impact of ~~the~~ vertical profile assumption ~~on the retrieval in the retrieval~~ of  $r_{\text{eff}}$ . A systematic deviation ~~of the retrieved cloud properties~~ is found between retrievals assuming a vertically homogeneous ~~cloud~~ compared to realistic cloud profiles. For ice clouds with decreasing  $r_{\text{eff}}$  ~~towards~~ towards the cloud top, retrievals assuming a vertically homogeneous cloud ~~will~~ result in an underestimation of  $r_{\text{eff}}$  ~~of~~ up to 1  $\mu\text{m}$ . The impact is larger for retrievals using

35 wavelengths with smaller absorption by cloud particles (e.g.,  $\lambda = 1240$  nm) ~~, when because the~~ lower cloud layers contribute more strongly to the ~~reflected upward~~ radiance. The analysis of vertical weighting function ~~also~~ shows, that each individual cloud layer ~~affects~~ has a contribution to the absorption imprinted in the ~~radiance reflected above cloud top upward radiance~~

with a weighting depending on the cloud profile itself and the chosen wavelength. Therefore, ~~the  $r_{\text{eff}}$  retrieved by this solar remote sensing this has to be kept in mind that the retrieved  $r_{\text{eff}}$  does not represent  $r_{\text{eff}}(z)$   $r_{\text{eff}}$  at a single layer-only cloud layer.~~

5 Instead, the retrieved  $r_{\text{eff}}$  ~~describes-represents~~ a bulk property of the entire cloud layer. The occurrence of ~~liquid water below cirrus significantly-low liquid water cloud~~ leads to an overestimation of the retrieved cirrus  $\tau$ , when the low ~~cloud layer is omitted in the retrieval algorithm. In such conditions, the liquid water cloud cloud is not considered in the forward simulation. The~~ vertical weighting function of the cirrus will change and ~~biases-bias~~ the retrieved cirrus  $r_{\text{eff}}$  ~~if, particularly when~~ the cirrus layer is ~~suffieiently~~ thin ( $\tau < 5$ ). ~~The-In this condition, the~~ radiation is transmitted

10 through the cirrus and reflected by the low liquid water cloud back to the cirrus. Consequently, the absorption ~~of radiation, and the vertical weighting~~ in the lower ~~cloud~~ layers is enhanced. For ~~typical~~ cirrus with decreasing particle size towards the cloud top, the retrieved cirrus  $r_{\text{eff}}$  becomes larger when a liquid water cloud occurs below the cirrus. When the cirrus  $\tau$  is sufficiently thick ( $\tau > 5$ ), the impact ~~is-reduceddecreases~~. The accuracy of the properties of the low liquid water cloud ~~also-strongly~~ determines the uncertainties of the retrieved cirrus properties. Underestimating the liquid water  $\tau$  will artificially increase cirrus

15  $\tau$ . When the liquid water  $r_{\text{eff}}$  ~~is-underestimatedunderestimates the actual value~~, the retrieved cirrus  $r_{\text{eff}}$  becomes larger than in reality. The opposite results are expected when the properties of the low liquid water cloud ~~are-overestimatedoverestimate the actual values~~.

The cloud properties retrieved by SMART and MODIS are compared for the two ~~selected~~ cloud cases. For the cirrus case, the normalized mean absolute deviation ~~of the retrieved cloud properties from SMART and MODIS using combination C1 and C2 is found to be about yields a value of up to~~ 1.2% for  $\tau$  and 2.1% for  $r_{\text{eff}}$ . The deviations are slightly larger than those found in the ~~comparisons-comparison~~ of upward radiance, showing that the errors are only slightly amplified by the non-linearity in the retrieval algorithm. ~~Cirrus-The cirrus~~  $\tau$  derived from the MODIS cloud product overestimate the ~~results from SMART and MODIS retrievals-retrieval results~~ because the MODIS cloud product algorithm assumes only a single cloud layer ~~for their operational retrievals~~. For the DCC case, the deviation is ~~found up to be about in the range of~~ 3.5% for  $\tau$  and 6.5% for  $r_{\text{eff}}$ .

5 In this case, the fast cloud evolution and larger 3-D radiative effects contribute ~~most to the to the deviations and~~ retrieval uncertainties. For both cloud cases, it is found that the particle size decreases towards the cloud top. A higher horizontal variability of  $r_{\text{eff}}$  is observed in the lower cloud layers, while in the upper layers the particle sizes are more homogeneous.

For the cirrus case, the retrieved  $r_{\text{eff}}$  are compared to in situ measurements. To allow the comparison of both methods, the vertical weighting function is considered. Using additional near-infrared wavelengths of SMART and MODIS increases the

10 information on particle size extracted from the spectral measurements and the vertical resolution of retrieved  $r_{\text{eff}}$ . The normalized mean absolute deviation between ~~in situ and retrieved-retrieved and in situ~~  $r_{\text{eff}}$  ranges between 1.5 – 10.3%, which falls within the standard deviation ~~with a correlation coefficient~~. ~~A robust correlation coefficient is obtained with a value~~ of 0.82. The variability of particle size distributions, the uncertainties of deriving  $r_{\text{eff}}$  from the in situ measurements, the presence of liquid water cloud below cirrus, and the uncertainties caused by unconstrained choice of ice crystal ~~shapes-habit~~ for the

15 retrievals are identified as the major contributors which can reveal the discrepancies between ~~in situ and retrieved-retrieved and in situ~~  $r_{\text{eff}}$ . The assumption of vertically homogeneous cloud in the retrieval algorithm has only a small impact on the retrieval results.

*Acknowledgements.* This work was supported by the Max Planck Society (MPG), the German Science Foundation (DFG) funding the SPP HALO 1294 and the grants of WE 1900/35-1 and VO 1504/4-1, the German Aerospace Center (DLR), and the FAPESP (Sao Paulo Research Foundation) grants 2009/15235-8 and 2013/05014-0. The HGF is acknowledged for supports under the contract number W2/W3-60). C. Mahnke and R. Weigel received funding by the German BMBF within the joint ROMIC-project SPITFIRE (01LG1205A). Trismono C. Krisna acknowledges the Ministry of Research, Technology and Higher Education of the Republic of Indonesia (RISTEKDIKTI) and the German Academic Exchange Service (DAAD) for the research grant under the scheme of Indonesia-German Scholarship Programme (IGSP). The entire ML-CIRRUS and ACRIDICON-CHUVA team is gratefully acknowledged for collaboration and support.

## 25 References

- Ackerman, S., Moeller, C., Strabala, K., Gerber, H., Gumley, L., Menzel, W., and Tsay, S.-C.: Retrieval of effective microphysical properties of clouds: A wave cloud case study, *Geophys. Res. Lett.*, 25, 1121–1124, 1998.
- Afchine, A., Rolf, C., Costa, A., Spelten, N., Riese, M., Buchholz, B., Ebert, V., Heller, R., Kaufmann, S., Minikin, A., Voigt, C., Zöger, M., Smith, J., Lawson, P., Lykov, A., Khaykin, S., and Krämer, M.: Ice particle sampling from aircraft – influence of the probing position on the ice water content, *Atmos. Meas. Tech.*, 2017, 1–23, doi:10.5194/amt-2017-373, 2017.
- Anderson, G., Clough, S., Kneizys, F., Chetwynd, J., and Shettle, E.: AFGL Atmospheric Constituent Profiles (0–120 km), Tech. Rep. AFGL-TR-86-0110, AFGL (OPI), Hanscom AFB, MA 01736, 1986.
- Baum, B. A., Yang, P., Heymsfield, A. J., Bansemer, A., Cole, B. H., Merrelli, A., Schmitt, C., and Wang, C.: Ice cloud single-scattering property models with the full phase matrix at wavelengths from 0.2 to 100  $\mu\text{m}$ , *J. Quant. Spectrosc. Radiat. Transfer*, 146, 123 – 139, doi:https://doi.org/10.1016/j.jqsrt.2014.02.029, electromagnetic and Light Scattering by Nonspherical Particles XIV, 2014.
- Baumgardner, D., Strapp, W., and Dye, J. E.: Evaluation of the Forward Scattering Spectrometer Probe. Part II: Corrections for Coincidence and Dead-Time Losses, *J. Atmos. Oceanic Technol.*, 2, 626–632, doi:10.1175/1520-0426(1985)002<0626:EOTFSS>2.0.CO;2, 1985.
- Berendes, T. A., Mecikalski, J. R., MacKenzie, W. M., Bedka, K. M., and Nair, U. S.: Convective cloud identification and classification in daytime satellite imagery using standard deviation limited adaptive clustering, *J. Geophys. Res.*, 113, D20 207, 2008.
- Bierwirth, E.: Airborne measurements of the spectral surface albedo over morocco and its influence on the radiative forcing of saharan dust, Ph.D. thesis, Johannes Gutenberg University Mainz, Germany, 2008.
- Braga, R. C., Rosenfeld, D., Weigel, R., Jurkat, T., Andreae, M. O., Wendisch, M., Pöhlker, M. L., Klimach, T., Pöschl, U., Pöhlker, C., Voigt, C., Mahnke, C., Borrmann, S., Albrecht, R. I., Molleker, S., Vila, D. A., Machado, L. A. T., and Artaxo, P.: Comparing parameterized versus measured microphysical properties of tropical convective cloud bases during the ACRIDICON–CHUVA campaign, *Atmos. Chem. Phys.*, 17, 7365–7386, doi:10.5194/acp-17-7365-2017, 2017.
- Brückner, M., Pospichal, B., Macke, A., and Wendisch, M.: A new multispectral cloud retrieval method for ship-based solar transmissivity measurements, *J. Geophys. Res. Atmos.*, 119, 11.338–11.354, doi:10.1002/2014JD021775, 2014.
- Cecchini, M. A., Machado, L. A. T., Andreae, M. O., Martin, S. T., Albrecht, R. I., Artaxo, P., Barbosa, H. M. J., Borrmann, S., Fütterer, D., Jurkat, T., Mahnke, C., Minikin, A., Molleker, S., Pöhlker, M. L., Pöschl, U., Rosenfeld, D., Voigt, C., Weinzierl, B., and Wendisch, M.: Sensitivities of Amazonian clouds to aerosols and updraft speed, *Atmos. Chem. Phys.*, 17, 10 037–10 050, doi:10.5194/acp-17-10037-2017, 2017.
- Chang, F. L. and Li, Z. Q.: A new method for detection of cirrus overlapping water clouds and determination of their optical properties, *J. Atmos. Sci.*, 62, 3993–4009, 2005.
- Dye, J. E. and Baumgardner, D.: Evaluation of the Forward Scattering Spectrometer Probe. Part I: Electronic and Optical Studies, *J. Atmos. Oceanic Technol.*, 1, 329–344, doi:10.1175/1520-0426(1984)001<0329:EOTFSS>2.0.CO;2, 1984.
- Ehrlich, A., Bierwirth, E., Istomina, L., and Wendisch, M.: Combined retrieval of Arctic liquid water cloud and surface snow properties using airborne spectral solar remote sensing, *Atmos. Meas. Tech.*, 10, 3215–3230, doi:10.5194/amt-10-3215-2017, 2017.
- Eichler, H., Ehrlich, A., Wendisch, M., Mioche, G., Gayet, J.-F., Wirth, M., Emde, C., and Minikin, A.: Influence of ice crystal shape on retrieval of cirrus optical thickness and effective radius: A case study, *J. Geophys. Res.*, 114, D19203, doi:10.1029/2009JD012215, 2009.



- 25 Emde, C., Buras-Schnell, R., Kylling, A., Mayer, B., Gasteiger, J., Hamann, U., Kylling, J., Richter, B., Pause, C., Dowling, T., and Bugliaro, L.: The libRadtran software package for radiative transfer calculations (version 2.0.1), *Geosci. Model Dev.*, 9, 1647–1672, doi:10.5194/gmd-9-1647-2016, 2016.
- Fisher, A.: Cloud and cloud-shadow detection in SPOT5 HRG imagery with automated morphological feature extraction, *Remote Sensing*, 6, 776–800, doi:10.3390/rs6010776, 2014.
- 30 Frey, W., Borrmann, S., Kunkel, D., Weigel, R., de Reus, M., Schlager, H., Roiger, A., Voigt, C., Hoor, P., Curtius, J., Krämer, M., Schiller, C., Volk, C. M., Homan, C. D., Fierli, F., Di Donfrancesco, G., Ulanovsky, A., Ravegnani, F., Sitnikov, N. M., Viciani, S., D’Amato, F., Shur, G. N., Belyaev, G. V., Law, K. S., and Cairo, F.: In situ measurements of tropical cloud properties in the West African Monsoon: Upper tropospheric ice clouds, Mesoscale Convective System outflow, and subvisual cirrus, *Atmos. Chem. Phys.*, 11, 5569–5590, doi:10.5194/acp-11-5569-2011, 2011.
- 35 Fricke, C., Ehrlich, A., Jäkel, E., Bohn, B., Wirth, M., and Wendisch, M.: Influence of local surface albedo variability and ice crystal shape on passive remote sensing of thin cirrus, *Atmos. Chem. Phys.*, 14, 1943–1958, doi:10.5194/acp-14-1943-2014, 2014.
- Fu, Q. and Liou, K.: Parameterization of the radiative properties of cirrus clouds, *J. Atmos. Sci.*, 50, 2008–2025, 1993.
- Gueymard, C. A.: The sun’s total and spectral irradiance for solar energy applications and solar radiation models, *Sol. Energy*, 76, 423–453, 2004.
- Hahn, J., Warren, G., London, J., Chervin, M., and Jenne, R.: Atlas of simultaneous occurrence of different cloud types over land, 1984.
- Heymsfield, A. J., Krämer, M., Luebke, A., Brown, P., Cziczo, D. J., Franklin, C., Lawson, P., Lohmann, U., McFarquhar, G., Ulanowski, Z., and Tricht, K. V.: Cirrus Clouds, *Meteorological Monographs*, 58, 2.1–2.26, doi:10.1175/AMSMONOGRAPHS-D-16-0010.1, 2017.
- 5 Holz, R. E., Platnick, S., Meyer, K., Vaughan, M., Heidinger, A., Yang, P., Wind, G., Dutcher, S., Ackerman, S., Amarasinghe, N., Nagle, F., and Wang, C.: Resolving ice cloud optical thickness biases between CALIOP and MODIS using infrared retrievals, *Atmos. Chem. Phys.*, 16, 5075–5090, doi:10.5194/acp-16-5075-2016, 2016.
- Jäkel, E., Walther, J., and Wendisch, M.: Thermodynamic phase retrieval of convective clouds: impact of sensor viewing geometry and vertical distribution of cloud properties, *Atmos. Meas. Tech.*, 6, 539–547, doi:10.5194/amt-6-539-2013, 2013.
- 10 Jensen, M. P. and Del Genio, A. D.: Radiative and microphysical characteristics of deep convective systems in the tropical western Pacific, *J. Appl. Meteor.*, 42, 1234–1254, 2003.
- Järvinen, E., Schnaiter, M., Mioche, G., Jourdan, O., Shcherbakov, V. N., Costa, A., Afchine, A., Krämer, M., Heidelberg, F., Jurkat, T., Voigt, C., Schlager, H., Nichman, L., Gallagher, M., Hirst, E., Schmitt, C., Bansemer, A., Heymsfield, A., Lawson, P., Tricoli, U., Pfeilsticker, K., Vochezer, P., Möhler, O., and Leisner, T.: Quasi-Spherical Ice in Convective Clouds, *J. Atmos. Sci.*, 73, 3885–3910, doi:10.1175/JAS-D-15-0365.1, 2016.
- 15 Kaufmann, S., Voigt, C., Jeßberger, P., Jurkat, T., Schlager, H., Schwarzenboeck, A., Klingebiel, M., and Thornberry, T.: In situ measurements of ice saturation in young contrails, *Geophys. Res. Lett.*, 41, 702–709, 2014.
- Kaufmann, S., Voigt, C., Jurkat, T., Thornberry, T., Fahey, D. W., Gao, R.-S., Schlage, R., Schäuble, D., and Zöger, M.: The airborne mass spectrometer AIMS – Part 1: AIMS-H<sub>2</sub>O for UTLS water vapor measurements, *Atmos. Meas. Tech.*, 9, 939–953, doi:10.5194/amt-9-939-2016, 2016.
- 20 Kim, D. Y. and Ramanathan, V.: Solar radiation budget and radiative forcing due to aerosols and clouds, *J. Geophys. Res.*, 113, D02 203, 2008.
- King, M., Tsay, S.-C., Platnick, S., Wang, M., and Liou, K.-N.: Cloud retrieval algorithms for MODIS: Optical thickness, effective particle radius, and thermodynamic phase, MODIS Algorithm Theoretical Basis Document, No. ATBD-MOD-05, 1997, 1997.

- 25 King, N. J., Bower, K. N., Crosier, J., and Crawford, I.: Evaluating MODIS cloud retrievals with in situ observations from VOCALS-REx, *Atmos. Chem. Phys.*, 13, 191–209, doi:10.5194/acp-13-191-2013, 2013.
- Klingebiel, M., de Lozar, A., Molleker, S., Weigel, R., Roth, A., Schmidt, L., Meyer, J., Ehrlich, A., Neuber, R., Wendisch, M., and Borrmann, S.: Arctic low-level boundary layer clouds: in situ measurements and simulations of mono- and bimodal supercooled droplet size distributions at the top layer of liquid phase clouds, *Atmos. Chem. Phys.*, 15, 617–631, doi:10.5194/acp-15-617-2015, 2015.
- 30 Korolev, A.: Limitations of the Wegener-Bergeron-Findeisen mechanism in the evolution of mixed-phase clouds, *J. Atmos. Sci.*, 64, 3372–3375, 2007.
- Lance, S., Brock, C. A., Rogers, D., and Gordon, J. A.: Water droplet calibration of the Cloud Droplet Probe (CDP) and in-flight performance in liquid, ice and mixed-phase clouds during ARCPAC, *Atmos. Meas. Tech.*, 3, 1683–1706, doi:10.5194/amt-3-1683-2010, 2010.
- Lane, T. P. and Sharman, R. D.: Intensity of thunderstorm-generated turbulence revealed by large-eddy simulation, *Geophys. Res. Lett.*, 41, 2221–2227, doi:10.1002/2014GL059299, 2014.
- 35 Larar, A. M., Smith, W. L., Zhou, D. K., Liu, X., Revercomb, H., Taylor, J. P., Newman, S. M., and Schlüssel, P.: IASI spectral radiance validation inter-comparisons: case study assessment from the JAIVEx field campaign, *Atmos. Chem. Phys.*, 10, 411–430, doi:10.5194/acp-10-411-2010, 2010.
- Liang, L. S., Di Girolamo, L., and Platnick, S.: View-angle consistency in reflectance, optical thickness and spherical albedo of marine water-clouds over the northeastern Pacific through MISR-MODIS fusion, *Geophys. Res. Lett.*, 36, L09 811, doi:10.1029/2008GL037124, 2009.
- 5 Lindsey, D. T., Hillger, D. W., Grasso, L., Knaff, J. A., and Dostalek, J. F.: GOES climatology and analysis of thunderstorms with enhanced 3.9- $\mu$  m reflectivity, *Mon. Wea. Rev.*, 134, 2342–2353, 2006.
- Liou, K.-N.: Influence of cirrus clouds on weather and climate processes: A global perspective, *Mon. Wea. Rev.*, 114, 1167–1199, 1986.
- Luo, Z. J., Jeyaratnam, J., Iwasaki, S., Takahashi, H., and Anderson, R.: Convective vertical velocity and cloud internal vertical structure: An A-Train perspective, *Geophys. Res. Lett.*, 41, 723–729, doi:10.1002/2013GL058922, 2013GL058922, 2014.
- 10 Lyapustin, A., Wang, Y., Xiong, X., Meister, G., Platnick, S., Levy, R., Franz, B., Korkin, S., Hilker, T., Tucker, J., Hall, F., Sellers, P., Wu, A., and Angal, A.: Scientific impact of MODIS C5 calibration degradation and C6+ improvements, *Atmos. Meas. Tech.*, 7, 4353–4365, doi:10.5194/amt-7-4353-2014, 2014.
- Mayer: Book Review: Barry, R.G., *The Late R. Chorley: Atmosphere, Weather and Climate. Eighth Edition.*, *Meteor. Z.*, 14, 79–80, 2005.
- McFarquhar, G. and Heymsfield, A.: Microphysical characteristics of three anvils sampled during the central equatorial pacific experiment, *J. Atmos. Sci.*, 53, 2401–2423, 1996.
- 15 McFarquhar, G. M. and Heymsfield, A. J.: The definition and significance of an effective radius for ice clouds, *J. Atmos. Sci.*, 55, 2039–2052, 1998.
- Mecikalski, J. R., Feltz, W. F., Murray, J. J., Johnson, D. B., Bedka, K. M., Bedka, S. T., Wimmers, A. J., Pavolonis, M., Berendies, T. A., Haggerty, J., Minnis, P., Bernstein, B., and Williams, E.: Aviation applications for satellite-based observations of cloud properties, convection initiation, in-flight icing, turbulence, and volcanic ash, *Bull. Amer. Meteor. Soc.*, 88, 1589–+, 2007.
- 20 Molleker, S., Borrmann, S., Schlager, H., Luo, B., Frey, W., Klingebiel, M., Weigel, R., Ebert, M., Mitev, V., Matthey, R., Woiwode, W., Oelhaf, H., Dörnbrack, A., Stratmann, G., Groöß, J.-U., Günther, G., Vogel, B., Müller, R., Krämer, M., Meyer, J., and Cairo, F.: Microphysical properties of synoptic-scale polar stratospheric clouds: in situ measurements of unexpectedly large HNO<sub>3</sub>-containing particles in the Arctic vortex, *Atmos. Chem. Phys.*, 14, 10 785–10 801, doi:10.5194/acp-14-10785-2014, 2014.

- 25 Mu, Q., Wu, A., Xiong, X., Doelling, D. R., Angal, A., Chang, T., and Bhatt, R.: Optimization of a Deep Convective Cloud Technique in Evaluating the Long-Term Radiometric Stability of MODIS Reflective Solar Bands, *Remote Sensing*, 9, 535, doi:10.3390/rs9060535, 2017.
- Nakajima, T. and King, M.: Determination of the optical thickness and effective particle radius of clouds from reflected solar radiation measurements. Part I: Theory, *J. Atmos. Sci.*, 47, 1878–1893, 1990.
- 30 Painemal, D. and Zuidema, P.: Assessment of MODIS cloud effective radius and optical thickness retrievals over the Southeast Pacific with VOCALS-REx in situ measurements, *J. Geophys. Res.*, 116, doi:10.1029/2011JD016155, d24206, 2011.
- Platnick, S.: Vertical photon transport in cloud remote sensing problems, *J. Geophys. Res.*, 105, 22 919–22 935, 2000.
- Platnick, S., King, M. D., Ackerman, S. A., Menzel, W. P., Baum, B. A., Riédi, J. C., and Frey, R. A.: The MODIS cloud products: Algorithms and examples from Terra, *IEEE Trans. Geosci. Remote Sens.*, 41, 459–473, 2003.
- 35 Platnick, S., Meyer, K. G., King, M. D., Wind, G., Amarasinghe, N., Marchant, B., Arnold, G. T., Zhang, Z., Hubanks, P. A., Holz, R. E., Yang, P., Ridgway, W. L., and Riedi, J.: The MODIS Cloud Optical and Microphysical Products: Collection 6 Updates and Examples From Terra and Aqua, *IEEE Trans. Geosci. Remote Sens.*, 55, 502–525, doi:10.1109/TGRS.2016.2610522, 2017.
- Rolland, P. and Liou, K.: Surface variability effects on the remote sensing of thin cirrus optical and microphysical properties, *J. Geophys. Res.*, 106, 22 965–22 977, doi:10.1029/2001JD900160, 2001.
- Rosenfeld, D. and Lensky, I. M.: Satellite-based insights into precipitation formation processes in continental and maritime convective clouds, *Bull. Amer. Meteor. Soc.*, 79, 2457–2476, 1998.
- Rozanov, V. and Kokhanovsky, A.: Semianalytical cloud retrieval algorithm as applied to the top altitude and the cloud geomet-
- 5 rical thickness determination from top-of-atmosphere reflectance measurements in the oxygen A band, *J. Geophys. Res.*, 109, doi:10.1029/2003JD004 104, 2004.
- Savtchenko, A., Kummerer, R., Smith, P., Gopalan, A., Kempler, S., and Leptoukh, G.: A-Train Data Depot: Bringing Atmospheric Measurements Together, *IEEE Transactions on Geoscience and Remote Sensing*, 46, 2788–2795, doi:10.1109/TGRS.2008.917600, 2008.
- Schumacher, C., Stevenson, S. N., and Williams, C. R.: Vertical motions of the tropical convective cloud spectrum over Darwin, Australia,
- 10 *Quart. J. Roy. Meteor. Soc.*, 141, 2277–2288, doi:10.1002/qj.2520, 2015.
- Schumann, U., Baumann, R., Baumgardner, D., Bedka, S. T., Duda, D. P., Freudenthaler, V., Gayet, J.-F., Heymsfield, A. J., Minnis, P., Quante, M., Raschke, E., Schlager, H., Vázquez-Navarro, M., Voigt, C., and Wang, Z.: Properties of individual contrails: a compilation of observations and some comparisons, *Atmos. Chem. Phys.*, 17, 403–438, doi:10.5194/acp-17-403-2017, 2017.
- Sherwood, S. C., Minnis, P., and McGill, M.: Deep convective cloud-top heights and their thermodynamic control during CRYSTAL-FACE,
- 15 *J. Geophys. Res.*, 109, doi:10.1029/2004JD004811, d20119, 2004.
- Shettle, E.: Comments on the use of LOWTRAN in transmission calculations for sites with the ground elevated relative to sea level, *Appl. Opt.*, 28, 1451–1452, 1989.
- Shupe, M. D. and Intrieri, J. M.: Cloud radiative forcing of the Arctic surface: The influence of cloud properties, surface albedo, and solar zenith angle, *J. Climate*, 17, 616–628, 2004.
- 20 Slingo, A.: Sensitivity of the Earths Radiation Budget to Changes in Low Clouds, *Nature*, 343, 49–51, 1990.
- Sohn, B.-J., Choi, M.-J., and Ryu, J.: Explaining darker deep convective clouds over the western Pacific than over tropical continental convective regions, *Atmos. Meas. Tech.*, 8, 4573–4585, doi:10.5194/amt-8-4573-2015, 2015.
- Sourdeval, O., C-Labonnote, L., Baran, A. J., and Brogniez, G.: A methodology for simultaneous retrieval of ice and liquid water cloud properties. Part I: Information content and case study, *Quart. J. Roy. Meteor. Soc.*, 141, 870–882, doi:10.1002/qj.2405, 2015.

- 25 Stamnes, K., Tsay, S.-C., Wiscombe, W., and Laszlo, I.: DISORT, a General-Purpose Fortran Program for Discrete-Ordinate-Method Radiative Transfer in Scattering and Emitting Layered Media: Documentation of Methodology, Tech. rep., Dept. of Physics and Engineering Physics, Stevens Institute of Technology, Hoboken, NJ 07030, 2000.
- Stephens, G. L. and Kummerow, C. D.: The remote sensing of clouds and precipitation from space: A review, *J. Atmos. Sci.*, 64, 3742–3765, 2007.
- 30 Strahler, A., Muller, J., and Members, M. S. T.: MODIS BRDF/Albedo product: Algorithm Theoretical Basis Document Version 5.0, 1999.
- Stubenrauch, C., Rossow, W., Kinne, S., Ackerman, S., Cesana, G., Chepfer, H., Di Girolamo, L., Getzewich, B., Guignard, A., Heidinger, A., et al.: Assessment of global cloud datasets from satellites: Project and database initiated by the GEWEX radiation panel, *Bull. Amer. Meteor. Soc.*, 94, 1031–1049, 2013.
- Sun, J.-Q., Xiong, X., Barnes, W. L., and Guenther, B.: MODIS reflective solar bands on-orbit lunar calibration, *IEEE Trans. Geosci. Remote Sens.*, 45, 2383–2393, 2007.
- 35 van Diedenhoven, B., Fridlind, A. M., Cairns, B., and Ackerman, A. S.: Variation of ice crystal size, shape, and asymmetry parameter in tops of tropical deep convective clouds, *J. Geophys. Res.*, 119, 11,809–11,825, doi:10.1002/2014JD022385, 2014.
- van Diedenhoven, B., Fridlind, A. M., Cairns, B., Ackerman, A. S., and Yorks, J. E.: Vertical variation of ice particle size in convective cloud tops, *Geophys. Res. Lett.*, 43, 4586–4593, doi:10.1002/2016GL068548, 2016GL068548, 2016.
- Voigt, C., Schlager, H., Ziereis, H., Kärcher, B., Luo, B., Schiller, C., Krämer, M., Popp, P., Irie, H., and Kondo, Y.: Nitric acid in cirrus clouds, *Geophys Res Lett*, 33, 2006.
- 5 Voigt, C., Jessberger, P., Jurkat, T., Kaufmann, S., Baumann, R., Schlager, H., Bobrowski, N., Giuffrida, G., and Salerno, G.: Evolution of CO<sub>2</sub>, SO<sub>2</sub>, HCl, and HNO<sub>3</sub> in the volcanic plumes from Etna, *Geophys. Res. Lett.*, 41, 2196–2203, 2014.
- Voigt, C., Schumann, U., Minikin, A., Abdelmonem, A., Afchine, A., Borrmann, S., Boettcher, M., Buchholz, B., Bugliaro, L., Costa, A., Curtius, J., Dollner, M., Dörnbrack, A., Dreiling, V., Ebert, V., Ehrlich, A., Fix, A., Forster, L., Frank, F., Fütterer, D., Giez, A., Graf, K., Groß, J.-U., Groß, S., Heimerl, K., Heinold, B., Hüneke, T., Järvinen, E., Jurkat, T., Kaufmann, S., Kenntner, M., Klingebiel, M., 10 Klimach, T., Kohl, R., Krämer, M., Krisna, T. C., Luebke, A., Mayer, B., Mertes, S., Molleker, S., Petzold, A., Pfeilsticker, K., Port, M., Rapp, M., Reutter, P., Rolf, C., Rose, D., Sauer, D., Schäfler, A., Schlage, R., Schnaiter, M., Schneider, J., Spelten, N., Spichtinger, P., Stock, P., Walser, A., Weigel, R., Weinzierl, B., Wendisch, M., Werner, F., Wernli, H., Wirth, M., Zahn, A., Ziereis, H., and Zöger, M.: ML-CIRRUS: The Airborne Experiment on Natural Cirrus and Contrail Cirrus with the High-Altitude Long-Range Research Aircraft HALO, *Bull. Amer. Meteor. Soc.*, 98, 271–288, doi:10.1175/BAMS-D-15-00213.1, 2017.
- 15 Wan, Z.: New refinements and validation of the collection-6 {MODIS} land-surface temperature/emissivity product, *Remote Sensing of Environment*, 140, 36 – 45, doi:10.1016/j.rse.2013.08.027, 2014.
- Wang, L., Qu, J. J., Xiong, X., Hao, X., Xie, Y., and Che, N.: A new method for retrieving band 6 of aqua MODIS, *IEEE Geoscience and Remote Sensing Letters*, 3, 267–270, doi:10.1109/LGRS.2006.869966, 2006.
- Wang, X., Liou, K. N., Ou, S. S. C., Mace, G. G., and Deng, M.: Remote sensing of cirrus cloud vertical size profile using MODIS data, *J. Geophys. Res.*, 114, doi:10.1029/2008JD011327, d09205, 2009.
- 20 Warren, S., Hahn, C., and London, J.: Simultaneous occurrence of different cloud types, *J. Climate Appl. Meteor.*, 24, 658–667, 1985.
- Weigel, K., Rozanov, A., Azam, F., Bramstedt, K., Damadeo, R., Eichmann, K.-U., Gebhardt, C., Hurst, D., Kraemer, M., Lossow, S., Read, W., Spelten, N., Stiller, G. P., Walker, K. A., Weber, M., Bovensmann, H., and Burrows, J. P.: UTLS water vapour from SCIAMACHY limb measurements V3.01 (2002–2012), *Atmos. Meas. Tech.*, 9, 133–158, doi:10.5194/amt-9-133-2016, 2016.

- 25 Wendisch, M. and Brenguier, J.-L.: *Airborne Measurements for Environmental Research – Methods and Instruments*, Wiley–VCH Verlag GmbH & Co. KGaA, Weinheim, Germany, Weinheim, Germany, ISBN: 978-3-527-40996-9, 2013.
- Wendisch, M. and Mayer, B.: Vertical distribution of spectral solar irradiance in the cloudless sky: A case study, *Geophys. Res. Lett.*, 30, 1183–1186, doi:10.1029/2002GL016529, 2003.
- Wendisch, M., Heintzenberg, J., and Bussemer, M.: Measurement-based aerosol forcing calculations: the influence of model complexity, *Meteor. Z.*, 10, 45–60, 2001.
- Wendisch, M., Pilewskie, P., Jäkel, E., Schmidt, S., Pommier, J., Howard, S., Jonsson, H. H., Guan, H., Schröder, M., and Mayer, B.: Airborne measurements of areal spectral surface albedo over different sea and land surfaces, *J. Geophys. Res.*, 109, Art. No. D08 203, doi:10.1029/2003JD004392, 2004.
- Wendisch, M., Pilewskie, P., Pommier, J., Howard, S., Yang, P., Heymsfield, A. J., Schmitt, C. G., Baumgardner, D., and Mayer, B.: Impact of cirrus crystal shape on solar spectral irradiance: A case study for subtropical cirrus, *J. Geophys. Res.*, 110, doi:10.1029/2004JD005294, d03202, 2005.
- Wendisch, M., Yang, P., and Pilewskie, P.: Effects of ice crystal habit on thermal infrared radiative properties and forcing of cirrus, *J. Geophys. Res.*, 112, D03 202, doi:10.1029/2006JD007899, 2007.
- Wendisch, M., Pöschl, U., Andreae, M. O., Machado, L. A. T., Albrecht, R., Schlager, H., Rosenfeld, D., Martin, S. T., Abdelmonem, A., Afchine, A., Araujo, A., Artaxo, R., Aufmhoff, H., Barbosa, H. M. J., Borrmann, S., Braga, R., Buchholz, B., Cecchini, M. A., Costa, A., Curtius, J., Dollner, M., Dorf, M., Dreiling, V., Ebert, V., Ehrlich, A., Ewald, F., Fisch, G., Fix, A., Frank, F., Fütterer, D., Heckl, C., Heidelberg, F., Hüneke, T., Jäkel, E., Järvinen, E., Jurkat, T., Kanter, S., Kästner, U., Kenntner, M., Kesselmeier, J., Klimach, T., Knecht, M., Kohl, R., Kölling, T., Krämer, M., Krüger, M., Krisna, T. C., Lavric, J. V., Longo, K., Mahnke, C., Manzi, A. O., Mayer, B., Mertes, S., Minikin, A., Molleker, S., Münch, S., Nillius, B., Pfeilsticker, K., Pöhlker, C., Roiger, A. E., Rose, D., Rosenow, D., Sauer, D., Schnaiter, M., Schneider, J., Schulz, C., de Souza, R. A. F., Spanu, A., Stock, P., Vila, D., Voigt, C., Walser, A., Walter, D., Weigel, R., Weinzierl, B., Werner, R., Yamasoe, M. A., Ziereis, H., Zinner, T., and Zöger, M.: The ACRIDICON-CHUVA campaign: Studying tropical deep convective clouds and precipitation over Amazonia using the new German research aircraft HALO, *Bull. Am. Meteorol. Soc.*, doi:10.1175/BAMS-D-14-00255.1, 2016.
- Wendisch, M., Brückner, M., Burrows, J. P., Crewell, S., Dethloff, K., Ebell, K., Lüpkes, C., Macke, A., Notholt, J., Quaas, J., Rinke, A., and Tegen, I.: Understanding causes and effects of rapid warming in the Arctic, *Eos*, 98, doi:10.1029/2017EO064803, 2017.
- 880 Werner, F., Siebert, H., Pilewskie, P., Schmeissner, T., Shaw, R. A., and Wendisch, M.: New airborne retrieval approach for trade wind cumulus properties under overlying cirrus, *J. Geophys. Res. Atmos.*, 118, 3634–3649, doi:10.1002/jgrd.50334, 2013.
- Wind, G., Platnick, S., King, M. D., Hubanks, P. A., Pavolonis, M. J., Heidinger, A. K., Yang, P., and Baum, B. A.: Multilayer Cloud Detection with the MODIS Near-Infrared Water Vapor Absorption Band, *J. Appl. Meteorol. Climatol.*, 49, 2315–2333, doi:10.1175/2010JAMC2364.1, 2010.
- 885 Wiscombe, W.: Improved Mie scattering algorithms, *Appl. Opt.*, 19, 1505–1509, 1980.
- Wolf, K., Ehrlich, A., Hüneke, T., Pfeilsticker, K., Werner, F., Wirth, M., and Wendisch, M.: Potential of remote sensing of cirrus optical thickness by airborne spectral radiance measurements at different sideward viewing angles, *Atmos. Chem. Phys.*, 17, 4283–4303, doi:10.5194/acp-17-4283-2017, 2017.
- Wylie, D., Jackson, D. L., Menzel, W. P., and Bates, J. J.: Trends in global cloud cover in two decades of HIRS observations, *J. Climate*, 18, 890 3021–3031, 2005.

- Xiong, X., Sun, J., Chiang, K., Xiong, S., and Barnes, W.: MODIS on-orbit characterization using the moon, *Sensors, Systems And Next-Generation Satellites Vi*, 4881, 299–307, 2003.
- Xiong, X. X. and Barnes, W.: An overview of MODIS radiometric calibration and characterization, *Adv. Atmos. Sci.*, 23, 69–79, 2006.
- 895 Yang, P., Bi, L., Baum, B. A., Liou, K.-N., Kattawar, G. W., Mishchenko, M. I., and Cole, B.: Spectrally consistent scattering, absorption, and polarization properties of atmospheric ice crystals at wavelengths from 0.2 to 100  $\mu\text{m}$ , *J. Atmos. Sci.*, 70, 330–347, doi:10.1175/JAS-D-12-039.1, 2013.
- Zhang, S., Xue, H., and Feingold, G.: Vertical profiles of droplet effective radius in shallow convective clouds, *Atmos. Chem. Phys.*, 11, 4633–4644, doi:10.5194/acp-11-4633-2011, 2011.
- 900 Zhang, Z. and Platnick, S.: An assessment of differences between cloud effective particle radius retrievals for marine water clouds from three MODIS spectral bands, *J. Geophys. Res.*, 116, doi:10.1029/2011JD016216, d20215, 2011.
- Zhang, Z. B., Platnick, S., Yang, P., Heidinger, A. K., and Comstock, J. M.: Effects of ice particle size vertical inhomogeneity on the passive remote sensing of ice clouds, *J. Geophys. Res.*, 115, D17 203, doi:10.1029/2010JD013835, 2010.