## Effect assessment of NO<sub>x</sub> and SO<sub>2</sub> control policies on acid components in precipitation from 2005

## to 2016 in China, based on satellite monitoring

Xiuying Zhang <sup>1,\*</sup>, Dongmei Chen <sup>2,3</sup>, Lei Liu <sup>1,4</sup>, Limin Zhao <sup>1</sup>, Wuting Zhang <sup>1</sup>


<sup>&</sup>lt;sup>1</sup> International Institute for Earth System Science, Nanjing University, Nanjing 210023, China

<sup>&</sup>lt;sup>2</sup> Department of Geography and Planning, Queen's University, Kingston, ONK7L 3N6, Canada


<sup>&</sup>lt;sup>3</sup> School of Geography and Remote Sensing, Nanjing University of Information Science & Technology, Nanjing, China

<sup>&</sup>lt;sup>4</sup> Jiangsu Center for Collaborative Innovation in Geographical Information Resource Development and Application, Nanjing 210023, China

<sup>\*</sup> Corresponding authors: Xiuying Zhang (<u>lzhxy77@163.com</u>)



**Fig.S1** Trend of percentage of urban with acid rains (%) from 2005 to 2016 (China Environmental Bulletin 2005 - 2016)



**Fig. S2** Spatial distribution of precipitation pH in 2005 and 2016 (China Environmental Bulletin 2005 and 2016)