

Dear Co-editor,

We thank for your technical comments. In this revised manuscript we have taken them into consideration, updated the affiliations and reference list and checked our spelling. The changes are highlighted.

A point-by-point reply to the comments:

5 **On page 16 line 5 you refer to "dynamic model" but I believe you had changed this to "kinetic model" elsewhere.**
This is corrected to "kinetic model".

On page 16 line 30-32. I wonder whether you can state more directly that this implies that k_{dec} dominate k_{loss} . It may be confusing to readers to have a term that contains the concentration of water but than state that water concentration is irrelevant.

10 This is a very good point. We added "under these conditions thermal decomposition dominates the loss mechanism of sCI and the reaction with water vapour is less important."

On page 25 line 28. I believe you are missing "the" between "that" and "condensation sink".

The is added.

15 **On p.26 line 29-30 (and page 28 line 27) . Could you provide a mechanistic suggestion why the SCI yield would depend on the SCI concentration or does this reflect experimental uncertainties?**

We added a sentence "The higher yield in the experiment with low SO_2 concentration is still within the experimental uncertainties." to clarify the situation.

Best regards,

Nina Sarnela on behalf of the authors

20

25

30

35

Measurement-model comparison of stabilized Criegee Intermediate and Highly Oxygenated Molecule production in the CLOUD chamber

Nina Sarnela¹, Tuija Jokinen¹, Jonathan Duplissy¹, Chao Yan¹, Tuomo Nieminen², Mikael
5 Ehni¹, Siegfried Schobesberger^{1,2,3}, Martin Heinritzi⁴, Sebastian Ehrhart^{4,a}, Katrianne
Lehtipalo^{1,5}, Jasmin Tröstl⁵, Mario Simon⁴, Andreas Kürten⁴, Markus Leiminger⁶, Michael J.
Lawler⁷, Matti P. Rissanen¹, Federico Bianchi¹, Arnaud P. Praplan⁸, Jani Hakala¹, Antonio
Amorim⁹, Marc Gonin¹⁰, Armin Hansel⁶, Jasper Kirkby^{4,11}, Josef Dommen⁵, Joachim Curtius⁴,
10 James N. Smith⁷, Tuukka Petäjä¹, Douglas R. Worsnop^{1,12}, Markku Kulmala¹, Neil M.
Donahue^{13,1} and Mikko Sipilä¹

¹ Institute for Atmospheric and Earth System Research / Physics, Faculty of Science, University of Helsinki, Box 64, 00014 Helsinki, Finland

²University of Eastern Finland, Department of Applied Physics, PO Box 1627, FI-70211 Kuopio, Finland

³Department of Atmospheric Sciences, University of Washington, 408 ATG Bldg, Box 351640, Seattle, WA 98195, USA

⁴Institute for Atmospheric and Environmental Sciences, Goethe University Frankfurt, Altenhöferallee 1, 60438 Frankfurt am Main, Germany

⁵Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, 5232 Villigen PSI, Switzerland

⁶University of Innsbruck, Institute for Ion Physics and Applied Physics, Technikerstraße 25, 6020 Innsbruck, Austria

⁷University of California, Irvine, Department of Chemistry, Irvine, CA 92697, USA

⁸Finnish Meteorological Institute, P.O. Box 503, 00101 Helsinki, Finland

⁹CENTRA, Faculdade de Ciencias da Universidade de Lisboa

¹⁰Tofwerk AG, 3600 Thun, Switzerland

¹¹CERN, CH-1211 Geneva, Switzerland

¹²Aerodyne Research, Inc., Billerica, MA 01821, USA

¹³Carnegie Mellon University Center for Atmospheric Particle Studies, 5000 Forbes Ave, Pittsburgh Pennsylvania, 15213, USA

^anow at: Max-Planck Institute of Chemistry, Atmospheric Chemistry Department, Hahn-Meitner-Weg 1, 55128 Mainz, Germany

Correspondence to: Nina Sarnela (nina.sarnela@helsinki.fi)

Abstract. Atmospheric oxidation is an important phenomenon, which produces large quantities of low-volatile compounds such as sulfuric acid and oxidised organic compounds. Such species may be involved in nucleation of particles and enhance their subsequent growth to reach the size of cloud condensation nuclei (CCN). In this study, we investigate α -pinene, the most abundant monoterpene globally, and its oxidation products formed through ozonolysis in the Cosmic Leaving OUtdoors Droplets (CLOUD) chamber at CERN (the European Organization for Nuclear Research). By scavenging hydroxyl radicals (OH) with hydrogen (H₂), we were able to investigate the formation of Highly Oxygenated Molecules (HOM) purely driven by ozonolysis, and study the oxidation of sulfur dioxide (SO₂) driven by stabilized Criegee Intermediates (sCI). We measured the concentrations of HOM and sulfuric acid with a chemical ionization atmospheric pressure interface time-of-flight (CI-APi-TOF) mass

spectrometer and compared the measured concentrations with simulated concentrations calculated with a kinetic model. We found molar yields in the range of 3.5 – 6.5% for HOM formation and 22 - 32% for the formation of stabilized Criegee Intermediates by fitting our model to the measured sulfuric acid concentrations. The simulated time evolution of the ozonolysis products was in good agreement with measured concentrations except that in some of the experiments sulfuric acid formation was faster than simulated. In those experiments the simulated and measured concentrations met when the concentration reached a plateau but the plateau was reached 20-50 minutes later in the simulations. The results shown here are consistent with the recently published yields for HOM formation from different laboratory experiments. Together with the sCI yields, these results help to understand atmospheric oxidation processes better and make the reaction parameters more comprehensive for broader use.

10 1. Introduction

Atmospheric new-particle formation begins when trace gases form small molecular clusters, which can grow to larger sizes through the condensation of vapours. When they have reached a large enough diameter, these particles can act as a cloud and ice condensation nuclei that may affect the optical properties of clouds or have other effects on climate and air quality such as decrease in visibility. A lot of effort have been put into identifying the vapours responsible for nucleation and growth of the particles. Various studies have identified sulfuric acid and low volatility organic compounds as the key compounds in atmospheric new particle formation (Bianchi et al., 2016; Kirkby et al., 2016; Riccobono et al., 2014; Riipinen et al., 2011; Sihto et al., 2006; Tröstl et al., 2016; Weber et al., 1997; Wehner et al., 2005). Other important aerosol precursors identified in several laboratory studies include ammonia and amines (Almeida et al., 2013; Ball et al., 1999; Murphy et al., 2007). Laboratory measurements (Almeida et al., 2013; Berndt et al., 2010, 2014b; Jen et al., 2014; Kirkby et al., 2011) and computational studies (Kurtén et al., 2008; Paasonen et al., 2012) suggest that ammonia and amines can enhance particle formation but it is challenging to estimate their importance in the ambient atmosphere without comprehensive measurements of their concentration in the atmosphere. Neutral sulfuric acid – amine clusters have been observed in the CLOUD chamber experiments (Kürten et al., 2014) but similar neutral clusters have not yet been detected in the atmosphere. Field studies suggest that iodine oxides could be the key compounds for new particle formation in coastal areas during periods when high tidal movements expose algae beds to sunlight (O'Dowd et al., 2002; Sipilä et al., 2016). However, these iodine oxides do not appear as abundantly in the atmosphere as sulfuric acid or low volatile organic vapours, so their importance seem to be limited to coastal areas.

Sulfuric acid is linked with new particle formation events all around the world (Kulmala et al., 2004). Gas phase sulfuric acid was previously thought to be formed solely via OH-radical oxidation of sulfur dioxide (SO_2) and dimethylsulfide (Lucas and Prinn, 2005). However, stabilized Criegee Intermediates (sCI), formed in a reaction between unsaturated hydrocarbons and ozone, are also capable of oxidising SO_2 into sulfuric acid. While sCI's (Chuong et al., 2004; Donahue et al., 2011; Drozd and Donahue, 2011) and their reactions with SO_2 (Cox and Penkett, 1971) have been investigated for decades, the potential of the atmospheric relevance of sCI+ SO_2 was demonstrated more recently (Berndt et al., 2012; Mauldin III et al., 2012; Welz et al., 2012). In the reaction between ozone and alkenes, a primary ozonide is formed which decomposes quickly to a carbonyl and a carbonyl oxide known as the Criegee Intermediate (Criegee, 1975). A fraction of the Criegee Intermediates can be collisionally stabilized to form sCI (Donahue et al., 2011; Herron et al., 1982). In the case of α -pinene and other endocyclic alkenes, both functional groups—the carbonyl and Criegee Intermediate—remain in the same

molecule. Recent studies indicate that the sCI can have a significant role in ambient sulfuric acid formation (Boy et al., 2013; Welz et al., 2012; Yao et al., 2014). Determining the reaction rate constants for sCI + SO₂ reactions has been challenging and the previous estimates have varied considerably due to the lack of direct measurements of sCI compounds (Johnson and Marston, 2008). Recent studies (Berndt et al., 2012; Mauldin III et al., 2012; Welz et al., 2012) with new experimental methods have shown up to four orders of magnitude higher reaction rate constants for the reaction between a sCI and SO₂ compared to previous estimates. Also differences between the reactivity of sCI derived from different alkenes and their reactivity towards SO₂, water and several other atmospheric compounds has been emphasized (Berndt et al., 2014a; Sipilä et al., 2014; Taatjes et al., 2013; Vereecken et al., 2012, 2014).

10 The other important reaction pathway associated with ozonolysis of alkenes, such as α -pinene, is the formation of Highly Oxygenated Molecules (HOM, Ehn et al., 2014; Kirkby et al., 2016; Tröstl et al., 2016). Crounse et al. (2013) suggested that autoxidation, in which the radicals produced after the initial oxidation are oxidised by atmospheric oxygen, plays an important role in the atmospheric oxidation of organic compounds. Organic radicals, including radicals formed when Criegee Intermediates decompose, will react with molecular oxygen (O₂) to form
15 a peroxy radical (RO₂). The RO₂ can subsequently undergo an intramolecular H-shift, which will be followed by subsequent O₂ addition to form a more oxidised RO₂. According to the mechanism introduced by Ehn et al. (2014), the RO₂ can undergo several additional reactions with O₂, which eventually leads to the formation of HOM, also referred to as Extremely Low Volatility Organic Compounds (ELVOC, Donahue et al., 2012; Ehn et al., 2014; Jokinen et al., 2015) or Highly Oxidised Multifunctional organic compound (HOM, Ehn et al., 2012). Here we
20 call them HOM, as it was recently recognized, that not all HOM necessarily are extremely low volatile (Tröstl et al., 2016). The RO₂ can also react with nitrogen monoxide (NO), hydroperoxyl radical (HO₂) or another RO₂, which can terminate the autoxidation reaction chain and form a closed shell product. The molar yield of HOM formed from α -pinene and ozone is reported to be around 3-7% (Ehn et al., 2014; Jokinen et al., 2015; Kirkby et al., 2016).
25 In Earth's atmosphere α -pinene is the most abundant monoterpene having yearly emissions of 50 Tg globally (Guenther et al., 1995; Seinfeld and Pankow, 2003) and around 80% of the emitted α -pinene undergo oxidation via ozonolysis (Griffin et al., 1999). The high yields of HOM acting as condensing vapours can explain a large portion of the formed secondary organic aerosol (SOA) at least in forested regions. At atmospheric pressure, ozonolysis of the endocyclic α -pinene generates sCI with a low but important yield, measured to be around 15%
30 (15% Drozd and Donahue, 2011), 15% \pm 7% (Sipilä et al., 2014)).

In this study we conducted pure ozonolysis experiments in which OH was removed by a scavenger in the CLOUD chamber facility at CERN (Kirkby et al., 2011; Duplissy et al., 2016) during the CLOUD7 campaign in fall 2012. We examined the formation of sulfuric acid originating from Criegee Intermediate oxidation and of HOM from α -pinene oxidation and compared the temporal trends of the measured to the modelled concentrations. The modelling of HOM concentration was based on the experimental yield terms obtained from recent studies by Ehn et al. (2014) and Jokinen et al. (2015) while the sulfuric acid concentration was modelled using the reaction coefficient and the yield term from the study by Sipilä et al. (2014). In addition, we calculated the yield terms for sCI and HOM formation in the CLOUD experiments by fitting our model to the measured sulfuric acid and HOM concentrations.

2. Methods

2.1 Experiments

We conducted the experiments in the CLOUD chamber, which is a 26.1 m^3 electro-polished stainless steel cylinder at CERN (Geneva, Switzerland, Kirkby et al., 2011; Duplissy et al., 2016). We compared our results to previous 5 experiments of α -pinene ozonolysis conducted in the Tropos Laminar Flow Tube (Berndt et al., 2005) and the Jülich Plant Atmosphere Chamber facility (Mentel et al., 2009) (Table 1). In contrast to these experimental facilities, the CLOUD chamber has a smaller wall loss rate (e.g. around $1.8 \times 10^{-3} \text{ s}^{-1}$ for sulfuric acid), which is similar to the condensation sink in relatively unpolluted ambient environments. This feature allows us to investigate nucleation and growth processes with precursors at atmospherically relevant concentrations.

10 **Table 1. Description of different experimental systems compared in this study**

| | TROPOS-LFT | JPAC | CLOUD |
|----------------------------|--|--|---|
| | Tropos Laminar Flow Tube, Leipzig, Germany (Berndt et al., 2005) | Jülich Plant Atmosphere Chamber, Jülich, Germany (Mentel et al., 2009) | The Cosmics Leaving Outdoor Droplets, Geneva, Switzerland (Kirkby et al., 2011) |
| Description | Laminar flow glass tube with 40s residence time | Borosilicate glass chamber with 45min residence time | Stainless steel chamber with 3h residence time |
| Volume | 0.025 m^3 | 1.45 m^3 | 26.1 m^3 |
| Temperature | 293 K | 289 K | 278 K |
| RH | 25-50% | 63% | 38% |
| Wall loss | 10-27% | $11 \times 10^{-3} \text{ s}^{-1}$ | $1.2-1.9 \times 10^{-3} \text{ s}^{-1}$ |
| Scavenger (for OH) | H ₂ , propane | CO | H ₂ |

For this study, only α -pinene ozonolysis experiments fulfilling certain conditions were selected:

15

- Only ozone, α -pinene and sulfur dioxide were added as precursors to the chamber
- H₂ was used as OH scavenger
- Ions were constantly removed from the chamber (i.e. neutral conditions)
- 38% relative humidity and 278 K temperature

We used two electrodes operating at voltages of $\pm 30 \text{ kV}$ inside the chamber to produce an electric field of 20 kV/m throughout the chamber which removed all the ions in order to maintain neutral conditions. All the 20 experiments were done at 278 K and the thermal insulation kept the temperature stable within 0.05 K (Duplissy et al., 2016). The relative humidity was kept at 38% during all the experiments. The synthetic air used in the chamber was provided from cryogenic liquid N₂ and O₂ (79:21, volume ratio) and 0.1% of H₂ was added to the air to scavenge all the hydroxyl radicals (OH) and prevent any OH-initiated reactions. The ozone mixing ratio was kept around 22 ppbv in all the experiments. Sulfur dioxide was added to the chamber at a mixing ratio around 70 ppbv 25

in four experiments and at a mixing ratio of 17 ppbv in one experiment. α -Pinene was supplied with mixing ratios varying between 80 pptv and 600 pptv from a temperature controlled evaporator using N_2 as a carrier gas. Two counter-rotating stainless steel fans are mounted inside the chamber to achieve efficient turbulent mixing of the gases and ions (Voigtländer et al., 2012). The total flow through the chamber is kept constant during the experiments.

We started the ozonolysis experiments with a constant concentration of SO_2 , O_3 and H_2 in the chamber (background measurement). Then we injected α -pinene into the chamber with a constant flow rate during the whole experiment (4-7 hours). In between the experiments, the chamber was cleaned by closing the α -pinene flow and flushing the chamber with pure air (mixture of evaporated liquid nitrogen and liquid oxygen). All particles formed were removed by repeated charging the particles and applying the high-voltage clearing electric field inside the chamber. The conditions of each experiment are shown in Table 2.

Table 2: The measured concentrations of precursor vapours (ozone, α -pinene and sulfur dioxide), formation rates at 2.5 nm, growth rates of sub-3nm particles and calculated yields for sCI and HOM during the experiments.

| | O_3 (ppbv) | α -pinene (pptv) | SO_2 (ppbv) | Formation rate ($cm^{-3} s^{-1}$) | Growth rate ($nm h^{-1}$) | sCI yield (%) | HOM yield (%) |
|---|-----------------|----------------------------|------------------|--|--------------------------------|---------------|---------------|
| 1 | 22 | 80 | 72 | 13.26 | 1.88 | 22 | 5 |
| 2 | 24 | 80 | 72 | 9.11×10^{-2} | 1.50 | 22 | 3.5 |
| 3 | 22 | 600 | 67 | 47.98 | 7.21 | 23 | 6 |
| 4 | 22 | 170 | 68 | 3.95 | 2.32 | 24 | 5.5 |
| 5 | 22 | 530 | 17 | 18.10 | 3.77 | 32 | 6.5 |

2.2 Instruments

A Proton Transfer Reaction Mass Spectrometer (PTR-MS, Ionicon Analytik GmbH, Lindinger et al., 1998) was used to measure the concentrations of volatile organic compounds (including α -pinene). The neutral particle size distribution of 2-40 nm particles was measured with a Neutral cluster and Air Ion Spectrometer (NAIS, Mirme and Mirme, 2013). The particle size distribution of 5 to 80 nm particles was measured with a nano Scanning Mobility Particle Sizer (nanoSMPS, Wang and Flagan, 1990) and the condensation sink due to particles in the chamber was calculated from the size distribution. Sulfur dioxide concentration was measured with a high sensitivity pulse fluorescence analyzer (model 43i-TLE; Thermo Fisher Scientific Inc), and ozone with a UV photometric ozone analyzer (model 49C, Thermo Environmental Instruments).

The gas-phase sulfuric acid and HOM were detected with a nitrate ion-based Chemical Ionization Atmospheric Pressure interface Time of Flight mass spectrometer (nitrate-Cl-APi-TOF, Tofwerk AG, Thun, Switzerland and Aerodyne Research Inc., USA, Jokinen et al., 2012; Junninen et al., 2010). A soft X-ray source (Hamamatsu L9490) was deployed to ionize nitric acid to nitrate ions ($(HNO_3)_{0.2}NO_3^-$), which were used as the reagent ions for the chemical ionization. The ionization method is selectively suited for detecting strong acids such as sulfuric acid

or methane sulfonic acid (Eisele and Tanner, 1993). In the case of the oxidised organic compounds, it requires molecules to have at least two hydroperoxy (OOH) groups or some other H-bond donating groups to be ionized (Hyttinen et al., 2015). A previous study of cyclohexene ozonolysis showed that in contrast to highly oxygenated products such as $C_6H_8O_7$ and $C_6H_8O_9$ (with three carbonyl groups and two and three hydroperoxy groups, respectively) products like $C_6H_8O_5$ (three carbonyl groups and one hydroperoxy group), could not be detected (Rissanen et al., 2014). However, in previous α -pinene experiments oxidised products with a O:C ratio of as low as 0.6 have been detected (Jokinen et al., 2015; Praplan et al., 2015).

The concentration of sulfuric acid was calculated according to equation 1, where a calibration coefficient c is applied on the count rates of the bisulfate ion and its cluster with nitric acid normalized to the sum of count rates of reagent ions (Jokinen et al., 2012). To obtain the calibration coefficient c , the instrument was calibrated for sulfuric acid with a calibration setup described by Kürten et al. (2012). The calibration constant was measured to be 5×10^9 molecules/cm³. Taking sample tube losses into account a value of 1.25×10^{10} molecules/cm³ was obtained for c .

$$[H_2SO_4] = \frac{HSO_4^- + (HNO_3)HSO_4^-}{NO_3^- + (HNO_3)NO_3^- + (HNO_3)_2NO_3^-} \times c \quad (1)$$

In the experiments the concentration of sulfuric acid clusters was low since there were no stabilizing agents such as amines or ammonia added into the chamber. Thus the vast majority of the sulfuric acid concentration was in form of a monomer, not in the clusters as “hidden sulfuric acid” (Rondo et al., 2016). At most, less than 2% of the total sulfuric acid concentration was involved in the clusters while most of the time no sulfuric acid clusters were detected.

2.3 Estimation of HOM sensitivity

In this study, we counted all the α -pinene oxidation products that were detected and identified with nitrate-Cl-APi-TOF as HOM. The total concentration of HOM was calculated by summing up the high resolution fitted signals of identified highly oxygenated compounds (see the full list of peaks in the Appendix). These compounds were detected in the range of 220 – 620 Th and their O:C ratios were between 0.6 and 1.3. Most of the elemental compositions found in the experiments were the same as have been published by Ehn et al. (2012) and Jokinen et al. (2014). The sum of signals was divided by reagent ion signals and multiplied by the same calibration constant that was used for sulfuric acid (Eq. (2)).

$$[HOM] = \frac{\sum HOM \cdot NO_3^-}{NO_3^- + (HNO_3)NO_3^- + (HNO_3)_2NO_3^-} \times c \quad (2)$$

Since we did not have a direct calibration method for HOM, we considered three additional terms, which may affect the detection of molecules before the calibration constant of sulfuric acid can be used (Eq. (2), (Kürten et al., 2014)).

$$[\text{HOM}] = \frac{k_{\text{SA}}}{k_{\text{HOM}}} \times \frac{T_{\text{SA}}}{T_{\text{HOM}}} \times \frac{e_{\text{SA}}}{e_{\text{HOM}}} \times \frac{\sum \text{HOM} \cdot \text{NO}_3^-}{\text{NO}_3^- + (\text{HNO}_3)\text{NO}_3^- + (\text{HNO}_3)_2\text{NO}_3^-} \times c \quad (3)$$

The first term $k_{\text{SA}}/k_{\text{HOM}}$ corrects for the difference in reaction rate between the HOM and the reagent ions compared to sulfuric acid and the reagent ions. In the chemical ionisation method, there is an excess of nitric acid in the drift tube, where the sample flow and reagent ions meet. The nitrate dimer, $\text{HNO}_3\text{NO}_3^-$, is an extremely stable cluster,

5 which means that if there are some other clusters forming with NO_3^- in the drift tube, they need to be even more stable than the nitrate dimer. As we can detect a large total signal of HOM-nitrate clusters, we can assume that they are very stable. If we assume that all the HOM that collide with nitrate ions in the drift tube form clusters and stick together subsequently, we get the lower limit of HOM concentration from our measurements. If all collisions would not in reality produce clusters or if some fraction of the clusters would decompose in the drift tube or inside
10 the high vacuum region of the TOF, the real concentrations of HOM would be higher than assumed by this method. Ehn et al. (2014) reported calculated collision limited reaction rates of $k_{\text{HOM}} = (1.5-2.8) \times 10^{-9} \text{ cm}^3 \text{ s}^{-1}$ for HOM and $k_{\text{SA}} = (1.5-2.5) \times 10^{-9} \text{ cm}^3 \text{ s}^{-1}$ for sulfuric acid. To achieve these values Ehn et al. used the formulation of Su and Bowers (Su and Bowers, 1973) and assumed some possible structures of HOM and calculated the collision frequencies of nitrate clusters $(\text{HNO}_3)_{0-2}\text{NO}_3^-$ with selected HOM and sulfuric acid. The collision limited reaction
15 rates are so close to each other that we approximated the term $k_{\text{SA}}/k_{\text{HOM}}$ to be 1.

The second term $T_{\text{SA}}/T_{\text{HOM}}$ describes the differences in the transmission efficiency of different sized molecules or clusters through the sampling line, as increasing size of the molecule or cluster implies smaller diffusivity. A third term $e_{\text{SA}}/e_{\text{HOM}}$ takes into account the mass discrimination effects inside the mass spectrometer. The total effect of the terms $T_{\text{SA}}/T_{\text{HOM}}$ and $e_{\text{SA}}/e_{\text{HOM}}$ was determined experimentally with a high resolution differential mobility

20 analyser (HR-DMA) method (Junninen et al., 2010). By this method, trioctylmethylammonium bis(trifluoromethylsulfonyl)imide particles were produced with an electrospray, and size ranges were selected with a high-resolution Vienna type Differential Mobility Analyzer (UDMA, Steiner et al., 2010) and the selected size range was guided to the APi-TOF. To calculate the transmission the signal in the mass spectrometer was divided by the signal in the electrometer. The transmission in the mass range between 90 and 600 Th varied so that the
25 largest difference compared to the transmission of sulfuric acid was 1.4-fold at 320 Th (7.3×10^{-4}). Since HOM could be measured over a wide mass range, the transmission varied between individual HOM molecules ($(6.4-10.4) \times 10^{-4}$). The averaged difference of the transmissions was around 30% so that the transmission of HOM signals was higher than the sulfuric acid signals, and this was taken into account in the concentration calculations by correcting the values according the transmission curve.

30 We estimated a systematic uncertainty of +50%/-33% for the sulfuric acid concentration (Kirkby et al., 2016). The estimation is based on the uncertainty of the sulfuric acid calibration and a comparison with the sulfuric acid concentration measured by an other CIMS instrument (independently measured sulfuric acid concentration at CLOUD experiments (Kürten et al., 2011)). For the HOM concentration the uncertainty is larger due to lack of a direct calibration method. We estimated an uncertainty of +80%/-45% for HOM concentrations taking the sulfuric
35 acid calibration, charging efficiency, mass dependent transmission efficiency calibration and sampling line losses into consideration (Jokinen et al., 2015; Kirkby et al., 2016). The uncertainty for HOM yield arises from the

uncertainties of α -pinene concentration, O₃ concentration, HOM wall loss rate and rate constants. This results in a mean estimated uncertainty in HOM yield of +100%/-60%.

2.3 The simulations of sCI and HOM concentrations

The temporal behaviour of the reaction products from monoterpene ozonolysis in the CLOUD chamber was simulated with a 0-dimensional kinetic model. The production of stabilized Criegee Intermediates was calculated from the measured α -pinene and ozone concentrations using a reaction rate coefficient of $8.05 \times 10^{-17} \text{ cm}^3 \text{s}^{-1}$ (Atkinson et al., 2006, updated data sheet can be found: http://iupac.pole-ether.fr/htdocs/datasheets/pdf/Ox_VOC8_O3_apinene.pdf). Since the temperature of CLOUD experiments was lower than in previous experiments done in TROPOS-LFT and JPAC (Table 1) we used a lower reaction rate coefficient ($8.66 \times 10^{-17} \text{ cm}^3 \text{s}^{-1}$

¹⁰ in JPAC experiments and $1.1 \times 10^{-16} \text{ cm}^3 \text{s}^{-1}$ in TROPOS-LFT experiments). The reaction of sCI with SO₂ is in competition with the reaction of sCI with water vapour, thus, three loss paths were taken into account for the sCI: (1) its reaction with sulfur dioxide (k_{sCI+SO_2}), (2) the thermal decomposition of sCI (k_{dec}) and (3) its reaction with water vapour (k_{sCI+H_2O}). The latter two reactions are included in the loss term k_{loss} (Eq. (4)) **and under these**

conditions thermal decomposition dominates the loss mechanism of sCI and the reaction with water vapour is less

¹⁵ **important.** The condensation sink, wall loss and dilution are negligible compared to the loss term k_{loss} . The reaction rate of sCI and water vapour has been found to strongly depend on the structure of the Criegee Intermediate (Berndt et al., 2014c; Huang et al., 2015) and for the monoterpene-derived sCIs, the relative rate coefficients k_{loss} / k_{sCI+SO_2} was found to be nearly independent of the relative humidity (Sipilä et al., 2014). The kinetic study of Huang et al. (2015) suggested that sCIs with more complicated substitution groups (such as α -pinene derived sCIs) react with ²⁰ water slowly but react with SO₂ quickly, thus, supporting the reaction parameters achieved by Sipilä et al. (2014). Other possible loss paths of sCI are considered to be negligible. The studies of Vereecken et al. (2012 and 2014) show that a high substitution of CI and/or the reaction partner results in strong steric hindrance between the substituents, which effectively inhibits reactions between them. Thus, reactions between monoterpene-derived sCI and SVOCs are not favorable.

$$25 \quad k_{loss} = k_{dec.} + (k_{(sCI+H_2O)} \times [H_2O]) \quad (4)$$

$$\frac{d[sCI]}{dt} = (Y_{sCI} \times k_{O_3+\alpha-pinene} \times [O_3] \times [\alpha-pinene]) - (k_{loss} \times [sCI]) - (k_{sCI+SO_2} \times [SO_2] \times [sCI]) \quad (5)$$

The concentration of sCI was calculated according to equation 5, in which the values of the reaction rate coefficient (k_{sCI+SO_2}), the sCI yield term (Y_{sCI}) and the loss term (k_{loss}) were taken from the TROPOS-LFT measurements (Sipilä et al., 2014). Those measurements were conducted at 50% relative humidity (RH) and the derived sCI yield from the reaction between α -pinene and ozone was determined to be 0.15 ± 0.07 and the ratio between the loss term and k_{sCI+SO_2} was $(2.0 \pm 0.4) \times 10^{12} \text{ molecules cm}^{-3}$. Sipilä et al. (2014) also found that in the case of α -pinene and limonene the ratio k_{loss} / k_{sCI+SO_2} was nearly independent of the relative humidity. Therefore we neglected the difference in RH of the experiments shown here (38%) compared to the experiments at TROPOS-LFT (50%). The temperature was 278 K in the CLOUD experiments whereas it was 293 K in the previous experiments. The influence of temperature on the H₂SO₄ formation from the gas-phase reaction of monoterpene-derived sCIs has not yet been investigated. It is very likely that k_{dec} is higher at higher temperatures, which would cause

underestimation of the sulfuric acid concentration in the CLOUD simulations, where we are using the loss term derived from experiments performed at higher temperature. In the sCI yield experiments performed with acetone oxide the temperature influence on the ratio of k_{dec} / k_{sCI+SO_2} was 2-fold when the temperature was increased by 10 K (Berndt et al., 2014c).

5 The lower and upper limits of the sCI concentration were modelled with the upper and lower values for the yield and loss term so that the lower limit was calculated with a yield of 8% and a k_{loss} / k_{sCI+SO_2} ratio of 2.4×10^{12} molecules cm^{-3} while the upper limit was calculated with a yield of 22% and a k_{loss} / k_{sCI+SO_2} ratio of 1.6×10^{12} molecules cm^{-3} (Eq. (5)). For the calculations we needed to separate the terms k_{loss} and k_{sCI+SO_2} from each other. As long as the ratio between the terms stays the same, the chosen values do not make a difference for the sulfuric acid concentration.

10 The concentration of sulfuric acid in the CLOUD chamber was modelled according to equation 6.

$$\frac{d[\text{H}_2\text{SO}_4]}{dt} = k_{sCI+SO_2} \times [\text{sCI}] \times [\text{SO}_2] - (\text{CS} + k_{wall_loss} + k_{dil}) \times [\text{H}_2\text{SO}_4] \quad (6)$$

15 As an OH scavenger was used in the experiments, the only formation pathway for sulfuric acid was assumed to be the reaction between the sCI and SO_2 . The production of sulfuric acid was calculated with the modelled sCI concentration, measured sulfur dioxide concentration and the reaction coefficient k_{sCI+SO_2} (Sipilä et al., 2014). The sulfur dioxide and ozone concentrations were kept constant during the experiments. Three loss processes were taken into account for sulfuric acid: the condensation sink (CS), the wall loss (k_{wall_loss}) and the dilution (k_{dil}). The lifetime of sulfuric acid with respect to wall loss in the CLOUD chamber has been measured to be around 550s (Almeida et al., 2013; Duplissy et al., 2016; Rondo et al., 2014). The dilution rate due to injection of makeup gases 20 into the chamber was $0.1 \times 10^{-3} \text{ s}^{-1}$.

25 The production rate of HOM in the CLOUD chamber (Eq. (7)) was calculated from the measured α -pinene and ozone concentrations, a reaction rate coefficient of $8.05 \times 10^{-17} \text{ cm}^3 \text{ s}^{-1}$ (Atkinson et al., 2006) and an experimentally derived yield term (Ehn et al., 2014; Jokinen et al., 2015). The yield of HOM from the reaction between α -pinene and ozone was reported in recent studies. Ehn et al. (2014) obtained a yield of $7 \pm 3.5\%$ in their experiments in the Jülich Plant Atmosphere Chamber while Jokinen et al. (2015) calculated a yield of 3.4% with an estimated uncertainty of $-1.7 \pm 3.4\%$ from the experiments done in the TROPOS-LFT and Kirkby et al. (2016) reported a yield of 2.9% for the CLOUD experiments with and without ions. The same loss paths were taken into account in the modelled HOM concentration as for the sulfuric acid concentration. The lifetime of HOM was measured to be around 900s which is longer than the lifetime of sulfuric acid (Kirkby et al., 2016). All the values used in the 30 modelling of sulfuric acid and HOM concentrations are shown in Table 3.

$$\frac{d[\text{HOM}]}{dt} = Y_{HOM} \times k_{O_3+\alpha-\text{pinene}} \times [\text{O}_3] \times [\alpha-\text{pinene}] - (\text{CS} + k_{wall_loss} + k_{dil}) \times [\text{HOM}] \quad (7)$$

Table 3: Reaction rates, loss terms and yields used in simulations.

| | |
|--|---|
| k_{loss} / k_{sCI+SO_2} | $(1.6-2.4) \times 10^{12}$ molecules cm^{-3} |
| $k_{O_3+\alpha\text{-pinene}}$ | 8.05×10^{-17} $\text{cm}^3 \text{s}^{-1}$ |
| Y_{sCI} (Sipilä et al. 2014) | 0.08-0.22 |
| $k_{wall\ loss\ (SA)}$ | 1.8×10^{-3} $\text{cm}^3 \text{s}^{-1}$ |
| $k_{wall\ loss\ (HOM)}$ | 1.1×10^{-3} $\text{cm}^3 \text{s}^{-1}$ |
| k_{dil} | 0.1×10^{-3} $\text{cm}^3 \text{s}^{-1}$ |
| Y_{HOM} (Ehn et al. 2014) | 0.035-0.105 |
| Y_{HOM} (Jokinen et al. 2014) | 0.017-0.068 |

3 Results

3.1 Reaction products from α -pinene ozonolysis

5 During the ozonolysis experiments of α -pinene, a simultaneous increase of the concentrations of sulfuric acid and HOM were observed. Several highly oxidised α -pinene oxidation products were observed between 220 and 620 Th (Fig. 1). All the HOM were detected as clusters with a nitrate ion (NO_3^-). As a result of the high cleanliness of the CLOUD chamber, the mass spectra consist mainly of the oxidation products and concentrations of contaminants were low. The most abundant HOM monomers, containing a C_{10} carbon skeleton, had an O:C ratio

10 between 0.7 and 1.1 whereas the most abundant HOM dimers, containing a C_{20} carbon skeleton, had an O:C ratio around 0.6-0.8. The highest concentrations were observed from compounds identified as $\text{C}_{10}\text{H}_{14}\text{O}_7$, $\text{C}_{10}\text{H}_{15}\text{O}_8$, $\text{C}_{10}\text{H}_{14}\text{O}_9$, $\text{C}_{10}\text{H}_{15}\text{O}_{10}$, $\text{C}_{10}\text{H}_{16}\text{O}_{10}$ and $\text{C}_{19}\text{H}_{28}\text{O}_{11}$, which represent the majority of the total concentration of HOM. These compounds have also been found to be abundant in the boreal forest, when analysing the naturally charged ions (Ehn et al., 2012). The rest of the compounds taken into account in the concentration calculation are listed in

15 Table A1.




Figure 1: HOM mass spectrum during an α -pinene ozonolysis experiment (C10 compounds in the upper panel and C20 compounds in the lower panel). The measured mass spectrum is depicted in black and the compounds identified as HOM are depicted in red. The elemental composition of the compounds with the highest concentrations are shown in the figure and the six most abundant compounds are labelled in bold face.

During most of the experiments, clear particle formation and growth was observed shortly after the α -pinene injection was started. In Figure 2 the particle size distribution and precursor vapour concentrations during an example ozonolysis experiment are shown. In this experiment the α -pinene injection started at noon and the sulfuric acid and HOM concentrations started to increase immediately. The particle growth above 3 nm can be seen approximately 45 minutes after the injection. While the concentration of α -pinene continued to increase, the sulfuric acid and HOM concentrations reach their steady-state concentrations after one to two hours. The sulfuric acid reaches its steady-state concentration slightly before the HOM concentration reaches its maximum value which is expected due to its faster wall loss rate.




Figure 2: Example of the size distribution (2-40 nm, measured by NAIS) of neutral particles (a) and the concentrations of sulfuric acid, HOM and α -pinene (b) during an ozonolysis experiment in the CLOUD chamber.

5

The method presented in Sec. 2.3 was used to describe the temporal behaviour of the total HOM concentration.

Since the total HOM is a sum of several molecules that are formed by the same autoxidation mechanism but

possibly via various different intermediate steps, the time evolution of individual HOM molecules can differ from one to another. The time evolution of sulfuric acid and the most abundant HOM was studied in detail using mass

10 spectra integrated over 30s. The time evolution of the experiment with 600 pptv of α -pinene, 22 ppbv of O_3 and 67 ppbv of SO_2 is shown in Fig. 3. In our studies sulfuric acid concentration started to increase first followed by the concentration of RO_2 ($C_{10}H_{15}O_8$ and $C_{10}H_{15}O_{10}$). The formation of closed shell monomers ($C_{10}H_{14}O_7$ and

15 $C_{10}H_{14}O_9$) started a few minutes after the RO_2 . The most oxidised closed shell monomer of the selected HOM ($C_{10}H_{16}O_{10}$) and the dimer ($C_{19}H_{28}O_{11}$) took more than ten minutes to start increasing. The time evolution of the

compounds might give us information about the formation of the molecules. The rapid formation of the radicals and $C_{10}H_{15}O_8$ and $C_{10}H_{15}O_{10}$ implies that they are formed via autoxidation in which the peroxy radical undergoes

oxidation by adding oxygen molecules stepwise. The selected dimer ($C_{19}H_{28}O_{11}$) formation starts clearly later, which supports the hypothesis that it forms from reaction of two RO_2 (Ehn et al., 2014; Jokinen et al., 2014). The

carbon number 19 can be explained by loss of CO from RO_2 (Jangiella et al., 2000; Rissanen et al., 2014), followed

20 by reaction with a 10-carbon RO_2 . The interesting feature in this data is that $C_{10}H_{16}O_{10}$ appears significantly later than most monomers, at the same time that the first dimer appears in the spectrum. This might indicate that this

more highly oxidised product is also formed via bi-molecular reaction of two RO_2 radicals. The time evolution was similar in all the experiments. In the experiments with low α -pinene (80 pptv), the concentrations of $C_{19}H_{28}O_{11}$ and $C_{10}H_{16}O_{10}$ were very low. It is also a possibility that the formation of dimers (and other compounds that appear

25 later in the measurements) starts already earlier but the concentrations are just below the detection limit.

Understanding the exact formation mechanisms of individual HOM compounds requires additional experiments and will be a topic of further studies.




Figure 3: The time evolution of bisulfate ion (green dotted line), total HOM (blue line) and the HOM signals

5 **of the highest concentrations with 30s time resolution in experiment with 600 ppt of α -pinene and 67 ppb of SO₂. RO₂ signals are shown with dashed lines. All HOM are detected as clusters with nitrate-ion. The black line shows the start of the experiment (i.e. α -pinene injection).**

In our experiments, we used H₂ to scavenge the OH. In the reaction of OH and H₂, water and H is produced and 10 subsequently H can react with O₂ to form HO₂ (Eq. (8) and (9)). HO₂ can then react with RO₂ ending its autoxidation process (Ehn et al., 2014). This means that in the presence of HO₂ the HOM concentration can be lower because the organic compounds that react with HO₂ are not oxidised further into highly oxidised products. In these experiments, we did not have an instrument capable of measuring less oxidised products from α -pinene ozonolysis. The relevance of these experiments to the atmosphere depends on the relative and absolute levels of 15 all species participating in the autoxidation process, including RO₂, HO₂, and NO in both the experiment and the atmosphere. Jokinen et al. (2015) also used H₂ to scavenge OH while Ehn et al. (2014) used CO, both of which produce also H and then HO₂ (Eq. (10)). Thus, these experiments and the yield terms determined from them are equally affected by HO₂. However, Jokinen et al. (2015) did also experiments with propane as OH scavenger,


which does not produce HO_2 , and found similar yields as with H_2 . This implies that HO_2 , produced by the scavenger reactions, does not significantly affect HOM formation.



3.2 The formation of sulfuric acid

The sulfuric acid formation in the CLOUD chamber was simulated as described in section 2.3. The measured steady-state concentrations varied between 4×10^6 and 2×10^7 molecules cm^{-3} (Fig. 4). Sulfuric acid concentrations were the highest in the experiments where also the α -pinene mixing ratio was the highest, around 600 pptv. The

10 steady-state concentration was reached in two hours in the experiment at high SO_2 concentrations (~70 ppbv) whereas in the experiment with the same amount of α -pinene but significantly lower sulfur dioxide concentration (17 ppbv), the steady-state was reached an hour later. In the other three experiments the α -pinene mixing ratio was clearly lower (80 pptv and 170 pptv) and the increase of sulfuric acid concentration took more time and continued throughout the whole experiment.



15

Figure 4: Measured (black dots) and modelled (green shade shows the concentration with uncertainty) sulfuric acid concentrations formed from the oxidation of SO_2 by sCI in the CLOUD chamber. The simulation with fitted yield term to the measured concentration is shown as a blue line and the α -pinene concentration is a grey line. Under conditions of atmospheric concentrations of OH (1×10^6 molecules cm^{-3})

the sulfuric acid concentrations would be significantly higher, around 6.3×10^8 molecules/cm³ in the experiment with 67 ppb of SO₂ and 1.6×10^8 molecules/cm³ in the experiment with 17 ppb of SO₂.

In the simulations, the minimum and maximum concentrations were calculated from the upper and lower limits of given k_{loss} / k_{sCI+SO_2} and sCI yield term in Sipilä et al. (2014). In the CLOUD experiments, the measured sulfuric

5 acid concentrations were at the upper range of the simulated concentrations in all the cases (Fig. 4). In the experiment where we had the least sulfur dioxide in the chamber, the measured concentration was slightly higher than the simulated one (measured 1.6×10^7 molecules cm⁻³, upper range simulated concentration 1.2×10^7 molecules cm⁻³). The sulfuric acid was formed in the fast reaction between SO₂ and sCI, thus, the formation of sulfuric acid was strongly dependent on the formation of sCI. We calculated yield terms for the sCI in CLOUD experiments by

10 fitting the model to the measured concentrations. When using a value of 1.6×10^{12} for the term k_{loss} / k_{sCI+SO_2} (lower end of the range given in Sipilä et al. 2014) the calculated yields of sCI were 22-24% for the experiments with higher concentration of SO₂ and 32% for the experiment with low SO₂. (Table 2). **The higher yield in the**

experiment with low SO₂ concentration is still within the experimental uncertainties.

In the study done in the TROPOS-LFT the sCI yield from α -pinene oxidation was determined to be 8-22% (Sipilä et al., 2014). The

15 simulated sulfuric acid concentration represent the measured concentrations well. When the α -pinene concentration was low, the measured α -pinene concentration was fluctuating which led to fluctuation in the simulated sulfuric acid concentration. In the experiment with a middle-range α -pinene concentration (170 pptv), the upper bound of the simulated time evolution matched perfectly with the measured concentrations, so that the trend in measured and simulated concentration was identical and the difference of simulated concentration from

20 measured concentration did not exceed 30%. In the other experiments the measured sulfuric acid concentration increased faster than the simulated concentration in the beginning but then stabilized at the upper level of the simulated concentrations. The difference is still small and mostly within the measurement uncertainty. In all the experiments, the simulated sulfuric acid concentration followed the measured concentration very well after 10000 s (166 min) and thus the discrepancy cannot be explained by only one term. The simulation can be modified to

25 match the measurements better if, for example, both the sCI yield term and condensation sink values are increased significantly (two-fold increase in both condensation sink and sCI yield). However, it seems unlikely that **the** condensation sink for sulfuric acid would have such a large error. As mentioned earlier, the influence of temperature on the H₂SO₄ formation from the gas-phase reaction of monoterpane-derived sCIs has not been investigated. It is likely that we underestimate the sulfuric acid concentration in the CLOUD experiments as we

30 are using the loss term derived from experiments performed at higher temperature.

To compare the sCI oxidation with ambient sulfuric acid formation, we calculated the sulfuric acid produced at typical ambient OH concentration for otherwise similar conditions as in these experiments. The sulfur dioxide concentration was high in most of the experiments and with atmospherically relevant concentration of OH (1×10^6 molecules cm⁻³) the sulfuric acid concentration would be around 6.3×10^8 molecules cm⁻³ (SO₂ 67 ppbv, reaction

35 rate constant 8.5×10^{-13} cm³s⁻¹ (Weber et al., 1996)). With lower SO₂ concentration (17 ppbv) the OH-produced sulfuric acid would be around 1.6×10^8 molecules cm⁻³. Thus, the sulfuric acid concentrations that resulted from sCI oxidation in these experiments were around 3% of what would be formed from OH oxidation in high SO₂ conditions and 10% in low SO₂ conditions at typical OH concentrations. It should be noted that in the atmosphere the mixture of gases is much more complex. In ambient conditions the α -pinene concentration is often less than

the concentration used in this calculation (600 pptv) but on the other hand in the atmosphere there are also other alkenes than α -pinene that can be oxidised to form sCI. Also the sulfur dioxide concentrations used in the experiments are relatively high for most of the atmosphere, such as rural areas (Mikkonen et al., 2011).

To get more insight in the sulfuric acid production with ambient concentrations we calculated 24 hour production of sulfuric acid from OH and sCI oxidation pathways. We used typical spring – summer time concentrations of precursors in boreal forest: measured OH concentrations (medians of event day concentrations from late March to early June, Petäjä et al., 2008), measured O_3 , SO_2 , (medians of concentrations from April to June in 2013, Smart-SMEAR: <https://avaa.tdata.fi/web/smart/smear>, Junninen et al., 2009), measured monoterpene concentrations (concentrations measured in July 2004, Rinne et al., 2005) and calculated condensation sink values (median of data from April to June 2013 Junninen et al., 2009; Kulmala et al., 2001). The 24 hour sulfuric acid productions were calculated with sCI yield of 22% and the results are shown in Fig. 5a. During the daytime the sulfuric acid produced by OH dominates but during night time both of the production pathways are important. In this example with conditions of boreal forest the SO_2 concentration is significantly lower than in our experiments (Fig 5d, around 0.1 ppb). Ozone concentrations are lowest during early morning being around 35 ppbv while the concentrations reach 43 ppbv in the evening. The importance of sCI in the sulfuric acid production strongly depends on the monoterpene concentrations. In this example the monoterpene concentration is highest during early hours and at that time the sulfuric acid concentration reaches 4×10^4 molecules/cm³. We calculated a sCI yield of 32% in our experiment with low SO_2 and if we use that yield term in the calculation the highest sulfuric acid concentration becomes 6×10^4 molecules/cm³.





Figure 5. Example of sulfuric acid concentration produced by OH and sCI in ambient boreal forest conditions (a). The precursor gas concentrations and condensation sink used are shown in plots b-e.

20

3.3 The formation of HOM

The HOM formation in the CLOUD chamber was simulated as described in section 2.3. The measured steady-state concentrations of HOM varied between 2×10^6 and 2×10^7 molecules cm^{-3} (Fig. 6). The respective α -pinene and ozone concentrations were atmospherically relevant, so these HOM concentrations are similar to those found

5 in ambient air (Sarnela et al., 2015; Yan et al., 2016). We calculated the HOM yield by fitting the model to the measured concentrations and obtained yields of 3.5-6.5% for the experiments in the CLOUD chamber (Table 2). In previous studies of α -pinene ozonolysis, yield terms for HOM formation have been experimentally determined. Ehn et al. (2014) measured a yield of $7 \pm 3.5\%$ and Jokinen et al. (2015) a yield of 3.4% with an estimated uncertainty of $-50\% / +100\%$. Kirkby et al. (2016) made a fit to both neutral and charged experiments and obtained 10 a yield of 2.9% for the CLOUD experiments. The yields calculated in this study are in good agreement with all previous studies and the simulated time evolution reproduce the measured concentrations very well. The highest difference between simulated and measured concentration was 40% but in most experiments the simulated and measured concentrations matched within 20% difference.



15 **Figure 6: Measured (grey dots) and simulated (shaded area show the concentration with uncertainty) HOM concentrations formed from ozonolysis of α -pinene in CLOUD-chamber. The simulation with fitted yield term to the measured concentration is shown as a black line. The modelled concentration with a yield term**

from Ehn et al. (2014) is shown in red shade while the concentration with the yield term from Jokinen et al. (2015) is shown in yellow shade. The overlapping area within the error estimates of these studies is coloured in orange shade.

5 4 Conclusions

In this study we conducted several α -pinene ozonolysis experiments in an ultraclean environment, the CLOUD-chamber. These experiments were designed to be OH-radical free, thus allowing to study the formation of Highly Oxygenated Molecules (HOM) from the ozonolysis of α -pinene. The other objective of this study was to observe the formation of sulfuric acid from the oxidation of SO_2 by stabilized Criegee Intermediates. Both HOM and sulfuric acid concentrations were experimentally measured with a high resolution time-of-flight mass spectrometer by utilizing a highly selective chemical ionization method. To estimate the molar yield of the HOM and the sCI-yield in our experiments, we used a 0-dimensional model with reaction parameters some of which were obtained from other recent publications on sCI and HOM formation (Ehn et al., 2014; Jokinen et al., 2015; Sipilä et al., 2014).

15 The formation of HOM was initiated immediately after an α -pinene injection into the chamber with a stable ozone concentration. We observed a consecutive formation of peroxy radicals, HOM monomer and dimer species, which is in agreement with previous studies conducted in both, a laminar flow tube and in a continuously stirred flow reactor (Ehn et al., 2014; Jokinen et al., 2014). The simulated time evolution of the HOM followed the measured concentrations very precisely and the calculated yields from several experiments were in the range of 3.5-6.5%.

20 The yields observed in the CLOUD chamber were within the range of previously published HOM yields for α -pinene ozonolysis (3.5-10.5% by Ehn et al., 2014, 1.7-6.8% by Jokinen et al., 2015, 1.2-5.8% Kirkby et al., 2016).

Sulfuric acid in the chamber was assumed to be solely produced via stabilized Criegee Intermediates reacting with the added SO_2 . The formation of sulfuric acid started promptly after the α -pinene injection and the associated sCI formation. The measured concentration increased quickly, in some experiments even faster than was expected from the simulations. With a high SO_2 concentration (70 ppbv), the sCI yields were measured to reach 22-24%, which are on the upper edge of the values found by Sipilä et al. (2014), i.e. $15\pm7\%$. When the SO_2 concentration was considerably lower (17 ppbv) the sCI yield was higher (32%). These results are not denying that OH is the main daytime oxidiser of sulfur dioxide. In the presence of OH the role of sCI in the formation of sulfuric acid is relatively small but in dark conditions there can be considerable sulfuric acid formation due to sCI. The results of this study emphasize the potential importance of stabilized Criegee Intermediates in sulfuric acid formation, also in the presence of water vapour. In this paper we introduce a way to simulate the ozonolysis products of α -pinene in a simple manner. The results indicate that the CLOUD experiments on α -pinene ozonolysis support the recently published chemistry of HOM and sCI formation, thus making the experimentally determined yield and loss terms more reliable for following modelling and theoretical use.

35 Acknowledgements

We would like to thank CERN for supporting CLOUD with technical and financial resources. We also thank Dr. Theo Kurtén for valuable input on ion-molecule collision frequencies. This research has received funding from

the EC Seventh Framework Programme (Marie Curie Initial Training Network ‘CLOUD-ITN’ no. 215072, MC-ITN ‘CLOUD-TRAIN’, no. 316662, and ERC-StG-ATMOGAIN (278277) and ERC-Advanced ‘ATMNUCLE’ grant no. 227463 and ERC-StG-GASPARCON (714621) **and ERC-StG-COALA (638703)**), the German Federal Ministry of Education and Research (project nos 01LK0902A and 01LK1222A), the Swiss National Science Foundation (project nos 200020_135307, 200020_152907, 20FI20_149002 and 200021_140663), the Academy of Finland Center of Excellence programme (grant no. 307331), the Academy of Finland (CoE project no. 1118615, LASTU project no. 135054), the Academy of Finland (296628 **and 299574**), the Nessling Foundation, the Austrian Science Fund (FWF; project no. J3198-N21), the EU’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie (no. 656994), the Swedish Research Council, Vetenskapsrådet (grant no. 2011-5120), the Portuguese Foundation for Science and Technology (project no. CERN/FP/116387/2010), the Presidium of the Russian Academy of Sciences and Russian Foundation for Basic Research (grants 08-02-91006-CERN and 12-02-91522-CERN), Dreyfus Award EP-11-117, the Davidow Foundation, the US National Science Foundation (grants AGS1136479, AGS1447056, AGS1439551 and CHE1012293), US Department of Energy (grant DE-SC00014469) and the FP7 project BACCHUS (grant agreement 603445).

References

Almeida, J., Schobesberger, S., Kürten, A., Ortega, I. K., Kupiainen-Määttä, O., Praplan, A. P., Adamov, A., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Donahue, N. M., Downard, A., Dunne, E., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Henschel, H., Jokinen, T., Junninen, H., Kajos, M., Kangasluoma, J., Keskinen, H., Kupc, A., Kurtén, T., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Leiminger, M., Leppä, J., Loukonen, V., Makhmutov, V., Mathot, S., McGrath, M. J., Nieminen, T., Olenius, T., Onnela, A., Petäjä, T., Riccobono, F., Riipinen, I., Rissanen, M., Rondo, L., Ruuskanen, T., Santos, F. D., Sarnela, N., Schallhart, S., Schnitzhofer, R., Seinfeld, J. H., Simon, M., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Tröstl, J., Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Virtanen, A., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Williamson, C., Wimmer, D., Ye, P., Yli-Juuti, T., Carslaw, K. S., Kulmala, M., Curtius, J., Baltensperger, U., Worsnop, D. R., Vehkämäki, H. and Kirkby, J.: Molecular understanding of sulphuric acid–amine particle nucleation in the atmosphere, *Nature*, 502, 359–363, doi:10.1038/nature12663, 2013.

Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J. and Troe, J.: IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, *Atmos. Chem. Phys.*, 6, 3625, 2006.

Ball, S. M., Hanson, D. R., Eisele, F. L. and McMurry, P. H.: Laboratory studies of particle nucleation: Initial results for H₂SO₄, H₂O, and NH₃ vapors, *J. Geophys. Res. Atmos.*, 104, 23709–23718, doi:10.1029/1999JD900411, 1999.

Berndt, T., Böge, O., Stratmann, F., Heintzenberg, J. and Kulmala, M.: Rapid formation of sulfuric acid particles at near-atmospheric conditions., *Science*, 307, 698–700, doi:10.1126/science.1104054, 2005.

Berndt, T., Stratmann, F., Sipilä, M., Vanhanen, J., Petäjä, T., Mikkilä, J., Grüner, A., Spindler, G., Lee Mauldin III, R., Curtius, J., Kulmala, M. and Heintzenberg, J.: Laboratory study on new particle formation from the reaction OH + SO₂: influence of experimental conditions, H₂O vapour, NH₃ and the amine tert-butylamine on the overall process, *Atmos. Chem. Phys.*, 10, 7101–7116, doi:10.5194/acp-10-7101-2010, 2010.

5 Berndt, T., Jokinen, T., Mauldin, R. L., Petäjä, T., Herrmann, H., Junninen, H., Paasonen, P., Worsnop, D. R. and Sipilä, M.: Gas-phase ozonolysis of selected olefins: The yield of stabilized criegee intermediate and the reactivity toward SO₂, *J. Phys. Chem. Lett.*, 3, 2892–2896, doi:10.1021/jz301158u, 2012.

Berndt, T., Voigtländer, J., Stratmann, F., Junninen, H., Mauldin, R. L., Sipilä, M., Kulmala, M. and Herrmann, H.: Competing atmospheric reactions of CH₂OO with SO₂ and water vapour, *Phys. Chem. Chem. Phys.*, 16, 10 19130–19136, doi:10.1039/C4CP02345E, 2014a.

Berndt, T., Sipilä, M., Stratmann, F., Petäjä, T., Vanhanen, J., Mikkilä, J., Patokoski, J., Taipale, R., Mauldin III, R. L. and Kulmala, M.: Enhancement of atmospheric H₂SO₄/H₂O nucleation: organic oxidation products versus amines, *Atmos. Chem. Phys.*, 14, 751–764, doi:10.5194/acp-14-751-2014, 2014b.

Berndt, T., Jokinen, T., Sipilä, M., Mauldin, R. L., Herrmann, H., Stratmann, F., Junninen, H. and Kulmala, M.: 15 H₂SO₄ formation from the gas-phase reaction of stabilized Criegee Intermediates with SO₂: Influence of water vapour content and temperature, *Atmos. Environ.*, 89, 603–612, doi:10.1016/j.atmosenv.2014.02.062, 2014c.

Bianchi, F., Tröstl, J., Junninen, H., Frege, C., Henne, S., Hoyle, C. R., Molteni, U., Herrmann, E., Adamov, A., 20 Bukowiecki, N., Chen, X., Duplissy, J., Gysel, M., Hutterli, M., Kangasluoma, J., Kontkanen, J., Kürten, A., Manninen, H. E., Münch, S., Peräkylä, O., Petäjä, T., Rondo, L., Williamson, C., Weingartner, E., Curtius, J., Worsnop, D. R., Kulmala, M., Dommen, J. and Baltensperger, U.: New particle formation in the free troposphere: A question of chemistry and timing, *Science*, 352, 1109–1112, doi:10.1126/science.aad5456, 2016.

Boy, M., Mogensen, D., Smolander, S., Zhou, L., Nieminen, T., Paasonen, P., Plass-Dülmer, C., Sipilä, M., Petäjä, T., Mauldin, L., Berresheim, H. and Kulmala, M.: Oxidation of SO₂ by stabilized Criegee intermediate (sCI) radicals as a crucial source for atmospheric sulfuric acid concentrations, *Atmos. Chem. Phys.*, 13, 3865–3879, 25 doi:10.5194/acp-13-3865-2013, 2013.

Chuong, B., Zhang, J. and Donahue, N. M.: Cycloalkene Ozonolysis: Collisionally Mediated Mechanistic Branching, *J. Am. Chem. Soc.*, 126, 12363–12373, doi:10.1021/ja0485412, 2004.

Cox, R. A. and Penkett, S. A.: Oxidation of Atmospheric SO₂ by Products of the Ozone–Olefin Reaction, *Nature*, 230, 321–322, doi:10.1038/230321a0, 1971.

30 Criegee, R.: Mechanism of Ozonolysis, *Angew. Chemie Int. Ed. English*, 14, 745–752, doi:10.1002/anie.197507451, 1975.

Donahue, N. M., Drozd, G. T., Epstein, S. A., Presto, A. A. and Kroll, J. H.: Adventures in ozoneland: down the rabbit-hole, *Phys. Chem. Chem. Phys.*, 13, 10848–10857, doi:10.1039/C0CP02564J, 2011.

Donahue, N. M., Kroll, J. H., Pandis, S. N. and Robinson, A. L.: A two-dimensional volatility basis set – Part 2: 35 Diagnostics of organic-aerosol evolution, *Atmos. Chem. Phys.*, 12, 615–634, doi:10.5194/acp-12-615-2012, 2012.

Drozd, G. T. and Donahue, N. M.: Pressure dependence of stabilized Criegee intermediate formation from a sequence of alkenes., *J. Phys. Chem. A*, 115, 4381–4387, doi:10.1021/jp2001089, 2011.

Duplissy, J., Merikanto, J., Franchin, A., Tsagkogeorgas, G., Kangasluoma, J., Wimmer, D., Vuollekoski, H., Schobesberger, S., Lehtipalo, K., Flagan, R. C., Brus, D., Donahue, N. M., Vehkamäki, H., Almeida, J., Amorim, 5 A., Barmet, P., Bianchi, F., Breitenlechner, M., Dunne, E. M., Guida, R., Henschel, H., Junninen, H., Kirkby, J., Kürten, A., Kupc, A., Määttänen, A., Makhmutov, V., Mathot, S., Nieminen, T., Onnela, A., Praplan, A. P., Riccobono, F., Rondo, L., Steiner, G., Tome, A., Walther, H., Baltensperger, U., Carslaw, K. S., Dommen, J., Hansel, A., Petäjä, T., Sipilä, M., Stratmann, F., Vrtala, A., Wagner, P. E., Worsnop, D. R., Curtius, J. and Kulmala, 10 M.: Effect of ions on sulfuric acid-water binary particle formation: 2. Experimental data and comparison with QC-normalized classical nucleation theory, *J. Geophys. Res. Atmos.*, 121, 1752–1775, doi:10.1002/2015JD023539, 2016.

Ehn, M., Kleist, E., Junninen, H., Petäjä, T., Lönn, G., Schobesberger, S., Dal Maso, M., Trimborn, A., Kulmala, 15 M., Worsnop, D. R., Wahner, A., Wildt, J. and Mentel, T. F.: Gas phase formation of extremely oxidized pinene reaction products in chamber and ambient air, *Atmos. Chem. Phys.*, 12, 5113–5127, doi:10.5194/acp-12-5113-2012, 2012.

Ehn, M., Thornton, J. A., Kleist, E., Sipilä, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F., Tillmann, R., Lee, B., Lopez-Hilfiker, F., Andres, S., Acir, I.-H., Rissanen, M., Jokinen, T., Schobesberger, S., Kangasluoma, J., Kontkanen, J., Nieminen, T., Kurtén, T., Nielsen, L. B., Jørgensen, S., Kjaergaard, H. G., Canagaratna, M., 20 Maso, M. D., Berndt, T., Petäjä, T., Wahner, A., Kerminen, V.-M., Kulmala, M., Worsnop, D. R., Wildt, J. and Mentel, T. F.: A large source of low-volatility secondary organic aerosol, *Nature*, 506, 476–479, doi:10.1038/nature13032, 2014.

Eisele, F. L. and Tanner, D. J.: Measurement of the gas phase concentration of H₂SO₄ and methane sulfonic acid and estimates of H₂SO₄ production and loss in the atmosphere, *J. Geophys. Res.*, 98, 9001–9010, doi:10.1029/93JD00031, 1993.

Griffin, R. J., Cocker, D. R., Flagan, R. C. and Seinfeld, J. H.: Organic aerosol formation from the oxidation of 25 biogenic hydrocarbons, *J. Geophys. Res.*, 104, 3555–3567, doi:10.1029/1998JD100049, 1999.

Guenther, A., Nicholas, C., Fall, R., Klinger, L., McKay, W. a., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J. and Zimmerman, P.: A global model of natural volatile organic compound emissions, *J. Geophys. Res.*, 100, 8873–8892, doi:10.1029/94JD02950, 1995.

Herron, J. T., Martinez, R. I. and Huie, R. E.: Kinetics and energetics of the criegee intermediate in the gas phase. I. The criegee intermediate in ozone-alkene reactions, *Int. J. Chem. Kinet.*, 14, 201–224, doi:10.1002/kin.550140302, 1982.

Huang, H.-L., Chao, W. and Lin, J. J.-M.: Kinetics of a Criegee intermediate that would survive high humidity and may oxidize atmospheric SO₂., *P. Natl. Acad. Sci. USA.*, 112, 10857–10862, doi:10.1073/pnas.1513149112, 30 2015.

Hyttinen, N., Kupiainen-Määttä, O., Rissanen, M. P., Muuronen, M., Ehn, M. and Kurtén, T.: Modeling the

Charging of Highly Oxidized Cyclohexene Ozonolysis Products Using Nitrate-Based Chemical Ionization., *J. Phys. Chem. A*, 119, 6339–6345, doi:10.1021/acs.jpca.5b01818, 2015.

Jagiella, S., Libuda, H. G. and Zabel, F.: Thermal stability of carbonyl radicals Part I. Straight-chain and branched C4 and C5 acyl radicals, *Phys. Chem. Chem. Phys.*, 2, 1175–1181, doi:10.1039/a909557h, 2000.

5 Jen, C. N., Hanson, D. R. and McMurry, P. H.: Towards Reconciling Measurements of Atmospherically Relevant Clusters by Chemical Ionization Mass Spectrometry and Mobility Classification/Vapor Condensation, *Aerosol Sci. Technol.*, 49, i-iii, doi:10.1080/02786826.2014.1002602, 2015.

Johnson, D. and Marston, G.: The gas-phase ozonolysis of unsaturated volatile organic compounds in the troposphere, *Chem. Soc. Rev.*, 37, 699–716, doi:10.1039/b704260b, 2008.

10 Jokinen, T., Sipilä, M., Junninen, H., Ehn, M., Lönn, G., Hakala, J., Petäjä, T., Mauldin, R. L., Kulmala, M. and Worsnop, D. R.: Atmospheric sulphuric acid and neutral cluster measurements using CI-APi-TOF, *Atmos. Chem. Phys.*, 12, 4117–4125, doi:10.5194/acp-12-4117-2012, 2012.

Jokinen, T., Sipilä, M., Richters, S., Kerminen, V.-M., Paasonen, P., Stratmann, F., Worsnop, D., Kulmala, M., Ehn, M., Herrmann, H. and Berndt, T.: Rapid Autoxidation Forms Highly Oxidized RO₂ Radicals in the 15 Atmosphere, *Angew. Chemie Int. Ed.*, 53, 14596–14600, doi:10.1002/anie.201408566, 2014.

Jokinen, T., Berndt, T., Makkonen, R., Kerminen, V.-M., Junninen, H., Paasonen, P., Stratmann, F., Herrmann, H., Guenther, A. B., Worsnop, D. R., Kulmala, M., Ehn, M. and Sipilä, M.: Production of extremely low volatile organic compounds from biogenic emissions: Measured yields and atmospheric implications., *P. Natl. Acad. Sci. USA.*, 112, 7123–7128, doi:10.1073/pnas.1423977112, 2015.

20 Junninen, H., Lauri, A., Keronen, P., Aalto, P., Hiltunen, V., Hari, P. and Kulmala, M.: Smart-SMEAR: On-line data exploration and visualization tool for SMEAR stations, *Boreal Environ. Res.*, 14, 447–457, 2009.

Junninen, H., Ehn, M., Petäjä, T., Luosujärvi, L., Kotiaho, T., Kostiainen, R., Rohner, U., Gonin, M., Fuhrer, K., Kulmala, M. and Worsnop, D. R.: A high-resolution mass spectrometer to measure atmospheric ion composition, *Atmos. Meas. Tech.*, 3, 1039–1053, doi:10.5194/amt-3-1039-2010, 2010.

25 Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagné, S., Ickes, L., Kürten, A., Kupc, A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S., Tsagkogeorgas, G., Wimmer, D., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Downard, A., Ehn, M., Flagan, R. C., Haider, S., Hansel, A., Hauser, D., Jud, W., Junninen, H., Kreissl, F., Kvashin, A., Laaksonen, A., Lehtipalo, K., Lima, J., Lovejoy, E. R., Makhmutov, V., Mathot, S., Mikkilä, J., Minginette, P., Mogo, S., Nieminen, T., Onnela, A., 30 Pereira, P., Petäjä, T., Schnitzhofer, R., Seinfeld, J. H., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Vanhanen, J., Viisanen, Y., Vrtala, A., Wagner, P. E., Walther, H., Weingartner, E., Wex, H., Winkler, P. M., Carslaw, K. S., Worsnop, D. R., Baltensperger, U. and Kulmala, M.: Role of sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation, *Nature*, 476, 429–433, doi:10.1038/nature10343, 2011.

Kirkby, J., Duplissy, J., Sengupta, K., Frege, C., Gordon, H., Williamson, C., Heinritzi, M., Simon, M., Yan, C., 35 Almeida, J., Tröstl, J., Nieminen, T., Ortega, I. K., Wagner, P., Adamov, A., Amorim, A., Bernhammer, A.-K., Bianchi, F., Breitenlechner, M., Brilke, S., Chen, X., Craven, J., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A.,

Fuchs, C., Guida, R., Hakala, J., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Kim, J., Krapf, M., Kürten, A., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Molteni, U., Onnela, A., Peräkylä, O., Piel, F., Petäjä, T., Praplan, A. P., Pringle, K., Rap, A., Richards, N. A. D., Riipinen, I., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Scott, C. E., Seinfeld, J. H., Sipilä, M., Steiner, G., Stozhkov, Y., Stratmann, F., 5 Tomé, A., Virtanen, A., Vogel, A. L., Wagner, A. C., Wagner, P. E., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P., Zhang, X., Hansel, A., Dommen, J., Donahue, N. M., Worsnop, D. R., Baltensperger, U., Kulmala, M., Carslaw, K. S. and Curtius, J.: Ion-induced nucleation of pure biogenic particles, *Nature*, 533, 521–526, doi:10.1038/nature17953, 2016.

10 Kulmala, M., Dal Maso, M., Mäkelä, J. M., Pirjola, L., Väkevä, M., Aalto, P., Miikkulainen, P., Hämeri, K. and O'Dowd, C. D.: On the formation, growth and composition of nucleation mode particles, *Tellus, Ser. B Chem. Phys. Meteorol.*, 53, 479–490, doi:10.1034/j.1600-0889.2001.d01-33.x, 2001.

15 Kulmala, M., Vehkamäki, H., Petäjä, T., Dal Maso, M., Lauri, A., Kerminen, V.-M., Birmili, W. and McMurry, P. H.: Formation and growth rates of ultrafine atmospheric particles: a review of observations, *J. Aerosol Sci.*, 35, 143–176, doi:10.1016/j.jaerosci.2003.10.003, 2004.

20 Kürten, A., Rondo, L., Ehrhart, S. and Curtius, J.: Performance of a corona ion source for measurement of sulfuric acid by chemical ionization mass spectrometry, *Atmos. Meas. Tech.*, 4, 437–443, doi:10.5194/amt-4-437-2011, 2011.

Kürten, A., Rondo, L., Ehrhart, S. and Curtius, J.: Calibration of a chemical ionization mass spectrometer for the measurement of gaseous sulfuric acid, *J. Phys. Chem. A*, 116, 6375–6386, doi:10.1021/jp212123n, 2012.

25 20 Kürten, A., Jokinen, T., Simon, M., Sipilä, M., Sarnela, N., Junninen, H., Adamov, A., Almeida, J., Amorim, A., Bianchi, F., Breitenlechner, M., Dommen, J., Donahue, N. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Hakala, J., Hansel, A., Heinritzi, M., Hutterli, M., Kangasluoma, J., Kirkby, J., Laaksonen, A., Lehtipalo, K., Leiminger, M., Makhmutov, V., Mathot, S., Onnela, A., Petäjä, T., Praplan, A. P., Riccobono, F., Rissanen, M. P., Rondo, L., Schobesberger, S., Seinfeld, J. H., Steiner, G., Tomé, A., Tröstl, J., Winkler, P. M., Williamson, C., Wimmer, D., Ye, P., Baltensperger, U., Carslaw, K. S., Kulmala, M., Worsnop, D. R. and Curtius, J.: Neutral molecular cluster formation of sulfuric acid–dimethylamine observed in real time under atmospheric conditions, *P. Natl. Acad. Sci. USA*, 111, 15019–15024, doi:10.1073/pnas.1404853111, 2014.

30 Kurtén, T., Loukonen, V., Vehkamäki, H. and Kulmala, M.: Amines are likely to enhance neutral and ion-induced sulfuric acid–water nucleation in the atmosphere more effectively than ammonia, *Atmos. Chem. Phys.*, 8, 4095–4103, doi:10.5194/acp-8-4095-2008, 2008.

Lindinger, W., Hansel, A. and Jordan, A.: On-line monitoring of volatile organic compounds at pptv levels by means of proton-transfer-reaction mass spectrometry (PTR-MS) medical applications, food control and environmental research, *Int. J. Mass Spectrom. Ion Process.*, 173, 191–241, doi:10.1016/S0168-1176(97)00281-4, 1998.

35 Lucas, D. D. and Prinn, R. G.: Parametric sensitivity and uncertainty analysis of dimethylsulfide oxidation in the clear-sky remote marine boundary layer, *Atmos. Chem. Phys.*, 5, 1505–1525, doi:10.5194/acp-5-1505-2005, 2005.

Mauldin III, R. L., Berndt, T., Sipilä, M., Paasonen, P., Petäjä, T., Kim, S., Kurtén, T., Stratmann, F., Kerminen, V.-M. and Kulmala, M.: A new atmospherically relevant oxidant of sulphur dioxide, *Nature*, 488, 193–196, doi:10.1038/nature11278, 2012.

5 Mentel, T. F., Wildt, J., Kiendler-Scharr, A., Kleist, E., Tillmann, R., Dal Maso, M., Fisseha, R., Hohaus, T., Spahn, H., Uerlings, R., Wegener, R., Griffiths, P. T., Dinar, E., Rudich, Y. and Wahner, A.: Photochemical production of aerosols from real plant emissions, *Atmos. Chem. Phys.*, 9, 4387–4406, doi:10.5194/acp-9-4387-2009, 2009.

10 Mikkonen, S., Romakkaniemi, S., Smith, J. N., Korhonen, H., Petäjä, T., Plass-Duelmer, C., Boy, M., McMurry, P. H., Lehtinen, K. E. J., Joutsensaari, J., Hamed, A., Mauldin III, R. L., Birmili, W., Spindler, G., Arnold, F., Kulmala, M. and Laaksonen, A.: A statistical proxy for sulphuric acid concentration, *Atmos. Chem. Phys.*, 11, 11319–11334, doi:10.5194/acp-11-11319-2011, 2011.

Mirme, S. and Mirme, A.: The mathematical principles and design of the NAIS – a spectrometer for the measurement of cluster ion and nanometer aerosol size distributions, *Atmos. Meas. Tech.*, 6, 1061–1071, doi:10.5194/amt-6-1061-2013, 2013.

15 Murphy, S. M., Sorooshian, A., Kroll, J. H., Ng, N. L., Chhabra, P., Tong, C., Surratt, J. D., Knipping, E., Flagan, R. C. and Seinfeld, J. H.: Secondary aerosol formation from atmospheric reactions of aliphatic amines, *Atmos. Chem. Phys.*, 7, 2313–2337, doi:10.5194/acp-7-2313-2007, 2007.

20 O'Dowd, C. D., Jimenez, J. L., Bahreini, R., Flagan, R. C., Seinfeld, J. H., Hämeri, K., Pirjola, L., Kulmala, M., Jennings, S. G. and Hoffmann, T.: Marine aerosol formation from biogenic iodine emissions., *Nature*, 417, 632–636, doi:10.1038/nature00775, 2002.

25 Paasonen, P., Olenius, T., Kupiainen, O., Kurtén, T., Petäjä, T., Birmili, W., Hamed, A., Hu, M., Huey, L. G., Plass-Duelmer, C., Smith, J. N., Wiedensohler, A., Loukonen, V., McGrath, M. J., Ortega, I. K., Laaksonen, A., Vehkamäki, H., Kerminen, V.-M. and Kulmala, M.: On the formation of sulphuric acid – amine clusters in varying atmospheric conditions and its influence on atmospheric new particle formation, *Atmos. Chem. Phys.*, 12, 9113–9133, doi:10.5194/acp-12-9113-2012, 2012.

Petäjä, T., Mauldin, III, R. L., Kosciuch, E., McGrath, J., Nieminen, T., Boy, M., Adamov, A., Kotiaho, T. and Kulmala, M.: Sulfuric acid and OH concentrations in a boreal forest site, *Atmos. Chem. Phys.*, 9, 7435–7448, doi:10.5194/acp-9-7435-2009, 2009.

30 Praplan, A. P., Schobesberger, S., Bianchi, F., Rissanen, M. P., Ehn, M., Jokinen, T., Junninen, H., Adamov, A., Amorim, A., Dommen, J., Duplissy, J., Hakala, J., Hansel, A., Heinritzi, M., Kangasluoma, J., Kirkby, J., Krapf, M., Kürten, A., Lehtipalo, K., Riccobono, F., Rondo, L., Sarnela, N., Simon, M., Tomé, A., Tröstl, J., Winkler, P. M., Williamson, C., Ye, P., Curtius, J., Baltensperger, U., Donahue, N. M., Kulmala, M. and Worsnop, D. R.: Elemental composition and clustering behaviour of α -pinene oxidation products for different oxidation conditions, *Atmos. Chem. Phys.*, 15, 4145–4159, doi:10.5194/acp-15-4145-2015, 2015.

35 Riccobono, F., Schobesberger, S., Scott, C. E., Dommen, J., Ortega, I. K., Rondo, L., Almeida, J., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Downard, A., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C.,

Franchin, A., Hansel, A., Junninen, H., Kajos, M., Keskinen, H., Kupc, A., Kurten, A., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Nieminen, T., Onnela, A., Petaja, T., Praplan, A. P., Santos, F. D., Schallhart, S., Seinfeld, J. H., Sipila, M., Spracklen, D. V., Stozhkov, Y., Stratmann, F., Tome, A., Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Wimmer, D., Carslaw, K. 5 S., Curtius, J., Donahue, N. M., Kirkby, J., Kulmala, M., Worsnop, D. R. and Baltensperger, U.: Oxidation Products of Biogenic Emissions Contribute to Nucleation of Atmospheric Particles, *Science*, 344, 717–721, doi:10.1126/science.1243527, 2014.

Riipinen, I., Pierce, J. R., Yli-Juuti, T., Nieminen, T., Häkkinen, S., Ehn, M., Junninen, H., Lehtipalo, K., Petäjä, T., Slowik, J., Chang, R., Shantz, N. C., Abbatt, J., Leaitch, W. R., Kerminen, V.-M., Worsnop, D. R., Pandis, S. 10 N., Donahue, N. M. and Kulmala, M.: Organic condensation: a vital link connecting aerosol formation to cloud condensation nuclei (CCN) concentrations, *Atmos. Chem. Phys.*, 11, 3865–3878, doi:10.5194/acp-11-3865-2011, 2011.

Rinne, J., Ruuskanen, T. and Reissell, A.: On-line PTR-MS measurements of atmospheric concentrations of volatile organic compounds in a European boreal forest ecosystem, *Boreal Env. Res.*, 10, 425–436, 2005.

Rissanen, M. P., Kurtén, T., Sipilä, M., Thornton, J. a, Kangasluoma, J., Sarnela, N., Junninen, H., Jørgensen, S., Schallhart, S., Kajos, M. K., Taipale, R., Springer, M., Mentel, T. F., Ruuskanen, T., Petäjä, T., Worsnop, D. R., Kjaergaard, H. G. and Ehn, M.: The formation of highly oxidized multifunctional products in the ozonolysis of cyclohexene., *J. Am. Chem. Soc.*, 136, 15596–15606, doi:10.1021/ja507146s, 2014.

Rondo, L., Kürten, A., Ehrhart, S., Schobesberger, S., Franchin, A., Junninen, H., Petäjä, T., Sipilä, M., Worsnop, 20 D. R. and Curtius, J.: Effect of ions on the measurement of sulfuric acid in the CLOUD experiment at CERN, *Atmos. Meas. Tech.*, 7, 3849–3859, doi:10.5194/amt-7-3849-2014, 2014.

Rondo, L., Ehrhart, S., Kürten, A., Adamov, A., Bianchi, F., Breitenlechner, M., Duplissy, J., Franchin, A., Dommen, J., Donahue, N. M., Dunne, E. M., Flagan, R. C., Hakala, J., Hansel, A., Keskinen, H., Kim, J., Jokinen, T., Lehtipalo, K., Leiminger, M., Praplan, A. P., Riccobono, F., Rissanen, M. P., Sarnela, N., Schobesberger, S., 25 Simon, M., Sipilä, M., Smith, J. N., Tomé, A., Tröstl, J., Tsagkogeorgas, G., Vaattovaara, P., Winkler, P. M., Williamson, C., Wimmer, D., Baltensperger, U., Kirkby, J., Kulmala, M., Petäjä, T., Worsnop, D. R. and Curtius, J.: Effect of dimethylamine on the gas phase sulfuric acid concentration measured by Chemical ionization Mass Spectrometry, *J. Geophys. Res. Atmos.*, 121, 3036–3049, doi:10.1002/2015JD023868, 2016.

Sarnela, N., Jokinen, T., Nieminen, T., Lehtipalo, K., Junninen, H., Kangasluoma, J., Hakala, J., Taipale, R., 30 Schobesberger, S., Sipilä, M., Larnimaa, K., Westerholm, H., Heijari, J., Kerminen, V.-M., Petäjä, T. and Kulmala, M.: Sulphuric acid and aerosol particle production in the vicinity of an oil refinery, *Atmos. Environ.*, 119, 156–166, doi:10.1016/j.atmosenv.2015.08.033, 2015.

Seinfeld, J. H. and Pankow, J. F.: Organic atmospheric particulate material., *Annu. Rev. Phys. Chem.*, 54, 121–140, doi:10.1146/annurev.physchem.54.011002.103756, 2003.

Sihto, S.-L., Kulmala, M., Kerminen, V.-M., Dal Maso, M., Petäjä, T., Riipinen, I., Korhonen, H., Arnold, F., Janson, R., Boy, M., Laaksonen, A. and Lehtinen, K. E. J.: Atmospheric sulphuric acid and aerosol formation: implications from atmospheric measurements for nucleation and early growth mechanisms, *Atmos. Chem. Phys.*,

6, 4079–4091, doi:10.5194/acp-6-4079-2006, 2006.

5 Sipilä, M., Jokinen, T., Berndt, T., Richters, S., Makkonen, R., Donahue, N. M., Mauldin III, R. L., Kurtén, T., Paasonen, P., Sarnela, N., Ehn, M., Junninen, H., Rissanen, M. P., Thornton, J., Stratmann, F., Herrmann, H., Worsnop, D. R., Kulmala, M., Kerminen, V.-M., Petäjä, T., Kurten, T., Paasonen, P., Sarnela, N., Ehn, M., Junninen, H., Rissanen, M. P., Thornton, J., Stratmann, F., Herrmann, H., Worsnop, D. R., Kulmala, M., Kerminen, V.-M. and Petäjä, T.: Reactivity of stabilized Criegee intermediates (sCIs) from isoprene and monoterpane ozonolysis toward SO₂ and organic acids, *Atmos. Chem. Phys.*, 14, 12143–12153, doi:10.5194/acp-14-12143-2014, 2014.

10 Sipilä, M., Sarnela, N., Jokinen, T., Henschel, H., Junninen, H., Kontkanen, J., Richters, S., Kangasluoma, J., Franchin, A., Peräkylä, O., Rissanen, M. P., Ehn, M., Vehkamäki, H., Kurten, T., Berndt, T., Petäjä, T., Worsnop, D., Ceburnis, D., Kerminen, V.-M., Kulmala, M. and O'Dowd, C.: Molecular-scale evidence of aerosol particle formation via sequential addition of HIO₃, *Nature*, 537, 532–534, doi:10.1038/nature19314, 2016.

15 Steiner, G., Attoui, M., Wimmer, D. and Reischl, G. P.: A Medium Flow, High-Resolution Vienna DMA Running in Recirculating Mode, *Aerosol Sci. Technol.*, 44, 308–315, doi: 10.1080/02786821003636763, 2010.

20 15 Su, T. and Bowers, M. T.: Ion-polar molecule collisions—effect of molecular size on ion-polar molecule rate constants., *J. Am. Chem. Soc.*, 95, 7609–7610, 1973.

20 Taatjes, C. A., Welz, O., Eskola, A. J., Savee, J. D., Scheer, A. M., Shallcross, D. E., Rotavera, B., Lee, E. P. F., Dyke, J. M., Mok, D. K. W., Osborn, D. L., Percival, C. J., Johnson, D., Marston, G., Harrison, R. M., Gäb, S., Hellpointner, E., Turner, W. V., Korte, F., Welz, O., Asatryan, R., Bozzelli, J. W., Taatjes, C. A., Taatjes, C. A., Beames, J. M., Liu, F., Lu, L., Lester, M. I., Vereecken, L., Harder, H., Novelli, A., Mauldin, R. L., Carlsson, P. T. M., Keunecke, C., Krüger, B. C., Maaß, M.-C., Zeuch, T., Berndt, T., Anglada, J. M., González, J., Torrent-Sucarrat, M., Kuwata, K. T., Hermes, M. R., Carlson, M. J., Zogg, C. K., Huang, H., Eskola, A. J., Taatjes, C. A., Osborn, D. L., Taatjes, C. A., Taatjes, C. A., Sweigart, D. A., Turner, D. W., Anglada, J. M., Bofill, J. M., Olivella, S., Solé, A., Sebbar, N., Bockhorn, H., Bozzelli, J. W., Donahue, N. M., Drozd, G. T., Epstein, S. A., Presto, A. A., Kroll, J. H., Kurtén, T., Donahue, N. M., Fenske, J. D., Hasson, A. S., Ho, A. W., Paulson, S. E., Jiang, L., Xu, Y. S., Ding, A. Z., Kurtén, T., Lane, J. R., Jørgensen, S., Kjaergaard, H. G., Jenkin, M. E., Saunders, S. M., Pilling, M. J., Alderdice, D. S., Dixon, R. N., Vereecken, L., Francisco, J. S., Nguyen, M. T., Nguyen, T. L., Ngan, V. T., Nguyen, H. M. T., Knizia, G., Adler, T. B., Werner, H.-J., Werner, H.-J., Peterson, K. A., Adler, T. B., Werner, H.-J., Hill, J. G., Mazumder, S., Peterson, K. A., et al.: Direct measurements of conformer-dependent reactivity of the Criegee intermediate CH₃CHOO., *Science*, 340, 177–180, doi:10.1126/science.1234689, 2013.

30 Tröstl, J., Chuang, W. K., Gordon, H., Heinritzi, M., Yan, C., Molteni, U., Ahlm, L., Frege, C., Bianchi, F., Wagner, R., Simon, M., Lehtipalo, K., Williamson, C., Craven, J. S., Duplissy, J., Adamov, A., Almeida, J., Bernhammer, A.-K., Breitenlechner, M., Brilke, S., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Gysel, M., Hansel, A., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Keskinen, H., Kim, J., Krapf, M., Kürten, A., Laaksonen, A., Lawler, M., Leiminger, M., Mathot, S., Möhler, O., Nieminen, T., Onnela, A., Petäjä, T., Piel, F. M., Miettinen, P., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Sengupta, K., Sipilä, M., Smith, J. N., Steiner, G., Tomè, A., Virtanen, A., Wagner, A. C., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P., Carslaw, K. S., Curtius, J., Dommen, J., Kirkby, J., Kulmala, M., Riipinen, I., Worsnop,

D. R., Donahue, N. M. and Baltensperger, U.: The role of low-volatility organic compounds in initial particle growth in the atmosphere, *Nature*, 533, 527–531, doi:10.1038/nature18271, 2016.

Vereecken, L., Harder, H. and Novelli, A.: The reaction of Criegee intermediates with NO, RO₂, and SO₂, and their fate in the atmosphere, *Phys. Chem. Chem. Phys.*, 14, 14682–14695, doi:10.1039/c2cp42300f, 2012.

5 Vereecken, L., Harder, H. and Novelli, A.: The reactions of Criegee intermediates with alkenes, ozone, and carbonyl oxides, *Phys. Chem. Chem. Phys.*, 16, 4039–4049, doi:10.1039/c3cp54514h, 2014.

Voigtländer, J., Duplissy, J., Rondo, L., Kürten, A. and Stratmann, F.: Numerical simulations of mixing conditions and aerosol dynamics in the CERN CLOUD chamber, *Atmos. Chem. Phys.*, 12, 2205–2214, doi:10.5194/acp-12-2205-2012, 2012.

10 Wang, S. C. and Flagan, R. C.: Scanning Electrical Mobility Spectrometer, *Aerosol Sci. Technol.*, 13, 230–240, doi:10.1080/02786829008959441, 1990.

Weber, R. J., Marti, J. J., McMurry, P. H., Eisele, F. L., Tanner, D. J. and Jefferson, A.: Measured atmospheric new particle formation rates: implications for nucleation mechanisms, *Chem. Eng. Commun.*, 151, 53–64, doi:10.1080/00986449608936541, 1996.

15 Weber, R. J., Marti, J. J., McMurry, P. H., Eisele, F. L., Tanner, D. J. and Jefferson, A.: Measurements of new particle formation and ultrafine particle growth rates at a clean continental site, *J. Geophys. Res.*, 102, 4375–4385, doi:10.1029/96JD03656, 1997.

Wehner, B., Petäjä, T., Boy, M., Engler, C., Birmili, W., Tuch, T., Wiedensohler, A. and Kulmala, M.: The contribution of sulfuric acid and non-volatile compounds on the growth of freshly formed atmospheric aerosols, 20 Geophys. Res. Lett., 32, doi:10.1029/2005GL023827, 2005.

Welz, O., Savee, J. D., Osborn, D. L., Vasu, S. S., Percival, C. J., Shallcross, D. E. and Taatjes, C. A.: Direct kinetic measurements of Criegee intermediate (CH₂OO) formed by reaction of CH₂I with O₂., *Science*, 335, 204–207, doi:10.1126/science.1213229, 2012.

25 Yan, C., Nie, W., Äijälä, M., Rissanen, M. P., Canagaratna, M. R., Massoli, P., Junninen, H., Jokinen, T., Sarnela, N., Häme, S. A. K., Schobesberger, S., Canonaco, F., Yao, L., Prévôt, A. S. H., Petäjä, T., Kulmala, M., Sipilä, M., Worsnop, D. R. and Ehn, M.: Source characterization of highly oxidized multifunctional compounds in a boreal forest environment using positive matrix factorization, *Atmos. Chem. Phys.*, 16, 12715–12731, doi:10.5194/acp-16-12715-2016, 2016.

30 Yao, L., Ma, Y., Wang, L., Zheng, J., Khalizov, A., Chen, M., Zhou, Y., Qi, L. and Cui, F.: Role of stabilized Criegee Intermediate in secondary organic aerosol formation from the ozonolysis of α -cedrene, *Atmos. Environ.*, 94, 448–457, doi:10.1016/j.atmosenv.2014.05.063, 2014.

Appendix A

Table A1: The elemental compositions and exact masses of most abundant isotopes of the HOM compounds that were added together to make the total HOM

| Elemental composition | Exact mass (Th) |
|--|-----------------|
| C ₇ H ₁₀ O ₄ NO ₃ ⁻ | 220.0463 |
| C ₅ H ₆ O ₆ NO ₃ ⁻ | 224.0048 |
| C ₅ H ₆ O ₇ NO ₃ ⁻ | 239.9997 |
| C ₈ H ₁₂ O ₇ NO ₃ ⁻ | 282.0461 |
| C ₈ H ₁₂ O ₈ NO ₃ ⁻ | 298.0416 |
| C ₁₀ H ₁₄ O ₇ NO ₃ ⁻ | 308.0623 |
| C ₉ H ₁₂ O ₈ NO ₃ ⁻ | 310.0416 |
| C ₁₀ H ₁₆ O ₇ NO ₃ ⁻ | 310.0780 |
| C ₈ H ₁₂ O ₉ NO ₃ ⁻ | 314.0365 |
| C ₁₀ H ₁₄ O ₈ NO ₃ ⁻ | 324.0572 |
| C ₁₀ H ₁₅ O ₈ NO ₃ ⁻ | 325.0651 |
| C ₉ H ₁₂ O ₉ NO ₃ ⁻ | 326.0365 |
| C ₁₀ H ₁₆ O ₈ NO ₃ ⁻ | 326.0729 |
| C ₉ H ₁₄ O ₉ NO ₃ ⁻ | 328.0521 |
| C ₁₀ H ₁₄ O ₉ NO ₃ ⁻ | 340.0521 |
| C ₉ H ₁₂ O ₁₀ NO ₃ ⁻ | 342.0314 |
| C ₁₀ H ₁₆ O ₉ NO ₃ ⁻ | 342.0678 |
| C ₁₀ H ₁₄ O ₁₀ NO ₃ ⁻ | 356.0471 |
| C ₁₀ H ₁₅ O ₁₀ NO ₃ ⁻ | 357.0549 |
| C ₉ H ₁₂ O ₁₁ NO ₃ ⁻ | 358.0263 |
| C ₁₀ H ₁₆ O ₁₀ NO ₃ ⁻ | 358.0627 |
| C ₁₀ H ₁₄ O ₁₁ NO ₃ ⁻ | 372.0420 |
| C ₉ H ₁₂ O ₁₂ NO ₃ ⁻ | 374.0212 |
| C ₁₀ H ₁₆ O ₁₁ NO ₃ ⁻ | 374.0576 |
| C ₁₀ H ₁₄ O ₁₃ NO ₃ ⁻ | 404.0318 |
| C ₁₅ H ₂₈ O ₁₂ NO ₃ ⁻ | 462.1464 |
| C ₁₇ H ₂₄ O ₁₁ NO ₃ ⁻ | 466.1202 |
| C ₁₇ H ₂₆ O ₁₁ NO ₃ ⁻ | 468.1359 |
| C ₁₈ H ₂₆ O ₁₁ NO ₃ ⁻ | 480.1359 |
| C ₁₄ H ₂₀ O ₁₅ NO ₃ ⁻ | 490.0686 |
| C ₁₉ H ₂₈ O ₁₁ NO ₃ ⁻ | 494.1515 |
| C ₂₀ H ₃₂ O ₁₁ NO ₃ ⁻ | 510.1828 |
| C ₁₇ H ₂₆ O ₁₄ NO ₃ ⁻ | 516.1206 |
| C ₂₀ H ₃₀ O ₁₂ NO ₃ ⁻ | 524.1621 |
| C ₁₉ H ₂₈ O ₁₃ NO ₃ ⁻ | 526.1414 |
| C ₁₈ H ₂₆ O ₁₄ NO ₃ ⁻ | 528.1206 |
| C ₁₈ H ₂₈ O ₁₄ NO ₃ ⁻ | 530.1363 |
| C ₁₇ H ₂₆ O ₁₅ NO ₃ ⁻ | 532.1155 |
| C ₂₀ H ₃₀ O ₁₃ NO ₃ ⁻ | 540.1570 |
| C ₂₀ H ₃₂ O ₁₃ NO ₃ ⁻ | 542.1727 |
| C ₁₇ H ₂₆ O ₁₆ NO ₃ ⁻ | 548.1105 |
| C ₂₀ H ₃₀ O ₁₄ NO ₃ ⁻ | 556.1519 |
| C ₁₈ H ₂₈ O ₁₆ NO ₃ ⁻ | 562.1261 |
| C ₂₀ H ₃₀ O ₁₅ NO ₃ ⁻ | 572.1468 |
| C ₂₀ H ₃₂ O ₁₅ NO ₃ ⁻ | 574.1625 |
| C ₂₀ H ₃₀ O ₁₆ NO ₃ ⁻ | 588.1418 |
| C ₁₈ H ₂₈ O ₁₈ NO ₃ ⁻ | 594.1159 |
| C ₂₀ H ₃₀ O ₁₈ NO ₃ ⁻ | 620.1316 |