

1 **Low Carbon Energy Generates Public Health Savings in**  
 2 **California**

3 Christina B. Zapata<sup>1</sup>, Chris Yang<sup>2</sup>, Sonia Yeh<sup>2</sup>, Joan Ogden<sup>2</sup>, Michael J. Kleeman<sup>1</sup>

4 <sup>1</sup> Department of Civil and Environmental Engineering, University of California – Davis, Davis, California, USA

5 <sup>2</sup> Institute of Transportation Studies, University of California – Davis, Davis, California, USA

6 Correspondence to: Michael J. Kleeman ([mjkleeman@ucdavis.edu](mailto:mjkleeman@ucdavis.edu))

7  
 8 **Supplementary Information**

9  
 10 **Table S1: Integrated GHG and co-benefit assessments that encompass California and examines criteria pollutant**  
 11 **emissions, air pollution simulations, health benefits, and valued health savings for 2050.**

Author	Region, resolution	Time-frame	Energy model, scenario, policy, economics, sectors	Emission Estimation	Climate/ Atmospheric Model	Short-lived Pollutant(s)	Additional Impacts analyzed
This study	California, 4 km x 4 km (~0.05° x 0.05°)	2050	2 scenarios from CA-TIMES: BAU, GHG-Step. System-wide cost minimization optimization. (Yang, Yeh et al. 2014, Yang, Yeh et al. 2015)	16km <sup>2</sup> spatial emissions scaling for each sector/mode/technology/fuel (fossil and alternative) based on literature, and existing current emission inventories	UCD/CIT Airshed (Kleeman, Cass et al. 1997, Kleeman and Cass 2001, Ying, Fraser et al. 2007)	PM <sub>2.5</sub> (speciation and size resolved), O <sub>3</sub>	Mortality (deaths, death rate), cost (US\$)
(Jacobson, Delucchi et al. 2014)	California, 0.6° x 0.5°	2020, 2030, 2050	2 scenarios: reference, 100% Wind, Water, Sunlight (WWS) by 2050	Measured California county 2010-2012 PM2.5 concentration and 8-hour ozone level	GATOR-GCMOM, Gas, Aerosol, Transport, Radiation, General Circulation, Mesoscale, and Ocean Model (Jacobson, Kaufman et al. 2007)	PM <sub>2.5</sub> , O <sub>3</sub>	Mortality (deaths), costs (\$US), job loss (jobs, earnings)
Garcia-Menendez, Saari et al. (2015)	US contiguous, 1.9° x 2.5°	2050, 2100	3 scenarios: reference, 4.5 W m <sup>-2</sup> and 3.7 W m <sup>-2</sup> stabilization to 2100	Emissions Prediction and Policy Analysis (EPPA) Model (Paltsev, Reilly et al. 2005)	CAM-Chem (Lamarque, Emmons et al. 2012)	O <sub>3</sub> , PM <sub>2.5</sub>	Years of life saved, Cost/Benefit (% GDP), % ton <sup>-1</sup>

13  
14

**Table S2 (cont.):**

Author	Region, resolution	Time-frame	Energy model, scenario, policy, economics, sectors	Emission Estimation	Climate/ Atmospheric Model	Short-lived Pollutant(s)	Additional Impacts analyzed
Shindell, Kyulenstierna et al. (2012)	Global, 4° x 5°	2050	manually selected policy measures based on GAINS	Greenhouse Gas and Air Pollution Interactions and Synergies (GAINS) model used for reductions (GAINS Development Team. International Institute for Applied Systems Analysis (IIASA) 2009)	ECHAM5-HAMMOZ (Pozzoli, Bey et al. 2008) and GISS PUCINI (Shindell, Faluvegi et al. 2006)	CH <sub>4</sub> , BC	mortality, crop loss, Cost (\$ tonne <sup>-1</sup> ), temperature, RF (W m <sup>-2</sup> ),
Smith, West et al. (2011) West, Smith et al. (2013)	Global, 2° x 2.5°/1000km <sup>2</sup>	2030, 2050, 2100	2 scenarios from Global Change Assessment Model (GCAM) : RCP4.5 scenario (Thomson, Calvin et al. 2011), global carbon price	ERB module in Mini Climate Assessment (MiniCAM) model (Brenkert, Smith et al. 2003)	MAGICC6 (Model for the Assessment of Greenhouse-gas Induced Climate Change) module in MiniCAM model (Brenkert, Smith et al. 2003, Meinshausen, Raper et al. 2011)	PM <sub>2.5</sub> , O <sub>3</sub>	Mortality, cost (\$/tonne)
van Aardenne, Dentener et al. (2010)	Global, 1x1 North America, Europe, India, China, 3x2 remaining Northern Hemisphere, 6x 4 Southern Hemisphere	2050	6 scenarios: climate/air policy alone or together with various country adoption from Prospective Outlook for the Long term Energy System (POLES) model (Russ, Wiesenthal et al. 2007)	Emission Database for Global Atmospheric Research (EDGAR) v4 and GAINS	TM5 (Tracer Model ver. 5) (Krol, Houweling et al. 2005)	PM, O <sub>3</sub>	Crop loss reduction (%), years life lost/ life expectancy (years/person), radiative forcing (W/m <sup>2</sup> )

15

16 Numerous regional and worldwide studies have examined how climate policies and energy scenarios impact air  
17 quality, health, and costs as far as 2050 (Table S1). However, the approach and assumptions associated with each  
18 study's energy/climate policy scenarios produce very different outcomes and insights. For instance, some studies  
19 produced future energy/climate policy scenarios that are heavily user defined, often on the basis that both pollutants  
20 be mitigated. Shindell, Kuylenstierna et al. (2012), used only measures that were "assumed to improve air quality"  
21 and mitigate both long-lived GHGs and short-lived criteria pollutants after ranking them by climate impact. The  
22 scenario by Jacobson, Delucchi et al. (2014) is defined by 100% wind, water, sunlight (100%WWS) renewable  
23 supply (and no fossil fuels and biofuels) versus one that is constrained to achieve 80% GHG emissions reduction  
24 such as CA-TIMES GHG-Step. These user-defined measures and energy constraints not only produce different  
25 energy projections, but air pollution/health outcomes as well. By permitting only low polluting or zero-emission  
26 technologies and policy measures, these scenarios are more heavily defined by the authors/modelers to be somewhat  
27 biased to be much cleaner and highly advantageous to human health. The extensive study by van Aardenne,  
28 Dentener et al. (2010) explores 6 scenarios with wider levels of air and/or climate policy, as well as the option of  
29 biofuel consumption, however, technology adoption is again largely dependent on user specified assumptions on  
30 shares of existing technologies. In comparison, this study did not predefine scenario conditions that ensured both  
31 criteria pollutants and GHGs would reduce.


32 In contrast, CA-TIMES (1) permits dirtier energy resources (such as biofuels) and measures that don't necessary  
33 reduce air pollution, enabling a more conservative pollution outcome, (2) reveals sectors/technology investments  
34 and fuel/energy that are most cost-effective rather than user defined choices (3) has more complex new/alternative  
35 technology choices, deployment costs, that simpler scenarios ignore. Additional differences exist in the  
36 development of the criteria emissions in CA-REMARQUE such as spatial (12 electric utility regions) resource mix  
37 growth projections based on SWITCH (Fripp 2012, Johnston, Mileva et al. 2013, Nelson, Mileva et al. 2013) 2050  
38 electricity generation projections, optimal biofacility siting locations that comply with potential air quality  
39 permitting restrictions (Tittmann, Parker et al. 2010, Parker 2012), and alternative fuel and regenerative braking  
40 emission rates based on alternative fuel emission literature.

41

42 **Table S3: Population growth rate between 2050 and 2010.**

<b>County</b>	<b>2050:2010 Population</b>	<b>County</b>	<b>2050:2010 Population</b>
<b>Alameda</b>	1.113	Orange	1.102
<b>Alpine</b>	0.985	Placer	1.562
<b>Amador</b>	1.184	Plumas	0.983
<b>Butte</b>	1.515	Riverside	1.747
<b>Calaveras</b>	1.321	Sacramento	1.452
<b>Colusa</b>	1.711	San Benito	1.479
<b>Contra Costa</b>	1.415	San Bernardino	1.594
<b>Del Norte</b>	1.127	San Diego	1.279
<b>El Dorado</b>	1.565	San Francisco	1.126
<b>Fresno</b>	1.619	San Joaquin	2.009
<b>Glenn</b>	1.351	San Luis Obispo	1.256
<b>Humboldt</b>	1.099	San Mateo	1.244
<b>Imperial</b>	1.863	Santa Barbara	1.194
<b>Inyo</b>	1.244	Santa Clara	1.205
<b>Kern</b>	2.209	Santa Cruz	1.153
<b>Kings</b>	1.706	Shasta	1.46
<b>Lake</b>	1.636	Sierra	1.125
<b>Lassen</b>	1.183	Siskiyou	1.161
<b>Los Angeles</b>	1.164	Solano	1.435
<b>Madera</b>	2.138	Sonoma	1.237
<b>Marin</b>	1.048	Stanislaus	1.673
<b>Mariposa</b>	1.264	Sutter	2.241
<b>Mendocino</b>	1.136	Tehama	1.551
<b>Merced</b>	1.941	Trinity	1.325
<b>Modoc</b>	1.119	Tulare	1.77
<b>Mono</b>	1.335	Tuolumne	1.118
<b>Monterey</b>	1.304	Ventura	1.207
<b>Napa</b>	1.354	Yolo	1.476
<b>Nevada</b>	1.425	Yuba	1.991

43  
44



46 Figure S1: 2050 population projection. The population is projected using 2010 census block group population  
 47 interpolated 16km<sup>2</sup> information, and projected for each age group within each county projections (California Department  
 48 of Finance. Demographic Research Unit 2014).

50 Both 4km domains combined constitute about 93 % of the state's projected population. The maximum population  
 51 density occurs in areas surrounding downtown Los Angeles and the San Francisco Financial District. The  
 52 population ages 35 and older represent roughly 57 % of the total all age group population for either domain.

53  
 54 **Table S4: 2050 population summary.**

Region Type	Region	Area (km <sup>2</sup> )	Population (persons)	Percentage of the CA Population (%)
<b>Domain Total</b>	CA statewide	162912.0	57,952,084	92.6 %
<b>Domain</b>	Northern/ Central CA	94464.00	22,946,980	36.7 %
<b>Basin</b>	San Francisco (SF)	14422.26	11,611,289	18.5 %
<b>Basin</b>	San Joaquin Valley (SJV)	46881.20	6,426,141	10.3 %
<b>Domain</b>	Southern CA	68448.00	35,005,104	55.9 %
<b>Basin</b>	South Coast (SC)	16773.50	26,069,030	41.6 %
<b>Basin</b>	San Diego (SD)	7756.350	4,825,970	7.7 %

55  
 56  
 57 **Table S5: California 8-hour O3 nonattainment areas and attainment dates as of June 16, 2016. Design values are listed**  
 58 **from designation in 2008-2010.**

Area Name	Design Values (ppm)	Classification	Expected Attainment Date	Number of Counties	2010 Population	Percent of Population
<b>Los Angeles-South Coast Air Basin</b>	0.112	Extreme	2031	4	15,719,485	45.4 %
<b>San Joaquin Valley</b>	0.104	Extreme	2031	8	3,842,365	11.1 %
<b>Riverside Co. (Coachella Valley)</b>	0.095	Extreme	2026	1	425,806	1.2 %
<b>Los Angeles-San Bernardino Counties (West Mojave Desert)</b>	0.099	Severe 15	2026	2	868,380	2.5 %
<b>Sacramento Metro</b>	0.095	Severe 15	2026	6	2,241,057	6.5 %
<b>Ventura County</b>	0.086	Serious	2020	1	823,262	2.4 %
<b>Kern Co (Eastern Kern)</b>	0.083	Moderate	2017	1	95,176	0.3 %
<b>San Diego County</b>	0.082	Moderate	2017	1	3,095,199	8.9 %
<b>Nevada Co. (Western part)</b>	0.079	Moderate	2017	1	82,107	0.2 %
<b>Imperial County</b>	0.078	Moderate	2017	1	174,528	0.5 %
<b>Mariposa County</b>	0.077	Moderate	2017	1	18,251	0.1 %
<b>San Francisco Bay Area</b>	0.080	Marginal	2015	9	6,973,020	20.1 %
<b>Chico (Butte County)</b>	0.079	Marginal	2015	1	220,000	0.6 %
<b>San Luis Obispo (Eastern San Luis Obispo)</b>	0.078	Marginal	2015	1	1,649	0.0 %
<b>Calaveras County</b>	0.077	Marginal	2015	1	45,578	0.1 %
<b>Tuscan Buttes</b>	0.076	Marginal	2015	1	0	0.0 %

## 61 **S1 PM<sub>2.5</sub> Species Concentration Results**

### 62 **S1.1 PM<sub>2.5</sub> Nitrogen containing species**

63 Table S3 summarizes population-weighted BAU PM<sub>2.5</sub> concentrations and changes induced by the GHG-Step  
64 scenario across different locations in California. The results in Table S3 are broken down by chemical species for  
65 an improved understanding of the different sources and atmospheric processes involved in the total particulate  
66 matter reductions. Nitrate (NO<sub>3</sub><sup>-</sup>) accounts for 17 % of the statewide population-weighted PM<sub>2.5</sub> mass concentrations  
67 in the BAU scenario (2<sup>nd</sup> most dominant chemical species) with contributions as high as 19 % for SD and as low as  
68 15 % for the SJV. Population-weighted nitrate concentrations decrease by 49 %-50 % in SD and the SJV in the  
69 GHG-Step scenario, with more modest reductions in San Francisco (-45 %) and Los Angeles (-34 %). Most of this  
70 nitrate reduction is associated with lower NO<sub>x</sub> emissions from avoided combustion through the adoption of electric  
71 power generated from solar or wind energy. Net NO<sub>x</sub> emissions from all land sources (excluding marine and  
72 aviation) are lower in the GHG-Step scenario. This trend is also reflected in atmospheric concentrations.  
73 Population-weighted gas-phase NO and NO<sub>x</sub> (NO+NO<sub>2</sub>) concentrations fall by 71 % and 46 % in the GHG-Step  
74 scenario (Table S5).

75 PM<sub>2.5</sub> ammonium (NH<sub>4</sub><sup>+</sup>) concentrations also decrease in the GHG-Step scenario which contributes strongly to  
76 overall PM<sub>2.5</sub> reductions, but this trend is not driven by the same source changes that affected nitrate. Ammonium  
77 (NH<sub>4</sub><sup>+</sup>) accounts for 0.57 μg m<sup>-3</sup> (6 %) of the population weighted mass in the BAU scenario with concentrations  
78 decreasing by 33 % - 50 % in the GHG-Step scenario. These reductions mainly stemmed from the improved  
79 management of animal waste in agricultural operations to produce biogas as a renewable energy source which had  
80 the added benefit of reducing ammonia emissions.

### 81 **S1.2 PM<sub>2.5</sub> Carbonaceous Species**

82 Elemental carbon (EC), primary and secondary organic aerosol (POA and SOA) contribute to 43 % of the statewide  
83 population-weighted PM<sub>2.5</sub> concentration in the BAU scenario making any changes to these species under the GHG-  
84 Step scenario extremely important for public health. Statewide average concentrations of PM<sub>2.5</sub> EC, POA, and SOA  
85 are predicted to decrease in the GHG-Step scenario but the effects are mixed, with both increases and decreases  
86 depending on location. There is a significant EC increase of 1.0 μg m<sup>-3</sup> in the GHG-Step scenario in the shipping  
87 lanes over the Pacific Ocean near southern California cities of Malibu, Santa Monica and south towards Manhattan  
88 Beach due to higher emissions from marine vessels but there is little population exposure associated with this  
89 change. All inland locations experience a significant decrease in EC concentrations with the largest reductions  
90 around the San Francisco Bay Area where population weighted EC concentration decreases by 38 % from a BAU  
91 value of 0.88 μg m<sup>-3</sup>. Statewide, population weighted EC concentrations decrease by 22 % accounting for a 10 %  
92 reduction in total population-weighted PM<sub>2.5</sub> mass.

93 Primary organic aerosol (POA) accounts for approximately 1/3 of PM<sub>2.5</sub> mass across all of California including large  
94 population centers in the 2050 BAU scenario, but the modest 12 % POA reductions under the GHG-Step scenario  
95 yield only a 13.9 % reduction in population-weighted PM<sub>2.5</sub> mass concentrations (3<sup>rd</sup> highest). POA reductions  
96 occur around emissions sources that are affected by GHG mitigation strategies such as natural gas reduction in the  
97 residential and commercial sectors. The population-weighted POA reductions for each basin in order of highest to  
98 lowest are SD, SJV, SC, and SF at 19 %, 16 %, 14 %, and 10 % respectively.

99 Secondary organic aerosol (SOA) contributes to 1.10 µg m<sup>-3</sup> (14 %) of population-weighted PM<sub>2.5</sub> mass in the 2050  
100 BAU scenario. Changes to population-weighted PM<sub>2.5</sub> SOA concentrations caused by the adoption of the GHG-  
101 Step scenario were not statistically significant over the majority of California and their individual contributions to  
102 PM<sub>2.5</sub> mass reduction were not analyzed further in the current study. Updated models of SOA production could  
103 change this finding but an exploration of this research topic is outside the scope of the current study.

### 104 **S1.3 PM<sub>2.5</sub> Sulfate**

105 California adopted low sulfur fuels in the decades after 1980, which greatly reduced PM<sub>2.5</sub> sulfate concentrations.  
106 Not surprisingly then, statewide population-weighted PM<sub>2.5</sub> sulfate (SO<sub>4</sub><sup>2-</sup>) concentrations are only 0.39 µg m<sup>-3</sup> in the  
107 2050 BAU scenario, which represents 5 % of PM<sub>2.5</sub> mass. Some point sources and marine vessels switch to low  
108 sulfur biofuels in the GHG-Step scenario, leading to significant decreases in sulfate concentrations in the immediate  
109 vicinity around those sources but much smaller changes across other areas. Overall, statewide population-weighted  
110 PM<sub>2.5</sub> sulfate concentrations decrease by 23 % yielding a 7 % reduction in population-weighted PM<sub>2.5</sub> mass in the  
111 GHG-Step scenario.

### 112 **S1.4 PM<sub>2.5</sub> Metals**

113 In the BAU scenario, copper (Cu), iron (Fe), and manganese (Mn) contribute to a relatively smaller percentage of  
114 PM<sub>2.5</sub> mass. Cu, Fe, and Mn concentrations align strongly with freeways and roads as well as some off-road areas.  
115 Population-weighted concentrations of Cu, Fe, and Mg have large relative reductions in the GHG-Step scenario (51  
116 % decrease) but still account for modest reductions in total PM<sub>2.5</sub> mass (9 % of the total decrease). Reductions  
117 occurred mainly along transportation corridors due to the widespread adoption of regenerative braking in electric  
118 vehicles and hybrid electric vehicles. Regenerative braking reduces brake wear emissions rich in Cu and Fe by  
119 approximately 59 %. Both of these metals are considered to be toxic in airborne particulate matter due to their  
120 contribution to oxidative stress.

121



122 **Table S5: Annual average population-weighted Business-as-Usual (BAU) concentration and percent change between BAU**  
123 **and GHG-Step scenario for each species and air basin jurisdiction. SJV, SD, SF, SC are the San Joaquin Valley, San**  
124 **Diego, San Francisco, and South Coast air basins, respectively. The statewide or CA value represents roughly 93 % of the**  
125 **population represented by both 4km high resolution northern and southern CA modeling domains. All annual average**  
126 **concentration differences between BAU and GHG-Step scenarios were found to have p-values less than 0.1 for all**  
127 **pollutants listed except for those in bold italics mainly SD and SC SOA and SD's 1-hour O<sub>3</sub>.**

Pollutant	Units	Avg.	BAU Concentration					GHG-Step Scenario Percentage Change				
			Air Basin				State	Air Basin				State
			SJV	SD	SF	SC	CA	SJV	SD	SF	SC	CA
Ozone (O <sub>3</sub> )	ppb	1-hr max.	55.9	56.8	50.1	55.2	54.1	-4.3 %	<b>+0.5 %</b>	+6.5 %	+5.1 %	+2.8 %
		8-hr max.	52.0	51.0	45.0	50.2	49.5	-3.5 %	+2.8 %	+9.5 %	+5.5 %	+3.9 %
Nitrogen Oxide (NO)	ppb	24-hr avg.	0.77	2.77	8.62	2.00	3.15	-50 %	-81 %	-82 %	-53 %	-71 %
Nitrogen Dioxide (NO <sub>2</sub> )	ppb	24-hr avg.	6.62	13.8	20.2	10.7	12.1	-35 %	-56 %	-46 %	-32 %	-39 %
Nitrogen Oxides (NO <sub>x</sub> )	ppb	24-hr avg.	7.38	16.5	28.8	12.7	15.3	-36 %	-60 %	-57 %	-35 %	-46 %
Sulfur Dioxide (SO <sub>2</sub> )	ppm	24-hr avg.	0.88	0.25	0.85	0.47	0.55	-6 %	-30 %	-20 %	-17 %	-16 %
PM <sub>0.1</sub> mass (PM <sub>0.1</sub> )	µg m <sup>-3</sup>	24-hr avg.	2.06	2.97	3.59	2.94	2.88	-35 %	-46 %	-35 %	-31 %	-34 %
PM <sub>2.5</sub> mass (PM <sub>2.5</sub> )	µg m <sup>-3</sup>	24-hr avg.	5.98	8.37	9.54	7.63	7.63	-20 %	-25 %	-19 %	-15 %	-18 %
PM <sub>2.5</sub> elemental carbon (EC)	µg m <sup>-3</sup>	24-hr avg.	0.38	0.70	0.88	0.59	0.59	-29 %	-34 %	-38 %	-8.0 %	-22 %
PM <sub>2.5</sub> primary organic aerosol (POA)	µg m <sup>-3</sup>	24-hr avg.	1.15	1.53	2.01	1.82	1.56	-16 %	-19 %	-10 %	-14 %	-12 %
PM <sub>2.5</sub> secondary organic aerosol (SOA)	µg m <sup>-3</sup>	24-hr avg.	1.03	1.18	0.85	1.19	1.10	-8.0 %	<b>+0.7 %</b>	-2.9 %	<b>+0.1 %</b>	-2.2 %
PM <sub>2.5</sub> nitrate (NO <sub>3</sub> <sup>-</sup> )	µg m <sup>-3</sup>	24-hr avg.	0.88	1.81	1.47	1.47	1.31	-50 %	-49 %	-45 %	-34 %	-39 %
PM <sub>2.5</sub> sulfate (SO <sub>4</sub> <sup>2-</sup> )	µg m <sup>-3</sup>	24-hr avg.	0.36	0.55	0.51	0.47	0.39	-13 %	-61 %	-23 %	-25 %	-23 %
PM <sub>2.5</sub> ammonium (NH <sub>4</sub> <sup>+</sup> )	µg m <sup>-3</sup>	24-hr avg.	0.44	0.76	0.67	0.63	0.57	-36 %	-50 %	-36 %	-30 %	-33 %
PM <sub>2.5</sub> copper, iron, magnesium (Cu, Fe, Mg)	µg m <sup>-3</sup>	24-hr avg.	0.16	0.35	0.23	0.28	0.25	-52 %	-53 %	-52 %	-51 %	-51 %
PM <sub>2.5</sub> metals	µg m <sup>-3</sup>	24-hr avg.	0.14	0.33	0.29	0.26	0.25	-19 %	-23 %	-17 %	-16 %	-17 %
PM <sub>2.5</sub> other	µg m <sup>-3</sup>	24-hr avg.	0.80	0.97	0.81	0.64	0.67	-2.5 %	-20 %	-1.9 %	-4.7 %	-3.1 %
PM <sub>2.5</sub> unknown	µg m <sup>-3</sup>	24-hr avg.	0.47	0.68	1.57	0.61	0.75	0.5 %	+2.6 %	+2.3 %	+1.4 %	+2.3 %
PM <sub>10</sub> mass (PM <sub>10</sub> )	µg m <sup>-3</sup>	24-hr avg.	7.85	10.9	12.4	9.04	9.60	-19 %	-21 %	-16 %	-14 %	-16 %

128  
129

## 130 **S2 PM Size Distribution Results**

131 Figure S2 summarizes predicted changes to the airborne particle size distribution in the GHG-Step scenario relative  
132 to the BAU scenario. Size distribution shifts are caused by changes to emissions sources (which each emit different  
133 particle size distributions) and changes to the rate of secondary PM formation (which condense on particles and  
134 change the size distribution).

### 135 **S2.1 Ultrafine particles ( $D_p < 0.1 \mu\text{m}$ )**

136  $\text{PM}_{0.1}$  mass (mass of particles with diameter  $D_p < 0.1 \mu\text{g m}^{-3}$ ) is substantially reduced in the GHG-Step scenario;  
137  $\text{PM}_{0.1}$  decreases are greater than any other size fraction. BAU  $\text{PM}_{0.1}$  concentrations are highest around large  
138 combustion sources associated with urban centers and many of these combustion sources are controlled through  
139 GHG mitigation measures. Peak  $\text{PM}_{0.1}$  concentrations over California in the BAU scenario are predicted to be  $5.8 \mu\text{g m}^{-3}$   
140  $-7.2 \mu\text{g m}^{-3}$  with GHG-Step reductions of about  $2.76 \mu\text{g m}^{-3}$  in SF,  $3.2 \mu\text{g m}^{-3}$  in SF and in San Bernardino.  
141  $\text{PM}_{0.1}$  decreases from a statewide population-weighted average of  $2.9 \mu\text{g m}^{-3}$  in the BAU scenario to  $1.9 \mu\text{g m}^{-3}$  (-34  
142 %) in the GHG-Step scenario. Regionally, population-weighted  $\text{PM}_{0.1}$  declines from 31 %-46 %, with minimal  
143 decline in SC and the highest decline in SD. The  $\text{PM}_{0.1}$  mass fraction of  $\text{PM}_{10}$  drops from 30 % to 23 % statewide.

### 144 **S2.2 Accumulation mode ( $0.1 < D_p < 2.5 \mu\text{m}$ )**

145 As expected, the accumulation mode has the highest BAU PM mass with a statewide population-weighted average  
146 of  $4.8 \mu\text{g m}^{-3}$  and regional concentrations ranging from  $3.9$ - $5.9 \mu\text{g m}^{-3}$ . Most regions have accumulation mode  
147 population-weighted concentrations that are roughly double the ultrafine and coarse mode concentrations. GHG  
148 mitigation strategies have less effect on the accumulation mode than on the ultrafine mode. Population-weighted  
149 concentrations of accumulation mode mass decrease by  $0.35 \mu\text{g m}^{-3}$  (-7.4 %) but their contribution to  $\text{PM}_{2.5}$  mass  
150 increases from 50 % in the BAU scenario to 54 % in the GHG-Step scenario. This statewide trend is reflected in all  
151 regions of California.

### 152 **S2.3 Coarse mode ( $D_p > 2.5 \mu\text{m}$ )**


153 Particles in the coarse mode ( $D_p > 2.5 \mu\text{m}$ ) are predicted to contribute to 20.5 % of the population-weighted  $\text{PM}_{10}$   
154 mass across California in the BAU scenario. The GHG-Step scenario is predicted to decrease concentrations of  
155 coarse mode particles by  $0.17 \mu\text{g m}^{-3}$  (-8.7 %), which is the smallest amount for any of the size ranges considered.

### 156 **S2.4 Particulate size distribution shift**

157 As the chemical composition of PM changes from fuel and energy consumption changes, the size distribution will  
158 also change. Smaller ultrafine particles of  $\text{PM}_{0.1}$  drop steeper than fine  $\text{PM}_{2.5-10}$  and coarse  $\geq \text{PM}_{10}$ . Since primary  
159 emissions from combustion have a size distribution profile that skews towards ultrafine PM, it explains why the  
160 ultrafine mode would reduce substantially. Reductions in the accumulation mode  $\text{PM}_{2.5-1}$  largely attributes to the

161 sulfate and nitrate particles that are largely caused gas-to-particle partitioning of NO<sub>x</sub> and SO<sub>x</sub> into the aqueous  
 162 phase. The coarse ≥ PM<sub>10</sub> decline likely attributes to the reduction of brake wear metals such as Cu, Mg, and Fe.

163 **Figure S1: Particulate size mode concentration for BAU scenario and GHG-Step–BAU scenario difference.**



164  
 165  
 166  
 167

**References**

168 Brenkert, A., A. Smith, S. Kim and H. Pitcher (2003). Model Documentation for the MiniCAM. U. S. D.  
 169 o. Energy. Pacific Northwest National Laboratory, Pacific Northwest National Laboratory, Richland,  
 170 Washington 99352.  
 171 California Department of Finance. Demographic Research Unit (2014). P-2: State and County Population  
 172 Projections - Race/Ethnicity and 5-Year Age Groups. 2010 through 2060. W. Schwarm. Sacramento, CA.  
 173 Fripp, M. (2012). "Switch: a planning tool for power systems with large shares of intermittent renewable  
 174 energy." *Environ Sci Technol* **46**(11): 6371-6378.  
 175 GAINS Development Team. International Institute for Applied Systems Analysis (IIASA) (2009).  
 176 GAINS Online: Tutorial for advanced users. *Version 1.0*. Schlossplatz 1, A-2361 Laxenburg, Austria,  
 177 International Institute for Applied Systems Analysis (IIASA).  
 178 Garcia-Menendez, F., R. K. Saari, E. Monier and N. E. Selin (2015). "U.S. Air Quality and Health  
 179 Benefits from Avoided Climate Change under Greenhouse Gas Mitigation." *Environ Sci Technol* **49**(13):  
 180 7580-7588.  
 181 Jacobson, M. Z., M. A. Delucchi, A. R. Ingrassia, R. W. Howarth, G. Bazouin, B. Bridgeland, K. Burkart,  
 182 M. Chang, N. Chowdhury, R. Cook, G. Escher, M. Galka, L. Han, C. Heavey, A. Hernandez, D. F.  
 183 Jacobson, D. S. Jacobson, B. Miranda, G. Novotny, M. Pellat, P. Quach, A. Romano, D. Stewart, L.  
 184 Vogel, S. Wang, H. Wang, L. Willman and T. Yeskoo (2014). "A roadmap for repowering California for  
 185 all purposes with wind, water, and sunlight." *Energy* **73**: 875-889.  
 186 Jacobson, M. Z., Y. J. Kaufman and Y. Rudich (2007). "Examining feedbacks of aerosols to urban  
 187 climate with a model that treats 3-D clouds with aerosol inclusions." *Journal of Geophysical Research*  
 188 **112**(D24).  
 189 Johnston, J., A. Mileva, J. H. Nelson and D. M. Kammen (2013). SWITCH-WECC. Data, Assumptions,  
 190 and Model Formulation. Berkeley, California, Renewable and Appropriate Energy Laboratory.  
 191 Kleeman, M. J. and G. R. Cass (2001). "A 3D Eulerian Source-Oriented Model for an Externally Mixed  
 192 Aerosol." *Environmental Science & Technology* **35**(24): 4834-4848.  
 193 Kleeman, M. J., G. R. Cass and A. Eldering (1997). "Modeling the airborne particle complex as a source-  
 194 oriented external mixture." *Journal of Geophysical Research: Atmospheres* **102**(D17): 21355-21372.

195 Krol, M., S. Houweling, B. Bregman, M. van den Broek, A. Segers, P. van Velthoven, W. Peters, F.  
196 Dentener and P. Bergamaschi (2005). "The two-way nested global chemistry-transport zoom model TM5:  
197 algorithm and applications." *Atmos. Chem. Phys.* **5**(2): 417-432.

198 Lamarque, J. F., L. K. Emmons, P. G. Hess, D. E. Kinnison, S. Tilmes, F. Vitt, C. L. Heald, E. A.  
199 Holland, P. H. Lauritzen, J. Neu, J. J. Orlando, P. J. Rasch and G. K. Tyndall (2012). "CAM-chem:  
200 description and evaluation of interactive atmospheric chemistry in the Community Earth System Model."  
201 *Geoscientific Model Development* **5**(2): 369-411.

202 Meinshausen, M., S. C. B. Raper and T. M. L. Wigley (2011). "Emulating coupled atmosphere-ocean and  
203 carbon cycle models with a simpler model, MAGICC6 – Part 1: Model description and calibration."  
204 *Atmospheric Chemistry and Physics* **11**(4): 1417-1456.

205 Nelson, D. J., A. Mileva, J. Johnston and P. D. Kammen (2013). Scenarios for Deep Carbon Emission  
206 Reductions from Electricity by 2050 in Western North America Using the SWITCH Power Electric  
207 Power Sector Planning Model. California's Carbon Challenge Phase II. California Energy Commission.  
208 310 Barrows Hall Berkeley, CA 94720-3050, University of California, Berkeley, Renewable and  
209 Appropriate Energy Laboratory Energy and Resources Group. **II**: 142.

210 Paltsev, S., J. M. Reilly, H. D. Jacoby, R. S. Eckaus, J. McFarland, M. Sarofim, M. Asadoorian and M.  
211 Babiker (2005). The MIT Emissions Prediction and Policy Analysis (EPPA) Model: Version 4. *MIT Joint*  
212 *Program on the Science and Policy of Global Change*. Joint Program on the Science and Policy of Global  
213 Change, 77 Massachusetts Avenue, MIT E40-428, Cambridge MA 02139-4307 (USA), Massachusetts  
214 Institute of Technology (MIT).

215 Parker, N. (2012). "Spatially Explicit Projection of Biofuel Supply for Meeting Renewable Fuel  
216 Standard." *Transportation Research Record: Journal of the Transportation Research Board* **2287**: 72-79.

217 Pozzoli, L., I. Bey, S. Rast, M. G. Schultz, P. Stier and J. Feichter (2008). "Trace gas and aerosol  
218 interactions in the fully coupled model of aerosol-chemistry-climate ECHAM5-HAMMOZ: 1. Model  
219 description and insights from the spring 2001 TRACE-P experiment." *Journal of Geophysical Research*  
220 **113**(D7).

221 Russ, P., T. Wiesenath, D. van Regemorter and J. Ciscar (2007). Global Climate Policy Scenarios for  
222 2030 and beyond. Analysis of Greenhouse Gas Emission Reduction Pathway Scenarios with the POLES  
223 and GEM-E3 models., European Commission. Joint Research Centre. Institute for Prospective  
224 Technological Studies.

225 Shindell, D., J. C. I. Kuylenstierna, E. Vignati, R. van Dingenen, M. Amann, Z. Klimont, S. C. Anenberg,  
226 N. Muller, G. Janssens-Maenhout, F. Raes, J. Schwartz, G. Faluvegi, L. Pozzoli, K. Kupiainen, L.  
227 Höglund-Isaksson, L. Emberson, D. Streets, V. Ramanathan, K. Hicks, N. T. K. Oanh, G. Milly, M.  
228 Williams, V. Demkine and D. Fowler (2012). "Simultaneously Mitigating Near-Term Climate Change  
229 and Improving Human Health and Food Security." *Science* **335**(6065): 183.

230 Shindell, D. T., G. Faluvegi, N. Unger, E. Aguilar, G. A. Schmidt, D. M. Koch, S. E. Bauer and R. L.  
231 Miller (2006). "Simulations of preindustrial, present-day, and 2100 conditions in the NASA GISS  
232 composition and climate model G-PUCCINI." *Atmos. Chem. Phys.* **6**(12): 4427-4459.

233 Smith, S. J., J. J. West and P. Kyle (2011). "Economically consistent long-term scenarios for air pollutant  
234 emissions." *Climatic Change* **108**(3): 619.

235 Thomson, A. M., K. V. Calvin, S. J. Smith, G. P. Kyle, A. Volke, P. Patel, S. Delgado-Arias, B. Bond-  
236 Lamberty, M. A. Wise, L. E. Clarke and J. A. Edmonds (2011). "RCP4.5: a pathway for stabilization of  
237 radiative forcing by 2100." *Climatic Change* **109**(1-2): 77-94.

238 Tittmann, P. W., N. C. Parker, Q. J. Hart and B. M. Jenkins (2010). "A spatially explicit techno-economic  
239 model of bioenergy and biofuels production in California." *Journal of Transport Geography* **18**(6): 715-  
240 728.

241 van Aardenne, J., F. Dentener, R. Van Dingenen, G. Maenhout, E. Marmer, E. Vignati, P. Russ, L. Szabo  
242 and F. Raes (2010). Climate and air quality impacts of combined climate change and air pollution policy  
243 scenarios. *JRC Scientific and Technical Reports*. Luxembourg: Publications Office of the European  
244 Union, European Commission. Joint Research Centre. Institute for Environment and Sustainability.

245 West, J. J., S. J. Smith, R. A. Silva, V. Naik, Y. Zhang, Z. Adelman, M. M. Fry, S. Anenberg, L. W.  
246 Horowitz and J. F. Lamarque (2013). "Co-benefits of Global Greenhouse Gas Mitigation for Future Air  
247 Quality and Human Health." Nat Clim Chang **3**(10): 885-889.  
248 Yang, C., S. Yeh, K. Ramea, S. Zakerinia, D. McCollum, D. Bunch and J. Ogden (2014). Modeling  
249 Optimal Transition Pathways to a Low Carbon Economy in California: California TIMES (CA-TIMES)  
250 Model. Davis, CA., Institute of Transportation Studies, University of California, Davis.  
251 Yang, C., S. Yeh, S. Zakerinia, K. Ramea and D. McCollum (2015). "Achieving California's 80%  
252 greenhouse gas reduction target in 2050: Technology, policy and scenario analysis using CA-TIMES  
253 energy economic systems model." Energy Policy **77**: 118-130.  
254 Ying, Q., M. P. Fraser, R. J. Griffin, J. Chen and M. J. Kleeman (2007). "Verification of a source-oriented  
255 externally mixed air quality model during a severe photochemical smog episode." Atmospheric  
256 Environment **41**(7): 1521-1538.  
257