Assessment of wood burning versus fossil fuel contribution to wintertime black carbon and carbon monoxide concentrations in Athens, Greece

Athina-Cerise Kalogridis¹, Stergios Vratolis¹, Eleni Liakakou², Evangelos Gerasopoulos², Nikolaos Mihalopoulos², Konstantinos Eleftheriadis¹

¹Institute of Nuclear & Radiological Sciences & Technology, Energy & Safety, National Centre of Scientific Research "Demokritos", Ag. Paraskevi, 15310, Greece

² Institute for Environmental Research and Sustainable Development, National Observatory of Athens, Metaxa & V. Pavlou, P. Penteli, 15236, Athens, Greece

10 Correspondence to: Athina-Cerise Kalogridis (akalogridi@ipta.demokritos.gr)

Abstract

5

The scope of this study was to estimate the contribution of fossil fuel and wood burning combustion to black carbon (BC) and carbon monoxide (CO) during wintertime, in Athens. For that purpose, in-situ measurements of equivalent Black Carbon (eBC) and CO were simultaneously conducted in a suburban and an urban background monitoring site in Athens during three months

- of winter 2014-2015. For the deconvolution of eBC into eBC emitted from fossil fuel (BC_{ff}) and wood burning (BC_{wb}), a method based on the spectral dependency of the absorption of pure black carbon and brown carbon was used. Thereafter, BC_{wb} and BC_{ff} estimated fractions were used along with measured CO concentrations in a multiple regression analysis, in order to quantify the contribution of each one of the combustion sources to the ambient CO levels. For an independent evaluation of the results, we additionally estimated the wood-burning and fossil fuel contribution to CO, calculated on the basis of their CO/NO_x emission
- 20 ratios. The results indicate that during wintertime BC and CO are mainly emitted by local sources within the Athens Metropolitan Area (AMA). Fossil fuel combustion, mainly from road traffic, is found to be the major contributor to both eBC in PM_{2.5} and CO ambient concentrations in AMA. However, wintertime wood burning makes a significant contribution to the observed eBC (of about 30%) and CO concentrations (on average, 11 % and 16 % of total CO in the suburban and urban background sites respectively). Both, BC and CO from biomass burning (BC_{wb} and CO_{wb}, respectively) present a clear diurnal

25 pattern with highest concentrations during night, supporting the local domestic heating as their main source.

1 Introduction

Air pollution, which is originating largely from combustion processes, is a very important environmental concern in Athens, like in other large urban agglomerations around the world. High population (3.75 million in the metropolitan area) and the confinement of commercial and industrial activities in a relatively small area (approximately 450 km²), has led to severe

- 30 environmental degradation. Over the years high loadings of atmospheric pollutants have been documented (Chaloulakou et al., 2005; Eleftheriadis et al., 1998, 2014; Kalabokas et al., 1999; Theodosi et al., 2011). Combustion processes used for transportation, power generation and other human activities produce a complex mixture of chemical pollutants (Cohen et al., 2004), which at any given location have characteristics depending on the relative contributions of the different sources of pollution and on the effects of the local geo-climatic factors. Black carbon (BC) aerosol and carbon monoxide (CO) are two
- 35 major products of incomplete combustion and are important atmospheric components because of their substantial impact on health (Ostro et al., 2015), including respiratory and cardiovascular effects, as well as on climate (Zanatta et al., 2016). BC refers to the absorbing components of soot and is the second most significant contributor to climate change (Andreae and

Gelencsér, 2006; Bond et al., 2013). On the other hand, CO strongly influences the oxidative capacity of the atmosphere (by reacting with the OH radicals), and thereby alters the lifetime of methane and other greenhouse gases (Seinfeld and Pandis, 2012). The potential adverse health and climate effects associated with exposure to high levels of BC and CO, motivates a thorough characterization of their emission from different sources. In urban environments, fossil fuel combustion is the major

- 5 source of BC and CO, mainly related to motor vehicle exhausts. However, biomass combustion, from forest fires (especially in summer; Diapouli et al., 2014), or from domestic heating (in winter) may also contribute significantly to their ambient levels. Improvement of air quality in Athens after measures adopted during the last decades are described by Kanakidou et al. (2011) and are in line with proposed mitigation strategies (Aleksandropoulou et al., 2012). Vrekoussis et al. (2013) used satellite observations over Athens for NO₂ and SO₂, to show that the economic crisis resulted in acceleration of the reduction of air
- 10 pollutants in Athens. In the recent years, a resurgence in the use of biofuels over more expensive fuels for heating has been observed in Europe (Denier van der Gon et al., 2015; Gonçalves et al., 2011), especially in Greece where the economic crisis has tripled the fossil fuel cost in a few years (Saffari et al., 2013). The technology of domestic wood burning in Athens is known to suffer from low burning efficiency. The extensive use observed in recent years resulted in considerable emissions of incomplete combustion, i.e. CO, hydrocarbons and soot particles during cold months. Paraskevopoulou et al.(2015), reported
- 15 the impact of wood combustion (dominant fuel for domestic heating) on air quality in Athens, as an almost 30 % increase of the contribution of particulate organic matter to the urban aerosol mass, from winter 2012 to winter 2013. At the same time, a long term analysis of EC concentrations in Athens (Paraskevopoulou et al., 2014) revealed a significant increase in wintertime EC in the period 2011-2013. Respectively, a significant increase in winter evening CO level has been reported for the period 2012-2015 by Gratsea et al., (2017) and attributed to the increase in wood burning use. On the other hand, during summertime
- 20 a consistent decrease is encountered in the last decade as a result of the simultaneous reduction in traffic and industrial activity due to the economic crisis in Greece (Diapouli et al., 2017b).

Even though biomass burning from domestic heating has been recognized as a main source of atmospheric pollutants in Southern Europe (Denier van der Gon et al., 2015; Giannoni et al., 2012; Gonçalves et al., 2011; Paglione et al., 2014; Saffari et al., 2013) emission estimates are still scarce and the associated uncertainty remains high. This is because wood consumption

- 25 statistics are difficult to obtain since wood is often non-commercial and emission factors vary greatly with wood type, combustion equipment and flame temperature. As a matter of fact, reported emission factors of PM for different types of residential combustion appliances range between 10-2000 mg per MJ (Mega Joule) of fuel burnt (Kocbach Bølling et al., 2009). In a similar way, CO emission factors from fireplaces and traditional or eco-labelled woodstoves range typically from 30-120 g kg⁻¹ (ratio of the mass of CO emitted related to the mass of the burnt fuel) (AIRUSE, 2015). The spatial variability of the
- 30 carbonaceous aerosol pollutants is also of great interest with respect to the health impacts of their major contributing sources in the urban PM_{10} .

In this context, the scope of the present study is to investigate, based on ambient measurements, the impact of biomass burning versus fossil fuel use on the air pollution observed in Athens during wintertime. For that purpose, three-months of continuous and simultaneous measurements of equivalent Black Carbon (eBC) by aethalometers and CO at a suburban and an urban

- 35 background site of Athens were analyzed and compared. The measured eBC was deconvoluted into two fractions using a model based on the different spectral dependencies of light absorption by pure black carbon (related to fossil fuel) and brown carbon (linked to wood-burning emissions) (Sandradewi et al., 2008). For simplicity we refer to these light absorbing carbon fractions as "BC" with an additional index for specifying the wood burning (BC_{wb}) or fossil fuel (BC_{ff}) origin. Two independent methods based on the relations between CO and co-products of combustion processes were used and compared for the estimation of CO
- 40 originated from traffic and wood combustion.

2.1 Sampling Sites

Simultaneous measurements of eBC and CO were performed at the National Center of Scientific Research (NCSR) Demokritos (DEM) and at the National Observatory of Athens (NOA), from December 6th 2014 until March 10th 2015 (Figure 1). The

- 5 campus of Demokritos is situated at the foot of Mount Hymmettus in Aghia Paraskevi and covers an area of 600 acres in a forest of pine trees. Situated at 7 km NNE from Athens centre (Triantafyllou et al., 2016), the GAW-Demokritos measurement station (37.99 N, 23.82 E, 270 m a.s.l) is considered representative of the suburban areas of the Athens Metropolitan Area (AMA). NOA's station is located at its central premises at Thission, in the center of Athens and on the top of Nymphs Hill (38.0 N, 23.7 E, 107 m a.s.l.). Its central setting, still relatively far (further than 500 m) from main traffic lines, can be
- 10 considered ideal for monitoring the air pollution urban background of Athens.

2.2 Measurements of aerosol light absorption and carbon monoxide

2.2.1 Aerosol light absorption and equivalent black carbon

The aerosol light absorption coefficient, b_{abs} , was retrieved at each station by means of a 7-wavelength (370, 470, 520, 590, 660, 880 and 950 nm) aethalometer (Magee Scientific Corp., Berkeley, CA 94703, USA) with a 5-min temporal resolution. At

- 15 DEM, the new generation AE33 aethalometer model was used, which provides a real-time compensation for multiple scattering in the filter matrix and loading effects using the DualSpot Technology® (Drinovec et al., 2015). The AE33 sampled aerosol through a PM_{2.5} cut-off inlet. At NOA, a Portable Aethalometer® Model AE42 was used and aerosol sampling was performed with no size-cut. As BC mainly contributes to PM₁ (Laborde et al., 2013; Wang et al., 2015), differences in the eBC concentrations due to the different aerodynamic diameters of sampled aerosols are expected to be negligible. Raw absorption coefficients at a given wavelength
- 20 λ ($b_{aeth,\lambda}$) from AE42 dataset were corrected from loading and scattering effects following the procedure introduced by Weingartner et al. (2003).

$$b_{abs,\lambda} = \frac{b_{ATN,\lambda}}{C_0 \times R(ATN)_{\lambda}} \tag{1}$$

$$R(ATN)_{\lambda} = \left(\frac{1}{f_{\lambda}} - 1\right) \times \frac{\ln(ATN_{\lambda}) - \ln 10}{\ln 50 - \ln 10} + 1$$
(2)

- As described in Eq. (1), a C_0 constant was used to correct for multiple scattering by the filter fibers and the scattering of the 25 aerosols embedded in the filter, whereas a R(ATN) function enabled to compensate the loading effect, i.e. the fact that the 25 attenuation increases as light absorbing particles accumulate in the filter (Eq. (2)). The value of 3.5 was used for C₀ as 26 recommended in Zanatta et al., (2016). The compensation parameter f_{λ} is a parameter that mainly depends on the single 27 scattering albedo of aerosol (SSA). Since no simultaneous scattering coefficient measurements were available during this 28 campaign, f_{λ} values given in Drinovec et al.(2015) for an urban site with a single scattering albedo of about 0.75 were used for 30 loading compensation at the urban background site.
- In order to investigate systematic differences between the processed outputs by the two types of instruments, an intercomparison exercise was performed between data from the new generation AE33-aethalometer compensated in real-time, and data from an AE31-Aethalometer (with identical measurement configuration and settings to the AE42) which operates continuously in parallel with the AE33 at DEM station. A two-month period of data was found reasonable for this exercise with
- 35 AE31 data compensated by post-processing using the Weingartner algorithm (N=8784). The results indicated a very good agreement between the absorption measurements from the AE33 and AE31 instruments after compensation, with R²=0.79, a small intercept of -0.15 (Mm⁻¹) and a slope of 0.97.

Eventually, aerosol light absorption coefficients were converted into mass concentration of the equivalent BC (eBC) as defined by Petzold et al. (2013). BC is historically defined from aethalometer measurements at 880 nm. At 880 nm, no significant difference in MAC at 880 nm between eBC originating from traffic or wood-burning emission is expected (Zotter et al., 2017). eBC mass concentration was derived in this study by multiplying the b_{abs} coefficient at 880 nm with a constant value of mass absorption cross-section (MAC) of 4.6 m² g⁻¹(determined from the comparison with simultaneous measurements at DEM of elemental carbon). An extensive description of EC/OC measurements at DEM is available in Diapouli et al. (2014). Assuming

5 an absorption Ångstrom exponent of 1.0, the MAC used here is 6.13 when adjusted to 637 nm. Our MAC value is at the lower limit of the values reported by Zanatta et al., (2016), for nine rural background stations across Europe (7.5-13.3 m² g⁻¹, calculated for 637 nm), and within the range of values reported by Hitzenberger et al., (2006) for an urban background site in Vienna (5.9 -7.5 at 637 nm).

10 2.2.2 Carbon monoxide and nitrogen oxides

Ambient CO mixing ratios were measured at DEM station at a time resolution of 1 Hz using a Cavity Ring-Down Spectroscopy analyzer (Model G2401, Picarro, CA, USA), which provides high resolution and low detection limit CO, CO_2 and CH_4 ambient mixing ratios in line with GAW standards. Air was pulled through a 5-m line at about 0.4 L min⁻¹, and water was removed from the sample using a NafionTM copolymer membrane dryer (http://www.permapure.com/resources/all-about-nafion-and-faq/). CO

15 measurements were obtained with a typical precision (one sigma) of about ± 4 ppbv in a 1-s measurement for concentrations ranging between 75 and 300 ppbv. Additionally, hourly measurements of NO_x were available from the monitoring station of the Greek Ministry of Environment and Energy (www.ypeka.gr) situated 300 m from DEM station.

CO and NO_x were determined at the NOA station with a 1-min, integration time using a Horiba APMA-360 series automatic gas analyser (NDIR technique, scale: 0-20 ppmv, lower detectable limit: 0.05 ppmv) and a Horiba APNA-360 series (chemiluminescence technique, scale: 0-1000 ppbv, lower detectable limit: 0.5 ppbv) respectively.

2.3 Source apportionment of black carbon from fossil fuel and wood burning combustion.

Source apportionment of the ambient BC concentrations was based on the method developed by Sandradewi et al. (2008) and successfully applied in several studies (Favez et al., 2009; Fourtziou et al., 2017; Fuller et al., 2014; Petit et al., 2014; Sciare et al., 2011). This model relies on the first assumption that the total absorption at a wavelength λ , $b_{abs}(\lambda)$, is a combination of absorption due to fossil fuel ($b_{abs}(\lambda)_{ff}$) and wood burning ($b_{abs}(\lambda)_{wb}$) aerosols:

 $b_{abs}(\lambda) = b_{abs}(\lambda)_{ff} + b_{abs}(\lambda)_{wb}$, (3) Secondly, it is based on the difference in the dependency of the absorption coefficient on wavelength assuming that absorption from fossil fuel and biomass burning emissions follow different spectral dependencies. The wavelength dependent absorption of light b_{abs} by aerosols is proportional to $\lambda^{-\alpha}$ where α is the absorption Ångström exponent such that:

$$30 \quad \frac{b_{abs}(\lambda_1)_{ff}}{b_{abs}(\lambda_2)_{ff}} = \left(\frac{\lambda_1}{\lambda_2}\right)^{-\alpha_{ff}},\tag{4}$$

And respectively:

20

25

$$\frac{b_{abs}(\lambda_1)_{wb}}{b_{abs}(\lambda_2)_{wb}} = \left(\frac{\lambda_1}{\lambda_2}\right)^{-\alpha_{wb}},\tag{5}$$

Light absorption measurements at $\lambda_1 = 470 \text{ nm}$ (*UV*), and $\lambda_2 = 950 \text{ nm}$ (*IR*) are used in this approach due to the fact that when compared to BC from fossil fuel combustion (BC_{ff}), wood burning aerosols (BC_{wb}) exhibit greater absorption in the near

- 35 ultraviolet. This enhanced absorption at near UV for wood burning aerosols is due to the presence of absorbing organic molecules, especially polycyclic aromatic hydrocarbons and humic-like substances (Hoffer et al., 2006). Even though different pairs of near-UV and near-IR wavelengths can be used, it is recommended to use the pair 470 nm versus 950 nm. The choice of 470 against 370 is even more critical as explained in Zotter et al (2017) since VOCs or other absorbing non-BC particles can interfere with measurements with the 370 nm channel of aethalometer. By solving equations 2-5, unique values of $b_{abs}(\lambda_{UV})_{wb}$,
- 40 $b_{abs}(\lambda_{IR})_{wb}, b_{abs}(\lambda_{UV})_{ff}$ and $b_{abs}(\lambda_{IR})_{ff}$, can be calculated, thus leading to the determination of BC_{wb} and BC_{ff}:

$$b_{abs}(\lambda_{UV})_{wb} = \frac{1}{1 - \left(\frac{\lambda_{UV}}{\lambda_{IR}}\right)^{-\alpha_{ff}} * \left(\frac{\lambda_{UV}}{\lambda_{IR}}\right)^{\alpha_{wb}}} \times \left(b_{abs}(\lambda_{UV}) - \left(\frac{\lambda_{UV}}{\lambda_{IR}}\right)^{-\alpha_{ff}} * b_{abs}(\lambda_{IR})\right), \tag{6}$$

$$b_{abs}(\lambda_{IR})_{wb} = \left(\frac{\lambda_{UV}}{\lambda_{IR}}\right)^{a_{wb}} * b_{abs}(\lambda_{UV})_{wb} , \qquad (7)$$

$$b_{abs}(\lambda_{UV})_{ff} = b_{abs}(\lambda_{UV}) - b_{abs}(\lambda_{UV})_{wb}, \tag{8}$$

$$b_{abs}(\lambda_{IR})_{ff} = b_{abs}(\lambda_{IR}) - b_{abs}(\lambda_{IR})_{wb} , \qquad (9)$$

5
$$BC_{ff} = \frac{b_{abs}(\lambda_{IR})_{ff}}{b_{abs}(\lambda_{IR})} \times EBC , \qquad (10)$$
$$BC_{wb} = EBC - BC_{ff} , \qquad (11)$$

The application of the model requires the selection of suitable Ångström exponents for fossil fuel ($\alpha_{\rm ff}$) and wood burning ($\alpha_{\rm wb}$), since one of the greatest uncertainties of the model is associated with the a priori assumed α values for both types of emissions. Reported Ångström exponents range between 0.8-1.1 for pure traffic. For wood burning a wider range of values has been

- 10 observed (0.9-3.5), even though α_{wb} equal to 2.0 has long been considered a typical value for wood burning aerosol (Favez et al., 2009; Fuller et al., 2014; Herich et al., 2011; Petit et al., 2014; Sciare et al., 2011). Recently, Zotter et al., (2017) recommended to use $\alpha_{ff}=0.9$ and $\alpha_{wb}=1.68$, obtained by fitting the model outputs (calculated with the absorption coefficients at 470 and 950 nm) against the fossil fraction of EC derived from 14C measurements. At DEM site, a previous study showed that values of a_{wb} below 1.7 were not appropriate for the specific site (Diapouli et al., 2017a). On top of that during fire events,
- 15 Ångström exponent values up to 2 have been observed at DEM. Taking into consideration these results, absorption Ångström exponents of 0.9 and 2.0 for pure traffic (α_{ff}) and wood burning (α_{wb}), respectively, were used in this study. It should also be noted that coal-burning organic aerosol is known to significantly absorb light at near UV wavelengths (Yang et al., 2009) and may thus interfere with $b_{abs}(\lambda_{UV})_{wb}$. Lignite coal is the single most important local energy source in Greece

(Kavouridis, 2008). However, interferences from coal use are expected to be very low, as the lignite-fired power plants are 20 located far away from Athens (>200 km distance).

2.4 Source apportionment of carbon monoxide from fossil fuel and wood burning combustion

25

The partitioning of CO ambient concentrations into different sources has been investigated in a limited number of studies and was mainly based on the variable isotopic composition of CO (Gros et al., 2002; Kato et al., 1999; Saurer et al., 2009). In the absence of isotopic analysis, we use two different models for the source apportionment, based on the correlations between CO and other combustion tracers.

2.4.1 Model 1: the CO-NO_x linear model.

The CO/NO_x ratio has been used in the past as a diagnostic to characterize different type of emission sources (Fujita et al., 1992; Ravindra et al., 2006; Saurer et al., 2009; Wahlina et al., 2001). It can serve as a useful tool for apportioning CO concentrations 30 at monitoring stations part of national networks where only regulated air pollutants are often measured. The CO-NOx linear model, introduced by Saurer et al. (2009), relies on the fact that both CO and NO_x are common products of combustion processes. Assuming that the only significant combustion processes in urban environments are traffic and wood burning for residential heating (in addition to the regional background, and a minor contribution by industrial processes), the concentrations

35 of NO_x and CO can be expressed as:

$$[NO_x] = [NO_x]_{bgd} + [NO_x]_{ff} + [NO_x]_{wb} , \qquad (12)$$

$$[CO] = [CO]_{bgd} + [CO]_{ff} + [CO]_{wb} , \qquad (13)$$

where $[X]_{ff}$, $[X]_{wb}$ represent the concentration of the tracer X resulting from fossil fuel (mainly traffic) and wood-burning, respectively, whereas $[X]_{bgd}$ represent the background concentration of X.

The CO-NO_x linear model is based on the distinct CO/NO_x ratios for the two emission sources, where the wood-burning emission ratio, r_{wb} , is much larger than the one for traffic, r_{ff} . Considering that photochemical processes do not substantially

5 affect the ambient concentrations of CO and NO_x in winter, the ratios of the concentrations can be regarded as approximately the same as their respective emission ratios, $[CO]_{ff}/[NOx]_{ff} \approx r_{ff}$ and $[CO]_{wb}/[NOx]_{wb} \approx r_{wb}$. Based on this assumption, we can consider that r_{ff} and r_{wb} are given, and as a consequence Eq. (12) can be rewritten as:

$$[NO_x] = [NO_x]_{bgd} + \frac{[CO]_{ff}}{r_{ff}} + \frac{[CO]_{wb}}{r_{wb}},$$
(14)

Equations 13 and 14 allow $[CO]_{ff}$ and $[CO]_{wb}$ to be determined. The concentration of CO originating from wood burning emissions can be expressed as:

$$[CO]_{wb} = \frac{r_{wb}}{r_{ff} - r_{wb}} \times \left[[CO]_{bgd} - [CO] + r_{ff} \left([NO_x] - [NO_x]_{bgd} \right) \right],$$
(15)

Defining the emission ratios r_{wb} and r_{ff} is a crucial step for the source apportionment of CO. The methodology used for their selection is presented in section 3.3.2

It is important here to mention the limitations of this model for CO apportionment. Firstly, it requires an *a priori* knowledge of the emission ratios r_{ff} and r_{wb} . Secondly, it is based on the hypothesis that the CO/NO_x ratio remains constant, while in fact it could be affected by photochemistry. CO is a long-lived species with an atmospheric lifetime of several days to several weeks; hence photochemical processes influence CO concentrations on a limited extent. In contrast NO_x, are much more reactive species. Consequently, any change in reactive nitrogen compounds, mainly by photochemistry, would alter the CO/NO_x ratio.

2.4.2 Model 2: CO-BCwb-BCff multiple linear regression model

- 20 The second model for CO source apportionment is based on the existing relation between the concentrations of CO and BC, introducing advantages in order to overcome the limitations of the previously presented CO-NO_x linear model. In a similar manner to model 1, considering both CO and BC exclusively produced by combustion processes and that in the urban environment the CO/BC ratios can be regarded as equivalent to their source emission ratio, the CO concentration can be
- 25 $[CO] = [CO]_{bdg} + r'_{ff} \times [BC]_{ff} + r'_{wb} \times [BC]_{wb}$, (16) where $r'_{ff} = [CO]_{ff} / [BC]_{ff}$, and $r'_{wb} = [CO]_{wb} / [BC]_{wb}$ the relevant emission ratios at the source. The difference of our approach in this second model resides in the way that this equation is solved. Unlike the CO-NO_x linear model, here *a priori*

30 Eq. (16).

expressed as:

10

Using r'_{ff} and r'_{wb} resulting from the model, the concentration of CO attributed to fossil fuel and wood burning sources can be estimated such that:

knowledge of r_{wb} and r_{ff} emission ratios is not required. Instead, BC_{ff} and BC_{wb} are known variables (determined previously using the method presented in section 2.2), and r'_{ff} and r'_{wb} can be calculated by a multiple linear regression model applied on

$$CO_{ff} = r'_{ff} \times [BC]_{ff} , \qquad (17)$$

and,

40

35 $CO_{wb} = r'_{wb} \times [BC]_{wb}$

Moreover, the hypothesis of negligible photochemical chemistry is met for the time scale considered in this study and for the long-lived species BC and CO, and therefore should not have a significant impact on their ambient ratio.

(18)

While multiple linear regressions are known techniques for source apportionment, they have not yet been applied to investigate wood burning contribution to CO using the aethalometer's model results. The considerable increase in measurements carried out using aethalometers makes this technique an interesting and very useful methodology for apportioning CO concentrations.

3. Results and Discussion

3.1 Levels and diurnal variations of black carbon and carbon monoxide

Statistical summary of eBC and CO levels at NOA and DEM stations, as well as their respective time series are displayed in Table 1 and Fig. 2, respectively. Median eBC and CO levels were, respectively, 2.3 and 1.7 times higher at NOA station

- 5 compared to DEM station. In particular, on days with stagnant atmospheric conditions (low wind speed), concentrations of both combustion tracers were up to 10 times higher at NOA compared to DEM. During days with more turbulence, the levels of eBC and CO were similar at both stations, as a result of intensive mixing and uniform horizontal pollutants' dispersion in the Athens valley. As shown in Fig. 3, eBC mass concentrations observed at NOA station are similar to previously reported values in various urban background sites of European highly populated cities, whereas eBC concentrations at DEM are of the same order of magnitude than in residential urban or suburban areas in Europe.
- Diurnal cycles of eBC and CO, as well as wind speed and temperature have been calculated as 1-h mean values and are shown in Fig. 4. eBC and CO exhibit similar diurnal variability. At both stations, maximum concentrations of eBC and CO occur during morning hours (between 08:00-09:00 a.m) and late evening (between 08:00-09:00 p.m), suggesting common emission sources. Based on traffic volume data (Grivas et al., 2012), a first peak in the emissions from transportation is expected around
- 15 08:00 when people commute to work, followed by a plateau from 08:00-18:00, and a secondary peak until 21:00, after when traffic is decreasing. Wood burning emissions from residential heating are expected to increase during the evening, when temperatures drop and people are back-home. As a result, the first peak in eBC and CO concentrations occurs during morning traffic rush hours, while the surface boundary layer is still shallow. At NOA peaks are quite more pronounced compared to DEM, suggesting that each site is under the influence of different small scale dynamics in the Athens valley (Tombrou et al.,
- 20 2007). It is also interesting to observe that the average minimum of concentrations at NOA occurs during midday due to a higher boundary layer height (BLH) and corresponding aerosol dilution during daytime, when both sampling sites are under the same well-mixed atmosphere, winds are stronger, and consequently the pollutants are more homogeneously dispersed in the metropolitan area. The second peak at NOA during night-time is the result of the Nocturnal Boundary Layer (NBL) formation, with the site (elevation 107 m) well within the nocturnal boundary layer (Kassomenos and Koletsis, 2005), leading to an
- 25 accumulation of atmospheric pollutants from combustion sources active at night. During the late hours of the night, the minimum of the 24-h concentrations are observed at the periphery of the basin (DEM), where during stagnant conditions advection from the urban pollution sources is reduced. Occasional downslope winds (katabatic winds) from Hymettus Mountain (Amanatidis et al., 1992) may enhance air exchange from outside the nocturnal boundary layer (NBL), at the same time induce a build up, and increase in concentrations at the Athens Basin.

30 **3.2 Source apportionment of BC and diurnal variability**

35

Following the deconvolution of BC (see section 2.3), on average BC_{wb} represents 33 % and 29 % of total eBC in PM, at DEM and NOA, respectively. BC_{ff} and BC_{wb} fractions comprise the background concentration of BC (BC_{bgd}). Nevertheless, it was estimated that BC_{bgd} , defined as the 1.25 percentile of the dataset (Kondo et al., 2006), is below 10% of the arithmetic mean concentration for both stations (Table 1). As a matter of fact, regional BC background concentrations are expected to be low compared to ambient levels in urban and suburban environments due to the low emission intensity of widespread sources.

- Wood burning contribution to total eBC is similar as in other European cities. Indeed, wintertime wood burning contribution of about 23-25 % has been reported for urban and suburban areas in Paris (Favez et al., 2009; Petit et al., 2014; Sciare et al., 2011), (24±11) % in Zurich downtown (Herich et al., 2011), and 23 % in London (Fuller et al., 2014).
- Figure 5 and Fig. 6 present the diurnal cycle of BC_{wb} , BC_{ff} , BC_{total} (=eBC) as well as the relative contribution of wood burning 40 aerosols to the total BC (*WB%*). BC_{wb}, as well as *WB%* show a clear diurnal trend, with values from 20-25 % early in the morning to peaks at 40 % during night-time, suggesting a large contribution of wood burning domestic sources spread over the

region of Athens, in addition to the enhancement of concentrations at ground level because of the suppression of the boundary layer height. Other sources like industry and power generation are considered negligible as at European scales both consume less than 1 % of the total amounts of wood used annually (Denier van der Gon et al., 2015; IEA, 2008). Fossil fuel source is nevertheless the main contributor to black carbon concentrations in both areas. In particular, during morning rush hours, it represents up to 70 % and 90 % of total eBC at DEM and NOA, respectively.

5

10

3.3 Source identification of CO

3.3.1 Using BCwb and BCff as tracers of fossil fuel and wood burning sources

The association between CO concentration, BC_{wb} and BC_{ff} were examined using multiple linear regressions. Regression analysis between CO, BC_{wb} and BC_{ff} are shown for both sites in Fig. 7. Regressions were carried out using 10-minutes averaged data which represented a sample size of 13259 and 7474 values for DEM and NOA, respectively. The best-fitted linear equation to observed data, and the partial regression coefficients of the model r'_0 , r'_{ff} and r'_{wb} were calculated so that:

$$[CO] = r'_0 + r'_{ff} \times [BC]_{ff} + r'_{wb} \times [BC]_{wb}, \qquad (18)$$

The model was run with no constraint for DEM, and the determined regression coefficients (r'_{ff} and r'_{wb} for DEM) were found with a relative standard deviation below 2 %. However, for NOA, a constraint was applied in order to achieve a solution mathematically and physically meaningful. This choice was made because of the simultaneous advection of aerosols, resulting in a significant correlation between BC_{wb} and B_{ff} (Fig. 7), thus making more difficult the separation of different sources based on their variability. Since the variability of the emission ratios is greater for wood burning emissions (the emission ratio strongly

depends on type of biofuel and appliances used), the choice was made to constrain the emission ratio of fossil fuel r'_{ff} . The r'_{ff} value for NOA was set identical to the one predicted by the model for DEM (i.e equal to 0.184 ppbv ng⁻¹ m³, see Eq. 19). This

- 20 is an approximation based on the assumption that BC and CO are chemically inert and their emission ratio r'_{ff} cannot differ significantly within the same urban area due to an uniform vehicle fleet mix (size distribution and age of vehicle fleet, environmental performance, driving behaviour etc.). We acknowledge that this assumption might introduce some level of uncertainty. The latter is difficult to be estimated with accuracy. A sensitivity analysis for the NOA emission ratios was made, based on the statistical error of determined r'_{ff} at DEM. r'_{wb} for NOA was re-calculated using not only a single constraint value,
- but a range of values from the lowest (0.184-0.00137) to the highest (0.184+0.00137) around the determined r'_{ff} value from the multilinear regression analysis at DEM. An uncertainty of 25% was finally estimated from this exercise for the calculated r'wb value at NOA. The results of the multiple regression analysis applied at the two sites are presented in Fig. 8.

$$CO_{ff-DEM}(ppbv) = (0.184 \pm 0.00137) \ x \ BC_{ff-DEM} \left(ng \ m^{-3} \right) \rightarrow (BC/CO)_{ff-DEM} = \ 5.4 \ ng \ m^{-3}/ppbv , \tag{19}$$

$$CO_{wb-DEM}(ppb) = (0.114 \pm 0.00216) \ x \ BC_{wb-DEM}(ng \ m^{-3}) \to (BC/CO)_{wb-DEM} = 8.8 \ ng \ m^{-3}/ppbv, \tag{20}$$

30
$$CO_{wb-NOA}(ppb) = (0.131 \pm 0.03275) \times BC_{wb-NOA}(ng m^{-3}) \rightarrow (BC/CO)_{wb-NOA} = 7.6 ng m^{-3}/ppbv,$$
 (21)

The resulting regression coefficients were applied to estimate the fraction of CO attributed to fossil fuel and biomass burning combustion sources whereas the intercept values (108.5 ± 0.64 and 146.8 ± 2.5 ppbv and for DEM and NOA, respectively) were regarded as the background concentrations of CO. The resulting background concentrations are in very good agreement with

- 35 those calculated as 1.25 percentile of the dataset (see Table 1). It is noteworthy to mention here that CO background levels are very significant with regard to the ambient concentrations, representing about 26% and 46% of the arithmetic mean concentration at NOA and DEM respectively. As a matter of fact, widespread natural sources of CO, such as plants, oceans and oxidation of hydrocarbons, in combination with its long atmospheric lifetime are known to maintain a significant background concentration even outside urban areas. These results show that the BC/CO ratio is higher for emission related to biomass
- 40 burning compared to fossil fuel combustion, which is consistent with literature values (Pan et al., 2012). The determined values for DEM and NOA for wood burning are very similar, with a (BC/CO)_{wb} ratio of 7.6-8.8 ng m⁻³ ppbv⁻¹. These values are also

in the low range of emission ratios found in the literature for both transport and domestic heating (using biofuel) sources (see Table 2).

The time series of the deconvolution of CO into three fractions, namely CO_{ff} , CO_{wb} and $CO_{background}$, are shown in Fig. 9. According to our results, the wood-burning fraction of CO, represents on average 11 % and 16 % of total CO for DEM and

5 NOA, respectively. In terms of concentrations, CO_{wb} ranges between 5-52 ppbv (≈ 25 ppbv on average) at DEM and between 2-406 ppbv (≈ 135 ppbv on average) at NOA (Table 3). During night-time (20:00-02:00), this contribution is estimated at 15% and 25% respectively.

Diurnal variabilities of CO_{ff} and CO_{wb} at each site are presented in Fig. 10 (*NB*: different vertical scales are used for each station). As expected, CO_{ff} presents similar variability with that of BC_{ff} , i.e a pronounced bimodal distribution, with higher

10 concentrations during rush hours. Along the same line, the diurnal variability of CO_{wb} shows a unimodal pattern with increasing concentrations after 18:00, due to the combination of enhanced wood burning emission which is a source more active during evening hours and lower ambient temperature and BLH, as discussed previously. Comparison with other European cities is limited by the very few number of studies investigating the carbon monoxide

concentrations sources. Saurer et al., (2009) used the stable isotope composition of CO (δ^{13} C and δ^{18} O) for the characterization of different CO sources at 3 sites in Switzerland during winter (along with other indicators for traffic and wood combustion

such as NO_x -concentration and aerosol light absorption at different wavelengths) and estimated the wood burning contribution to night-time CO concentrations at 70%, 49% and 29% for a village site dominated by domestic heating, a site close to a motorway and a rural site respectively. These differences reflect the spatial variability in the wood burning use within the same region depending on the type of site, as well as between countries depending on the country and heating practices.

20 3.3.2 Comparison the CO-BC_{wb}-BC_{ff} linear model vs the CO-NO_x linear model

The results of the multiple-regression model were compared with those from the CO-NO_x linear model. As presented in section 2, an *a priori* knowledge of the emission ratio of CO/NO_x from traffic (r_{ff}) and wood burning (r_{wb}) emissions is required for the CO-NO_x model and therefore the choice of their value is of major importance.

- In Fig. 11, the scatter plot of CO versus NO_x is shown for the data recorded at DEM and NOA. The data set is mostly comprised between two well defined slopes and the range of values of CO/NO_x observed in actual air samples is 7-25 ppbv ppbv⁻¹ (after subtraction of the CO_{bgd} concentration) for DEM and NOA stations and depends on the contribution of each of the sources. As applied in other studies for similar purposes (Rodríguez and Cuevas, 2007; Saurer et al., 2009), these slopes were estimated using two best fit lines, the first to the points aligned in the lower edge of CO versus NO_x scatter charts, and the second one to the points aligned in the upper edge respectively. More precisely, they have been calculated by fitting only data below the 10th
- 30 percentile and above the 90th percentile respectively of CO/NO_x data. Knowing that lowest ratios are obtained when traffic emissions dominate (and when the contribution of woodburning is insignificant), an r_{ff} value of 7 ppbv ppbv⁻¹ was estimated. Higher ratios were obtained typically late in the night-time, when traffic emission decreased and domestic heating increased. However, as we do not expect a contribution of domestic heating close to 100 % at any time of the day in Athens, it is impossible to estimate with accuracy the r_{wb} ratio based solely on this data. Nevertheless, based on "wood burning" lines from Fig. 11, and
- 35 assuming that emission ratios from wood burning are similar between NOA and DEM, we estimate a r_{wb} ratio for the area of Athens, larger than 25 ppbv ppbv⁻¹.

A sensitivity analysis of the CO-NO_x model was performed using the experimentally determined r_{ff} (at 7 ppbv ppbv⁻¹) and by varying the r_{wb} . Measurements performed directly at the emission source and close to the chimney exhausts during controlled wood-burning experiments, indicated ratios in the range of 50-150 ppbv ppbv⁻¹(Albinet et al., 2015; Nalin, 2014). As a result,

40 the choice was made to vary the r_{wb} parameter from 50 to 150 ppbv ppbv⁻¹. The results of the analysis are presented in Table 4. The influence of increasing r_{wb} at a constant r_{ff} resulted in a relatively minor reduction in the calculated contribution from woodburning of 2-3 %. A good correlation was found between both models, with coefficients of determination R² of 0.52 and 0.85 at DEM and NOA, respectively. The wood burning contribution to CO estimated using the $CO-NO_x$ linear model is higher compared to the estimations from $CO-BC_{ff}-BC_{wb}$ model, of a factor of about 1.5 for NOA, which can be considered as "acceptable" given the uncertainties associated with both models. A higher overestimation compared to model 2 of a factor of about 2.4 is found at DEM suburban station, which could be explained by the fact that the site is characterized by more aged

5

air masses compared to the urban background NOA station. Consequently, ambient CO/NO_x ratios might differ more significantly from emission ratios at DEM suburban station. These results suggest that $CO-NO_x$ linear model probably overestimate wood burning contribution to CO, especially in environments characterized by aged air masses were photochemical loss of NOx cannot be considered as negligible.

4. Conclusion

- 10 In this study we performed a comprehensive field campaign at two surface stations in a suburban and a central area of Athens during winter 2014-2015 in order to investigate the impact of fossil fuel and biomass combustion on the urban air quality. We report measurements of particulate black carbon and CO and NO_x gaseous components performed simultaneously at the monitoring station of Demokritos (DEM), representative of suburban areas of the Athens Metropolitan Area and at the National Observatory of Athens (NOA), typical of urban background conditions. More precisely, black carbon particles were
- 15 concurrently measured using two 7-wavelength aethalometers, whereas mixing ratios of CO were measured with an infrared absorption analyzer at NOA and by wavelength-scanned cavity ring down spectrometry at DEM. The median BC concentrations were 528 ng m⁻³ at DEM and 1198 ng m⁻³ at NOA. In a similar way, median CO mixing ratios were 195 ppbv at DEM and 324 ppbv at NOA. These differences have been explained by the location of the two sites with respect to the proximity from sources and local atmospheric dynamics in the Athens valley. Both BC and CO displayed a clear
- 20 bimodal diurnal pattern, in which morning peaks were observed due to morning inversion and rush-hour traffic, while evening peaks were attributed to combustion sources (evening traffic rush-hour, residential heating) combined with the effects of a shallow nocturnal boundary layer. Highest concentrations were observed during low wind speeds, suggesting that both combustion products were not related to regional transport but instead originated from sources within Athens.
- Source apportionment of BC was carried out using a model based on the absorbance spectral differences of black carbon (related to fossil fuel) and brown carbon (related to biomass burning). Our results suggest that even though fossil fuel combustion is the major contributor to BC in PM₁₀, woodburning makes an important contribution of about 30% to wintertime BC concentrations at both sites (on average 33 % at DEM and 29 % at NOA, respectively, but this difference lies within the uncertainty range of the calculations). BC from biomass burning displayed a clear unimodal diurnal pattern with the highest concentrations during night, confirming that its main source was local domestic heating.
- 30 As both datasets showed significant BC and CO correlations, we used observations of CO mixing ratios along with the fraction of BC_{wb} and BC_{ff} to quantify the percentage of observed CO which originates from fossil fuel and wood burning sources. This analysis led to the conclusion that the wood-burning fraction of CO from local emissions, represents on average 11-16 % of total CO in Athens during wintertime. The method proposed here for the source apportionment of CO was compared to a previously reported method based on the CO-NO_x ratios. From our results, it appears that the CO-NO_x linear model over-
- 35 estimates the contribution of wood burning to CO concentrations, especially in environments characterized by aged air masses, likely due to the fact that the hypothesis of negligible photochemical loss of NO_x is not always met.

Acknowledgements

Financial support from the EnTeC FP7 Capacities programme (REGPOT-2012-2013-1, FP7, ID: 316173), is kindly acknowledged. Thanks are due to prof. Jean Sciare for the loan of AE42 aethalometer, we acknowledge financial support by

the KRIPIS/NSRF2007-2013 project and thank NOA team (Drs V. Psilloglou and M. Lianou) for the operation and maintenance of NOA's site at Thission. This study contributes to ChArMEx work package 1 on Emissions and Sources.

References

AIRUSE: Emission factors for biomass burning. Action B4. AIRUSE LIFE 11 ENV/ES/584., 2015.

Albinet, A., Favez, O., Collet, S., Petit, J.-E., Nalin, F., Aujay, R., Verlhac, S., Dermingny, A., Fievet, A., Karsoski, N., Dubois, P., Golly, B., Besombes, J.-L., Pelletier, C., Sarda-Esteve, R., Gros, V., Bonnaire, N. and Sciare, J.: Emission dynamic from residential wood combustion: PM, BC, OM, VOCs and levoglucosan, in Handbook EAC, Milan., 2015.

5

25

35

Aleksandropoulou, V., Eleftheriadis, K., Diapouli, E., Torseth, K. and Lazaridis, M.: Assessing PM10 source reduction in urban agglomerations for air quality compliance, J. Environ. Monit., 14(1), 266–278, doi:10.1039/C1EM10673B, 2012.

Alves, C., Carvalho, A. and Pio, C.: Mass balance of organic carbon fractions in atmospheric aerosols, J. Geophys. Res. Atmospheres, 107(D21), 8345, doi:10.1029/2001JD000616, 2002.

10 Amanatidis, G. T., Papadopoulos, K. H., Bartzis, J. G. and Helmis, C. G.: Evidence of katabatic flows deduced from a 84 m meteorological tower in Athens, Greece, Bound.-Layer Meteorol., 58(1–2), 117–132, doi:10.1007/BF00120754, 1992.

Andreae, M. O. and Gelencsér, A.: Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols, Atmos Chem Phys, 6(10), 3131–3148, doi:10.5194/acp-6-3131-2006, 2006.

- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B.,
 Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang,
 S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P.,
 Shindell, D., Storelvmo, T., Warren, S. G. and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, J. Geophys. Res. Atmospheres, 118(11), 5380–5552, doi:10.1002/jgrd.50171, 2013.
- Denier van der Gon, H. A. C., Bergström, R., Fountoukis, C., Johansson, C., Pandis, S. N., Simpson, D. and Visschedijk, A. J.
 H.: Particulate emissions from residential wood combustion in Europe revised estimates and an evaluation, Atmos Chem Phys, 15(11), 6503–6519, doi:10.5194/acp-15-6503-2015, 2015.

Diapouli, E., Popovicheva, O., Kistler, M., Vratolis, S., Persiantseva, N., Timofeev, M., Kasper-Giebl, A. and Eleftheriadis, K.: Physicochemical characterization of aged biomass burning aerosol after long-range transport to Greece from large scale wildfires in Russia and surrounding regions, Summer 2010, Atmos. Environ., 96(Supplement C), 393–404, doi:10.1016/j.atmosenv.2014.07.055, 2014.

Diapouli, E., Kalogridis, A.-C., Markantonaki, C., Vratolis, S., Fetfatzis, P., Colombi, C. and Eleftheriadis, K.: Annual Variability of Black Carbon Concentrations Originating from Biomass and Fossil Fuel Combustion for the Suburban Aerosol in Athens, Greece, Atmosphere, 8(12), 234, doi:10.3390/atmos8120234, 2017a.

Diapouli, E., Manousakas, M., Vratolis, S., Vasilatou, V., Maggos, T., Saraga, D., Grigoratos, T., Argyropoulos, G., Voutsa,
D., Samara, C. and Eleftheriadis, K.: Evolution of air pollution source contributions over one decade, derived by PM10 and
PM2.5 source apportionment in two metropolitan urban areas in Greece, Atmos. Environ., 164, 416–430, doi:10.1016/j.atmosenv.2017.06.016, 2017b.

Drinovec, L., Močnik, G., Zotter, P., Prévôt, A. S. H., Ruckstuhl, C., Coz, E., Rupakheti, M., Sciare, J., Müller, T., Wiedensohler, A. and Hansen, A. D. A.: The "dual-spot" Aethalometer: an improved measurement of aerosol black carbon with real-time loading compensation, Atmos Meas Tech, 8(5), 1965–1979, doi:10.5194/amt-8-1965-2015, 2015.

Favez, O., Cachier, H., Sciare, J., Sarda-Estève, R. and Martinon, L.: Evidence for a significant contribution of wood burning aerosols to PM2.5 during the winter season in Paris, France, Atmos. Environ., 43(22–23), 3640–3644, doi:10.1016/j.atmosenv.2009.04.035, 2009.

Fourtziou, L., Liakakou, E., Stavroulas, I., Theodosi, C., Zarmpas, P., Psiloglou, B., Sciare, J., Maggos, T., Bairachtari, K.,
Bougiatioti, A., Gerasopoulos, E., Sarda-Estève, R., Bonnaire, N. and Mihalopoulos, N.: Multi-tracer approach to characterize domestic wood burning in Athens (Greece) during wintertime, Atmos. Environ., 148(Supplement C), 89–101, doi:10.1016/j.atmosenv.2016.10.011, 2017.

Fuller, G. W., Tremper, A. H., Baker, T. D., Yttri, K. E. and Butterfield, D.: Contribution of wood burning to PM10 in London, Atmos. Environ., 87, 87–94, doi:10.1016/j.atmosenv.2013.12.037, 2014.

45 Giannoni, M., Martellini, T., Bubba, M. D., Gambaro, A., Zangrando, R., Chiari, M., Lepri, L. and Cincinelli, A.: The use of levoglucosan for tracing biomass burning in PM2.5 samples in Tuscany (Italy), Environ. Pollut., 167(Supplement C), 7–15, doi:https://doi.org/10.1016/j.envpol.2012.03.016, 2012. Gonçalves, C., Alves, C., Fernandes, A. P., Monteiro, C., Tarelho, L., Evtyugina, M. and Pio, C.: Organic compounds in PM2.5 emitted from fireplace and woodstove combustion of typical Portuguese wood species, Atmos. Environ., 45(27), 4533-4545, doi:10.1016/j.atmosenv.2011.05.071, 2011.

Gratsea, M., Liakakou, E., Mihalopoulos, N., Adamopoulos, A., Tsilibari, E. and Gerasopoulos, E.: The combined effect of 5 reduced fossil fuel consumption and increasing biomass combustion on Athens' air quality, as inferred from long term CO measurements, Sci. Total Environ., 592, 115-123, doi:10.1016/j.scitotenv.2017.03.045, 2017.

Grivas, G., Cheristanidis, S. and Chaloulakou, A.: Elemental and organic carbon in the urban environment of Athens. Seasonal and diurnal variations and estimates of secondary organic carbon, Sci. Total Environ., 414, 535-545, doi:10.1016/j.scitotenv.2011.10.058, 2012.

Gros, V., Jöckel, P., Brenninkmeijer, C. A. M., Röckmann, T., Meinhardt, F. and Graul, R.: Characterization of pollution events 10 observed at Schauinsland, Germany, using CO and its stable isotopes, Atmos. Environ., 36(17), 2831–2840, doi:10.1016/S1352-2310(02)00144-9, 2002.

Herich, H., Hueglin, C. and Buchmann, B.: A 2.5 year's source apportionment study of black carbon from wood burning and fossil fuel combustion at urban and rural sites in Switzerland, Atmospheric Meas. Tech., 4(7), 1409–1420, doi:10.5194/amt-4-1409-2011, 2011.

15

20

30

Hitzenberger, R. and Tohno, S.: Comparison of black carbon (BC) aerosols in two urban areas - concentrations and size distributions, Atmos. Environ., 35(12), 2153–2167, doi:10.1016/S1352-2310(00)00480-5, 2001.

Hitzenberger, R., Petzold, A., Bauer, H., Ctyroky, P., Pouresmaeil, P., Laskus, L. and Puxbaum, H.: Intercomparison of Thermal and Optical Measurement Methods for Elemental Carbon and Black Carbon at an Urban Location, Environ. Sci. Technol., 40(20), 6377-6383, doi:10.1021/es051228v, 2006.

Hoffer, A., Gelencsér, A., Guyon, P., Kiss, G., Schmid, O., Frank, G. P., Artaxo, P. and Andreae, M. O.: Optical properties of humic-like substances (HULIS) in biomass-burning aerosols, Atmos Chem Phys, 6(11), 3563–3570, doi:10.5194/acp-6-3563-2006, 2006.

IEA: International Energy Agency: World Energy Statistics and Balances (2008 Edition), 2008.

25 Invernizzi, G., Ruprecht, A., Mazza, R., De Marco, C., Močnik, G., Sioutas, C. and Westerdahl, D.: Measurement of black carbon concentration as an indicator of air quality benefits of traffic restriction policies within the ecopass zone in Milan, Italy, Atmos. Environ., 45(21), 3522–3527, doi:10.1016/j.atmosenv.2011.04.008, 2011.

Kanakidou, M., Mihalopoulos, N., Kindap, T., Im, U., Vrekoussis, M., Gerasopoulos, E., Dermitzaki, E., Unal, A., Kocak, M., Markakis, K., Melas, D., Kouvarakis, G., Youssef, A. F., Richter, A., Hatzianastassiou, N., Hilboll, A., Ebojie, F., Wittrock, F., von Savigny, C., Burrows, J. P., Ladstaetter-Weissenmayer, A. and Moubasher, H.: Megacities as hot spots of air pollution in the East Mediterranean, Atmos. Environ., 45(6), 1223–1235, doi:10.1016/j.atmosenv.2010.11.048, 2011.

Kassomenos, P. A. and Koletsis, I. G.: Seasonal variation of the temperature inversions over Athens, Greece, Int. J. Climatol., 25(12), 1651-1663, doi:10.1002/joc.1188, 2005.

Kato, S., Akimoto, H., Bräunlich, M., Röckmann, T. and Brenninkmeijer, C. A. M.: Measurements of stable carbon and oxygen isotopic compositions of CO in automobile exhausts and ambient air from semi-urban Mainz, Germany, Geochem. J., 33(2), 35 73-77, doi:10.2343/geochemj.33.73, 1999.

Kavouridis, K.: Lignite industry in Greece within a world context: Mining, energy supply and environment, Energy Policy, 36(4), 1257–1272, 2008.

Kocbach Bølling, A., Pagels, J., Yttri, K. E., Barregard, L., Sallsten, G., Schwarze, P. E. and Boman, C.: Health effects of 40 residential wood smoke particles: the importance of combustion conditions and physicochemical particle properties, Part. Fibre Toxicol., 6, 29, doi:10.1186/1743-8977-6-29, 2009.

Kondo, Y., Komazaki, Y., Miyazaki, Y., Moteki, N., Takegawa, N., Kodama, D., Deguchi, S., Nogami, M., Fukuda, M., Miyakawa, T., Morino, Y., Koike, M., Sakurai, H. and Ehara, K.: Temporal variations of elemental carbon in Tokyo, J. Geophys. Res. Atmospheres, 111(D12), D12205, doi:10.1029/2005JD006257, 2006.

Laborde, M., Crippa, M., Tritscher, T., Jurányi, Z., Decarlo, P. F., Temime-Roussel, B., Marchand, N., Eckhardt, S., Stohl, A., 45 Baltensperger, U., Prévôt, A. S. H., Weingartner, E. and Gysel, M.: Black carbon physical properties and mixing state in the European megacity Paris, Atmos Chem Phys, 13(11), 5831–5856, doi:10.5194/acp-13-5831-2013, 2013.

Nalin, F.: PAHs and their nitrated and oxygenated derivatives (NPAHs and OPAHs) emitted by wood combustion: Original analytical development and study of their evolution from emission point to ambient air, PhD dissertation, Universita degli studi di torino, Universita degli studi di torino, July., 2014.

Paglione, M., Saarikoski, S., Carbone, S., Hillamo, R., Facchini, M. C., Finessi, E., Giulianelli, L., Carbone, C., Fuzzi, S.,
Moretti, F., Tagliavini, E., Swietlicki, E., Eriksson Stenström, K., Prévôt, A. S. H., Massoli, P., Canaragatna, M., Worsnop, D. and Decesari, S.: Primary and secondary biomass burning aerosols determined by proton nuclear magnetic resonance (1H-NMR) spectroscopy during the 2008 EUCAARI campaign in the Po Valley (Italy), Atmos Chem Phys, 14(10), 5089–5110, doi:10.5194/acp-14-5089-2014, 2014.

Pan, X. L., Kanaya, Y., Wang, Z. F., Taketani, F., Tanimoto, H., Irie, H., Takashima, H. and Inomata, S.: Emission ratio of
carbonaceous aerosols observed near crop residual burning sources in a rural area of the Yangtze River Delta Region, China, J.
Geophys. Res. Atmospheres, 117(D22), D22304, doi:10.1029/2012JD018357, 2012.

Paraskevopoulou, D., Liakakou, E., Gerasopoulos, E., Theodosi, C. and Mihalopoulos, N.: Long-term characterization of organic and elemental carbon in the PM2.5 fraction: the case of Athens, Greece, Atmos Chem Phys, 14(23), 13313–13325, doi:10.5194/acp-14-13313-2014, 2014.

15 Paraskevopoulou, D., Liakakou, E., Gerasopoulos, E. and Mihalopoulos, N.: Sources of atmospheric aerosol from long-term measurements (5 years) of chemical composition in Athens, Greece, Sci. Total Environ., 527–528, 165–178, doi:10.1016/j.scitotenv.2015.04.022, 2015.

Petit, J.-E., Favez, O., Sciare, J., Canonaco, F., Croteau, P., Močnik, G., Jayne, J., Worsnop, D. and Leoz-Garziandia, E.: Submicron aerosol source apportionment of wintertime pollution in Paris, France by double positive matrix factorization
(PMF2) using an aerosol chemical speciation monitor (ACSM) and a multi-wavelength Aethalometer, Atmos Chem Phys, 14(24), 13773–13787, doi:10.5194/acp-14-13773-2014, 2014.

Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, S.-M., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A. and Zhang, X.-Y.: Recommendations for reporting "black carbon" measurements, Atmos Chem Phys, 13(16), 8365–8379, doi:10.5194/acp-13-8365-2013, 2013.

25 Reche, C., Querol, X., Alastuey, A., Viana, M., Pey, J., Moreno, T., Rodríguez, S., González, Y., Fernández-Camacho, R., de la Rosa, J., Dall'Osto, M., Prévôt, A. S. H., Hueglin, C., Harrison, R. M. and Quincey, P.: New considerations for PM, Black Carbon and particle number concentration for air quality monitoring across different European cities, Atmos Chem Phys, 11(13), 6207–6227, doi:10.5194/acp-11-6207-2011, 2011.

Rodríguez, S. and Cuevas, E.: The contributions of "minimum primary emissions" and "new particle formation enhancements"
to the particle number concentration in urban air, J. Aerosol Sci., 38(12), 1207–1219, doi:10.1016/j.jaerosci.2007.09.001, 2007.

Saffari, A., Daher, N., Samara, C., Voutsa, D., Kouras, A., Manoli, E., Karagkiozidou, O., Vlachokostas, C., Moussiopoulos, N., Shafer, M. M. and others: Increased biomass burning due to the economic crisis in Greece and its adverse impact on wintertime air quality in Thessaloniki, Environ. Sci. Technol., 47(23), 13313–13320, 2013.

Sandradewi, J., Prévôt, A. S. H., Szidat, S., Perron, N., Alfarra, M. R., Lanz, V. A., Weingartner, E. and Baltensperger, U.:
Using Aerosol Light Absorption Measurements for the Quantitative Determination of Wood Burning and Traffic Emission Contributions to Particulate Matter, Environ. Sci. Technol., 42(9), 3316–3323, doi:10.1021/es702253m, 2008.

Saurer, M., Prévôt, A. S. H., Dommen, J., Sandradewi, J., Baltensperger, U. and Siegwolf, R. T. W.: The influence of traffic and wood combustion on the stable isotopic composition of carbon monoxide, Atmospheric Chem. Phys., 9(9), 3147–3161, doi:10.5194/acp-9-3147-2009, 2009.

40 Sciare, J., d'Argouges, O., Sarda-Estève, R., Gaimoz, C., Dolgorouky, C., Bonnaire, N., Favez, O., Bonsang, B. and Gros, V.: Large contribution of water-insoluble secondary organic aerosols in the region of Paris (France) during wintertime, J. Geophys. Res. Atmospheres, 116(D22), D22203, doi:10.1029/2011JD015756, 2011.

Seinfeld, J. H. and Pandis, S. N.: Atmospheric chemistry and physics: from air pollution to climate change, John Wiley & Sons., 2012.

45 Tombrou, M., Dandou, A., Helmis, C., Akylas, E., Angelopoulos, G., Flocas, H., Assimakopoulos, V. and Soulakellis, N.: Model evaluation of the atmospheric boundary layer and mixed-layer evolution, Bound.-Layer Meteorol., 124(1), 61–79, doi:10.1007/s10546-006-9146-5, 2007. Triantafyllou, E., Diapouli, E., Tsilibari, E. M., Adamopoulos, A. D., Biskos, G. and Eleftheriadis, K.: Assessment of factors influencing PM mass concentration measured by gravimetric & beta attenuation techniques at a suburban site, Atmos. Environ., 131, 409–417, doi:10.1016/j.atmosenv.2016.02.010, 2016.

Verma, R. L., Sahu, L. K., Kondo, Y., Takegawa, N., Han, S., Jung, J. S., Kim, Y. J., Fan, S., Sugimoto, N., Shammaa, M. H.,
Zhang, Y. H. and Zhao, Y.: Temporal variations of black carbon in Guangzhou, China, in summer 2006, Atmos Chem Phys, 10(14), 6471–6485, doi:10.5194/acp-10-6471-2010, 2010.

Vrekoussis, M., Richter, A., Hilboll, A., Burrows, J. P., Gerasopoulos, E., Lelieveld, J., Barrie, L., Zerefos, C. and Mihalopoulos, N.: Economic crisis detected from space: Air quality observations over Athens/Greece, Geophys. Res. Lett., 40(2), 458–463, doi:10.1002/grl.50118, 2013.

10 Weingartner, E., Saathoff, H., Schnaiter, M., Streit, N., Bitnar, B. and Baltensperger, U.: Absorption of light by soot particles: determination of the absorption coefficient by means of aethalometers, J. Aerosol Sci., 34(10), 1445–1463, doi:10.1016/S0021-8502(03)00359-8, 2003.

Yang, M., Howell, S. G., Zhuang, J. and Huebert, B. J.: Attribution of aerosol light absorption to black carbon, brown carbon, and dust in China – interpretations of atmospheric measurements during EAST-AIRE, Atmos Chem Phys, 9(6), 2035–2050, doi:10.5194/acp-9-2035-2009, 2009.

15

20

30

Zanatta, M., Gysel, M., Bukowiecki, N., Müller, T., Weingartner, E., Areskoug, H., Fiebig, M., Yttri, K. E., Mihalopoulos, N., Kouvarakis, G., Beddows, D., Harrison, R. M., Cavalli, F., Putaud, J. P., Spindler, G., Wiedensohler, A., Alastuey, A., Pandolfi, M., Sellegri, K., Swietlicki, E., Jaffrezo, J. L., Baltensperger, U. and Laj, P.: A European aerosol phenomenology-5: Climatology of black carbon optical properties at 9 regional background sites across Europe, Atmos. Environ., 145, 346–364, doi:10.1016/j.atmosenv.2016.09.035, 2016.

Zhang, Q. J., Beekmann, M., Drewnick, F., Freutel, F., Schneider, J., Crippa, M., Prevot, A. S. H., Baltensperger, U., Poulain, L., Wiedensohler, A., Sciare, J., Gros, V., Borbon, A., Colomb, A., Michoud, V., Doussin, J.-F., Denier van der Gon, H. A. C., Haeffelin, M., Dupont, J.-C., Siour, G., Petetin, H., Bessagnet, B., Pandis, S. N., Hodzic, A., Sanchez, O., Honoré, C. and Perrussel, O.: Formation of organic aerosol in the Paris region during the MEGAPOLI summer campaign: evaluation of the volatility-basis-set approach within the CHIMERE model, Atmospheric Chem. Phys., 13(11), 5767–5790, doi:10.5194/acp-13-

volatility-basis-set approach within the CHIMERE model, Atmospheric Chem. Phys., 13(11), 5767–5790, doi:10.5194/acp-13-5767-2013, 2013.

Zotter, P., Herich, H., Gysel, M., El-Haddad, I., Zhang, Y., Močnik, G., Hüglin, C., Baltensperger, U., Szidat, S. and Prévôt, A. S. H.: Evaluation of the absorption Ångström exponents for traffic and wood burning in the Aethalometer-based source apportionment using radiocarbon measurements of ambient aerosol, Atmos Chem Phys, 17(6), 4229–4249, doi:10.5194/acp-17-4229-2017, 2017.

December 2014	and 10th Mar	ch 2015.					
	Arithmetic	Geometric	Median	90 th	10 th	1.25 th	Number of
	$mean \pm stdv$	mean		percentile	percentile	percentile**	datapoints
							(10-min
							averaged)
eBC (DEM)	656±519	497	528	1265	179	71	13259
ng.m ⁻³							
eBC (NOA)	2655±3554	1372	1198	6963	352	57	7474
ng.m ⁻³							
CO (DEM)	214±95	199	195	315	125	101	13259
ppbv							
CO (NOA)	555±570	404	324	1282	196	143	12127
ppbv							
*NOx (DEM)	6.17±3.47		5.56	9.93	3.66	3.13	2238
NOx (NOA)	29.6±42.7	12.4	11.4	84.5	1.9	1.2	12127

Table 1. Statistical summary of eBC and CO concentrations at DEM and NOA stations for the period between 6th December 2014 and 10th March 2015.

^{*}NO_x measurements were available from the monitoring station of the Greek Ministry of Environment and Energy (www.ypeka.gr) situated 300 m from DEM station

5 **Background concentration levels estimated as in Kondo et al., 2006

Table 2. BC/CO (ngC m⁻³ ppbv⁻¹) ratios derived from emission factors found in the literature and from ambient measurements in Athens using the multiple regression model.

	Transport	Domestic heating	Reference		
	(biofuel)				
Derived from Emissio	on factors				
	3.1-11.5 (gasoline)	8.7-26.6	Verma et al., 2010		
	1.3-55 (diesel)				
Derived from ambien	t measurements (multiple regress	sion model)			
DEM station	5.4	8.8	This study		
NOA station	5.4 (fixed)	7.6	This study		

10 Table 3. Statistical summary of calculated CO_{wb} and CO_{ff} concentrations, as well as CO_{wb}(%) using the multiple regression model (model 1) at DEM and NOA stations.

		DEM			NOA	
	CO _{wb} (ppbv)	CO _{ff} (ppbv)	CO _{wb} (%)	CO _{wb} (ppbv)	CO _{ff} (ppbv)	$CO_{wb}(\%)$
mean (±stdv)	25.1 (25.7)	80.2 (65.5)	11 (9)	134.5 (230.6)	297.3 (366.9)	16.3 (14.3)
90 th percentile	51.9	156.7	21	405.9	731.0	37.1
10 th percentile	4.6	19.8	3.0	1.9	50.8	1.0

	r_t	r_{wb}	Mean CO _{wb} /CO	Slope of regression
	(ppbv/ppbv)	(ppbv/ppbv)	(%)	between (CO _{wb}) _{model1}
				and (CO _{wb}) _{model2}
DEM				
station				
Test 1	7	50	29%	2.4 (R ² =0.52)
Test 2	7	100	28%	2.2 (R ² =0.52)
Test 3	7	150	26%	2.1 (R ² =0.52)
NOA				
station				
Test 1	7	50	34%	1.6 (R ² =0.85)
Test 2	7	100	31%	1.51 (R ² =0. 85)
Test 3	7	150	30%	1.48 (R ² =0. 85)

Table 4. Sensitivity test of the CO-NO_x linear model for a constant emission ratio for traffic (r_t) and a variable emission ratio for wood burning and comparison of model 1 (linear) versus model 2 (multilinear).



Figure 1: Regional map along with 3D satellite map of the Athens Metropolitan area (black dashed rectangle), and photos of the NCSR Demokritos (DEM) campus in Aghia Paraskevi (Athens suburban, red star), and of the National Observatory of Athens (NOA) at Thiseion (Athens centre, blue star).

10



Figure 2: Time series of 10-minutes averaged CO and eBC concentrations measured at NOA and DEM monitoring stations from 6th December 2014 until 10th march 2015.



Figure 3: Levels of black carbon (mean±stdv) reported for various European cities, (blue) in traffic-influenced sites, such as London (year 2009, Reche et al., 2011), Milan (Summer 2010, Lisbon (June 2000, Alves et al., 2002), Barcelona(year 2009, Reche et al., 2011), Paris (summer 2009, Zhang et al., 2013) and Helsinki (winter 2000, Hitzenberger and Tohno, 2001); (red) in urban background/residential areas such as Milan (Summer 2010, Invernizzi et al., 2011), London (year 2009, Reche et al., 2011) and Lugano (year 2009, Reche et al., 2011) (green) a suburban area of Paris (J aborde et al., 2013) along with the results from this study at NOA

10 (year 2009, Reche et al., 2011), (green) a suburban area of Paris (Laborde et al., 2013), along with the results from this study at NOA and DEM stations in winter 2014-2015.



Figure 4: (Top) Diurnal trends of BC and CO concentrations at NOA and DEM monitoring stations (Local Time = UTC). Vertical bars show standard deviation to the mean value. (Bottom) Diurnal trends of wind speed and temperature measured at the meteorological station of Demokritos. Box plots indicate the 10th, 25th, 50th and 90th percentile, whereas markers represent the mean values.



Figure 5: Diurnal cycle of BCwb, BCff, BCtotal and wood burning contribution to total BC (WB, defined as BCwb/BCtotal*100) at DEM station.

5



Figure 6: Diurnal cycle of BC_{wb}, BC_{ff}, BC_{total} and wood burning contribution to total BC (WB, defined as BC_{wb}/ BC_{total}*100) at NOA station.

19



Figure 7: Correlation plots between (left) CO and BC_{ff}, (middle) CO and BC_{wb}, (right) BC_{wb} and BC_{ff} for DEM (a) and NOA (b) stations.



Figure 8: Best-fit linear correlations between CO and a combination of BCff and BCwb for DEM (a) and NOA station (b).



Figure 9: Time series of the calculated CO_{ff}, CO_{wb} and CO_{bgd} concentration at (a) DEM and (b) NOA stations.



5 Figure 10: Diurnal variations of CO_{ff} and CO_{wb} at DEM and NOA stations.



Figure 11: Scatter plot for CO and NO_x data from NOA (blue) and DEM (red) along with theoretical lines based on pure traffic and pure wood combustion CO/NO_x emission ratios.