

## **Response to referees' comments**

Journal: ACP

Title: Seasonal variation and light absorption property of carbonaceous aerosol in a typical glacier region of the southeastern Tibetan Plateau

5 Author(s): Hewen Niu et al.

MS No.: acp-2017-865

MS Type: Research article

Iteration: Major Revision

Special Issue: Study of ozone, aerosols and radiation over the Tibetan Plateau

10 (SOAR-TP) (ACP/AMT inter-journal SI)

We thank the referee and editor for their careful review, insightful comments and suggestions, which greatly helped improve our manuscript. Below are our point-by-point responses and detailed corrections/revisions made to the manuscript

15 accordingly.

Report #2

Suggestions for revision

20 Usually, the term “EC” should be used here instead of BC as thermal-optical method was applied here. I strongly suggest use EC in this paper. And I noticed that TSP samples were collected, I doubt that carbonate carbon (CC) may be important in coarse fractions. CC will influence on the results of OC and EC as well as the estimated SOC. SOC estimation has a large uncertainty when biomass burning  
25 emissions were important. I suggest removing SOC part as such an estimation may be not convincing.

Response: We agree with the referee that the refractory constituent of the carbonaceous aerosols determined using the thermal-optical method should be called

30 “elemental carbon” rather than “black carbon”, which describes the light-absorbing

part determined using optical methods or in aerosol-climate models. Following the suggestions, we have replaced “BC” with “EC” when describing our measurements and analysis in the revised paper (with “BC” kept for the general discussion and the modeling part).

- 5 According to the suggestion, we have also removed the SOC-related materials (analyses, results and figures) from the revised manuscript.

#### Comments from editor Xiaobin Xu

- 10 Comments to the Author:

Dear authors:

Your revised manuscript has been reviewed by two referees. Additional revisions are suggested by referee #2. Please improve your manuscript further by considering the suggestions of referee #2 and also following comments from me:

15

1. Page 5 line: Ssandradewi et al., 2008 is not given in the references list. A thorough check of citations is needed.

Response: We have added (Sandradewi et al., 2008) to the reference list, and

- 20 thoroughly checked the citations in the manuscript.

2. Page 6 line 9 and page 13 line 8: change "monsoons season" to "monsoon season".

Response: Corrected.

25

3. Page 6 lines 11-15: I do not think you should give elevations and coordinates of that accurate, in particular on the glaciers.

Response: Following the suggestion, we have reduced the number of significant

- 30 digits for the elevations and coordinates.

4. Page 6 line 30: change "atmospheric gas" to "air".

Response: Done.

5

5. Page 7 lines 2-3: "using a stable vacuum pump, which had a good quality and performance (e.g., running steadily with low noise level)." It is better to give pump's model and manufacturer information instead of these vague expressions.

10 Response: We have added the pump's model and manufacturer information in the revised manuscript as the following:

"..., and each sample was collected for 24 h using a Thomas pump (2628TE32, America)."

15 6. Page 11 lines 7-10: What do you mean by "photochemical ability of OC" and "little material available for external-coating of BC"?

Response: The phrase "photochemical ability of OC" in the previous manuscript was a mistake, which has been corrected in the revised manuscript. We meant to  
20 say that relatively weak solar radiation in winter would result in less production of organic compounds through photochemical reactions.

We have revised the original sentence to "Lower ratio of OC/EC in the other seasons (winter, pre-monsoon, post-monsoon) was probably due to the less photochemical production of secondary organic compounds as coating material on  
25 EC particles (Knox et al., 2009; Cappa et al., 2012)" in the revised manuscript.

7. Page 11 and lines 19-21: How can this happen?

Response: The monsoon season is the peak tourism season and winter season is the  
30 off-season in GHZ. Thus the amount of vehicle emissions from touring buses and private vehicles had seasonal difference, which we believe played an important role

in the seasonal variations of OC/EC ratios at the sampling site. Since we don't have direct observational evidence to support this, we have revised the statement to: "Seasonal differences in vehicle emissions from touring buses and private vehicles in the GHZ basin might have played an important role in the seasonal variations of

5 OC/EC ratios".

8. Fig. 2 and some other figures: there are obviously some data gaps. You should not fill the gaps with lines.

10 Response: The data gaps are now unfilled with lines in the revised manuscript (i.e. Fig. 2, Fig. 4, and Fig. 5).

9. Figs. 9 and 10: displaying MAE data repeatedly in all plots is not necessary. If you are discussing the correlations of MAE with other quantities, it is better to show the  
15 data as scatter plots with regression results.

Response: This is an excellent suggestion. We have now shown the data as scatter plots with regression results in the revised manuscript. The following are included as new figures (Figs. 9 and 10).

20




Fig. 9




Fig. 10

## References

- Cappa, C. D. et al.: Radiative absorption enhancements due to the mixing state of atmospheric black carbon, *Science*, 31, 337, 1078-1081, 2012.
- 5 Knox, A., Evans, G. J., Brook, J. R., Yao, X., Jeong, C. H., Godri, K. J., Sabaliauskas, K., and Slowik, J. G.: Mass absorption cross-section of ambient black carbon aerosol in relation to chemical age, *Aerosol Sci. Technol.* 43, 522-532, 2009.

10

15

20

25

30

# **Seasonal variation and light absorption property of carbonaceous aerosol in a typical glacier region of the southeastern Tibetan Plateau**

Hewen Niu<sup>a</sup>, Shichang Kang<sup>a,\*</sup>, Hailong Wang<sup>b,\*</sup>, Rudong Zhang<sup>b,c</sup>, Xixi Lu<sup>d</sup>, Yun Qian<sup>b</sup>,

5 Rukumesh Paudyal<sup>a</sup>, Shijin Wang<sup>a</sup>, Xiaofei Shi<sup>e</sup>, Xingguo Yan<sup>f</sup>

<sup>a</sup> State Key Laboratory of Cryospheric Science, Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, Lanzhou 730000, China.

<sup>b</sup> Atmospheric Sciences and Global Change Division, Pacific Northwest National Laboratory

10 (PNNL), Richland, WA 99352, USA

<sup>c</sup> Institute for Climate and Global Change Research, School of Atmospheric Sciences, Nanjing University, Nanjing 210093, China; Collaborative Innovation Center of Climate Change, Jiangsu Province, China.

<sup>d</sup> Department of Geography, National University of Singapore, 1 Arts Link, Singapore 117570,

15 Singapore

<sup>e</sup> College of Earth Environmental Sciences, Lanzhou University, Lanzhou 730000, China.

<sup>f</sup> College of Geography and Environmental Science, Northwest Normal University, Lanzhou 730030, China

\* Corresponding authors: shichang.kang@lzb.ac.cn; Hailong.Wang@pnnl.gov

20

25

## **Abstract:**

Deposition and accumulation of light-absorbing carbonaceous matters on glacier surfaces can alter the energy balance of glaciers. In this study, two years (December 2014 to December 2016) of continuous observations of carbonaceous aerosols in 5 glacierized region of Mt. Yulong and Ganhaizi (GHZ) basin are analyzed. The average elemental carbon (**EC**) and organic carbon (OC) concentrations were  $1.51 \pm 0.93$  and  $2.57 \pm 1.32 \mu\text{g m}^{-3}$ , respectively. Although the annual mean OC/**EC** ratio was  $2.45 \pm 1.96$ , monthly mean **EC** concentrations during the post-monsoon season were even higher than OC in the high altitudes (approximately 5000 m asl) of Mt. Yulong.

10 Strong photochemical reactions and local tourism activities were likely the main factors inducing high OC/**EC** ratios in the Mt. Yulong region during the monsoon season. The mean mass absorption efficiency (MAE) of **EC**, measured for the first time in Mt. Yulong, at 632 nm with a thermal-optical carbon analyzer under the filter-based method, was  $6.82 \pm 0.73 \text{ m}^2 \text{ g}^{-1}$ , comparable with the results from other

15 studies. Strong seasonal and spatial variations of **EC** MAE were largely related to the OC abundance. Source attribution analysis using a global aerosol-climate model, equipped with a black carbon (**BC**) source tagging technique, suggests that East Asia emissions, including local sources, have the dominant contribution (over 50%) to annual mean near-surface BC in the Mt. Yulong area. There is also a strong seasonal variation in the regional source apportionment. South Asia has the largest contribution

20 in BC emission during the pre-monsoon season, while East Asia dominates the monsoon season and post-monsoon season. Results in this study have great implications for accurately evaluating the influences of carbonaceous matter on glacial melting and water resource supply in glacierization areas.

## 1. Introduction

Carbonaceous aerosols play an important role in Earth's climate system and energy budget (Bond et al., 2007, 2013; Schuckmann et al., 2016). It has sophisticated/complex effects on the human health and living species (Jerret et al., 5 2005), visibility (Park et al., 2003), atmospheric radiative balance (Bond et al., 2013; Schuckmann et al., 2016), and the surface albedo of snow and ice (Gertler et al., 2016; Kaspari et al., 2014; Niu et al., 2017a, b). Carbonaceous aerosol is an aggregate of thousands of species with various thermal, physicochemical and optical properties (Andreae and Gelencsér, 2006; Cheng et al., 2011a). In the atmosphere, carbonaceous 10 aerosols affect the radiative balance by absorbing and scattering solar radiation and affecting the properties of clouds (IPCC, 2013; Lohmann and Feichter, 2005; Carslaw et al., 2010). In the cryosphere, deposition of carbonaceous matter on snow and glaciers reduces the surface spectral albedo (snow darkening) (Flanner et al., 2009; Doherty et al., 2013; Qian et al., 2015; Niu et al., 2017a) and accelerates snow/glacial 15 melting (Hansen and Nazarenko, 2004; Xu et al., 2009a).

Carbonaceous matter in smoke/emissions from biomass burning and fossil fuel combustion has been identified as the typical atmospheric pollutant since historical period (Brimblecombe, 1987). Recently, scientific attention has shifted from the role of carbonaceous matter as atmospheric pollutant to its influence as one of driving 20 factors of climate change (Andreae, 1995; Andreae and Gelencsér, 2006; Hansen et al., 2005; Ramanathan et al., 2005). Some model simulations proposed that the radiative forcing of black carbon (BC) is comparable to that of methane (Chung and Seinfeld, 2005; Jacobson, 2004), suggesting that BC may be the second most important warming agent (only after CO<sub>2</sub>) in terms of direct radiative forcing (Jacobson, 2001). 25 Ding et al. (2016) found that BC particles play a key role in modifying/heating the planetary boundary layer (PBL) and enhancing the haze pollution, called the “dome effect” of BC, and suggested an urgent need for reducing BC emissions to mitigate the extreme haze pollution in megacities in China. In addition, high concentrations of absorbing aerosols (e.g., BC, brown carbon, and/or dust) over eastern China during 30 winter and spring were found to be the major reason for the observed recent warming

trend (Yu et al., 2001). BC in snow can increase the surface air-temperature by approximately 1.0 °C over the Tibetan Plateau (TP) and reduce spring snow cover (Qian et al., 2011). Generally, sampled carbonaceous aerosols can be divided into elemental carbon (EC) and organic carbon (OC) using thermal/optical reflectance (TOR) method (Cao et al., 2010; Chow et al., 1993). EC is also known as BC when measured using optical methods or in aerosol-climate modeling studies (e. g., Cheng et al., 2011a; Ming et al., 2013; Xu et al., 2009b; Wang et al., 2014; Wang et al., 2015). Moreover, in the low-latitude and high-elevation areas, extensive incoming solar radiation and large amount of carbonaceous aerosol deposited on snowpack and glaciers result in surface albedo reduction and the retreat of glaciers in the TP, and further affect Asian hydrological cycle and monsoon climate (Qian et al., 2011; Qu et al., 2014; Li et al., 2016a). This is closely related to water resources for a large population of local habitants in South Asia (Ramanathan et al., 2005, 2007). Therefore, it is rather important and necessary to carry out carbonaceous aerosol study in glacierization regions.

The mass absorption efficiency (MAE,  $\text{m}^2 \text{ g}^{-1}$ ) is a typical parameter characterizing the optical (or light absorbing) properties of aerosols. For EC particles, it is determined by the mass concentration ( $\mu\text{g m}^{-3}$ ) and absorption coefficient ( $b_{\text{abs}}$ ,  $\text{Mm}^{-1}$ ) of EC (Lioussse et al., 1993; Cheng et al., 2011b), where  $b_{\text{abs}}$  is the cross section of EC available to absorb light (Bond and Bergstrom 2006a; Knox et al., 2009). MAE of EC is usually estimated using quartz filter based methods, which detect the change in the light transmittance through a quartz filter due to the presence of EC particles (Sharma et al., 2002; Knox et al., 2009; Cheng et al., 2011a). Bond and Bergstrom (2006a) suggested a mass-normalized MAE of  $7.5 \pm 1.2 \text{ m}^2 \text{ g}^{-1}$  at 550 nm for uncoated aerosol particles. However, Ram and Sarin (2009) studied the  $b_{\text{abs}}$  and MAE of EC in aerosols sampled at three typical sites in India, and they found a distinct spatio-temporal variability in  $b_{\text{abs}}$  and EC MAE on a regional scale. Discrepancies are sometimes inevitable for the filter based techniques related to aerosol-filter interactions (Cheng et al., 2011a; Sandradewi et al., 2008). Moreover, the MAE of EC can be largely influenced by the aerosol mixing state (Bond et al., 2006b; Doran et al., 2007;

Jacobson, 2001; Schnaiter et al., 2005). It was proposed that non-carbon species (e.g., sulphate, nitrate) can increase the MAE value of EC (Knox et al., 2009) because the coating by other components can focus light into the EC core of the particle (Bergstrom et al., 1982; Cheng et al., 2011a). Enhancement of MAE by coating can be 5 described in terms of absorption amplification that is largely independent of coating thickness (Schnaiter et al., 2005; Knox et al., 2009). Therefore, it is quite necessary to make a further analysis on EC MAE to reduce the uncertainties in evaluating light absorption of carbonaceous aerosols.

Recently, in addition to EC, it has been recognized that certain OC components in 10 carbonaceous aerosol can also absorb light (Andreae and Gelencsér, 2006; Cheng et al., 2011a). However, light absorption by OC has not yet been taken into consideration in many climate models, e.g., various versions of the Community Earth System Model (CESM) (Flanner and Zender, 2006; Wang et al., 2013; Qian et al., 2015; Liu et al., 2016), which causes certain uncertainties in precisely evaluating climate/radiative 15 forcing of carbonaceous aerosol in the atmosphere and snow/ice. The radiative forcing of carbonaceous aerosol remains one of the great challenges in climate simulation (Jacobson, 2001; IPCC, 2013).

In this study, total suspended particulates (TSP) were simultaneously collected at 20 two remote stations on Yulong Snow Mountain (abbreviated Mt. Yulong), in the southeast fringe of the TP. Small scale spatial and seasonal variations of atmospheric carbonaceous matter are examined, and the corrected EC MAE is calculated to evaluate light-absorbing property of carbonaceous aerosol in the typical glacierization area. Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) is used to retrieve 25 total attenuated backscatter and sub-type information of smoke plumes to help on source attribution of carbonaceous aerosols in Mt. Yulong.

## 2. Material and methods

### 2.1 Study area and TSP sampling

The study area, Mt. Yulong ( $26^{\circ}59' - 27^{\circ}17'$  N,  $100^{\circ}04' - 100^{\circ}15'$  E), is the southernmost glaciated mountain in the Eurasian continent (Fig. 1). The Baishui 30 glacier ( $27^{\circ}06'$  N;  $100^{\circ}11'$  E) on Mt. Yulong is a typical temperate glacier that has

large energy fluxes, particularly at lower snow-covered elevations. The climate of Mt. Yulong is typically affected by the Indian summer monsoon and East Asian summer monsoon (Nie et al., 2017) in the monsoon season (June-September), and characterized by other three distinct seasons: post-monsoon (Oct. - Nov.), winter (Dec. 5 - Feb.), pre-monsoon (Mar. - May) season (Chen et al., 2015; Cong et al., 2015b; Bonasoni et al., 2010; Niu et al., 2013). Annual precipitation in Mt. Yulong occurs considerably (~80%) in the monsoon season (Fig. 2). There is little precipitation in winter when the westerly jet dominates (Liu et al., 2017).

One of the TSP sampling sites is at the elevation of 4510 m asl, which is close to 10 the location of the upper station ( $27^{\circ}06' \text{ N}$ ,  $100^{\circ}11' \text{ E}$ ) of tourism cableway on Mt. Yulong. Ganhaizi (GHZ) basin, the other TSP sampling site ( $27^{\circ}06' \text{ N}$ ,  $100^{\circ}15' \text{ E}$ ), is located on the east side and the foot of Mt. Yulong, at an elevation of 3054 m asl. It is separated from the urban area. However, it becomes an increasingly popular tourist destination, with a geological museum and a golf course. Moreover, there is a 15 spacious parking lot and a tourist dining-center in GHZ. Besides the major emissions from tourist vehicles, there are some other limited pollution sources, such as agricultural waste burning, (open fire) biomass burning, and crustal aerosols (Niu et al., 2014, 2016) near the study sites. These two sampling sites are located on the southeast fringe of the TP, away from urban areas, and thus considered as typical 20 remote areas in the Northern Hemisphere (Li et al., 2016a) and ideal observation areas for atmospheric environment in glacierization region.

The TSP samples analyzed in this study were collected from December 2014 to December 2016 at these two sites using a particulate sampling apparatus (TH150-A, Wuhan Tianhong INST Group). The sampling apparatuses were placed 15 m above 25 the ground, away from the surface dust and any specific pollutant sources. The quality assurance of this apparatus (TH150-A) is demonstrated by the difference between manually calculated volume of gas and automatically recorded value. The volume of air was usually calculated automatically by the apparatus. The air (and suspended particulates) was sampled at a flow rate of  $100 \text{ L min}^{-1}$  with an accuracy of  $\pm 2.5\%$ , 30 and each sample was collected for 24 h using a Thomas pump (2628TE32, America).

The sampling interval for each TSP sample was 6 days. The number of aerosol samples collected at Mt. Yulong and GHZ was 117 and 120, respectively. The collected TSP samples were loaded on 90 mm (in diameter) pre-combusted (heating at 550 °C for 6 h in an oven) quartz fibre filters (Whatman Corp.).

5 After sampling, the quartz filters were wrapped with an aluminum foil and were kept in a refrigerator at 4 °C in the Yulong Glacier Station of Chinese Academy of Sciences in Lijiang city, and then were transported to the State Key Laboratory of Cryospheric Science, Northwest Institute of Eco-Environment and Resources in Lanzhou for chemical analysis.

## 10 2.2 Sample analysis

OC and EC on the quartz filters were analyzed using a Desert Research Institute (DRI) Model 2001 thermal-optical reflectance carbon analyzer (Atmoslytic Inc., Calabasas, California) to determine the EC mass and optical attenuation (ATN) (Chow et al., 2001, 2004; Cheng et al., 2011a; Li et al., 2016b; Niu et al., 2017a). We adopted 15 the improved U.S. Interagency Monitoring of Protected Visual Environments (IMPROVE)-A thermal/optical reflectance protocol (Niu et al., 2017b). Every filter was analyzed for a portion of carbon in a 0.502 cm<sup>2</sup> punch. A temperature peak (550 °C) was designed to reduce measuring time that EC exposed in the catalyzing atmosphere. The applied heating conditions permitted the separation of EC portions in 20 O<sub>2</sub> (2%) and helium (98%) atmosphere and OC portions in a helium atmosphere (Wang et al., 2015; Niu et al., 2017a). The residence time of each heating step was determined by the stabilization of the carbon signal.

## 2.3 Calculation methods of ATN, MAE and POC

### 2.3.1 ATN calculation

25 The ATN is calculated based on the transmittance signal during the filter analysis, defined as

$$ATN = \ln\left(\frac{I_0}{I}\right)$$

Where,  $I_0$  and  $I$  are the transmittance signal after and before the thermal/optical analysis (Ram and Sarin, 2009). Lights passing through a particle-loaded and a blank

filter were simultaneously measured in the determination of ATN by the (thermal-optical reflectance) carbon analyzer. ATN determined by the carbon analyzer is similar to the Aethalometer (Cheng et al., 2011a). Previous studies have demonstrated that ATN of blank quartz fibre filters averaged  $0.00 \pm 0.01$ , suggesting 5 that the ATN of loaded quartz filter could generally be ascribed to the existence of light-absorbing carbon (Yang et al., 2009; Cheng et al., 2011a).

### 2.3.2 MAE calculation

The mass absorption efficiency (MAE) is calculated as:

$$MAE = \frac{ATN}{EC_s} \times 10^2 \times \frac{1}{C}$$

10 Where,  $EC_s$  ( $\mu\text{g C cm}^{-2}$ ) is the filter loading amount of EC, which is directly measured from the thermal optical analysis. The filter-based determination of light absorption has many artifacts, though the various scattering effects can be properly corrected by an empirical factor,  $C$ . A value of  $C = 3.6$  was proposed for the internally mixed atmospheric aerosol when employing thermal-optical analysis method in several 15 studies (Weingartner et al., 2003; Doran et al., 2007; Ram and Sarin, 2009; Cheng et al., 2011a). The same empirical factor was also used in the optical measurement by the Aethalometer (Ram and Sarin, 2009). There are many factors (e.g., measurement methods, mixed states) accounting for the discrepancy of MAE. The corrected equation of MAE calculation (i.e., corrected for the multiple scattering effects) 20 performed in this study has greatly diminished the uncertainties (around 15%). By converting previously published MAE values (Ram and Sarin, 2009) to the equivalent-MAE, Cheng et al (2011a) has found that the equivalent-MAE was much lower in the regions severely affected by biomass burning (e.g., Allahabad, India).

The dependence of light attenuation (ATN) detected at 632 nm on EC loading ( $EC_s$ ) 25 is shown in Fig. 3, to identify the artifact relevant to filter-based measurements. As a result, ATN and  $EC_s$  correlate well ( $R^2=0.83$ ) with a slope ( $K$ ) of  $0.08 \text{ m}^2 \text{ g}^{-1}$  and an intercept ( $b$ ) of  $0.35 \text{ m}^2 \text{ g}^{-1}$  for our samples at the Mt. Yulong sampling site. Strong correlation ( $R^2=0.62$ ) between ATN and  $EC_s$  also exists for the GHZ sampling site, with  $K=0.08 \text{ m}^2 \text{ g}^{-1}$  and  $b=0.45 \text{ m}^2 \text{ g}^{-1}$ . Strong dependence of ATN on  $EC_s$  has been

found in the present study, and artifact associated with filter-based method was not identified and thus can be neglected.

### 2.3.3 Primary OC (or POC) calculation

$$OC_{pri} = (EC \cdot (OC/EC)_{min}, OC_{tot})_{min}$$

- 5 where  $(OC/EC)_{min}$  is the minimum OC/EC ratio in a specific set of data, the same sets of samples as used in this study.  $OC_{pri}$  indicates the POC content.

## 2.4 Model experiment

We use a global aerosol-climate model, the Community Atmosphere Model version 5 (CAM5) equipped with a BC (or EC) source tagging technique (Wang et al., 2013, 10 2014; Zhang et al., 2015), to help estimate the source attribution of BC in the Mt. Yulong area. The 4-mode modal aerosol scheme of CAM5 recently developed by Liu et al. (2016) is used here, in which BC and primary OC particles are emitted into a primary-carbon mode. Then they grow through condensation of gas-phase precursors (e.g., sulfuric and organic gases) and move to the accumulation size mode, where 15 hygroscopic aerosol particles, including carbonaceous aerosols, are subject to wet removal by precipitation.

The CAM5 experiment is conducted for five years (2010-2014) at horizontal grid spacing of  $1.9^\circ \times 2.5^\circ$  and 30 vertical levels. The sea surface temperatures and sea ice concentrations are prescribed with observations, and winds are constrained with 20 reanalysis from NASA Modern Era Retrospective-Analysis for Research and Applications (MERRA) (Rienecker et al., 2011; Ma et al., 2013). Monthly mean anthropogenic and open fire emissions (Hoesly et al., 2018; van Marle et al., 2017), including primary OC and BC, used in the simulation come from the recently released datasets for the Coupled Model Intercomparison Project Phase 6 (CMIP6), which are 25 only available up to year 2014. Therefore, the model experiment is not designed to simulate the two-year observations of aerosols in TP, but rather for a recent time period (2010-2014) to estimate the mean source attributions and seasonal variations of near-surface BC concentrations.

## 3. Results and discussion

- 30 3.1 Characteristics of the carbonaceous aerosols

Temporal variations of carbonaceous matter measured from Mt. Yulong are shown in Fig. 4. Distinct seasonal differences presented during the sampling time period. The winter season and post-monsoon season had higher concentrations of carbonaceous matter in the TSP during the two years, which is consistent for EC, OC, and POC.

5 However, the OC/EC ratio showed an opposite seasonal contrast. Monsoon season in 2016 had the lowest carbonaceous matter contents in the two-year time period (Fig. 4), whereas the concentrations of OC, EC, and POC in the monsoon and pre-monsoon season in 2015 presented relatively high and low values, respectively. This is uncharacteristic, compared to the general seasonal variations of atmospheric

10 chemistry (Kang et al., 2004, 2007; Niu et al., 2013, 2016). It is quite likely that frequent rainfall events with occasional dust (e.g., Dong et al., 2011; Niu et al., 2014) from anthropogenic activities (Shrestha et al., 2000) during the monsoon in 2015 are responsible for this unusual phenomenon (i.e. relatively high content in monsoon season in 2015). Note that the minimum value of OC/EC ratio is used in the POC

15 calculation. It varies greatly among different seasons (i.e., 0.38, 0.71, 0.42 and 0.35 for winter, pre-monsoon, monsoon, and post-monsoon in Mt. Yulong, respectively), so seasonal minimum values are used to estimate POC concentrations in the corresponding seasons.

In comparison to Mt. Yulong, the trends of carbonaceous matter from the GHZ basin presented more distinct seasonal variations (Fig. 5), i.e., the monsoon season regularly had the lowest concentrations of EC, OC, and POC. The OC/EC ratio was consistently opposite to that, e.g., higher values of OC/EC appeared in monsoon seasons. Lower ratio of OC/EC in the other seasons (winter, pre-monsoon, post-monsoon) was probably due to the less photochemical production of secondary

20 organic compounds as coating material on EC particles (Knox et al., 2009; Cappa et al., 2012). In addition, seasonal changes in EC and OC sources (e.g., biomass burning vs. fossil fuel combustion) might play an important role for the variations of OC/EC ratios. An obviously higher EC and OC concentrations were found in the post-monsoon season at the Mt. Yulong site, and in winter season at the GHZ site (Fig.

25 4 and 5) when wet removal by precipitation is inefficient. This suggests the

importance of seasonal changes in sources (Carrico et al., 2003; Cong et al., 2015b; Wan et al., 2017). In addition, OC/**EC** ratio was usually employed to evaluate the combustion fuel sources. Previous studies reported that the global mean of OC/**EC** by biomass burning was higher than fossil fuel burning (Bond et al., 2004; Cao et al., 5 2010; Liousse et al., 1996). Seasonal differences in vehicle emissions from touring buses and private vehicles in the GHZ basin might have played an important role in the seasonal variations of OC/**EC** ratios.

Table 1 summarizes statistical results of **EC**, **OC**, and POC concentrations in atmospheric aerosol from Mt. Yulong. The annual mean **EC** concentration was  $1.53 \pm 1.49 \mu\text{g m}^{-3}$ , with the sampled values ranging from 0.02 to  $6.83 \mu\text{g m}^{-3}$  during the study period (December 2014-Decmber 2016). The post-monsoon season had the highest **EC** content of  $3.51 \pm 1.20 \mu\text{g m}^{-3}$ , ranging from 1.22 to  $5.80 \mu\text{g m}^{-3}$ . The annual average OC concentration was  $1.65 \pm 1.14 \mu\text{g m}^{-3}$ , with the lowest concentrations in the monsoon season ( $1.34 \pm 0.90 \mu\text{g m}^{-3}$ ) and the highest 10 concentrations in the post-monsoon season (OC:  $2.57 \pm 1.30 \mu\text{g m}^{-3}$ ). Similar seasonal differences were also found in other areas such as Lumbini, Nepal (Wan et al., 2017), Nepal Climate Observatory-Pyramid (NCO-P) (Bonasoni et al., 2010), Kanpur, India (Ram et al., 2012), and Delhi, India (Mandal et al., 2014). Moreover, monthly averaged **EC** and OC concentrations were analyzed for Mt. Yulong (Fig. 6). It shows 15 that from the monsoon season to the post-monsoon season (particularly from September to December), **EC** concentrations were higher than that of OC. A large amount of biomass burning emissions in the high atmosphere (around 5000 m asl) in Mt. Yulong were probably transported from distant source regions which can be determined by the analysis of CALIOP retrievals and source apportionment of 20 carbonaceous aerosols (see section 3.4). In addition, we performed trajectory analysis with the HYSPLIT model. From the monsoon season to post monsoon season, the trajectories of air mass reaching the sampling location changed much. In the monsoon season, the air mass (and pollutants) mainly originated from southwest and southeast monsoons, while in the post monsoon season it mainly came from the west (Fig. 7).

30 The annual mean OC/**EC** ratio was found to be highest in the monsoon season and

lowest in winter. The aging process of EC (or soot) resulting from photochemical oxidation by molecular O<sub>2</sub> and the photooxidation of OC (Han et al., 2012) were likely involved and increased OC/EC ratio. Photochemical reaction can change their physical and chemical properties from the original molecule of the substance (alkyne C-H ( $\equiv$ C-H) and aromatic C-H (Ar-H)) (Kirchner et al., 2000; Cain et al., 2010). When this occurs, these molecules tend to form a new structure by combining with each other or with other molecules (carbonyl C=O and ether C-O) (Daly and Horn, 2009; Cain et al., 2010; Nieto-Gligorovski et al., 2008), which may change the state of OC, EC or OC/EC ratios. The photochemical oxidation by O<sub>2</sub> under sunlight is an important aging process for EC (Han et al., 2012). Statistical results of EC, OC, and POC concentrations in aerosol from the GHZ site are shown in Table 2. The annual average EC concentration,  $1.50 \pm 0.90 \mu\text{g m}^{-3}$ , is comparable with that from Mt. Yulong ( $1.53 \pm 1.49 \mu\text{g m}^{-3}$ ). However, the annual average concentration of OC,  $3.50 \pm 1.50 \mu\text{g m}^{-3}$ , was greatly higher than that from Mt. Yulong. Remote sources are likely to have a similar impact on aerosols over the two sites that are fairly close to each other. Therefore, the additional OC in the GHZ basin was more likely contributed by local sources such as fossil fuel (vehicle) emissions associated with the frequent and intense tourism activities, which is totally different from that of Mt. Yulong. Maximum seasonal mean EC and OC concentration occurred in the winter season (EC:  $2.42 \pm 0.74 \mu\text{g m}^{-3}$ , OC:  $4.09 \pm 1.70 \mu\text{g m}^{-3}$ , respectively) in the GHZ basin, and their lowest concentrations conventionally occurred in monsoons season (EC:  $0.75 \pm 0.30 \mu\text{g m}^{-3}$ , OC:  $2.80 \pm 0.80 \mu\text{g m}^{-3}$ , respectively). The seasonal variation of carbonaceous aerosols found in the GHZ basin was different from that found in Mt. Yulong. It is likely related to the distinct elevation difference (nearly 1,500 m) and different amount of local human activities (here mainly referring to tourism related activities) between the Mt. Yulong and the GHZ sampling sites. The GHZ site location is close to a parking lot for private vehicles and touring buses and a visitor service center that involves food cooking. These tourism activities can contribute to local emissions of carbonaceous aerosols and precursor gases for OC (Borrego et al., 2000; Cong et al., 2015a; Shi et al., 2017). However, we don't have direct

observational evidence to support this.

The OC/**EC** ratios at the two sites were relatively low, and they have distinct seasonal variations and spatial differences over the Mt. Yulong region. The annual average value was  $2.06 \pm 3.38$  for Mt. Yulong, with the highest value occurred in the 5 pre-monsoon season ( $3.67 \pm 5.70$ ) and the lowest value in the post-monsoon season ( $0.79 \pm 0.40$ ) (Table 1). Monthly variation of the average OC/**EC** ratio was determined by the relative concentrations of **EC** and OC in aerosol, for example, the lowest OC/**EC** ratio occurred in post-monsoon was due to substantially high **EC** concentrations in that season (Fig. 6). The annual mean OC/**EC** ratio in samples from 10 the GHZ basin was  $2.90 \pm 1.80$ , while the monsoon season had the highest value ( $4.40 \pm 2.00$ ) and the winter season had the lowest value ( $1.93 \pm 1.58$ ) (Table 2). Previous studies suggested that OC/**EC** ratios from biofuel and biomass burning emissions are generally higher than those from fossil fuel combustion (Cao et al., 2013; Ram et al., 2012; Cong et al., 2015b; Wan et al., 2017). Strong photochemical reactions (due to 15 extensive solar radiation) (Fig. 2) and tourism activities in the monsoon season (is the peak season for tourism) were likely the main factors that result in relatively high OC/**EC** ratios in GHZ. In addition, high OC/**EC** ratios are closely related to the OC concentration ( $2.80 \pm 0.80 \mu\text{g m}^{-3}$ ) in the monsoon season in GHZ site (Table 2) when extensive photochemical processes were proceeded (e.g., Antony et al., 2011; 20 Schneidemesser et al., 2009).

### 3.2 Optical properties of **EC**

The corrected MAE values of **EC** at 632 nm (calculated using the equation in section 2.3.3) were  $7.38 \pm 1.01$  and  $6.25 \pm 0.46 \text{ m}^2 \text{ g}^{-1}$  for Mt. Yulong and GHZ samples, respectively. The **EC** MAE has distinct seasonal variations, with the peak of 25 **EC** MAE values in the pre-monsoon and monsoon seasons at Mt. Yulong and GHZ, respectively (Fig. 8). The high MAE values suggest an enhancement of MAE (or absorption amplification) by external-coating with OC (Cheng et al., 2011a; Knox et al., 2009; Schnaiter et al., 2005). Strong seasonal and spatial differences of **EC** MAE values in Mt. Yulong and GHZ largely related to OC abundance. Furthermore, 30 correlation analysis among MAE and OC, EC<sub>s</sub> concentrations, OC/**EC** and POC/OC

ratios in Mt. Yulong and GHZ aerosols were performed. Strong correlations were found between EC MAE and OC/EC, POC/OC ratios ( $R^2 = 0.55$  and  $0.40$ , respectively,  $p < 0.01$ ), as well as  $EC_s$  ( $R^2 = 0.58$ ,  $p < 0.01$ ) in Mt. Yulong aerosol (Fig. 9). While the coefficient of determination  $R^2 = 0.57$  ( $p < 0.01$ ) for MAE and OC/EC ratio,  $R^2 = 0.47$  ( $p < 0.01$ ) for MAE and POC/OC ratio, and  $R^2 = 0.69$  ( $p < 0.01$ ) for MAE and  $EC_s$  in GHZ aerosol (Fig. 10). As expected, correlation between MAE and OC is weak for aerosols at Mt. Yulong ( $R^2 = 0.18$ ,  $p < 0.01$ ) and GHZ ( $R^2 = 0.11$ ,  $p < 0.01$ ). The strong correlations between MAE and OC/EC, POC/OC ratio appear to be influenced by the abundance of OC and other particulate (e. g., sulfate) (Omar et al., 1999). The availability of POC for external-coating is responsible for the variations of mean EC MAE (Knox et al., 2009). Values of atmospheric EC MAE are also dependent on the extent of internal-mixing of the EC with other substances (Cappa et al., 2012; Schnaiter et al., 2005). Atmospheric EC light absorption is linearly proportional to the EC concentration since EC particles are small enough (Schwarz et al., 2013).

Many previous studies have quantified the EC MAE values at various sites (Bond and Bergstrom, 2006a; Cheng et al., 2011a; Knox et al., 2009; Ram and Sarin, 2009; Li et al., 2016c). However, large uncertainties exist among different calculation approaches. Measurement methods of ATN and  $EC_s$  (various temperature protocols) definitely affect the EC MAE (Cheng et al., 2011a, b; Li et al., 2016c). In addition, brown carbon (BrC) appearing in the particle mixture can decrease the EC MAE (Jeong et al., 2004; Hecobian et al., 2010). BrC is less absorptive comparing with pure EC (Cheng et al., 2011a). BrC emitted from biomass burning considerably lower MAE values (Jeong et al., 2004). Whereas coating by organic aerosol or mixing-state can enhance the MAC values (Knox et al., 2009; Zhang et al., 2008). Higher absorption cross-sections result from coating processes in the atmosphere (Andreae and Gelencsér, 2006; Bond et al., 2013). EC particles in aerosol, due to condensation processes and/or cloud processing, acquire non-absorbing coatings (mainly sulfate and OC), which lead to absorption enhancements (Andreae and Gelencsér, 2006; Fuller et al., 1999; Schnaiter et al., 2005).

### 3.3 Controls of carbonaceous matter components

Our results show that carbonaceous matter (**EC**, OC) in aerosol exhibited a discernable small-scale spatial variation between Mt. Yulong and GHZ. In addition to the elevation difference, other potential factors such as the more tourism activities near GHZ than at Mt. Yulong could partly account for the difference.

In addition, inter-annual differences of carbonaceous aerosol from Mt. Yulong and GHZ were also distinct (Fig. 11). The annual mean concentrations of carbonaceous matter collected in 2016 were lower than those in 2015 for the two sites (Fig 11a, b), which could be partly due to the strict mitigation measures that improved the local atmospheric environment in the Mt. Yulong region. For example, the amount of soot emissions (2.44 Tg C) in 2016 were reduced by 21.76% compared to 2015 in Yunnan province (<http://www.zhb.gov.cn/>), where Mt. Yulong locates. Other factors likely contributed to the inter-annual difference as well. The amount of precipitation is important to determine aerosol wet removal from the atmosphere during the transport (e.g., Wang et al., 2013). The stronger precipitation in 2016 than in 2015 at Mt. Yulong and GHZ (Fig. 11a, b) partly explains the smaller carbonaceous aerosols mass concentrations in 2016. The average OC/**EC** ratios also decreased in 2016 compared to 2015, whereas the mean POC/OC ratios have no obvious difference between the two years in both Mt. Yulong and GHZ (Fig. 11).

We also compare atmospheric **EC** concentrations in Mt. Yulong with other (results derived from TSP samples) interested areas. It shows that **EC** concentrations in Mt. Yulong aerosol were relatively low and, among the compared values, most of **EC** concentrations were within the range of 1.0-2.5  $\mu\text{g m}^{-3}$  (Fig. 12), while some of values were extremely low (close to 0.5  $\mu\text{g m}^{-3}$ ) or high (above 6.0  $\mu\text{g m}^{-3}$ ). The low **EC** concentrations were typically found in the TP, e.g., QOMS (i. e. Qomolangma (Mt. Everest) Station for Atmospheric and Environmental Observation), TP (Cong et al., 2015b), pre-monsoon season in Mt Yulong. Whereas the **EC** concentrations in Agra, India ( $6.1 \pm 0.83 \text{ g m}^{-3}$ ) (Pachauri et al., 2013) were almost three folds of values found in Mt. Yulong. Agra and Lumbini have been identified as regions in the world that are highly affected by biomass burning (Wan et al., 2017). A large amount of

carbonaceous aerosols emitted from those regions can reach Mt. Yulong by crossing the Himalayas (e.g., Lüthi et al., 2014), which substantially influence the Mt. Yulong region.

### 3.4 Source apportionments of carbonaceous aerosols

5     Aerosol vertical distributions from CALIOP retrievals often reveal that smoke plume could reach approximately 6 km (Fig. 13), which is higher than most of the mountains and mountain glaciers in the Himalaya regions. Some typical CALIOP/CALIPSO transects, having strong backscattering signal (i.e., 532 nm total attenuated backscatter), show spatially continuous atmospheric pollutant layers  
10 from the ocean all the way to Mt. Yulong (Fig. 13), indicating a penetration of smoke plume toward the inland of the TP. Carbonaceous aerosols is an important anthropogenic driving force of the observed changes in the high elevations ( $> 5000$  m asl) and remote regions (such as the TP and Himalayas region) (Lau et al., 2010; Ramanathan and Carmichael, 2008). It was reported that the pre-monsoon season is  
15 the major vegetation-fire period in the foothill areas of the southern Himalaya (Vadrevu et al., 2012; Putero et al., 2014), and the winds surrounding the Himalayas and TP could facilitate the transport of carbonaceous matter from South Asia to the Himalayas (Cong et al., 2015a; Dong et al., 2017b; Lau et al., 2010).

We analyzed the CAM5 model results to quantify the source attributions of BC in  
20 the Mt. Yulong area. BC emissions from each of the four source regions in the surrounding area (i.e., South Asia, East Asia, Middle East and Southeast Asia) are explicitly tracked. Figure 14 shows the annual and seasonal mean relative contributions from the tagged source regions. The two sampling sites locate in the same model grid box, as marked in the figure. The modeled near-surface BC is  
25 predominately (more than 90%) from South Asia and East Asia. East Asia has a dominant contribution in the monsoon, post-monsoon and winter seasons, while South Asia dominates in the pre-monsoon season (Table 3). As discussed by Wang et al.  
(2015), circulation patterns during the monsoon and non-monsoon seasons largely determine the seasonal variations in the transport of aerosols from the different major  
30 sources to the southeastern TP. Strong precipitation during the monsoon season can

substantially remove atmospheric BC during the transport, especially, from South Asia. Although smoke plumes can sometimes be lifted over the natural block of the Himalayas, they have relatively less important contribution to the surface than to the upper-level BC concentrations. According to our climate model results, emissions 5 (2010-2014) from East Asia, including local sources accounted in the emission dataset, have a dominant contribution to the near-surface BC at the Mt. Yulong sites during the monsoon (78%) and post-monsoon (68%) seasons, as well as the winter season (53%). The seasonal changes in source apportionment also have an implication on the cause of variations in OC/BC ratios over the southeastern TP (e.g., Wang et al., 2015).

#### 10 4. Conclusions and remarks

Carbonaceous aerosols from the Mt. Yulong region and GHZ basin were measured to investigate the small-scale spatio-temporal variations and light-absorbing properties. Results of first two years of continuous observations show that the annual mean EC 15 and OC concentrations in aerosol were  $1.51 \pm 0.93$  and  $2.57 \pm 1.32 \mu\text{g m}^{-3}$ , respectively.

Concentrations of carbonaceous matter displayed distinct seasonal differences, with the lowest content found in monsoon season and the highest concentration in winter season. Monthly mean EC concentrations in aerosols from monsoon to post-monsoon season were higher than OC, a large impact from biomass burning emissions in the 20 high atmosphere (approximately 5000 m asl) of Mt. Yulong. The seasonal differences of carbonaceous matter found in GHZ basin were different with that in Mt. Yulong, distinct elevation difference and different degrees of human (or tourism related) activities between the two sites were the main reasons account for the discrepancy. Furthermore, high carbonaceous matter associated with OC in GHZ basin was mainly 25 contributed from vehicle emissions. Therefore, there was a discernable spatial difference in the concentrations of carbonaceous matter in this glacierized region. Moreover, inter-annual differences of carbonaceous aerosols in Mt. Yulong and GHZ were also distinct. The annual mean concentrations of carbonaceous matter in 2016 30 were lower than those in 2015, partly indicating the improvement of local air quality in the Mt. Yulong region.

The annual mean OC/**EC** ratio was  $2.45 \pm 1.96$  in Mt. Yulong, with the highest value in monsoon season ( $4.4 \pm 2.0$ ) and the lowest in winter ( $0.79 \pm 0.4$ ). Strong photochemical reactions and local tourism activities in monsoon season were likely the main factors resulting in relatively high OC/**EC** ratios in the Mt. Yulong region,  
5 particularly in GHZ basin.

**EC** MAE was quantified using a thermal-optical carbon analyzer, and was measured at 632 nm under the quartz filter-based method. The corrected mean **EC** MAE at 632 nm was  $6.82 \pm 0.73 \text{ m}^2 \text{ g}^{-1}$  in the Mt. Yulong region, comparable with the results in other studies. The strong correlations were found between EC MAE and  
10 POC/OC, OC/**EC**, and EC<sub>s</sub> in aerosol. Obvious seasonal variations and discernable spatial difference of **EC** MAE in the study area were largely related to the OC abundance. The enhancement of MAE was mainly due to external-coating of OC and/or mixing state (internally mixed).

To quantitatively estimate the source apportionment of **EC** (or BC) in the Mt.  
15 Yulong area, we used a global aerosol-climate model, in which BC emissions from four regions (i.e., South Asia, East Asia, Middle East and Southeast Asia) are explicitly tracked. The five-year (2010-2014) mean results show that East Asia has the largest contribution (52%) to the annual mean near-surface BC concentration in this area, followed by South Asia (43%). There is a quite strong seasonal variation in the  
20 source apportionment. East Asia has a dominant contribution in BC emission in the monsoon and post-monsoon seasons, while South Asia dominates in the pre-monsoon season.

**The authors declare that they have no conflict of interest.**

25

### Acknowledgements

This work was supported by the National Natural Science Foundation of China (41601071, 41721091, 41630754), and Key Research Program for Frontier Science of Chinese Academy of Sciences (QYZDY-SSW-DQC021, QYZDJ-SSW-DQC039).  
30 The independent program of SKLCS (SKLCS-ZZ-2017) and the "Light of West

China" Program (Y62992), and China Postdoctoral Science Foundation (2016T90963, 2015M582725) and Youth Talent Program of CAREERI CAS (Y551C11001). R. Zhang acknowledges support from NSFC (41605041), Jiangsu Provincial Science Fund (BK20160621), Fundamental Research Funds for the Central Universities (020714380020) and International Postdoctoral Exchange Fellowship (20160046). H. Wang and Y. Qian acknowledge support from the U.S. Department of Energy (DOE), Office of Science, Biological and Environmental Research as part of the Earth System Modeling program. The Pacific Northwest National Laboratory (PNNL) is operated for DOE by Battelle Memorial Institute under contract DE-AC05-76RLO1830. The model simulations were performed using PNNL Institutional Computing resources.

## References

- Andreae, M. O.: Climatic effects of changing atmospheric aerosol levels, in: World Survey of Climatology, Future Climates of the World, edited by: Henderson-Sellers, A., Elsevier, Amsterdam, Vol. 16, pp. 341-392, 1995.
- Andreae, M. O. and Gelencsér, A.: Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols, *Atmos. Chem. Phys.*, 6, 3131-3148, doi:10.5194/acp-6-3131-2006, 2006.
- Antony, R., Mahalinganathan, K., Thamban, M.: Organic carbon in Antarctic snow: spatial trends and possible sources, *Environ. Sci. Technol.*, 45, 9944-9950, 2011.
- Bergstrom, R. W., Ackerman, T. P., and Richards, L. W.: Optical properties of particulate elemental carbon. Particulate carbon: atmospheric life cycle, edited by: Wolff, G. T. and Klimisch, R. L., Plenum Press, New York, USA, 43-51, 1982.
- Bonasoni, P., Laj, P., Marinoni, A., Sprenger, M., Angelini, F., Arduini, J., Bonafe, U., Calzolari, F., Colombo, T., Decesari, S., Di Biagio, C., Di Sarra, A., Evangelisti, F., Duchi, R., Facchini, M., Fuzzi, S., Gobbi, G., Maione, M., Panday, A., Roccato, F., Sellegri, K., Venzac, H., Verza, G., Villani, P., Vuillermoz, E., Cristofanelli, P.: Atmospheric brown clouds in the Himalayas: first two years of continuous observations at the Nepal Climate Observatory-Pyramid (5079 m), *Atmos. Chem.*

Phys. 10, 75150-7531, 2010.

Bond, T. C., Streets, D. G., Yarber, K. F.: A technology-based global inventory of black and organic carbon emissions from combustion, J. Geophys. Res.-Atmos., 109(14), D14203. DOI:10.1029/2003JD003697, 2004.

5 Bond, T. C., and Bergstrom, R. W.: Light absorption by carbonaceous particles: an investigative review, Aerosol Sci. Technol. 40:27-67, 2006a.

Bond, T. C., Habib, G., and Bergstrom, R. W.: Limitations in the enhancement of visible light absorption due to mixing state, J. Geophys. Res., 111, D20211, doi:10.1029/2006JD007315, 2006b.

10 Bond, T. C., Bhardwaj, E., Dong, R., Jogani, R., Jung, S. K. Roden, C., Streets, D. G. and Trautmann, N. M.: Historical emissions of black and organic carbon aerosol from energy-related combustion, 1850-2000, Global Biogeochem. Cycles, 21(2), doi:10.1029/2006GB002840, 2007.

Bond, T. C., et al.: Bounding the role of black carbon in the climate system: A  
15 scientific assessment, J. Geophys. Res.-Atmos., 118, 5380-5552,  
doi:10.1002/jgrd.50171, 2013.

Borrego, C., Tchepel, O., Barros, N., Miranda, A.: Impact of road traffic emissions on air quality of the Lisbon region, Atmos. Environ., 34, 4683-4690, 2000.

20 Bou, K. D., Flamant, C., Cuesta, J.: Dust emission and transport associated with a Saharan depression: February 2007 case, J. Geophys. Res., 115, D0\_0H27.  
<http://dx.doi.org/10.1029/2009JD012390>, 2010.

Brimblecombe, P.: The big smoke: a history of air pollution in London since medieval times, London, New York, Methuen, 185 p., 1987.

Cain, J. P., Gassman, P. L., Wang, H., Laskin, A.: Micro-FTIR study of soot chemical  
25 composition-evidence of aliphatic hydrocarbons on nascent soot surfaces, Phys Chem. Chem. Phys., 12(20), 5206-5218, 2010.

Cao, J. J., Tie, X., Xu, B. Q., Zhao, Z., Zhu, C., Li G., Liu S.: Measuring and modeling black carbon (BC) contamination in the SE Tibetan Plateau, J. Atmos. Chem., 67, 45-60, 2010.

30 Cao, J. J., Zhu, C. S., Tie, X. X., Geng, F. H., Xu, H. M., Ho, S. S. H., Wang, G. H.,

- Han, Y. M., Ho, K.F.: Characteristics and sources of carbonaceous aerosols from Shanghai, China, *Atmos. Chem. Phys.*, 13, 803-817, 10.5194/acp-13-803-2013, 2013.
- Cappa, C. D. et al.: Radiative absorption enhancements due to the mixing state of atmospheric black carbon, *Science*, 31, 337, 1078-1081, 2012.
- Carrico, C. J., Bergin, M. H., Shrestha, A. B.: The importance of carbon and mineral dust to seasonal aerosol properties in the Nepal Himalaya, *Atmos. Environ.*, 37, 2811-2824, 2003.
- Carslaw, K. S., Boucher O., Spracklen D. V., Mann G. W., Rae J. G. L., Woodward S.,  
10 Kulmala M.: A review of natural aerosol interactions and feedbacks within the Earth system, *Atmos. Chem. Phys.*, 10, 1701-1737, 2010.
- Chen, P. F, Kang, S. C, Li, C. L., Rupakheti, M., et al.: Characteristics and sources of polycyclic aromatic hydrocarbons in atmospheric aerosols in the Kathmandu Valley, Nepal, *Sci. of the Tot. Environ.*, 538, 86-92, 2015.
- 15 Cheng, Y., He, K. B., Zheng, M., Duan, F. K., Du, Z. Y., Ma, Y. L., Tan, J. H.: Mass absorption efficiency of elemental carbon and water-soluble organic carbon in Beijing, China, *Atmos. Chem. Phys.*, 11, 11497-11510, 2011a.
- Cheng, Y., He, K. B., Duan, F. K., Zheng, M., Du, Z. Y., Ma, Y. L., Tan, J. H.:  
Ambient organic carbon to elemental carbon ratios: Influences of the measurement  
20 methods and implications, *Atmos. Environ.*, 45, 2060-2066, 2011b.
- Chow, J. C., Watson, J. G., Pritchett, L. C., Pierson, W. R., Frazier, C. A., and Purcell,  
R. G.: The DRI Thermal/Optical Reflectance carbon analysis system: description,  
evaluation and applications in U.S. air quality studies, *Atmos. Environ.*, 27A,  
1185-1201, 1993.
- 25 Chow, J. C, Watson, J. G., Crow, D.: Comparison of IMPROVE and NIOSH carbon measurements, *Aerosol Sci. Technol.*, 34, 23-34, 2001.
- Chow, J. C., Watson, J. G., Chen, L. W. A., et al.: Equivalence of elemental carbon by thermal/optical reflectance and transmittance with different temperature protocols,  
*Environ. Sci. Technol.* 38, 4414-4422, 2004.
- 30 Chung, S. H., Seinfeld, J. H.: Climate response of direct radiative forcing of

anthropogenic black carbon, J. Geophys. Res., 110, D11102, doi:10.1029/2004JD005441, 2005.

5 Cong, Z. Y., Kawamura, K., Kang, S. C., et al.: Penetration of biomass-burning emissions from South Asia through the Himalayas: new insights from atmospheric organic acids, *Sci. Reports*, 5, 9580, Doi:10.1038/srep09580, 2015a.

Cong, Z. Y., Kang, S. C., Kawamura, K.: Carbonaceous aerosols on the south edge of the Tibetan Plateau: concentrations, seasonality and sources, *Atmos. Chem. Phys.*, 15, 1573-1584, 2015b.

10 Daly, H. M., Horn, A. B.: Heterogeneous chemistry of toluene, kerosene and diesel soots, *Phys. Chem.*, 11(7), 1069-1076, 2009.

Ding, A. J., Huang, X., Nie, W., Sun, J. N., Kerminen, V., Petäjä T., Su, H., Cheng, Y. F., Yang, X. Q., Wang, M. H., Chi, X. G., Wang, J. P., Virkkula, A., Guo, W. D., Yuan, J., Wang, S. Y., Zhang, R. J., Wu, Y. F., Song, Y., Zhu, T., Zilitinkevich, S., Kulmala, M., and Fu, C. B.: Enhanced haze pollution by black carbon in megacities 15 in China, *Geophys. Res. Lett.*, 43, 2873-2879, 2016.

Doherty, S. J., Grenfell, T. C., Forsström, S.: Observed vertical redistribution of black carbon and other insoluble light-absorbing particles in melting snow, *J. Geophys Res-Atmos.*, 118, 5553-5569, 2013.

20 Dong, Z. W., Li, Z. Q., Ross, E., Wu, L. H., Zhou, P.: Temporal characteristics of mineral dust particles in precipitation of Urumqi River Valley in Tian Shan, China: a comparison of alpine site and rural site, *Atmos. Res.*, 101, 294-306, 2011.

Dong, Z. W., Kang, S. C., Qin, D. H., Qin, X., Yan, F. P., Du, W. T., Wei, T.: Temporal and diurnal analysis of trace elements in the Cryospheric water at remote Laohugou basin in northeast Tibetan Plateau, *Chemosphere*, 171, 386-398, 2017a.

25 Dong, Z. W., Kang, S. C., Guo, J. M., Wang, X. J., Qin, D. H.: Composition and mixing states of brown haze particle over the Himalayas along two transboundary south-north transects. *Atmos. Environ.*, 156, 24-35, 2017b.

Doran, J. C., Barnard, J. C., Arnott, W. P., Cary, R., Coulter, R., Fast., J. D., Kassianov, E. I., Kleinman, L., Laulainen, N. S., Martin, T., Paredes-Miranda, G., Pekour, M. 30 S., Shaw, W. J., Smith, D. F., Springston, S. R., and Yu, X. Y.: The T1-T2 study:

evolution of aerosol properties downwind of Mexico city, *Atmos. Chem. Phys.*, 7, 1585-1598, 2007.

Flanner, M. G. and Zender, C. S.: Linking snowpack microphysics and albedo evolution, *J. Geophys. Res.-Atmos.*, 111, D12208, doi:10.1029/2005jd006834, 5 2006.

Flanner, M. G., Zender, C. S., Hess, P. G., Mahowald, N. M., Painter, T. H., Ramanathan, V., Rasch P. J.: Springtime warming and reduced snow cover from carbonaceous particles, *Atmos. Chem. Phys.* 9, 2481-2497, doi:10.5194/acp-9-2481- 2009, 2009.

10 Fuller, K. A., Malm, W. C., and Kreidenweis, S. M.: Effects of mixing on extinction by carbonaceous particles, *J. Geophys. Res.*, 104, 15941-15954, 1999.

Gertler, C. G., Puppala, S. P., Panday, A., Stumm, D., Shea, J.: Black carbon and the Himalayan cryosphere: a review. *Atmos. Environ.* 125, 404-417, 2016.

15 Han, C., Liu, Y. C., Ma, J. Z., He, H.: Key role of organic carbon in the sunlight-enhanced atmospheric aging of soot by O<sub>2</sub>, *Proc. Natl. Acad. Sci. USA*, 109 (52), 21250-21255, 2012.

Hansen, J., and Nazarenko, L.: Soot climate forcing via snow and ice albedos, *Proc. Natl. Acad. Sci. USA*, 101(2), 423-428, doi:10.1073/pnas.2237157100, 2004.

20 Hansen, J., Sato, M., Ruedy, R., et al.: Efficacy of climate forcings, *J. Geophys. Res.*, 110, D18104, doi:10.1029/2005JD005776, 2005.

Hecobian, A., Zhang, X., Zheng, M., Frank, N., Edgerton, E. S., Weber, R. J.: Water-soluble organic aerosol material and the light-absorption characteristics of aqueous extracts measured over the Southeastern United States, *Atmos. Chem. Phys.*, 10, 5965-5977, 2010.

25 Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J. I., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.: Historical (1750-2014) anthropogenic emissions of reactive gases and aerosols from the Community Emission Data System (CEDS), *Geosci. Model Dev.*, 11, 369-408, 2018.

- IPCC, 2013.: Summary for Policymakers. In: Climate Change 2013: The Physical  
Science Basis. Contribution of Working Group I to the Fifth Assessment Report of  
the Intergovernmental Panel on Climate Change, (Stocker, T. F., Qin, D., Plattner,  
G. K., et al.: Cambridge University Press, Cambridge, United Kingdom and New  
5 York: NY, USA. 129-234, 2013.)
- Jacobson, M. Z.: Strong radiative heating due to the mixing state of black carbon in  
atmospheric aerosols, *Nature*, 409, 695-697, 2001.
- Jacobson, M. Z.: Climate response of fossil fuel and biofuel soot, accounting for  
soot's feedback to snow and sea ice albedo and emissivity, *J. Geophys. Res.*, 109,  
10 D21201, doi:10.1029/2004JD004945, 2004.
- Jeong, C. H., Hopke, P. K., Kim, E., and Lee, D. W.: The comparison between  
thermal-optical transmittance elemental carbon and Aethalometer black carbon  
measured at multiple monitoring sites, *Atmos. Environ.*, 38, 5193-5204, 2004.
- Jerret, M., Burnett, R. T., Ma, R., Pope, C. A., Krewski, D., Newbold, K. B., Thurston,  
15 G., Shi, Y., Finkelstein, N., Calle, E. E., and Thun, M. J.: Spatial analysis of air  
pollution and mortality in Los Angeles, *Epidemiology*, 16, 727-736, 2005.
- Kang, S. C., Mayewski, P. A., Qin, D. H., Snead, S. R., Ren, J. W. and Zhang, D. Q.:  
Seasonal differences in snow chemistry from the vicinity of Mt. Everest, central  
Himalayas, *Atmos. Environ.*, 38, 2819-2829, 2004.
- 20 Kang, S. C., Zhang, Q. Q., Kaspari, S. S., Qin, D. H., Cong, Z. Y., Ren, J. W. and  
Mayewski, P. A.: Spatial and seasonal variations of elements composition in Mt.  
Everest (Qomolangma) snow/firn, *Atmos. Environ.*, 41, 7208-7218, 2007.
- Kaspari, S. D., Painter, T. H., Gysel, M., Skiles, S. M., Schwikowski, M.: Seasonal  
and elevational variations of black carbon and dust in snow and ice in the  
25 Solu-Khumbu, Nepal and estimated radiative forcings, *Atmos. Chem. Phys.*, 14,  
8089-8103, 2014.
- Kirchner, U., Scheer, V., Vogt, R.: FTIR spectroscopic investigation of the mechanism  
and kinetics of the heterogeneous reactions of NO<sub>2</sub> and HNO<sub>3</sub> with soot, *J. Phys.  
Chem. A*, 104, 8908-8915, 2000.
- 30 Knox, A., Evans, G. J., Brook, J. R., Yao, X., Jeong, C. H., Godri, K. J., Sabaliauskas,

- K., and Slowik, J. G.: Mass absorption cross-section of ambient black carbon aerosol in relation to chemical age, *Aerosol Sci. Technol.*, 43, 522-532, 2009.
- Lau, W. K. M., Kim, M. K., Lee, W. S.: Enhanced surface warming and accelerated snow melt in the Himalayas and Tibetan Plateau induced by absorbing aerosols, 5 *Environ. Res. Lett.*, 5, 025204, doi:10.1088/1748-9326/5/2/025204, 2010.
- Li, C. L., Yan, F. P., Kang, S. C., et al.: Light absorption characteristics of carbonaceous aerosols in two remote stations of the southern fringe of the Tibetan Plateau, *China Atmos. Environ.*, 143, 79-85, 2016a.
- Li, C. L. et al.: Sources of black carbon to the Himalayan-Tibetan Plateau glaciers, 10 *Nat. Commun.*, 7, 12574, doi: 10.1038/ncomms12574, 2016b.
- Li, C. L., Chen, P. F., Kang, S. C., Yan, F. P., Hu, Z. F., Qu, B.: Concentrations and light absorption characteristics of carbonaceous aerosol in PM<sub>2.5</sub> and PM<sub>10</sub> of Lhasa city, the Tibetan Plateau, *Atmos. Environ.*, 127, 340-346, 2016c.
- Lioussse, C., Cachier, H., and Jennings, S. G.: Optical and thermal measurements of 15 black carbon aerosol content in different environments: variation of the specific attenuation cross-section, sigma ( $\sigma$ ), *Atmos. Environ.*, 27A, 1203-1211, 1993.
- Lioussse, C., Penner, J. E., Chuang, C.: A global three-dimensional study of carbonaceous aerosols, *J. Geophys. Res-Atmos.*, 101(D14), 19411-19432, 1996.**
- Liu, X., Ma, P.-L., Wang, H., Tilmes, S., Singh, B., Easter, R. C., Ghan, S. J., and 20 Rasch, P. J.: Description and evaluation of a new four-mode version of the Modal Aerosol Module (MAM4) within version 5.3 of the Community Atmosphere Model, *Geosci. Model Dev.*, 9, 505-522, <https://doi.org/10.5194/gmd-9-505-2016>, 2016.
- Liu, J. B., Röhland, K. M., Chen, J. H., Xu, Y. Y., Chen, S. Q., Chen, Q. M., Huang, W., Xu, Q. H., Chen, F. H., Smol, J. P.: Aerosol-weakened summer monsoons 25 decrease lake fertilization on the Chinese Loess Plateau, *Nature Climate Change*, DOI:10.1038/NCLIMATE3220, 2017.
- Lohmann, U. and Feichter, J.: Global indirect aerosol effects: a review, *Atmos. Chem. Phys.*, 5, 715-737, 2005.
- Lüthi, Z. L., Skerkak, B., Kim, S. W., et al.: Atmospheric brown clouds reach the 30 Tibetan Plateau by crossing the Himalayas, *Atmos. Chem. Phys.*, 14, 28105-28146,

2014.

- Ma, P. L., Rasch, P. J., Wang, H., Zhang, K., Easter, R. C., Tilmes, S., Fast, J. D., Liu, X., Yoon, J. H., and Lamarque, J. F.: The role of circulation features on black carbon transport into the Arctic in the Community Atmosphere Model version 5 (CAM5), *J. Geophys. Res.-Atmos.*, 118, 4657-4669, doi:10.1002/jgrd.50411, 2013.
- Mandal, P., Saud, T., Sarkar, R., Mandal, A., Sharma, S. K., Mandal, T. K., and Bassin, J. K.: High seasonal variation of atmospheric C and particle concentrations in Delhi, India, *Environ. Chemi. Lett.*, 12, 225-230, 10.1007/s10311-013-0438-y, 2014.
- Ming, J., Wang, P. L., Zhao, S. Y., Chen, P. F.: Disturbance of light-absorbing aerosols on the albedo in a winter snowpack of Central Tibet, *J. of Environ. Sci.*, 25(8), 1601-1607, 2013.
- Nie, J., Garzione, C., Su, Q., Liu, Q., Zhang, R., Heslop, D., Necula, C., Zhang, S., Song, Y., Luo, Z.: Dominant 100,000-year precipitation cyclicity in a late Miocene lake from northeast Tibet, *Sci. Adv.* 3, e1600762. 2017.
- Nieto-Gligorovski, L., et al.: Interactions of ozone with organic surface films in the presence of simulated sunlight: Impact on wettability of aerosols, *Phys. Chem. Chem. Phys.*, 10(20), 2964-2971, 2008.
- Niu, H. W., He, Y. Q., Zhu, G. F., Xin, H. J., Du, J. K., Pu, T., Lu, X. X.: Environmental implications of the snow chemistry from Mt Yulong, southeastern Tibetan Plateau, *Quaternary International*, 313-314, 168-178, 2013.
- Niu, H. W., He, Y. Q., Lu, X. X., Shen, J.: Chemical composition of rainwater in the Yulong Snow Mountain region, Southwestern China, *Atmos. Res.*, 144, 195-206, 2014.
- Niu, H. W., He, Y. Q., Kang, S. C., Lu, X. X.: Chemical compositions of snow from Mt. Yulong, southeastern Tibetan Plateau, *J. Earth Syst. Sci.*, 125(2), 403-416, 2016.
- Niu, H. W., Kang, S. C., Shi, X. F.: In-situ measurements of light-absorbing impurities in snow of glacier on Mt. Yulong and implications for radiative forcing estimates, *Sci. of the Tot. Environ.*, 581-582, 848-856, 2017a.
- Niu, H. W., Kang, S. C., Zhang, Y. L., Shi, X. F.: Distribution of light-absorbing

- impurities in snow of glacier on Mt. Yulong, southeastern Tibetan Plateau, Atmos. Res., 197, 474-484, 2017b.
- Omar, A. H., Biegalski, S., Larson, S. M., Landsberger, S.: Particulate contributions to light extinction and local forcing at a rural Illinois site, *Atmos. Environ.*, 33(17), 2637-2646, 1999.
- Pachauri, T., Singla, V., Satsangi, A.: Characterization of carbonaceous aerosols with special reference to episodic events at Agra, India, *Atmos. Res.*, 128, 98-110, 2013.
- Park, R. J., Jacob, D. J., Chin, M., Martin, R. V.: Sources of carbonaceous aerosols over the United States and implications for natural visibility, *J. Geophys. Res.*, 108(D12), 4355, doi:10.1029/2002JD003190, 2003.
- Putero, D. et al.: Influence of open vegetation fires on black carbon and ozone variability in the southern Himalayas (NCO-P, 5079 m a.s.l.), *Environ. Pollut.*, 184, 597-604, 2014.
- Qian, Y., Flanner, M. G., Leung, L. R., Wang, W.: Sensitivity studies on the impacts of Tibetan Plateau snowpack pollution on the Asian hydrological cycle and monsoon climate, *Atmos. Chem. Phys.*, 11, 1929-1948, 2011.
- Qian, Y., Yasunari, T., Doherty, S. J.: Light-absorbing particles in snow and ice: Measurement and modeling of climatic and hydrological impact, *Adv. Atmos. Sci.*, 32, 64-91, 2015.
- Qu, B., Ming, J., Kang, S. C., Zhang, G. S., Li, Y. W., Li, C. D., Zhao, S. Y., Ji, Z. M., Cao, J. J.: The decreasing albedo of the Zhadang glacier on western Nyainqntanglha and the role of light-absorbing impurities, *Atmos. Chem. Phys.*, 14, 11117-11128, 2014.
- Ram, K. and Sarin, M. M.: Absorption coefficient and site-specific mass absorption efficiency of elemental carbon in aerosols over urban, rural, and high-altitude sites in India, *Environ. Sci. Technol.*, 43, 8233-8239, 2009.
- Ram, K., Sarin, M. M., and Tripathi, S. N.: Temporal trends in atmospheric PM<sub>2.5</sub>, PM<sub>10</sub>, elemental carbon, organic carbon, water-soluble organic carbon, and optical properties: impact of biomass burning emissions in the Indo-Gangetic Plain, *Environ. Sci. Technol.*, 46, 686-695, 10.1021/es202857w, 2012.

- Ramanathan, V., Chung, C., Kim, D., Bettge, T., Buja, L., Kiehl, J. T., Washington, W. M., Fu, Q., Sikka, D. R., and Wild, M.: Atmospheric brown clouds: Impacts on South Asian climate and hydrological cycle, *Proc. Natl. Acad. Sci. USA*, 102, 5326-5333, 2005.
- 5 Ramanathan, V., Li, F., Ramana, M. V., Praveen, P. S., Kim, D., Corrigan, C. E., Nguyen, H., Stone, E. A., Schauer, J. J., Carmichael, G. R., Adhikary, B., Yoon, S. C.: Atmospheric brown clouds: hemispherical and regional variations in long-range transport, absorption, and radiative forcing, *J. Geophys. Res. Atmos.*, 112, 2007.
- Ramanathan, V., Carmichael, G.: Global and regional climate changes due to black carbon, *Nature Geoscience*, 1(4), 221-227, doi: 10.1038/ngeo156, 2008.
- 10 Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, R., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G-K, Bloom, S., Chen, J., Collins, D., Conaty, A., da Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick, A. G., Sienkiewicz, M., and Woollen, J.: MERRA: NASA's modern-era retrospective analysis for research and applications, *J. Climate*, 24, 3624-3648, 2011.
- 15 Sandradewi, J., Prévot, A. S. H., Weingartner, E., Schmidhauser, R., Gysel, M., and Baltensperger, U.: A study of wood burning and traffic aerosols in an Alpine valley using a multi-wavelength Aethalometer, *Atmos. Environ.*, 42, 101-112, 2008.
- Schnaiter, M., Linke, C., Mohler, O., Naumann, K. H., Saathoff, H., Wagner, R., Schurath, U., and Wehner, B.: Absorption amplification of black carbon internally mixed with secondary organic aerosol, *J. Geophys. Res.*, 110, D19204, 2005.
- 20 Schneidemesser, E. V., Schauer, J. J., Hagler, G. S. W.: Concentrations and sources of carbonaceous aerosol in the atmosphere of Summit, Greenland, *Atmos. Environ.*, 43, 4155-4162, 2009.
- Schwarz, J. P., Gao, R. S., Perring, A. E., Spackman, J. R., and Fahey, D. W.: Black carbon aerosol size in snow, *Sci. Reports*, 3, 1356. doi:10.1038/srep01356, 2013.
- 25 Schuckmann, K. V., Palmer, M. D., Trenberth, K. E.: An imperative to monitor Earth's energy imbalance, *Nature Climate Change*, 6, DOI: 10.1038/NCLIMATE2876,

2016.

- Sharma, S., Brook, J. R., Cachier, H., Chow, J., Gaudenzi, A., and Lu, G.: Light absorption and thermal measurements of black carbon in different regions of Canada, *J. Geophys. Res.*, 107, 4771, doi:10.1029/2002JD002496, 2002.
- 5 Shi, X. F., Niu, H. W., He, Y. Q.: Characteristics of rainwater chemistry in Lijiang-Yulong Snow Mountain, *Environmental Chemistry* 36(5), 994-1002, 2017. (in Chinese with English abstract).
- Shrestha, A. B., Wake, C. P., Dibb, J. E., Mayewski, P., Whitlow, S.: Seasonal variations in aerosol concentrations and compositions in the Nepal Himalaya, 10 *Atmos. Environ.* 34, 3349-3363, 2000.
- Vadrevu, K. P. et al.: Vegetation fires in the Himalayan region - Aerosol load, black carbon emissions and smoke plume heights, *Atmos. Environ.*, 47, 241-251, 2012.
- van Marle, M. J. E., Kloster, S., Magi, B. I., Marlon, J. R., Daniau, A. L., Field, R. D., Arneth, A., Forrest, M., Hantson, S., Kehrwald, N. M., Knorr, W., Lasslop, G., Li, 15 F., Mangeon, S., Yue, C., Kaiser, J. W., and van der Werf, G. R.: Historic global biomass burning emissions based on merging satellite observations with proxies and fire models (1750-2015), *Geosci. Model Dev.*, 10, 3329-3357, 2017.
- Wan, X., Kang, S. C., Li, Q. L., Rupakhetiet, D., Zhang, Q. G., Guo, J. M., Chen, P. F.: Organic molecular tracers in the atmospheric aerosols from Lumbini, Nepal, in 20 the northern Indo-Gangetic Plain: Influence of biomass burning. *Atmos. Chem. Phys.*, 17, 8867-8885, 2017.
- Wang, H., Easter, R. C., Rasch, P. J., Wang, M., Liu, X., Ghan, S. J., Qian, Y., Yoon, J.-H., Ma, P.-L., and Vinoj, V.: Sensitivity of remote aerosol distributions to representation of cloud-aerosol interactions in a global climate model, *Geosci. Model Dev.*, 6, 765-782, doi:10.5194/gmd-6-765-2013, 2013.
- 25 Wang, H., Rasch, P. J., Easter, R. C., Singh, B., Zhang, R., Ma, P. L., Qian, Y., Ghan, S. J., and Beagley, N.: Using an explicit emission tagging method in global modeling of source-receptor relationships for black carbon in the Arctic: Variations, sources, and transport pathways, *J. Geophys. Res.-Atmos.*, 119, 12, 888-12,909, doi:10.1002/ 2014JD022297, 2014.

- Wang, M., Xu, B. Q., Cao, J. J., et al.: Carbonaceous aerosols recorded in a southeastern Tibetan glacier: analysis of temporal variations and model estimates of sources and radiative forcing, *Atmos. Chem. Phys.*, 15, 1191-1204, 2015.
- Weingartner, E., Saathoff, H., Schnaiter, M., Streit, N., Bitnar, B., and Baltensperger,  
5 U.: Absorption of light by soot particles: determination of the absorption coefficient  
by means of Aethalometers, *J. Aerosol Sci.*, 34, 1445-1463, 2003.
- Xu, B. Q., and Coauthors: Black soot and the survival of Tibetan glaciers, *Proc. Natl.  
Acad. Sci. USA*, 106(52), 22 114-22 118, doi: 10.1073/pnas.0910444106, 2009a.
- Xu, B. Q., Wang, M., Joswiak, D. R., Cao, J. J., Yao, T. D., Wu, G. J., Yang, W., Zhao,  
10 H. B.: Deposition of anthropogenic aerosols in a southeastern Tibetan glacier, *J.  
Geophys. Res.*, 114, D17209, doi:10.1029/2008JD011510, 2009b.
- Yang, M., Howell, S. G., Zhuang, J. and Huebert, B. J.: Attribution of aerosol light  
absorption to black carbon, brown carbon, and dust in China - interpretations of  
atmospheric measurements during EAST-AIRE, *Atmos. Chem. Phys.*, 9,  
15 2035-2050, doi:10.5194/acp-9-2035-2009, 2009.
- Yu, S. C., Saxena, V. K., and Zhao, Z.: A comparison of signals of regional  
aerosol-induced forcing in eastern China and the southeastern United States,  
*Geophys. Res. Lett.*, 28, 713-716, 2001.
- Zhang, R., Wang, H., Qian, Y., Rasch, P. J., Easter, R. C., Ma, P.-L., Singh, B., Huang,  
20 J., and Fu, Q.: Quantifying sources, transport, deposition, and radiative forcing of  
black carbon over the Himalayas and Tibetan Plateau, *Atmos. Chem. Phys.*, 15,  
6205-6223, <https://doi.org/10.5194/acp-15-6205-2015>, 2015.
- Zhang, T., Claeys, M., Cachier, H., Dong, S. P., Wang, W., Maenhaut, W., and Liu, X.  
D.: Identification and estimation of the biomass burning contribution to Beijing  
25 aerosol using levoglucosan as a molecular marker, *Atmos. Environ.*, 42, 7013-7021,  
2008.
- Zhao, Z., Cao, J., Shen, Z., Xu, B., Zhu, C., Chen, L. W. A., Su, X., Liu, S., Han, Y.,  
and Wang, G.: Aerosol particles at a high-altitude site on the Southeast Tibetan  
Plateau, China: implications for pollution transport from South Asia, *J. Geophys.  
30 Res.-Atmos.*, 118, 11360-11375, 2013.

Table 1

Statistical results of EC, OC, and POC concentrations ( $\mu\text{g m}^{-3}$ ) and OC/EC ratios in aerosol from Mt. Yulong during Dec. 2014-Dec. 2016.

	Annual (n=120)		Winter (n=36)		Pre-monsoon (n=34)		Monsoon (n=33)		Post-monsoon (n=17)	
	Mean $\pm$ SD	Range	Mean $\pm$ SD	Range	Mean $\pm$ SD	Range	Mean $\pm$ SD	Range	Mean $\pm$ SD	Range
OC	1.65 $\pm$ 1.14	0.07-5.96	1.75 $\pm$ 0.8	0.51-3.66	1.37 $\pm$ 1.2	0.07-5.73	1.34 $\pm$ 0.9	0.24-4.6	2.57 $\pm$ 1.3	0.99-5.96
EC	1.53 $\pm$ 1.49	0.02-6.83	1.81 $\pm$ 1.5	0.02-6.83	0.55 $\pm$ 0.4	0.03-1.77	1.25 $\pm$ 1.2	0.04-3.9	3.51 $\pm$ 1.2	1.22-5.8
OC/EC	2.06 $\pm$ 3.38	0.35-20.9	1.45 $\pm$ 1.1	0.38-5.55	3.67 $\pm$ 5.7	0.7-22.9	1.85 $\pm$ 1.8	0.42-8.3	0.79 $\pm$ 0.4	0.35-2.27
POC	0.53 $\pm$ 0.53	0.009-2.4	0.63 $\pm$ 0.5	0.008-2.39	0.19 $\pm$ 0.1	0.01-0.62	0.43 $\pm$ 0.4	0.01-1.3	1.24 $\pm$ 0.4	0.42-2.04

Table 2

Statistical results of EC, OC, and POC concentrations ( $\mu\text{g m}^{-3}$ ) and OC/EC ratios in aerosol from GHZ during Dec. 2014-Dec. 2016.

	Annual (n=116)		Winter (n=27)		Pre-monsoon (n=47)		Monsoon (n=28)		Post-monsoon (n=12)	
	Mean $\pm$ SD	Range	Mean $\pm$ SD	Range	Mean $\pm$ SD	Range	Mean $\pm$ SD	Range	Mean $\pm$ SD	Range
OC	3.5 $\pm$ 1.5	1.1-10	4.09 $\pm$ 1.7	2.3-9.9	3.7 $\pm$ 1.7	1.3-10	2.8 $\pm$ 0.8	1.1-4.7	3.1 $\pm$ 1.1	2.0-5.6
EC	1.5 $\pm$ 0.9	0.15-4.5	2.42 $\pm$ 0.74	1.1-4.2	1.4 $\pm$ 0.7	0.28-4.5	0.75 $\pm$ 0.3	0.15-1.5	1.58 $\pm$ 0.9	0.63-3.2
OC/EC	2.9 $\pm$ 1.8	0.87-8.8	1.93 $\pm$ 1.58	0.87-8.9	2.72 $\pm$ 1.3	1.1-8.3	4.4 $\pm$ 2.0	1.7-8.5	2.36 $\pm$ 0.9	1.1-3.9
POC	1.3 $\pm$ 0.8	0.13-4	2.11 $\pm$ 0.7	0.9-3.7	1.25 $\pm$ 0.6	0.25-4.0	0.67 $\pm$ 0.3	0.13-1.3	1.4 $\pm$ 0.8	0.56-2.9

Table 3 Fractional contribution (%) from four major source regions including south Asia (SAS), East Asia (EAS), Southeast Asia (SEA), and Middle East (MDE) to BC surface concentrations over the Mt. Yulong region in winter (December-February), pre-monsoon (March-May), monsoon (June-September), post-monsoon (October-November), and all months during the model simulation time period (2010-2014).

Source region	Winter	Pre-monsoon	Monsoon	Post-monsoon	Annual
SAS	40.8	67.63	17.12	27.14	42.49
EAS	53.76	25.37	77.96	68	52.1
SEA	3.03	2.52	3.73	3.18	3.02
MDE	1.06	2.39	0.47	0.67	1.02






Fig. 1 Location of Mt. Yulong in the Tibetan Plateau and the sampling sites at Mt. Yulong and Ganhaizi basin.



**Fig. 2** Time series of meteorological parameters (air-temperature, net radiation, pressure, rainfall, relative humidity) at Mt. Yulong from December 2014 to December 2015. Monsoon and non-monsoon seasons (including winter, pre-monsoon, post-monsoon seasons) are divided by vertical lines.




Fig. 3 Dependence of optical attenuation (ATN) detected at 632 nm on the EC loading ( $EC_s$ ) for (a) the Mt. Yulong sampling site and (b) the GHZ sampling site. Results of linear regression are displayed with  $K$  as the slope and  $b$  as the intercept.





Fig. 4 Seasonal variations of EC, OC, and POC concentrations and OC/EC ratio from Mt. Yulong during Dec. 2014-Dec. 2016.



**Fig. 5** Seasonal variations of EC, OC, and POC concentrations and OC/EC ratio from GHZ basin during Dec. 2014-Dec. 2016.





Fig. 6 Monthly averaged EC, OC concentrations and OC/EC ratios from Mt. Yulong.




Fig. 7 Seven-day backward trajectory analysis with HYSPLIT model (a) in the monsoon and (b) post monsoon seasons during the study period (Source ★ at 27.01 N, 100.20 E). The trajectories of air mass in the plot were the average of a few episodes. The two heights are the elevation of Mt. Yulong and GHZ, respectively.





Fig. 8 Histogram of (a) annual mean and (b) seasonal mean MAE values of atmospheric EC from Mt. Yulong and GHZ sampling sites during December 2014-December 2016.



**Fig. 9** Regression analysis for scatter plot between EC MAE and POC/OC, OC/EC ratios, and EC<sub>s</sub>, OC concentrations for aerosols sampled at Mt. Yulong.




Fig. 10 Regression analysis for scatter plot between EC MAE and POC/OC, OC/EC ratios, and EC<sub>s</sub>, OC concentrations for aerosols sampled at GHZ.




Fig. 11 Inter-annual differences of OC, EC, and POC concentrations at (a) Yulong and (b) GHZ site. Precipitation at Mt. Yulong and GHZ sites in different years was also plotted in (a) and (b). The OC/EC, and POC/OC ratios are shown in (c, d).




Fig. 12<sup>1</sup> Comparison of EC concentrations in aerosols from Mt. Yulong and other areas of interest.

---

<sup>1</sup> Annual mean EC concentrations at Mt. Yulong were rather low ( $\sim 1.5 \text{ g m}^{-3}$ ), while most of the compared EC concentrations (counted by number of measurements) were within the range of  $1.0\text{--}2.5 \text{ g m}^{-3}$ .




Fig. 13 CALIPSO retrieved backscatter signal at 532 nm (upper panel) and aerosol sub-type information (bottom panel) on 15 March, 2015. The Mt. Yulong region was covered by a thick aerosol layer (mainly consisting of polluted smoke and dust) in the high atmosphere (above 6000 m asl), likely transported far from their source regions. CALIPSO profiles were obtained from (<http://www-calipso.larc.nasa.gov>). The CALIOP-derived reflectivity is usually taken as an indicator to reflect the structure of atmospheric layers since it depends on mass concentration and optical properties of atmospheric aerosol (Bou et al., 2010; Dong et al., 2017a). The topography is outlined by a solid red-line. Suspended dust and aerosol pollutants are in orange and red.




Fig. 14 Annual and seasonal (winter, pre-monsoon, monsoon and post-monsoon) mean relative contributions of emissions in four tagged source regions, including South Asia, East Asia, Southeast Asia and Middle East, to near-surface BC concentrations. The black box in each panel marks the grid box where the Yulong and GHZ sites locate.