

## Anonymous Referee #1

This is a very interesting study on PAR measurements. There are not too many studies on PAR long term measurements and this study is presenting a lot of new and interesting aspects of global and diffuse par. I would suggest the publication of the manuscript after some minor revisions.

*We thank the reviewer for the positive comments.*

The abstract is overloaded with quantified results and is difficult to follow in the current form. Please shorten it and provide only the main quantified findings in fewer paragraphs.

*The abstract was shortened as suggested.*

Page 2, line 53 needs a reference.

*The reference was added.*

Page 3, line 83 add months to the covered dataset period (e.g. Jan 2002 - Dec 2016).

*The months were added.*

Page 4, line 07 and page 6 lines 68-69 are there any additional uncertainty information (e.g. for hourly MFRSR data) about the angular response in order to exclude data for  $SZA > 80$  degrees?

*The study by Mizoguchi et al. (2010) shows that above  $80^\circ$  the cosine error may be of the order of 50%. Although the estimated error on the global irradiance is expected to be smaller, it may still be significant (of the order of 15-20% on the hourly mean).*

Page 4, lines 20 and 28 provide a sentence explaining to the basics the Langley plot method with a relevant reference and mention the additional AOD uncertainty.

*A short description was added.*

Maybe Sections 3 and 4 can be under the same Section with potential title "Global and diffuse PAR irradiances". Then, the current section 3 would be 3.1 under the title "All sky conditions" and the current section 4 would be 3.2 under the title "Cloud-free conditions", while the sub-sections will be renamed accordingly (3.1.1, 3.2.1 and 3.2.2 respectively).

*The sections were rearranged as suggested.*

Section 5 is recommended to be renamed as "Sensitivity analysis"

*We do not agree with this suggestion. Maybe only part of section 5 might fall under this title. We have preferred to leave the title "Cloud effects" to this section.*

Add grid lines in Fig. 1.

*Gridlines were added to figure 1.*

link of the long term series with a discussion and details of the calibration series results and uncertainty

*This comment is not clear.*

Since Lampedusa is severely affected by dust intrusions it would be important to discuss also aerosol attenuation in addition to Cloud attenuation. If the paper is focusing only on cloud effects this have to be mentioned clearly in the discussion or/and in the title. If there will be another manuscript describing the aerosol effects then this has to be mentioned too. Otherwise you can dedicate at least a paragraph on the issue including results of a number of publications on aerosol/dust effects on solar radiation at the site, that authors of this manuscript have investigated in depth in the past.

*We are preparing a second paper dedicated to the aerosol influence on PAR at Lampedusa. This was very shortly stated in the paper (page 14, lines 44-45); we have better addressed this aspect in the introduction and in the conclusions.*

I would point out more in the conclusions the significance of this study (and PAR and diffuse PAR series, seasonal analysis e.t.c.) for agricultural related applications.

*We have added a comment on this topic. At the same time, since measurements used in the paper are obtained in a marine environment, we have also emphasized the potential interest for the estimate of primary production and biological processes occurring in the Mediterranean.*

happy holidays and wishes for a happy new year, to all authors and the editor.

*We thank the reviewer and wish the best for the current year.*

## Anonymous Referee #2

This manuscript covers efficiently the topic of PAR from the point of view of statistics information and I suggest to be published in the frame of this special issue. I am quite satisfied with the work and the discussion done about the performance of the used instruments and my overall impression is that the authors have a well-established background about the topic. Eventhough I could accept the manuscript as it is, I would like to make a few suggestions to the authors in order to improve it.

*We thank the reviewer for the positive comments and for the suggestions.*

Page 2, first paragraph (lines 46-49) and second paragraph (lines 50-53): References are needed

*References were added in the first and second paragraph of page 2.*

Figure 1: As I can understand the real slope and the theoretical slope (linear fit) differ more at low zenith angles where the PAR values are higher. Add an analysis about the dependence of the ratio PAR to PAR<sub>L</sub> versus solar zenith angle.

*We have checked that the linear relationship holds for different values of the solar zenith angle. There is some change in the slope with the selected solar zenith angle intervals; the differences are small (the slope changes between 0.99 and 1.07 when different ranges of solar zenith angles are considered) and the overall uncertainty remains within the stated estimates. It must also be pointed out that part of the large spread found for values of PAR above  $50 \text{ W m}^{-2}$  is due to the role of small time differences between the two observations during cloudy periods. We have redrawn figure 1 removing the fit, which is not used in the analysis and is not discussed in the text.*

Page 6, line 74: "This interannual variability is larger for the diffuse than for the global PAR". Quantify both variabilities and then conclude the above sentence.

*The interannual variability of the annual mean global and diffuse PAR was quantified and discussed.*

Page 6, lines 76-78: "while diffuse PAR displays a more articulated seasonal evolution, with a first maximum in April and a secondary in June". Give a physical explanation for the two maxima.

*This effect is primarily due to the aerosol seasonal evolution. A sentence was added in the text.*

Figure 11 is useless and complicated. It should be removed.

*We have preferred to leave figure 11. Although it is complicated, is the basis for the discussion of several events which characterize the interannual variability of the cloud radiative effects. We believe that this discussion is useful to interpret the dataset, and would not be supported without the figure.*

figure 12 & 13: Since you apply linear regression in order to find the dependence, I suggest to apply multilinear regression for gaining clearer results and safer conclusions.

*We thank the reviewer for the suggestion. We have applied a multilinear regression with respect to the three cloud properties. We have replaced figure 12 and the associated discussions with the results of the new analysis.*

In the future, I hope that the authors should publish a work expanding and combining the findings of the current manuscript with theoretical data (radiation transfer model's outputs). It will be easier to qualify and quantify the factors affecting PAR.

*We thank the reviewer for the suggestion. We are now writing a second paper dealing with the effects of aerosol, based on the cloud-free PAR and aerosol optical properties. The analysis incorporating radiation transfer modelling will be the object of future studies.*

The manuscript follows with the relevant changes marked in red.

# A long-term time series of global and diffuse photosynthetically active radiation in the Mediterranean: interannual variability and cloud effects

Pamela Trisolino<sup>1,2</sup>, Alcide di Sarra<sup>1</sup>, Fabrizio Anello<sup>3</sup>, Carlo Bommarito<sup>3</sup>, Tatiana Di Iorio<sup>1</sup>, Daniela Meloni<sup>1</sup>, Francesco Monteleone<sup>3</sup>, Giandomenico Pace<sup>1</sup>, Salvatore Piacentino<sup>3</sup>, and Damiano Sferlazzo<sup>4</sup>.

<sup>1</sup> ENEA, Laboratory for Observations and Analyses of Earth and Climate, Roma, Via Anguillarese 301, 00123, Italy

<sup>2</sup> Department of Ecological and Biological Sciences, University of Tuscia, Viterbo, Largo dell'Università snc, 01100, Italy

<sup>3</sup> ENEA, Laboratory for Observations and Analyses of Earth and Climate, Palermo, via Principe di Granatelli 24, 90139 Italy.

<sup>4</sup> ENEA, Laboratory for Observations and Analyses of Earth and Climate, Lampedusa, Contrada Capo Grecale, 92010, Italy

Correspondence to: [pamela.trisolino@gmail.com](mailto:pamela.trisolino@gmail.com)

**Abstract.** Measurements of global and diffuse photosynthetically active radiation (PAR) have been carried out on the island of Lampedusa, in the central Mediterranean Sea, since 2002. PAR is derived from observations made with multi filter rotating shadowband radiometers (MFRSRs) by comparison with a freshly calibrated PAR sensor and by relying on the on-site Langley plots. In this way, a long-term calibrated record covering the period 2002-2016 is obtained and is presented in this work.

The monthly mean global PAR peaks in June, with about  $160 \text{ W m}^{-2}$ , while the diffuse PAR reaches  $60 \text{ W m}^{-2}$  in spring or summer. The global PAR displays a clear annual cycle with a semi amplitude of about  $52 \text{ W m}^{-2}$ . The diffuse PAR annual cycle has a semi amplitude of about  $12 \text{ W m}^{-2}$ .

A simple method to retrieve the cloud-free PAR global and diffuse irradiances in days characterized by partly cloudy conditions has been implemented and applied to the dataset. This method allows to retrieve the cloud-free evolution of PAR, and to calculate the cloud radiative effect. CRE, for downwelling PAR. The cloud-free monthly mean global PAR reaches  $175 \text{ W m}^{-2}$  in summer, while the diffuse PAR peaks at about  $40 \text{ W m}^{-2}$ .

The cloud radiative effect, CRE, on global and diffuse PAR is calculated as the difference between all-sky and cloud-free measurements. The annual average CRE is about  $-14.7 \text{ W m}^{-2}$  for the global, and  $+8.1 \text{ W m}^{-2}$  for the diffuse PAR. The smallest CRE is observed in July, due to the high cloud-free conditions frequency. Maxima (negative for the global, and positive for the diffuse component) occur in March-April and in October, due to the combination of elevated PAR irradiances and high occurrence of cloudy conditions. Summer clouds appear to be characterized by a low frequency of occurrence, low altitude, and low optical thickness, possibly linked to the peculiar marine boundary layer structure. These properties also contribute to produce small radiative effects on PAR in summer.

32 The cloud radiative effect has been de-seasonalized to remove the influence of annual irradiance variations. The monthly  
33 mean normalized CRE for global PAR can be well represented by a multi-linear regression with respect to monthly cloud  
34 fraction, cloud top pressure, and cloud optical thickness, as determined from satellite MODIS observations. The behaviour  
35 of the normalized CRE for diffuse PAR can not be satisfactorily described by a simple multi-linear model with respect to the  
36 cloud properties, due to its non linear dependency, in particular on the cloud optical depth. The analysis suggests that about  
37 77 % of the global PAR interannual variability may be ascribed to clouds variability in winter.

## 38 **1 Introduction**

39 The solar radiation comprised between 400 and 700 nm is defined photosynthetically active radiation (PAR, McCree, 1972),  
40 because it is involved in primary production and photosynthetic processes. The knowledge of PAR reaching the Earth's  
41 surface is essential for the determination of biomass production and for the understanding of terrestrial and marine  
42 vegetation physiology (Monteith, 1994; Field et al., 1998).

43 The spectral distribution of solar radiation and PAR, as well as the fraction that reaches the surface, is modified by the  
44 atmospheric components through absorption and scattering by clouds and aerosols, and through absorption by ozone and  
45 other minor gases, as a function of the solar zenith angle, and surface albedo. Beside solar zenith angle, clouds are the main  
46 modulators of PAR (e.g., Alados et al., 1996).

47 Terrestrial and marine vegetation may respond in complex ways to PAR, due to combination of photosynthesis, photo-  
48 inhibition, and photo-damage (e.g., Dimier et al., 2009). The repartition between direct and diffuse PAR, which depends on  
49 the solar zenith angle, atmospheric properties, and albedo, also affects primary production (e.g., Gu et al., 2002; Mercado et  
50 al., 2009, Min, 2005). The determination of these two components is very important for the understanding and description  
51 of photosynthetic processes.

52 Despite its importance, few direct measurements of PAR are carried out routinely, especially for the diffuse component.  
53 These measurements are particularly lacking over the sea, where very few measurements are available. This applies also to  
54 the Mediterranean, which is characterized by high levels of solar (e.g., Hatzianastassiou et al., 2005) and photosynthetically  
55 active radiation (e.g., Hader et al., 2008). Due to the complexity of the basin and of the occurring interactions, and the high  
56 anthropogenic pressure, long-term measurement programs have been started in the Mediterranean with the aim of  
57 understanding the regional climate, underlying mechanisms, and impacts. Basin-wide experiments like the Chemistry and  
58 Aerosol Mediterranean Experiment (dedicated to chemistry-climate interactions; Mallet et al., 2016), the Marine Ecosystem  
59 Response in the Mediterranean Experiment (addressing the marine ecosystem; the Mermex group, 2011), and the Hydrology  
60 Cycle in Mediterranean Experiment (dedicated to the water cycle, Drobinski et al., 2014), were started. Long-term  
61 observations play an important role within these experiments, for the understanding of variability and of the definition of  
62 conditions occurring during intensive measurement campaigns. This study is dedicated at discussing the long-term  
63 behaviour of PAR in the central Mediterranean based on observations made at the Atmospheric Observatory on the island of

64 Lampedusa, in the central Mediterranean. Lampedusa is one of the long-term supersites of ChArMEx (Mallet et al., 2016),  
65 and an atmospheric site of HYMEX. Moreover, the determination of PAR in the marine environment is crucial for the  
66 quantification of the biomass production and the understanding of ecosystem processes. Thus, this study provides basic  
67 information related with different aspects of the regional Mediterranean climate.

68 Different approaches have been used to derive PAR: measurements with dedicated sensors (e.g., Alados and Alados-  
69 Arboledas, 1999), estimates from satellite observations (e.g., Frouin and Pinker, 1995), as a fraction of the broadband solar  
70 irradiance (e.g., Meek et al., 1984; Alados et al., 1996), or using empirical expressions (Udo and Aro, 2000). In a previous  
71 work (Trisolino et al., 2016), we developed a simple method to obtain calibrated global and diffuse PAR irradiances from  
72 measurements made with a multi-filter rotating shadowband radiometer, MFRSR. The method relies on narrowband  
73 measurements of global and diffuse irradiance in 4 bands within the PAR spectral interval, and on the possibility to  
74 frequently obtain on-site calibrations with the Langley plot method, without interruptions of the routine measurements.

75 In this work, we apply the method by Trisolino et al. (2016) to MFRSR measurements made at Lampedusa and derive a  
76 long-term series of global and diffuse PAR. The dataset covers the period January 2002-December 2016 and is used to  
77 investigate the behaviour and variability of global and diffuse PAR, and to quantify the influence of clouds on these  
78 components. The role of aerosols, which also play a large role in modulating PAR in the Mediterranean, will be discussed in  
79 a follow-up paper. Section 2 describes the site and the instruments used in this study. The long-term time series of global  
80 and diffuse PAR is presented in section 3 for all-sky and cloud-free conditions. Cloud effects are discussed in section 4.

## 81 **2 Site and instruments**

82 The instruments used for this study are Li-190 PAR sensors and MFRSR radiometers installed at the ENEA (Italian National  
83 Agency for New Technologies, Energy, and Sustainable Economic Development) Atmospheric Observatory on the island of  
84 Lampedusa (35.5° N, 12.6° E; <http://www.lampedusa.enea.it/>), in the central Mediterranean Sea. The Observatory is located  
85 along the North-Eastern promontory of Lampedusa, at about 45 m above mean sea level. The instruments are installed on  
86 the roof of the main Observatory building, at about 15 m from the Northern cliff of the island; the instruments field of view  
87 is almost totally devoid of obstacles. Lampedusa is a small island (about 20 km<sup>2</sup> surface area) relatively far (> 100 km) from  
88 continental regions and with a very limited impact from local sources. A large set of instruments for the characterization of  
89 regional climate and relevant parameters (radiation, greenhouse gases, aerosol, water vapour, clouds, ozone, etc.; see, e.g.,  
90 Ciardini et al., 2016) is operational at the Lampedusa Atmospheric Observatory, which contributes to the World  
91 Meteorological Organization Regional Global Atmosphere Watch Network.

92 Measurements of PAR global irradiance were started in Lampedusa with a Li-Cor 190 radiometer in June 2004.  
93 The PAR sensor consists of a diffuser, a visible bandpass interference filter, and a Si photodiode, and measures  
94 the down-welling PAR irradiance. According to Ross and Sulev (2000) the systematic spectral errors of the Li-  
95 190 do not exceed 1 %; this sensor also display a good angular response, with a maximum deviation from the

96 ideal cosine response of 7 % at 80°, and larger deviations for larger angles (Akitsu et al., 2017). Different Li-190  
97 PAR sensors have been operational at Lampedusa; only data from two freshly calibrated instruments in two  
98 periods are used in this study, essentially to derive regression parameters which allow to obtain PAR irradiances  
99 from MFRSR signals, as in Trisolino et al. (2016). Two hundred Watt quartz tungsten halogen lamps traceable to  
00 the U.S. National Institute of Standards and Technology (NIST) have been used during the calibration by the  
01 manufacturer. The estimated uncertainty is less than  $\pm 5$  % (typically  $\pm 3$  %). The PAR irradiance measured with  
02 a Li-190,  $PAR_L$ , is used as reference determination during a period of 6-7 months following the factory  
03 calibration.

04  
05 The Multi-filter Rotating Shadowband Radiometer uses six narrow-band and one broadband channels with a  
06 single input diffuser and different interference-filter and photodiode detectors, to measure global and diffuse  
07 components of the solar irradiance. The six narrow-bands have a bandwidth (FWHM) of about 10 nm and are  
08 centred at the wavelengths of about 415, 500, 615, 673, 879, and 940 nm. With an automatic rotating shadow-  
09 band, the instruments measures the diffuse component of the irradiance. This allows to derive the irradiance  
10 direct component as the difference between global and diffuse measurements (Harrison, 1994). MFRSR  
11 measurements at Lampedusa were started in 2001, and different instruments have been used in different periods.  
12 The data used in this study are from MFRSR #424, operational from 2001 to 2011, and from MFRSR #586,  
13 which was installed in 2011 and has been operational since then. The central wavelengths of the 4 bands within  
14 the PAR spectral range for the two MFRSR radiometers are reported in Table 1. MFRSR data are calibrated on  
15 site with the Langley plot method, and the aerosol optical depth is derived routinely at several wavelengths. **The**  
16 **Langley plot methods allows to derive estimates of the signal that would be measured by the instrument at zero**  
17 **air mass. This signal is derived by extrapolating the curve of the logarithm of the signal when plotted versus the**  
18 **air mass, and good estimates are obtained in days with stable atmospheric optical depths. Thus, the zero air mass**  
19 **signal may be derived from ordinary measurements, by selecting days characterized by cloud-free and low**  
20 **aerosol variability conditions.** Further details on the MFRSR radiometers operating at Lampedusa and on the  
21 applied calibration method are given by di Sarra et al. (2015). In addition to MFRSR, a Cimel Sun-photometer  
22 which is part of the Aerosol Robotic Network (AERONET, Holben et al., 1998) is operational at Lampedusa.  
23 The MFRSR data are corrected for the aerosol forward scattering for cases of large particles, and an aerosol  
24 optical depth record constituted by the integrated dataset of MFRSR and AERONET data is obtained (di Sarra et  
25 al., 2015).

26 A calibration of the PAR sensors at least every two years is recommended by the manufacturer. Using the method  
27 by Trisolino et al. (2016) it is possible derive global and diffuse PAR irradiances with a good accuracy, using  
28 MFRSR measurements and taking advantage of the MFRSR in situ calibration with the Langley plot method.  
29 Two sets of regression coefficients needed to derive PAR from MFRSR measurements in the 4 bands falling in  
30 the PAR range were derived. The first set was derived as in Trisolino et al. (2016), in the period 2 March-30  
31 May, 2007 by comparing MFRSR #424 signals with the irradiance measured with a freshly calibrated PAR, #  
32 Q36970. The second sets was derived for MFRSR #586, by comparing its signals with PAR # Q100313  
33 measurements in the period 19 September-21 December 2015. It must be noted that two different sets of  
34 coefficients are needed, due to the somewhat different central wavelengths of the filters of the two instruments.  
35 Li-190 # Q36970 was calibrated at the factory in November 2006, and Li-190 # Q100313 in May 2015.

36 Following Trisolino et al. (2016), the PAR irradiance was derived by linearly combining the calibrated MFRSR  
37 signals in the 4 channels falling in the PAR spectral range. The following relationship is used to infer calibrated  
38 PAR measurements:

$$39 \text{ PAR} = c_1 S_1 + c_2 S_2 + c_3 S_3 + c_4 S_4, \quad (1)$$

40 in which  $S_1$ ,  $S_2$ ,  $S_3$ , and  $S_4$  are the signals at channels centred at about 415, 500, 615, and 675 nm, and  $c_1$ ,  $c_2$ ,  $c_3$ ,  
41 and  $c_4$  are coefficients. The coefficients are obtained by minimizing the difference between PAR and PAR<sub>L</sub>. The  
42 two sets of coefficients for the two MFRSR radiometers used in this study are reported in table 2.

43 As discussed in Trisolino et al. (2016), the estimated uncertainty on the MFRSR-derived PAR is about 5 % for  
44 the global, and 9 % for the diffuse irradiance.

45 MFRSR-derived diffuse PAR irradiances were compared with radiative transfer model (MODTRAN 5.3)  
46 calculations in Trisolino et al. (2016), because direct measurements of diffuse PAR were not available. A Li-190  
47 sensor, #101552, was installed on a sun-tracker for diffuse PAR measurements at Lampedusa on 18 February  
48 2016, and has been operational since then. PAR #101552 was calibrated at the factory in December 2015. Thus,  
49 we have the opportunity to verify the estimated values of diffuse PAR against diffuse PAR<sub>L</sub>. Overlapping data  
50 between 19 February to 25 May 2016 are used in the comparison. The scatterplot and the fit between 5-minute  
51 averages of diffuse PAR derived from MFRSR and diffuse PAR<sub>L</sub> are shown in figure 1 for all atmospheric  
52 conditions.

53 The root mean squared difference between the two datasets is around 9 %. If we sum quadratically the absolute  
54 uncertainty on PAR measurement (3-5 %), we have a total estimated uncertainty of about 10 %. This value is in  
55 agreement with the uncertainty estimate based on the comparison with the radiative transfer model reported by  
56 Trisolino et al. (2016).

57 By comparing  $PAR_L$  from different Li-190 sensors obtained with the original factory calibration with PAR  
58 inferred from MFRSR measurements, it is possible to determine the long-term variation of the Li-190 sensitivity.  
59 All different PAR sensors display a long-term decrease of the instrumental sensitivity. This decrease, calculated  
60 over at least 2 years, is between 0.6 and 2.3 %/year. This is consistent with the results by Mizoguchi et al.  
61 (2010), who found a degradation by 1-3 % after 1 year of field operation for different PAR sensors. Our data  
62 show in some cases a faster decline, up to 15 %/year, for specific sensors over shorter periods (few months),  
63 which appears to take place preferably during the first months of use. Similar reductions were observed by Blum  
64 and Schwank (1985) and, more recently, by Akitsu et al. (2017). This behaviour suggests that a frequent  
65 calibration and redundancy in the measurements may be required to obtain accurate PAR observations. The  
66 possibility to rely on on-site routine calibrations, as is the case for the method used in this study, guarantees a  
67 better long-term accuracy than obtainable from standard PAR observations.  
68

### 69 **3 Global and diffuse PAR irradiances**

#### 70 **3.1 All sky conditions**

71 The time series of monthly mean global and diffuse component of photosynthetically active radiation from  
72 MFRSR measurements over the period 2002-2016 is shown in fig. 2. The MFRSR data are acquired every 15 s;  
73 hourly, and then daily averages are calculated from the 15 s measurements. Monthly means are calculated from  
74 daily values for those months with at least 27 daily average values. Data acquired at solar zenith angles larger  
75 than  $80^\circ$ , affected by a significant cosine response error (e.g., Mizoguchi et al., 2010; Akitsu et al., 2017), are  
76 excluded. This limitation on the solar zenith angle produces an underestimate of the daily PAR by about  $1.4 \pm 0.5$   
77  $Wm^{-2}$  for the global component, and  $1.1 \pm 0.3 Wm^{-2}$  for the diffuse component.

78 Figure 2 shows the monthly mean global and diffuse PAR as derived from MFRSR observations. The two data  
79 series display a typical annual cycle with a summer maximum, driven by the annual course of the solar zenith  
80 angle, with significant interannual variability. The year-to-year changes of the annual mean values are of about 4  
81  $W m^{-2}$  for both global and diffuse PAR; the relative interannual changes are thus much larger for diffuse than for  
82 global PAR.

### 83 3.1.1 Mean annual cycles

84 Figure 3 shows the annual evolution of monthly median, percentiles, maxima and minima of global and diffuse  
85 PAR. Global PAR shows an evident annual cycle with single summer maximum, while diffuse PAR displays a  
86 more articulated seasonal evolution, with a first maximum in April and a secondary in June. This behaviour of  
87 diffuse PAR is primarily due to the large role played by atmospheric aerosols, mainly during the summer months,  
88 when cloud-free conditions are frequent. The role of the aerosol will be discussed in a separate study.

89 Alados et al. (1996) show the annual evolution of global PAR in Almeria ( $36.83^{\circ}$  N,  $2.41^{\circ}$  W), at a latitude close  
90 to that of Lampedusa, but on land. The differences between the monthly median values of the two datasets are  
91 generally within 10 %. Largest differences are found in March.

92 Yu et al. (2015) show ground based measurements of daily global PAR at several sites of the Surface Radiation,  
93 SURFRAD, network in United States for year 2011. Goodwin Creek ( $34.3^{\circ}$  N,  $89.9^{\circ}$  W, 98 m a.m.s.l.) and  
94 Desert Rock ( $36.6^{\circ}$  N,  $116.0^{\circ}$  W, 1007 m a.m.s.l.) are located in the same latitude band of Lampedusa. Despite  
95 they represent very different environments, PAR measurements are in reasonable agreement with those of  
96 Lampedusa: differences with respect to Lampedusa are about 3 % in summer and 4 % in winter for Desert Rock,  
97 and 12 % in summer and 3 % in winter for Goodwin Creek.

98 Jacovides et al. (2004) report global PAR observations made over three years (September 1997-May 2000) in  
99 Athalassa, Cyprus, in the Eastern Mediterranean. The daily values of PAR in Cyprus are in good agreement with  
00 those at Lampedusa, with differences between maximum mean values of about 4 % in summer and 8 % in winter.  
01 Very few studies are dedicated to the annual evolution of the diffuse PAR and, to our knowledge, no direct  
02 measurements over the sea or on small islands are available.

03 Measurements of diffuse PAR at latitudes similar to that of Lampedusa were reported by Dye (2004) and Wang  
04 et al. (2017). Dye et al. (2004) shows measurements of diffuse PAR at the Southern Great Plains Atmospheric  
05 Radiation Measurement, ARM, program site ( $36.6^{\circ}$  N,  $97.5^{\circ}$  W; 314 m) between 1 March and 31 July 2000.  
06 Differences between the monthly mean diffuse PAR at Southern Great Plains and Lampedusa are between 7 % in  
07 July to 19 % in June.

08 Wang et al. (2017) applied various methods to estimate the diffuse PAR to observations made in Canada and in  
09 USA with Delta-T BF3 sensors. Two of the sites used in this study are located at the same latitude of Lampedusa  
10 (US-fuf,  $35.09^{\circ}$  N,  $111.76^{\circ}$  W, and US-fwf,  $35.45^{\circ}$  N,  $111.77^{\circ}$  W) but are at about 2200 m above mean sea level  
11 (2180 m and 2270 m, respectively) and in different terrestrial environments (woody savannas and grassland,

12 respectively). The monthly diffuse PAR at Lampedusa is 25-54 % of that at US-fwf, and 17-52 % of that at US-  
13 fuf. This large difference is primarily due to altitude and albedo, and possibly clouds.

14 Differences in altitude, surface properties and albedo, aerosol, and clouds are expected to play a large role in the  
15 diffuse PAR modulation, and the agreement found between sites at similar latitude may be somewhat incidental.

16 After solar zenith angle, clouds are the main modulation factor, which are expected to act differently on the  
17 global and diffuse PAR components.

18 In general, if we exclude global irradiance enhancements which may occur with scattered cloud conditions, the  
19 global irradiance for cloudy conditions is expected to be always lower than for cloud-free sky. Conversely, the  
20 diffuse irradiance is larger than in cloud-free conditions for thin and moderate clouds, and is smaller for thick  
21 clouds. Thus, the relationship between cloudiness and diffuse PAR is expected to be more complex than for the  
22 global component.

23 Figure 4 shows the monthly mean occurrence of cloud free conditions (cloud cover approximately lower than 1  
24 okta and direct radiation unobstructed; see section 4.1 for more details on the cloud-free identification scheme)  
25 between 9 and 14 UTC. The monthly median and quartiles are calculated over the 2002-2016 time interval,  
26 considering only months with at least 27 days with measurements.

27 The largest interquartile range of the global PAR irradiance is observed to occur in April and August. This is due  
28 to the combination of elevated PAR levels and a higher cloud occurrence than in June and July.

29 Largest median values of diffuse PAR are found in April. The monthly diffuse PAR median is lower in summer  
30 than in April, although the solar zenith angle is higher, due to the higher rate of occurrence of cloud-free  
31 conditions. In addition to the cloud occurrence, cloud properties are expected to play a significant role in  
32 determining the amount of global and diffuse PAR reaching the surface.

33 The average diffuse-to-global PAR ratio is  $0.39 \pm 0.08$ , and displays a marked seasonal variation, with maxima  
34 around 0.5 in winter and minima at about 0.25-0.3 in summer.

35 The monthly time series shown in figure 2 was fit with the following simple oscillatory function:

36  $f(t) = a + b \sin [\omega (t + \varphi)]$  (2)

37 with the aim of quantitatively determining the main parameters of the PAR evolution. The coefficients  $a$ ,  $b$ , and  
38  $\varphi$ , retrieved for the global and diffuse PAR are reported in table 3, while the fitting curve is also included in  
39 figure 2. The variable  $\omega$  is  $2\pi/12$  and  $t$  is in months.

40 As discussed in section 2, our time series is based on the determination of weighting coefficients for the signals  
41 of two MFRSR instruments against two freshly calibrated PAR sensors. The two determinations took place in

42 2007 and 2015. Unfortunately, we do not have overlapping reliable PAR observations in 2011, when MFRSR  
43 #424 stopped working and was replaced by MFRSR #586. Thus, the series relies on two independent  
44 calibrations of PAR sensors, whose nominal accuracy is between 3 and 5 %. Thus, we can not exclude that a step  
45 change in calibration, within the nominal accuracy of the used sensors, took place in 2011. Since there is no  
46 independent verification with a higher degree of accuracy, we have preferred to avoid calculating trends on the  
47 time series. Thus, we did not include a term describing the trend in  $f(t)$ .  
48 As expected, the correlation coefficient between data and the fitting function is higher for global than for diffuse  
49 PAR. The ratio between the annual semi amplitude,  $b$ , and the long-term average is larger for the global than for  
50 the diffuse PAR (0.55 against 0.36).

## 51 **3.2 Cloud-free conditions**

52 As discussed above, clouds play a large role in modulating the PAR evolution and variability. The determination  
53 of PAR in cloud free conditions is necessary to quantitatively assess the role played by clouds. Thus, we have  
54 developed a method to infer monthly mean cloud-free global and diffuse irradiances, which may be used to  
55 determine the cloud radiative effect in the PAR spectral range.  
56

### 57 **3.2.1 Determination of cloud-free PAR**

58 The determination of cloud-free PAR irradiances is based on the identification of cloud-free conditions.  
59 Measurements of MFRSR radiometers are used to select cloud-free periods. The algorithm developed by Biavati  
60 et al. (2004) is an adaptation of the method by Long and Ackerman (2000) to the characteristic conditions and  
61 properties of the aerosol observed at Lampedusa. This algorithm, briefly described by Meloni et al. (2007),  
62 allows to determine periods with no clouds, also with high values of aerosol optical depth. The method is very  
63 selective and allows identifying sky conditions virtually devoid of clouds. A comparison with sky imager  
64 pictures shows that the algorithm selects cases with cloud cover  $\leq 1$  okta and Sun unobstructed by clouds. This  
65 method has been used to select conditions suitable for the determination of aerosol radiative effects, for which the  
66 cloud radiative influence needs to be negligible (see, e.g., di Sarra et al., 2008; Di Biagio et al., 2010).

67 With the aim of estimating the cloud-free irradiance during partly cloudy days we applied to PAR the method  
68 proposed by Long and Ackerman (2000) for the shortwave radiation. The following function

$$69 \quad F(\theta) = A(\cos\theta)^B \quad (3)$$

70 is fit to the PAR data in cloud-free intervals, and is used to estimate the cloud-free expected irradiance during  
71 cloudy intervals. The variable  $\theta$  is the solar zenith angle and  $A$  is the cloud-free irradiance for a solar zenith angle  
72 of  $0^\circ$ ; the fitting curve includes the effects of the atmosphere constituents (changes in atmospheric pressure,  
73 aerosol, and absorbing gases) which affect PAR and are assumed to vary slowly, and possible residual  
74 instrumental calibration inaccuracies. The  $B$  coefficient includes possible effects of the cosine response of the  
75 instrument. The contribution of water vapour to the modulation of PAR is negligible; this is an advantage in  
76 applying this fitting procedure to PAR data with respect to the whole shortwave range.

77 Figure 5 shows the measured PAR and the derived fit for a day with only a portion with cloud-free conditions.  
78 The coefficients are determined for every partly cloudy day. The Long and Ackerman method requires some  
79 minimum number of cloud-free measurements over a significant range of solar zenith angle. The method is not  
80 applicable to days with persistent overcast conditions, or when few short cloud-free intervals are present. Thus,  
81 only a selection of cases is retained in the cloud-free dataset. These are selected by visually looking at the fitting  
82 curve and eliminating the days with incomplete or inadequate fits.

83

### 84 **3.2.2 Long-term cloud-free PAR irradiances**

85 We obtain a long-term time series of cloud-free PAR estimates using the method described in section 4.1. The  
86 monthly mean values for cloud-free conditions are calculated from daily means with at least 11 daily values.  
87 During the winter months also averages calculated with at least 5 values are accepted for diffuse PAR, due to the  
88 low number of available days for which it is possible to estimate the cloud-free irradiance. The monthly mean  
89 time series of global and diffuse irradiance is shown in fig. 2 together with those obtained in all sky conditions.

90 As expected, the global irradiance is always larger than for all sky conditions, while the opposite occurs for the  
91 diffuse PAR. The average diffuse-to-global PAR ratio is  $0.237 \pm 0.035$ ; the seasonal variation is much smaller  
92 than for all-sky conditions.

93 As for the PAR measurements in all sky conditions, the time series was fit with expression (1). The fitting  
94 coefficients for cloud-free series are reported in table 4.

95 The values of the annual semi-amplitudes are very close for all-sky and cloud-free conditions, while the long-  
96 term average is larger for cloud-free conditions for global PAR, and larger for all-sky conditions for diffuse PAR.

97 The ratio between the annual cycle semi-amplitude and the long-term average is about 0.45 for both the global

98 and the diffuse components for cloud-free conditions. The ratio between the diffuse and the global PAR long-  
99 term averages is 0.24; this is significantly smaller than for all-sky conditions.

00

### 01 **3.2.3 Mean annual cycles**

02 The average annual evolution of global and diffuse cloud-free PAR is shown in fig. 6. The PAR interannual  
03 variability in this case is primarily attributable to aerosol. As discussed by di Sarra et al. (2015) and by Di Iorio  
04 et al. (2009), the aerosol distribution at Lampedusa is characterized by a significant annual cycle, essentially  
05 modulated by the occurrence of desert dust events, which generally occur in spring and summer (Meloni et al.,  
06 2007). The day-to-day variability is largest in spring and summer, and smallest in winter. This behaviour is  
07 evident in the variability of both global and diffuse PAR. Previous studies carried out at Lampedusa have shown  
08 that this aerosol variability largely affects UV (di Sarra et al., 2002; Meloni et al., 2005; Casasanta et al., 2011),  
09 solar radiation (di Sarra et al., 2008; Di Biagio et al., 2010) and, for desert dust particles, infrared radiation (di  
10 Sarra et al., 2011; Meloni et al., 2015).

11 Figure 7 shows the comparison between the monthly average for all-sky and for cloud-free conditions. The  
12 difference between the two curves is the monthly mean cloud radiative effect, CRE, which is produced by clouds  
13 on downwelling global and diffuse PAR at Lampedusa. The reader is reminded that this analysis is based on a  
14 selective cloud screening, and partially cloudy or short cloud-free intervals are classified as cloudy by our  
15 algorithm.

## 16 **4 Cloud effects**

17 The differences between all-sky and cloud-free global and diffuse downwelling irradiances are shown in figure 8.  
18 As expected, CRE is negative (a reduction of down-welling PAR) for global, and positive (an increase) for  
19 diffuse PAR. The long-term average cloud radiative effect is about  $-14.7 \text{ W m}^{-2}$  for the global component, and  
20  $+8.1 \text{ W m}^{-2}$  for the diffuse PAR. The interannual variability is significant.

21 Extended summer periods with small values of CRE ( $>-7 \text{ W m}^{-2}$  for the global component) occur in 2003, 2008,  
22 2009, 2012, 2013, and 2015. It is interesting to note that elevated summer temperatures were recorded at  
23 Lampedusa during the summer seasons of 2003, 2009, 2012, and 2015 (Ciardini et al., 2016). Some of these  
24 events (e.g., 2003, 2015) were recorded at the regional level.

25 Extended time intervals with large cloud effects occurred in several periods (early 2003, 2005, 2006, 2010),  
26 while cases with large cloud effects over limited periods are also present (late 2007, early 2013 and 2014, mid  
27 2016). Some of these negative peaks appear to correspond with cold winters at Lampedusa (e.g., in 2003, 2005,  
28 2006, 2009, 2012, 2013, and 2015; Ciardini et al., 2016).

29 Figure 9 shows the annual evolution of the monthly median CRE and its variability, for the global and diffuse  
30 component. The absolute value of the diffuse CRE displays an annual evolution similar, but opposite in sign, to  
31 the global. The absolute value of CRE is smallest during months characterized by dominant cloud-free  
32 conditions, generally in late summer, and shows a maximum generally between March and April, and sometimes  
33 a secondary maximum in October. For the diffuse CRE the maximum variability takes place in September, and  
34 the largest CRE median in April.

35 Pyrina et al. (2015) investigated the effect of clouds on the shortwave radiation in the Mediterranean basin in the  
36 period 1984-2007 based on satellite observations and radiative transfer model calculations. The seasonal patterns  
37 of the surface CRE in the shortwave,  $CRE_{SW}$ , is similar to that for CRE of global PAR in fig. 10, with maximum  
38 values in April-May and a secondary maximum in October. The secondary maximum in October is less  
39 pronounced in the analysis by Pyrina et al. than at Lampedusa. Data by Pyrina et al. show a significant  
40 variability of  $CRE_{SW}$  throughout the basin, and particularly in the North-South direction, except in October.

41 Pyrina et al. (2015) also show that the radiative effects depend on the cloud type.

42 We have investigated this aspect by extracting monthly average cloud properties retrieved by the Moderate  
43 Resolution Imaging Spectroradiometer, MODIS, on board Aqua over a region  $2^{\circ} \times 2^{\circ}$  around Lampedusa. Level 3  
44 monthly mean Cloud Optical Thickness (COT), Cloud Fraction (CF) and Cloud Top Pressure (CTP) determined  
45 from July 2002 to December 2016 are used (MOD08\_M3\_6). Monthly averages from daytime observations from  
46 MODIS Aqua, whose daytime passage over Lampedusa is close to solar noon, have been selected.

47 Level 3 MODIS products are based on a cloud mask for the identification of cloudy pixels, with a resolution of 5  
48 km.

49 Cloud Optical Thickness is derived from multispectral reflectance measurements compared with parametrizations  
50 obtained from a theoretical model, simulated in function of 3 different geometrical angles. The look-up table of  
51 the theoretical model results contains reflectance values, effective radii, and surface albedo (King et al., 1997).

52 The cloud fraction is calculated as the ratio between the number of cloudy pixels and the total number of pixels in  
53 the image.

54 CTP is determined using the radiance measurements in the CO<sub>2</sub> absorbing region from 13.3 to 14.2 μm using a  
55 CO<sub>2</sub>-slicing technique. An infrared channel at 11 μm is also used for optically thick and lower-level clouds  
56 (Platnick et al., 2015).

57 Figure 10 shows the annual evolution and variability of monthly values of COT, CF, and CTP. A significant co-  
58 variance, in particular of CF and CTP, is apparent, with lowest clouds and smallest CF in July, when also a  
59 minimum of COT is observed. Thus, cloud properties vary seasonally, with lower, although less frequent and  
60 thinner, clouds occurring in summer. The annual evolution of CF and of the fraction of cloud-free conditions of  
61 fig. 4 are also highly consistent. Contrarily to expectations, low level clouds are associated, especially in July,  
62 with low values of COT. This may be related, in addition to prevailing synoptic conditions, to the seasonal  
63 evolution of the planetary boundary layer at Lampedusa, which is dominated by the thermal difference between  
64 the sea surface temperature and the atmosphere. This produces strongest temperature inversions during summer  
65 (see e.g., Pace et al., 2012), with the possible formation of thin low level clouds at the top of the inversion layer,  
66 which is usually located within 200 m of the sea surface. The frequency of occurrence and properties of clouds in  
67 summer favour high level of PAR and, similarly, UV radiation (e.g., Meloni et al., 2005). As discussed by  
68 Becagli et al. (2013), these elevated radiation levels, together with a shallow marine mixed layer, stress the  
69 marine biological cells; these mechanisms lead to increased di-methyl sulphide emissions and high values of the  
70 phytoplankton productivity index (rate of photosynthetic carbon assimilation per weight of chlorophyll a) in  
71 summer, while the chlorophyll amount displays a maximum in winter.

72 The largest cloud radiative effects occurring in the transition seasons appear to be associated with moderate  
73 values of COT (except in November, when the median is higher and the minimum-to-maximum COT excursion  
74 is largest) and relatively high clouds.

75 CRE is modulated, in addition to clouds, by the annual evolution of the photosynthetic irradiance. To reduce the  
76 effect of the seasonal changes in irradiation we have defined as normalized cloud radiative effect, NCRE, the  
77 ratio between CRE and the cloud-free PAR irradiance of the corresponding month. NCRE is between about -0.5  
78 and 0 for the global PAR, and between 0 and +0.8 for the diffuse PAR, with larger relative variations for diffuse  
79 than for global PAR.

80 Figure 11 shows the time series of global and diffuse NCRE together with cloud fraction, cloud optical thickness,  
81 and cloud top pressure determined by MODIS. The variability of cloud properties helps explaining the  
82 interannual variability of NCRE, in particular for global PAR.

83 Winters 2009-2010 and 2015-2016 are characterized by a less pronounced seasonal minimum of global NCRE,  
84 which is associated with low values of COT in these seasons compared to the other years. The winter 2009-2010

85 clouds produce also the largest effect on the diffuse NCRE. The COT seasonal cycle appears to be attenuated in  
86 2007, with an anticipation of the winter maximum already in 2006 and a lacking summer minimum. This seems  
87 to produce a smaller winter negative peak of NCRE, but does not seem to affect the summer maximum, probably  
88 because of the annual evolution of CF and the prevailing role of cloud-free conditions. The largest NCRE for the  
89 month of April is also found in 2007. This is also the largest CRE of the whole record, and appears to be due to  
90 elevated values of COT in this season compared with the other years; CTP and CF do not show significant  
91 anomalies in this season.

92 Several winter seasons display minima of the global NCRE less than -0.3. Two of the three largest NCRE  
93 negative peaks, in early 2006 and 2014, coincide with maxima of both COT and CF. The other peaks, in early  
94 2003, December 2004, early 2008 and 2012, coincide with maxima of one between COT and CF.

95 *A multi-linear regression of the form*

$$96 \text{NCRE}_c = A \text{CF} + B \text{COT} + C \text{CTP}$$

97 (4)

98 has been applied to the data.  $\text{NCRE}_c$  is the NCRE value estimated by the regression model. The A, B, and C  
99 coefficients have been derived by least square minimization of the differences between  $\text{NCRE}_c$  and NCRE.  
100 Table 5 reports the coefficients obtained for the global and diffuse PAR NCRE, and the average percent weight  
101 of each term in expression 4) with respect to  $\text{NCRE}_c$ . Figure 12 shows the scatterplots between  $\text{NCRE}_c$  and  
102 NCRE for the global and diffuse PAR. The mean bias, correlation coefficient, and root mean squared deviation  
103 (RMSD) between  $\text{NCRE}_c$  and NCRE are indicated in the graphs. The multi-linear model performs well for  
104 global NCRE, while is not capable of describing the behaviour of the diffuse NCRE. This is expected, due to the  
105 non linear dependency between diffuse PAR irradiance and COT discussed in section 3.1.1.

106 The coefficients in table 5 show that the correlation between global NCRE and both CF and COT is negative,  
107 while is positive between global NCRE and CTP.

108 Cloud fraction is the property which most largely affects global NCRE, and explains about 65% of its variability.  
109 COT also plays an important role (41%), while the contribution by CTP is minor. This simple model suggests  
110 that cloud properties might be used as predictors for the global NCRE. Conversely, this is not the case for the  
111 diffuse NCRE. This confirms that direct measurements of diffuse PAR are very critical for the understanding of  
112 photosynthetic and climate-related effects.

113 Given the low correlation coefficient and large RMSD between diffuse  $\text{NCRE}_c$  and NCRE, the coefficients and  
114 contributions for the diffuse component reported in table 5 must be looked with caution. The correlation,  
115 although low, between diffuse NCRE and CF and CTP is positive, while is negative between NCRE and COT.

16  
17  
18  
19  
20  
21

The monthly mean time series was rearranged in order to remove the seasonality and better investigate the interannual variability. The long-term monthly mean annual cycle was calculated from the datasets, as the average of all values for each month (all values for January, then for February and so on) of the record. The de-seasonalized time series is thus calculated as:

$$P_{ij} = \frac{PAR_{ij} - \overline{PAR}_i}{\overline{PAR}_i} \quad (5)$$

22  
23 where  $\overline{PAR}_i$  is the long-term monthly mean for month  $i$  ( $i=1, \dots, 12$ ), and  $PAR_{ij}$  is the monthly mean value for  
24 month  $i$  and year  $j$ .  $P_{ij}$  is thus the fractional change in global PAR with respect to its long-term monthly mean.  
25 Similarly, a deseasonalized value of NCRE,  $N_{ij}$ , was calculated.

26 Figure 13 shows the scatterplot between de-seasonalized PAR and NCRE. The variations of  $P_{ij}$  are within  $\pm 20$   
27 %, while  $N_{ij}$  varies within about  $\pm 80$  %. The best linear fit is also shown in figure 13. The correlation coefficient  
28 is relatively low when all data are included in the scatterplot. Data were also sorted according to the season, over  
29 an extended winter (months of November, December, January, February, and March) and an extended summer  
30 (months of May, June, July, August, and September) period, and the corresponding scatter plots are included in  
31 figure 13. A much stronger correlation is found for the winter months, when NCRE is largest and appears to  
32 explain about 77 % of the PAR interannual variability. The slope of the  $P_{ij}$  versus  $N_{ij}$  fitting line in summer is 4  
33 times smaller than in winter; the range of  $P_{ij}$  values is within  $\pm 0.2$  in winter, and  $\pm 0.1$  in summer. The correlation  
34 is much weaker in summer, with clouds explaining only 31 % of the PAR interannual variability. This is  
35 attributed to the high occurrence of cloud free conditions, and the larger role of atmospheric aerosols, which  
36 typically display a maximum in summer.

## 37 **5 Summary and conclusions**

38 This analysis is based on a new method to estimate global and diffuse component of PAR from calibrated  
39 MFRSR measurements (Trisolino et al., 2016). The method is applied to the long-term MFRSR record at the  
40 ENEA Atmospheric Observatory on the island of Lampedusa. As a complement to the previous study, direct  
41 measurements of diffuse PAR with a Li-190 sensor on a solar tracker were used to verify the reliability of  
42 MFRSR diffuse PAR estimates. The estimated uncertainty is in agreement with previous results obtained from  
43 the comparison with radiative transfer model calculations.

44 Main results of this study may be summarized as follows:

45 1. the long-term global and diffuse PAR record in the central Mediterranean covers the period 2002-2016.  
46 Measurements are being continued, and the record will be expanded in the future. The long-term mean is 95 and  
47 35  $\text{W m}^{-2}$  for global and diffuse PAR, respectively. The mean annual cycle semi-amplitude is about 55 % and 36  
48 % of the long-term mean for the global and diffuse PAR, respectively. Interannual variations of global and  
49 diffuse PAR appear to be essentially related with the clouds frequency of occurrence. The cloud occurrence is  
50 very low in summer, especially in July.

51 2. The daily PAR irradiance for cloud-free conditions was estimated also during partially cloudy days by fitting  
52 the cloud-free data with an analytical function which depends on the solar zenith angle. In this way, cloud-free  
53 global and diffuse PAR time series were derived. The cloud-free long-term mean global PAR is 110  $\text{W m}^{-2}$ ,  
54 about 16 % more than in all-sky conditions. Conversely, the cloud-free diffuse PAR long-term mean PAR is 26  
55  $\text{W m}^{-2}$ , about 24 % less than in all-sky conditions. The cloud-free annual cycle semi-amplitude is 45 % for global  
56 PAR, and 47 % for diffuse PAR. The differences in the annual semi-amplitude-to-long-term mean ratio with  
57 respect to the all-sky conditions are essentially driven by changes in the long-term mean values.

58 3. By using all-sky and cloud-free PAR data we have calculated the cloud radiative effect, CRE, for global and  
59 diffuse PAR. Global and diffuse CRE display a similar annual evolution, but opposite sign, positive for diffuse  
60 and negative for global PAR. The long-term mean CRE is  $-14.7 \text{ W m}^{-2}$  for global PAR and  $+8.1 \text{ W m}^{-2}$  for  
61 diffuse PAR. The CRE seasonal evolution is characterized by a primary maximum in late winter/early spring,  
62 and a secondary maximum in autumn. The largest global PAR monthly CRE is  $-31 \text{ W m}^{-2}$  in April 2007, while  
63 the smallest is generally observed in July, in correspondence with the lowest occurrence of clouds. The largest  
64 diffuse PAR CRE is recorded in April 2010, with  $+18 \text{ W m}^{-2}$ .

65 4. The CRE has been scaled by taking into account the monthly cloud-free estimated PAR to remove the  
66 influence of the solar zenith angle annual course. The normalized CRE has been associated with cloud properties  
67 (fraction, optical thickness, and top pressure) as derived from space with MODIS. **A simple multi-linear model**  
68 **has been applied to relate the deseasonalized NCRE and cloud properties. For global PAR, NCRE is correlated**  
69 **positively with respect to cloud top pressure, and negatively with cloud fraction and optical thickness. Cloud**  
70 **fraction appears as the most relevant parameter influencing NCRE. The multi-linear model is conversely not**  
71 **appropriate to describe the relationship linking diffuse PAR NCRE with cloud properties.** The cloud influence on  
72 diffuse PAR is more difficult to disentangle, due to the stronger dependency on cloud properties. Clouds are  
73 expected to produce an increase or a decrease of diffuse PAR depending on their optical thickness.

74 5. De-seasonalized global PAR data were related with de-seasonalized NCRE. The correlation analysis between  
75 these two parameters suggests that cloud properties explain about 77 % of the interannual global PAR variability  
76 in winter, and about 31 % in summer.

77 6. The large values of global PAR found in summer appear to be due to the combination of low cloud frequency  
78 of occurrence and low cloud optical thickness. The summer clouds are also characterized by a high top pressure,  
79 suggesting that low level thin clouds, possibly forming at the top of the summer temperature inversion, play a  
80 large role. Also due to these cloud properties, high PAR levels are reached in summer. The high PAR and UV  
81 radiation has been observed to stress marine phytoplankton in this region of the Mediterranean (Becagli et al.,  
82 2013).

83 This study contributes to provide new information about the behaviour of global and diffuse PAR in the  
84 Mediterranean, and in the marine environment. **These data, as well the methodology, may be also relevant for**  
85 **agricultural related applications, and for estimates of ocean primary production.** Information on diffuse PAR is  
86 particularly relevant and difficult to be found in the literature.

87 As a prosecution of this study, we plan to better investigate the variability of cloud-free PAR, in particular with  
88 respect to the role of atmospheric aerosol, which play an important role in the Mediterranean radiative processes.

## 89 **Acknowledgements**

90 This study has been partly supported by the Italian Ministry for University and Research through the NextData  
91 and RITMARE projects. We acknowledge the MODIS mission scientists and associated National Aeronautics  
92 and Space Administration personnel for the production of the data used in this research effort. MODIS data were  
93 downloaded from the Giovanni online data system, developed and maintained by the NASA Goddard Earth  
94 Sciences Data and Information Services Center. This study contributes to ChArMEx WP-4 on Aerosol-Cloud-  
95 Radiation-Climate and WP-6 on Variability and Trends.

## 96 **References**

- 97 Akitsu, T., Nishida Nasahara, K., Hirose, Y., Ijima, O. and Kume, A.: Quantum sensors for accurate and stable  
98 long-term photosynthetically active radiation observations, *Agricul. Forest Meteorol.*, 237, 171-183, 2017.
- 99 Alados, I. and Alados-Arboledas, L.: Validation of an empirical model for photosynthetically active radiation,  
00 *Int. J. Climatol.*, 19, 1145–1152, 1999.

01 Alados, I., Foyo-Moreno, I. and Alados-Arboledas, L.: Photosynthetically Active Radiation: measurements and  
02 modelling, *Agr. Forest. Meteorol.*, 78, 121-131, 1996.

03 Becagli, S., Lazzara, L., Fani, F., Marchese, C., Traversi, R., Severi, M., di Sarra, A., Sferlazzo, D., Piacentino,  
04 S., Bommarito, C., Dayan, U. and Udisti, R.: Relationship between methanesulfonate (MS-) in atmospheric  
05 particulate and remotely sensed phytoplankton activity in oligo-mesotrophic central Mediterranean Sea, *Atmos.*  
06 *Environ.*, 79, 681-688, 2013.

07 Biavati, G. and di Sarra, A.: Individuazione dei periodi di cielo sereno da misure di radiazione solare globale e  
08 diffusa, Technical report ENEA, RT/2004/66/CLIM., 2004.

09 Blum, H. and Schwank, O.: The stability of calibration factors of sensors for photosynthetically active radiation  
10 used in wet conditions, *Agricul. Forest Meteorol.*, 34, 77-82, 1985.

11 Casasanta, G., di Sarra, A., Meloni, D., Monteleone, F., Pace, G., Piacentino, S. and Sferlazzo, D.: Large aerosol  
12 effects on ozone photolysis in the Mediterranean, *Atmos. Environ.* 45, 3937-3943, 2011.

13 Ciardini, V., Contessa, G. M., Falsaperla, R., Gómez-Amo, J. L., Meloni, D., Monteleone, F., Pace,  
14 G., Piacentino, S., Sferlazzo, D. and di Sarra, A.: Global and Mediterranean climate change: a short summary,  
15 *Ann. Ist. Super. Sanità*, 52(3),325-337, doi: 10.4415/ANN\_16\_03\_04, 2016.

16 Di Biagio, C., di Sarra, A. and Meloni, D.: Large atmospheric shortwave radiative forcing by Mediterranean  
17 aerosol derived from simultaneous ground-based and spaceborne observations, and dependence on the aerosol  
18 type and single scattering albedo, *J. Geophys. Res.*, 115, D10209, 2010.

19 Di Iorio, T., di Sarra, A., Sferlazzo, D. M., Cacciani, M., Meloni, D., Monteleone, F., Fuà, D. and Fiocco, G.:  
20 Seasonal evolution of the tropospheric aerosol vertical profile in the central Mediterranean and role of desert  
21 dust, *J. Geophys. Res.*, 114, D02201, doi:10.1029/2008JD010593, 2009.

22 Dimier, C., Saviello, G., Tramontano, F., and Brunet, C.: Comparative ecophysiology of the xanthophyll cycle in  
23 six marine phytoplanktonic species, *Protist*, 160, 397–411, 2009.

24 di Sarra, A., Cacciani, M., Chamard, P., Cornwall, C., DeLuisi, J. J., Di Iorio, T., Disterhoft, P., Fiocco, G., Fuà,  
25 D. and Monteleone, F.: Effects of desert dust and ozone on the ultraviolet irradiance at the Mediterranean island  
26 of Lampedusa during PAUR II, *J. Geophys. Res.-Atmos.*, 107, 2-1-2-14, 2002.

27 di Sarra, A., Pace, G., Meloni, D., De Silvestri, L., Piacentino, S., Monteleone, F.: Surface shortwave radiative  
28 forcing of different aerosol types in the central Mediterranean, *Geophys. Res. Lett.*, 35, L02714,  
29 doi:10.1029/2007GL032395, 2008.

30 di Sarra, A., Di Biagio, C., Meloni, D., Monteleone, F., Pace, G., Pugnaghi, S. and Sferlazzo M. D.: Shortwave  
31 and longwave radiative effects of the intense Saharan dust event of 25-26 March 2010 at Lampedusa  
32 (Mediterranean Sea), *J. Geophys. Res.-Atmos.*, 116, D23209, 2011.

33 di Sarra, A., Sferlazzo, D., Meloni, D., Anello, F., Bommarito, C., Corradini, S., De Silvestri, L., Di Iorio, T.,  
34 Monteleone, F., Pace, G., Piacentino, S. and Pugnaghi, S.: Empirical correction of multifilter rotating  
35 shadowband radiometer (MFRSR) aerosol optical depths for the aerosol forward scattering and development of a  
36 long-term integrated MFRSR-Cimel dataset at Lampedusa, *Appl. Optics*, 54, 2725-2737, 2015.

37 Drobinski, P., Ducrocq, V., Alpert, P., Anagnostou, E., Béranger, K., Borga, M., Braud, I., Chanzy, A., Davolio,  
38 S., Delrieu, G., Estournel, C., Filali Boubrahmi, N., Font, J., Grubisic, V., Gualdi, S., Homar, V., Ivančan-Picek,  
39 B., Kottmeier, C., Kotroni, V., Lagouvardos, K., Lionello, P., Llasat, M. C., Ludwig, W., Lutoff, C., Mariotti, A.,  
40 Richard, E., Romero, R., Rotunno, R., Roussot, O., Ruin, I., Somot, S., Taupier-Letage, I., Tintoré, J., Uijlenhoet,  
41 R. and Wernli, H: HyMeX: a 10-year multidisciplinary program on the Mediterranean water cycle, *Bull. Amer.*  
42 *Meteorol. Soc.*, 95, 1063-1082, 2014.

43 Dye, D. G.: Spectral composition and quanta-to-energy ratio of diffuse photosynthetically active radiation under  
44 diverse cloud conditions, *J. Geophys. Res.*, 109, d10203, doi:10.1029/2003JD004251, 2004.

45 Field, C. B., Behrenfeld, M. J., Randerson, J. T. and Falkowski, P.: Primary production of the biosphere:  
46 Integrating terrestrial and oceanic components, *Science*, 281(5374), 237–240, 1998.

47 Frouin, R. and Pinker, R. T.: Estimating Photosynthetically Active Radiation (PAR) at the Earth's surface from  
48 satellite observations, *Remote Sens. Environ.* 51, 98-107, 1995.

49 Gu, L., Baldocchi, D., Verma, S. B., Black, T. A., Vesala, T., Falge, E. M. and Dowty, P. R.: Advantages of  
50 diffuse radiation for terrestrial ecosystem productivity, *J. Geophys. Res.-Atmos.*, 107(D6),  
51 doi:10.1029/2001JD001242, 2002.

52 Hader, D. P., Lebert, M., Schuster, M., del Ciampo, L., Helbling, E. W. and McKenzie, R.: ELDONET – A  
53 decade of monitoring solar radiation on five continents, *Photochem. Photobiol.*, 83, 1348-1357, 2007.

54 Harrison, L., Michalsky, J. and Berndt, J.: Automated multifilter rotating shadow-band radiometer: an instrument  
55 for optical depth and radiation measurements, *Appl. Optics*, 33, 5118-5125, 1994.

56 Hatzianastassiou, N., Matsoukas, C., Fotiadi, A., Pavlakis, K. G., Drakakis, E., Hatzidimitriou, D. and Vardavas,  
57 I.: Global distribution of Earth's surface shortwave radiation budget, *Atmos. Chem. Phys.*, 5, 2847-2867, 2005.

58 Holben, B. N., Eck, T. F., Slutsker, I., Tanré, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y.  
59 J., Nakajima, T., Lavenu, F., Jankowiak, I. and Smirnov, A.: AERONET - A Federated Instrument Network and  
60 Data Archive for Aerosol Characterization. *Remote Sens. Environ.*, 66,1–16, 1998.

61 Jacovides, C. P., Timvios, F. S., Papaioannou, G., Asimakopoulos, D. N. and Theofilou, C. M.: Ratio of PAR to  
62 broadband solar radiation measured in Cyprus, *Agricul. Forest Meteorol.*, 121, 135-140, 2004.

63 King, M. D., Tsay, S. C., Platnick, S. E., Wang, M. and Liou, K. N.: Cloud retrieval algorithms for MODIS:  
64 Optical thickness, effective particle radius, and thermodynamic phase. MOD Theoretical Basis Document No.  
65 ATBD-MOD-05, MOD06-Cloud Product. Goddard Space Flight Center, 1997.

66 Long, C. N. and Ackerman, T. P.: Identification of clear skies from broadband pyranometer measurements and  
67 calculation of downwelling shortwave cloud effects, *J. Geophys. Res.*, 105(D12), 15609-15626, 2000.

68 Mallet, M., Dulac, F., Formenti, P., Nabat, P., Sciare, J., Roberts, G., Pelon, J., Ancellet, G., Tanré, D., Parol, F.,  
69 Denjean, C., Brogniez, G., di Sarra, A., Alados-Arboledas, L., Arndt, J., Auriol, F., Blarel, L., Bourriane, T.,  
70 Chazette, P., Chevaillier, S., Claeys, M., D'Anna, B., Derimian, Y., Desboeufs, K., Di Iorio, T., Doussin, J.-F.,  
71 Durand, P., Féron, A., Freney, E., Gaimoz, C., Goloub, P., Gómez-Amo, J. L., Granados-Muñoz, M. J., Grand,  
72 N., Hamonou, E., Jankowiak, I., Jeannot, M., Léon, J.-F., Maillé, M., Mailler, S., Meloni, D., Menut, L.,  
73 Momboisse, G., Nicolas, J., Podvin, T., Pont, V., Rea, G., Renard, J.-B., Roblou, L., Schepanski, K.,  
74 Schwarzenboeck, A., Sellegri, K., Sicard, M., Solmon, F., Somot, S., Torres, B., Totems, J., Triquet, S., Verdier,  
75 N., Verwaerde, C., Waquet, F., Wenger, J., and Zapf, P.: Overview of the Chemistry-Aerosol Mediterranean  
76 Experiment/Aerosol Direct Radiative Forcing on the Mediterranean Climate (ChArMEx/ADRIMED) summer  
77 2013 campaign, *Atmos. Chem. Phys.*, 16, 455-504, doi:10.5194/acp-16-455-2016, 2016.

78 McCree, K. J.: Test of current definitions of Photosynthetically Active Radiation against leaf photosynthesis data.  
79 *Agr. Meteorol.*, 10, 443-453, doi:10.1016/0002-1571(72)90045-3, 1972.

80 Meek, D. W., Hatfield, J. L., Howell, T. A., Idso, S. B. and Reginato, R. J.: A generalized relationship between  
81 photosynthetically active radiation and solar radiation, *Agron J.*, 76(6),  
82 doi:10.2134/agronj1984.00021962007600060018x, 1984.

83 Meloni, D., di Sarra, A., Di Iorio, T. and Fiocco, G.: Influence of the vertical profile of Saharan dust on the  
84 visible direct radiative forcing, *J. Quant. Spectrosc. Ra.*, 93, 397-413, 2005.

85 Meloni, D., di Sarra, A., Biavati, G., DeLuisi, J. J., Monteleone, F., Pace, G., Piacentino, S. and Sferlazzo, D. M.:  
86 Seasonal behavior of Saharan dust events at the Mediterranean island of Lampedusa in the period 1999–2005,  
87 *Atmos. Environ.*, 41, 3041–3056, doi:10.1016/j.atmosenv.2006.12.001, 2007.

88 Meloni, D., Junkermann, W., di Sarra, A., Cacciani, M., De Silvestri, L., Di Iorio, T., Estellés, V., Gómez-Amo,  
89 J. L., Pace, G. and Sferlazzo, D. M.: Altitude-resolved shortwave and longwave radiative effects of desert dust in  
90 the Mediterranean during the GAMARF campaign: Indications of a net daily cooling in the dust layer, *J.*  
91 *Geophys. Res.*, 120, 3386-3407, 2015.

92 Mercado, L. M., Lloyd, J., Dolman, A. J., Sitch, S. and Patino, S.: Modelling basin-wide variations in Amazon  
93 forest productivity – Part 1: Model calibration, evaluation and upscaling functions for canopy photosynthesis,  
94 *Biogeosciences*, 6, 1247–1272, 2009.

95 Mermex group: Marine ecosystems’ responses to climatic and anthropogenic forcings in the Mediterranean,  
96 *Progress Oceanogr.*, 91, 97–166, 2011

97 Min, Q.: Impacts of aerosols and clouds on forest-atmosphere carbon exchange, *J. Geophys. Res. Atmos.*, 110,  
98 D06203, 2005.

99 Mizoguchi, Y., Ohtani, Y., Aoshima, T., Hirakata, A., Yuta, S., Takanashi, S., Iwata, H. and Nakai, Y.:  
00 Comparison of the characteristics of five quantum sensors, *Bulletin of FFPRI*, 9, 113-120, 2010.

01 Monteith, J. L.: Validity of the correlation between intercepted radiation and biomass, *Agric. For. Meteorol.* 68, 213–220,  
02 1994.

03 Pace, G., Cremona, G., di Sarra, A., Meloni, D., Monteleone, F., Sferlazzo, D. and Zanini, G.: Continuous  
04 vertical profiles of temperature and humidity at Lampedusa island, *Proceed. of the 9th Intern. Symp. on*  
05 *Tropospheric Profiling*, Cimini D., Di Girolamo P., Marzano F. S., and Rizi V. (Eds.), doi:10.12898/ISTP9prc,  
06 L'Aquila, Italy, September 2012.

07 Platnick, S., King, M. D., Meyer, K. G., Wind, G., Amarasinghe, N., Marchant, B., Arnold, G. T., Zhang, Z.,  
08 Hubanks, P. A., Ridgway, B., Riedi, J.: MODIS Cloud Optical Properties: User Guide for the Collection 6 Level-  
09 2 MOD06/MYD06 Product and Associated Level-3 Datasets, 141 pp., NASA Goddard, Greenbelt, MD, USA,  
10 2015.

11 Pyrina, M., Hatzianastassiou, N., Matsoukas, C., Fotiadi, A., Papadimas, C. D., Pavlakis, K. G. and Vardavas, I.:  
12 Cloud effects on the solar and thermal radiation budgets of the Mediterranean basin, *Atmos. Res.*, 152, 14-28,  
13 2015.

14 Ross, J. and Sulev, M.: Sources of errors in measurements of PAR, *Agr. Forest. Meteorol.*, 100, 103-125, 2000.

15 Trisolino, P., di Sarra, A., Meloni, D. and Pace, G.: Determination of global and diffuse photosynthetically active  
16 radiation from a multifilter shadowband radiometer, *Appl. Optics.*, 55, 8280-8286, 2016.

17 Udo, S. O. and Aro, T. O.: New empirical relationships for determining global PAR from measurements of global  
18 solar radiation, infrared radiation or sunshine duration, *Int. J. Climatol.*, 20, 1265–1274, 2000.

19 Wang, L., Hu, B., Kisi, O., Zounemat-Kermani, M. and Gong, W.: Prediction of diffuse photosynthetically active  
20 radiation using different soft computing techniques, *Q. J. Roy. Meteor. Soc.*, 143, 2235-2244,  
21 doi:10.1002/qj.3081, 2017.

22 Yu, X., Wu, Z., Jiannng, W. and Guo, X.: Predicting daily photosynthetically active radiation from global solar  
23 radiation in the Contiguous United States, *Energ. Convers. Manage.*, 89, 71-82, 2015.

24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

40

41

42

**Table 1. Nominal central wavelengths (in nm) of the 4 band used in the PAR estimate for the two MFRSR radiometers. The FWHM bandwidth is about 10 nm for all channels.**

Band number	MFRSR #424	MFRSR #586
1	415.6	414.7
2	495.7	495.5
3	614.6	613.6
4	672.8	672.1

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66  
67  
68  
69  
70  
  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92

**Table 2. Coefficients, in  $W/m^2$ , of the linear fit between PAR and MFRSR calibrated signals for the 4-band linear model.**

<b>MFRSR s/n</b>	$c_1$	$c_2$	$c_3$	$c_4$
<b>424</b>	96.09	2.30	-28.94	271.5
<b>586</b>	240.50	-218.23	91.58	252.432

93  
94  
95  
96  
97  
98  
99  
00  
01  
02  
03  
04  
05  
06  
07  
08  
09  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19

**Table 3. Coefficients of the fit, number of data and correlation coefficient for the global and diffuse monthly PAR time series.**

	<b>a (W m<sup>-2</sup>)</b>	<b>b (W m<sup>-2</sup>)</b>	<b>φ (months)</b>	<b>n</b>	<b>r<sup>2</sup></b>
<i>Global PAR</i>	95.2	52.4	-3.3	161	0.976
<i>Diffuse PAR</i>	34.7	12.5	-3.0	161	0.798

20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

**Table 4. Coefficients of the fit, number of data and correlation coefficient for the cloud-free global and diffuse monthly PAR time series.**


	<b>a (W m<sup>-2</sup>)</b>	<b>b (W m<sup>-2</sup>)</b>	<b>φ (months)</b>	<b>n</b>	<b>r<sup>2</sup></b>
<i>Global PAR</i>	109.9	-49.2	-3.1	158	0.975
<i>Diffuse PAR</i>	26.2	12.2	-3.4	131	0.816

47  
 48  
 49  
 50  
 51  
 52  
 53  
 54  
 55  
 56  
 57  
 58  
 59  
 60  
 61  
 62  
 63  
 64  
 65  
 66  
 67  
 68  
 69  
 70  
 71

**Table 5. Coefficients of the multilinear regression and statistics between NCRE and clouds properties (clouds optical thickness, clouds fraction, and clouds top height). Percentages in brackets represent the contribution of each term of the multilinear regression to the NCRE variability.**

	<i>A</i>	<i>B</i>	<i>C</i>	<i>BIAS</i>	<i>RMSD</i>
<b>NCRE, global PAR</b>	-0.005 (41%)	-0.215 (65%)	$8.48 \times 10^{-6}$ (-6%)	0.0028	0.039
<b>NCRE, diffuse PAR</b>	-0.008 (-43%)	0.501 (88%)	$1.35 \times 10^{-4}$ (55%)	-0.105	0.209


72  
73  
74  
75  
76



77  
78  
79  
80  
81  
82  
83  
84  
85  
86

**Fig. 1. Scatterplot between 5-minute averages of PAR and PAR<sub>L</sub> for the period 19 February-25 May, 2016 at Lampedusa. The linear fit is also shown.**


87  
88  
89  
90



91  
92  
93  
94  
95  
96  
97  
98  
99  
00  
01  
02  
03  
04

**Fig. 2. Time series of monthly mean global (black lines) and diffuse (green lines) PAR for cloud-free (dashed lines) and all sky conditions (continuous lines). Fitting curves (see text) are shown by thin lines for the cloud-free time series.**


05  
06  
07  
08  
09  
10  
11  
12  
13



14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

**Fig. 3. Annual evolution of median, lower and upper quartiles, maxima and minima of global (left) and diffuse (right) monthly average PA**


31  
32  
33  
34  
35  
36



37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

**Fig. 4. Monthly statistics of the fraction of cloud-free sky between 9 and 14 UTC. Only months with at least 27 days with measurements in the period 2002-2016 are considered. Minimum, lower quartile, median, upper quartile, and maximum are indicated (from the bottom to the top) for each month. The number of monthly mean values used for each month is displayed above the top axis.**


48  
49  
50  
51  
52  
53  
54



55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68

**Fig. 5. PAR measured during 4 June, 2016, a partially cloudy day (black line). The cloud-screened data are shown as black dots, and the cloud-free fitting function is shown in red.**


69  
70  
71  
72  
73  
74  
75  
76  
77  
78



79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94

**Fig. 6. Annual evolution of global (left) and diffuse (right) cloud-free PAR; maxima, upper quartile, lower quartile, and minima of the monthly mean are displayed.**


95  
96  
97  
98  
99  
00  
01  
02



03  
04  
05  
06  
07  
08  
09  
10  
11  
12  
13  
14  
15

**Fig. 7. Global (black) and diffuse (green) monthly average PAR in all-sky (solid lines) and in cloud-free (dashed lines) conditions.**


16  
17  
18  
19  
20  
21  
22  
23  
24  
25



26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38


**Fig. 8. Time series of monthly mean cloud radiative effect for global (black) and diffuse (green) PAR, calculated as the difference between all-sky condition and cloud-free monthly mean PAR (see text). Horizontal lines are mean values of global and diffuse PAR, respectively.**

39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52




53  
54  
55  
56  
57  
58


**Fig. 9. Annual evolution of global (left) and diffuse (right) PAR CRE. Maxima, upper quartile, median, lower quartile, and minima of the monthly values are displayed.**



59




60



61

62  
63  
64  
65


**Fig. 10. Annual evolution of monthly averages of Cloud Optical Thickness (top), Cloud Fraction (middle), and Cloud Top Pressure as derived from MODIS Aqua between July 2002 and December 2016. Maximum, upper quartile, median, lower quartile, and minimum are displayed for each month.**



66

67 Fig. 11. Time series of monthly normalized cloud radiative effects for global (black) and diffuse (green)  
68 PAR, and cloud top pressure (top graph, red dashed line), cloud fraction (middle, red dashed line) and  
69 cloud optical thickness (bottom, red dashed line). Cloud properties are derived from MODIS Aqua  
70 observations.


71  
72  
73  
74  
75  
76  
77  
78



79  
80  
81  
82  
83  
84  
85  
86  
87

**Fig. 12. Scatterplot between the normalized cloud radiative effect (NCREc) calculated from the multilinear regression and the normalized cloud radiative effect (NCRE). Graphs for global (left; 151 data points) and diffuse (right; 104 data points) PAR are shown.**

88  
89  
90  
91  
92  
93  
94  
95



96  
97  
98  
99  
00

**Fig. 13. Scatterplot between deseasonalized global PAR and deseasonalized NCRE for all data (left), for the months November-March (middle), and for the months May-September (right). Least square fitting lines are plotted in red; the best fit equation, squared correlation coefficients, and number of data points are shown in each graph.**