

Interactive comment on “Non-polar organic compounds in aerosols in a typical city of Eastern China: Size distribution, gas-particle partitioning and tracer for PM_{2.5} source apportionment” by Deming Han et al.

Anonymous Referee #1

Received and published: 5 February 2018

This study analyzed the molecular composition of non-polar organic compounds (NPOCs) in PM_{2.5} and their size distributions at Jiujiang city. The target NPOCs include n-alkanes, PAHs, and hopanes. Diagnostic ratios and PMF model were applied to the compositional data to evaluate the sources and atmospheric processing of PM_{2.5}. In general, this work is well organized and written. However, I still think this work lacks novelty, and would not recommend this manuscript to be accepted for publication at Atmos. Chem. Phys., although a lot of chemical and data analysis work have been done.

[Printer-friendly version](#)

[Discussion paper](#)



General comments:

1. PMF model was utilized to apportion PM2.5 components to factors/sources. However, the author did not provide any information about the method in the manuscript or supporting information. Which version of PMF model (PMF2 or EPA PMF 5.0) was used for source apportionment? How did the author determine the factor number? How did the author deal with the missing values or measurements below detection limit? Measurement uncertainty was required for PMF input, where were these data from or how were they calculated? Are there any uncertainty analysis related to the PMF modeling? Are the PMF results valid?
2. Page 23, line 449-457. This manuscript stated that the PMFP and PMFT profiles are similar, and should be attributed to the fact that the major NPOCs are enriched in particle phase. This might not apply for factor 6, 7 and 8 (Figure 8f, g and h). The impact of G-P partitioning process will mainly affect the factors highly loaded with low molecular weight species. So the author might need to discuss the impacts of G-P partitioning on these factors.
3. From the title, it seems that the manuscript focused on the size distribution, G-P partitioning of NPOCs, and the application of NPOCs on source apportionment of PM2.5. While this study did not measure the gas-phase NPOCs, the gas-particle partitioning is only simulated basing on Pankow's theory, and could not be validated. As such, it might not be appropriate to put G-P partitioning in the title, or we can say "G-P partitioning simulation", or "the impacts of G-P partitioning on source apportionment". Size distribution was measured for NPOCs, which should be related to other parts of this manuscript. For example, does the size distribution help to explain the PMF results?
4. Diagnostic ratios of n-alkanes, PAHs and hopanes were intensively used to evaluate the sources of NPOCs in previous work. The criteria of diagnostic ratios are qualitative and confusing.

Besides the above comments, the lack of enough novelty is the main issue for this work.



The size distribution and diagnostic ratios of NPOCs in typical Chinese cities were intensively investigated (Bi et al., 2005; Zhou et al., 2005; Wang et al., 2009a, b, 2011; Duan et al., 2012). The impacts of G-P portioning of semi-volatile organic compounds (SVOCs) on PMF source apportionment have been observed and validated by Xie et al. (2013, 2014), and the method of using gas + particle phase SVOCs have been intensively applied in PMF source apportionment studies (Gao et al., 2015; Wang et al., 2016; Zhai et al., 2016). Moreover, this work did not provide any new phenomenon or viewpoints that add our knowledge on size distribution or G-P partitioning of NPOCs, or sources apportionment using NPOCs data.

References

Bi, X., Sheng, G., Peng, P.a., Chen, Y., Fu, J., 2005. Size distribution of n-alkanes and polycyclic aromatic hydrocarbons (PAHs) in urban and rural atmospheres of Guangzhou, China. *Atmospheric Environment* 39, 477-487.

Duan, J., Tan, J., Wang, S., Chai, F., He, K., Hao, J., 2012. Roadside, Urban, and Rural comparison of size distribution characteristics of PAHs and carbonaceous components of Beijing, China. *Journal of Atmospheric Chemistry* 69, 337-349.

Wang, G., Kawamura, K., Xie, M., Hu, S., Gao, S., Cao, J., An, Z., Wang, Z., 2009b. Size-distributions of n-alkanes, PAHs and hopanes and their sources in the urban, mountain and marine atmospheres over East Asia. *Atmos. Chem. Phys. J1 - ACP* 9, 8869-8882.

Wang, G., Kawamura, K., Xie, M., Hu, S., Cao, J., An, Z., Waston, J.G., Chow, J.C., 2009a. Organic Molecular Compositions and Size Distributions of Chinese Summer and Autumn Aerosols from Nanjing: Characteristic Haze Event Caused by Wheat Straw Burning. *Environmental Science & Technology* 43, 6493-6499.

Wang, G., Kawamura, K., Xie, M., Hu, S., Li, J., Zhou, B., Cao, J., An, Z., 2011. Selected water-soluble organic compounds found in size-resolved aerosols collected

[Printer-friendly version](#)

[Discussion paper](#)



from urban, mountain and marine atmospheres over East Asia. *Tellus B* 63, 371-381.

Wang, Q., Feng, Y., Huang, X., Griffith, S.M., Zhang, T., Zhang, Q., Wu, D., Yu, J.Z., 2016. Nonpolar organic compounds as PM_{2.5}. 5 source tracers: Investigation of their sources and degradation in the Pearl River Delta, China. *Journal of Geophysical Research: Atmospheres* 121.

Gao, B., Wang, X.-M., Zhao, X.-Y., Ding, X., Fu, X.-X., Zhang, Y.-L., He, Q.-F., Zhang, Z., Liu, T.-Y., Huang, Z.-Z., Chen, L.-G., Peng, Y., Guo, H., 2015. Source apportionment of atmospheric PAHs and their toxicity using PMF: Impact of gas/particle partitioning. *Atmospheric Environment* 103, 114-120.

Xie, M., Barsanti, K.C., Hannigan, M.P., Dutton, S.J., Vedal, S., 2013. Positive matrix factorization of PM_{2.5} - eliminating the effects of gas/particle partitioning of semivolatile organic compounds. *Atmospheric Chemistry and Physics* 13, 7381-7393.

Xie, M., Hannigan, M.P., Barsanti, K.C., 2014. Impact of Gas/Particle Partitioning of Semivolatile Organic Compounds on Source Apportionment with Positive Matrix Factorization. *Environmental Science & Technology* 48, 9053-9060.

Zhai, Y., Li, P., Zhu, Y., Xu, B., Peng, C., Wang, T., Li, C., Zeng, G., 2016. Source Apportionment Coupled with Gas/Particle Partitioning Theory and Risk Assessment of Polycyclic Aromatic Hydrocarbons Associated with Size-Segregated Airborne Particulate Matter. *Water, Air, & Soil Pollution* 227, 44.

Zhou, J., Wang, T., Huang, Y., Mao, T., Zhong, N., 2005. Size distribution of polycyclic aromatic hydrocarbons in urban and suburban sites of Beijing, China. *Chemosphere* 61, 792-799.

Interactive comment on *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2017-908>, 2018.

