

1 **Secondary Sulfate is Internally Mixed with Sea Spray Aerosol**
2 **and Organic Aerosol in the Winter-Spring Arctic**

3 Rachel M. Kirpes¹, Amy L. Bondy¹, Daniel Bonanno², Ryan C. Moffet², Bingbing Wang^{3,4},
4 Alexander Laskin^{3†}, Andrew P. Ault^{1,5}, Kerri A. Pratt^{1,6}

5
6
7 ¹Department of Chemistry, University of Michigan, Ann Arbor, Michigan

8 ²Department of Chemistry, University of the Pacific, Stockton, California

9 ³Environmental Molecular Sciences Laboratory, Pacific Northwest National Laboratory,
10 Richland, Washington

11 ⁴State Key Lab of Marine and Environmental Science & College of Ocean and Earth Sciences,
12 Xiamen University, Xiamen, China

13 ⁵Department of Environmental Health Sciences, University of Michigan, Ann Arbor, Michigan


14 ⁶Department of Earth & Environmental Sciences, University of Michigan, Ann Arbor, Michigan

15 ‡Current affiliation: Department of Chemistry, Purdue University, West Lafayette, Indiana

16 *Correspondence to:* Kerri A. Pratt (prattka@umich.edu), Andrew P. Ault (aulta@umich.edu)

17

18



19

20 **Figure S1:** Size-resolved number fractions of particle types for additional sample periods not
 21 shown in Figure 7. Particles were sorted into 16 log size bins.

22