

Answers to Referee#1's comments

Thanks a lot for your time and comments. Our point-to-point answers are listed below the comments in blue italic.

Comments on the manuscript "Anomalous holiday precipitation over southern China" by Zhang et al. (2018)

I am glad that the authors accepted my previous advice and revised the manuscript accordingly to some degree. I am fully satisfied with the scientific findings presented. However, I still cannot recommend its publication at this phase. There are two major issues.

First, since the authors no longer attributed precipitation changes to aerosol loading changes, why did they take so much space to talk about aerosol changes during the CSF? The rationales should be given clearly to avoid confusing or misleading readers. Currently, that part seems quite irrelevant to the major topic, especially for the Abstract part.

[Answer] We completely removed the part of aerosol changes during the CSF. Now the discussion is only about time-lag correlation between the temperature and anomalous cyclone. We just discussed a little about aerosol in the end. Because Chinese Spring Festival (CSF) is a cultural event that is only related to human beings. Many studies show that air pollution reduces significantly during the CSF holiday. The factors which induce the anomalous atmospheric circulation during the CSF holiday need further research and aerosol is the most likely factor affecting atmospheric physics during the CSF holiday.

Second, the authors should spend some time on the English writing in the next version. The manuscript now is too verbose, informal and sometimes confusing or misleading, and has many grammar errors. Below, I list a few examples and hope the authors can check and revise the whole manuscript accordingly.

[Answer] Thank you for your careful checking. All the suggestions have been taken. And we checked the whole manuscript and improved the English writing. We also used the academic English editing from American Journal Experts (<https://www.aje.com/>) and the editorial certificate is behind. In addition, before submitting the paper to Atmospheric Chemistry and Physics, we used the English language editing service from Springer Nature (<https://authorservices.springernature.com/>).

1. What's the relationship between Chinese Spring Festival (CSF), CSF holiday and Lunar New Year's Day? It could be clear for Chinese readers, but not necessarily for readers from other cultures. The authors should explain it clearly at the beginning of the Abstract, or readers may have difficulties understanding results below.

[Answer] Thank you for your suggestion. We modified the Abstract to explain Chinese Spring Festival (CSF), CSF holiday and Lunar New Year's Day.

In revision, it reads: "The Chinese Spring Festival (CSF, also known as the Chinese New Year or Lunar New Year) is the most important festival in China. Lunar New Year's Day (LNYD) is the first day of the Lunar New Year. Traditionally, the CSF holiday begins a couple of days before LNYD and ends on lantern day, lasting for approximately 2 weeks"

2. In 'this manuscript reports that during the holidayshas been significantly reduced', I think 'holidays' should be replaced with 'holiday'; 'significantly reduced' compared with what period? A better statement could be 'the precipitation during the CSF holiday over southern China tends to be lower than days before and after the holiday'.

[Answer] Taken. We also checked the whole manuscript and replaced 'holidays' to 'holiday' in the same cases.

3. The authors has a mixed use of present and past tenses in describing their results, such as 'the Δ RH showed an evident', 'the Δ RH vertical profiles displays ...', 'low cloud cover decreased ...', 'LCC also shows a...'. They had better stick to one throughout the manuscript.

[Answer] Modified. We revised the manuscript and now only use present tense in describing our results.

4. When citing papers, the authors also should pay attention to proper and consistent use of tenses. For example, Page 3, line 3: 'PM2.5 in Beijing-Tianjin-Hebei were reduced' vs. Page 3, line 19: 'there are significant negative ...'. Both are observational results published in previous studies, why use different tenses?

Page 2, line 6-9, the two adjacent sentences both start with 'Recent'. Better use a different transition word.

[Answer] All taken.

5. Page 2, line 20 'there are also discernible weekly cycles' in what aspect?

[Answer] We added 'in meteorological parameters' in this sentence.

6. Page 2, line 20-25, these sentences are too verbose. They can be shortened as 'there are also discernible weekly cycles in meteorological parameters. For example, Gong et al. (2007) found that there are robust weekly co-variations in the surface solar radiation, total cloud cover, maximum temperature, and relative humidity in China, where the weekdays tend to have lower total cloud cover and relative humidity but higher surface solar radiation and temperature than the weekends'. Try to condense your words.

[Answer] We have modified these sentences.

7. Page 2, line 26, 'Choi et al. (2008) found that the summertime cloudiness shows a bell-shaped curve between midweek and weekend'. This sentence is confusing. What parameter shows the bell shape in what way? I checked the paper, where the bell-shaped curve referred to interdecadal variations. I could not get this information or even got wrong information from the authors' saying.

[Answer] Yes, the bell-shaped curve is the description to interdecadal variations. To avoid confusing, we replaced this sentence with another paper's results.

In revision, it reads:" In Korea, Kim et al. (2009) found that there are more cloudiness and less insolation for Wednesday-Thursday and less cloudiness and more insolation for Monday-Tuesday. Furthermore, weekly periodicities are enhanced especially in autumn, more than 2–3 times as great as those of the annual mean with long-term surface measurements of meteorology (1975–2005)."

Reference:

Kim, B.-G., Choi, M.-H., and Ho, C.-H.: Weekly periodicities of meteorological variables and their possible association with aerosols in Korea, Atmospheric Environment, 43(38): 6058-6065, doi: 10.1016/j.atmosenv.2009.08.023, 2009.

8. Page 3, line7: 'for example, on' → such as

[Answer] Done.

9. Page 3, line 10: 'The emission may differ among these events' differ in what aspect? The sentence tells me nothing.

[Answer] We modified the sentence to 'The emission may differ in the magnitude and type among these events'.

10. Remove all 'etc.' from the manuscript. It is too informal for a scientific paper.

[Answer] We checked the whole manuscript and removed all 'etc'.

11. Page 4, the last line, 'Lunar New Year's Days (LNYD)', LNYD→LNYDs; replace all 'New Year's Day' afterward with 'LNYD'

[Answer] All taken.

12. Page 5, line 14, everyday → every day.

[Answer] Done.

13. Page 5, line 17, 'whose lunar date is' → 'whose lunar dates are'; 'and calculated the average' grammar error

[Answer] Modified.

13. Page 16, line 6, this sentence should go into the introduction section.

[Answer] We think you may be referring to this sentence 'The CSF is a cultural tradition that is directly related to the daily activity of human beings, particularly economic activities.' in Page 15, line 6.

We have moved this sentence to the fourth paragraph of the introduction section.

I could not check errors and verbose sentences throughout the paper and do not ensure no other errors in places between what I list above. The authors should do it and revise the manuscript accordingly.

[Answer] Thank you very much and we have checked the whole manuscript.



EDITORIAL CERTIFICATE

This document certifies that the manuscript listed below was edited for proper English language, grammar, punctuation, spelling, and overall style by one or more of the highly qualified native English speaking editors at American Journal Experts.

Manuscript title:

Anomalous holiday precipitation over southern China

Authors:

Jiahui Zhang, Dao-Yi Gong, Rui Mao, Jing Yang, Ziyin Zhang, Yun Qian

Date Issued:

October 9, 2018

Certificate Verification Key:

EF64-5711-A073-12CF-172P



This certificate may be verified at www.aje.com/certificate. This document certifies that the manuscript listed above was edited for proper English language, grammar, punctuation, spelling, and overall style by one or more of the highly qualified native English speaking editors at American Journal Experts. Neither the research content nor the authors' intentions were altered in any way during the editing process. Documents receiving this certification should be English-ready for publication; however, the author has the ability to accept or reject our suggestions and changes. To verify the final AJE edited version, please visit our verification page. If you have any questions or concerns about this edited document, please contact American Journal Experts at support@aje.com.

Nature Research Editing Service Certification

This is to certify that the manuscript titled Anomalous holiday precipitation over the southern China was edited for English language usage, grammar, spelling and punctuation by one or more native English-speaking editors at Nature Research Editing Service. The editors focused on correcting improper language and rephrasing awkward sentences, using their scientific training to point out passages that were confusing or vague. Every effort has been made to ensure that neither the research content nor the authors' intentions were altered in any way during the editing process.

Documents receiving this certification should be English-ready for publication; however, please note that the author has the ability to accept or reject our suggestions and changes. To verify the final edited version, please visit our verification page. If you have any questions or concerns over this edited document, please contact Nature Research Editing Service at support@as.springernature.com.

Manuscript title: Anomalous holiday precipitation over the southern China

Authors: Jiahui Zhang, Dao-Yi Gong, Rui Mao, Jing Yang, Ziyin Zhang, Yun Qian

Key: 1C72-7131-81E9-AB98-A43P

This certificate may be verified at secure.authorservices.springernature.com/certificate/verify.

Nature Research Editing Service is a service from Springer Nature, one of the world's leading research, educational and professional publishers. We have been a reliable provider of high-quality editing since 2008.

Nature Research Editing Service comprises a network of more than 900 language editors with a range of academic backgrounds. All our language editors are native English speakers and must meet strict selection criteria. We require that each editor has completed or is completing a Masters, Ph.D. or M.D. qualification, is affiliated with a top US university or research institute, and has undergone substantial editing training. To ensure we can meet the needs of researchers in a broad range of fields, we continually recruit editors to represent growing and new disciplines.

Uploaded manuscripts are reviewed by an editor with a relevant academic background. Our senior editors also quality-assess each edited manuscript before it is returned to the author to ensure that our high standards are maintained.

Anomalous holiday precipitation over southern China

Jiahui Zhang¹, Dao-Yi Gong¹, Rui Mao¹, Jing Yang¹, Ziyin Zhang², Yun Qian³

¹State Key Laboratory of Earth Surface Process and Resource Ecology/Academy of Disaster Reduction and Emergency Management, Beijing Normal University, Beijing, 100875, China

5 ²Environmental Meteorology Forecast Center of Beijing-Tianjin-Hebei, China Meteorological Administration, Beijing, 100081, China

³Pacific Northwest National Laboratory, Climate Physics Group, Washington, 99352, USA

Correspondence to: Dao-Yi Gong (gdy@bnu.edu.cn)

Abstract: The Chinese Spring Festival (CSF, also known as the Chinese New Year or Lunar New Year) is the most important festival in China. Lunar New Year's Day (LNYD) is the first day of the Lunar New Year. Traditionally, the CSF holiday begins a couple of days before LNYD and ends on lantern day, lasting for approximately 2 weeks. In this manuscript, Officially, this holiday lasts approximately one week. Based on the long-term station observations from 1979 to 2012, this manuscript reports that during the holidays, the precipitation during the holiday over southern China (108°E-123°E and 21°N-33°N, 155 stations) tends to be lower than that before and after the holiday has been significantly reduced. The mean precipitation frequency anomalies from the fourth day to the sixth day after Lunar New Year's Day LNYD (i.e., days [+4, +6]) were found to decrease by -7.4%. At the same time Simultaneously, the daily precipitation amounts experienced a reduction of -0.62 mm d⁻¹ during days [+2, +5]. The holiday precipitation anomalies are strongly linked to the relative humidity (ΔRH) and cloud cover. The station observations of the ΔRH showed an evident decrease from day +2 to day +7, and a minimum appeared on days [+4, +6], with a mean of -3.9%. The ΔRH vertical profile displays a significant drying below approximately 800 hPa. Between 800 hPa and 1000 hPa, the mean ΔRH is -3.9%. The observed station daytime low cloud cover (LCC) evidently decreased by -6.1% during days [+4, +6]. Meanwhile, the ERA-Interim daily LCC also shows a comparable reduction of -5.0%. The anomalous relative humidity is mainly caused by the decreased water vapor in the lower-middle troposphere. Evident negative specific humidity anomalies persist from day -3 to day +7 in the station observations. The average specific humidity anomaly for days [+4, +6] is -0.73 g kg⁻¹. When the precipitation days are excluded, the anomaly remains significant at being -0.46 g kg⁻¹. A significant deficit of water vapor vapour deficit is observed in the lower troposphere below 700 hPa. Between 800 hPa and 1000 hPa, the mean specific humidity dropped by -0.70 g kg⁻¹. This drier lower-middle troposphere is due to anomalous northerly winds, which are tightly closely related to the cyclonic circulation anomaly over the northwestern Pacific. The time-lag correlation demonstrates that approximately one week after a lower temperature occurs over eastern China, a stronger cyclone is observed over the western Pacific. Authors have analyzed the concurrent holiday aerosol anomaly and their time lag correlations with the atmospheric circulation. Their strongest correlations occur when the PM_{2.5} leads by 9 to 6 days, with correlation coefficients varying from +0.52 to +0.57. And it is found that the East Asian aerosol concentrations during the CSF decrease evidently, the largest reduction occurring

带格式的: 突出显示

on days [-3, -1]. At the same time, a concurrent cooling is observed in the lower troposphere, which enhances the cold advection over East Asia. The strength of the anomalous cyclone on days [+4,+6] is statistically correlated with PM₁₀ concentration over East China. Their strongest correlations occur when the PM₁₀ leads by -9 to -6 days, with correlation coefficients varying from +0.52 to +0.57. The possible mechanism needs further clarification throughby elaborate observation analysis and numerical modeling.

带格式的: 突出显示

1 Introduction

Human activities have important impacts on weather/climate at diverse spatial/temporal scales by causing changes in Earth's atmosphere in the amounts of greenhouse gases, aerosols and cloudiness in the Earth's atmosphere. The changes of these factors alter the energy balance of the Earth-atmosphere system. In particular, Particularly the large uncertainty involving the clouds, precipitation, and radiation makes it difficult to accurately estimate the contributions of these factors to the long-term climate changes (Boucher et al., 2013). Recent estimation indicates that in the fast-developing China, anthropogenic emissions may have contributed 10%±4% of the current global radiative forcing (Li et al., 2016a). Many relatedRecent studies have noted that short-term air pollution associated with intense economical/political/cultural events has discernible influences on local and/or regional weather worldwide, where air pollution caused by regular or occasional anthropogenic emissions significantly impacts atmospheric physics and chemistry (e.g., Zhao et al., 2006; Sanchez-Lorenzo et al., 2012; Williams et al., 2015). The involved cloud-precipitation processes and interactions over monsoonal Asia remains among the most challenging topics (Li et al., 2016b).

Alternating weekdays and weekends may be the most apparent regular activities of human beings. The so-called weekly cycle effects of atmospheric parameters, which are strongly related to the aforementioned regular human activity, have been widely reported globally (Gordon, 1994; Cervený and Balling, 1998; B äumer et al., 2008). At first, these studies of the weekly cycles were focused on cities and industrialized areas. Later, it was reportedly, that the weekly cycles of atmospheric variables are not only local phenomena confined within urban areas but are also large-scale phenomena that are closely related to regular human activities (Gordon, 1994). Over developed Europe, for example, B äumer et al. (2008) analyzed the station data from the Aerosol Robotic Network over central Europe and reported that the aerosol optical thickness is higher midweek than on the weekend. In the rapidly industrializing Asia, there are also discernible weekly cycles in meteorological parameters. For example, Gong et al. (20062007) found that the weekend effect in the diurnal temperature range (DTR) has the opposite signal between winter and summer, where wintertime DTR tends to have a positive weekend effect (i.e., larger DTR on weekend days compared to weekdays), which is in association with increased maximum temperature and total irradiance but decreased relative humidity, whereas summertime DTR displays a much stronger and significantly negative weekend effect. the meteorological parameters in China experience significant weekly cycles. There are robust co-variations

带格式的: 突出显示

~~in the surface solar radiation, total cloud cover, maximum temperature, and relative humidity. During the weekdays, the total cloud cover and relative humidity are lower, while the surface solar radiation and maximum temperature are higher. In contrast, the higher cloud cover and humidity and the lower radiation and maximum temperature appear on the weekends.~~ Evident weekly cycles are also identified in other parameters, such as wind speed, precipitation frequency (Gong et al., 2007)
5 and the diurnal temperature range (Gong et al., 2006) and thunderstorm frequency (Yang et al., 2016). In Korea, Kim et al. (2009) found that there are more cloudiness and less insolation for Wednesday-Thursday and less cloudiness and more insolation for Monday-Tuesday. Furthermore, weekly periodicities are enhanced especially in autumn, more than 2–3 times as great as those of the annual mean with long-term surface measurements of meteorology (1975–2005).
10 ~~In South Korea, Choi et al. (2008) found that the summertime cloudiness shows a bell-shaped curve between midweek and weekend.~~ The mechanisms of these weekly cycles are not yet fully understood, though their strong co-variations with human activities (e.g., air pollutants) are widely measured widely. For Gong et al. (2006; 2007) supposed that the phases of weekly cycle may depend on the different climatological circumstances. Wang et al. (2012) emphasized that during the autumn, the semi-direct aerosol effects likely play a dominant role in the weekly cycle of cloud cover in southeastern China. Yang et al. (2016) proposed the importance of ~~the~~ human emission types.

15 In addition, there are occasional, intense human activities (e.g., societal and cultural events) ~~that, which~~ also change atmospheric environments (particularly the air quality). For example, Xin et al. (2012) found that the concentrations of PM_{2.5} in Beijing-Tianjin-Hebei area are were reduced by approximately 50% with the implementation of emission controls during the 2008 Beijing summer Olympic Games ~~in the summer of 2008~~. Similar actions were also implemented when China hosted the 2014 Asia-Pacific Economic Cooperation meeting ~~in 2014~~ and held the 2015 ~~held the~~ national military parade ~~in~~
20 2015; as a result, the air quality is was considerably improved over northern China (Chen et al., 2015; Wang et al., 2015; Xu et al., 2015; Wen et al., 2016; Xu et al., 2017). In contrast, setting off fireworks during holiday celebrations can produce severe air pollution around ~~eross~~ the world; ~~such as for example, on~~ the Fourth of July holiday in America (Carranza et al., 2001), the New Year's fireworks in Mainz (Drewnick et al., 2006), ~~during~~ the Diwali festival in India (Singh et al., 2014) and, during the Chinese Spring Festival (CSF) (Tang et al., 2016), ~~etc.~~ The emissions may differ in magnitude and type
25 among these events (Wang et al., 2007; Zhang et al., 2010; Tang et al., 2016). Comparing with weekly cycles, the possible meteorological co-variations in association with these occasional, intense events are less studied.

Notably, ~~e that~~ the CSF holiday repeatedly occurs annually every year in January-February across the whole country and is the most important holiday in China. Moreover, the CSF is a cultural tradition that is directly related to the daily activity of human beings, particularly economic activities. ~~H~~The human emissions during the CSF holidays decreases evidently (Lin and McElroy 2011; Gong et al., 2014). Some studies have tried to identify the possible covariations-changes in atmospheric physics and weather anomalies accompanying with these local and regional occasional society/cultural events. Travis et al. (2002) noted a significant increase in DTR. As an example, in the United States, the diurnal temperature range increased significantly during the three days after 11 September 2001 when civil aviation planes were grounded. ~~(Travis et al., 2002).~~

带格式的: 非突出显示

Gong et al. (2014) analyzed the temperature anomalies during the CSF holidays ~~from~~during 2001-2012 and found that there are significant negative anomalies of approximately -0.81°C in eastern China, wherein atmospheric feedbacks likely play a role in enhancing ~~the~~ cooling.

The purpose of the present study is to investigate the possible changes ~~of~~ precipitation over southern China during the CSF holidays. The paper is organized as follows. The data and methods are presented in Section 2. ~~The r~~Results, including the anomalous precipitation frequency and amount, the relevant changes in humidity, cloud cover and temperature, and the anomalous atmospheric water vapor and circulation, are described in Section 3. The possible relevance of these results to ~~the human activity~~holiday aerosols is discussed in Section 4. ~~Finally~~In addition, a summary is presented in Section 5 ~~presents a summary~~.

2 Data and methods

2.1 Data and study area

The daily precipitation data analyzed in the present study ~~are~~were obtained from the China Meteorological Administration observation archives. The daily precipitation amounts as well as the precipitation types are recorded. There are six kinds of readings, namely, (a) no precipitation; (b) trace precipitation (daily precipitation readings of less than the measurable 0.1 mm); (c) precipitation from fog, frost and dew; (d) precipitation from pure snow; (e) precipitation from snow and rain; and (f) precipitation from rainfall. Because the frequency of snowfall events is low in the warm area of southern China and the precipitation from fog and dew is considerably small (accounting for 0.14% of our data), in this study, we considered only the precipitation amounts and ignored the precipitation types. This dataset consists of 917 stations, and the earliest observations start in the 1950s. Here, we ~~confined~~our analysis period to 1979-2012, during which China experienced ~~a~~rapid development and ~~immense amounts~~massive of air pollutant emissions ~~of air~~ pollutants. Since we aim to investigate the precipitation around the ~~CSF holiday~~Lunar New Year's Day, in this study, only the daily precipitation from January to February is employed. After excluding the 319 stations with missing values, we retained the 598 stations without missing data during 1979-2012.

Prior to this analysis, we ~~have to~~determined the target area based on the precipitation climatology. Because the Chinese Lunar New Year occurs in January or February, here, we investigated the climatological means of the precipitation amounts as well as the precipitation ~~frequency~~days based on the observations from 598 stations. As shown in Figure 1a, in the boreal winter, the number of precipitation days is much greater in southern China than in ~~northern China~~the north. A maximum of >35 days occurs between approximately $25\text{-}30^{\circ}\text{N}$ and $102\text{-}115^{\circ}\text{E}$, while the precipitation frequency ~~north of~~ 35°N is less than 15 days ~~north of~~ 35°N . ~~Simultaneously~~At the same time, the major precipitation belt, with precipitation amounts > 50 mm, also appears over the southeast of mainland China, ~~which is~~being generally consistent with the spatial features of the frequency patterns (Figure 1a). Although the number of precipitation days in the Sichuan Basin is high, the

precipitation amounts are not ~~assuch~~ large. In addition, ~~the~~ Sichuan Basin has a unique climate due to its special basin topography. Therefore, we did not include ~~the Sichuan Basin~~ in our study area. In addition, for 319 stations north of 33°N, the average precipitation days and amounts are 12.8 days and 9.9 mm, respectively. ~~It would be difficult to obtain a statistically meaningful signal in northern China due to a small precipitation frequency and amount. Too small precipitation frequency and amount in northern China would be difficult to statistically yield a meaningful signal.~~ Finally, we confined our study area to 108°E-123°E and 21°N-33°N (as indicated by the black dashed rectangle in Figure 1). Within this area, there are 155 available meteorological stations. Among these stations, the minimum number of precipitation days (15.7 days) appeared at the Zhumadian station (33°N, 114°E), and the minimum precipitation amount (15.3 mm) appeared at the Ankang station (32°N, 109°E). Meanwhile, the maximum precipitation days (41.6 days) and maximum precipitation amount (214.5 mm) ~~arewere~~ found at Wugang (26°N, 110°E) and Nanyue (27°N, 112°E), respectively. On average, the precipitation frequencies and amounts during the period of 1979-2012 are 30.3 days and 124 mm, respectively. In addition to the daily precipitation, we also analyzed other meteorological variables, including the mean temperature and relative humidity.

The atmospheric circulation and moisture in the troposphere ~~arewere~~ taken from the ERA-Interim reanalysis datasets (Dee et al., 2011), which have a 1.5° spatial resolution and span the period of 1979-2012. We utilized surface parameters, such as the cloud cover and total column water vapor. In addition, the impacts of ~~the~~ air temperature, relative humidity, and horizontal winds on pressure levels ~~arewere~~ also investigated. Only the data at step 0 for 00:00, 06:00, 12:00, and 18:00 UTC ~~arewere~~ obtained and subjected to analysis. Averaging these 4 times yields the daily mean.

2.2 Analysis method

We employed a temporal composite analysis to investigate the possible changes in the daily precipitation and the statistical significance. This ~~analysis iswas~~ carried out via the following steps. First, we identified all Lunar New Year's Days (LNYDs) from 1979 to 2012. As listed in Table 1, the ~~New Year's Days~~LNYDs all fall within late January to mid-February. During our analysis period, the earliest ~~New Year's Day~~LNYD is on 22 January, and the latest occurs on 20 February. These LNYDs are denoted as day 0. The days before and the day after the LNYD are denoted as day -1 and day +1, respectively. Accordingly, two days before and after the LNYD are denoted as day -2 and day +2, ~~and so onete~~.

Second, we estimated the anomalies from the climate means. When dealing with continuous variables (such as the temperature, humidity, atmospheric circulation, and water vapor), we calculated the long-term means for each calendar day from 1 January to 28 February. Then, the means ~~arewere~~ subtracted from the daily observations to determine the corresponding anomalies. This process is ~~the~~ same as that used in Gong et al. (2014). In contrast to these continuous variables, precipitation occurs ~~quite~~ randomly. For a specific day, ~~such assay~~ day 0, we counted the number of precipitation days (N) during the ~~entirewhole~~ data period from 1979-2012. Thus, the precipitation frequency (hereafter denoted ~~to~~ as F) is deduced by:

$$F = \frac{N}{34} \times 100\%, \quad (1)$$

To remove the randomness of precipitation, we used a 3-day window. For example, precipitation occurring on the previous day (day -1) and the day after (day +1) is considered to occur on day 0. The above method is also used to calculate the climatic values of F_i which are based on Gregorian calendar days. For a station from 1st January to 31st March, every day has a precipitation frequency. Then, with this climatic sequence, we used a 7-day window filter to reserve the changes for more than 7 days because the typical synoptic time scale is approximately 7 days. For a station, on a specific day, such as day 0, all precipitation frequencies whose lunar dates are day 0 were found from the climatic sequence. Then, we calculated their average. The departure of F from the average is the anomalous frequency anomaly (ΔF). The final result is the average of all stations' anomalies.

3 Results

3.1 Significant reduction of precipitation

We computed the daily anomalies of the precipitation frequencies for each of the stations. To determine how the precipitation might change before and after New Year's Day LNYD, we considered days from -12 to +12. This method is similar to that of the work of Gong et al. (2014). We also tried to use a longer time window and found that the conclusions remain the same. To focus on southern China, we computed the means for all 155 stations within the target area (Figure 1). The regional mean ΔF is shown in Figure 2a. Although the ΔF before the LNYD is generally positive, the salient feature is the ΔF reduction after the LNYD. The mean for day +3 to day +7 is -5.5%, and a minimum of -9.4% occurs on day +5. Note that the largest reductions occur on days [+4, +6], where the averaged anomalies are as low as -7.4%. Precipitation is a natural phenomenon. Thus, its frequency anomaly should be spatiotemporally random for a certain day from 1979-2012 among all the stations over this large region. The standard deviation of the stations' ΔF would provide information concerning further uncertainties. A smaller standard deviation suggests that similar anomalies tend to be observed, thus indicating a robust signal, whereas a larger standard deviation implies diverse changes. We found that the standard deviation of the days [+3, +7] is approximately $\pm 3.7\%$, which is being smaller than the standard deviation of most days before the LNYD. Note that the means may be biased by outliers or skewness. Thus, to determine the diverseness of these frequency anomalies among the 155 stations, we also computed the medians and the lower and upper percentiles. In Figure 2a, the ranges of the 10th and 90th percentiles are plotted as the error bars and are shown with the medians. The 90th percentiles for the days [+4, +6] are clearly -0.9%, -3.9%, and -1.7% respectively, all being significantly below 0. At the same time, Simultaneously, the medians are generally similar to the means; for example, from days [+3 to, day +7], the means (-3.1%, -6.5%, -9.4%, -6.3%, and -2.2%, respectively) and medians (-3.6%, -7.0%, -9.8%, -7.0%, and -1.8%, respectively) are almost identical, likely suggesting that the majority of the anomalies tend to be negative and that the means are not skewed by large departures or outliers.

We further investigated the statistical significances of the precipitation anomalies by employing a Monte-Carlo approach (e.g., Stjern, 2011; Wang et al., 2012). Here, the Monte-Carlo test ~~is~~was performed by randomly rearranging ~~the~~ sequences of the lunar calendar days. For each experiment, we first generated the random sequence and then estimated the frequencies for each lunar day station by station. In addition, the regional anomaly mean ~~was~~is calculated as described above. We repeated these steps 1000 times and obtained the 10th and 90th percentiles of these experiments. By comparing ~~the~~ observations with these Monte-Carlo-yielded percentiles, we ~~may~~ight estimate whether the frequency anomalies are beyond those of random chance. If the original observations are lower than the 10th percentile, these measurements are significant at the 0.1 level.

For the regional mean frequency anomalies, there are few cases where the simulated anomaly is smaller than the observations during the holidays. Only 1 of 1000 random Monte-Carlo experiments is below the observation on day +5. On days +4 and +6, there are 10 and 14 cases, ~~respectively~~, with values smaller than the observation. On days +3 and +7, there are slightly more ~~values smaller than the observation~~ (135 and 190, respectively). We found that the 10th percentiles during days +4 to +6 are -3.6%, -3.4% and -3.5%, respectively. Note that the observations (-6.5%, -9.4% and -6.3%) are all below the corresponding 10th percentiles of the Monte-Carlo experiments. Therefore, we may conclude that the observed ΔF is significant at the 0.1 level from days +4 to ~~day~~+6. We also examined the medians of the 155 stations and found a similar result. ~~For~~om days +4, +5 and ~~to~~+6, there are only 9, 1 and 7 cases, ~~respectively~~, with values below the observations. The observed medians during days +4, +5 and ~~to~~+6 are -7.0%, -9.8% and -7.0%, ~~respectively~~, which are all below the Monte-Carlo 10th percentiles (-3.7%, -3.6% and -3.6%, respectively).

~~Further, w~~We ~~further~~ investigated the possible changes of daily precipitation amounts. Here, all trace precipitation records are excluded. The composite method is similar to that for frequency. In other words, for a specific lunar calendar day, the anomalous amount is calculated as the mean daily precipitation amount minus the climate reference, where the climate reference is estimated based on the Gregorian calendar. Then, we averaged the anomalies for all stations to obtain a regional mean. Figure 2b shows the results. Interestingly, from day +2 to day +5, the amounts experience continuously negative anomalies with a mean of -0.62 mm d⁻¹. In particular, a minimum of -1.0 mm d⁻¹ occurs on day +3. During days +2, +4, and +5, the anomalies are similar (being -0.54, 0.52, and -0.48 mm d⁻¹, respectively). Differing from the precipitation frequency, the amounts ~~rang~~inges from the 10th-90th percentiles are obviously larger. In addition, during days [+2, +5], the upper bounds all exceed zero and imply larger uncertainties. Thus, the gradual decrease in the amount after the LNYD is likely consistent with the significant reduction in precipitation frequency. Note that the Monte-Carlo test suggests a high confidence of the negative amount anomalies during the holidays. Of 1000 Monte-Carlo experiments, there are 83, 14, 75 and 70 cases with values smaller than the observations during days [+2, +5]. The Monte-Carlo 10th percentiles for days [+2, +5] are -0.47, -0.45, -0.42 and -0.40 mm d⁻¹, respectively, whereas the corresponding observations of -0.54, -0.95, -0.52 and -0.48 mm d⁻¹ are all smaller. We also examined the median of the 155 stations by employing the Monte-Carlo test. The results ~~are~~is similar. There are 70, 16, 66 and 43 experimental values that are smaller than those observed. The medians of the observation for these days (-0.63, -0.97, -0.65 and -0.70 mm d⁻¹, respectively) are all smaller than the corresponding Monte-Carlo 10th

percentiles (-0.56, -0.54, -0.52 and -0.47, respectively). Therefore, the means and medians of the amount anomalies during days [+2, +5] are significant at the 0.1 level. Note that the large error bars in Figure 2b may imply somewhat larger diversities/differences of the daily precipitation amounts among the 155 stations. The Monte-Carlo test likely provides a more reasonable estimation of the significance. With the above analysis, we may conclude that there is a significant reduction in the precipitation frequency and a decrease in the daily precipitation amount after the LNYD.

3.2 Spatial distribution of the holiday precipitation anomalies

Although the precipitation reduction after the LNYD is evident in the composite analysis based on the regional means (Figure 2), this feature might differ among stations. In this subsection, we investigated the spatial distribution of the precipitation anomalies. A significant frequency reduction occurs on days [+4, +6], and a significant amount reduction appears on days [+2, +5]. Here, we computed the mean frequency anomalies during days [+4, +6] and the mean amount anomalies during days [+2, +5] for each station. The statistical significance is also estimated using the Monte-Carlo test at each station. The method used is similar to that described in Section 3.1. Here, the Monte-Carlo test iswas performed for every station by randomly rearranging the lunar calendar, and the simulation iswas repeated 1000 times. The 10th (90th) and 20th (80th) percentiles arewere taken as the criteria. For the precipitation frequency, the mean value is regarded as significant at the 0.1 (or the 0.2) level only when the anomalies in all three days (i.e., days +4, +5 and +6) arewere significant at the 0.1 (or 0.2) level. Similarly, when the daily precipitation amounts on all four individual days (from day +2 to day +5) are significant, their mean is regarded as a significant anomaly. The results are shown in Figure 3.

Figure 3a shows that the majority of the ΔF have negative signs. Among the 155 stations, only 7 stations have positive signs, and none of these positive anomalies are significant. Of the 148 stations with ΔF reductions, 57 stations are significant at the 0.1 level, and 108 are significant at the 0.2 level. Most stations located in the north and east of study area have atthe significant ΔF reduction. The ΔF minimum reaches -13.5% at the Sansui station (27°N, 108°E). ConsideringLooking at the anomaly's magnitude of the anomaly, there are 55 stations with ΔF reductions exceeding -9%. Among these 55 stations, the anomalies for 45 stations are significant at the 0.1 level, and those at the other 10 stations are significant at the 0.2 level.

Unlike the frequency, the spatial features of the amount anomalies differ among stations (Figure 3b), which is consistent with the relatively larger span of the 10th-90th percentiles, as shown in Figure 2b. Nevertheless, there are still 117 stations with decreased amounts during days [+2, +5]. Among these stations, 27-stations showed significant decreases significant at the 0.2 level, and 8 stations showed those-significant decreases at the 0.1 level. The largest reduction is -2.41 mm d⁻¹, which appearsed at the Shouxian station (32°N, 116°E).

Based on the above analysis of the temporal composites and spatial distributions, we found that the-significant reductions of precipitation frequency and daily precipitation amount occur regionally within an approximate one-week period after the LNYD.

3.3 Relative humidity and cloud anomalies

Relative humidity (RH) is the most important factor that directly controls/influences precipitation. In this subsection, we investigated the possible changes in the relative humidities/humidity associated with the holiday precipitation decreases. Here, we examined the station data to analyze the relative humidity by making composites. Note that the method of estimating the anomalies is different from that for precipitation frequencies. The composites of the relative humidity (as well as those of other meteorological parameters analyzed in the following sections) are based on anomalies (Δ RH), which are obtained by subtracting the 1979-2012 long-term mean from the daily relative humidity values.

For each lunar calendar day, all 34 daily anomalies are collected and subjected to a t -test with the null hypothesis of their mean being not significantly different from zero at the 0.05 (or 0.1) level. In Figure 4, we plotted the means during days [-12, +12]. To facilitate a comparison, the range of one standard error estimated from the 34 anomalies is shown via error bars. The Δ RH experiences an evident decrease from day +2 to +7, with a mean value of -2.5%. The driest value is -4.6% at day +5. The mean Δ RH of the lowest anomalies for days [+4, +6] is -3.9%, which is significant at the 0.1 level as estimated from a left-tail t -test. Note that the drying (wetting) may be the result of less (more) precipitation, as demonstrated in the previous sections. For clarity, we also computed the Δ RH using only the data from no-rain days. We found that the Δ RH for no-rain cases shows a similar temporal feature (figure not shown). The lowest values still occur on days [+4, +6], with anomalies of -0.97%, -0.69% and -0.38%, respectively. However, these anomalies are not statistically significant. This result could be caused by the relatively smaller data sample and exclusion of the precipitation days. The sample numbers of no rain days is 48.9% of that of all days. In any case, the similar drying features on no-rain days may provide support for the theory that the lower Δ RH could be a cause of the precipitation reduction and may even help enhance the drying.

We also examined the spatial distribution of the Δ RH. The station means of the Δ RH during days [+4, +6] are plotted in Figure 5. All 155 stations show negative anomalies, with a maximum and a minimum of -0.6% and -8.1%, respectively. When estimating the statistical significance, we considered the mean for days [+4, +6]. Thus, we did not consider the significance of each of the three days. If the resulting anomaly is significant at the 0.05 (or 0.1) level as calculated by a t -test, this station is regarded as significant during days [+4, +6]. As shown in Figure 5, there are 70 (98) stations that are significant at the 0.05 (0.1) level. Additionally, we investigated the spatial features of no-rain days and found that most Δ RH values during days [+4, +6] are also negative (the number of station days is 136, accounting for approximately 88% of the total stations), and the lowest anomaly is -7.6% (figure not shown). Again, this result suggests a drying atmosphere near the surface during the New Year's holiday.

For the precipitation of stratus precipitation during the winter season, the relative humidity in the middle-lower-middle troposphere is more important than the surface humidity. The precipitation reduction could occur with a drier and upper atmosphere. Then, we then investigated the Δ RH in the lower-middle level troposphere using the ERA-Interim reanalysis datasets of the pressure levels from 1979 to 2012. Here, we selected 99 grid points between 21°N-33°N and 108°E-123°E. We examined the Δ RH values at each level from 1000 hPa to 500 hPa. Generally, the results are similar, showing drying

tendencies throughout, as expected. Note that the Δ RH values in the low levels are more evident. At 1000 hPa, the Δ RH ~~shows~~ a continuous reduction during days [+3, +6], with a mean decrease of -3.8%. The negative anomalies on these days are significant at the 0.1 level. The largest reduction occurs on day +5, being -5.2% (figure not shown). In Figure 4, we ~~plotted~~ the regional mean Δ RH at 850 hPa. A similar reduction is observed during days [+2, +6], where the lowest anomalies are still found on days [+4, +6]. In addition, the values on days +4 and +5 (-4.0% and -5.7%) are both significant at the 0.1 level. In the middle-upper troposphere, the Δ RH during the holidays is not evident.

The spatial distributions of Δ RH at 1000 and 850 hPa during days [+4, +6] ~~were~~ are also analyzed. At 1000 hPa, the majority of the significantly negative Δ RH values ~~were~~ are located over southern China and the neighboring western Pacific, with a northeast-southwest distribution. The results at 850 hPa are plotted in Figure 5, and only the significant values (above the 0.1 level) are shown. At 850 hPa, the significant region is ~~spatially smaller~~ spatially but has a greater magnitude than that at 1000 hPa. Over the land, the majority of the anomalies are between -4% and -6% at 1000 hPa, whereas the values are between -6% and -10% at 850 hPa. The minimum grid point value at 1000 hPa is -6.5%, while the minimum is as low as -9.7% at 850 hPa.

The drying in the lower troposphere can be more clearly seen in the relative humidity profiles. The vertical profile of the Δ RH during days +4 to +6, as averaged over all 99 grids in southern China, is shown in Figure 6. Because the meaningful changes occurred in the lower troposphere, here we plotted only the Δ RH values below the 500 hPa level. Below 500 hPa, the anomalies all have negative signs. In addition, the significant Δ RH values appear below approximately the 800 hPa level (Figure 6a). The negative anomaly at 850 hPa is -3.9%, and that at 1000 hPa is -4.2%. The mean for these layers (below 800 hPa) is -3.9%. Generally, the significant drying of the relative humidity appears in the ~~lower~~ middle ~~troposphere level~~, which physically agrees with the precipitation reduction.

As the relative humidity at the ~~lower~~ middle level decreases, consistent changes in clouds should occur, particularly in the low-level cloud coverage (LCC). To confirm this, we first analyzed the surface-observed cloud data, which are from the Global Surface Weather Dataset obtained from the China Meteorological Administration. In this dataset, the observations are taken 4 times per day (00:00, 06:00, 12:00, 18:00 UTC) during the years 1980-2012, except ~~for~~ during 2000, which is not available. Only those days with all 4 records are included. Of the 304 stations in our study area, we selected 137 stations with overall data availabilities above 30%. When deriving the anomalies, we applied the same method as that used for Δ RH. The results show that both the total cloud cover and LCC ~~reduces~~, especially the LCC experiences a significant decrease during the New Year's holiday. It seems the reduction of total cloud cover is contributed by the decrease of ~~low-cloud cover~~ LCC. The mean anomaly of LCC during days [+1, +5] is -2.6%. In addition, a minimum of -3.1% appears on day +5, which is significant at the 0.1 level. We should note that sometimes the LCC cannot be completely distinguished from the middle cloud cover in the station observations. Moreover, the cloud cover changes rapidly and might depend on the observer. At nighttime, the uncertainty of the observations is much greater. We would expect more reliable observations during the daytime. The mean LCC of 00:00 and 06:00 UTC is computed and regarded as the daytime observation. As

shown in Figure 4b, the daytime LCC shows an evident reduction during days [+4, +6]. The values of ~~each of the~~ three days are ~~each of~~ significant at the 0.1 level, as estimated using a left-tail *t*-test. The mean for the three days is -6.1%, and a minimum of -9.7% occurs on day +5. Compared to the climate daytime cloud cover of 78% during January to March in southern China, the LCC reduction during the New Year's holiday has a considerably large magnitude. Figure 5b shows the spatial distribution of the daytime LCC values during days [+4, +6]. For a total of 137 stations, 119 (86.9%) stations showed negative signs. In addition, there are 8 (27) stations with values significant at the 0.05 (0.1) level, as estimated via a both-tail *t*-test. A minimum of -16.8% occurs at the Fuyang station (33°N, 115°E).

To assess the robustness of the cloud cover changes, the ~~cloud cover of the~~ ERA-Interim data ~~about cloud cover are~~ was investigated. We found that the high cloud cover and ~~the~~ middle cloud cover showed no significant changes (figure not shown). Interestingly, there is a significant reduction in the LCC during the ~~CSF~~ New Year's holidays. Figure 4d shows the temporal variations of the ERA-Interim LCC. The LCC clearly shows an outstanding reduction during days [+4, +6]. The largest reduction, i.e., -5.9%, appears on day +5. The values from day +4 to day +5 are all significant at the 0.1 level, as calculated by a left-tail *t*-test. Averaging over days [+4, +6], the mean value is -5.0%. This magnitude is comparable to the station-based daytime LCC anomaly (-6.1%). The spatial distribution of the LCC anomalies for the ERA-Interim data during days [+4, +6] is shown in Figure 5d. The LCC reduction center is located in the eastern region and has a magnitude between approximately -8% and -12%. This reduction is almost identical to that of the station observations (c.f., Figure 5b, d). Note that in the ERA-Interim data, the cloud height is defined according to the corresponding sigma level. The low clouds correspond to sigma 0.8-1.0. A significant decrease of the relative humidity occurs below 800 hPa (Figure 6a), which physically agrees with the negative LCC anomalies.

20 3.4 Temperature and water vapor anomalies

The negative ΔRH during the holidays might be caused by anomalous temperatures, ~~by~~ the water vapor or both, depending on certain conditions. In this subsection, we analyzed the changes of the temperature and water vapor associated with the holiday precipitation reduction. First, we examined the temporal changes of the temperature over southern China using the same composite analysis as that used for the relative humidity. We found that there is ~~a~~ continuous cooling of the daily mean temperatures from day -3 to day +6, which are all significant at the 0.1 level, as estimated by a left-tail *t*-test (figure not shown). The mean temperature anomaly of these days is -1.12°C. The maximum and minimum temperature anomalies are -0.70°C and -1.42°C for days +6 and +3, respectively. Precipitation often causes a low temperature. To exclude the possible influence of precipitation on temperature, we also examined the daily temperatures for no-rain days and found a similar temperature anomaly of -1.22°C for days +1 to +6. In addition, there is a similar minimum of -1.48°C. These values for the no-rain days are significant at the 0.05 level.

Note that Gong et al. (2014) ~~have~~ reported negative temperature anomalies over the whole of eastern China around the LNYD when analyzing the shorter data period of 2001-2012. They found that the most significant decreases appears on days

-3 to +2, with a mean value of -0.81°C . However, the temperature beyond this period over the southern China, however, ~~is~~ was not addressed in their analysis. As demonstrated in the previous sections, the most significant reductions of the ΔF and ΔRH are observed on days [+4, +6]. Thus, we further computed the mean of the temperature anomalies (ΔT) for days [+4, +6] and estimated their statistical significance using a t -test for each of the stations. The results show that all 155 stations have negative anomalies (figure not shown). The maximum ΔT of -0.28°C occurs at the Fengjie station (31°N , 109°E), and a minimum of -2.14°C appears at the Jiuxian station (25°N , 118°E). The anomalies for 116 stations are statistically significant at the 0.1 level, and those for 95 stations are significant at the 0.05 level. We also investigated the temperature anomalies during days [+4, +6] using no-rain days and found similar results. All stations continue to show negative anomalies, and more stations are significant (134 stations exceed the 0.1 level, and 117 stations exceed the 0.05 level).

The above spatial distribution suggests that temperature anomalies are unlikely at the local scale. For clarity, we also investigated the vertical profiles and the spatial distributions of ΔT using the ERA-Interim pressure level data. The mean ΔT values below the 500 hPa level during days [+4, +6] are averaged for all 99 grid points over southern China and plotted in Figure 6b. The significant negative temperature anomalies below 700 hPa are evident, where all values are as low as $< -1^{\circ}\text{C}$. The mean for these layers is -1.37°C . To determine whether the cooling of the lower-middle troposphere is a regional-scale phenomenon, we further analyzed the spatial distribution of ΔT . Note that the minimum of -1.56°C occurs at the 850 hPa level in Figure 6b. Here, we computed the 850 hPa ΔT during days [+4,+6] for each of the grid points (figure not shown), revealing that the majority of the study area experiences a significant cooling (exceeding -1°C) that is almost identical to that from the surface station observations. Based on these analyses, we may conclude that during the holidays (particularly, days [+4, +6]), there is an anomalous temperature cooling over southern China from the surface to the middle troposphere (below 500 hPa).

Cooling favors condensation and precipitation. If the water vapor content remains a constant, a cooler temperature should result in a higher ΔRH . During the period of 1979-2012, over southern China, the climatic mean temperatures for days [+4, +6] at 1000 hPa and 850 hPa are 9.3°C and 3.3°C , respectively. Meanwhile, the specific humidities at 1000 hPa and 850 hPa are 5.5 g kg^{-1} and 4.4 g kg^{-1} , respectively. According to the Clausius-Clapeyron equation (Murray, 1967), the climatological relative humidities are, respectively, 75.4% at 1000 hPa and 77.5% at 850 hPa. The observed coolings of -1.15°C at 1000 hPa and -1.56°C at 850 hPa (Figure 6b) could result in corresponding ΔRH increases of +6.0% and +9.1%, respectively. This contradicts the observed negative ΔRH anomalies (c.f., Figure 5). Therefore, the cooling temperature is not a direct factor causing the drier ΔRH and the precipitation reduction.

Alternatively, the water vapor should be responsible for the anomalous ΔRH . We analyzed the station specific humidity (SH). Here, the SH is estimated from the surface pressure, relative humidity and temperature, where the Tetens formula is employed to estimate the saturated vapor pressure over water (Murray, 1967). Figure 7a shows the specific humidity anomalies (ΔSH) before and after the LNYD. Evident negative anomalies persist from days -3 to +7. The lowest values appear at approximately days [+4, +5]. Note that the negative ΔSH s during days +2 to +6 are significant at the 0.05 level,

with a mean of -0.71 g kg^{-1} . A minimum of -0.82 g kg^{-1} occurs on day +4. In addition, the average ΔSH for days [+4, +6] is -0.73 g kg^{-1} . We repeated the composite analyses for no-rain days and found that the ΔSH has ~~ve the~~ similar continuous reductions after ~~LNYD~~~~New Year's Day~~ (Figure 7b). The ΔSH s of days [+2, +5] are significant at the 0.1 level, and their mean is -0.43 g kg^{-1} . The minimum of -0.54 g kg^{-1} also appears on day +4, while the mean from day +4 to +6 is -0.46 g kg^{-1} , being smaller than the values estimated for all days. ~~Simultaneously~~~~At the same time~~, the spatial distribution of ΔSH during days [+4, +6] reveals a regional-scale reduction over the study area (Figure 8). All stations clearly show negative anomalies, both when all days are analyzed and when only no-rain days are analyzed. In the former instance, the ΔSH values for the 140 stations ~~are~~~~is~~ statistically significant at the 0.05 level, and in the latter case, the number of stations with statistically significant values is 102.

~~Further, we~~~~further~~ investigated the ΔSH values in the ~~lower~~~~middle~~ troposphere using the ERA-Interim reanalysis data. The ΔSH at 1000 hPa (850 hPa) during days [+2, +6] are all significant at the 0.05 (0.1) level, with a mean of -0.75 (-0.65) g kg^{-1} . It is clear that the ΔSH s at 1000 hPa and 850 hPa display features similar to those of the surface observations in both ~~their~~ temporal variations and ~~the~~ magnitudes of ~~the~~ negative anomalies during the holidays (c.f., Figure 7). Furthermore, we investigated the spatial distributions of ΔSH on days [+4, +6]. The evident ΔSH reduction covers ~~almost the whole~~~~nearly~~ all of southern China, with minimums extending from southeastern China to the western North Pacific, south of approximately 30°N (Figure 8). The anomaly center lies between 110°E - 130°E and 20°N - 30°N .

The drying of the ΔSH in the ~~lower~~~~middle~~ troposphere is more obvious, as seen in the vertical profile (Figure 6c). ~~The~~ significant negative ΔSH values appear ~~in the levels~~ below 700 hPa. Below 800 hPa, the ΔSH values are all ~~less than~~~~below~~ -0.50 g kg^{-1} . The mean for these layers (800 hPa to 1000 hPa) is -0.70 g kg^{-1} . Similarly, we also estimated the variations of the relative humidity corresponding to these ΔSH s. The ΔSH s at 1000 hPa and 850 hPa are -0.78 g kg^{-1} and -0.69 g kg^{-1} , respectively (Figure 6c). If the temperature remains unchanged, these values would reduce the relative humidity by -10.9% and -12.3%, respectively. Obviously, the ~~reduction of the~~ water vapor ~~reduction~~ in the lower-middle troposphere plays the dominant role in causing anomalous relative humidity~~yies~~, low-level clouds and precipitations.

The total column water vapor (TCWV), or the precipitable water, is also a large-scale factor that correlates well with precipitation (e.g., Qian et al., 2009). The drying from the surface to the mid-troposphere is suggestive of the reduction in the TCWV. For clarification, we performed a composite analysis of the ERA-Interim TCWV. Unsurprisingly, the ΔTCWV experiences a continuous reduction from day +1 to day +5 (Figure 9a). The ΔTCWV are significant at the 0.05 level during days [+2, +5], with a mean of -1.93 kg m^{-2} . In addition, the largest negative ΔTCWV values are observed on days +4 and +5 (2.43 kg m^{-2} and 2.36 kg m^{-2} , respectively). The spatial distribution of the ΔTCWV values during days [+4, +6] is shown in Figure 9b. The TCWV decreases by approximately -1.00 kg m^{-2} or more over most of ~~the~~ southern China. The minimum is located over the neighboring western North Pacific, between 120°E - 130°E and 20°N - 30°N . Meanwhile, positive ΔTCWV s with somewhat smaller magnitudes appear to the east (over approximately 150°E - 175°E and 20°N - 30°N). Generally, the reduction in the TCWV over the western North Pacific and eastern China is physically consistent with the negative

anomalies of ~~the~~ SH over southern China. The observed decrease in precipitable water during the holidays is unfavorable for precipitation.

Based on the above analysis, we may conclude that, although the relative humidity (as well as the relevant cloud cover and precipitation) would benefit from the cooling temperatures, its reduction during days [+4, +6] is strongly dominated by the drying water vapor in the lower troposphere. Therefore, the reduction of ~~the~~ SH is likely to contribute to the decrease of precipitation during the New Year's holiday.

3.5 Water vapor budget and atmospheric circulation

Our analysis demonstrates that in association with the significant precipitation reduction over southern China, an anomalous negative departure of the SH occurs. One further question is ~~that~~ what causes such water vapor deficits during the New Year's holiday? Here, we discuss ed the possible factors relevant to the atmospheric column water vapor balance as well as their individual contributions. Regional changes of the column water vapor are essentially related to two components. One is the budget between evaporation and precipitation, and the other is the budget of moisture inflow and outflow from the horizontal air motion crossing the four lateral boundaries. We investigated ed the corresponding anomalies separately during the holidays.

First, we computed the differences between the evaporation and ~~the~~ total precipitation (evaporation minus precipitation, i.e., E-P) using the ERA-Interim data from 00:00 and 12:00 UTC. Note that both evaporation and precipitation are forecasted, and the accumulated variables are collected at step 12. We analyzed the E-P anomaly (mm, equivalent to kg m^{-2}) during the holidays (figure not shown). From days +1 to +5, the E-P anomalies are all positive values with a mean of 0.85 mm. In addition, the anomalies on days +3, +4 and +5 are all significant at the 0.05 level. The maximum of the E-P anomaly (+1.25 mm) appears on day +4. We also investigated the evaporation independently and found it that the evaporation is enhanced during the holidays, with the mean of the evaporation anomalies reaching +0.39 mm (figure not shown). Notably, ~~We should note that~~ the positive E-P anomalies occur concurrently with the precipitation reductions. The E-P anomalies are likely dominated by precipitation changes. Although the positive E-P anomaly implies a net gain, this effect cannot account for the total net loss of the column water vapor.

Second, we investigated the column water vapor budget contributed by the horizontal transport. Here, we simply estimated the moisture flux across the borders (i.e., the box in Figure 1). The vertically integrated moisture transport (Q) crossing the border is defined as follows:

$$Q = \int_L \left[\frac{1}{g} \int_{500\text{hPa}}^{1000\text{hPa}} q \cdot V dp \right] dl_{\perp} \quad (2)$$

where V is the horizontal wind vector, g is the gravitational acceleration, q is the SH and L is the length of the border.

Moisture transport above 500 hPa is ignored because its contribution to the total column moisture transport is relatively small, and large winds tend to cause errors in the upper troposphere, where the SH is quite small (Qian et al., 2009). Along the northern and southern borders, we computed the meridional transport, while along the eastern and western borders, we

computed only the zonal transport. The summation of the meridional and zonal transports yields the net budget. The results shows ~~that~~ the net water budgets are negative from day +1 to day +5 (figure not shown), with a mean of $-2.9 \times 10^7 \text{ kg s}^{-1}$. Note that the anomalies of days +1, +3 and +4 are all significant at the 0.05 level. The lowest water vapor budget is $-4.4 \times 10^7 \text{ kg s}^{-1}$ (~~approximately~~ $-1.96 \text{ kg m}^{-2} \text{ day}^{-1}$), occurring on day +4. This simple calculation suggests that the horizontal moisture transport dominates the anomalous variations of the total net water vapor budget.

Because the negative anomaly in the water vapor budget may be caused by a reduced inflowing water vapor transport or by an increased outflowing transport, we investigated the transport crossing of each of the boundaries during the holidays. We found that the zonal transports crossing the eastern and western borders are negligible (figure not shown). In contrast, the water vapor transport at the southern boundary has the greatest magnitude (Figure 9). The southern border shows continuous negative anomalies from day +1 to day +5, and the largest departure ~~of~~ $-5.5 \times 10^7 \text{ kg s}^{-1}$; occurs on day +4. The anomalies during days +2 to +5 are significant at the 0.05 level and have a mean of $-4.4 \times 10^7 \text{ kg s}^{-1}$. Thus, the negative transport crossing the southern border implies an enhanced outflow, and is largely responsible for the net loss of the column water vapor during ~~the~~ holidays.

The water vapor transport over this large area should be closely related to the regional atmospheric circulation. Previous studies have emphasized the importance of ~~the~~ anomalous atmospheric circulation over the western Pacific in modulating ~~the~~ precipitation over southern China by influencing ~~the~~ moisture transport and convergence (e.g., He et al. 2007; Li et al., 2013; among others). To elaborate on the details of the regional atmospheric circulation around the New Year's holiday, we investigated the tropospheric horizontal winds over eastern Asia. The vertical mean horizontal winds averaged from 1000 hPa to 700 hPa are plotted in Figure 9b. Over southeastern China, there are ~~clear~~-dominant northerly wind anomalies ~~clearly~~. The northerly winds flow toward the east over the western Pacific, south of approximately 25°N , and then, ~~the winds~~ turn northward. Consistent with this cyclonic circulation anomaly, eastern China and the neighboring western Pacific experience remarkable reductions in the column water vapor. This pattern remains stable when ~~looking at~~ considering each ~~of the~~ pressure levels in the mid-troposphere. For example, at the 850 hPa level, the anomalous cyclone is more evident, and the strong northerly winds over southeastern China are greater than the vertical means (Figure 9c). Interestingly, the precipitation reduction in ~~the~~ southern China is well captured in the ERA-Interim reanalysis, and such ~~a~~ reduction is physically consistent with the large-scale atmospheric circulation change. We repeated the composite analysis of the atmospheric circulation using the NCEP/NCAR Reanalysis I and NCEP-DOE Reanalysis II datasets and found a similar pattern (figures not shown). Employing the same ERA-Interim data but a shorter data length (2001-2012), Gong et al. (2014) also reported a similarly anomalous circulation pattern over East Asia around the Chinese New Year. This anomalous cyclone is likely a robust signal of the holiday weather. Generally, this anomalous cyclone plays a dominant role in bringing stronger northerly wind, causing ~~less~~drier humidityes and lower water vapor ~~values~~, ~~which~~ finally results ~~ing~~ in less precipitation over southeastern China.

4 Discussion: time-lag correlation between the temperature and anomalous cyclone on co-variations in holiday air pollution

4.1 Air pollution reduction over eastern China

The CSF is a cultural tradition that is directly related to the daily activity of human beings, particularly economic activities. The concurrent phenomena observed in atmospheric physics and weather likely imply links between these systems. We suspect the holiday anthropogenic emission may act as an intermediate connection. Air pollutants were widely reported to considerably decrease during this holiday. For example, Tan et al. (2009) investigated the 1994–2006 observations from 13 monitoring sites around the Taipei metropolitan area and found that the concentrations of NO_x , CO, nonmethane hydrocarbon, SO_2 and PM_{10} were lower during the New Year period than during the non New Year periods. Using three different approaches considering the thermal power generation, satellite retrieval products and statistical and modeling attributing, Lin and McElroy (2011) estimated that the economic slowdown during the celebrations of CSF was responsible for a notable reduction in anthropogenic emission. According to their estimations for 2005, 2007–2008 and 2010, the CSF contributes a NO_x emission reduction of 12%. In addition, the 2009 CSF contributed a reduction of 10%. These estimations are comparable to those given in the work of Gong et al., (2014), in which they reported a PM_{10} reduction derived from 323 surface station measurements over eastern China and estimated the magnitude of the PM_{10} concentration decrease to be approximately 9.24% for the days -4 to +5 (excluding day 0 to rule out the New Year's Eve firework emissions).

Note that the boreal winter is the season when the air pollution concentration reaches its peak over China. Thus, we investigated the climatic mean and holiday changes of the aerosol optical depth (AOD). Here, the AOD data were obtained from the Moderate Resolution Imaging Spectroradiometer (MODIS) data sets produced by using the AOD 550 Dark Target Deep Blue algorithm and combined by the MOD08_D3 and MYD08_D3 product (Platnick et al., 2015a; Platnick et al., 2015b). These daily level 3 gridded atmospheric data are available from 2002 to 2012 at a 1-degree spatial resolution. As shown in Figure 1, the climatic mean AOD during January and February for eastern China is remarkably greater than those of the surrounding areas and has two maximum centers, one appearing to the north (i.e., the northern plains) and the other appearing to the west (namely, over Sichuan Basin). Concurrently, a considerably high AOD appears over southern China, ranging from 0.4 to 0.8. For comparison, the spatial distribution of PM_{10} concentrations in January–February is also provided. Generally, the PM_{10} concentration in the north is higher than that in the south. However, the pollution in southern China is also severe in the winter.

To better understand the aerosol changes during the holidays, we simply investigated the frequency distribution of the AOD for days [-1, +5]. During this period, a significant net water vapor reduction is observed. Because frequent cloudy weather causes a smaller clear-day AOD sample, it is difficult to calculate a meaningful daily AOD anomaly. Alternatively, we randomly selected days [-15, -11] to compute the preholiday period AOD frequency distribution for comparison. We also tested other time period, like days [-12, -8], and the result is robust. The frequencies for the holiday and preholiday periods were both calculated based on the original AOD data using a bin width of 0.2 (Figure 10). The ambient atmosphere during

带格式的: 突出显示

the holidays is much cleaner than that of the preholiday period. On average, the AOD means for days [-15, -11] and days [+1, +5] are 0.67 and 0.58 for southern China, respectively. During the holidays, the magnitude of AOD reduction is shown to be approximately 13.4%. The improvement of air quality is well demonstrated by the frequency changes shown in Figure 10. The frequencies of the cases in which $AOD > 0.6$ are 50.4% and 36.6% for days [-15, -11] and days [+1, +5], respectively. This result shows a decrease of 13.8%. Note that the largest frequency reduction (approximately 6.7%) occurs in the bin of 0.6–0.8. Additionally, the daily PM_{10} data were also analyzed for comparison and were calculated from the daily air pollution index (API) (Gong et al., 2007). There are 113 stations in our study area. Similar to the AOD, the PM_{10} frequency distribution was calculated with a bin width of $20 \mu\text{g m}^{-3}$ for both the holidays and pre-holidays. The frequency of PM_{10} concentrations greater than $100 \mu\text{g m}^{-3}$ during the holiday period is 19.8%, while the frequency for the same bin during the preholiday period is 41.7%. There is a 21.9% reduction in the high PM_{10} concentration cases during the CSF holidays. Not only do the frequencies change but the mean PM_{10} concentrations also evidently decrease. The mean PM_{10} concentrations for days [-15, -11] and days [+1, +5] are $98.9 \mu\text{g m}^{-3}$ and $71.5 \mu\text{g m}^{-3}$, respectively.

In addition to southern China, eastern Asia shows a marked drop in air pollution concentrations (figure not shown). We investigated the mean AOD and PM_{10} values of eastern China (108°E–123°E and 21°N–42°N). The AOD values decrease from 0.60 during days [-15, -11] to 0.58 during days [+1, +5]. The frequency of cases with $AOD > 0.6$ is 33.3% during the holiday period, while the frequency for the preholiday period is 38.1%. Additionally, the mean PM_{10} concentrations for days [-15, -11] and days [+1, +5] are $112.7 \mu\text{g m}^{-3}$ and $85.2 \mu\text{g m}^{-3}$, respectively. The reduction in high pollution cases is notable. The proportion of $PM_{10} > 100 \mu\text{g m}^{-3}$ records declined from 47.9% to 30.1%. The simultaneous and anomalous variations of the pollution concentrations of PM_{10} in southern and eastern China can be clearly seen in their temporal composites, as shown in Figure 10c. The significant negative anomalies around the LNYD clearly appear in both eastern and southern China, and their lowest departures both occur a couple of days before New Year's Day. For eastern China, the mean value for days [-4, -1] is $-12.26 \mu\text{g m}^{-3}$, which is approximately 14.71% smaller than the climatic background concentration ($83.35 \mu\text{g m}^{-3}$). In southern China, the mean of the anomalies during days [-4, -1] is $-9.07 \mu\text{g m}^{-3}$, being 12.63% lower than that of the background ($71.84 \mu\text{g m}^{-3}$). These robust AOD and PM_{10} signals clearly demonstrate that around the LNYD, eastern China experiences a relatively cleaner ambient atmospheric environment although the climatic background air pollution level in January and February is the highest of all seasons.

Note the PM_{10} composites might be biased by their trends and outliers. To address this question, we repeated the analysis based on yearly normalized data. Here all PM_{10} data are rescaled according to maximum minus minimum range year by year. The normalized data show that the frequency of PM_{10} greater than 0.2 during days [-15, -11] is 44.1% while the frequency for the same bin during days [+1, +5] is 23.9%. The mean value for days [-4, -1] is 16% lower than climate mean. This agrees well with above analysis.

We further investigated the radiation from MERRA2. Here we employed two MERRA2 products, one runs with assimilation aerosols and the other without (Gelaro et al., 2017). We analysed the longwave radiation at top of atmosphere by making composites same as that for temperature. Results are shown in Figure 11. It is clear that significant negative

anomalies appear on days [-4, -1] and [+3, +6] in both the MERRA2 products with/without assimilation aerosols. The means for these two short periods are respectively -1.77 and -1.57 Wm^{-2} in no aerosol MERRA2 dataset. Meanwhile, the mean longwave radiation anomalies are -1.79 and -1.58 Wm^{-2} in MERRA2 with aerosols. ERA Interim has similar results (Figure 11). The analysed radiation datasets in MERRA2 and ERA Interim are both under clear sky condition. Top of atmosphere longwave radiation reflects equivalent temperature of air column, smaller top of atmosphere upwelling longwave radiation means a cooler beneath atmosphere. Top of atmosphere net radiation anomalies during holidays are also negative due to longwave radiation (figure not shown). The negative longwave and net radiation at top of atmosphere and temperature after the New Year's Day would be related to the cooling in association with the anomalous cyclone, whereas the simultaneous influence of atmospheric circulation on the cooling and negative longwave/net radiation in days [-4, -1] should be small since there is no significantly anomalous circulation. Generally, the radiation anomalies seem reasonably consistent with the observed temperature changes.

4.2 Time lag correlation between PM_{10} concentration and the anomalous cyclone

There are increasing numbers of works addressing the climate consequences and feedbacks of human air pollutions with focus on precipitation. For example, Zhao et al. (2006) analyzed 40 years' observations of precipitation, satellite aerosol data and meteorological sounding data over eastern China and suggested a positive feedback mechanism (i.e., more aerosols cause less precipitation, which in turn leads to more aerosols) that will induce an acceleration of the reduction of precipitation in eastern China. Particularly, the observed changes of light rains are likely related to aerosols. Qian et al. (2009) employed long term observational data from 1956-2005 and found that both the frequency and amount of light rain have decreased over eastern China. Their numerical simulations suggest that the light precipitation tends to be suppressed due to a higher cloud droplet number condensation and smaller droplet sizes under polluted conditions. Some studies suggested that aerosols induce surface cooling and weaken land-sea thermal contrast, thereby weakening monsoon circulations and the resultant precipitation changes (e.g., Wang et al., 2013; Ye et al., 2013; Guo et al., 2013; among others). Nevertheless, we note a few studies addressed the synoptic scale circulation anomalies. For example, Jones et al. (2004) suggested the dust aerosol heating in northern Africa plays an important role in changing the amplitudes of the downstream easterly waves over a couple of days. Ding et al. (2013) reported that the pollution caused by agricultural burning and fossil fuel combustion in some provinces over East China results in significant modification of regional daily temperature and precipitation through pollution boundary layer dynamics and aerosol radiation cloud feedbacks. Numerical experiments with cloud microphysical processes show that Asian pollution invigorates weather activities along the North Pacific storm tracks (Zhang et al., 2007; Wang et al., 2014).

Note that in the observations, the holiday precipitation reduction is strongly related to the significant drying of the SH, which is caused by anomalous northerly winds. In fact, an anomalous cyclone dominates East Asia and the western Pacific region during the CSF holidays (Figure 9). The anomalous northerly winds over southern China are just located in the rear side of the anomalous cyclonic circulation. The anomalous cyclone is likely correlated to the temperature cooling during the CSF

带格式的: 缩进: 首行缩进: 0 字符

holiday. Gong et al. (2014) reported that during days [-4, 0], there is no anomalous cyclone over East Asia, but simultaneously, the temperature cools significantly. These phenomena can be observed in both the long-term (1979-2012) and short-term (2001-2012) data periods. The anomalous cyclone appears in the troposphere after LNYD, moves eastward, and disappears after approximately 12 days. This result is highly consistent with our analysis, as demonstrated in Figure 9.

5 On days [-3, -1], the greatest temperature cooling in eastern China was observed (c.f., Gong et al., 2014, Figures 2 and 4). As shown in Figure 10, the lower-middle troposphere shows no evident northerly wind anomaly over East Asia or the western Pacific before LNYD. However, there is significant cooling in the lower-middle troposphere, as indicated by the negative 1000-500 hPa thickness anomalies during days [-4, 0]. The cooling center is located between 100°E-130°E and 30°N-40°N. In this case, the negative temperature anomalies can result in anomalous cold advection due to the climatic
10 northerly winds. This condition is helpful for constructive baroclinic interaction between the upper and lower troposphere and favors the midlatitude cyclone system (Hakim et al., 2003). We would expect a mature cyclone anomaly approximately one week later if the cyclone develops from days [-3, -1].

We suppose that air pollution reduction around the holiday period. With respect to this issues, two facts are likely indicative of their relationship.

15 Firstly, there is a time lag between the PM₁₀ reduction and the anomalous cyclone. Gong et al. (2014) reported that during days [-4, 0], there is no anomalous cyclone over East Asia, but at the same time, the temperature cools significantly. These phenomena can be observed in both the long term (1979-2012) and short term (2001-2012) data periods. The anomalous cyclone appears after New Year's Day in the troposphere, moves eastward, and disappears after approximately 12 days. This result is highly consistent with our analysis, as demonstrated in Figure 9. The largest aerosol reduction occurs
20 on days [-3, -1]. At the same time, the greatest temperature cooling in eastern China was observed (c.f., Gong et al., 2014, figures 2 and 4). As shown in Figure 12, the lower middle troposphere shows no evident northerly wind anomaly over East Asia and the western Pacific before New Year's Day. However, there is a significant cooling in the lower middle troposphere, as indicated by the negative 1000-500 hPa thickness anomalies during days [-4, 0]. The cooling center is located between 100°E-130°E and 30°N-40°N. In this case, the negative temperature anomalies can result in anomalous cold
25 advection due to the climatic northerly winds. This condition is helpful for constructive baroclinic interaction between the upper and lower troposphere and favouring midlatitude cyclone system (Hakim et al., 2003). We would expect a mature cyclone anomaly approximate one week later, if it is true that the cyclone develops from days [-3, -1].

Secondly, the intensity of the cyclone is significantly correlated with the temperature cooling preceding air pollution. We computed the time-lag correlation between the holiday cyclone and the regional mean temperature. Because the
30 anomalous cyclone is most robust during days [+4, +6], we selected the corresponding center (140°E-145°E and 25°N-35°N) to calculate a regional mean for the 500 hPa height. In eastern China, there are 394 available meteorological stations. We averaged these stations' the daily temperature anomalies of these stations to derive a regional mean time series and then computed their correlation with that of the 500 hPa height on day +5. To suppress the noise and identify a stable signal, we used a 3-day window when computing the means, i.e., the data on a specific day as well as those of the day before and after

带格式的: 非突出显示

带格式的: 突出显示

are averaged. The results are plotted in Figure 11; note that a time-lag of 0 refers to day +5, a time-lag of -2 refers to day +3, and so on. The results show that the cyclone is closely related to the temperature anomalies that occurred approximately 5 days previously. A maximum correlation of 0.52 was observed when temperature leads the pressure by 4 to 5 days. And it is statistically significant at the 0.1 level. These significant positive correlations suggest that when the temperature is lower around LNYD, the intensity of the anomalous cyclone on days [+4, +6] over the western Pacific is likely to be stronger. Thus, the anomalous cyclone is significantly related to the continental temperature that occurred approximately one week previously. The regional temperature over southern China experiences a similar cross-correlation with the following cyclone (Figure 11). We also computed the cross-correlations using the 31 PM₁₀ collocate stations. The features are almost identical (Figure 13). All these correlations are statistically significant at the 0.1 level. It should be pointed out that these time-lag correlations should not be explained by the natural 1-2 week processes. The natural synoptic system's occurrence and phases are random in time. Here, we prepared the atmospheric correlation/temperature time series according to the lunar calendar dates. If there is a natural cyclone around the CSF holiday, its random phase should be offset by other cyclones when all years are put together. Anyway, these results demonstrate that a stronger cyclone over the western Pacific on days [+4, +6] is often accompanied by lower preceding temperature that occurs approximately one week previously over eastern China.

CSF is a cultural event that is only related to human beings. Many studies show that air pollution reduces significantly during the CSF holiday. For example, Tan et al. (2009) investigated the 1994-2006 observations from 13 monitoring sites around the Taipei metropolitan area and found that the concentrations of NO_x, CO, nonmethane hydrocarbon, SO₂ and PM₁₀ are lower during the New Year period than during the non-New Year periods. Using three different approaches considering the thermal power generation, satellite retrieval products and statistical and modeling attribution, Lin and McElroy (2011) estimated that the economic slowdown during the celebrations of CSF is responsible for a notable reduction in anthropogenic emissions. According to their estimations for 2005, 2007, 2008 and 2010, the CSF contributes a NO_x emission reduction of 12%. In addition, the 2009 CSF contributes a reduction of 10%. These estimations are comparable to those given in the work of Gong et al. (2014), in which they reported a PM₁₀ reduction derived from 323 surface station measurements over eastern China and estimated the magnitude of the PM₁₀ concentration decrease to be approximately -9.24% for the day -4 to day +5 (excluding day 0 to rule out the New Year's Eve firework emissions). Aerosol is the most likely factor affecting atmospheric physics during the CSF holiday.

In addition, it should be noted that the northerly winds are helpful in cooling the temperature. We think that when considering the simultaneous temperature changes in association with short period aerosol anomalies, the atmospheric feedback would be ignorable. However, during a moderate period (such as >3 days to one week), the atmospheric feedback is likely discernible. The mechanism about how the concurrent aerosol reduction influences temperature and further affects circulation needs more elaborate observation and model simulation.

PM₁₀ concentrations. Because the anomalous cyclone is most robust during days [+4, +6], we selected its corresponding center (140°E-145°E and 25°N-35°N) to calculate a regional mean for the 500 hPa height. Then, we computed its correlation

带格式的: 突出显示

with the daily mean PM_{10} anomalies over eastern China for each of day around the LNYD. The daily PM_{10} data were taken from Gong et al. (2014). Note that the PM_{10} observations are highly variable. To reduce the possible uncertainty of the correlation, we employed only those PM_{10} stations with >70% of the daily data available during the period from 2001–2012. To suppress the noise and identify a stable signal, we further used a 3-day window when computing the means, i.e., the data on a specific day as well as those of the day before and after are averaged. On New Year's Eve, there is a sharp PM_{10} peak due to intense firework burning (Figure 10c). Therefore, the data on New Year's Day are intentionally excluded. The results are plotted in Figure 13; note that a time lag of 0 refers to day +5, a time lag of -2 refers to day +3, etc. The simultaneous correlation between the regional mean PM_{10} and the cyclonic pressure is quite weak (only 0.2). Interestingly, their correlations increase substantially as the PM_{10} leading time grows, until approximate 10 days. The strongest correlations occur when the PM_{10} leads by -9 to -6 days, with correlation coefficients varying from +0.52 to +0.57. The maximum appears for a time lag of -9. It reveals the correlation is the best when there is a 9-day time lag between PM_{10} and circulation during holidays. Such correlation between the year-to-year variation of circulation and PM_{10} reflects that year-to-year variation has a fixed time lag of 9 days, not a relationship on the synoptic scale. All these correlations are statistically significant at the 0.1 level. As indicated previously, around the holidays, the PM_{10} experiences similar changes over eastern China. Such a time lag relationship should be found for the sub-regional PM_{10} data. We repeated the cross-correlation analysis using only the mean PM_{10} values over southern China. The largest correlations also occur when the PM_{10} values lead by 6 to 9 days, as expected. Note that 9-day and a 6-day leading times correspond to days -4 and -1, respectively. During these days, there are remarkable PM_{10} reductions (Figure 10c). Their significant positive correlations suggest that when the PM_{10} concentration is lower during days [-4, -1], the intensity of the anomalous cyclone on days [+4, +6] over the western Pacific is likely to be stronger. Thus, the anomalous cyclone is significantly related to the continental PM_{10} concentration that occurred approximate one week previously.

We further computed the cross-correlations of the regional mean temperatures with the anomalous cyclone on day +5. In eastern China, there are 394 available meteorological stations. We averaged these stations' daily temperature anomalies to derive a regional mean time series and then computed its correlation with that of the 500 hPa height on day +5. The results show that the cyclone is closely related to the temperature anomalies that occurred approximate 5 days previously. A maximum correlation of 0.52 was observed when temperature leads the pressure by 4 to 5 days. We also computed the cross-correlations using the 31 PM_{10} collocate stations. The features are almost identical (Figure 13). Similar to the PM_{10} reduction, the holiday temperatures cool over eastern China. Therefore, the regional temperature over southern China experiences a similar cross-correlation with the following cyclone (figure not shown). It should be pointed out that these time lag correlations should not be explained by the natural 1–2-week processes. Because the natural synoptic system's occurrence and phases are randomly in time. Here we prepared the atmospheric correlation/temperatures time series according to the lunar calendar dates. If there is a natural cyclone around the New Year's Day, its random phase should be offset by other cyclones when put all years together. Anyway, these results demonstrate that a stronger cyclone over the western Pacific on days [+4, +6] is often accompanied by a lower preceding PM_{10} concentration and cooler temperatures over eastern China

that occurred approximate one week previously. In addition, it should be pointed out that the northerly winds is helpful reducing the PM_{10} concentration and cooling the temperature. We think that when considering the simultaneous temperature changes in association with short period aerosol anomaly, the atmospheric feedback would be ignorable. However, during a moderate period (such as >3 days to one week) the atmospheric feedback is likely discernible. In Figure 13, we show the correlation peak of the temperature lagging the PM_{10} by a couple of days, which should be indicative of such a feedback. In a word, although our analysis has not proven the cause and effect connection between the holiday air pollution and cyclone, the diagnostics and correlations likely shed lights on their links.

5 Summary

Briefly, the major findings of our analysis are summarized as follows:

The long-term station precipitation data from 1979 to 2012 ~~are~~ analyzed with a focus on the possible changes during the CSF holidays. We found that the precipitation frequency over southern China experiences a significant holiday reduction. The largest reduction occurs on days [+4, +6], with a mean anomaly of -7.4% and a minimum anomaly of -9.4% on day +5. At the same time, the daily precipitation amounts from day +2 to day +5 show continuous negative anomalies, and their mean is -0.62 mm d^{-1} . The Monte-Carlo test implies that the holiday-related frequency and amount anomalies are significantly different from random occurrences. The spatial distributions of the mean frequency anomalies for days [+4, +6] clearly show that a clear reduction appears across the whole southern China. Among the 155 stations, negative anomalies were observed at 148 stations.

The holiday precipitation anomalies are strongly linked to the relative humidity and cloud cover. The station ΔRH shows an evident decrease from day +2 to +7, and the lowest anomalies appear on days [+4, +6], with a mean of -3.9%. When all precipitation days are excluded, the ΔRH shows similar decreases, where the lowest values also occur on days [+4, +6], with anomalies of -0.97%, -0.69% and -0.38%, respectively. The ΔRH vertical profile demonstrates a the significantly drying under approximately 800 hPa. The ERA-Interim reanalysis data revealed that the negative anomaly at 850 hPa is -3.9% and that at 1000 hPa is -4.2%. The mean ΔRH of the layers between 850 hPa and 1000 hPa is -3.9%. The negative anomalies in the lower troposphere are consistent with the significant decreases in the LCC. The daytime station LCC shows an evident reduction during days [+4, +6], with a mean of -6.1%. Meanwhile, the daily ERA-Interim LCC also displays a notable reduction during days [+4, +6]. The corresponding mean is -5.0%, and a minimum of -5.9% appears on day +5. This magnitude as well as its spatial distribution are comparable to that of the station-based daytime LCC anomaly.

The anomalous relative humidity is found to be mainly caused by thea drying of the water vapor in the lower-middle troposphere over southern China during the holidays. Evident negative SH anomalies persist from days -3 to +7 in the station observations. The lowest values appear on approximately days [+4, +5]. The average ΔSH for days [+4, +6] is -0.73 g kg^{-1} . When the precipitation days are excluded, the ΔSH shows a similar, continuous reduction after LNYDNew Year's Day. A minimum of -0.54 g kg^{-1} appears on day +4, while the mean for days [+4, +6] is -0.46 g kg^{-1} . Significant drying-of-the-water

vapor drying is observed for the whole-entire lower troposphere below 700 hPa. Below 800 hPa, the Δ SH values are all less than below -0.50 g kg^{-1} . The mean value between 800 hPa and 1000 hPa is -0.70 g kg^{-1} . The reduction of the water vapor reduction in the lower-middle troposphere likely plays the dominant role in causing anomalous relative humiditys, low-level clouds and precipitation. This water vapor deficit results from the anomalous meridional horizontal moisture transport.

5 During the holidays, a large-scale cyclonic circulation appears over the western Pacific, which brings anomalous northerly winds to East Asia and leads to the negative water vapor flux in the troposphere.

We calculated the correlations between holiday aerosol-temperature and the anomalous cyclones. As the observation PM_{10} and satellite AOD demonstrated, the aerosol concentrations around the CSF decrease evidently. In eastern China, the largest reduction of PM_{10} occurs on days [-3, -1]. A simultaneous cooling was observed (e.g., Gong et al., 2014). Cross-
10 correlation demonstrates that approximately one week after a lower PM_{10} concentration and a cooler temperature over eastern China, a stronger cyclone is observed over the western Pacific. The cooling is likely to lead to anomalous cold advection, which provides a favourable condition for the midlatitude cyclone system. The cyclone brings anomalous northerly wind to East Asia, reduces the atmospheric SH, and consequently results in less precipitation over southern China. The mechanisms need further clarification by elaborate observation analysis and numerical modeling.

15 **Acknowledgements.** This research was supported by projects NSFC-41621061, and NSFC-40975043. The meteorological data were obtained from the China Meteorological Administration. ERA-Interim reanalysis data used in this study were obtained from ECMWF at <http://www.ecmwf.int/>. The API data were obtained from the Chinese Ministry of Environmental Protection. Satellite data used in this study were obtained from MODIS Web of NASA at <https://modis.gsfc.nasa.gov/>. The MERRA-2 reanalysis dataset is publicly available online through the Goddard Earth Sciences Data and Information Services Center (GES DISC; <http://disc.sci.gsfc.nasa.gov/mdisc/>).

References

- 25 B äumer, D., Rinke, R., and Vogel, B.: Weekly periodicities of Aerosol Optical Thickness over Central Europe - evidence of an anthropogenic direct aerosol effect, *Atmospheric Chemistry and Physics*, 8(1), 83-90, doi:10.5194/acp-8-83-2008, 2008.
- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S. K., Sherwood, S., Stevens, B., and Zhang, X.-Y., Clouds and Aerosols, in:
30 *Climate Change 2013: The Physical Science Basis, Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.

K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, UK and New York, NY, USA, 571–657, 2013.

Carranza, J.E., Fisher, B.T., Yoder, G.D., and Hahn, D.W.: On-line analysis of ambient air aerosols using laser-induced breakdown spectroscopy, *Spectrochimica Acta Part B: Atomic Spectroscopy*, 56(6), 851-864, doi: 10.1016/S0584-8547(01)00183-5, 2001.

Cerveny, R.S. and Balling, R.C.: Weekly cycles of air pollutants, precipitation and tropical cyclones in the coastal NW Atlantic region, *Nature*, 394, 561-563, doi: 10.1038/29043, 1998.

Chen, C., Sun, Y.L., Xu, W.Q., Du, W., Zhou, L.B., Han, T.T., Wang, Q.Q., Fu, P.Q., Wang, Z.F., Gao, Z.Q., and Worsnop, D. R.: Characteristics and sources of submicron aerosols above the urban canopy (260 m) in Beijing, China, during the 2014 APEC summit, *Atmospheric Chemistry and Physics*, 15(22), 12879-12895, doi:10.5194/acp-15-12879-2015, 2015.

Choi, Y. S., Ho, C. H., Kim, B. G., and Hur, S. K.: Long-term variation in midweek/weekend cloudiness difference during summer in Korea, *Atmospheric Environment*, 42(28), 6726-6732, doi: 10.1016/j.atmosenv.2008.05.039, 2008.

Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137(656), 553-597, doi:10.1002/qj.828, 2011.

Ding, A. J., Fu, C.B., Yang, X.Q., Sun, J.N., Petäjä T., Kerminen, V. M., Wang, T., Xie, Y., Herrmann, E., Zheng, L.F., Nie, W., Liu, Q., Wei, X.L., and Kulmala, M.: Intense atmospheric pollution modifies weather: A case of mixed biomass burning with fossil fuel combustion pollution in eastern China, *Atmospheric Chemistry and Physics*, 13, 10545-10554, doi:10.5194/acp-13-10545-2013, 2013.

Drewnick, F., Hings, S.S., Curtius, J., Eerdekens, G., and Williams, J.: Measurement of fine particulate and gas-phase species during the New Year's fireworks 2005 in Mainz, Germany, *Atmospheric Environment*, 40(23), 4316-4327, doi: 10.1016/j.atmosenv.2006.03.040, 2006.

Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M. and Zhao, B.: The modern era retrospective analysis for research and applications, version 2 (MERRA-2), *Journal of Climate*, 14, 5419-5454, doi:10.1175/JCLI-D-16-0758.1, 2017.

Gong, D.-Y., Guo, D., and Ho, C.-H.: Weekend effect in diurnal temperature range in China: Opposite signals between winter and summer, *Journal of Geophysical Research: Atmospheres*, 111, D18113, doi:10.1029/2006JD007068, 2006.

Gong, D.-Y., Ho, C.-H., Chen, D., Qian, Y., Choi, Y.-S., and Kim, J.: Weekly cycle of aerosol - meteorology interaction over China, *Journal of Geophysical Research: Atmospheres*, 112, D22202, doi:10.1029/2007JD008888, 2007.

- Gong, D.-Y., Wang, W., Qian, Y., Bai, W., Guo, Y., and Mao, R.: Observed holiday aerosol reduction and temperature cooling over East Asia, *Journal of Geophysical Research: Atmospheres* 119(11), 6306-6324, doi:10.1002/2014JD021464, 2014.
- Gordon, A.H.: Weekdays warmer than weekends?, *Nature*, 367, 325-326, doi: 10.1038/367325b0, 1994.
- 5 ~~Guo, L., Highwood, E.J., Shaffrey, L.C., and Turner, A.G.: The effect of regional changes in anthropogenic aerosols on rainfall of the East Asian Summer monsoon, *Atmospheric Chemistry Physics*, 13, 1521-1534, doi:10.5194/acp-13-1521-2013, 2013.~~
- He, X., Ding, Y., He, R., He, J., and Li, Q.: Analysis on anomalous precipitation in southern China during winter monsoons, *Journal of Meteorological Research*, 21(4), 385-396, 2007.
- 10 Hakim, G., Joly, A., Ayrault, F., Malardel, and S., Cyclone, in: *Encyclopedia of Atmospheric Sciences*, vol. 2, edited by J. R. Holton, J. A. Curry, and J. A. Pyle, Academic Press, London, pp. 589-615, 2003.
- ~~Jones, C., Mahowald, N., and Luo, C.: Observational evidence of African desert dust intensification of easterly waves, *Geophysical Research Letters*, 31, L17208, doi: 10.1029/2004GL020107, 2004.~~
- 15 ~~Kim, B.-G., Choi, M.-H., and Ho, C.-H.: Weekly periodicities of meteorological variables and their possible association with aerosols in Korea, *Atmospheric Environment*, 43(38): 6058-6065, doi: 10.1016/j.atmosenv.2009.08.023, 2009.~~
- Li, B., Gasser, T., Ciais, P., Piao, S., Tao, S., Balkanski, Y., Hauglustaine, D., Boisier, J.-P., Chen, Z., Huang, M., Li, L.Z., Li, Y., Liu, H., Liu, J., Peng, S., Shen, Z., Sun, Z., Wang, R., Wang, T., Yin, G., Yin, Y., Zeng, H., Zeng, Z., and Zhou, F.: The contribution of China's emissions to global climate forcing, *Nature*, 531, 357-361, doi: 10.1038/nature17165, 2016a.
- 20 Li, H., Lin, Z., and Song, Y.: Analysis of the possible factors that influence March precipitation anomalies over South China, *Chinese Journal of Atmospheric Sciences*, 37(3), 719-730, doi: 10.3878/j.issn.1006-9895.2012, 2013. (in Chinese)
- Li, Z., Lau, W.K.M., Ramanathan, V., Wu, G., Ding, Y., Manoj, M.G., Liu, J., Qian, Y., Li, J., Zhou, T., Fan, J., Rosenfeld, D., Ming, Y., Wang, Y., Huang, J., Wang, B., Xu, X., Lee, S.S., Cribb, M., Zhang, F., Yang, X., Zhao, C., Takemura, T., Wang, K., Xia, X., Yin, Y., Zhang, H., Guo, J., Zhai, P.M., Sugimoto, N., Babu, S.S., and Brasseur, G.P.: Aerosol and monsoon climate interactions over Asia, *Reviews of Geophysics*, 54, 866-929, doi:10.1002/2015RG000500, 2016b.
- 25 Lin, J.T. and McElroy, M.B.: Detection from space of a reduction in anthropogenic emissions of nitrogen oxides during the Chinese economic downturn, *Atmospheric Chemistry and Physics*, 11(15), 8171-8188, doi: 10.5194/acp-11-8171-2011, 2011.
- Murray, F.W.: On the Computation of Saturation Vapor Pressure, *Journal of Applied Meteorology*, 6, 203-204, 1967.
- 30 ~~Platnick, S., et al.: MODIS Atmosphere L3 Daily Product, NASA MODIS Adaptive Processing System, Goddard Space Flight Center, USA: http://dx.doi.org/10.5067/MODIS/MOD08_D3.006, 2015a.~~
- ~~Platnick, S., et al.: MODIS Atmosphere L3 Daily Product, NASA MODIS Adaptive Processing System, Goddard Space Flight Center, USA: http://dx.doi.org/10.5067/MODIS/MYD08_D3.006, 2015b.~~

带格式的: 英语(英国)

带格式的: 缩进: 左侧: 0 厘米, 首行缩进: 0 字符

- Qian, Y., Gong, D.-Y., Fan, J., Leung, L.R., Bennartz, R., Chen, D., and Wang, W.: Heavy pollution suppresses light rain in China: Observations and modeling, *Journal of Geophysical Research: Atmospheres*, 114(D7), doi: 10.1029/2008JD011575, 2009.
- Sanchez-Lorenzo, A., Laux, P., Hendricks Franssen, H.-J., Calbó J., Vogl, S., Georgoulias, A.K., and Quaas, J.: Assessing large-scale weekly cycles in meteorological variables: a review, *Atmospheric Chemistry and Physics*, 12(13), 5755-5771, doi: 10.5194/acp-12-5755-2012, 2012.
- Singh, B.P., Srivastava, A.K., Tiwari, S., Singh, S., Singh, R.K., Bisht, D.S., Lal, D.M., Singh, A.K., Mall, R.K., and Srivastava, M.K.: Radiative Impact of Fireworks at a Tropical Indian Location: A Case Study, *Advances in Meteorology*, Article ID 197072, 8, doi: 10.1155/2014/197072, 2014.
- Stjern, C.W.: Weekly cycles in precipitation and other meteorological variables in a polluted region of Europe, *Atmospheric Chemistry and Physics*, 11(9), 4095-4104, doi: 10.5194/acp-11-4095-2011, 2011.
- Tan, P.-H., Chou, C., Liang, J.-Y., Chou, C. C.K., and Shiu, C.-J.: Air pollution “holiday effect” resulting from the Chinese New Year, *Atmospheric Environment*, 43(13): 2114-2124, doi: 10.1016/j.atmosenv.2009.01.037, 2009.
- Tang, M., Ji, D.-S., Gao, W.K., Yu, Z.-W., Chen, K., and Cao, W.: Characteristics of air quality in Tianjin during the Spring Festival period of 2015, *Atmospheric and Oceanic Science Letters*, 9(1), 15-21, doi: 10.1080/16742834.2015.1131948, 2016.
- Travis, D.J., Carleton, A.M., and Lauritsen, R.G.: Climatology: Contrails reduce daily temperature range, *Nature*, 418, 601, doi: 10.1038/418601a, 2002.
- ~~Wang, T., Wang, H.J., Otterå O.H., Gao, Y.Q., Suo, L.L., Furevik, T., and Yu, L.: Anthropogenic agent implicated as a prime driver of shift in precipitation in eastern China in the late 1970s, *Atmospheric Chemistry and Physics*, 13, 12,433-12,450, doi: 10.5194/acp-13-12433-2013, 2013.~~
- Wang, W., Gong, D., Zhou, Z., and Guo, Y.: Robustness of the aerosol weekly cycle over Southeastern China, *Atmospheric Environment*, 61, 409-418, doi: 10.1016/j.atmosenv.2012.07.029, 2012.
- ~~Wang, Y., Wang, M., Zhang, R., Ghan, S.J., Lin, Y., Hu, J., Pan, B., Levy, M., Jiang, J.H., and Molina, M.J.: Assessing the effects of anthropogenic aerosols on Pacific storm track using a multiscale global climate model, *Proceedings of the National Academy of Sciences of the United States of America*, 111, 19, 6894-6899, doi: 10.1073/pnas.1403364111, 2014.~~
- Wang, Y., Zhuang, G., Xu, C., and An, Z.: The air pollution caused by the burning of fireworks during the lantern festival in Beijing, *Atmospheric Environment*, 41(2), 417-431, doi: 10.1016/j.atmosenv.2006.07.043, 2007.
- Wang, Z., Li, Y., Chen, T., Li, L., Liu, B., Zhang, D., Sun, F., Wei, Q., Jiang, L., and Pan, L.: Changes in atmospheric composition during 2014 APEC conference in Beijing, *Journal of Geophysical Research: Atmospheres*, 120(24), 12695-12707, doi: 10.1002/2015JD023652, 2015.

- Wen, W., Cheng, S., Chen, X., Wang, G., Li, S., Wang, X., and Liu, X.: Impact of emission control on PM_{2.5} and the chemical composition change in Beijing-Tianjin-Hebei during the APEC summit 2014, *Environmental Science and Pollution Research*, 23(5), 4509-4521, doi: 10.1007/s11356-015-5379-5, 2016.
- Williams, A. P., Schwartz, R. E., Iacobellis, S., Seager, R., Cook, B. I., Still, C. J., Husak, G., and Michaelsen, J.:
5 Urbanization causes increased cloud-base height and decreased fog in coastal southern California, *Geophysical Research Letters*, 42, 1527-1536, doi: 10.1002/2015GL063266, 2015.
- Xin, J., Wang, Y., Wang, L., Tang, G., Sun, Y., Pan, Y., and Ji, D.: Reductions of PM_{2.5} in Beijing-Tianjin-Hebei urban agglomerations during the 2008 Olympic Games, *Advances in Atmospheric Sciences*, 29(6), 1330-1342, doi: 10.1007/s00376-012-1227-4, 2012.
- 10 Xu, W., Song, W., Zhang, Y., Liu, X., Zhang, L., Zhao, Y., Liu, D., Tang, A., Yang, D., Wang, D., Wen, Z., Pan, Y., Fowler, D., Collett Jr., J.L., Erisman, J.W., Goulding, K., Li, Y., and Zhang, F.: Air quality improvement in a megacity: implications from 2015 Beijing Parade Blue Pollution control actions, *Atmospheric Chemistry and Physics*, 17(1), 31-46, doi: 10.5194/acp-17-31-2017, 2017.
- Xu, W. Q., Sun, Y. L., Chen, C., Du, W., Han, T. T., Wang, Q. Q., Fu, P. Q., Wang, Z. F., Zhao, X. J., Zhou, L. B., Ji, D. S.,
15 Wang, P. C., and Worsnop, D. R.: Aerosol composition, oxidation properties, and sources in Beijing: results from the 2014 Asia-Pacific Economic Cooperation summit study, *Atmospheric Chemistry and Physics*, 15(23), 13681-13698, doi: 10.5194/acp-15-13681-2015, 2015.
- Yang, X., Li, Z., Liu, L., Zhou, L., Cribb, M., and Zhang, F.: Distinct weekly cycles of thunderstorms and a potential
20 connection with aerosol type in China, *Geophysical Research Letters*, 43(16), 8760-8768, doi:10.1002/2016GL070375, 2016.
- ~~Ye, J., Li, W., Li, L., and Zhang, F.: "North drying and south wetting" summer precipitation trend over China and its potential linkage with aerosol loading, *Atmospheric Research*, 125-126, 12-19, doi:10.1016/j.atmosres.2013.01.007, 2013.~~
- Zhang, M., Wang, X., Chen, J., Cheng, T., Wang, T., Yang, X., Gong, Y., Geng, F., and Chen, C.: Physical characterization
25 of aerosol particles during the Chinese New Year's firework events, *Atmospheric Environment*, 44(39), 5191-5198, doi: 10.1016/j.atmosenv.2010.08.048, 2010.
- ~~Zhang, R., Li, G., Fan, J., Wu, D.L., and Molina, M.J.: Intensification of Pacific storm track linked to Asian pollution, *Proceedings of the National Academy of Sciences*, 104(13), 5295-5299, doi: 10.1073/pnas.0700618104, 2007.~~
- 30 Zhao, C., Tie, X., and Lin, Y.: A possible positive feedback of reduction of precipitation and increase in aerosols over eastern central China, *Geophysical Research Letters*, 33, L11814, doi:10.1029/2006GL025959, 2006.

5

10

15

20

25 **Table 1:** The Lunar New Year's Days from 1979 to 2012

Year	Date	Year	Date
1979	28 January	1996	19 February
1980	16 February	1997	7 February
1981	5 February	1998	28 January
1982	25 January	1999	16 February
1983	13 February	2000	5 February
1984	2 February	2001	24 January
1985	20 February	2002	12 February

1986	9 February	2003	1 February
1987	29 January	2004	22 January
1988	17 February	2005	9 February
1989	6 February	2006	29 January
1990	27 January	2007	18 February
1991	15 February	2008	7 February
1992	4 February	2009	26 January
1993	23 January	2010	14 February
1994	10 February	2011	3 February
1995	31 January	2012	23 January






Figure 1: Spatial distribution of January- February precipitation days (a) and precipitation amount (b) for the period of 1979-2012. -Spatial distribution of the January- February averaged Aerosol Optical Depth (AOD) for the period of 2002-2012 (c) and PM₁₀ concentration for the period of 2001-2012 (d) over eastern Asia. The study area is shown in a black dashed rectangle.



10 **Figure 2:** The anomalies of precipitation frequency (a) and amount (b) computed based on 155 stations. The means are shown as gray bars, and the median; and the range of the 10-90th percentiles are also plotted for comparison.



10 **Figure 3:** Anomalies of the precipitation frequency (a) and amount (b). The significances are estimated using a Monte-Carlo approach; stations with circles and dots denote that all days have values significant at the 0.1 and 0.2 levels, respectively.




Figure 4: Station observational relative humidity anomalies (ΔRH) estimated from all available days (a) and the daytime low cloud cover (LCC) anomalies (b) during the holidays from 1979-2012. ERA-Interim ΔRH at 850 hPa (c) and the daily LCC anomalies (d) during the holidays. Standard error is shown as error bars.

10

15

20





Figure 5: Station anomalies of relative humidity estimated from all available days (a) and daytime low cloud cover (b) during days +4 to +6. The significance is estimated using a t -test; stations with circles and dots denote values are significant at the 0.05 and 0.1 level, respectively. Spatial distribution of ERA-Interim ΔRH at 850 hPa (c) and the daily LCC anomalies (d) during days +4 to +6. Only the significant values (at the 0.1 level) are plotted.

10

15

5



10 **Figure 6:** The vertical profiles of the relative humidity (a), temperature (b) and specific humidity (c) anomalies below 500 hPa. The significance level is estimated using a left-tail *t*-test, and the values significant at the 0.05 level are indicated with red dots.

15

20




Figure 7: Means of station-specific humidity anomalies (ΔSH) during the holidays from 1979-2012 as estimated from all available days (a) and no-rain days (b). The corresponding ERA-Interim ΔSH at 1000 hPa (c) and 850 hPa (d) are also plotted.





Figure 8: Spatial distribution of the observational specific humidity anomalies (Δ SH) during days [+4, +6] as estimated from all available days (a) and no-rain days (b). The spatial distributions of the ERA-Interim ASH at 1000 hPa (c) and 850 hPa (d) during days [+4, +6]. Only the significant values (above the 0.05 level) are plotted.






Figure 9: (a) Temporal anomalies of the total column water vapor (ΔTCWV) during the holidays. (b) Spatial distribution of the ΔTCWV (in color shading with unit of kg m^{-2}) and the 700-1000 hPa mean horizontal wind anomalies (in vectors with unit of m s^{-1}) during days [+4, +6]. Stipples denote significant ΔTCWV (at the 0.05 level). Only the anomalous wind vectors significant at the 0.05 level are plotted. (c) Spatial distribution of the mean anomalies of the total precipitation (ΔTP) (in color shading with unit of mm) and the significant (at the 0.05 level) 850 hPa mean horizontal wind anomalies (in vectors with unit of m s^{-1}) during days [+1, +5].




Figure 10: Frequency distributions for the AOD (a) and PM_{10} concentration (b) during days [-15, -11] (red dashed line) and days [+1, +5] (blue solid line) over southern China. (c) The temporal anomalies of the PM_{10} concentration in eastern and southern China. Only years with more than 50% PM_{10} station data available are employed for anomalous composites.

5




Figure 11: Top-of-atmosphere (TOA) longwave radiation anomalies around holidays from MERRA2 (a) and ERA-Interim (b) under clear sky condition.




Figure 102: The mean horizontal wind anomalies at 1000-500 hPa (in vectors with unit of m s^{-1}) and the 1000-500 hPa thickness anomalies (in color shading) during days [-4, 0]. Only significant (at the 0.1 level) winds and thickness are plotted. The long-term mean horizontal winds between 1000-500 hPa during days [-4, 0] are shown together as the streamlines for comparison. A red dashed rectangle (140 E-145 E and 25 N-35 N) indicates the central location of the anomalous cyclone which is manifest in day+5.




Figure 113: (a) Correlations of the 500 hPa geopotential heights over the anomalous cyclone center (140°E-145°E and 25°N-35°N) on day+5 with the PM_{10} concentration regional mean temperature over eastern/southern China in varying lead/lag days. Temperatures from all available 394 stations and that from 31 PM_{10} collocated stations are plotted together for comparison in varying lead/lag days. PM_{10} for eastern and southern China are computed separately. (b) Same as in (a) but for the regional mean temperature over eastern China. Temperatures from all available 394 stations and that from 31 PM_{10} collocated stations are plotted together for comparison.

带格式的: 题注

