

1 **Response to Comments of Reviewers of Manuscript:**

2

3 **Measurements of spectral irradiance during the solar**
4 **eclipse of 21 August 2017: reassessment of the effect of**
5 **solar limb darkening and of changes in total ozone**

6

7

8 **Table of Contents**

9

10	Response to comments by Editor, on 13 March 2019	2
11	Response to comments by Anonymous Referee #1	11
12	Response to comments by Anonymous Referee #2	29
13	Response to comments by Forrest M. Mims III, posted on 24 November 2018	31
14	Response to 2nd comments by Forrest M. Mims III, posted on 21 December 2018	37
15	References	45
16		
17	A marked-up manuscript version of the manuscript is provided from page 48 onward.	

Response to comments by Editor, on 13 March 2019

We thank the editor for his encouraging comments regarding the new version of our manuscript and are pleased to hear that we have satisfactorily responded to the comments by the referees.

All references to figures and other material (e.g., page and line numbers) provided in the following response to the editor's comments refer to the version of our manuscript that we have uploaded on 12 March 2019.

Comment by Editor

Concerning the statements about the diffuse to direct ratio. Reading the paper it is not clear to me from the measurements if this is increasing or not during the eclipse. Looking at figure 10, what it looks like is that direct to global ratio is slightly decreasing at all wavelengths close to the eclipse time. This means that the diffuse to global ratios has to slightly increase. Is the above statement more or less correct ?

Authors' Response

We presume that the question is with respect to Fig. 11, not Fig. 10, of the latest version of the manuscript. Yes, Figure 11 (and in particular Panel b of that figure) indicates that the direct-to-global ratio is decreasing during the eclipse period. However, the minimum ratios are not observed at the time near totality but about 30 minutes before totality (see also P18, L1). Since $\text{global irradiance} = \text{direct irradiance} + \text{diffuse irradiance}$, this implies that the diffuse-to-direct and diffuse-to-global ratios increase during the eclipse period. As described in the manuscript (P18, L6-10 and P24, 8-11), the available evidence suggests that the decrease in the direct-to-global ratio and the concomitant increase in the diffuse-to-direct ratio are caused by aerosols, not by eclipse effects. However, we acknowledge (P24, L10-11) that "aerosol effects are difficult to decouple from processes initiated by the occlusion of the Sun or artifacts of the algorithm used to calculate the direct irradiance from shadowband data" and our assessment is therefore subject to uncertainties. On the other hand, according to 3D radiative transfer (3DRT) calculations by Emde and Mayer (2007), eclipse effects that affect direct and diffuse radiation differently are smaller than 1 % for times 10 minutes away from totality and smaller than 4 % for times 105 seconds away from totality (i.e., the shortest time to totality available from our shadowband measurements) (P24, L13-15). These calculations support our assessment that changes in the direct-to-global ratio were caused by aerosols.

1 In Sect. 8.2. (P24, L17-22), we stated that theoretical calculations [using 3D radiative
2 transfer models]

3 “disagree with the results by Zerefos et al. (2000; 2001), who suggest that the erythema
4 (sunburning) diffuse irradiance was declining at a slower rate than the erythema direct
5 irradiance during the solar eclipse observed in Thessaloniki, Greece, on 11 August 1999.
6 The largest difference was observed at the time of the eclipse maximum when 88% of the
7 solar disk was obscured and the diffuse irradiance was reduced 30 % less than the direct
8 irradiance. The available evidence from our observations and model calculations suggests
9 that the large change of the diffuse-to-direct ratio reported by Zerefos et al. (2000; 2001)
10 may have been a measurement artifact.”

11
12 Considering that “our observations” refer to the direct-to-global ratio while the results by Zerefos
13 et al. (2000) refer to the diffuse-to-direct ratio, we performed additional calculations so that our
14 results can be directly compared with those by Zerefos et al. (2000). Specifically, we calculated
15 the diffuse-to-direct ratio at 315 nm (the wavelength most representative of the erythema range)
16 from our measurements and compared this ratio with the corresponding ratio calculated with our
17 radiative transfer model. Model results were corrected for the bias between measurement and
18 model in the same way as the direct-to-global ratios shown in Fig. 11 of the manuscript.

19
20 Panel a of Fig. E1 below compares the measured and modeled diffuse-to-direct ratio while Panel
21 b shows the ratio of measurement and model, analogously to Fig. 11. Panel a can be directly
22 compared with Fig. 3a of Zerefos et al. (2000), which is also reproduced on the next page. The
23 following can be learned from the comparison of our results and those by Zerefos et al. (2000):

- 24 ■ While the measured diffuse-to-direct ratio increases during both eclipses, the increase in
25 our data is more broad and peaks about 30 minutes before the time of totality. In contrast,
26 data shown by Zerefos et al. (2000) are more “pointy” and peak at the time of the eclipse
27 maximum, which is 11:05 according to Zerefos et al. (2000).
- 28 ■ According to our data, the maximum diffuse-to-direct ratio is 31 %, while the same value
29 estimated from Fig. 3a of Zerefos et al. (2000) is 50 %.




Figure E1: Panel (a): ratio of diffuse-to-direct spectral irradiance. Measurements are shown as line. The bias-corrected model is indicated with a broken line. Panel (b): Ratio of the measured and modeled direct-to-global ratios, corrected for the bias between measurement and model.




Figure 3a of Zerefos et al. (2000), showing the ratio of diffuse-to-direct erythemal solar irradiance for August 11, 1999, measured at Thessaloniki (solid line) and estimated from model calculations (dashed line).

1 **Change to manuscript**

2 Considering that aerosol introduce uncertainty in our assessment of whether or not the eclipse had
3 a significant impact on the direct-to-global and diffuse-to-direct ratios, we weakened the
4 statements where we compare our results with those by Zerefos et al. (2000, 2001).

5
6 In Sect. 8.2., we replaced the paragraph quoted above (P24, L17-22) with:

7 “These theoretical results disagree with the results by Zerefos et al. (2000; 2001),
8 who suggest that the erythema (sunburning) diffuse irradiance was declining at a
9 slower rate than the erythema direct irradiance during a solar eclipse observed in
10 Thessaloniki, Greece, on 11 August 1999. The largest difference was observed at
11 the time of the eclipse maximum when 88 % of the solar disk was obscured and
12 the ratio of diffuse to direct irradiance was increased by approximately 50 %
13 relative to the change expected from calculations with a one-dimensional radiative
14 transfer model. For comparison, the increase in the diffuse-to-direct ratio
15 calculated at 315 nm from our measurements was only 31 % and did not peak at
16 the time of totality. The available evidence from our observations and 3DRT
17 model calculations suggests that measurement artifact may have contributed to the
18 large change of the diffuse-to-direct ratio reported by Zerefos et al. (2000; 2001).”

19
20 **Comment by Editor**

21 There was a lot of arguing against the non use of direct sun measurements and the choice of the
22 305nm and 340nm (well, it was not actually a choice but what was available..). In addition, there
23 is a number of other processes like the channel width (10nm FWHM) procedure to retrieve the
24 1nm spectrum and finally the FOV. FOV of the shadowband is difficult to simulate. And for sure
25 it is larger than an “ideal” FOV of 0.5 degrees or even a 1.5-2.5 degree FOV that most of direct
26 sun pointing AOT measuring instruments use. Correction techniques and calibration principles
27 can improve a lot the AOT results but in the case of the eclipse things are much more
28 complicated, especially at (low signal) lower wavelengths.

29 Could you comment on the above ?

Authors' Response

We agree that it would have been great if our (unfunded) measurement campaign would have also include Sun-pointing instruments with narrow FOV. Such measurements would have afforded to calculate AODs with lower uncertainty and would have also helped to corroborate our global and diffuse shadowband measurements. While we did not have the luxury of deploying additional instrument, we tried our best to keep the uncertainties of our measurements at low as possible, and documented these uncertainties. For example,

- we developed a new method (Section S1 of Supplement) to convert our 10 nm FWHM measurements to 1 nm FWHM measurements and calculated the uncertainty of spectral irradiances calibrated with this method (Section S.1.3 of Supplement),
- used a relatively new data analysis method to minimize uncertainties from the relatively large FOV of the shadowband (see P5, L26 – P6, L9),
- and used an instrument that has low noise levels (and high signal-to-noise ratios), which allowed measurements with sufficient precision at all times, with the exception of measurements of the two 305 nm channels during totality (Sect. 7.5 and 8.4).

All these elements have been discussed in the manuscript and we believe that no changes to the manuscript are necessary to further emphasize these aspects. Furthermore, we do not hide that there are limitations to our measurements and suggests (P26, L10-13) that remaining questions could be settled “by performing LD-corrected measurements of TOC with different instrument types during one of the upcoming solar eclipses.”

Change to manuscript

None.

Comment by Editor

Finally, the ozone issue is a big puzzle for me. Kazadzis, Kazantzidis, 2 papers of Zerefos and Groebner studies all pointed an artificial drop of total column ozone. On the contrary you find an increase. the limb darkening corrections and explanations of all the previous papers can be improved as you say. But the fact remains that a number (about 10) of instruments at different locations (Brewer and NILUs) performing automatic measurements are providing the same artificial decline of ozone.

Would it be possible to explain why this is not the case here ?

Is it the wavelength choice ?

Authors' Response

First, we like to point out that the LD effect increases with decreasing wavelength below the “Balmer break” at about 364.6 nm. Hence, if only a small sliver of the Sun is visible, the extraterrestrial spectrum is more reduced at shorter wavelengths than at longer wavelengths. This is evident both from Fig. 7 of our paper and from Fig. 6 of Neckel (2005). If no correction for LD is applied, measurements at a short wavelength are therefore more suppressed during an eclipse than during the no-eclipse case, which should result in *larger* TOCs retrievals. So our results are consistent with theory. If some instruments report *smaller* TOCs during an eclipse, this would either mean that TOCs have indeed decreased during an eclipse (which is very unlikely, see Sect. 8.3) or measurements are affected by a systematic error (either in the raw data or in the data processing routine), which would be larger than the LD effect and goes in the opposite direction.

Let us discuss results from previous eclipses:

▪ Bojkov (1968)

As mentioned in the manuscript (P25, L3-5), the LD correction applied to Dobson spectrophotometer data from the solar eclipse of 20 May 1966 discussed of Bojkov (1968) reduces the TOC by up to 6 %, in good agreement with our calculations. Fig. 3 of the paper by Bojkov (1968) is reproduced below.




Fig. 3. Changes in total ozone calculated by C_z wavelength measurement (-----), and the same after dark-limb correction (x x x).

1 It shows that the TOC (derived from the Dobson C wavelength pair; 311.2 nm / 332.3
2 nm) was increased close to the eclipse maximum (as in our observations) and that the LD
3 correction reduces the effect. Unfortunately, the paper is very short and there are no
4 useful details on the LD correction. Still, this is a valuable example from Dobson
5 measurements that are by and large consistent with our data.

6
7 ■ **Gröbner et al., 2017**

8 Gröbner et al. (2017) retrieved TOC from Brewer spectrophotometers using a weighted
9 mean of the direct solar irradiance measurements at 310.1 nm, 313.5 nm, 316.8 nm, and
10 320.0 nm. They observed a systematic decrease of up to 15 DU during a partial eclipse of
11 20 March 2015. According to Gröbner et al. (2017), this decrease can be explained to a
12 large extent by LD. Their results “demonstrate that the largest contribution to the
13 observed apparent total column ozone changes of the Brewer spectrophotometer are due
14 to the CLV [aka LD] variations and the relative spectral changes between the specific
15 wavelengths used by the Brewer spectrophotometer.”

16
17 In contrast to our measurements, which are based on pairs of wavelengths (e.g., 305 nm /
18 340 nm), Brewer retrievals depend on four wavelengths (or sometimes five wavelengths
19 as described by Kazadzis et al. (2007)) and the calculation of a weighted mean. This
20 method could conceivably result in the spurious decrease of 15 DU reported by Gröbner
21 et al. (2017). Since we have not operated Brewers and do not know the details of the
22 algorithm, it is beyond the scope of our paper to calculate LD corrections for Brewer
23 TOC retrievals.

24
25 ■ **Kazadzis et al. (2007), Zerefos et al. (2000; 2001)**

26 Kazadzis et al. (2007) and Zerefos et al. (2000; 2001) attribute this apparent reduction in
27 TOC to contamination of the Brewer’s direct measurements by diffuse radiation in the
28 instrument’s field of view (P24, L25-28). Since we are not experts in Brewer
29 measurements we cannot determine whether this assertion is correct. However, we
30 suspect that the combination of different wavelengths, as described by Gröbner et al.
31 (2017), may have been the dominant factor in causing the apparent reduction in TOC. It
32 would be interesting to test whether this apparent reduction would turn into a increase if
33 the Brewer measurements reported by Kazadzis et al. (2007) were reprocessed using a
34 pair of wavelength (as done by us and Bojkov (1968)) instead of using the standard

Brewer algorithm. Again, such a undertaking is beyond the scope of our paper.

▪ **Kazantzidis et al (2007)**

Kazantzidis et al (2007) report on TOC measurements performed during the solar eclipse of 29 March 2006 using NILU-UV filter radiometers, which measure global irradiance. TOCs retrieved by Kazantzidis et al. (2007) using the 305 nm / 320 nm wavelength pair increase as the visible fraction of the sun decreases from ~60% to ~20% up to about 5 DU (Fig. 6 of Kazantzidis et al. (2007)). These increases are by and large consistent with our results. However, as the visible fractions of the sun decreases further, the derived TOCs decrease. The decline is small (<5 DU) for two sites, but quite large (~45 DU) at Kastelorizo. Kazantzidis et al. (2007) attribute this decrease to an artifact in their irradiance measurements and cite Zerefos et al. (2001) and Kazadzis et al. (2007). Considering that both authors use Brewer direct measurements, we don't think that these publications are applicable to NILU-UV measurements, so the drop in TOC (although very small for 2 out of 3 sites) is indeed a mystery.

In summary, while we cannot explain in detail the patterns of TOC measurements performed by other groups in the past, we conclude that the increase in TOC seen in our measurements is consistent with theory. Furthermore, the results by Gröbner et al. (2017) suggest that the decrease in Brewer TOC measurements during an eclipse could partially be caused by the fact that ozone is calculated from four wavelengths rather than two wavelengths as in the case of our results. For example, by combining two wavelength pairs, the effect of aerosols is reduced in standard Dobson retrievals using the AD “double-pair” (Basher, 1982, Bernhard et al., 2004). However, in the case of an eclipse, this method could conceivably lead to incorrect results because the wavelength dependence from solar LD is incorrectly attributed to aerosols. At this point, this explanation is just an untested hypothesis, but the fact that the LD effect calculated by Gröbner et al. (2017) has a similar pattern than the TOC variations measured by two Brewer spectrophotometers during the solar eclipse of 20 March 2015 suggests that this hypothesis warrants further examination. However, this is beyond the scope of this paper.

Change to manuscript

We do not know with confidence why most studies in the past report artificial drops in TOC during solar eclipses and do not want to go too far into speculation. However, the hypothesis is intriguing that ozone retrievals with four wavelengths instead of two could potentially lead to

1 systematic errors because LD and aerosol effects become mixed. We therefore added the
2 following sentence to Sect. 8.3:

3 “We speculate that the observed drop in Brewer TOC measurements during an eclipse is
4 partly caused by the fact that Brewer data are typically calculated from four instead of
5 two wavelengths, as it is the case for our results. Using more than two wavelengths has
6 the advantage that aerosol effects can be reduced (Basher, 1982). However, this method
7 could potentially lead to errors if wavelength-dependent changes in irradiance
8 caused by the LD effect are incorrectly attributed to aerosols.”

Response to comments by Anonymous Referee #1

We thank the referee for his or her comments, which we have addressed as follows:

Comment by Referee

Paragraph 7.4: I am referring firstly to this paragraph because the paper is becoming quite confusing at this point. If the authors conclude that the LD parameterization by Waldmeier is too simple, particularly in UV range, why this is being discussed previously in the estimation of total O₃? It is recommended to move this paragraph to 7.1 and then discuss the results based on those findings on LD parameterizations. In this concept, paragraph 7.3 should follow the new 7.1. In figure 10, the effect of the eclipse is masked by the dominant effect of changing solar zenith angle. The authors are asked to subtract the solar zenith angle effect by using the AOD and O₃ measurements just at the end (or start) of the eclipse (as inputs in model calculations) and, then, discuss the effect of the eclipse and the changes on aerosols and ozone (these paragraph should follow now).

Authors' Response

We have changed the sequence of Sect. 7 as suggest by the referee. The sequence is now:

- 7.1 Effect of solar limb darkening
- 7.2 Direct-to-global ratio
- 7.3 Aerosol optical depth
- 7.4 Total ozone column
- 7.5 Measurements near and during totality

We have also changed the sequence of Sect. 8 (Discussion) to be consistent with Sect. 7. The new sequence of Sect. 8 is now:

- 8.1 Magnitude of solar limb darkening
- 8.2 Variations of direct-to-global ratio
- 8.3 Total ozone column variations during an eclipse
- 8.4 Validation of GUVIS-3511 measurements

Following the suggestion by the referee, we have compared the measured direct-to-global ratio with the corresponding modeled ratio and have added a new panel to Figure 10 (now Figure 11), which shows the ratio of measurement and model. The text was modified to describe the new panel, see below.

Change to manuscript

The following was added to Sect. “Direct-to-global ratio” (now Section 7.2)

“The measured direct-to-global ratio was compared with the modeled direct-to-global ratio using the same method as that employed in Sect. 7.1. In brief, the measured direct-to-global ratio was divided by the modeled direct-to-global ratio. To correct for the difference between measurement and model, linear functions were again constructed to match the ratios of measurement and model at the times of the 1st and 4th contact. Finally, the ratios were normalized by dividing with these linear functions. The resulting bias-corrected ratios of the measured and modeled direct-to-global ratio are shown in Fig. 11b and are denoted $R_{DG}(\lambda_i)$.

Before 15:00, the measured direct-to-global ratio is smaller than 0.03 at wavelengths in the UV-B and values of $R_{DG}(\lambda_i)$ are subject to large uncertainties at these wavelengths. Before 15:20, $R_{DG}(\lambda_i)$ varies between 0.55 and 0.87 for wavelengths in the visible range, indicating that the measured direct-to-global ratio is 13 to 45 % below the modeled one. These low values can be attributed to increased aerosol loading prior to the start of the eclipse. After the start of the eclipse (1st contact), values of $R_{DG}(\lambda_i)$ start to drop to reach a local minimum at 16:52, approximately 30 minutes before the eclipse maximum. The decline has a clear wavelength dependence with wavelengths in the UV-B decreasing the most (up to 20 %), followed by wavelengths in the visible (up to 15%) and IR (up to 3.5%). After this local minimum, $R_{DG}(\lambda_i)$ slowly increases during the remainder of the eclipse.

The change of $R_{DG}(\lambda_i)$ over the period of the eclipse could either be caused by processes initiated by the occlusion of the Sun or by variability from aerosols. Aerosols effects are more likely because any eclipse effects should conspicuously peak at the time of totality, not 30 minutes earlier. The wavelength dependence of $R_{DG}(\lambda_i)$ is also characteristic for aerosol effects (Sect. 7.3). In addition, the minimum values of $R_{DG}(\lambda_i)$ during the eclipse are well within the

range of values observed prior to the eclipse, which could be unambiguously attributed to aerosols.”




Fig. 1. Panel (a): ratio of direct-to-global spectral irradiance. Measurements are shown as heavy lines. Thin lines connect measurements at the start of end of the eclipse and are drawn to guide the eye. Panel (b): Ratio of the measured and modeled direct-to-global ratios, corrected for the bias between measurement and model. Numbers “(1)” and “(2)” in the legend indicate the channel number for channels equipped with identical wavelengths.

In Section 8.1., the following paragraph:

“The direct-to-global ratios shown in Fig. 10 increase with time as expected from the decrease in SZA. Deviations from a straight line between the 1st and 4th contact are less than 0.04. It is difficult to determine whether this deviation is caused by processes initiated by the occlusion of the Sun, by variability from aerosols, or artifacts of the

algorithm to calculate the direct irradiance from shadowband data. We therefore consider the deviation of 0.04 as an upper limit for the eclipse effect.”

was removed and replaced with:

“Relative to model calculations for the unoccluded Sun, direct-to-global ratios during the eclipse decreased by up to 20 % in the UV-B, 15 % in the visible and 3.5 % in the IR (Fig. 1b). The largest decrease was observed about 30 minutes before totality. The timing of this decrease plus its spectral dependence suggest that changes in aerosol amounts are the main driver for the observed drop in the ratio. However, aerosol effects are difficult to decouple from processes initiated by the occlusion of the Sun or artifacts of the algorithm used to calculate the direct irradiance from shadowband data.”

Note that this paragraph is now part of Section 8.2.

Comment by Referee

Page 2, lines 15-20: Kazantzidis et al (2007) were measuring with NILU-UVs and calculated total ozone, too. They reported a slight increase when the visible part of the sun was more than 20% and decreased significantly as the eclipse progressed.

Authors' Response and change to manuscript

The reference to Kazantzidis et al. (2007) has been added to the introduction in support of the following statements:

“study wavelength-dependent changes in spectral irradiance”

“short-term and longer-lasting fluctuations in the total ozone column (TOC)”

Furthermore, we have added the following to Sect. 8.3. of the Discussion:

“Finally, Kazantzidis et al (2007) discuss TOC measurements performed with NILU-UV filter radiometers at several locations in Greece during the total solar eclipse of 29 March 2006. They did not observe any periodic fluctuations in TOC and only report a small increase in TOC of about 5 DU as the visible fraction of the Sun decreases from ~60% to ~20%. This small change could be caused by incomplete correction of the LD effect.”

Comment by Referee

Figure 3 and relevant text: please provide a figure only with the UV wavelengths, no logarithmic axes. From the literature it seems that the bandwidth of 305nm channel is quite wide: it is more

- 1 than 10nm even at full width at half maximum. In this case, please discuss the possible
- 2 implications when measuring with this high bandwidth.

Authors' Response and change to manuscript

Figure 3 was plotted separately for the UV, visible and IR wavelength range on a non-logarithmic x-axes. The new figure plus raw data in text format will be made available as Supplements.

The following was added to Section 3:

“The spectral bandwidth of all channels is approximately 10 nm full width at half maximum (FWHM) with the exception of the two channels at 305 nm, which have a bandwidth of 18.5 nm.”

The following was added to the Caption of Fig. 3:

“A version of the figure, plotted separately for UV, visible, and IR wavelengths, is part of the Supplement.”

The following was added to Sect. 5.1:

“Aerosol optical depth was not calculated for the two 305 nm channels because of the large bandwidth of these channels and the strong interference with ozone absorption at this wavelength. Both factors lead to large uncertainties.”

As described in Section 5.2., TOCs are calculated using look-up tables that are based on response-weighted global irradiance, i.e., the spectral irradiance weighted with the spectral response functions shown in Fig. 3. TOC calculations therefore take the large bandwidth of the 305 nm channels into account.

The impact of the relatively large bandwidth of the 305 nm channels on spectral irradiance calibrations is implicitly addressed by the vicarious calibration method described in Sect. S1.2 of the Supplement. Uncertainties to derive spectral irradiance at 1 nm resolution from response-weighted irradiance are discussed in Section S1.3 of the Supplement.

Comment by Referee

Page 5, lines 15-24: please provide/add some sentences about the performance of this method on estimating the direct and the shadowband corrected spectral irradiances.

Authors' response and change to manuscript

The following was added to the paragraph in question:

“The uncertainty of our method was estimated by Witthuhn et al. (2017). AOD can be retrieved with an uncertainty of 0.02 for all channels within a 95 % confidence interval.”

Comment by Referee

Figure 4 and relevant text: What is the expected cosine error in UV wavelengths? Are there any measurements? Are the measurements in UV channels (used for this study) cosine corrected? How significant this effect is expected to be during the solar eclipse (when the direct component of solar irradiance is minimized)?

Authors' Response

The cosine error in the UV has also been measured but results were omitted from Fig. 4 because these measurements are very noisy due to the low output of the FEL used for the characterization in the UV. In general, the cosine error of wavelengths between 305 and 380 nm is similar to that at 395 nm because scattering properties of PTFE (Teflon) deteriorate only towards longer wavelengths but not towards shorter wavelengths. We have now generated a composite figure showing cosine errors in the UV in the upper panel and those for the visible and IR ranges in the lower panel:




Fig. 2: Cosine error of the GUVIS-3511 radiometer in one azimuthal plane at wavelengths between 305 and 395 nm (Panel a) and 395 and 1020 nm (Panel b). Measurements performed at the orthogonal plane are similar. Measurements at wavelengths below 395 nm are affected by noise due the low UV output of the incandescent lamp used for the characterization.

The new figure will be included in the Supplement. Measurements of UV channels used in our study are cosine error corrected. Corrections of channels between 305 and 380 nm are based on the cosine error of the 395 nm channel to avoid spurious corrections caused by the noisy cosine response data. The cosine error correction takes the measured direct-to-global ratio into account. The accuracy of the cosine error correction method is therefore not affected by the solar eclipse.

Change to manuscript

The following was added to Sect. 3:

“Measurements of the cosine error at wavelengths below 395 nm are affected by noise due the low UV output of the incandescent lamp used for the characterization.

1 Corrections for the cosine error of UV channels described further below are based on the
2 measured cosine error at 395 nm.”

3

4 The following was added to the caption of Fig. 4:

5 “(A figure with cosine errors in the UV is provided in the Supplement.)”

6

7 The following was added to Sect. 4:

8 “Both direct and global irradiance measurements were corrected for the cosine error of
9 the instrument’s collector as described by Morrow et al. (2010), based on the measured
10 cosine error (Fig. 4) and the ratio of direct and global irradiance extracted from the
11 shadowband measurements.”

12

13 **Comment by Referee**

14 Figure 8: Aerosol Optical Depth (AOD) around 305nm is not depicted in figures 8a and 8b. Why?
15 Do you think that the LD correction (8b) by Pierce is valid also for the lower UV wavelengths? If
16 yes, how this decrease of AOD in these wavelengths can be explained? There is an assumption
17 that this is not a measurement artifact but an unknown absorber. Which type of absorber could
18 change the expected AOD values only at 314.2, 319.4nm (and 442.4nm!)? Moreover, what would
19 be the AOD estimations if you apply the Neckel LD parameterization?

20

21 **Authors’ Response and change to manuscript:**

22 The uncertainty in calculating AOD at 305 nm is far too high to be useful. As already mentioned
23 earlier, we have added the following to Sect. 5.1 to make this clear:

24 “Aerosol optical depth was not calculated for the two 305 nm channels because of the
25 large bandwidth of these channels and the strong interference with ozone absorption at
26 this wavelength. Both factors lead to large uncertainties.”

27

28 Yes, we believe that the LD correction by Pierce is valid also for the lower UV wavelengths
29 because the difference in global irradiance between measurement and theory is also reasonable at

305 nm (Fig. 11d of original manuscript). Furthermore, the LD corrections by Pierce and Neckel (which were derived from independent datasets) agree to within 1.8% at 305 nm.

We do not know with certainty why the AODs at 314.3, 319.4, and 442.4 nm are below the AODs expected from the Ångström parameterization, which was derived using channels between 340 and 1020 nm, excluding the channel at 940 nm, as stated in the manuscript. These low values could be caused by absorbing aerosols released by the near-by fires or measurement errors, but this is speculation. If the low values were due to measurement errors, one would expect that the discrepancies for the three wavelengths observed at 16:02:58 and 19:00:30 are of similar magnitude because measurements at both times were processed in the same way. The fact that these discrepancies are larger at 19:00:30 suggests that they are caused by real changes in aerosol properties. While these drops could be caused by an unknown absorber, we removed this suggestion from the text because we don't have evidence that such an absorbing aerosol or gas was indeed present.

Considering the uncertainty with respect to the cause of the low AODs at the three wavelengths, we changed the last sentence of the paragraph to:

“As measurements at the two time were processed in the same way it seems unlikely that the discrepancies are caused by a measurement artifact.”

With respect to the question “Moreover, what would be the AOD estimations if you apply the Neckel LD parameterization?” we note that AODs shown in the figure were calculated for times before the start (16:02:58) and after the end (19:00:30) of the eclipse. Hence, no LD parameterization was applied or is necessary.

Comment by Referee

Paragraph 7.2: It is quite surprising that although AOD can be estimated by direct GUV measurements, this is not happening for the total O₃ amount as well, despite that direct measurements (divided by global ones) are presented in figure 10. Moreover, the 305/340 wavelength ratio methodology to derive total O₃ is based on model calculations as a function of O₃ and solar zenith angles but under cloud-free skies e.g. specifically defined direct and diffuse components of solar irradiance. This is not the solar eclipse case. And this is accounted by the authors. However, how valid are the LD parameterizations on CHANNEL irradiances when it is known that the direct/diffuse ratio has significant spectral sensitivity? And how repeatable will be


1 their results if the they use: a) direct irradiances, b) different channel ratios, e.g. 305/320,
2 312/340? The authors here should acknowledge a couple of very crucial facts: 1) the 305/340
3 wavelength ratio of channel irradiances is a well-known method that can be used for total O3
4 estimations but it is accompanied by significant uncertainties (aerosol optical depth and scattering
5 properties, cloudiness, ozone profile, direct/diffuse irradiance etc), 2) the comparison with ozone
6 values or previous studies derived from instruments measuring the direct irradiance should be
7 done under the acknowledgement that these measurements are correct not only because they are
8 the standard ones but also because they are correct in terms of physics and the best in terms of
9 overall uncertainty.

11 **Authors' Response**

12 First, we like to emphasize that this paper is not about the most accurate or appropriate method to
13 calculate TOC. TOC measurements were mainly included in the manuscript (i) to illustrate that
14 TOC measurements during an eclipse are affected by the solar limb darkening (LD) effect and
15 that errors arising from this effect can be greatly reduced with appropriate LD corrections, and (ii)
16 to determine whether gravity waves generated by the Moon's shadow can cause oscillations in
17 TOC as suggest by several authors in the past. In our opinion, both questions were sufficiently
18 addressed with the results presented in the original manuscript. Please see also our response to the
19 comments by Forrest M. Mims.

21 Motivated by the referee's comments, we have calculated total ozone now also from global
22 irradiance of the wavelength pairs 305 / 313 and 313 / 340. Results are presented in Fig. 3. Fig. 3a
23 shows results without LD correction. TOCs for the 305 / 340 pair (blue) are identical with the
24 "uncorrected" data shown in Fig. 9 of the original manuscript. Ozone values for the 313 / 340 are
25 *lower* than those of the 305 / 340 pair and exhibit a *negative* slope as a function of time. In
26 contrast, ozone values for the 305 / 313 pair are *larger* than those of the 305 / 340 pair and exhibit
27 a *positive* slope. These biases are likely caused by a small systematic error in measurements of
28 the 313 nm channel. When measurements of this channels are scaled with a factor of 0.95 and
29 ozone retrievals repeated, TOCs derived from the three wavelengths pairs are consistent to within
30 ± 2.5 DU outside the period of the eclipse (Fig. 3b). A systematic error of 5% is well within the
31 expanded uncertainty of 7.3% of the 313 nm channel (see table S1 of Supplement.). Using this
32 adjustment, measurements of all pairs agree with OMI data to within 5 DU (1.7%) (red line in
33 Fig. 3).

1



2

3 Fig. 3. Total ozone column derived from global irradiance measurements of the GUVIS-3511 radiometer
 4 using different wavelength pairs. Panel (a): comparison of ozone retrievals with the 305 / 340 pair (blue),
 5 the 313 / 340 pair (orange), and the 305 / 313 pair (green). Panel (b): same as Panel (a) but measurements
 6 of the 313 nm channel were scaled with a factor of 0.95 before calculating ozone. Panel (c): TOC retrievals
 7 for the 313 / 340 pair using either no correction (blue), or the LD-corrections by Waldmeier (grey), Pierce
 8 (cyan), and Neckel (green). Panel (d): same as Panel (c) but for the 305 / 313 pair. Measurements of the
 9 313 nm channel were scaled by 0.95 for all retrievals shown in Panels (c) and (d). Long-dashed lines
 10 indicate the start and end times of the eclipse and the short-dashed line indicates the time at totality. Note
 11 that y-scales of the various panels differ.

12

13 TOC data shown in Fig. 3c and 3d were corrected for the LD-effect similar to the TOC data
 14 derived from the 305 / 340 pair shown in Fig. 9 of the original manuscript. As in that figure,
 15 TOCs calculated for the 313 / 340 and 305 / 313 pairs (Fig. 1c and 1d, respectively) spike at the
 16 time of totality when no LD correction is applied. The LD correction by Waldmeier has only a
 17 marginal effect on this result. In contrast, LD corrections using the parameterizations by Pierce
 18 and Neckel greatly decrease this spurious spike.

19

20 Calculations of TOCs from global irradiance using the pairs 305 / 313 and 313 / 340 therefore
 21 confirm the conclusions reached from results of the 305 / 340 nm pair that were presented in the
 22 original manuscript:

- 1 ▪ There are no periodic oscillations in the TOC, either before or after totality, that could be
- 2 attributed to the effects of bow waves from the Moon's shadow.
- 3 ▪ The LD corrections by Pierce and Neckel greatly reduce the spurious spike in TOCs
- 4 during totality resulting from the LD effect. In contrast, the correction that is based on the
- 5 parameterization by Waldmeier is too small.
- 6 ▪ The spurious TOC spike is larger for the 313 / 340 pair than for the 305 / 340 pair even
- 7 though the relative difference in the LD effect is smaller for the two wavelengths used in
- 8 the former pair. This observation is due to the fact that small errors in measurements at
- 9 313 nm have a larger effect on TOC than at 305 nm because the absorption coefficient of
- 10 ozone is much smaller at 313 than 305 nm. The effect of ozone absorption on the 305 /
- 11 340 ratio therefore outweighs the LD effect.
- 12 ▪ There is no evidence that aerosol effects have an important effect on our TOC
- 13 measurements. While we agree with the referee's assessment that the method of
- 14 calculating TOCs from ratios of global irradiance at 305 and 340 nm "is accompanied by
- 15 significant uncertainties (aerosol optical depth and scattering properties, cloudiness,
- 16 ozone profile, direct/diffuse irradiance etc)," we believe that these uncertainties are of
- 17 little relevance for the conclusions reached in our paper. Specifically:
 - 18 ○ AODs were characterized and used for the lookup tables for ozone retrievals. The
 - 19 good consistency of TOC retrievals for different wavelength pairs (after adjusting
 - 20 for the bias of measurements at 313 nm) confirms that aerosol effects were
 - 21 addressed adequately.
 - 22 ○ There were no clouds during the period of the measurements that could have
 - 23 skewed our TOC retrievals.
 - 24 ○ The ozone profile (through the Umkehr effect) becomes only important for SZAs
 - 25 > 75° (i.e., outside the range of SZAs occurring during the eclipse).
 - 26 ○ Changes in the direct-to-global irradiance agree to within 2.5% for the
 - 27 wavelength range 305 - 340 nm, except for a short period centered about totality
 - 28 (Fig. 1 above). So our look-up tables, which were calculated for no-eclipse
 - 29 conditions, are also suitable for calculating TOCs during the eclipse.

30

31 We did not calculate TOCs from measurements of direct irradiance because such retrievals have a

32 comparatively large uncertain when direct irradiance in the UV-B (in particular 305 nm) is

33 calculated from shadowband data. In contrast to wavelengths in the UV-A and visible, where

34 direct irradiance under clear skies with moderate aerosol loading makes up a large fraction of the

global irradiance, this is not true for wavelengths in the UV-B, in particular at large SZAs (see Fig. 1a). When using a shadowband, direct irradiance in the UV-B is calculated as the difference of two large numbers (global and diffuse irradiance), and the uncertainty of the result is therefore higher than those of TOCs derived from global irradiance.

Regarding the referee's comment:

“the comparison with ozone values or previous studies derived from instruments measuring the direct irradiance should be done under the acknowledgement that these measurements are correct not only because they are the standard ones but also because they are correct in terms of physics and the best in terms of overall uncertainty.”

While we agree that measuring TOC from direct irradiance is the more straightforward method (as only Beer-Lambert's law is involved), the referee's assertion that “these measurements are correct” is not appropriate. There are many error sources that can affect these measurements such as pointing errors, calibration errors, non-linear sensor response, stray light from diffuse radiation that is detected with the direct beam (a problem affecting Brewer measurements as mentioned by several papers cited in our manuscript), uncertainty of the filter functions and their center wavelengths, etc. Considering that we do not have the raw data that other authors have collected to calculate TOCs during historical ellipses we do not want to speculate on the uncertainty of those measurements and their magnitude relative to our results.

As explained above, adding TOC data derived from different wavelength pairs would not change our conclusions about the effect of the solar eclipse on short-term TOC fluctuations. However, results obtained with the 305 / 313 and 313 / 340 pairs are valuable because they confirm the results from the 305 / 340 pair discussed in the manuscript. In order to keep the length of the manuscript within reasonable limits while also respecting the referee's suggestion, we added these new results to the Supplement with references provided in the manuscript.

Change to manuscript

The following was added to Sect. 5.2:

“The following three wavelength pairs were used: (1) $\lambda_L = 340$ nm, $\lambda_S = 305$ nm; (2) $\lambda_L = 340$ nm, $\lambda_S = 313$ nm; (3) $\lambda_L = 313$ nm, $\lambda_S = 305$ nm.
[...]

1 “While TOCs could also be derived from direct irradiances, these measurements are not
2 discussed here due to the relatively large uncertainty to calculate direct irradiance from
3 shadowband data at wavelengths in the UV-B (280–315 nm), in particular at large
4 SZAs.”

5

6 The following was added to Sect: 7.4:

7 “Results obtained from the 305 / 340 wavelength pair are discussed below. Similar results
8 calculated with the 305 / 313 and 313 / 340 wavelength pairs are presented in Sect. S2 of
9 the Supplement.”

10 [...]

11 Results obtained with the 305 / 313 and 313 / 340 wavelength pairs (Sect. S2 of the
12 Supplement) corroborate these findings.”

13

14 **Change to Supplement:**

15 Calculations of TOC from the 305 / 313 and 313 / 340 wavelength pairs have been added to the
16 Supplement. Please see Section S2 of the new version of the Supplement for details.

17

18 **Comment by Referee**

19 Figure 9 and relevant text: Kazantzidis et al (2007), when using the 305/320 wavelength ratio (in
20 order to reduce the effect of spectral effect of the eclipse on direct and diffuse irradiances)
21 reported very similar results to those derived from yours when using the 305/340 wavelength
22 ratio AND applying the Neckel or Pierce parameterization. Surprisingly, this paper is referenced
23 only for the comparison of measured and model irradiances. However, the results of this paper for
24 ozone, irradiances and irradiance ratios vs eclipse percentage are not referenced, although it is
25 based on results from 8 narrowband multi-channel NILU-UV6 radiometers.

26 **Authors’ response and change to manuscript**

27 As mentioned already earlier, we have now added the following to the manuscript to compare our
28 ozone measurements with those reported by Kazantzidis et al (2007):

29 “Finally, Kazantzidis et al (2007) discuss TOC measurements performed with NILU-UV
30 filter radiometers at several locations in Greece during the total solar eclipse of 29 March
31 2006. They did not observe any periodic fluctuations in TOC and only report a small

1 increase in TOC of about 5 DU as the visible fraction of the Sun decreases from ~60% to
2 ~20%. This small change could be caused by incomplete correction of the LD effect.”
3

4 Furthermore, we have added the following to Sect. 8.2 (now Sect. 8.1) where we compare the
5 magnitude of solar LD used in our paper with that used by other authors:

6 “Kazantzidis et al.(2007) have analyzed ratios of global spectral irradiance (305 nm / 380
7 nm, 312 nm / 380 nm, 340 nm / 380 nm, and PAR / 380 nm) that were measured with
8 NILU-UV filter radiometers at three locations in Greece during the total solar eclipse of
9 29 March 2006. These measured ratios were compared with theoretical predictions based
10 on the algorithm by Koepke et al. (2001) and the LD parameterization by Waldmeier. As
11 the eclipse progressed, the model underestimated the measured spectral effect, capturing
12 only half of the observed change. For example, measured ratios of spectral irradiances at
13 340 and 380 nm were 10 % lower close to totality compared to similar ratios calculated
14 for the 1st and 4th contact. The theoretical calculation only predicted a decrease of 5%.
15 Our results suggests that discrepancies between the measured and modeled ratios
16 reported by Kazantzidis et al. (2007) can partly be attributed to limitations of the LD
17 parameterization by Waldmeier used in their model.”
18

19 **Comment by Referee**

20 Paragraph 8.1: the authors seem to have a point here. In order to better understand the
21 similarities/differences with previous studies, a more detailed information is needed apart from
22 direct/diffuse ratios: the measured wavelengths and the eclipse percentages should be provided.
23 Moreover, the theoretical calculations from Emde and Mayer are quite capable to estimate the
24 global irradiance (when normalized 5 minutes before totality) at 380 nm but maybe irradiance is
25 significantly underestimated at 312 nm (Kazantzidis et al., 2007, figure 7 and relevant text). This
26 affects directly the diffuse component. Of course, also this result is sensitive to factors like
27 surface albedo, ozone profile and the dynamic range of the measuring system. All these factors
28 should be mentioned.

29
30 **Authors' response**

31 We added the following sentence to Section 8.1. (now Sect. 8.2), which now specifies the
32 measured wavelengths (i.e. erythral irradiance), eclipse percentage (88%), location
33 (Thessaloniki), and time (11 August 1999):

1 “However, these results disagree with the results by Zerefos et al. (2000; 2001), who
2 suggest that the erythema (sunburning) diffuse irradiance was declining at a slower rate
3 than the erythema direct irradiance during the solar eclipse observed in Thessaloniki,
4 Greece, on 11 August 1999. The largest difference was observed at the time of the
5 eclipse maximum when 88% of the solar disk was obscured and the diffuse irradiance
6 was reduced 30 % less than the direct irradiance.”

7
8 With regards to factors such as “surface albedo, ozone profile and the dynamic range of the
9 measuring system” that affect irradiance close to totality, we note that we concluded our original
10 manuscript with the sentence:

11 “During totality, the irradiance at the surface will become also more sensitive to the
12 topography (e.g., the mountains surrounding the measurement sites), surface albedo (and
13 its spectral dependence), and the distribution of ozone in the atmosphere (the ozone
14 profile). These aspects will be discussed in a follow-on publication.”

15 The “follow-on publication” mentioned here is close to completion and will be submitted soon.
16 The factors enumerated by the referee will be discussed in detail therein.

17
18 **Comment by Referee**

19 Page 22-23: This comment refers to the ozone issue, described in detail by the authors. From my
20 point of view, some (or much?) of these differences could be attributed on the measuring methods
21 and the selected pairs of wavelengths. As mentioned before, a decrease of (uncorrected for LD
22 effect) ozone retrievals has been reported by Kazantzidis et al (2007) for the same eclipse when
23 using 8 NILU-UVs. As it was stated earlier in this review, the authors could strongly defend their
24 findings if they will come up with the same results when using other wavelength pairs and the
25 direct GUV irradiances. Unfortunately, this measuring campaign is not accompanied by more
26 instruments.

27
28 **Authors’ response**

29 One important “take-home message” from our paper is that accurate LD corrections are necessary
30 for accurate TOC retrievals. Basically all papers that have discussed TOC measurements for
31 previous eclipses use a LD correction that is based on a parameterization that is too simple for
32 wavelengths in the UV. In addition to TOC retrievals using the 305 / 340 wavelength pair, we
33 have now also calculated TOCs from the 305 / 313 and 313 / 340 wavelength pairs (Fig. 3

1 above). These new calculation corroborate our initial findings that the spurious peak in TOC
2 measurements caused by LD effect can be greatly reduced by using either the LD
3 parameterization by Neckel or Pierce. We agree with the referee's assessment that the presence of
4 additional instrument during "our" eclipse could have strengthened our conclusions further. We
5 hope that a multi-instrument campaign can be organized during an upcoming eclipse to get
6 closure on some of the remaining questions raised in our paper.

Response to comments by Anonymous Referee #2

We thank the referee for his or her comments, which we have addressed as follows:

Comment by Referee

A further examination on the effect of the above (mainly the FOV) issues have to be considered when comparing results of this and older papers using different principles of measurements. For example the definition of the diffuse (or direct) irradiance calculated using a shadow (band) that has spectral, solar zenith angle, (and in this case) also sun- dimensions dependent, apparent shadow dimensions compared with the instrument diffuser, could impact the presented results.

Authors' Response

First we like to point out that total ozone column (TOC) was calculated from measurements of global spectral irradiance, not direct solar irradiance as indicated by the referee's introductory statements. The FOV of the shadowband is therefore irrelevant for TOC retrievals. Please see our response to the comments by Forrest M. Mims III regarding TOC calculations.

Shadowband measurements were only used to determined aerosol optical depth and the direct-to-global ratio. The shadowband obstructs a wedge of the sky with a width of 15° when the band is vertical and 13° when it is horizontal. These relative large angles may suggest that the instrument is not able to adequately remove the contribution of circumsolar radiation when calculating the direct irradiance, which would also lead to systematic errors in aerosol optical depth (AOD) retrievals. However, this notion is not correct due to the unique way of shadowband operation and data analysis. Other shadowband radiometer typically use two measurements on either side of the Sun to correct for the signal lost from the portion of the sky that is obscured by the shadowband. For example, side-band measurements of Multifilter Rotating Shadowband Radiometer (MFRSR) are typically performed at 9° from the center of the Sun (Krotkov et al., 2005). Hence, the increase of diffuse radiation towards the Sun is often not corrected adequately. In contrast, direct measurements and AOD retrievals of GUVIS-3511 measurements are based on measurements at high sampling rate (15 Hz) whereby the band moves slowly and continuously over the instrument. The processing algorithm determines the irradiance from direct Sun plus the unshaded portion of the sky by analyzing several seconds of measurements when the shade from the band is close to the diffuser but not in contact with it. The algorithm is described in detail by

Morrow et al., (2010) and Witthuhn et al. (2017). The algorithm compensates for the effect of the relatively large FOV, resulting in AOD data that have similar accuracy than that of traditional shadowband radiometers.

Change to manuscript

The following text will be added to Sect. 3 of the manuscript:

“The band obstructs a wedge of the sky with a width of 15° when the band is vertical and 13° when it is horizontal. [...] The algorithm compensates for the effect of the relatively large width of the shadowband, resulting in AOD data that have similar accuracy than those of traditional shadowband radiometers, which use measurements on either side of the Sun to correct for the portion of the sky that is obscured by the band. For example, side-band measurements of Multifilter Rotating Shadowband Radiometer (MFRSR) are typically performed at 9° from the center of the Sun (Krotkov et al., 2005) and therefore may not adequately measure circumsolar radiation. Our algorithm alleviates this problem.”

Comment by Referee

What do you mean by: P6, L10 The calibration was further optimized for the conditions (solar zenith angle, TOC, AOD, etc.) at the measurement site.

Authors' Response

As described in the Supplement of the manuscript, the instrument measures “response-weighted irradiance”. The conversion to spectral irradiance requires knowledge of the solar zenith angle, TOC, AOD, etc., and uncertainties in these parameters results in uncertainty in the conversion factor. These issues are discussed at length in the Supplement.

Change to manuscript

The following will be added after the sentence in question:

“Details of this optimization are provided in the Supplement.”

Response to comments by Forrest M. Mims III, posted on 24 November 2018

We thank Mr. Mims for his good comments, which have helped to improve our manuscript.

Comment by reviewer

While the results in this paper are certainly intriguing, there are significant differences between the instrument employed by the authors and the TOPS instrument employed by Mims and Mims. The author's instrument uses filters having a FWHM bandpass of 10 nm, while TOPS has filters with a 5-nm bandpass FWHM. TOPS also measured column ozone at 300nm and 305 nm, which is much more sensitive to ozone variations than the wavelengths used by the authors. TOPS is also a direct sun instrument that can provide measurements in a few seconds, while the author's instrument is a full-sky C1 device with an exceptionally long 2-minute scan time. As we have learned from comparisons with an EPA Brewer placed at our site, the much faster scan time provided by TOPS provides higher resolution results and avoids errors caused by aerosol changes that can occur during minute-duration scans. Moreover, TOPS often detects subtle changes in the ozone column missed by Dobson and Brewer instruments, which both require considerably more time for an ozone measurement. Before our findings are ruled out by this paper, we feel that the authors should point out the very significant instrumental differences, especially the filter wavelengths, the filter bandpasses and the time required per scan. In each of these cases, TOPS offers superior performance when compared with their instrument. Thus, the findings of subtle waves in the ozone layer by TOPS cannot be so quickly discounted by this paper. I close by observing that TOPS uncovered a calibration drift in the Nimbus-7 Total Ozone Mapping Spectrometer (TOMS) (Satellite Monitoring Error, Nature 361, 1993). TOPS evolved into Microtops and then Microtops II. All these instruments provide results in close agreement with Brewers and Dobsons at the Mauna Loa Observatory. Thank you for considering the points made herein. Forrest M. Mims III fmimsiii@yahoo.com Interactive comment on Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2018-1048>, 2018.

Authors' Response

As described in Section 5.2. of the manuscript, the total ozone column (TOC) was derived from measurements of response-weighted global irradiance measured by the GUVis-3511. Specifically, ratios of measurement at 340 and 305 nm were compared with similar ratios in a look-up table that was calculated with a radiative transfer model as a function of SZA and TOC. The look-up table was calculated by taking the response functions of the instrument (Fig. 3 of manuscript) and observing conditions (e.g., aerosol optical depth, (AOD)) into account. The method of calculating TOC from measurements of global irradiance (instead of direct irradiance as it is typically done for Dobson, Brewer, TOPS, and Microtops instruments) was first proposed by Stamnes et al. (1991). We found that the accuracy of TOCs derived from global irradiance is similar to that of data from Dobson instruments or satellite (TOMS, OMI) observations if the look-up table takes local conditions into account (ozone profile, albedo, elevation, etc.) (Bernhard et al., 2005b). For example, this study uncovered systematic errors in Dobson measurements associated with approximations in the standard Dobson retrieval method, which subsequently helped to better understand the limitation of Dobson measurements. We have further validated the method for GUV instruments (Bernhard et al., 2005a). Based on this work we believe that our TOC measurements are not of inferior quality compared to TOPS measurements and provide further evidence below.

Ability to detect small changes in TOC

As described in the manuscript (P23, L26ff.), our data do not indicate oscillations in TOC that may have been triggered by bow waves from the Moon's shadow. In contrast, Zerefos et al. (2000) reported peak-to-peak variation in TOC of about 1 %. while Zerefos et al. (2007) reported a peak-to-peak amplitude of 2.0–3.5 %. Finally Mims and Mims (1993) describe a peak-to-peak amplitude of up to 5 DU (1.7 %). We show in the following that our method is capable of detecting fluctuations of this magnitude.

By analyzing the values of our ozone look-up table, we determined that a 1 % change in the ratio of the response-weighted global irradiance at 340 and 305 nm results in a TOC change of 1.1 DU at the start of the eclipse. At the end of the eclipse (when the SZA is smaller) a 1 % change in the ratio leads to a change of 1.6 DU. As described in the manuscript, ozone calculations are based on the average of 45 seconds (not 2 minutes as stated by the reviewer) of global spectral irradiance measurements that are sampled at 1 Hz. By calculating the standard deviation of these samples

1 and using standard error propagation, we calculated an uncertainty (confidence interval of 95 %)
2 for the ratio of measurements at 340 and 305 nm of 0.13 % for the start and 0.06 % for the end of
3 the eclipse. By combining these uncertainty estimates with the sensitivity of the TOC to changes
4 in this ratio, we determined that our measurements are able to detect changes in ozone of 0.14 DU
5 at the start and 0.10 DU at the end of the eclipse. Since the average TOC on 21 August 2017 was
6 about 298 DU, these absolute changes translate to relative changes of 0.05 % and 0.03 %,
7 respectively. Our method is therefore well capable to detect changes of the magnitude of 1 to 3.5
8 % reported by Zerefos et al. (2000, 2007) and Mims and Mims (1993).
9

10 We agree with the reviewer that measurements at 300 nm are more sensitive to changes in ozone
11 than measurements at 305 nm. However, measurements at 300 nm are also noisier than
12 measurements at 305 nm because the irradiance at 300 nm is smaller than that at 305 nm by
13 factors of 10 (end of eclipse) to 30 (start of eclipse). Moreover, small errors in the
14 characterization (e.g., center wavelength) of the filters functions have a larger effect at shorter
15 wavelength. Since we do not have access to a TOPS or Microtops, we cannot determine whether
16 these instruments are really superior to the GUVis-3511 in determining the TOC as the reviewer
17 asserts. (Of note, the shortest wavelength of a Dobson is 305.5 nm and if measurements at 300
18 nm would be of great advantage, these instruments would likely use a shorter wavelength.) In any
19 case, our analysis above illustrates that our instrument is sensitive enough for detecting bow-
20 waved induced TOC variations of the proposed magnitude.
21

22 Effect of aerosols on ozone retrieval

23 The look-up table for TOC retrievals was calculated with the same model parameters that were
24 used to convert response weighted irradiance measurements to spectral irradiance as described in
25 the Supplement to this publication. Specifically, aerosol extinction was parameterized with
26 Ångström's turbidity formula by setting the Ångström coefficients α and β to 2.10 and 0.0394,
27 respectively. These values were derived from the instrument's measurement of direct solar
28 irradiance at 19:00:30 after the end of the eclipse. At the start (16:02:58; see Fig. 8b of
29 manuscript), the Ångström coefficients were 1.96 and 0.0570, respectively. To quantify the effect
30 of changing aerosol conditions on TOC retrievals, we modeled the global spectral irradiance for
31 16:02:58 using either the Ångström coefficients used for the TOC look-up table or the
32 coefficients applicable to this time. The difference in AOD changed the ratio of global spectral

1 irradiance at 340 and 305 nm by only 0.12 %. Using the same sensitivity factor discussed above,
2 we determined that the resulting bias in TOC is 0.13 DU or 0.04 %.

3
4 Limb-darkening corrected AODs (Fig. 8a), suggest that the largest AOD during the eclipse
5 occurred at 16:52 when the Ångström coefficients were 1.95 and 0.0788, respectively. (This
6 estimate is somewhat uncertain due to the uncertainty of the limb-darkening correction.) We
7 calculated the aerosol effect in a similar way as before and conclude that the elevated AOD at
8 16:52 increases the retrieved TOC values by 1.5 DU or 0.5 %. Again, this value is considerably
9 smaller than the fluctuations reported by Zerefos et al. (2000, 2007). Yet, aerosols can explain 1.5
10 DU of the 5.0 DU increase in limb-darkening corrected TOC measurements that can be seen in
11 Fig 8a of the manuscript between 16:00 and 17:00.

12
13 We also like to point out that TOC retrievals from global irradiance is less sensitive to changes in
14 AOD than retrievals using the direct beam (such as those utilized by the TOPS instruments)
15 because photos that are removed from the direct beam by aerosols contribute to the global
16 irradiance. So direct measurements are not necessarily better.

17 18 Sampling frequency

19 TOC measurements discussed in the manuscript are available at a rate of one value every 2
20 minutes. It is therefore not possible to determine whether the moon's shadows resulted in
21 fluctuations in TOC on a shorter time scale. However, oscillations reported by Zerefos et al.
22 (2000) had a period of 20 minutes and those reported by Zerefos et al. (2007) had periods ranging
23 between 30 and 40 minutes. Zhang et al. (2017) reported ionospheric bow waves, which
24 manifested themselves as electron content disturbances, with a period of about 25 minutes during
25 the solar eclipse of 21 August 2017. If these bow waves had also affected the ozone layer, these
26 oscillations should have been detectable with our sampling frequency. Finally, Mims and Mims
27 (1993) report that TOC measurements taken during the total solar eclipse of 11 July 1991 show
28 three fluctuations, which began 700 seconds after the third contact with durations of 378, 270 and
29 432 seconds, respectively. It is curious that these oscillations had a much shorter duration than
30 those reported by Zerefos et al., (2000, 2007) and Zhang et al. (2017). In any case, oscillations of
31 this duration should still shown up in our 2-minute data considering the low noise in our data
32 discussed above and illustrated in Fig. 9, in particular between the 3rd and 4th contact.

Concluding remarks

We did not “rule out” the findings of the paper by Mims and Mims (1993) as stated by the reviewer. Instead, we simply stated that our data do not support the observation by Zerefos (2000; 2007) and Mims and Mims (1993), and concluded that the question of whether or not bow waves from the Moon’s shadow can lead to variations in TOC is still up for debate. We suggested that this debate could be settled by performing limb-darkening-corrected measurements of TOC with different instrument types during one of the upcoming solar eclipses. If such measurements were to show fluctuations in TOC with the same magnitude and timing, the effect of bow waves on TOC could be convincingly demonstrated. We still believe that this is a reasonable path forward and might stimulate future research.

Changes to manuscript

The following text was added to Sect. 5.2 of the manuscript:

“By analyzing the sensitivity of TOC to changes in the irradiance ratio and by quantifying the noise in the measurements of the 305 and 340 nm channel we determined that our measurements are able to detect changes in ozone of 0.14 DU (0.05 %) at the start and 0.10 DU (0.03 %) at the end of the eclipse at 95 % confidence level for constant aerosol conditions. The additional uncertainty in TOC values due to changing AODs was determined to be 1.5 DU (0.5 %).”

In Sect. 7.1. (now Sect. 7.3), the following sentence:

“Results corrected for the LD effect (thick lines in Fig. 8a) indicate that the AOD was monotonically decreasing over the period of the eclipse (from 0.41 to 0.32 at 319 nm and from 0.05 to 0.04 at 1018 nm) without a spurious spike near the time of totality.”

was replaced with:

“Results corrected for the LD effect (thick lines in Fig. 8a (now Fig. 12a)) indicate that the AOD was increasing between the 1st contact and 16:52 (from 0.41 to 0.58 at 319 nm and from 0.05 to 0.07 at 1018 nm) and then monotonically decreasing to 0.34 at 319 nm and 0.04 at 1018 nm at the end of the eclipse. Corrected results do not have a spurious spike near the time of totality.”

In Sect. 7.2 (now Sect. 7.4), the following sentence:

“(We have no explanation for the increase in TOC of about 5 DU between 16:00 and 17:00 other than natural variability)”

1 was replaced with:

2 “The increase in TOC of about 5 DU between 16:00 and 17:00 can partly be explained by
3 increasing AODs over this period, which cause a high-bias in the ozone retrievals.”
4

5 In Sect. 8.3, the following sentence:

6 “Bow waves with wavelengths between of 300 and 400 km and a period of about 25
7 minutes have indeed been observed during the solar eclipse of 21 August 2017 (Zhang et
8 al., 2017).”

9 was replaced with:

10 “Ionospheric bow waves with wavelengths between of 300 and 400 km and a period of
11 about 25 minutes, which manifested themselves as electron content disturbances, have
12 indeed been observed during the solar eclipse of 21 August 2017 (Zhang et al., 2017). If
13 these bow waves had affected the ozone layer, these oscillations should have been
14 detectable with our sampling frequency of one TOC value every two minutes.”
15

16 Also in Sect. 8.3, the following sentence:

17 “These small variations are well within the natural variability of the TOC.”

18 was replaced with:

19 “These small variations are well within the natural variability of the TOC and the
20 uncertainty of our TOC retrieval of 1.5 DU (0.5 %), which is mainly caused by the effect
21 of changing aerosols on ozone calculations.”

Response to 2nd comments by Forrest M. Mims III, posted on 21 December 2018

General remarks to comment by reviewer

We thank Mr. Mims for his additional comments but feel that many remarks are beyond the scope of the paper. Our paper is about the importance of applying corrections for solar limb darkening when observing a solar eclipse; the question of whether or not bow waves from the Moon's shadow may result in fluctuations in total ozone column (TOC); and the question of whether or not the ratio of direct-to-diffuse irradiance changes appreciably during the period of a solar eclipse (excluding the period near totality). The paper is NOT about the best, most accurate, or most precise method to measure TOC.

In our response to the first comments by the reviewer (posted on 24 November 2018), we provided new evidence that our method of measuring TOC during the eclipse is precise enough for detecting potential changes in TOC from bow waves. In brief, we concluded that the noise in our measurements is low enough for detecting relative changes in TOC of larger than 0.05 %. In addition, we calculated that changing aerosol concentrations during the time of our observations result in an additional uncertainty in TOC of 1.5 DU or 0.5 %. Even if our uncertainty estimates were too optimistic, for example due to an unknown systematic error in our TOC retrieval method, it would be highly unlikely that variations in our calculated TOC values would anti-correlate with real variations in TOC triggered by bow waves such that the resulting TOC measurements after the 3rd contact become basically flat, with a variation of only ± 1 DU or ± 0.33 % (see Fig. 9 of original manuscript). For comparison, the peak-to-peak amplitude attributed to bow waves reported by Zerefos et al. (2007) was 2.0–3.5 %, and the peak-to-peak amplitude reported by Mims and Mims (1993) was 1.7 %.

In conclusion, we cannot rule out that the Moon's shadow led to variations of TOC in the order of ± 0.3 % during the eclipse observed by us, but note that this upper limit is considerably lower than the fluctuations reported by Zerefos et al. (2007) and Mims and Mims (1993).

Changes to manuscript

In the abstract, the following sentence:

1 “In contrast to results of observations from earlier solar eclipses, no fluctuations in TOC
2 were observed that could be attributed to gravity waves.”
3 will be replaced with:

4 “In contrast to results of observations from earlier solar eclipses, no fluctuations in TOC
5 were observed that could be unambiguously attributed to gravity waves.”
6

7 In Sect. 8.3., the following sentence:

8 “Our data do not support the observation by Zerefos (2000; 2007) and Mims and Mims
9 (1993) that bow waves from the Moon’s shadow lead to oscillations in TOC.”

10 will be replaced with:

11 “Our data do not support the observation by Zerefos (2007) and Mims and Mims (1993)
12 that bow waves from the Moon’s shadow may lead to oscillations in TOC with a peak-to-
13 peak amplitude exceeding 1.5 %.”
14

15 **Comment by reviewer:**

16 TWO KEY POINTS: The authors have made several important revisions to their paper, but they
17 simply must remove their erroneous assertions to the effect that: (1) the signal at 300 nm is noisy,
18 which suggests poor measurements by TOPS (which used 300 nm and 305 nm) and (2) full-sky
19 measurements of the ozone layer are “similar” to the direct sun measurements employed by
20 Dobsons (and TOPS, Microtops, Brewers and Pandoras). These inappropriate claims and their 2-
21 minute measurement time support their general assertions that raises doubts about the papers by
22 me and others.
23

24 **Authors’ Response**

25 Regarding (1): We do not assert anywhere in the paper that signals of the TOPS instruments are
26 noisy or that measurements of this instrument are of poor quality. In fact, we do not even mention
27 “TOPS” in the manuscript. The discussion of the noise characteristics of the TOPS instrument is
28 only part of the reviewer’s first post and our response. It is and will not be part of the paper.
29

30 Regarding (2): We do not state in the manuscript that “full-sky measurements of the ozone layer
31 are ‘similar’ to the direct sun measurements employed by Dobsons (and TOPS, Microtops,
32 Brewers and Pandoras)”. We do not compare the accuracy of our TOC measurements with that of
33 other methods. Like in the case of (1), the discussion on the quality of the different methods of
34 measuring ozone is only part of the reviewer’s first post and our response.

Changes to manuscript

None.

Comment by reviewer:

1. OPTIMUM WAVELENGTH SELECTION FOR MEASURING TOTAL COLUMN OZONE DURING A SOLAR ECLIPSE: The authors claim the 300-nm minimum employed by TOPS provides a noisy signal. They erroneously observe that: “Of note, the shortest wavelength of a Dobson is 305.5 nm and if measurements at 300 nm would be of great advantage, these instruments would likely use a shorter wavelength.” The 305.5-nm Dobson minimum wavelength was selected due to the use of the instrument across a wide band of latitudes. However, there is ample signal at 300 nm at my site (29.9 N), as demonstrated by the instrument’s detection of an error in NASA’s Nimbus-7 TOMS ozone instrument (Mims, Nature, 1993). The 300 nm signal was especially strong at 22 N during the 1991 solar eclipse reported in my paper in GRL. The authors might be right about the 300-nm signal at their northerly location. But they cannot compare what might have been a noisy 300 nm signal at their northerly 44.36 N site to the much higher amplitude 300-nm signal at the 22 N site for the eclipse I measured. Furthermore, DeLuisi and others have demonstrated that wavelengths below 305 nm provide more accurate ozone measurements than higher wavelengths. (This explains why TOPS found the satellite error.)

Authors’ Response

Similar to the last comment, we do not discuss the TOPS instrument in our paper. The quote in the reviewer’s comment above is again from our response to the reviewer’s first post. Also, we do not discuss the best wavelengths to be used to calculate TOC for the location of “our” eclipse. We simply state in the manuscript that TOCs were calculated from the measurements of the GUVIS-3511’s channels at 305 and 340 nm.

Changes to manuscript

None.

Comment by reviewer:

Of special concern is this from the author’s abstract: “The total ozone column (TOC) was derived from measurements of global irradiance at 306 and 340nm.” While the 306 nm minimum is appropriate for 44.36 N, the 340 nm upper wavelength is far too high, for it allows for significant

aerosol errors. The closely-spaced 300 nm and 305 nm wavelengths of TOPS nearly eliminated the aerosol error, which explains this instrument's excellent accuracy when compared with hundreds of satellite measurements and an EPA Brewer for 60 days at my site. Consider this abstract by Saunders et al in High-Precision Atmospheric Ozone Measurements Using ... trial spectral irradiances between 290 and 305 nm (JGR Atmospheres 1984 <https://doi.org/10.1029/JD089iD04p05215>) "Abstract "It is shown theoretically that many errors are significantly less when determining atmospheric ozone thicknesses from measurements of solar terrestrial spectral irradiance in the wavelength region between 290 and 305 nm as compared to the to the 305- to 340-nm region employed by the Dobson spectrophotometer. In order to test this conclusion experimentally, an elaborate set of state-of-the-art measurements have been made in the shorter wavelength region in Gainesville, Florida, between June 13 and June 18, 1980. Details of these measurements, including an extensive error analysis, are presented and indicate that such short-wavelength measurements, particularly between 295 and 305 nm, can be used to detect long-term changes of atmospheric ozone with an uncertainty not exceeding 1%. Observing conditions restricted the Gainesville measurements to zenith angles of less than 35°. Further investigations are required to determine the shortest wavelength that can be used at significantly greater zenith angles."

Authors' Response

We agree with the reviewer that aerosols will lead to systematic errors in TOC measurements if the wavelengths used for the retrieval are far apart. However, we have estimated the uncertainty in our TOC retrievals for the period of interest and have concluded that variations in aerosols during the period of the eclipse cause an uncertainty in TOC of only 1.5 DU or 0.5 %. We note that this uncertainty refers to the precision of TOC measurements (the metric of relevance to the paper), not absolute accuracy, which could be worse. However, our TOC retrievals agree to within 3 DU (or 1%) with OMI, suggesting that also the accuracy of our data is within acceptable limits.

The discussion of whether or not TOC measurements should be based on wavelengths in the 290 to 305 nm range is of little relevance to the paper because the GUVIS-3511 radiometer has no channels with wavelengths below 305 nm. In addition, while TOC measurements using wavelengths shorter than 305 nm could indeed be more accurate, as the JGR quoted by the reviewer suggests, we like to point out that a higher accuracy can only be achieved if the filters of the instrument in question are well characterized. No filter instrument measures at exactly a

1 nominal wavelength and all real instruments (including the TOPS) use filters with a finite
2 bandpass. If a hypothetical channel that is supposed to measure at 300 nm measures in fact at
3 300.5 nm, TOC errors will result. We are not implying that this is the case for the TOPS
4 instrument, but just like to point out that performing ozone measurements with channels at 300
5 and 305 nm can potentially lead to errors that could be comparable in magnitude to those
6 affecting measurements using channels that are farther apart, and as a result are more sensitive to
7 aerosols. Again, we cannot, and do not want to, assess the quality of TOPS measurements
8 because we do not use data of this instrument in the paper and are not familiar with its
9 characteristics.

10
11 The sentence “The total ozone column (TOC) was derived from measurements of global
12 irradiance at 306 and 340nm.” in the abstract does not judge whether this wavelength selection is
13 the most appropriate. It simply states what was done and we see no reason to change it.

14
15 **Changes to manuscript**

16 None.

17
18 **Comment by reviewer:**

19 2. FULL-SKY VS. DIRECT SUN TOTAL OZONE MEASUREMENTS: The authors state in
20 their paper: “The method of calculating TOC from measurements of global irradiance (instead of
21 direct irradiance as it is typically done for Dobson, Brewer, TOPS, and Microtops instruments)
22 was first proposed by Stamnes et al. (1991). We found that the accuracy of TOCs derived from
23 global irradiance is similar to that of data from Dobson instruments or satellite (TOMS, OMI)
24 observations if the look-up table takes local conditions into account (ozone profile, albedo,
25 elevation, etc.) (Bernhard et al., 2005b).” The authors suggest that global irradiance provides
26 TOC data “. . . similar to that of data from Dobson instruments or satellite. . . .” But the authors
27 provide no data or citations to support this assertion, while leaving open the counter suggestion
28 that Dobsons (and TOPS, Brewers and Pandoras) could be replaced by much simpler instruments
29 that measure global irradiance and require no tracking and pointing.

30
31 **Authors’ Response**

32 We do not state in our paper:

33 “The method of calculating TOC from measurements of global irradiance (instead of
34 direct irradiance as it is typically done for Dobson, Brewer, TOPS, and Microtops

instruments) was first proposed by Stamnes et al. (1991). We found that the accuracy of TOCs derived from global irradiance is similar to that of data from Dobson instruments or satellite (TOMS, OMI) observations if the look-up table takes local conditions into account (ozone profile, albedo, elevation, etc.) (Bernhard et al., 2005b).”

as asserted in the comment by the reviewer above. This quote is taken from our comment to the first post of the reviewer. Instead, we simply state in the paper:

“This method was first proposed by Stamnes et al. (1991) and was validated for GUV instruments by Bernhard et al. (2005a).”

We are puzzled by the reviewer’s comment:

“The authors suggest that global irradiance provides TOC data “. . . similar to that of data from Dobson instruments or satellite. . . .” But the authors provide no data or citations to support this assertion, while leaving open the counter suggestion that Dobsons (and TOPS, Brewers and Pandoras) could be replaced by much simpler instruments that measure global irradiance and require no tracking and pointing.”

because the paper and the response to the post of the reviewer does include two citations (i.e., Bernhard et al., 2005a, and Bernhard et al., 2005b) where TOC measurements from global irradiance are compared with Dobson direct measurements and satellite (TOMS, OMI) observations.

In Bernhard et al., 2005b, we conclude that:

“When Dobson measurements are corrected for the temperature dependence of the ozone absorption cross section and accurate air mass calculations are implemented, data from the three instruments agree with each other to within $\pm 2\%$ on average and show no significant dependence on SZA or total ozone.”

The “three instruments” quoted above refer to the Dobson; our SUV-100 spectroradiometer, which measures global irradiance; and TOMS. The reviewer may not agree with this conclusion, but it is false to assert that “the authors provide no data or citations to support this assertion”.

We further like to point out that the method of retrieving TOC from spectra of global irradiance measured by SUV-100 radiometers has been published by Bernhard et al. (2003). This paper also includes an uncertainty estimate of the method. The abstract ends with:

“On average, the new algorithm generates ozone values in spring 2.2 % lower than TOMS observations and 1.8 % higher than Dobson measurements. From the uncertainty

1 budget and the comparison with TOMS and Dobson it can be concluded that ozone
2 values retrieved from global UV spectra have a similar accuracy as observations with
3 standard instrumentation used for ozone monitoring.”
4

5 **Changes to manuscript**

6 In Sect. 5.2., we will replace

7 “This method was first proposed by Stamnes et al. (1991) and was validated for
8 GUV instruments by Bernhard et al. (2005a)”

9 with

10 “The method of calculating TOC from measurements of global irradiance was
11 first proposed by Stamnes et al. (1991) and was further validated by Bernhard et
12 al. (2003; 2005b). The application of the method to GUV instruments was
13 described by Bernhard et al. (2005a).”
14

15 **Comment by reviewer:**

16 Of course, this is not the case. The authors have gone much, much too far in suggesting their
17 global, full-sky data is “similar” to direct sun measurements made by the recognized standard for
18 nearly a century and all other ozone instruments. I have always been intrigued by the prospect of
19 accurate ozone retrievals by pairs of global UV measurements at closely-space wavelengths. I
20 urge the authors to prepare a detailed paper about their claim that global TOC measurements are
21 “similar” to those by traditional direct sun instruments. An ideal comparison would be with
22 Brewers, which measure both direct sun and global TOC. Meanwhile, it is inappropriate to make
23 an unsupported claim about “similar” results.
24

25 **Authors’ Response**

26 The reviewer suggests to prepare a detailed paper about our claim that global TOC measurements
27 are “similar” to those by traditional direct sun instruments. This paper has already been written.
28 In fact, there are two: Bernhard et al. (2003; 2005b) and we therefore don’t agree with the
29 reviewer’s conclusion that it is inappropriate to make an unsupported claim about “similar”
30 results.

31 We do not advocate in any of our papers that the established network of Dobson and Brewer
32 instrument should be replaced with instruments measuring global irradiance. The “global
33 irradiance method” is only sufficiently accurate if the ozone profile is known with sufficient

accuracy, in particular for large solar zenith angles. So the method is not as independent as direct measurements that rely on Beer-Lambert's law and hence do not require knowledge of the profile. However, when there is cloud cover, the direct method cannot be used and Stamnes et al. (1991) showed that the "global" method is equally accurate as the zenith sky method used by Dobsons when the Sun is obstructed by clouds.

As a side note, because TOC retrievals from global irradiance depend on the ozone profile, the profile can in fact be determined from such measurements using a variant of the Umkehr method. We have recently published this "Global-Umkehr method" (Bernhard et al., 2017). Again, we not propose that this method is superior to the standard Umkehr method, which relies on zenith sky observations. The paper is just a prove of concept and makes the Umkehr method available to locations with global irradiance measurements. The abstract of Bernhard et al. (2017) concludes with "Total ozone columns (TOCs) calculated from the retrieved profiles agree to within 0.7 ± 2.0 % ($\pm 1\sigma$) with TOCs measured by the Ozone Monitoring Instrument on board the Aura satellite." This demonstrates again the good accuracy of the TOC retrievals from global measurements if its done right.

Changes to manuscript

None.

Finally, we like to conclude that most of the discussion above is of little relevance to our paper. The only important question is whether or not our TOC retrievals from global irradiance measurements of the GUVIS-3511 are of sufficient precision to detect fluctuations in TOC that could be triggered by bow waves. Using a sensitivity analysis, we concluded that this is the case. Based on our results, we cannot rule out that there is an effect with an amplitude of ± 1 DU or ± 0.33 %. We simply conclude that we did not observe fluctuations in TOC that could be unambiguously attributed to gravity waves. It is possible that the magnitude of gravity wave effects varies from eclipse to eclipse, explaining the discrepancy in the data by us, Zerefos et al. (2007) and Mims and Mims (1993), but this is speculation.

Changes to manuscript

None.

References

- Basher, R. E., Review of the Dobson spectrophotometer and its accuracy, WMO Global Ozone Res. Monit. Proj. Rep. 13, World Meteorol. Organ., Geneva, Switzerland, 1982.
- Bernhard, G., Booth, C. R. and McPeters R. D., Calculation of total column ozone from global UV spectra at high latitudes, *J. Geophys. Res.*, 108(D17), 4532, doi:10.1029/2003JD003450, 2003.
- Bernhard, G., Booth, C. R., and Ehramjian, J. C.: Real-time ultraviolet and column ozone from multichannel ultraviolet radiometers deployed in the National Science Foundation's ultraviolet monitoring network, *Opt. Eng.*, 44(4), 041011-1, <https://doi.org/10.1117/1.1887195>, 2005a.
- Bernhard, G., Evans, R. D., Labow, G. J., and Oltmans, S. J.: Bias in Dobson total ozone measurements at high latitudes due to approximations in calculations of ozone absorption coefficients and air mass, *J. of Geophys. Res.*, 110, D10305, <https://doi.org/10.1029/2004JD005559>, 2005b.
- Bernhard, G., Petropavlovskikh, I., and Mayer, B.: Retrieving vertical ozone profiles from measurements of global spectral irradiance, *Atmos. Meas. Tech.*, 10(12), 4979-4994, <https://doi.org/10.5194/amt-10-4979-2017>, 2017.
- Bojkov, R. D.: The ozone variations during the solar eclipse of 20 May 1966, *Tellus*, 20(3), 417–421, <https://doi.org/10.1111/j.2153-3490.1968.tb00382.x>, 1968.
- Emde, C., and Mayer, B: Simulation of solar radiation during a total eclipse: a challenge for radiative transfer, *Atmos. Chem. Phys.*, 7(9), 2259–2270, <https://doi.org/10.5194/acp-7-2259-2007>, 2007.
- Gröbner, J., Kazadzis, S., Kouremeti, N., Doppler, L., Tagirov, R. and Shapiro, A. I.: Spectral solar variations during the eclipse of March 20th, 2015 at two European sites. In AIP Conference Proceedings, 1810(1), p. 080008, <https://doi.org/10.1063/1.4975539>, 2017.
- Hooker, S. B., Bernhard, G., Morrow, J. H., Booth, C. R., Comer, T., Lind, R. N., and Quang, V.: Optical Sensors for Planetary Radiant Energy (OSPRey): calibration and Validation of Current and Next-Generation NASA Missions., NASA Goddard Space Flight Center, NASA/TM–2011–215872, 2012.
- Kazantzidis, A., Bais, A. F., Emde, C., Kazadzis, S., and Zerefos, C. S.: Attenuation of global ultraviolet and visible irradiance over Greece during the total solar eclipse of 29 March 2006, *Atmos. Chem. Phys.*, 7(23), 5959–5969, <https://doi.org/10.5194/acp-7-5959-2007>, 2007.

- 1 Koepke, P., Reuder, J., and Schween, J.: Spectral variation of the solar radiation during an
2 eclipse, *Meteorol. Z.*, 10(3), 179–186, <https://doi.org/10.1127/0941-2948/2001/0010-0179>,
3 2001.
- 4 Krotkov, Nickolay A., Pawan K. Bhartia, Jay R. Herman, James R. Slusser, Gordon J. Labow,
5 Gwendolyn R. Scott, George T. Janson, Tom Eck, and Brent N. Holben. "Aerosol ultraviolet
6 absorption experiment (2002 to 2004), part 1: ultraviolet multifilter rotating shadowband
7 radiometer calibration and intercomparison with CIMEL sunphotometers." *Optical
8 Engineering* 44, no. 4 (2005): 041004.
- 9 Mims III, F. M. and Mims, E. R.: Fluctuations in column ozone during the total solar eclipse of
10 July 11, 1991, *Geophys. Res. Lett.*, 20(5), 367–370, <https://doi.org/10.1029/93GL00493>,
11 1993.
- 12 Morrow, J. H., Hooker, S. B., Booth, C. R., Bernhard, G., Lind, R. N., and Brown, J. W.:
13 Advances in measuring the apparent optical properties (AOPs) of optically complex waters,
14 NASA/TM–2010–215856, National Aeronautics and Space Administration, Goddard Space
15 Flight Center, 2010.
- 16 Neckel, H.: Analytical Reference Functions $F(\lambda)$ for the Sun's Limb Darkening and Its Absolute
17 Continuum Intensities ($\lambda\lambda$ 300 to 1100 m). *Solar Physics*, 229(1), 13–33, 2005.
- 18 Stamnes, K., Slusser, J., and Bowen, M.: Derivation of total ozone abundance and cloud effects
19 from spectral irradiance measurements, *Appl. Opt.*, 30(30), 4418–4426,
20 <https://doi.org/10.1364/AO.30.004418>, 1991.
- 21 Witthuhn, J., Deneke, H., Macke, A., and Bernhard, G.: Algorithms and uncertainties for the
22 determination of multispectral irradiance components and aerosol optical depth from a
23 shipborne rotating shadowband radiometer, *Atmos. Chem. Phys.*, 10(2), 709–730,
24 <https://doi.org/10.5194/amt-10-709-2017>, 2017.
- 25 Zerefos, C. S., Balis, D. S., Meleti, C., Bais, A. F., Tourpali, K., Kourtidis, K., K. Vanicek, K.,
26 Cappellani, F., Kaminski, U., Colombo, T., and Stübi, R.: Changes in surface solar UV
27 irradiances and total ozone during the solar eclipse of August 11, 1999, *J. Geophys. Res.*,
28 105(D21), 26463–26473, <https://doi.org/10.1029/2000JD900412>, 2000.
- 29 Zerefos, C. S., Balis, D. S., Zanis, P., Meleti, C., Bais, A. F., Tourpali, K., D. Melas, D., Ziomas
30 I., Galani E., Kourtidis, K., Papayannis A., and Gogosheva Z.: Changes in surface UV solar
31 irradiance and ozone over the Balkans during the eclipse of August 11, 1999, *Adv. Space
32 Res.*, 27(12), 1955–1963, [https://doi.org/10.1016/S0273-1177\(01\)00279-4](https://doi.org/10.1016/S0273-1177(01)00279-4), 2001.
- 33 Zerefos C. S., Gerasopoulos E., Tsagouri I., Psiloglou B. E., Belehaki A., Herekakis T., Bais A.,
34 Kazadzis S., Eleftheratos C., Kalivitis N., and Mihalopoulos N.: Evidence of gravity waves

1 into the atmosphere during the March 2006 total solar eclipse, *Atmos. Chem. Phys.*, 7(18),
2 4943–4951, <https://doi.org/10.5194/acp-7-4943-2007>, 2007.
3 Zhang, S.-R., Erickson, P. J., Goncharenko, L. P., Coster, A. J., Rideout, W., and Vierinen, J.:
4 Ionospheric bow waves and perturbations induced by the 21 August 2017 solar eclipse,
5 *Geophys. Res. Lett.*, 44(24), <https://doi.org/10.1002/2017GL076054>, 2017.

Measurements of spectral irradiance during the solar eclipse of 21 August 2017: reassessment of the effect of solar limb darkening and of changes in total ozone

Germar Bernhard¹, Boyan Petkov²

¹Biospherical Instruments Inc., San Diego, CA 92110, USA

²Institute of Atmospheric Sciences and Climate (ISAC) of the Italian National Research Council (CNR), I-40129 Bologna, Italy

Correspondence to: Germar Bernhard (bernhard@biospherical.com)

Abstract. Measurements of spectral irradiance between 306 and 1020 nm were performed with a GUVis-3511 multi-channel filter radiometer at Smith Rock State Park, Oregon, during the total solar eclipse of 21 August 2017. The radiometer was equipped with a shadowband, allowing to separate the global (sun and sky) and direct components of solar radiation. Data were used to study the wavelength-dependent changes of solar irradiance at Earth’s surface. Results were compared with theoretical predictions using three different parameterizations of the solar limb darkening (LD) effect, which describes the change of the solar spectrum from the Sun’s center to its limb. Results indicate that the LD parameterization that has been most widely used during the last 15 years underestimates the LD effect, in particular at UV wavelengths. The two alternative parameterizations are based on two independent sets of observations from the McMath-Pierce Solar Telescope. When these parameterizations are used, the observed and theoretical LD effects agree to within 4 % for wavelengths larger than 400 nm and occultation of the solar disk of up to 97.8 %. Maximum deviations for wavelengths between 315 and 340 nm are 7 %. These somewhat larger differences compared to the visible range may be explained with varying aerosol conditions during the period of observations. Aerosol optical depth (AOD) and its wavelength dependence was calculated from measurements of direct irradiance. When corrected for the LD effect, AOD decreases over the period of the eclipse: from 0.41 to 0.34 at 319 nm and from 0.05 to 0.04 at 1018 nm. These results show that AODs can be accurately calculated during an eclipse if the LD effect is corrected. The total ozone column (TOC) was derived from measurements of global irradiance at 306 and 340 nm. Without correction for the LD effect, the retrieved TOC increases by 20 DU between the 1st and 2nd contact of the eclipse. With LD correction, the TOC remains constant to within natural variability (± 2.6 DU or ± 0.9 % between 1st and 2nd contact and ± 1.0 DU or ± 0.3 % between 3rd and 4th contact). In contrast to results of observations from earlier solar eclipses, no fluctuations in TOC were observed that could be unambiguously attributed to gravity waves, which can be triggered by the supersonic speed of the Moon’s shadow across the atmosphere. Furthermore, systematic changes in the ratio of direct and global irradiance that could be attributed to the solar eclipse were not observed, in agreement with results of three-dimensional radiative transfer models. Our results advance the understanding of the effects of solar LD on the spectral

Deleted: monotonically

Deleted: 2

Deleted: . This finding agrees

Deleted: but contradicts reports from earlier observations, which indicate that the diffuse-to-direct ratio may change by 30 %.

irradiance at Earth’s surface, the variations of ozone during an eclipse, and the partitioning of solar radiation in direct and diffuse components.

1 Introduction

A total solar eclipse could be observed on 21 August 2017 across the United States, from Oregon in the West to South Carolina in the East (Pasachoff, 2017). We performed measurements of global (direct Sun plus sky) and diffuse (sky only) spectral irradiance during this event with a GUVis-3511 multi-channel radiometer, which was equipped with a shadowband to separate the two components. Data collected during this campaign can be used for studying processes initiated in the atmosphere by the passing of the Moon’s shadow and for validating three-dimensional radiative transfer (3DRT) calculations that simulate the irradiance at Earth’s surface during the period of totality (Emde and Mayer, 2007). Similar observations have been used during previous solar eclipses to study wavelength-dependent changes in spectral irradiance (Blumthaler et al., 2006; Kazadzis et al., 2007; [Kazantzidis et al., 2007](#)); the change in the contribution of the diffuse irradiance to the global irradiance over the period of the eclipse (Zerefos et al., 2000, 2001); short-term and longer-lasting fluctuations in the total ozone column (TOC) (Antón et al., 2010; Kazadzis et al., 2007; [Kazantzidis et al., 2007](#); Mateos et al., 2014; Mims and Mims, 1993; Zerefos et al., 2000; 2001; 2006); variations in the NO₂ column (Adams et al., 2010); and comparison of measurements during totality with 3DRT model results (Kazantzidis et al., 2007). We revisit some of these issues by focusing on the wavelength dependence of solar limb darkening (LD), ozone observations, and the changing direct and diffuse radiation. For example, some observations of changes in the TOC during an eclipse are contradictory. Increasing TOCs were reported for measurements from Dobson spectrophotometers (Bojkov, 1968) and NILU-UV multifilter instruments (Antón et al., 2010) while measurements with Brewer photometers generally decrease as a eclipse progresses (Kazadzis et al., 2007). These differences cannot be explained with a real change in TOC but must be an artifact from either measurement or data processing.

Our measurements were also used to assess the quality of data of the GUVis-3511 radiometer, which is a relatively new instrument type. For this purpose, a new method to calibrate the instrument was developed and is described in detail in the Supplement to this publication. Data from the campaign are available for download and can, for example, be used to study variations of radiation during totality with 3DRT simulations.

2 Location and local conditions

The total eclipse was observed at Smith Rock State Park, located in Oregon between Bend and Madras. The partial eclipse started at 16:06:27 UT (1st contact, local time of 9:06:27 Pacific Daylight Time (PDT)) and ended at 18:41:02 UT (4th contact, 11:41:02 PDT). Totality occurred for one minute and 25 seconds, between 17:19:42 (2nd contact, 10:19:42 PDT) and

17:21:07 UT (3rd contact, 10:21:07 PDT). All times from here onward refer to UT. The GUVis-3511 radiometer was set up at 44.362950° N and 121.139667° W, 867 m above sea level. The site was approximately 37 km south of the centerline of the Moon's shadow (e.g., the line where the duration of totality is the longest). The ground surrounding the instrument was covered by dry grass. The park's prominent mountain peaks were mostly in the west, north, and northwest of the instrument (Fig. 1), and extended up to 12.5° above the horizon. The average extension was 5.75°. By assuming isotropic sky radiance distribution, we determined that objects above the horizon reduced the (cosine-weighted) diffuse irradiance by about 1.1 %. Data were not corrected for this effect.



Fig. 1. Panorama as seen from instrument location. Mountains are located in the west (W), north (N), and northeast, and extend up to 12.5° above the horizon. Trees restrict the horizon in eastern (E) direction by up to 9°. There is little (<5°) obstruction towards the south (S). The green arrow marks the mountain used for calibrating the elevation scale (red lines).

The state of Oregon experienced a total of 1069 reported wildfires in 2017, burning a total area of 451,863 acres or 1829 km² (https://en.wikipedia.org/wiki/2017_Oregon_wildfires). There were several active wildfires in the vicinity of the observation site and the air was filled with smoke and aerosols during the days preceding the eclipse. However, a few hours before the start of the eclipse, the wind direction changed and the aerosol loading decreased substantially. Aerosol optical depths (AOD) measured shortly before and after the eclipse period are discussed in Sect. 7.3. The sky was free of clouds in the direction of the Sun with small clouds lingering only near the horizon.

3 Instrumentation

Measurements were performed with a GUVis-3511 multi-channel filter radiometer (Seckmeyer et al., 2010) designed and built by Biospherical Instruments Inc (BSI). The system was set up on a sturdy tripod (Fig. 2) and powered by a 12 V dry-cell car battery and a sine-wave inverter.

The instrument was equipped with 18 channels with the following nominal wavelengths (nm): 305, 305, 313, 320, 340, 340, 380, 395, 412, 443, 490, 532, 555, 555, 665, 875, 940, and 1020. The spectral bandwidth of all channels is approximately 10 nm full width at half maximum (FWHM) with the exception of the two channels at 305 nm, which have a bandwidth of 18.5 nm. Spectral response functions of these channels are shown in Fig. 3 and were used for the calibration of the instrument (Sect. 4 and Supplement). Data from a 19th channel measuring photosynthetic active radiation (PAR) were not used in this study. Each channel uses a hard-coated, ion-assisted deposition interference filter plus bandpass filters for

Deleted: spectral bandwidths of approximately 10 nm and

additional out-of-band rejection that are coupled to a “microradiometer” (Morrow et al., 2010). Microradiometers consists of a silicon photodiode, three-stage preamplifier, 24 bit analogue-to-digital converter, microprocessor, and an addressable digital port. Data streams from the 19 microradiometers were combined and transmitted via a USB interface to a laptop computer. Measurements at 305, 340, and 555 nm were performed with two channels, respectively, using either a standard production photodiode (model S1226 from Hamamatsu) or an alternative photodiode (model S12698 from Hamamatsu) for evaluation purposes. The instrument’s internal temperature was stabilized to 40 ± 0.5 °C.



Fig. 2. Setup of the GUVIS-3511 radiometer at Smith Rock State Park. The radiometer (white cylinder), BioSHADE assembly and BioGPS are mounted on a tripod. Shown next to the instrument is the laptop for recording the data, the instrument's control unit, and a black case containing the power source.




Fig. 3. Normalized spectral response functions of the GUVIS-3511 radiometer discussed in this study. Functions were measured with BSI’s spectral tester (Bernhard et al., 2005a) and corrected for the finite bandpass of the tester using a simple

deconvolution routine also described by Bernhard et al. (2005a). Color coding is identical to that in Fig. 8. Note that the x-axis is logarithmic to emphasize the UV range. [A version of the figure, plotted separately for UV, visible, and IR wavelengths, is part of the Supplement.](#)

Deleted: Fig. 11

The filtered microradiometers point at the center of an irradiance collector, which features a composite diffuser made of layers of generic and porous polytetrafluoroethylene (PTFE) sheets (Hooker et al., 2012). This design leads to relatively small cosine errors (Seckmeyer and Bernhard, 1992) also in the infrared, where the scattering properties of traditional PTFE diffusers are typically degraded. The directional response of the collector is virtually independent of wavelength for wavelengths smaller than 800 nm; for larger wavelengths, the response is slightly lower (Fig. 4). [Measurements of the cosine error at wavelengths below 395 nm are affected by noise due the low UV output of the incandescent lamp used for the characterization. Corrections for the cosine error of UV channels described further below are based on the measured cosine error at 395 nm.](#) Cosine errors of all channels are smaller than $\pm 5\%$ for incidence angles smaller than 78° . The error in measuring isotropic radiation is smaller than 0.3 % for wavelengths smaller than 800 nm, and -1.8% (worst case) for the 1020 nm channel.

Deleted: (Seckmeyer and Bernhard, 1992)




Fig. 4. Cosine error of the GUVis-3511 radiometer in one azimuthal plane at wavelengths between 395 and 1020 nm. Color coding is identical to that of Fig. 8. Measurements performed at the orthogonal plane are similar. Measurements at wavelengths below 395 nm are affected by noise and are not shown for clarity. [\(A figure with cosine errors in the UV is provided in the Supplement.\)](#)

Deleted: due the low UV output of the incandescent lamp used for the characterization, and

The radiometer was equipped with a computer-controlled shadowband, called BioSHADE (Morrow et al., 2010). [The band obstructs a wedge of the sky with a width of \$15^\circ\$ when the band is vertical and \$13^\circ\$ when it is horizontal.](#) The band can either be “stowed” below the instrument’s diffuser for measuring global spectral irradiance or move slowly over the instrument to collect data for calculating the direct spectral irradiance from the solar beam with an algorithm developed for the BioSHADE (Morrow et al., 2010; Witthuhn et al., 2017). In brief, the algorithm first determines the spectral irradiance when the centers of diffuser, shadowband, and Sun are collinear. This is the spectral irradiance from the sky minus the portion of the sky that is shaded by the band. Next, the spectral irradiance from the direct Sun plus the unshaded portion of the sky is obtained by

Deleted: Hooker

Deleted: 2

analyzing several seconds of measurements when the shade from the band is close to the diffuser but not in contact with it.

These measurements are then combined with the directional test data (Fig. 4) and measurements of global spectral irradiance to calculated cosine-error-corrected global and direct spectral irradiances. [The algorithm compensates for the effect of the relatively large width of the shadowband, resulting in AOD data that have similar accuracy than those of traditional shadowband radiometers, which use measurements on either side of the Sun to correct for the portion of the sky that is obscured by the band. For example, side-band measurements of Multifilter Rotating Shadowband Radiometer \(MFRSR\) are typically performed at 9° from the center of the Sun \(Krotkov et al., 2005\) and therefore may not adequately measure circumsolar radiation. Our algorithm alleviates this problem. The uncertainty of our method was estimated by Witthuhn et al. \(2017\). AOD can be retrieved with an uncertainty of 0.02 for all channels within a 95 % confidence interval.](#)

Deleted: Fig. 3

For observing the solar eclipse, the band was programmed to perform one shadowband sweep every two minutes, between 14:36 and 19:30. Within each two-minute window, this configuration resulted in 45 seconds of global spectral irradiance measurements (with the shadowband stowed and applying a sampling rate of 1 Hz) and 75 seconds of data during which the band rotated by 180°. The sampling rate was set to 15 Hz during these “sweeps” to adequately resolve the short period when the band’s shadow moves over the diffuser. The system is also equipped with a GPS receiver, termed BioGPS, which was used to determined the geoposition specified in Sect. 2.

4 Calibration

The GUVis-3511 radiometer was calibrated using a new method described in the Supplement of this publication. Calibrated measurements report the solar spectral irradiance at the instrument’s collector at a spectral resolution of 1 nm. More specifically, calibrated measurement resemble measurements of a hypothetical spectroradiometer with a slit function $s(\lambda)$, where $s(\lambda)$ is a triangular function with a bandwidth of 1 nm FWHM. For the channels with nominal wavelengths of 305 and 313 nm, the instrument was vicariously calibrated against measurements of a SUV-100 spectrometer, which has a spectral resolution of 1 nm. All other channels were calibrated using a calibration lamp traceable to a standard issued by the U.S. National Institute of Standard and Technology. The calibration methods takes into account that the bandwidth of GUVis-3511 channels is about 10 nm FWHM [\(18.5 nm for the channels at 305 nm\)](#). The calibration was further optimized for the conditions (solar zenith angle, TOC, AOD, etc.) at the measurement site. [Details of this optimization are provided in the Supplement.](#) Results indicated that it is advantageous to calibrate the channels with nominal wavelengths of 305 and 313 nm for spectral irradiance at 306 and 315 nm, respectively, and measurements reported in the following are therefore referenced to these two wavelengths. [Both direct and global irradiance measurements were corrected for the cosine error of the instrument’s collector as described by Morrow et al. \(2010\), based on the measured cosine error \(Fig. 4\) and the ratio of direct and global irradiance extracted from the shadowband measurements.](#) Finally, an uncertainty budget was established. Expanded ($k = 2$, equivalent to a confidence interval of 95 %) uncertainties for the 305 and 313 nm channels are 7.5 and 7.3

Deleted: full width at half maximum (

Deleted:)

%, respectively. Expanded uncertainties of all other channels, with the exception of the 940 nm channel, are 2.7 %. An uncertainty budget for the 940 nm channel is not provided here because of the large effect of water vapor on measurements at this wavelength. Measurements of this channel should therefore be interpreted with caution.

5 Secondary data products

5 Secondary data products derived from the measurements of the GUVIS-3511 include AOD and TOC.

5.1 Aerosol optical depth

Aerosol optical depth τ_a was calculated from calibrated direct measurements and Beer-Lambert's law:

$$\tau_a = \frac{-1}{\mu_a} \left[\ln \left(\frac{\rho \hat{E}_{S_d}(\lambda_i)}{\hat{E}_0(\lambda_i)} \right) + \tau_r \mu_r + \tau_o \mu_o \right], \quad (1)$$

where

$\hat{E}_{S_d}(\lambda_i)$ is the response-weighted direct irradiance measured by the instrument (see Supplement for definition),

$\hat{E}_0(\lambda_i)$ is the extraterrestrial solar spectrum as defined in Sect. S1.1.1 of the Supplement at 1 astronomical unit (AU), weighted with the response functions of the GUVIS-3511,

ρ scales the extraterrestrial solar spectrum from 1 AU to the Sun-Earth distance applicable to 21 August 2018,

τ_r is the Rayleigh optical depth calculated by Eq. (30) of Bodhaine et al (1999),

μ_r is the Rayleigh airmass factor, calculated according to Kasten and Young (1989),

τ_o is the ozone optical depth derived from the ozone absorption cross section by Bass. and Paur (1985) and assuming a TOC of 298 DU,

μ_o is the ozone airmass factor calculated by Eq. (11) of Bernhard et al. (2005b) assuming an ozone layer height of 22 km, and

μ_a is the aerosol airmass factor calculated by Eq. (11) of Bernhard et al. (2005b) assuming an aerosol layer 2 km above the surface.

Aerosol optical depth τ_a is reported at the centroid wavelength (Seckmeyer et al., 2010) of each channel. [Aerosol optical depth was not calculated for the two 305 nm channels because of the large bandwidth of these channels and the strong interference with ozone absorption at this wavelength. Both factors lead to large uncertainties.](#)

5.2 Total ozone column

The TOC during the eclipse was derived from measurements of response-weighted global irradiance $\hat{E}_{S_g}(\lambda_i)$ measured by the GUVIS-3511 (see Supplement for definition of $\hat{E}_{S_g}(\lambda_i)$). Specifically, ratios of $\hat{E}_{S_g}(\lambda_L) / \hat{E}_{S_g}(\lambda_S)$ were compared with similar ratios in a look-up table that was calculated with a RT model (also described in the Supplement) as a function of SZA and TOC. The following three wavelength pairs were used: (1) $\lambda_L = 340$ nm, $\lambda_S = 305$ nm; (2) $\lambda_L = 340$ nm, $\lambda_S = 313$ nm; (3) $\lambda_L = 313$ nm, $\lambda_S = 305$ nm. This method was first proposed by Stamnes et al. (1991) and was further validated by Bernhard et al. (2003; 2005b). The application of the method to GUV instruments was described by Bernhard et al. (2005a). By analyzing the sensitivity of TOC to changes in the irradiance ratio and by quantifying the noise in the measurements of the 305 and 340 nm channel (wavelength pair (1)), we determined that our measurements are able to detect changes in ozone of 0.14 DU (0.05 %) at the start and 0.10 DU (0.03 %) at the end of the eclipse at 95 % confidence level for constant aerosol conditions. The additional uncertainty in TOC values due to changing AODs was determined to be 1.5 DU (0.5 %). While TOCs could also be derived from direct irradiances, these measurements are not discussed here due to the relatively large uncertainty to calculate direct irradiance from shadowband data at wavelengths in the UV-B (280–315 nm), in particular at large SZAs.

Deleted: and was validated for

6 Celestial calculations

The interpretation of measurements performed during the solar eclipse requires very accurate calculations of the relative distance between Sun and Moon and angular diameter of both celestial bodies. Such calculations are provided by the HORIZONS on-line solar system data and ephemeris computation service of the Jet Propulsion Laboratory (<https://ssd.jpl.nasa.gov/?horizons>). From the many parameters made available through this system in time intervals of one minute, we used the elevation, azimuth, and angular radius of Sun (θ_S , φ_S , δ_S) and Moon (θ_M , φ_M , δ_M), calculated the angular distance ξ between Sun and Moon via

$$\xi = \arccos[\sin(\theta_S) \sin(\theta_M) + \cos(\theta_S) \cos(\theta_M) \cos(\varphi_S - \varphi_M)], \quad (2)$$

and interpolated these measurements to the times of our measurements (Fig. 5). The eclipse begins and ends when $\xi = \delta_S + \delta_M$, and the maximum eclipse is observed when ξ is at its minimum. The start (1st contact), “maximum eclipse”, and end (4th contact) times calculated with this method were compared with solar eclipse data calculated with an online calculator provided by the Astronomical Applications Department of the U.S. Naval Observatory (USNO) at <http://aa.usno.navy.mil/data/docs/Eclipse2017.php>. Results of our calibration and those by the USNO are compared in Table

1 and agree to within ± 3 seconds, giving confidence in our calculations. Note that the period between 1st contact and maximum is about 6.5 minutes shorter than the period between maximum and 4th contact.




Fig. 5. Panel (a): Apparent angular radii of Sun and Moon on 21 August 2018. The apparent radius of the Sun is virtually constant over time while the radius of the Moon varies considerably during the day due to the rotation of the Earth, which changes the distance between two hypothetical observers standing on the Earth and the Moon. Panel (b): Corresponding solar and lunar elevation and azimuth angles. Note the conversion of the curves at the time of maximum eclipse (short-dashed vertical line). Long-dashed lines indicate the times of the 1st and 4th contact.

10

Table 1. Comparison of Eclipse calculations by us and USNO.

	Time (UT)		Difference (seconds)	Solar zenith (°)	Solar Azimuth (°)
	Our calculation	USNO			
Start partial eclipse (1 st contact)	16:06:24	16:06:27	−3	60.6	102.7
Start total eclipse (2 nd contact)		17:19:42		48.4	118.9
Maximum eclipse	17:20:27	17:20:24	3	48.3	119.1
End total eclipse (3 rd contact)		17:21:07		48.2	119.3
End partial eclipse (4 th contact)	18:41:05	18:41:02	3	37.4	143.6

The fraction of the solar disk seen by the GUVis-3511 and the change of the extraterrestrial spectrum as function of time during the progression of the eclipse were calculated with the algorithm by Koepke et al. (2001). Koepke et al. (2001) uses a simple analytical formula to describe the wavelength dependence of solar LD, which only considers the temperature at the Sun’s surface (Waldmeier, 1941). We also use two other parameterizations of the LD functions that are based on data

15

collected by the McMath-Pierce Solar Telescope of the National Solar Observatory on Kitt Peak. Specifically, the function $\Gamma_\lambda(r)$ defined in Eq. (2.1) of Koepke et al. (2001) was replaced with the polynomial

$$P(\lambda, r) = \sum_{k=0}^5 a_k(\lambda) \cos^k(\psi(r)), \quad (3)$$

where the angle of incidence ψ is defined in Fig. 6 and the coefficients $a_k(\lambda)$ were derived from measurements at Kitt Peak. The quantity $\Gamma_\lambda(r)$ used by Koepke et al. (2001) depends on the distance r between a point on the solar disk and the

5 center of Sun, relative to the radius of the Sun, R_S (Fig. 6). The angle ψ was calculated from r using basic trigonometry.




Fig. 6. Limb darkening geometry. The Sun is centered at point O and has the radius R . The observer is at point P at a distance R_{SE} from the center of the Sun, and is looking at point S on the surface of the Sun. From the point of view of the observer, S is at an angle δ_r relative to the center of the Sun, and the Sun's limb is at angle δ_S . The angular radius of the solar disk as viewed by the observer is R_S and the angular radius associated with S is r . ψ is the angle between the Sun's normal at S, and the line between S and P.

Two sets of coefficients $a_k(\lambda)$ were used in this study. The first set was developed by Pierce and Slaughter (1977) ('PS') and Pierce et al. (1977) ('PSW') using measurements collected between 1974 and 1976 near a minimum of solar activity. These coefficients are tabulated in Table IV of PS (wavelength range 303.327–729.675 nm) and Table IV of PSW (740.46–1046.7 nm). The second set was published by Neckel (2005), using measurements performed in 1986 and 1987 near the following solar minimum and these coefficients are listed in Table I of Neckel (2005) (300–1100 nm). Both sets of data were

collected with the observatory's large vertical spectrograph, which consists of a double-monochromator with prism pre-disperser. Measurements across the solar disk were taken by stopping the telescope drive to let the Sun pass across the entrance port of the radiometer.

We note that different parameterizations are used by Neckel (2005) for the wavelength ranges 300.00–372.98 nm, 385.00–422.57 nm, and 422.57–1100 nm, but no parameterization is provided for the range of 372.98–385.00 nm because this range is affected by the Balmer “jump” in the absorption of the hydrogen atom resulting in a large uncertainty in the solar LD function in this wavelength range. We compare our measurements at 380 nm with the parameterization developed for the 385.00–422.57 nm range.

Fig. 7 compares results of the three parameterizations, referred to in the following as the parameterizations by Waldmeier, Pierce, and Neckel. Fig. 7a is based on the parameterization by Pierce and shows ratios of the extraterrestrial solar spectral irradiance during the eclipse to the spectral irradiance that would be expected if the Sun were a uniformly bright star. Results are plotted versus the fraction of the Sun's area that is visible (0 = Sun completely occluded by Moon, 1 = Sun completely visible). The LD effect is strongly dependent on wavelength: when the Sun is almost completely eclipsed, the spectral irradiance at 306 nm is reduced to about 22 % of the intensity that would be expected from purely geometrical considerations, whereas the spectral irradiance at 1020 nm is only reduced to about 67 %. Figure 9a also indicates that the LD effect is slightly positive when 66 % or more of the Sun's area are visible. In these cases, the Moon occludes the darker, outer parts of the Sun, and the average intensity of the visible area is therefore larger than that of a hypothetical, uniformly bright star.

Fig. 7b shows similar results but is based on the parameterization by Waldmeier. For wavelengths larger than 500 nm, the LD effect is similar to that calculated with the parameterization by Pierce. For smaller wavelengths, results derived with the parameterization by Waldmeier indicate a much weaker LD effect.

Fig. 7c shows the ratio of the LD effect calculated with the parameterization by Neckel and Pierce. Results for the two parameterizations agree to within ± 2.3 % when 5 % or more of the Sun is visible. The largest difference of 7.6 % is observed at 395 nm when only 0.6 % of the Sun are still visible. On the other hand, results between the Waldmeier and Pierce parameterizations differ by as much as a factor of 1.78 (Fig. 7d).




Fig. 7. Comparison of the solar LD effect calculated with the parameterizations by Pierce, Neckel and Waldmeier. Top row: Ratios of the extraterrestrial solar spectral irradiance during the eclipse to the spectral irradiance from a hypothetical Sun without LD, plotted versus the fraction of the Sun that is visible. Panels (a) and (b) are based on the parameterizations by Pierce and Waldmeier, respectively. Bottom row: Ratios of LD effect calculated by the parameterizations by Neckel and Pierce (Panel c) and by Waldmeier and Pierce (Panel d). The vertical broken line indicates the minimum fraction of 0.022 (2.2 % of the Sun's disk) that could be assessed with our measurements.

7 Results

In this section, we [validate the solar LD parameterizations by the three methods discussed in Sect. 6, present the ratio of direct-to-global irradiance](#), show the AOD and TOC derived from our measurements before, during and after the eclipse, and compare measurements near and during totality with reconstructed measurements for the same period but without the Moon occulting the Sun.

7.1 Effect of solar limb darkening

Theoretical predictions of the solar LD effect described in Sect. 6, ~~were compared with the LD effect derived from our~~ measurements (Fig. 8). To account for the change in SZA, data were first compared with results of the RT model described

Deleted: present the ratio of direct-to-global irradiance, validate the solar LD parameterizations by the three methods discussed in Sect. 6,

Deleted: 5

in the Supplement. Model input parameters (e.g., AOD) are based on the measurement at 19:00:30 (shortly after the end of the eclipse) as described in Sect. 7.3. The only model parameter changing as a function of time is the SZA. Fig. 8a shows a comparison of measurements of global spectral irradiance (symbols) and these model calculations (thin lines). The agreement between measurement and model for times outside the period of the eclipse is reasonable, but not perfect. The ratio of measurement and model is shown in Fig. 8b. At 19:00:30 (i.e., the time relevant for the input parameters of the RT calculations) the ratio varies between 0.958 and 1.027, except for the channels at 313 and 320 nm (where the ratio is 0.932), and at 940 nm (ratio of 1.139) due to the uncertainty of the water vapor column affecting this wavelength. The discrepancy between measurement and model at the start of the eclipse is generally larger, likely due to the change in AOD, which was not considered by the model. To correct for the difference between measurement and model, linear functions were constructed to match the ratio of measurement and model at the times of the 1st and 4th contact (thin lines in Fig. 8b). This correction assumes that changes in the atmospheric constituents (aerosols, ozone, etc.) led to a linear change in the atmospheric transmission over the period of the eclipse. Next, ratios of measurement and model were normalized by dividing with these linear functions. The resulting bias-corrected ratios represent the LD effect derived from our measurements (symbols in Fig. 8c). Thin lines in that figure show the theoretical LD effect calculated with the parameterization by Pierce. The insert of Fig. 8c shows both datasets for the time between 17:01:00 and 17:04:30, indicating that there is almost perfect agreement for wavelengths larger than 400 nm, but measurements at UV wavelengths are below the theoretical prediction.

Lastly, the measured LD effect was divided by the theoretical data and the resulting ratios are shown in Fig. 8d. For wavelengths larger than 400 nm, measurement and theory agree better than ± 4.0 %. When excluding the period between 16:44 and 17:00, which was affected the most by aerosols during the duration of the eclipse, the agreement is to within ± 2.5 %. For times outside of ± 7 minutes from totality, measurement and theory for 380 and 395 nm also agree within this range. The larger discrepancy closer to totality is likely caused by the uncertainty of the theoretical prediction due to the Balmer “jump” affecting these wavelengths. The discrepancy between measurement and theory for wavelengths below 380 nm (306–340 nm) becomes larger than 3 % as totality is approached, with the smallest wavelengths showing the largest change (e.g., the ratio is 0.90 for 306 nm at 17:01). For these wavelengths, the discrepancy is generally larger before totality than after totality. We suspect that this asymmetry is caused by changing aerosols and note that the measured LD effect is lower than the theoretical prediction both for wavelengths affected by ozone absorption (306, 315, and 320 nm) and wavelengths not affected by ozone absorption (340 nm). Hence, the larger discrepancy at these UV wavelengths (and the asymmetry relative to totality) cannot solely be caused by ozone absorption in the atmosphere. Results for wavelengths that were measured independently by two channels (306, 340, and 555 nm) are virtually indistinguishable, suggesting that calibrations were applied consistently and that both types of photodiodes used for these channels led to practically identical results.

Deleted: likely

Deleted: do not know

Deleted: the reason for

Deleted: but




Fig. 8. Comparison of the solar LD effect derived from measurements of global spectral with theoretical predictions based on the parameterizations by Pierce. Panel (a): measurements of global spectral irradiance (symbols) and results of the RT calculations described in Sect. S1.1.1 of the Supplement (lines). Only every 20th data point is plotted for clarity. Panel (b): ratio of measurement and model (symbols). Lines connect the ratios at the times of the 1st and 4th contact, which are indicated by the vertical long-dashed lines. Panel (c): Ratio of measurement and model, corrected for the bias between measurement and model (symbols). The insert shows this ratio for times between 17:01:00 and 17:04:30. The theoretical LD effect calculated with the parameterizations by Pierce is also shown and indicated with lines. Panel (d): ratio of biased-corrected measurements (symbols in panel (c)) and theoretical prediction (lines in panel (c)). Ratios for the alternative channels at 306, 340, and 555 nm are indicated with broken lines and are virtually indistinguishable from those using the standard photodiode, which are indicated by solid lines. The time of maximum eclipse is indicated by the short-dashed black line. The dashed gray line specifies the time of 19:00:30 for which the AOD used by the model calculations was derived. The legend on top of the figure indicates spectral irradiance in nm.

Fig. 9 compares the measured LD effect with the theoretical prediction based on the parameterization by Neckel instead of Pierce. Results shown in Fig. 8d and Fig. 9 are very similar and differences are within the uncertainties of the measurements and the change of the atmospheric extinction due to the variability of the aerosol load. Despite the similarity of the two figures, there is one difference worth noting. For 380 and 395 nm, discrepancies calculated for the parameterization by Neckel are larger than those derived from the parameterizations by Pierce. This is not surprising considering that Neckel does not provide coefficients for the wavelength range of 372.98–385.00 nm because of the uncertainty from the Balmer jump. As described in Sect. 5, our parameterization for the LD effect at 380 nm is based on Neckel’s coefficients for the 385.00–422.57 nm range and therefore subject to uncertainties.




Fig. 9. Similar to Fig. 8d, but theoretical LD effect parameterized based on Neckel instead of Pierce.

Deleted: (2005)

Fig. 10 shows the ratio of the LD effect calculated with the parameterizations by Neckel and Pierce. The figure is based on the same data as those used for Fig. 7c, but are plotted versus time instead of the fraction of the Sun not obstructed by the Moon. Results of the parameterization are virtually identical if 60 % or more of the Sun is visible. Discrepancies increase as the Sun becomes more and more occluded and are largest at 380 and 395 nm, with a maximum difference of about 3 %. There is generally no systematic dependence on wavelength.




Fig. 10. Ratio of the LD effect calculated with the parameterizations by Neckel and Pierce. The color coding is identical with that used in Fig. 8. The dashed line indicates the fraction of the Sun's disk that is visible and is referenced to the right axis. Vertical dashed lines indicate the 1st and 4th contact.

5

Lastly, when we compared our observations with theoretical prediction based on the parameterization by Waldmeier, observation were lower than theory by up to 37 % at 306 nm, 24 % at 340nm, and 19 % at 412 nm. For longer wavelengths, differences were in the ± 8 % range. These differences are well outside the uncertainty of our measurements, confirming that the parameterization by Waldmeier is too simple, in particular in the UV range.

10 7.2 Direct-to-global ratio

Fig. 11a shows the ratio of direct spectral irradiance $\bar{E}_{S_d}(\lambda_i)$ (derived from shadowband data) and global spectral irradiance $\bar{E}_{S_g}(\lambda_i)$ (see Supplement for definition of symbols). The general upward slope is due to the change in SZA between morning and noon, and the steep slope between 15:15 and 15:45 is due to a decrease in aerosols as discussed in Sect. 7.3. Thin lines in Fig. 11a connect measurements at the start of end of the eclipse and are drawn to guide the eye. Note that direct measurements are not available between 200 s before the 2nd contact and 105 seconds after the 3rd contact.

Deleted: 7

Deleted: indicated by Fig. 12

15

The measured direct-to-global ratio was compared with the modeled direct-to-global ratio using the same method as that employed in Sect. 7.1. In brief, the measured direct-to-global ratio was divided by the modeled direct-to-global ratio. To correct for the difference between measurement and model, linear functions were again constructed to match the ratios of measurement and model at the times of the 1st and 4th contact. Finally, the ratios were normalized by dividing with these linear functions. The resulting bias-corrected ratios of the measured and modeled direct-to-global ratio are shown in Fig. 11b and are denoted $R_{DG}(\lambda_i)$.

20




Fig. 11. Panel (a): ratio of direct-to-global spectral irradiance. Measurements are shown as heavy lines. Thin lines connect measurements at the start/end of the eclipse and are drawn to guide the eye. Panel (b): Ratio of the measured and modeled direct-to-global ratios, corrected for the bias between measurement and model. Numbers “(1)” and “(2)” in the legend indicate the channel number for channels equipped with identical wavelengths

Deleted: R

Before 15:00, the measured direct-to-global ratio is smaller than 0.03 at wavelengths in the UV-B and values of $R_{DG}(\lambda_i)$ are subject to large uncertainties at these wavelengths. Before 15:20, $R_{DG}(\lambda_i)$ varies between 0.55 and 0.87 for wavelengths in the visible range, indicating that the measured direct-to-global ratio is 13 to 45 % below the modeled one. These low values can be attributed to increased aerosol loading prior to the start of the eclipse. After the start of the eclipse

(1st contact), values of $R_{DG}(\lambda_i)$ start to drop to reach a local minimum at 16:52, approximately 30 minutes before the eclipse maximum. The decline has a clear wavelength dependence with wavelengths in the UV-B decreasing the most (up to 20 %), followed by wavelengths in the visible (up to 15%) and IR (up to 3.5%). After this local minimum, $R_{DG}(\lambda_i)$ slowly increases during the remainder of the eclipse.

The change of $R_{DG}(\lambda_i)$ over the period of the eclipse could either be caused by processes initiated by the occlusion of the Sun or by variability from aerosols. Aerosols effects are more likely because any eclipse effect should conspicuously peak at the time of totality, not 30 minutes earlier. The wavelength dependence of $R_{DG}(\lambda_i)$ is also characteristic for aerosol effects (Sect. 7.3). In addition, the minimum values of $R_{DG}(\lambda_i)$ during the eclipse are well within the range of values observed prior to the eclipse, which could be unambiguously attributed to aerosols.

7.3 Aerosol optical depth

AOD was calculated with Eq. (1) from direct spectral irradiance, which in turn was derived from data collected during shadowbanding. The standard algorithm computing AODs does not take into account that a solar eclipse is occurring and (incorrectly) attributes the reduced direct irradiance during the eclipse to increased extinction of radiation by aerosols. The result (thin lines in Fig. 12a) is the large spike in AOD, peaking at the time of totality. Calculations were repeated by first scaling the direct spectral irradiance with the LD effect using the parameterization by Pierce (Sect. 6) and then feeding the scaled irradiances in the AOD algorithm. Results corrected for the LD effect (thick lines in Fig. 12a) indicate that the AOD was increasing between the 1st contact and 16:52 (from 0.41 to 0.58 at 319 nm and from 0.05 to 0.07 at 1018 nm) and then monotonically decreasing to 0.34 at 319 nm and 0.04 at 1018 nm at the end of the eclipse. Corrected results do not have a spurious spike near the time of totality.

AOD measurements before and after the eclipse were used to characterize the aerosol loading of the atmosphere. Before 15:45, AODs were variable due to aerosols from nearby wildfires. Fig. 12b shows AOD as a function of wavelength shortly before (16:02:58) and after the eclipse (19:00:30). AODs were somewhat lower after the eclipse compared to the start. An Ångström function of the type $\tau_a = \beta \lambda^{-\alpha}$, where the wavelength λ is provided in μm was fitted to the data using channels between 340 and 1020 nm, excluding the channel at 940 nm. Values of the Ångström coefficient were $\alpha = 1.96$ and $\beta = 0.057$ for the measurement at 16:02:58, and $\alpha = 2.1$ and $\beta = 0.0394$ for the measurement at 19:00:30. The second set of coefficients was used for the RT calculations discussed in Sect. S1.1.1 of the Supplement. Ångström exponents α in the order of 2.0 may seem large, however, O’Neil et al. (2002) have demonstrated that values of $\alpha = 2$ are typical for aerosols

Deleted: Measured ratios tend to be below these lines by ratio values of up to 0.04, and these low ratio values

Deleted: (A change by 0.04 is well within the range of aerosol effects occurring before 15:30).

Deleted: s

Deleted: as it was the case for the spurious peaks in AOD (Fig. 12) and TOC (Fig. 13) measurements

Deleted: Note that direct measurements are not available between 200 s before the 2nd contact and 105 seconds after the 3rd contact.

monotonically

Formatted: Superscript

over the period of the eclipse

(from 0.41 to

2

from 0.05 to

) without

somewhat

originating from forest fires. Deviations of the measured AODs from the Ångström parameterizations shown in Fig. 12b are smaller than 0.02, with the exception of the AODs at 314.3, 319.4, and 442.4 nm. Measured AODs are up to 0.15 smaller than the parameterization at these wavelengths. The reason for these discrepancies is unknown, however, differences between AODs at the three wavelengths relative to AODs at adjacent wavelengths are larger at 19:00:30 than 16:02:58. As measurements at the two time were processed in the same way it seems unlikely that the discrepancies are caused by a measurement artifact.

Deleted: , suggesting that the cause is an unknown absorber and not due to a measurement artifact




Fig. 12. Aerosol optical depth. Panel (a): AOD measured by the GUVIS-3511 radiometer at Smith Rock as a function of time. Thin lines indicate AOD measured without LD correction while thick lines indicate AOD corrected using the LD parameterization by Pierce. Long-dashed lines indicate the start and end times of the eclipse and the short-dashed line indicates the time at totality. Without LD correction, the reduced irradiance during the eclipse period is incorrectly attributed to aerosols, leading to a spurious spike at the time of totality. With LD correction, the effect of the eclipse is not noticeable in the AOD data. The legend indicates the centroid wavelengths of the GUV's channels. Panel (b): Aerosol optical depth as a function of wavelength shortly before (16:02:58) and after the eclipse (19:00:30). Dashed lines indicate Ångström function fits to the data based on measurements of the AODs between 339.8 and 1017.6 nm.

Deleted: barely

7.4 Total ozone column

Total ozone column was calculated from GUVIS-3511 measurements as described in Sect. 5.2. [Results obtained from the 305 nm / 340 nm wavelength pair are discussed below. Similar results calculated with the 305 nm / 313 nm and 313 nm / 340 nm wavelength pairs are presented in Sect. S2 of the Supplement.](#) Data were first processed without taking into account that the solar LD effects reduces the extraterrestrial spectrum more strongly in the UV-B than at 340 nm. Processing was repeated by scaling the raw data with the three parameterizations of the LD effect described in Sect. 6. Fig. 13 compares these results with measurements by OMI. (Of note, OMI data for 21 August 2018 are not available for the pixel that contains our measurement site. However, the average of OMI measurement within $\pm 1^\circ$ in latitude and $\pm 5^\circ$ in longitude is 297.9 DU with a standard deviation of 2.1 DU.) The TOC measured by the GUVIS-3511 was 293 DU shortly before the start of the eclipse and 294 DU shortly after the end, suggesting that the eclipse had no lasting effect on atmospheric ozone concentrations. Measured TOCs without LD correction spiked during the eclipse, resulting in 316 DU shortly before totality. When correcting the data using the LD parameterization by Waldmeier, the height of the peak is reduced to 312 DU (a 1.3 % reduction). When using either the parameterizations by Neckel or Pierce, the effect of the eclipse on measured TOCs almost disappears, in particular for the period between the 3rd and 4th contact. [The increase in TOC of about 5 DU between 16:00 and 17:00 can partly be explained by increasing AODs over this period, which cause a high-bias in the ozone retrievals.](#) The results shown in Fig. 13 can be explained with the wavelength dependence of LD. The stronger reduction of the extraterrestrial irradiance at 305 nm compared to 340 nm caused by the LD effect is interpreted by the algorithm as a larger TOC. Once the LD effect is removed, the TOC remains constant to within ± 2.6 DU or ± 0.9 %. [Results obtained with the 305 nm / 313 nm and 313 nm / 340 nm wavelength pairs \(Sect. S2 of the Supplement\) corroborate these findings.](#)

Deleted: of the 305 and 340 nm channels

Deleted: of the two channels

Deleted: nm

Deleted: (We have no explanation for the increase in TOC of about 5 DU between 16:00 and 17:00 other than natural variability).

Deleted: 6





Fig. 13. Total ozone column measured by GUVIS-3511 and OMI. For “uncorrected” data, the spike in TOC during the period of the eclipse is an artifact that can be attributed to solar LD. The effect is only marginally reduced when using the LD correction based on the parameterization by Waldmeier. By using either the parameterizations by Pierce or Neckel, variability in TOC during the eclipse is reduced to ± 2.6 DU (± 0.9 %).

7.5 Measurements near and during totality

Fig. 14 shows measurements between about 1.5 minutes before the 2nd contact and about 2 minutes after the 3rd contact. The measurement protocol was not changed during this period and the radiometer was therefore shadowbanding during a large part of totality (between 17:20:05 and 17:21:11). Periods when shadowbanding was active are indicated by grey shading in

- 5 Fig. 14. At 17:18:53 (before 2nd contact) and 17:22:53 (after 3rd contact) the shadowband blocked the direct Sun, leading to a strong decreases of the measured spectral irradiance. In contrast, the effect of the shadowband is barely noticeable during totality, although the small dip in the spectral irradiance between 17:20:10 and 17:20:40 can be attributed to the blocking of sky light by the band. Shadowband data are considerably noisier than global irradiance measurements because of the difference in integration times (1 second for global irradiance and 1/15 second for shadowband measurements). To
- 10 compensate for this effect, shadowband data shown in Fig. 14 are a running average of 15 measurements. In addition, shadowband measurements were scaled with a cosine error correction factor calculated for isotropic incident radiation, but were not corrected for the shading of skylight by the band.



- 15 Fig. 14. Spectral irradiance measured between about 1.5 minutes before the 2nd contact and about 2 minutes after the 3rd contact. The color coding is identical with that used in Fig. 8. Grey shading indicates times of shadowbanding. The times of the 2nd and 3rd contact are indicated by dot-dashed lines and the short-dashed line marks the time of maximum totality.

- Measurements of global spectral irradiance are available between 17:19:15, through the 2nd contact, and until 17:20:20. These measurements were cosine-error corrected without taking into account the change in the direct-to-global ratio at the beginning of totality. Despite this approximation, there is no obvious discontinuity at 17:20:20 when global irradiance measurements transition into shadowband data. During the last half-minute before the 2nd contact, spectral irradiances in the UV and visible decrease by about one order for magnitude. The decrease in the IR is considerably larger with a change of about 2 and 2.5 orders of magnitude at 1020 nm and 940 nm, respectively. The wavelength dependence of these changes can be explained with the wavelength-dependence of Rayleigh scattering, which makes it less likely for photons entering the
- 20

atmosphere outside the Moon’s shadow to travel to the area of the umbra. The long pathlength also amplifies absorption by gases (ozone, NO₂, H₂O) and aerosols. Spectral irradiance at 940 nm, which is affected by water vapor absorption, decreases therefore more rapidly than spectral irradiance at 1020 nm, which is absorbed much less. Measurements at 306 nm are below the detection limit during totality.

The difference of the global spectral irradiance observed during totality relative to the unoccluded Sun was quantified with the data shown in Fig. 8c. Specifically, the bias-corrected ratio of measurement and model averaged over the period of 17:19:50 and 17:20:00 (i.e., the last 10 seconds of global spectral irradiance measurements during totality before the start of the shadowbanding sequence) were compared to the respective ratio at the 4th contact, which is one by design. Fig. 15 shows the so-calculated reduction factors. Factors decrease strongly between 315 and 340 nm, followed by a gradual increase from 340 nm to 1020 nm. Spectral irradiance is reduced by a factors of 18,700 at 315 nm, 7,000 at 340 nm, 18,340 at 665 nm, 136,650 at 940 nm and 46,150 at 1020 nm. The relatively high factors in the UV-B and at 940 nm are caused by ozone and water vapor absorption, respectively. The average reduction for the channels in the visible (412–665 nm) is 11,460. For comparison, the irradiance from the full Moon is about a factor of 400,000 smaller than that of the Sun (Macdonald, 2012). Hence, the visible irradiance during totality is still a factor of 35 larger than the irradiance from the full Moon. This estimate is in agreement with empirical observations. For example, the light during totality is still bright enough to perceive colors while this is barely possible with moonlight.




Fig. 15. Factor quantify the reduction of spectral irradiance during totality relative to spectral irradiance from the unoccluded Sun. Measurements of the two channels at 306 nm are below the detection limit during totality and the factors calculated for two channels (connected by a dashed line) are therefore greatly different and meaningless. Factors for the other wavelengths with two channels (340 and 555 nm) are virtually identical. The reduction factor for 940 nm is 136,650.

8 Discussion

Some of the findings of this paper are in conflict with results from similar publications discussing solar eclipses that have occurred in the past. These pertain to ~~the~~ magnitude of ~~the~~ LD correction, ~~changes of the diffuse-to-direct ratio observed during an eclipse,~~ and ~~TOC variations during an eclipse.~~ The three issues are discussed below, followed by a discussion of the value of eclipse data for validating the performance of radiometers.

Deleted: changes of the diffuse-to-direct ratio observed during an eclipse,

Deleted: the

8.1 Magnitude of solar limb darkening

The excellent agreement (better than 4.0 % overall and better than 2.5 % excluding the period ~~most~~ affected by aerosols) between the solar LD effect derived from our measurements and theoretical predictions for wavelengths above 400 nm suggests that the parameterizations of LD by Pierce and Neckel are correct to within the measurement uncertainty for this wavelength range. The larger discrepancies below 400 nm ~~are likely caused by aerosol effects and~~ systematic errors in these parameterization. Specifically, Neckel uses different coefficients for the wavelengths ranges 300–372.98 nm, 385.00–422.57 nm and 422.57–1100 nm because of the Balmer jump. The electronic transitions from the 2nd atomic shell to shells higher than 6 occur at wavelengths between 410.2 (transition 2 > 6) and 364.6 nm (Balmer break). The hypothesis of possible systematic errors in the LD data by Neckel is supported by the relatively large noise in data at UV wavelengths presented by Neckel (2005), which formed the basis of LD parameterizations used here.

Deleted: could be

Deleted: due to

Deleted: On the other hand, the ratio of the measured and theoretical LD effect (Fig. 8) shows some asymmetry for wavelengths below 380 nm and these could be caused by measurement artifacts of unknown origin. ¶

Deleted: also

Our results ~~indicate that the~~ LD parameterization by Waldmeier is ~~too simple to accurately describe the change of the~~ radiance across the solar disk, in particular in the UV range. Several recent papers (e.g., Blumthaler et al., 2006; Emde and Mayer, 2007; Kazadzis et al., 2007; ~~Kazantzidis et al., 2007~~) have used the method by Koepke et al. (2001) combined with the LD darkening parameterization by Waldmeier. Some conclusions in those papers should therefore be reconsidered. For example, Kazadzis et al. (2007) calculated the TOC during the total eclipse of 29 March 2006 at Kastelorizo, Greece, from direct irradiance measurements of a Brewer spectrophotometer. By combining the method by Koepke et al. (2001) with the parameterization by Waldmeier (1941), they calculated negligible LD corrections of less than 0.01 % for their ozone retrievals. (Of note, the maximum correction using the parameterization by Waldmeier for our data (grey line in Fig. 13) is 1.3 %, which is much larger than the correction of 0.01 % quoted by Kazadzis et al. (2007). The reason for this discrepancy is unknown, however, the wavelengths used in our TOC retrieval (306 and 340 nm) are different from the Brewer wavelengths, which range between 306.2 and 320.1 nm.)

Deleted:

~~Kazantzidis et al. (2007) have analyzed ratios of global spectral irradiance (305 nm / 380 nm, 312 nm / 380 nm, 340 nm / 380 nm, and PAR / 380 nm) that were measured with NILU-UV filter radiometers at three locations in Greece during the total solar eclipse of 29 March 2006. These measured ratios were compared with theoretical predictions based on the algorithm by Koepke et al. (2001) and the LD parameterization by Waldmeier. As the eclipse progressed, the model~~

underestimated the measured spectral effect, capturing only half of the observed change. For example, measured ratios of spectral irradiances at 340 and 380 nm were 10 % lower close to totality compared to similar ratios calculated for the 1st and 4th contact. The theoretical calculation only predicted a decrease of 5 %. Our results suggests that discrepancies between the measured and modeled ratios reported by Kazantzidis et al. (2007) can partly be attributed to limitations of the LD parameterization by Waldmeier used in their model.

8.2 Variations of direct-to-global ratio

Relative to model calculations for the unoccluded Sun, direct-to-global ratios during the eclipse decreased by up to 20 % in the UV-B, 15 % in the visible and 3.5 % in the IR (Fig. 11b). The largest decrease was observed about 30 minutes before totality. The timing of this decrease plus its spectral dependence suggest that changes in aerosol amounts are the main driver for the observed drop in the ratio. However, aerosol effects are difficult to decouple from processes initiated by the occlusion of the Sun or artifacts of the algorithm used to calculate the direct irradiance from shadowband data.

Deleted: The direct-to-global ratios shown in Fig. 11 increase with time as expected from the decrease in SZA. Deviations from a straight line between the 1st and 4th contact are less than 0.04. It is difficult to determine whether this deviation is caused by processes initiated by the occlusion of the Sun, by variability from aerosols, or artifacts of the algorithm to calculate the direct irradiance from shadowband data. We therefore consider the deviation of 0.04 as an upper limit for the eclipse effect.

According to theoretical calculations by Emde and Mayer (2007), eclipse effects that affect direct and diffuse radiation differently are smaller than 1 % for times 10 minutes away from totality and smaller than 4 % for times 105 seconds away from totality (i.e., the shortest time to totality available from our shadowband measurements). These theoretical calculations also agree with the conclusion by Sharp et al., (1971) that “sky light may be considered as attenuated sunlight up to at least 99.8 % obscuration.” These theoretical results disagree with the results by Zerefos et al. (2000; 2001), who suggest that the erythmal (sunburning) diffuse irradiance was declining at a slower rate than the erythmal direct irradiance during the solar eclipse observed in Thessaloniki, Greece, on 11 August 1999. The largest difference was observed at the time of the eclipse maximum when 88 % of the solar disk was obscured and the ratio of diffuse to direct irradiance was increased by approximately 50 % relative to the change expected from calculations with a one-dimensional radiative transfer model. For comparison, the increase in the diffuse-to-direct ratio calculated at 315 nm from our measurements was only 31 % and did not peak at the time of totality. The available evidence from our observations and 3DRT model calculations suggests that measurement artifact may have contributed to the large change of the diffuse-to-direct ratio reported by Zerefos et al. (2000; 2001).

Deleted: Our results as well as t

Deleted: However, t

Deleted: in the UV is

Deleted: direct

Deleted: . When 90 % of the Sun are occluded (i.e., the maximum coverage for the eclipse discussed by Zerefos et al. (2000)),

Deleted: is

Deleted: reduced 30 % less than direct irradiance.

Deleted: was a measurement artifact.

8.3 Total ozone column variations during an eclipse

In contrast to the increase that is apparent in our uncorrected TOC measurements shown in Fig. 13, the ozone retrievals by Kazadzis et al. (2007) show a decrease in TOC during the eclipse. Kazadzis et al. (2007) attribute this apparent reduction in

TOC to contamination of the Brewer’s direct measurements by diffuse radiation in the instrument’s field of view, quoting a previous conclusion by Zerefos et al. (2000; 2001). We find this conclusion surprising considering that changes in the direct-to-diffuse ratio during an eclipse are small according to theoretical calculations by Emde and Mayer (2007), with the exception of a short (< 5 min) period before and after the period of totality. We speculate that the observed drop in Brewer TOC measurements during an eclipse is partly caused by the fact that Brewer data are typically calculated from four instead of two wavelengths, as it is the case for our results. Using more than two wavelengths has the advantage that aerosol effects can be reduced (Basher, 1982). However, this method could potentially lead to errors if wavelength-dependent changes in irradiance caused by the LD effect are incorrectly attributed to aerosols. For example, using a LD correction method based on the NLTE Spectral Synthesis code (Tagirov et al. 2016), Gröbner et al. (2017) provide evidence that the spurious drop in Brewer TOC measurements is primarily due to LD effects. Of note, the LD correction applied to Dobson spectrophotometer data from the solar eclipse of 20 May 1966 discussed of Bojkov (1968) reduces the TOC (calculated from the Dobson C pair) by up to 6 %, in good agreement with our calculations.

Chimonas (1970) hypothesized that the Moon’s shadow, as it travels at supersonic speeds across the atmosphere during a solar eclipse, induces “bow waves” (a type of gravity waves) due to sudden cooling of the atmosphere at altitudes between 10 and 60 km, where the ozone layer normally converts solar UV radiation to heat. Ionospheric bow waves with wavelengths between 300 and 400 km and a period of about 25 minutes, which manifested themselves as electron content disturbances, have indeed been observed during the solar eclipse of 21 August 2017 (Zhang et al., 2017). If these bow waves had affected the ozone layer, these oscillations should have been detectable with our sampling frequency of one TOC value every two minutes.

Using data measured during the solar eclipse of 11 August 1999, Zerefos et al. (2000) observed an oscillation with a period of 20 minutes in erythemal irradiance before and after the time of the eclipse with a peak-to-peak variation of about 1 %. Zerefos et al. (2000) attributed these oscillation to fluctuations in total ozone caused by eclipse-induced gravity waves. A variation of ± 0.5 % in erythemal irradiance would require a variation in total ozone of similar magnitude (McKenzie et al., 2011). In a similar analysis, using data from the 29 March 2006 solar eclipse, Zerefos et al. (2007) reports oscillations in TOC with a peak-to-peak amplitude of 2.0–3.5 % and periods ranging between 30 and 40 minutes. However, these measurements also indicate an unrealistic drop in the TOC from 295 to 225 DU between the beginning and the maximum of the eclipse, followed by an increase to 300 DU. This artifact is attributed to the contamination of direct irradiance measurements (which are used for the ozone retrieval) by diffuse radiation. It is concerning that the magnitude of this artifact (75 DU or 25 %) exceeds the magnitude attributed to the bow wave by a factor of 10. Mims and Mims (1993) report that TOC measurements taken during the total solar eclipse of 11 July 1991 show fluctuations in ozone with a peak-to-peak amplitude of up to 5 DU (1.7 %), which began 700 seconds after the third contact. However also these observations were affected by incomplete LD correction, resulting in TOC changes of 26 DU (about 9 %) over the course of the eclipse.

Deleted: does virtually not change

Deleted: the

Deleted: (both

Deleted: our measurements and the

Deleted:)

Deleted: U

Deleted: additional

Deleted: systematic bias

Deleted: Bow waves with wavelengths between of 300 and 400 km and a period of about 25 minutes have indeed been observed during the solar eclipse of 21 August 2017 (Zhang et al., 2017).¶

Deleted: (sunburning)

Deleted: Finally,

Finally, Kazantzidis et al (2007) discuss TOC measurements performed with NILU-UV filter radiometers at several locations in Greece during the total solar eclipse of 29 March 2006. They did not observe any periodic fluctuations in TOC and only report a small increase in TOC of about 5 DU as the visible fraction of the Sun decreases from ~60% to ~20%. This small change could be caused by incomplete correction of the LD effect.

Our data do not support the observation by Zerefos (2000; 2007) and Mims and Mims (1993) that bow waves from the Moon's shadow lead to oscillations in TOC with a peak-to-peak amplitude exceeding 1.5 %. For example, between 17:36 and 18:36, LD-corrected TOC data decreased by 1 DU (0.33 %) with no obvious oscillations. Between 18:38 and the end of the measurements at 19:32, TOC remained constant to within 0.3 DU (0.10 %). These small variations are well within the natural variability of the TOC and the uncertainty of our TOC retrieval of 1.5 DU (0.5 %), which is mainly caused by the effect of changing aerosols on ozone calculations. The lack of an effect is not surprising considering that diurnal variations in ozone are small. For example, at the Mauna Loa observatory (19.5° S), the day/night difference of the ozone profile is smaller than ± 2 % for altitudes up to 45 km where 99 % of the ozone column is located (Parrish et al., 2014). Since day/night changes below and above 30 km have a different sign and partly cancel each other, the effect on TOC is small. It would be surprising if TOC variations over the relatively short period of a solar eclipse driven by bow waves were larger than variations caused by the diurnal cycle. Despite the results by Zerefos (2000; 2007) and Mims and Mims (1993), we feel that the question of whether or not bow waves from the Moon's shadow can lead to variations in TOC is still up for debate. This debate could be settled by performing LD-corrected measurements of TOC with different instrument types during one of the upcoming solar eclipses. If such measurements were to show fluctuations in TOC with the same magnitude and timing, the effect of bow waves on TOC could be convincingly demonstrated.

Deleted: negligible

Deleted: driven

Deleted: ¶

8.4 Validation of GUVis-3511 measurements

The GUVis-3511 is a relatively new instrument and the measurements discussed here present a unique opportunity to examine some aspects of the instrument's performance. For example, as radiation levels decrease and increase by more than four orders of magnitude over the course of the eclipse (Fig. 15), the linearity of the measurements can be assessed. The GUVis-3511 uses a 24-bit amplifier with three gains, providing a dynamic range of over 11 orders of magnitude. Measurements of channels with wavelength larger than 400 nm change by 8.5 orders of magnitude between dark measurements (collector of the instrument capped) and full sunlight, and at some point, have to switch between "high" and "medium" gains to extend the ADC's dynamic range of about 7 orders of magnitude. Non-linearities have been noted for legacy instruments when the gain changes and may manifest itself as steps in the ratio of the instrument's measurements and a smoothly varying reference. Because every channel changes the gain at a different time, such step-changes would also occur at different times. While ratios of measurement and theory shown in Fig. 8d and Fig. 9 do exhibit some step changes of up to 0.5 % (e.g., when global measurements resume after shadowbanding), such steps occur simultaneously for all channels.

We estimate from this analysis that the non-linearity caused by gain changes are smaller than 0.1 % and there is also no evidence of non-linearity at other times.

Measurements during totality (Fig. 14) suggest that the detection limit of the GUVis-3511 radiometer is about $10^{-4} \mu W cm^{-2} nm^{-1}$. This detection limit is comparable with that of scanning spectroradiometers with PMT detector, which are considered the most accurate instruments for observing the solar irradiance at Earth's surface (Seckmeyer et al., 2001). While multi-channel filter radiometer such as the GUVis-3511 do not have the wavelength resolution of these spectroradiometers, they have a superior sampling rate than these instruments (e.g., 1 Hz versus five minutes for one spectrum), and are therefore better suited to study fast-changing phenomena such as the rapid change in irradiance close to the 2nd or 3rd contact.

9 Conclusions and outlook

Measurements of spectral irradiance during the total solar eclipse of 21 August 2017 were used to validate parameterizations of solar LD and to assess the effects of the eclipse on [the ratio of direct-to-global irradiance](#), AOD, [and](#) TOC. For wavelengths longer than 400 nm, the change of global spectral irradiance over the period of the eclipse agrees to within $\pm 4.0\%$ with theoretical predictions that were derived from two independent parameterizations of solar LD by Pierce and Neckel. When excluding a 16-minute period, which was likely affected by [excess amounts of](#) aerosols, the agreement is to within $\pm 2.5\%$. Between 315 and 400 nm, differences between observation and theory are smaller than 5.5 % and increase to about 9 % at 306 nm. In the visible and infrared ranges, these differences are similar in magnitude before and after totality. For wavelengths between 306 and 340 nm, the difference is larger between the 1st and 2nd contact than between the 3rd and 4th contact. [Most of this asymmetry is likely](#) caused by changing aerosol conditions.

Deleted: , and the ratio of direct-to-global irradiance.

Deleted: A part of

Deleted: could be

Several recent papers (e.g., Blumthaler et al., 2006; Emde and Mayer, 2007; Kazadzis et al., 2007) have used the LD parameterization by Waldmeier, which does not consider the wavelength dependence of hydrogen absorption in the Sun's photosphere. When we compared our observations with data derived from this parameterization, observation were lower than theory by up to 37 % at 306 nm and 19 % at 412 nm. For longer wavelengths, differences [were](#) in the $\pm 8\%$ range. These differences are well beyond the uncertainty of our observations, suggesting that the parameterization by Waldmeier is too simple and should not be used.

Deleted: are

Results shown in Fig. 13 illustrate that accurate TOC measurements during a solar eclipse can only be achieved if the wavelength-dependence of LD is appropriately corrected. Without correction, TOC values [derived from the 305 nm / 340 nm wavelength pair](#) increase with increasing solar occultation and are too high by 23 DU (or 8.2 %) shortly before the 2nd contact. Using the LD correction by Waldmeier, this bias is reduced to 19 DU. When data are corrected using the parameterization by either Pierce or Neckel, TOC measurements stay constant to within ± 2.6 DU during the duration of the

eclipse. Our conclusion that the LD correction is important contradicts findings of several recent papers, which determined that the LD effect is either less than 0.01 % (Blumthaler et al., 2006; Kazadzis et al., 2007), 1 % (Zerefos et al., 2000), or less than 1.6 % (Kazantzidis et al., 2007). As a result, TOC measurements reported in these papers show spurious changes over the course of an eclipse.

Our data do not support the conclusions by Zerefos (2000; 2007) that gravity or bow waves, which are set off by the Moon’s shadow moving at supersonic over the atmosphere, lead to variations the TOC. For example, variations in TOC are smaller than ± 1.0 DU (± 0.33 %) between the 3rd and 4th contact and smaller than ± 0.15 DU (± 0.05 %) between the 4th contact and the end of our measurements. These variations are well within the natural variability and the uncertainty of our measurements.

There is no clear evidence that the solar eclipse drives changes in the ratio of direct-to-global irradiance for the period accessible from our measurements (i.e., up to 200 s before the 2nd contact and 105 seconds after the 3rd contact). The observed decline in this ratio (up to 20% in the UV-B, 15% in the visible, and 3.5% in the IR) can be explained with variability from aerosols stemming from wildfires. Invariance of this ratio is expected from theory (Emde and Mayer 2007), but contradicts results by Zerefos et al. (2000), who determined that the ratio of diffuse-to-direct erythral irradiance increased by 50 % during the eclipse studied by Zerefos et al. (2000), peaking at the time of the eclipse maximum.

Deleted: small changes

Deleted: The i

Deleted: radiation

Deleted: s gradually

Deleted: 3

During totality, the global spectral irradiance was decreased by factors of 18,700 at 315 nm, 7,000 at 340 nm, 18,340 at 665 nm, 136,650 at 940 nm and 46,150 at 1020 nm. The average reduction for channels in the visible (412–665 nm) is 11,460, which is about a factor of 35 smaller than the ratio of about 400,000 between the irradiance from the Sun and the full Moon.

Results also confirmed that the GUVis-3511 radiometer is an excellent instrument for observing the fast-changing radiation levels during an eclipse with a detection limit of 10^{-4} $\mu\text{W cm}^{-2} \text{ nm}^{-1}$. By implementing a new calibration method developed for the specific conditions during the eclipse, measurements with low uncertainty (e.g., 2.7 % for wavelengths between 320 and 875 nm) can be achieved. Comparisons between measurement and theory did not indicate any signs of non-linearity over the dynamic range of about four orders of magnitude between full sunlight and totality.

Lastly, data from the campaign are available as Supplement and can be used to study the rapid change in irradiance during the transition from a partial to a total eclipse with the help of 3DRT models like that developed by Emde and Mayer (2007). During this transition, the radiation field changes fundamentally as the Sun’s direct component is removed and the illumination becomes entirely diffuse. During totality, the irradiance at the surface will become also more sensitive to the topography (e.g., the mountains surrounding the measurement sites (Fig. 1)), surface albedo (and its spectral dependence),

and the distribution of ozone in the atmosphere (the ozone profile). These aspects will be discussed in a follow-on publication.

Data availability

Calibrated data and data used to draw Fig. 3, Fig. 8 and Fig. 14 are available as a Supplement.

5 Author contribution

GB designed and executed the measurements, performed the data analysis, and wrote the manuscript.

BP contributed to the implementation of the solar limb darkening parameterizations by Pierce and Neckel and contributed to writing the manuscript.

Competing interests

10 GB is employed by Biospherical Instruments Inc, which is also the manufacturer of the GUVis-3511 radiometer described in this paper.

Acknowledgements. GB thanks Biospherical Instrument Inc. for supporting this activity and Anne Hoppe for her encouragement to travel to Oregon and her help in setting up the experiment. We thank Bjørn Johnsen from the Norwegian
15 Radiation Protection Authority ([now Norwegian Radiation and Nuclear Safety Authority](#)) for performing spectral response measurements of the GUVis-3511 radiometer used in this campaign. We also thank Bernhard Mayer and Paul Ockenfuß from the Ludwig-Maximilians-Universität München, Munich, Germany, for discussing our results and for their valuable suggestions. [Lastly, we thank two anonymous reviewers and Forrest M. Mims III for their comments, which greatly improved the manuscript.](#)

20 References

Adams, C., McLinden, C. A., Strong, K., and Umlenski, V.: Ozone and NO₂ variations measured during the 1 August 2008 solar eclipse above Eureka, Canada with a UV-visible spectrometer, J. Geophys. Res., 115(D19), <https://doi.org/10.1029/2010JD014424>, 2010.

25 Antón, M., Serrano, A., Cancillo, M. L., Vaquero, J. M., and Vilaplana J. M.: Solar irradiance and total ozone over El Arenosillo (Spain) during the solar eclipse of 3 October 2005, J. Atmos. Sol.-Terr. Phy., 72(9–10), 789–793, <https://doi.org/10.1016/j.jastp.2010.03.025>, 2010.

- Basher, R. E., Review of the Dobson spectrophotometer and its accuracy, WMO Global Ozone Res. Monit. Proj. Rep. 13, World Meteorol. Organ., Geneva, Switzerland, 1982.
- Bass, A. and Paur, R. J.: The ultraviolet cross sections of ozone: 1, The measurement, in Atmospheric Ozone, edited by C. Zerefos and A. Ghazi, pp. 606–616, D. Reidel, Norwell, Mass, 1985.
- 5 Bernhard, G., Booth, C. R., and McPeters, R.D.: Calculation of total column ozone from global UV spectra at high latitudes. *J. Geophys Research*, 108(D17), 4532, doi:10.1029/2003JD003450, 2003.
- Bernhard, G., Booth, C. R., and Ehranjian, J. C.: Version 2 data of the National Science Foundation's Ultraviolet Radiation Monitoring Network – South Pole, *J. Geophys. Res.*, 109, D21207, doi:10.1029/2004JD004937, 2004.
- Bernhard, G., Booth, C. R., and Ehranjian, J. C.: Real-time ultraviolet and column ozone from multichannel ultraviolet radiometers deployed in the National Science Foundation's ultraviolet monitoring network, *Opt. Eng.*, 44(4), 041011-1, 10 https://doi.org/10.1117/1.1887195, 2005a.
- Bernhard, G., Evans, R. D., Labow, G. J., and Oltmans, S. J.: Bias in Dobson total ozone measurements at high latitudes due to approximations in calculations of ozone absorption coefficients and air mass, *J. of Geophys. Res.*, 110, D10305, 15 https://doi.org/10.1029/2004JD005559, 2005b.
- Blumthaler, M., Bais, A., Webb, A., Kazadzis, S., Kift, R., Kouremeti, N., Schallhart, B., and Kazantzidis, A.: Variations of solar radiation at the Earth's surface during the total solar eclipse of 29 March 2006. In: Remote Sensing of Clouds and the Atmosphere XI, 6362, p. 63620F, International Society for Optics and Photonics, 20 https://doi.org/10.1117/12.6896302006, 2006.
- Bodhaine, B. A. and Dutton, E. G.: A long-term decrease in Arctic haze at Barrow, Alaska, *Geophys. Res. Lett.*, 20, 947–950, https://doi.org/10.1029/93GL01146, 1993.
- Bojkov, R. D.: The ozone variations during the solar eclipse of 20 May 1966, *Tellus*, 20(3), 417–421, https://doi.org/10.1111/j.2153-3490.1968.tb00382.x, 1968.
- Chimonas, G. and Hines, C. O.: Atmospheric gravity waves induced by a solar eclipse, 2. *J. Geophys. Res.*, 76(28), 7003–7005, https://doi.org/10.1029/JA075i004p00875, 1971.
- 25 Emde, C., and Mayer, B: Simulation of solar radiation during a total eclipse: a challenge for radiative transfer, *Atmos. Chem. Phys.*, 7(9), 2259–2270, https://doi.org/10.5194/acp-7-2259-2007, 2007.
- Gröbner, J., Kazadzis, S., Kouremeti, N., Doppler, L., Tagirov, R. and Shapiro, A. I.: Spectral solar variations during the eclipse of March 20th, 2015 at two European sites. In AIP Conference Proceedings, 1810(1), p. 080008, 30 https://doi.org/10.1063/1.4975539, 2017.
- Hooker, S. B., Bernhard, G., Morrow, J. H., Booth, C. R., Comer, T., Lind, R. N., and Quang, V.: Optical Sensors for Planetary Radiant Energy (OSPRey): calibration and Validation of Current and Next-Generation NASA Missions., NASA Goddard Space Flight Center, NASA/TM–2011–215872, 2012.
- Kasten, F. and Young, A. T.: Revised optical air mass tables and approximation formula, *Appl. Opt.*, 28(22), 4735–4738, https://doi.org/10.1364/AO.28.004735, 1989.

- Kazadzis, S., Bais, A., Blumthaler, M., Webb, A., Kouremeti, N., Kift, R., Schallhart, B., and Kazantzidis, A.: Effects of total solar eclipse of 29 March 2006 on surface radiation, *Atmos. Chem. Phys.*, 7(22), 5775–5783, <https://doi.org/10.5194/acp-7-5775-2007>.
- Kazantzidis, A., Bais, A. F., Emde, C., Kazadzis, S., and Zerefos, C. S.: Attenuation of global ultraviolet and visible irradiance over Greece during the total solar eclipse of 29 March 2006, *Atmos. Chem. Phys.*, 7(23), 5959–5969, <https://doi.org/10.5194/acp-7-5959-2007>, 2007.
- Koepke, P., Reuder, J., and Schween, J.: Spectral variation of the solar radiation during an eclipse, *Meteorol. Z.*, 10(3), 179–186, <https://doi.org/10.1127/0941-2948/2001/0010-0179>, 2001.
- Krotkov, N. A., Bhartia, P. K., Herman, J. R., Slusser, J. R., Labow, G. J., Scott, G. R., Janson, G., T. Eck, T. and Holben, B. N.: Aerosol ultraviolet absorption experiment (2002 to 2004), part 1: ultraviolet multifilter rotating shadowband radiometer calibration and intercomparison with CIMEL sunphotometers, *Opt. Eng.*, 44(4), 041004-1 - 041004-17, 2005.
- Mateos, D., Antón, M., and Vaquero, J. M.: Influence of solar eclipse of November 3rd, 2013 on the total ozone column over Badajoz, Spain., *J. Atmos. Sol.-Terr. Phys.*, 112, 43–46, <https://doi.org/10.1016/j.jastp.2014.02.005>, 2014.
- Macdonald, L.: Our Sun, In: *How to Observe the Sun Safely*, pp. 1–16, Springer, New York, https://doi.org/10.1007/978-1-4614-3825-0_1, 2012.
- McKenzie, R. L., Aucamp, P. J., Bais, A. F., Björn, L. O., Ilyas, M., and Madronich, S.: Ozone depletion and climate change: impacts on UV radiation, *Photochem. Photobiol. Sci.*, 10(2), 182–198, <http://doi.org/10.1039/C0PP90034F>, 2011.
- Mims III, F. M. and Mims, E. R.: Fluctuations in column ozone during the total solar eclipse of July 11, 1991, *Geophys. Res. Lett.*, 20(5), 367–370, <https://doi.org/10.1029/93GL00493>, 1993.
- Morrow, J. H., Hooker, S. B., Booth, C. R., Bernhard, G., Lind, R. N., and Brown, J. W.: Advances in measuring the apparent optical properties (AOPs) of optically complex waters, NASA/TM–2010–215856, National Aeronautics and Space Administration, Goddard Space Flight Center, 2010.
- Neckel, H.: Analytical Reference Functions $F(\lambda)$ for the Sun's Limb Darkening and Its Absolute Continuum Intensities ($\lambda\lambda$ 300 to 1100 m). *Solar Physics*, 229(1), 13–33, <https://doi.org/10.1007/s11207-005-4081-z>, 2005.
- O'Neill, N. T., Eck, T. F., Holben, B. N., Smirnov, A., Royer, A., and Li, Z.: Optical properties of boreal forest fire smoke derived from Sun photometry. *J. Geophys. Res.*, 107(D11), 4125, <https://doi.org/10.1029/2001JD000877>, 2002.
- Parrish, A., Boyd, I. S., Nedoluha, G. E., Bhartia, P. K., Frith, S. M., Kramarova, N. A., Connor, B. J., Bodeker, G. E., Froidevaux, L., Shiotani, M., and Sakazaki, T.: Diurnal variations of stratospheric ozone measured by ground-based microwave remote sensing at the Mauna Loa NDACC site: measurement validation and GEOSCCM model comparison, *Atmos. Chem. Phys.*, 14(14), 7255–7272, doi:10.5194/acp-14-7255-2014, 2014.
- Pasachoff, J. M.: The Great Solar Eclipse of 2017, *Sci Am.*, 317(2), 54–61, <https://10.1038/scientificamerican0817-54>, 2017.

- Pierce, A. K., and Slaughter, C. D.: Solar limb darkening, *Solar Physics*, 51(1), <https://doi.org/10.1007/BF00240442>, 25-41, 1977.
- Pierce, A. K., Slaughter, C. D., and Weinberger, D.: Solar limb darkening in the interval 7404–24 018 Å, II. *Solar Physics*, 52(1), 179–189, <https://doi.org/10.1007/BF00935800>, 1977.
- 5 Seckmeyer, G. and Bernhard, G.: Cosine error correction of spectral UV-irradiances. In *Atmospheric Radiation*, Vol. 2049, pp. 140–152, International Society for Optics and Photonics, <https://doi.org/10.1117/12.163505>, 1993.
- Seckmeyer, G., Bais, A., Bernhard, G., Blumthaler, M., Booth, C. R., Disterhoft, P., Erikson, P., McKenzie, R. L., Miyauchi, M., and Roy, C.: Instruments to measure solar ultraviolet radiation, Part 1: Spectral instruments, *World Meteorological Report* 125, WMO TD 1066, 2001.
- 10 Seckmeyer, G., Bais, A., Bernhard, G., Blumthaler, M., Johnsen, B., Lantz, K., and McKenzie, R.: Instruments to measure solar ultraviolet radiation, Part 3: Multi-channel filter instruments. *World Meteorological Organization Global Atmosphere Watch*, Report 190, WMO TD 1537, 2010.
- Sharp, W. E., Silverman, S. M., Lloyd, J. W. F.: Summary of sky brightness measurements during eclipses of the sun, *Appl. Opt.* 10(6), 1207–1210, <https://doi.org/10.1364/AO.10.001207>, 1971.
- 15 Stamnes, K., Slusser, J., and Bowen, M.: Derivation of total ozone abundance and cloud effects from spectral irradiance measurements, *Appl. Opt.*, 30(30), 4418–4426, <https://doi.org/10.1364/AO.30.004418>, 1991.
- Tagirov, R. V., Shapiro, A. I., and Schmutz, W.: NESSY: NLTE spectral synthesis code for solar and stellar atmospheres. *Astron. Astrophys.*, 603(A27), DOI: 10.1051/0004-6361/201628574, 2017.
- Waldmeier, M.: *Ergebnisse und Probleme der Sonnenforschung*, *Probleme der kosmischen Physik*, 264 pp., Akad. Verlagsges. Becker & Erler, Leipzig, Germany, 1941.
- 20 Witthuhn, J., Deneke, H., Macke, A., and Bernhard, G.: Algorithms and uncertainties for the determination of multispectral irradiance components and aerosol optical depth from a shipborne rotating shadowband radiometer, *Atmos. Chem. Phys.*, 10(2), 709–730, <https://doi.org/10.5194/amt-10-709-2017>, 2017.
- Zerefos, C. S., Balis, D. S., Meleti, C., Bais, A. F., Tourpali, K., Kourtidis, K., K. Vanicek, K., Cappellani, F., Kaminski, U., Colombo, T., and Stübi, R.: Changes in surface solar UV irradiances and total ozone during the solar eclipse of August 11, 1999, *J. Geophys. Res.*, 105(D21), 26463–26473, <https://doi.org/10.1029/2000JD900412>, 2000.
- 25 Zerefos, C. S., Balis, D. S., Zanis, P., Meleti, C., Bais, A. F., Tourpali, K., D.Melas, D., Ziomas I., Galani E., Kourtidis, K., Papayannis A., and Gogosheva Z.: Changes in surface UV solar irradiance and ozone over the Balkans during the eclipse of August 11, 1999, *Adv. Space Res.*, 27(12), 1955–1963, [https://doi.org/10.1016/S0273-1177\(01\)00279-4](https://doi.org/10.1016/S0273-1177(01)00279-4), 2001.
- 30 Zerefos C. S., Gerasopoulos E., Tsagouri I., Psiloglou B. E., Belehaki A., Herekakis T., Bais A., Kazadzis S., Eleftheratos C., Kalivitis N., and Mihalopoulos N.: Evidence of gravity waves into the atmosphere during the March 2006 total solar eclipse, *Atmos. Chem. Phys.*, 7(18), 4943–4951, <https://doi.org/10.5194/acp-7-4943-2007>, 2007.

Deleted: (

Deleted:)

Zhang, S.-R., Erickson, P. J., Goncharenko, L. P., Coster, A. J., Rideout, W., and Vierinen, J.: Ionospheric bow waves and perturbations induced by the 21 August 2017 solar eclipse, *Geophys. Res. Lett.*, 44(24), <https://doi.org/10.1002/2017GL076054>, 2017.