

Supplement to
"Indications for a potential synchronization
between the phase evolution of the
Madden-Julian oscillation and the solar 27-day
cycle"

Christoph G. Hoffmann and Christian von Savigny

Contents

1	Analysis repeated with the epoch window centered around the solar extremum	2
2	Analysis repeated with equal weights	5
3	Examples of sawtooth fits for nonoptimal filtering conditions	8

1 Analysis repeated with the epoch window centered around the solar extremum




Figure S1: As Fig. 6 in the manuscript, but based on the analysis variant, in which the epochs cover ± 14 days around solar extremum. See Sect. 4.2 in the manuscript for details.




Figure S2: As Fig. 7 in the manuscript, but based on the analysis variant, in which the epochs cover ± 14 days around solar extremum. See Sect. 4.2 in the manuscript for details.




Figure S3: As Fig. 8 in the manuscript, but based on the analysis variant, in which the epochs cover ± 14 days around solar extremum. See Sect. 4.2 in the manuscript for details.




Figure S4: As Fig. 10 in the manuscript, but based on the analysis variant, in which the epochs cover ± 14 days around solar extremum. See Sect. 4.2 in the manuscript for details.




Figure S5: As Fig. 11 in the manuscript, but based on the analysis variant, in which the epochs cover ± 14 days around solar extremum. See Sect. 4.2 in the manuscript for details.

2 Analysis repeated with equal weights




Figure S6: As Fig. 6 in the manuscript, but based on the analysis variant, in which each data point was equally weighted for the calculation of the deviation. See Sect. 4.2 in the manuscript for details.




Figure S7: As Fig. 7 in the manuscript, but based on the analysis variant, in which each data point was equally weighted for the calculation of the deviation. See Sect. 4.2 in the manuscript for details.




Figure S8: As Fig. 8 in the manuscript, but based on the analysis variant, in which each data point was equally weighted for the calculation of the deviation. See Sect. 4.2 in the manuscript for details.




Figure S9: As Fig. 10 in the manuscript, but based on the analysis variant, in which each data point was equally weighted for the calculation of the deviation. See Sect. 4.2 in the manuscript for details.




Figure S10: As Fig. 11 in the manuscript, but based on the analysis variant, in which each data point was equally weighted for the calculation of the deviation. See Sect. 4.2 in the manuscript for details.

3 Examples of sawtooth fits for nonoptimal filtering conditions




Figure S11: As Fig. 4 in the manuscript, but for QBO westerly phase conditions. The experimental setup corresponds with an MJO strength threshold of 1.1 to the weakest similarity of the QBO experiments (maximum of the blue curve in Fig. 6 of the manuscript). Nevertheless, the sawtooth shape is still rudimentarily present, albeit highly insignificant as 93% of the Monte Carlo runs resulted in lower deviations (see manuscript for details).




Figure S12: As Fig. 4 in the manuscript, but not filtered for seasons. The experimental setup corresponds with an MJO strength threshold of 1.7 to the weakest similarity of the all-season experiments (maximum of the red curve in Fig. 7 of the manuscript). Also here, the sawtooth shape is still rudimentarily present, albeit highly insignificant as 77% of the Monte Carlo runs resulted in lower deviations (see manuscript for details).




Figure S13: As Fig. 4 in the manuscript, but for summer conditions. The experimental setup corresponds with an MJO strength threshold of 0.5 to the weakest similarity of the season experiments (maximum of the magenta curve in Fig. 7 of the manuscript).




Figure S14: As Fig. 4 in the manuscript, but based on the RMM instead of the OMI index. The experimental setup corresponds with an MJO strength threshold of 1.2 to the weakest similarities of the RMM experiments (local maximum of the red curve in Fig. 10 of the manuscript). Also here, some sawtooth-like structure is still rudimentarily present, albeit highly insignificant as 47% of the Monte Carlo runs resulted in lower deviations (see manuscript for details).