

Comments (in *italics*), responses, and marked-up revised manuscript

General Comment from Anonymous Referee #1

The manuscript reports single-particle analysis from size-selected aerosol samples collected at two sites in the Amazon basin, ATTO forest site and Manaus city center, in the wet season of 2012. Results from the forest site samples confirm previous observations reported in the literature, with a predominance of SOA particles in the smaller size ranges. Results from the Manaus samples attest the influence of local urban emissions, as well as the influence of regional biogenic emissions and sea salt. In my opinion, the discussion of the results from Manaus, which is the novelty brought by this manuscript, could be improved. The manuscript is reasonably well written, but there are some redundancies that could be omitted.

Response: We thank the reviewer for his/her positive evaluation of our work and valuable comments. We revised the manuscript as much as possible respecting the reviewer's comments.

Comments to the text from Anonymous Referee #1

1) General comment on the introduction: I suggest restructuring the order of the contents. First, present general characteristics of the Amazon region and its importance, mentioning the ATTO site and the Manaus city. Then, focus on aerosol sources in the wet season, and previous studies on single-particle characterization in Amazonia.

Response: As suggested, the introduction was revised (see the revised introduction).

2) Line 65: interaction with? Maybe it would be better to replace “interaction” with “dynamical processes”.

Response: Done as suggested.

3) Line 65: aerosols are airborne by definition, so I suggest omitting “airborne” here.

Response: Done as suggested.

4) Lines 182-188: *These sentences, near the beginning of the results section, refer to a previous study, right? It might create a confusion between your current results and previous studies. Please reformulate, to make clear what your contribution is.*

Response: The sentences were modified mainly by inserting the phrase, “In a previous study (Pöschl, et al., 2010)”, to avoid the possible confusion. The sentence, “With the exception of the reacted sea-salt particles probably from the Atlantic Ocean as well as the abundant ammonium sulfate aerosols, the particle types observed in this study are comparable to their study”, was given (lines: 191-196).

5) Line 351: *Replace “nascent airborne” by “fresh”.*

Response: Done as suggested.

6) Line 377: *Replace “size” by “diameter”.*

Response: Done as suggested.

7) Line 394: *there is a typo here (90%).*

Response: Done as suggested.

8) Lines 439-445: *It is not necessary to describe the figure and its numbers in words, like here. I suggest that you go straight to the interpretation of the figure results.*

Response: Revised as suggested (lines: 458-462, 471-473, 476-477).

9) Section 3.2: *I recommend that you omit redundancies in this section. You already discussed the particle formation processes and its contribution to CCN and IN, there is no need to repeat it. For example, in lines 450-453 and 474-480.*

Response: As recommended, the redundancies were omitted.

Specific Comments from Anonymous Referee #1

1) Lines 77-78: There is not a unique pathway to produce SOA in Amazonia. Condensation of low volatility organic species can also occur at the surface of PBA, for example.

Response: In order to avoid such impression, a “mainly” word was added – “SOA particles are mainly formed through the condensation of biogenic organic compounds onto biogenic K-rich salt particles emitted from the forest”.

2) Lines 84-87: This is true for biomass burning aerosols in Amazonia. This sentence does not fit well into this paragraph, since you are talking mostly about biogenic aerosols.

Response: The reviewer’s comment is right. This part is deleted without deteriorating the context.

3) Lines 97-98: This sentence is unclear. First, you might say that Fraund et al. (2017) used cluster analysis to identity particle types. Second, what do you mean by "anthropogenic elements"?

Response: The reviewer is right as the sentence is not clear for people who are not familiar with the paper. Without deteriorating the context, the sentence was omitted.

4) Lines 107-108: One third of what? Mass? Number? PM or OOA (oxygenated organic aerosols)? Please clarify.

Response: Thank the reviewer for the comment. The sentence was modified as “where one third of the potential SOA would be of an urban origin”.

5) Lines 109-116: Compared to previous works, I would say that the main novelty of this study is the analysis of aerosol samples from the Manaus city center. Emphasize that.

Response: As suggested, the sentence, “Especially, single-particle characterization of aerosols collected at a Manaus city center has been scarce.” was added (lines: 75-76).

6) Line 127: Please provide more details about the city sampling site. Was it located in the southern or northern part of the city? What was the approximate distance to busy roads/avenues/streets? Do you know if vehicular types were mostly light duty, heavy duty, or both? Is there potential influence of industrial or power plant emissions? The Manaus urban plume is very heterogeneous. You can find more information at de Sá et al., 2018 and Medeiros et al., 2017.

Response: More detailed description is given – “In Manaus, the sampling site is situated in the central part of the city (S 03° 05.753’, W 59° 59.419’), which is at a representative urban region influenced by electricity production based on fuel oil, diesel, and natural gas, biogenic emissions from the surrounding forest, and mostly by light duty (using gasoline and ethanol) vehicle traffic. Heavy vehicles that use diesel account to less than 10 % of the urban fleet (Medeiros et al., 2017). The location is nearby a small parking area and around 200 m away from the intersection of four busy avenues, with frequent diurnal traffic jam on weekdays.” (lines: 126-132).

7) Line 132: information on sampling dates and sample names would be better suited in a table.

Response: As suggested by the reviewer, information on samples and meteorological parameters (temp., RH, rainfall, and meteorological conditions during the samplings) were given in Table 1.

8) Line 135: How were the samples conserved during transport and storage? Semivolatile organic compounds can volatilize when subjected to temperature variations. Do you think that this is an important issue in your measurements?

Response: During the transport or storage of the samples, semivolatile organic compounds can volatilize as the reviewer pointed out. However, more critically, the particles are under high vacuum during the EPMA measurements, which makes the detection of semivolatile components impractical. The EPMA measurements are not for semivolatile components in aerosols.

9) Line 141: You must provide more information about the backtrajectories calculation. What was the input meteorological database that you used? In which spatial resolution? How many trajectories were calculated for each aerosol sample?

Response: As suggested, a sentence was added – “In the HYSPLIT calculation, meteorological data output from the Global Data Assimilation System (GDAS) using GDAS1 data with a horizontal resolution of 1° corresponding to ~ 100 km × 100 km and 23 vertical layers was used, which was reported to provide plausible backward trajectory analysis (Su et al., 2015).” (lines: 148-151).

10) Lines 263-264: Is this true (ammonium sulfate dominant over ammonium bisulfate) both for Manaus and ATTO? Is it possible to provide a percentage of the relative amounts of sulfate in each

form (ammonium sulfate, ammonium bisulfate, sulfuric acid)? How does it compare to other metropolises in the world?

Response: Raman spectroscopy used in this work detected the ammonium sulfate in the samples, but no ammonium bisulfate and sulfuric acid. The Raman analysis was for the qualitative analysis rather than the quantitative one.

11) Line 274: I recommend citing Saturno et al. (2018), about the influence of African volcanic emissions on sulfate concentrations at the ATTO site.

Response: As suggested by the reviewer, Saturno et al. (2018) was cited – “including that from the African volcanic emissions (Saturno et al., 2018)” (lines: 284-285).

12) Lines 287-289: You discussed the potential sources of SO₄ and NH₄ in the Amazon forest, based on previous studies. How about the potential sources in Manaus city? You may refer to Medeiros et al., 2017 and de Sá et al., 2018. Also, you may refer to the literature produced on this subject for other metropolises in the world.

Response: A sentence was added for major ammonium sulfate sources in the urban environment – “In the urban environment, anthropogenic ammonium sulfate is mainly formed by gaseous reactions among SO₂ emitted from coal-fired plants and industrial activities, NH₃ emitted from human and animal activities and fertilization in the fields, and oxidants (e.g., O₃ and OH radical) (Li et al., 2016; Geng et al., 2017).” (lines: 301-304).

13) Lines 343-345: Here you could hypothesize on the potential sources of nitrate in Manaus.

Response: As suggested, a phrase was added – “The nitrates for the Manaus samples may be formed from nitrogen oxides emitted from the vehicles and coal-fired power plants (Geng et al., 2014, 2017; Li et al., 2016).” (lines: 359-361).

14) Line 364-366: Yes, biomass burning emissions reduce significantly in the Amazonian wet season. However, advection from Africa brings biomass burning aerosols, besides Saharan dust, to the Amazon region in March-April. See, for example, Baars et al., 2011. So, part of the observed K-salts could be associated with biomass burning aerosols from Africa.

Response: Thank the reviewer for the comment. A phrase was added – “although it was reported that a strong biomass burning smoke was transported from Africa to South America during the wet season (Baars et al., 2011) so that a part of the observed K-salts could be associated with biomass burning aerosols from Africa.” (lines: 382-385).

15) Line 405: You may cite a reference for the term "brown carbon". It could be either Andreae et al., 2006 or Laskin et al., 2010.

Response: Done as suggested.

16) Lines 467-469: A single backtrajectory is insufficient to prove the occurrence of African aerosol advection. You may add a sentence recognizing that.

Response: As suggested, a phrase was added – “although a single backtrajectory cannot confirm the African origin of the aged mineral dust particles.” (lines 482-483).

17) Lines 497-499: The reasoning is not strong. Part of the SOA observed at Manaus can be locally produced. You would need a biogenic tracer to make this point. A better indication of the influence of forest emissions in Manaus is the 20% of PBA in sample SM3, stage 4, provided that the sampling site was not in a city park. The influence of the forest over the city is an interesting question, and I suggest that you include a sentence about it in the abstract.

Response: It was stated in the original manuscript that a part of SOA is from local sources. However, the SOA and AS contents of samples SM1 and SM2 are clearly higher than that of sample SM3, so that the reasoning can be defensible. A sentence suggested by the reviewer was added in the abstract – “Based on the different contents of SOA, ammonium sulfate, and EC particles among the samples collected in Manaus, a considerable influence of the rainforest over the city was observed.” (lines: 44-46).

18) In general, I recommend that you improve the discussion on Manaus results. It is a metropolis in the middle of a tropical forest, a very unique situation. How does your findings compare to other metropolis in the world? What is the proportion of SOA, PBA and soot in other cities?

Response: I agree with the reviewer in that Manaus is a very unique place. However, it is not feasible to compare the current results with those in other metropolis in the world. We could not find literature data of SOA, PBA, and soot in other cities collected in the similar way as this

work. I would like to mention that this single-particle characterization of aerosols has some difficulty in comparing with the results from bulk analyses.

19) Lines 502-503: Was there anything special during the sampling SM3? Did you notice increased traffic activity compared to the other days?

Response: Indeed, only sample SM3 was collected during a regular working day, which is given in the revised text – “Samples SM1 and SM2 were collected during and just after a national holiday, respectively, when all the institutions (private and public) were closed during that day so that the traffics were quite low, similar to a weekend or vacation period. Sample SM3 was collected during a regular working day, so that sample SM3 is the only sample actually exposed to the high traffic of light vehicles in the area.” (lines: 508-512).

20) Line 650: Please update the reference, since it moved from ACPD to ACP.

Response: Done as suggested.

21) Figure 9: I suggest that you group the columns by size range, for a better comprehension of the plots. Also, you might include information on particle size ranges, either in the x-axis label, or in the figure caption.

Response: Done as suggested.

General Comments from Anonymous Referee #2

The manuscript provides information of individual particles in the Amazon basin using a combination of microscopy and infrared techniques, which is an unprecedented approach to the problem. This is the main achievement of the study from the analytic viewpoint. With respect to the location, particles are from the pristine ATTO Tower and urban pollution from the big city of Manaus. If ATTO has been extensively reported in the scientific literature, it is not the case for the city of Manaus, so the manuscript comes to contribute to the knowledge with respect to the properties of aerosols from Manaus.

Response: We thank the reviewer for the positive evaluation of our work.

Specific comments from Anonymous Referee #2

This referee does agree to all comments posted by the other reviewer with respect to explore better the Manaus database. I would add to this the poor discussion with respect to the weak point of the manuscript is with respect to a better of the atmospheric condition as a whole. For example, the characterization of the meteorology (lines 138 - 144) was extremely poor. Nothing was written on the synoptic situation. Hysplit is useful but it does not consider wet removal adequately thus a 10 day back trajectory is far from enough to provide a good information about meteorology. To say that RH was above 55% in the Amazon is useless, it is almost all the time from January - May above it. So, were the measured days ordinary? Anything different?

Response: Detailed meteorological conditions were added in Table 1. Based on the meteorological parameters during the samplings, the sampling days do not seem abnormal.

With respect to the discussion about the emissions from Manaus, only 9 lines (100 - 108) were written about it with very few information. So, the city is industrial but the author did not mention that these industries are not great emitters, they are basically assembling industries. The main source of pollutants is the light vehicles fleet. Poor information was also provided with respect to the sampling site in Manaus. Also there were a mention about downwind transport of pollution from Manaus that confuses the reader (lines 104-108) because ATTO is upwind and barely get any influence from the city. The focus of this is far from any downwind issues.

Response: The reviewer is correct as the industries in the region do not emit a large amount of pollutants, whereas power plants, refinery, and vehicle fleet are mainly responsible for the atmospheric emissions and the vehicle fleet is the main source. Light vehicles powered by gasoline, ethanol, or both account to the majority of the transportation fleet in the city. The information was added to the text (p. x; lines 126-132).

As suggested by the reviewer, the confusing downwind-upwind part was deleted – “Based on an investigation on particulate matter during the wet season, oxidized organic components were significantly observed at Manaus sites (de Sá et al., 2018)” (lines 105-108).

Comments to the text from Anonymous Referee #2

The text is very well written. I would add the following corrections:

- 1. tar ball > tarball (several locations in the text).*
- 2. Figures 6, 8, 10, 11: fix the weird position of the axis label "keV".*
- 3. Figure 9: put dates instead of sample label in X-axis.*

4. *Figure 11 (caption): tar call > tarball.*

5. *Figure 13 (caption): "Relative abundance" to "Relative MASS abundance"*

Response: Done as suggested, except (i) comment #3, for which Fig. 9 was modified as suggested by Reviewer #1 and (ii) comment #5, for which *"Relative abundance" was replaced by "Relative number abundance"*.

Single-particle characterization of aerosols collected at a remote site in the Amazonian rainforest and an urban site in Manaus, Brazil

Li Wu¹, Xue Li¹, HyeKyeong Kim¹, Hong Geng², Ricardo H. M. Godoi³, Cybelli G. G. Barbosa³, Ana F. L. Godoi³, Carlos I. Yamamoto⁴, Rodrigo A. F. de Souza⁵, Christopher Pöhlker⁶, Meinrat O. Andreae^{6,7}, and Chul-Un Ro^{1,*}

¹ Department of Chemistry, Inha University, Incheon, 402-751, Korea;

² Institute of Environmental Science, Shanxi University, Taiyuan 030006, China;

³ Environmental Engineering Department, Federal University of Parana-UFPR, Curitiba, PR, Brazil;

⁴ Chemical Engineering Department, Federal University of Paraná-UFPR, Curitiba, PR, Brazil;

⁵ Amazonas State University, Superior School of Technology, Manaus, Amazonas, Brazil;

⁶ Multiphase Chemistry & Biogeochemistry Departments, Max Planck Institute for Chemistry, 55020 Mainz, Germany;

⁷ Department of Geology and Geophysics, King Saud University, Riyadh, Saudi Arabia.

Abstract

In this study, aerosol samples collected at a remote site in the Amazonian rainforest and an urban site in Manaus, Brazil, were investigated on a single particle basis using a quantitative energy-dispersive electron probe X-ray microanalysis (ED-EPMA). Twenty-three aerosol samples were collected in four size ranges (0.25-0.5, 0.5-1.0, 1.0-2.0, and 2.0-4.0 μm) during the wet season in 2012 at two Amazon basin sites: 10 samples in Manaus, an urban area; and 13 samples at an 80-m high tower, located at the Amazon Tall Tower Observatory (ATTO) site in the middle of the rainforest, 150 km northeast of Manaus. The aerosol particles were classified into nine particle types based on the morphology on the secondary electron images (SEIs) together with the

* Corresponding author. Tel.: +82 32 860 7676; fax: +82 32 874 9207

E-mail address: curo@inha.ac.kr (C.-U. Ro)

elemental concentrations of 3,162 individual particles: (i) secondary organic aerosols (SOA), (ii) ammonium sulfate (AS), (iii) SOA and AS mixtures, (iv) aged mineral dust, (v) reacted sea-salts, (vi) primary biological aerosol (PBA), (vii) carbon-rich or elemental carbon (EC) particles, such as soot, tarball, and char, (viii) fly ash, and (ix) heavy metal (HM, such as Fe, Zn, Ni, and Ti)-containing particles. In submicron aerosols collected at the ATTO site, SOA and AS mixture particles were predominant (50-94% in relative abundance) with SOA and ammonium sulfate comprising 73-100%. In supermicron aerosols at the ATTO site, aged mineral dust and sea-salts (37-70%) as well as SOA and ammonium sulfate (28-58%) were abundant. PBAs were observed abundantly in the PM₂₋₄ fraction (46%), and EC and fly ash particles were absent in all size fractions. The analysis of a bulk PM_{0.25-0.5} aerosol sample from the ATTO site using Raman microspectrometry and attenuated total reflection Fourier transform infrared spectroscopy showed that ammonium sulfate, organics, and minerals are the major chemical species, which is consistent with the ED-EPMA results. In the submicron aerosols collected in Manaus, either SOA and ammonium sulfate (17-80%) or EC particles (6-78%) were dominant depending on the samples. In contrast, aged mineral dust, reacted sea-salt, PBA, SOA, ammonium sulfate, and EC particles comprised most of the supermicron aerosols collected in Manaus. The SOA, ammonium sulfate, and PBAs were mostly of a biogenic origin from the rainforest, whereas the EC and HM-containing particles were of an anthropogenic origin. Based on the different contents of SOA, ammonium sulfate, and EC particles among the samples collected in Manaus, a considerable influence of the rainforest over the city was observed. Aged mineral dust and reacted sea-salt particles, including mineral dust mixed with sea-salts probably during long-range transatlantic transport, were abundant in the supermicron fractions at both sites. Among the aged mineral dust and reacted sea-salt particles, sulfate-containing ones outnumbered those containing nitrates and sulfate+nitrate in the ATTO samples. In contrast, particles containing sulfate+nitrate were comparable in number to particles containing sulfate only in the Manaus samples, indicating the different sources and formation mechanisms of secondary aerosols, i.e., the predominant presence of sulfate at the ATTO site from mostly biogenic emissions and the elevated influences of nitrates from anthropogenic activities at the Manaus site.

1. Introduction

The Amazonian rainforest is regarded as one of the primitive continental regions and

atmospheric aerosol particles over the region are expected to be influenced minimally by anthropogenic activities, particularly during the wet season (Andreae, 2007; Martin et al., 2010b; Chen et al., 2015). The unique near-natural conditions during the wet season make it an ideal place to understand the occurrence, nature, origin, and transport of aerosol particles, which can directly scatter and absorb solar radiation and indirectly serve as cloud condensation nuclei (CCN) and/or ice nuclei (IN), to better predict the additional anthropogenic effects on aerosol particles, and to help determine their influences on the environment, climate, and human health (Artaxo et al., 2013). The Amazon Tall Tower Observatory (ATTO) consists of several observatory towers built in the middle of the Amazon rainforest for a continuous and detailed study of biota-atmosphere interactions (Andreae et al., 2015). Manaus, the capital of Amazonas state, is a large city located in the northern region of Brazil with more than 2 million inhabitants in an area of 11,401 km² (IBGE, 2017). The city, which is surrounded by the largest tropical rainforest, has a large industrial zone, a port area at the Rio Negro, an energy matrix based on fuel oil, diesel, and natural gas, and a growing automotive fleet (Martin et al., 2010a, 2016). Consequently, the pollution plume from Manaus can act as a laboratory for examining the perturbations in natural processes (Martin et al., 2016). Only a few studies examined airborne particles over the Amazon rainforest and nearby urban sites simultaneously (Fraund et al., 2017; Martin et al., 2016). Therefore, there is still little information on the urban vs. ecosystem influences. Especially, single-particle characterization of aerosols collected at a Manaus city center has been scarce.

Many studies have been performed on the aerosol characteristics in the Amazon basin, but the formation and dynamic processes of Amazonian aerosols are not completely understood (Andreae et al., 2015; Martin et al., 2010a, 2016; Fraund et al., 2017; Fan et al., 2018). The Amazon Basin atmosphere is near-pristine during the wet season, whereas biomass burning prevails during the dry season (Andreae et al., 2007; Pöschl et al., 2010; Artaxo et al., 2013; Pöhlker et al., 2018). Based on a long-term study, it was reported that aerosol particles in the coarse fraction are relatively constant in concentrations through the wet and dry seasons, whereas the aerosol particle levels in the fine fraction differ due to the predominant influence of biomass burning during the dry season (Artaxo et al., 2013; Moran-Zuloaga et al., 2018). Scanning electron microscopy/energy dispersive X-ray spectrometry (SEM/EDX) studies categorized the Amazonian aerosols mainly as secondary organic aerosol (SOA) particles, sulfates/chlorides, primary biological aerosol (PBA) particles, mineral dust, sea salts (fresh and/or aged), and pyrogenic carbon particles within the

different size fractions (Krejci et al., 2005; Pöschl et al., 2010). Over the Amazonian rainforest, SOA particles are **mainly** formed through the condensation of biogenic organic compounds onto biogenic K-rich salt particles emitted from the forest and are predominant in the fine fraction, which are important for CCN (Pöhlker et al., 2012) and can also affect the potential of mineral particles when acting as an organic coating (Möhler et al., 2008). Under high relative humidity conditions, nano- and micrometer SOA particles with a dominance of α -pinene and isoprene as their precursors can remain in the liquid phase (Bateman et al., 2016), which further enhances the formation of SOA as well as the oxygen-to-carbon (O/C) ratios. Hence, the study of this particle type can help elucidate some of the atmospheric interactions (Lin et al., 2014). The atmosphere in the Amazon Basin is also rich in PBA particles (Andreae, 2007; Artaxo et al., 1998, 2013; Martin et al., 2010a). Their unique morphology and elemental compositions of major C and O with minor S, K, P, Na, N, Cl, and/or Mg obtained by SEM/EDX are characteristic of individual PBAs like fungal spores (China et al., 2016). PBA particles can contribute to CCN after being transported to cloud formation altitudes by strong convection (Artaxo et al., 2013). At the ATTO site, single particle analysis by a combination of scanning transmission X-ray microscopy/near edge X-ray absorption fine structure spectroscopy (STXM/NEXAFS) and SEM/EDX highlighted the dominance of biological **particles and the abundance of biogenic SOA and the presence of C, N, O, P and K are characteristic of aerosols at the area (Fraund et al., 2017). Based on an investigation on particulate matter during the wet season, oxidized organic components were significantly observed at Manaus sites (de Sá et al., 2018), where one third of the potential SOA would be** of an urban origin (Palm et al., 2018).

In this study, twenty-three aerosol samples collected at the ATTO site and at an urban site in Manaus during the wet season in 2012 were examined on a single particle basis using a quantitative energy-dispersive electron probe X-ray microanalysis (ED-EPMA), which provided information on the morphology and chemical compositions of aerosols containing both light and heavy elements. This paper presents the different characteristics of the aerosols collected at the rainforest and in Manaus.

2. Experimental section

2.1. Samples

During the wet season in 2012, aerosol samples were collected at two sampling sites on the

Amazon basin, i.e., ATTO and a central area of Manaus. The ATTO site (S 02° 08.647' W 58° 59.992') is situated in the Uatumã Sustainable Development Reserve, approximately 150 km northeast of Manaus (Fig. 1). This is a multidisciplinary research site of an international joint project between Brazil and Germany for continuous monitoring of the biological, physical, and chemical functions of the Amazon rainforest to answer questions related to climate change (Andreae et al., 2015). Aerosol sampling was performed at an 80-m-height walk-up tower at the ATTO site. In Manaus, the sampling site is situated in the central part of the city (S 03° 05.753', W 59° 59.419'), which is at a representative urban region influenced by electricity production based on fuel oil, diesel, and natural gas, biogenic emissions from the surrounding forest, and mostly by light duty (using gasoline and ethanol) vehicle traffic. Heavy vehicles that use diesel account to less than 10 % of the urban fleet (Medeiros et al., 2017). The location is nearby a small parking area and around 200 m away from the intersection of four busy avenues, with frequent diurnal traffic jam on weekdays. Particles were collected at a 2 m height above ground level. The aerosol samples were collected on TEM grids (Ted Pella Inc., USA, Ted Pella Inc., Carbon/Formvar 200 mesh Cu grid, 35–70 nm thickness) using a five stage Battelle impactor (the cut-off diameters are 0.25, 0.5, 1, 2, and 4 μm for stages 1-5, respectively) at the ATTO and Manaus sites on April 1, 16, 17, and 18 and May 1, 2, and 3 (the four- and three-day samples were collected at the ATTO and Manaus sites, which are notated as samples SA1-SA4 and SM1-SM3, respectively). On each date, the sampling started around noon (local time) and lasted for approximately 100 min. The individual particles collected on stages 1-3 ($\text{PM}_{0.25-0.5}$, $\text{PM}_{0.5-1.0}$, and $\text{PM}_{1.0-2.0}$) for each sample and on an additional stage 4 ($\text{PM}_{2.0-4.0}$) for the SA4 and SM3 samples were examined.

During the sampling period, the temperature was in the range of 22 to 32°C and the relative humidity was above 55%. On April 16 and May 1 and 2, rain events occurred within the previous 24 hours prior to sampling. Detailed information on sampling dates and times and the meteorological conditions during the samplings are given in Table 1. The ten-day backward air-mass trajectories were obtained using the Hybrid Lagrangian Single-Particle Integrated Trajectory (HYSPLIT) model from the NOAA Air Resources Laboratory's web server (<http://www.arl.noaa.gov/ready/hysplit4.html>), as shown in Fig. 2. In the HYSPLIT calculation, meteorological data output from the Global Data Assimilation System (GDAS) using GDAS1 data with a horizontal resolution of 1° corresponding to ~ 100 km x 100 km and 23 vertical layers was

used, which was reported to provide plausible backward trajectory analysis (Su et al., 2015). All samples were influenced by transatlantic air masses at a 1000 m receptor height and the Manaus site was influenced mainly by the surrounding rainforest at 500 m and 100 m heights.

2.2. EPMA measurements and data analysis

Low-Z particle EPMA was carried out by SEM (JSM-6390, JEOL) equipped with an Oxford Link SATW ultrathin window EDX detector, which has a spectral resolution of 133 eV for Mn K_{α} X-rays. The X-ray spectra were recorded using INCA Energy software. To achieve the optimal experimental conditions, such as the low background level in the X-ray spectra and good sensitivity for low-Z element analysis, an accelerating voltage of 10 kV, a beam current of 0.5 nA, and a measuring time of 20 s were used. X-ray spectral data acquisition for individual particles was carried out manually in point analysis mode, i.e., the electron beam was focused at the center of each particle, and X-rays were acquired while the beam remained fixed on this single spot. The secondary electron images (SEIs) and X-ray spectra of an overall 3,162 individual particles for the ATTO and Manaus samples were examined. As the TEM grids are thin (35-70 nm thickness), strong X-rays from the Al or Cu metal stub commonly used in the SEM/EDX measurement would be a problem when the TEM grid substrate is placed on it. A home-made sample holder (Fig. 3(a)) for the TEM grid samples was used to avoid interference from the metal stub, resulting in X-ray spectra of bare TEM grids, which showed only C and O X-ray peaks from their carbon/Formvar thin-film, a Cu-L peak caused by lateral scattering from the Cu bars of TEM grids, and a Si peak from an impurity, as shown in Fig. 3(b). The net X-ray intensities for the chemical elements were obtained by non-linear, least-square fitting of the spectra using the AXIL program (Vekemans et al., 1994). Although the characteristic X-ray intensities of C and O were low for the bare TEM grids, determination of the C and O concentrations for individual particles on the TEM grids was performed using a methodology based on the Monte Carlo calculation technique to correct for the interfering X-ray peaks of C and O emitted from the TEM grid, which provided reliable quantification results when applied to the quantification of standard (sub)micron particles, such as CaCO_3 , CaSO_4 , Na_2SO_4 , and SiO_2 . On the other hand, electron beam-sensitive particles, such as NaNO_3 , $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$, and ammonium sulfate, provided deviating quantification results (Geng et al., 2010). As the Cu-L and Si X-ray intensities from the bare TEM grids are quite small (< 20 cps) under these measurement conditions, the two peaks could be neglected safely during the

quantification procedure.

3. Results and discussion

3.1. Particle types observed in samples collected at the ATTO and Manaus sites

In this study, the analyzed particles were classified based on their X-ray spectral and SEI data, where 9 different particle types were observed in the samples collected at the ATTO and Manaus sites during the wet season in 2012; i.e., (i) SOA, (ii) ammonium sulfate (AS) particles, (iii) SOA and AS mixture particles, (iv) aged (reacted) sea-salt, (v) aged mineral dust, (vi) PBA particles, (vii) carbon-rich or elemental carbon (EC) particles such as soot, tarball, and char or coal dust, (viii) fly ash particles, and (ix) heavy metal-containing (HM) particles. In a previous study (Pöschl, et al., 2010), five types of aerosols, such as (i) SOA droplets, (ii) SOA-inorganic mixture particles where the inorganics are mostly sulfates and/or chlorides, (iii) PBA, (iv) mineral dust, and (v) pyrogenic carbon particles, were reported based on single particle analysis for aerosol samples collected at a remote site north of Manaus, Brazil during the 2008 wet season (3-13 March) using SEM/EDX. The pure SOA droplets dominated in the nucleation and Aitken modes, whereas the pure SOA, SOA-inorganic mixture particles, and pyrogenic carbon particles dominated in accumulation mode (Pöschl, et al., 2010). With the exception of the reacted sea-salt particles probably from the Atlantic Ocean as well as the abundant ammonium sulfate aerosols, the particle types observed in this study are comparable to their study. Figs. 4 and 5 present typical field SEIs for submicron and supermicron aerosol particles collected at the ATTO and Manaus sites, respectively, where the chemical species comprising each particle is indicated. Ammonium sulfate and SOA particles are dominant in the sub- and super-micron aerosol fractions collected at the ATTO site with some mineral particles and aged sea-salts in the supermicron fractions, whereas the aerosol samples collected at the Manaus site are composed of various types of particles of anthropogenic and/or natural origin. Figs. S1-S7 of Supporting Information present typical SEIs for all the samples with identified chemical species on the SEIs, which helps briefly illustrate the different features of the samples collected at the ATTO and Manaus sites. The characteristics of the particle types observed in the ATTO and Manaus samples are described in the following.

3.1.1. Secondary organic aerosol (SOA) particles

In this study, SOA particles were observed frequently in both the ATTO and Manaus

samples, even though pure SOA particles were rare and most of them were mixed internally with other species, such as ammonium sulfate, K-rich salt, reacted sea-salts, etc. The SOA particles over the Amazon rainforest are formed by the oxidation of biogenic volatile organic compounds (Jimenez et al., 2009; Hallquist et al., 2009; Martin et al., 2010a; Andreae et al., 2018) and are the major constituents of particulate matter (PM), particularly for submicron ambient PM (Pöschl, et al., 2010; Martin et al., 2010b; Chen et al., 2015). In the SEI images, pure SOA droplet particles appear gray in contrast and have a circular shape, as shown in Fig. 6(a). As TEM grid films (with 90% C content) and SOAs are composed mainly of carbon and oxygen, the SOA aerosols appear gray on the TEM grids because of their similar secondary and backscattered electron yields to those of the TEM grid (Goldstein et al., 2003; Maskey et al., 2010). As TEM grids are hydrophobic due to the thin carbon layer over the Formvar film, the aqueous droplet aerosols appear circular on the TEM grids (Eom et al., 2014; Maskey et al., 2010), suggesting that SOAs were collected as aqueous droplets at the time of particle sample collection. Recent studies also reported that most submicron SOA particles in the Amazon basin are water soluble organic aerosols (WSOAs) rather than semi-solid or solid aerosols under the background conditions that are typically met during the wet season (Bateman et al., 2016, 2017). The X-ray spectrum of a typical pure SOA, as shown in Fig. 6(a), showed considerably higher levels of the C X-ray peak intensity compared to that from the Formvar/carbon film of the TEM grids, resulting in the unambiguous identification of SOA particles based on their SEIs and X-ray spectral data.

3.1.2. Ammonium sulfate (AS) particles

Ammonium sulfate particles were observed abundantly in the ATTO and Manaus samples, mostly as mixtures with secondary organics. Ammonium sulfate particles appear bright and crystalline on the SEIs before the X-ray measurements, both for pure airborne and standard ammonium sulfate particles, as shown in Figs. 6(b) and 6(f), respectively. The standard ammonium sulfate particles were deposited on TEM grids by the nebulization of a 0.1 M ammonium sulfate solution. As shown in the inset in Figs. 6(b) and 6(f), after the X-ray measurements, they show somewhat darkened SEIs with black holes, due to electron beam damage, at the places where the electron beam hits. As the ammonium sulfate particles are electron beam-sensitive (Geng et al., 2010; Worobiec et al., 2003; Huang and Turpin, 1996), their X-ray spectral signature is the presence of a significant S X-ray peak, as shown for both pure airborne and standard particles on

the TEM grids. The N X-ray peak was often not detected, particularly for small particles because the NH_4^+ moiety is especially prone to damage by electron beams.

Ambient urban and rural sulfates act as a sink for ammonia, of which the sources are largely animal waste, fertilizer application, soil release, and industrial emissions. The most common form is ammonium sulfate. On the other hand, if ammonia is scarce in the air, sulfates would be in more acidic forms, such as NH_4HSO_4 or H_2SO_4 (Millstein et al., 2008). The acidic $\text{NH}_4\text{HSO}_4/\text{H}_2\text{SO}_4$ particles have been reported to be more hygroscopic than pure ammonium sulfate (Pósfai et al., 1998). Hence, they can be spread over the collecting substrate (Formvar/carbon film). In addition, acidic sulfate particles can have unique halo rings in their morphology (Buseck and Pósfai, 1999). The crystalline structure of the ammonium sulfate-containing particles observed in this study suggests that they are sulfates fully neutralized with ammonia. In addition, the Raman spectra of airborne particles exhibiting this morphology were obtained on a single particle basis to confirm that they are ammonium sulfate. As shown in Fig. 7(a), the Raman peak at 975 cm^{-1} of the airborne particle is characteristic of ammonium sulfate (Ling and Chan, 2007), which was also confirmed by Raman spectroscopy on standard ammonium sulfate particles. Characteristic Raman peaks for NH_4HSO_4 , K_2SO_4 , $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$, CaSO_4 , Na_2SO_4 , and $\text{MgSO}_4 \cdot x\text{H}_2\text{O}$ ($x = 1-7, 11$) were reported to be at 1010 and 1042, 983, 1008, 1014 and 1025, 992, and 984-1046 cm^{-1} , respectively (Fung and Tang, 1988; Wang et al., 2006; Mabrouk et al., 2013; Prieto-Taboada et al., 2014). The sloping baseline in the Raman spectrum of the airborne particle was attributed to the fluorescence from organics, indicating the presence of organic compounds (Sobanska et al., 2012), probably from SOA. For aerosols collected on Ag foil at the ATTO site on June 10, 2014, ammonium sulfate is the major species with some organics and minerals such as kaolinite, for the bulk aerosols in the size range of 0.25 – 0.5 μm , based on their X-ray, attenuated total reflectance-FTIR (ATR-FTIR), and Raman spectra (Fig. 7(b)). A study of the samples collected in the central Amazon Basin during the wet season from February to March 2008 reported that ammonium was not sufficient to fully neutralize sulfates so that ammonium bisulfate would be present in the Amazon rainforest (Chen et al., 2015), whereas other studies reported that sulfates are sufficiently neutralized with ammonia in the fine and coarse fractions during both the wet and dry seasons (Andreae et al., 2015; Martin et al., 2010b; Fuzzi et al., 2007; Mace et al., 2003). The different results may be due to different sampling places and seasons. In this study, ammonium sulfate is dominant over ammonium bisulfate.

Previous studies have shown that the sulfate aerosols over the Amazon forest are predominantly from marine and terrestrial biogenic sources, with comparable contributions from marine and terrestrial biogenic emissions (Andreae et al., 1990). Sulfate originates from biogenic sources in the rainforest, i.e., dimethyl sulfide (DMS), H_2S , and CS_2 emitted by plants and microorganisms, which can be oxidized to sulfate (Andreae et al., 1990, 2015; Martin et al., 2010b). The rainforest ecosystem in the central Amazon can act as a source of DMS to the atmosphere throughout the year (Jardine et al., 2015). Several studies have reported that marine DMS transported from the Atlantic Ocean contributes significantly to the sulfate levels in the Amazon basin (Gregory et al., 1986; Andreae et al., 1990; Martin et al., 2010a). In addition, there is some sulfate from long-range transport across the Atlantic including that from the African volcanic emissions (Saturno et al., 2018) and minor upwind anthropogenic sources.

The nitrogen cycle is essential for organisms and some bacteria to fix the gaseous N_2 in the air to NH_4^+ for their own biosynthetic processes (Kim and Rees, 1994; Bazzaz, 1998; Kellerhals et al., 2010). In addition, some microorganisms produce enzymes to release nitrogen as NH_4^+ during the nitrogen mineralization process, which is important in tropical rain forest soils, where dead plants and animal matter accumulate continuously (Wright, 1996; Neill et al., 1999). A high level of NH_4^+ in tropical rain contributes significantly to the nitrogen influx in the rainforest soils (Jordan et al., 1982). The NH_4^+ species can be evaporated as gaseous NH_3 from surface soils, particularly from leaf litter, resulting in a strong NH_3 emission source as well as stomatal NH_3 emission of plants as another natural source in forest ecosystems (Sutton et al., 2009, 2013; Hansen et al., 2017). On the other hand, NH_4^+ species in rainforest soils might become airborne immediately after rainfall, similar to the way that airborne organic particles are produced directly from soils by raindrop impaction (Wang et al., 2016). The wetness of forest surfaces is significant in controlling both the deposition and emission of atmospheric NH_3 (Hansen et al., 2015). As ammonium sulfate-containing particles were also observed abundantly in the samples collected at Manaus site, they were influenced strongly by the surrounding Amazonian forest and/or generated by anthropogenic activities in the urban environment. In the urban environment, anthropogenic ammonium sulfate is mainly formed by gaseous reactions among SO_2 emitted from coal-fired plants and industrial activities, NH_3 emitted from human and animal activities and fertilization in the fields, and oxidants (e.g., O_3 and OH radical) (Li et al., 2016; Geng et al., 2017).

3.1.3. SOA and AS mixture particles

In this study, most airborne submicron SOAs were observed to be internally mixed with ammonium sulfate, particularly for the samples collected at the ATTO site. Figure 6(c) shows that a typical SOA and AS mixture particle has the crystalline, bright ammonium sulfate moiety in the center surrounded by circular, grey organic species. The circular morphology of the organic species strongly suggests that the organic species are SOAs, as stated above. As efflorescence and deliquescence relative humidity (ERH and DRH) of ammonium sulfate species are 30-40% and 80%, respectively (Yeung and Chan, 2010) and the ambient RH was always above 55% during sampling for the ATTO and Manaus samples, the ammonium sulfate would be mostly in aqueous droplets at the time of sample collection, rather than in crystalline form, as indicated by their overall circular shape. When the particle samples were under dry conditions either during sample storage or in the vacuum chamber of the SEM instrument, the ammonium sulfate species crystallized, resulting in core-shell structures of organic and inorganic mixture aerosols. When the ambient RH is low enough to make the ammonium sulfate species crystallize in the atmosphere, the organic and inorganic mixture aerosols would similarly be present as core-shell structures.

Some of the SOA and AS mixture particles were also mixed with K-salts. As shown in Figs. 6(d) and 6(e), their morphology was similar to that of the SOA and AS mixture particles, but their X-ray spectra revealed the presence of K and an elevated S level compared to those of the SOA and AS mixture particles, suggesting that the K is associated mostly with SO_4^{2-} . The shoulder Raman peak of the airborne ammonium sulfate particle at 982 cm^{-1} (Fig. 7(a)), which is indicative of the K_2SO_4 moiety (Mabrouk et al., 2013), also suggests that the K-salts are most probably K_2SO_4 . A previous study reported that small K-salt-rich particles can act as seeds for SOA formation in the Amazon basin and K-salts are present ubiquitously in Amazonian SOAs with their content being higher in the morning hours and for smaller SOAs (Pöhlker et al., 2012). On the other hand, among the 843 submicron SOA and/or ammonium sulfate particles collected at the ATTO site, only 31% contained K-salts, which is probably because the samples were collected in the afternoon and/or the analyzed particles were larger than $0.25\text{ }\mu\text{m}$ so that the K-salt content may be below the detection limit of EDX ($\sim 0.1\text{ wt. }\%$). In the Manaus samples, a total of 199 submicron SOA and/or ammonium sulfate particles were observed, of which approximately 40% contained K-salts, suggesting that the Manaus samples were influenced strongly by the surrounding rainforest as supported by the backward trajectories (Figs. 2(e)-(g)), where K-salts may be mostly of a biogenic

origin in the rainforest. The organic moiety is often mixed internally with aged sea-salts, mineral dust, and PBAs, which will be described below.

3.1.4. Mineral dust particles

The typical mineral dust particles include aluminosilicate, quartz (SiO_2), calcite (CaCO_3), dolomite ($\text{CaMg}(\text{CO}_3)_2$), and TiO_2 (Geng et al., 2009, 2011). They appear irregular and bright on the SEIs (Fig. 8). Various types of mineral particles from Saharan dust contribute significantly to the nutrient cycles in the Amazon rainforest (Talbot et al., 1990; Abouchami et al., 2013; Rizzolo et al., 2017). Mineral dust tends to provide reactive surfaces for heterogeneous reactions with trace atmospheric gases, such as SO_2 and NO_x , leading to chemical modifications of the particles that ultimately affect the atmospheric chemical balance and photochemical cycle (Sullivan et al., 2007; Chen et al., 2011). Modification of the physicochemical properties of particles can alter their optical, chemical, and hygroscopic properties (Sullivan et al., 2007; Geng et al., 2014). If some components in them (particularly Ca-containing species) react with airborne SO_2 and NO_x in the presence of moisture or with “secondary acids”, such as H_2SO_4 , HNO_3 , and HCl , they are regarded as reacted or aged ones. The reacted/aged ones can contain either nitrates, sulfates, or both (Geng et al., 2014, 2017; Li et al., 2016). In the Amazon samples, almost all the mineral dust particles were aged ones, as shown in Fig. 8, where an aluminosilicate and a carbonate/silica mixture particle containing sulfate are shown. The S-containing aged mineral dust particles outnumbered the N- and both N- and S-containing ones for the ATTO samples, whereas they were comparable to both the N- and S-containing ones for the Manaus samples, as shown in Fig. 9(a), indicating the predominance of sulfates over nitrates for the reaction of mineral dust particles at the ATTO site and somewhat significant influence of nitrates at the Manaus site. The nitrates for the Manaus samples may be formed from nitrogen oxides emitted from the vehicles and coal-fired power plants (Geng et al., 2014, 2017; Li et al., 2016). The mineral particles might be mixed or covered with SOA and/or ammonium sulfate and gradually become aged under a high RH in the rainforest. Some mineral particles were also mixed with sea salt particles during their transport to the Amazonian area across the Atlantic Ocean.

3.1.5. Reacted (aged) sea-salts

The fresh sea-salt particles can react with gas-phase sulfur and nitrogen oxides to contain

sulfate and nitrate, respectively (ten Brink, 1998). During the process, chlorine may be removed completely if the reaction is complete (Laskin et al., 2003). All sea-salt particles observed in the ATTO and Manaus samples were reacted ones. Figure 10(a) shows the X-ray spectrum, atomic concentration, and SEI of a typical aged sea salt particle, where the presence of Na, Mg, and Cl indicates its marine origin (Geng et al., 2014) and the presence of S and C indicates that it is mixed with sulfates and organics. The irregular and somewhat bright SEI is typical of the aged sea-salt particles. As shown in Fig. 9(b), S-containing sea-salts outnumbered N- and both N- and S-containing ones for the ATTO samples, indicating the predominance of sulfates over nitrates for the reaction of sea-salt particles. The S-containing particles are comparable in abundance to both the N- and S-containing ones at the Manaus site (Fig. 9(b)). The sea-salt particles may also become mixed with ammonium sulfate over the rainforest and become S-containing ones. Among overall 275 reacted sea-salts containing sulfate/nitrate and organics, 71% of them were mixed with K-salts, as shown in Fig. 10(b). The presence of K-salts in the reacted sea-salt particles indicates that the mixing of the K-salts of biogenic origin would happen in the rainforest during long-range transport because of the minimal biomass burning influence during the wet season, although it was reported that a strong biomass burning smoke was transported from Africa to South America during the wet season (Baars et al., 2011) so that a part of the observed K-salts could be associated with biomass burning aerosols from Africa. In addition, several elongated CaSO_4 particles, as shown in Fig. 10(c), were detected in both ATTO and Manaus samples, all of which contain a small amount of Na. Their elongated shape and the presence of Na strongly indicates that they were from the sea, possibly the Atlantic Ocean, not from the soil (Eom et al., 2016).

3.1.6. Primary biological aerosol (PBA) particles

PBA particles like fungal spores can be identified easily based on their unique morphology and the presence of their characteristic chemical elements (Geng et al., 2009; Martin et al., 2010a; Pöschl et al., 2010). The PBA particle shown in Fig. 11(a) has a unique oval morphology and the majority of C together with the characteristic small amounts of N, P, S, Cl, and K. PBA particles are relatively large (diameter $> 2.0 \mu\text{m}$) so that they are abundant in stage 4 samples ($2.0 \mu\text{m} < \text{diameter} < 4.0 \mu\text{m}$), particularly at the ATTO site. Figure 12 shows two image fields of stage 4 samples collected at the ATTO and Manaus sites, where the PBA particles have various types of morphology and many of them are mixed with SOA. The abundant observation of PBA particles

in the stage 4 sample of the Manaus site suggests the transport of the PBA particles from the rainforest to the urban area. Supermicron PBA particles were reported to be abundant over the Amazon (China et al., 2016; Moran-Zuloaga et al., 2017; Gilardoni et al., 2011). PBA particles can be pollen, bacteria, fungal and fern spores, viruses, and fragments of plants and animals emitted directly from the rainforest, showing a range of morphologies, and comprise the largest fraction of the coarse mass (Martin et al., 2010a). PBA particles appear to be the most efficient and abundant ice nuclei (Pöschl et al., 2010; Tobo et al., 2013; Haga et al., 2014). In addition, the release of nano- and submicron particles from fungal spores under high relative humidity can contribute to new particle formation and potentially affect cloud formation in the Amazon Basin (China et al., 2016).

3.1.7. Carbon-rich particles from combustion sources

Carbon-rich particles, such as soot, tarballs, and char or coal dust, which contain more than 90% C and O with the C content being dominant over O in low-Z particle EPMA analysis (Geng et al., 2009, 2010, 2014), were observed frequently in the Manaus samples, whereas they were rare in the ATTO samples. Based on the characteristic morphology of carbon-rich particles, soot aggregates of fractal-like chain structures (Fig. 11(b)), tarballs of separate spherules (Fig. 11(c)), and chars of irregular-shaped carbon (Fig. S5, SM1-2) could be differentiated straightforwardly from each other (Geng et al., 2010, 2014; Li et al., 2016). The soot aggregates formed via a vaporization-condensation mechanism during the combustion processes vary in size from sub to several micrometers (Chen et al., 2005, 2006). Once airborne, the complex microstructure of the soot aggregates may provide active sites for the deposition of organics and other chemical species, such as sulfates (Pósfai et al., 1999; Zhang et al., 2008), as revealed by the presence of S in Fig. 11(b). This results in aged soot aggregates that become compact with considerable restructuring and shrinkage (Zhang et al., 2008). Tarballs, which are a type of brown carbon (Andreae and Gelencsér, 2006; Laskin et al., 2015), usually have high C, N, and O contents with a spherical morphology (Fig. 11(c)), strongly indicating their formation during biomass combustion processes (Pósfai et al., 2003, 2004). Char appears compact and irregular in the SEIs; it is often mixed with minor inorganic species, such as K and S, and is regarded as the incomplete combustion remnants of liquid or solid carbonaceous fuel materials that have undergone carbonization during combustion (Chen et al., 2006). Only one soot particle was observed in all the ATTO samples,

whereas soot, tarball, and char or coal dust particles were abundant in the submicron Manaus samples, suggesting that the ATTO samples are barely affected by the anthropogenic carbon-rich particles generated in Manaus.

3.1.8. Fly ash particles

As shown in Fig. 11(d), fly ash particles are glassy spheres, composed mainly of O, Si, and Al with minor components, such as Fe and Ca, which can be identified clearly by their spherical shape and bright contrast on SEIs. The fly ash particles are different from tarballs having only C and O signals in their X-ray spectra, even though both are generated during the combustion processes (Geng et al., 2017). They were observed only in the Manaus samples, reflecting their anthropogenic origin.

3.1.9. Heavy metal-containing particles

Heavy metal-containing (HM) particles, such as Ni-, Ti-, Zn-, and Fe-containing ones, appear bright and irregular on SEM images, as shown in Figs. 11(e) and (f), and were observed mostly in the fine fraction with more than a half of them being Fe-containing particles both in the ATTO and Manaus samples. The Fe-containing particles in the ATTO samples were observed to be associated with SOA (and ammonium sulfate), as shown in Fig 11(f), indicating its mixing with the species of a biogenic emission origin. Sahara mineral dust has been reported to be essential for the nutrient cycles in the Amazon rainforest because many types of minerals are carried and transported into the rainforest, in which Fe is one of the important micronutrients in a Fe-limited rainforest (Rizzolo et al., 2017). Among all the mineral dust particles observed in the samples, approximately 40% of them contain Fe. The HM particles can also be of anthropogenic origin: emitted from the streets or road surface as brake dust, road paint particles, diesel exhaust particles, construction materials, and/or car catalyst materials (Qiao et al., 2016).

3.2. Relative abundances of particle types observed in the ATTO and Manaus samples

Figure 13 shows the relative abundance of the nine different particle types observed in the ATTO and Manaus samples. In the stage 1 samples (0.25-0.5 μm size) of SA1-SA4 collected at the ATTO site, almost all the particles were SOA and AS mixtures. In the stage 2 samples (0.5-1.0 μm size), SOA and AS mixture particles were dominant for the SA2 and SA3 samples. In the stage

2 samples of samples SA1 and SA4, SOA and AS mixture particles were most abundant, followed by pure ammonium sulfates, aged mineral dust, and pure SOA particles. In the stage 2 samples of SA1-SA4, the summed contents of SOA and ammonium sulfate were 73%, 99%, 85%, and 82%, respectively, suggesting that SOA and AS are the predominant species in submicron aerosols collected at the ATTO site. The observation of abundant submicron SOAs, which constitute a significant fraction of fine aerosol mass during the wet season at the rainforest, has been reported (Chen et al., 2015; Gilardoni et al., 2011). Although ammonium sulfates were reported to be present in significant quantities in the Amazon basin (Andreae et al., 2015; Chen et al., 2015; Fraund et al., 2017), this study emphasizes the observation of the predominant submicron ammonium sulfates mixed with SOA.

In the stage 2 samples, aged mineral dust and sea-salts for the SA1 sample, reacted sea salts for the SA3 sample, and aged mineral dust and sea-salts for the SA4 sample were significantly observed, suggesting that the samples from outside the Amazon rainforest have different influences because the mineral dust and sea-salt particles were all aged ones. The influences from outside were observed more clearly for supermicron aerosols at the ATTO site. In the stage 3 samples (1.0-2.0 μm size) of SA1-SA4, reacted sea-salt particles and aged mineral dust particles were abundantly observed, although the summed relative abundances of SOA and ammonium sulfate were 28%, 43%, 58%, and 50%, respectively, indicating that SOA and ammonium sulfate are abundant even in supermicron ATTO aerosols. As all the mineral dust particles were aged, they are not of local origin, and the observation of a high content of aged mineral dust particles in sample SA1 highlights the importance of long-range transatlantic transport (see the fast-moving air-masses from the Atlantic Ocean for the SA1 sample (Fig. 2(a) although a single backtrajectory cannot confirm their African origin of the aged mineral dust particles). Many studies have examined the influence of the Saharan dust particles over the Amazon rainforest region, starting with the measurements made during the ABLE-2B campaign (Talbot et al., 1990; Swap et al., 1992). Mineral dust is imported most frequently to the rainforest in March and April (Martin et al., 2010a; Moran-Zuloaga et al., 2018), which increases the ground-based soil dust element levels significantly (Artaxo et al., 2013). In this study, approximately 70% and 90% of the reacted sea-salt/mineral dust particles in the ATTO and Manaus samples, respectively, were either mixed or coated with organic matter in the Amazon basin and/or during the long-range transport.

PBA particles, which are from the rainforest, are sometimes observed in the stage 3 samples

for the SA1-SA4 samples. In the stage 4 sample (2.0-4.0 μm size) of the SA4 sample, the most abundant particle type was PBA, followed in order by reacted sea-salt, SOA, aged mineral dust, and ammonium sulfate, where both PBA, SOA, and ammonium sulfate particles of a local origin and the reacted sea-salt and mineral dust particles from the outside are considerably present. In summary, the aerosols collected at the ATTO site are mostly SOA and ammonium sulfate of a local origin in the submicron fraction, although some of the submicron sulfate are of marine and distant origin, whereas aerosols of both local and distant origins are significant in the supermicron fraction.

The aerosols collected at the Manaus site were diverse compared to those at the ATTO site. As shown in Fig. 13, for the stage 1 and 2 samples, SOA and ammonium sulfate particles, including their mixture, were the major components for the SM1 and SM2 samples, whereas they were relatively less abundant for the SM3 sample. As SOA and ammonium sulfate particles can be from the surrounding rainforest areas in addition to local anthropogenic sources, samples SM1-SM3 collected at the Manaus site appear to be influenced from the outside in the order of samples SM1 > SM2 > SM3. In addition, considerable amounts of submicron carbonaceous particles were observed, such as soot, char, and tarballs, which are of a local origin. The most abundant submicron aerosols for sample SM3 were carbonaceous ones, indicating that the local influence to the samples is in the order of SM3 > SM2 > SM1. Samples SM1 and SM2 were collected during and just after a national holiday, respectively, when all the institutions (private and public) were closed during that day so that the traffics were quite low, similar to a weekend or vacation period. Sample SM3 was collected during a regular working day, so that sample SM3 is the only sample actually exposed to the high traffic of light vehicles in the area.

In supermicron Manaus aerosols, PBA particles, aged mineral dust, and reacted sea-salts in addition to SOA and carbonaceous particles are abundant. In the stage 3 aerosols of the SM1 sample, the most abundant particles were reacted sea-salts, followed by aged mineral dust, SOA, ammonium sulfate, and PBA particles, which also indicates the strong influence on the SM1 sample from the outside. In stage 3 aerosols of the SM2 sample, the most abundant particles were SOA, followed in order by aged mineral dust, reacted sea-salt, PBA, carbonaceous particles, and ammonium sulfate, which also indicates the strong influence on the SM2 sample from the surrounding rainforest areas. In the stages 3 and 4 aerosols of the SM3 sample, the most abundant particles were aged mineral dust (36% and 52% for stages 3 and 4, respectively), followed by carbonaceous particles, PBA, SOA, and reacted sea-salt. As the aged sea-salt contents were

relatively low (10% and 4% for stages 3 and 4, respectively), most of the aged mineral dusts appear to be of a local origin. Fly ash and heavy metal-containing particles of an anthropogenic local origin were considerable in the Manaus samples, i.e., they were observed in the range of 3-7% relative abundances in the Manaus samples. In particular, soot particles of an anthropogenic origin were observed ubiquitously in all the Manaus samples. Among the samples, the aerosols collected at the Manaus site were different. The SM1 sample was influenced most strongly from the outside, including the surrounding rainforest and transatlantic transport. The SM2 sample has some influences by local sources as well as from the outside. The SM3 sample contains mainly aerosols of an anthropogenic local origin in the submicron fraction and some influences from the outside in the supermicron fraction. Figures 2(e)-(f) show that the backward trajectories at heights of 100-m and 500-m are further from the outside in the order of SM1 > SM2 > SM3, which is consistent with the characteristics of submicron and supermicron aerosols of the SM1-SM3 samples.

Conclusions

In this study, aerosol samples collected in the Amazonian rainforest and Manaus, Brazil during the 2012 wet season were investigated on a single particle basis using low-Z particle EPMA. The aerosol particles were classified into nine particle types based on their morphology on SEIs together with the elemental concentrations of a total of 3,162 individual particles: (i) secondary organic aerosols (SOA), (ii) ammonium sulfate (AS) particles, (iii) SOA and AS mixture particles, (iv) aged mineral dust, (v) reacted sea-salts, (vi) primary biological aerosol (PBA) particles, (vii) carbon-rich or elemental carbon (EC) particles such as soot, tarball, and char, (viii) fly ash particles, and (ix) heavy metal (HM, such as Fe, Zn, Ni, and Ti)-containing particles. For submicron aerosols collected at the ATTO site, the SOA and AS mixture particles were predominant (50-94% in relative abundance) with the summed contents of SOA and ammonium sulfate being 73-100%. In contrast, in the supermicron aerosols at the ATTO site, aged mineral dust and sea-salt (37-70%) as well as SOA and ammonium sulfate (28-58%) were abundant. PBAs were observed abundantly in the PM₂₋₄ fraction (46%), and EC and fly ash particles were absent in all the fractions. An analysis of a bulk PM_{0.25-0.5} aerosol sample collected at the ATTO site using RMS and ATR-FTIR showed that ammonium sulfate, organics, and minerals are the major chemical species, which is consistent with the EPMA results.

In the submicron aerosols collected in Manaus, either SOA and ammonium sulfate (17-

80%) or EC particles (6-78%) were dominant, depending on the samples. The supermicron aerosols collected in Manaus consisted mainly of aged mineral dust, reacted sea-salts, PBA, SOA, ammonium sulfate, and EC particles. SOA, ammonium sulfate, and PBAs were mostly of a biogenic origin and EC and HM-containing particles were of an anthropogenic origin. The aged mineral dust and reacted sea-salt particles, including mineral dust mixed with sea-salts probably during long-range transatlantic transport, were abundant in the supermicron fractions at both sites. The submicron aerosol at the ATTO site was influenced mainly by the emission from the rainforest and its supermicron aerosols showed additional contributions from long-range transport, including the Atlantic Ocean and Sahara Desert, whereas the aerosols collected in Manaus showed different local and outside contributions among the samples. Among all the aged mineral dust and reacted sea-salt particles, sulfate-containing ones outnumbered those containing nitrates and both nitrate and sulfate in the ATTO samples, whereas N and S containing particles were comparable to sulfate-only ones in the Manaus samples, indicating the different sources and formation mechanisms of secondary aerosols, i.e., the predominant presence of sulfate at the ATTO site from biogenic emissions and elevated influences of nitrates from anthropogenic activities at the Manaus site.

Author contributions

LW, RHMG, and CR designed the experiment and LW, XL, and HKK carried out the measurements and analyzed the data. CGGB, CIY, RAFDS, and CP organized and performed the samplings. LW, HG, RHMG, CGGB, AFLG, and CR interpreted the observations and LW, HG, RHMG, CGGB, MOA, and CR drafted the paper.

Acknowledgement

This study was supported by Basic Science Research Programs through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science, and Technology (NRF-2018R1A2A1A05023254). The work of M. O. Andreae and C. Pöhlker was supported by the German Max Planck Society (MPG). For the operation of the ATTO site, we acknowledge the support by the German Federal Ministry of Education and Research (BMBF contract 01LB1001A), the Brazilian Ministério da Ciência, Tecnologia e Inovação (MCTI/FINEP contract 01.11.01248.00), and the Coordination for the Improvement of Higher Education Personnel (CAPES) for scholarship funding (investigation) as well as the Amazon State University (UEA),

FAPEAM, LBA/INPA, and SDS/CEUC/RDS-Uatumã. A special thanks to Claudomiro Mauricio da Silva for the support during sampling.

References

- Abouchami, W., N  the, K., Kumar, A., Galer, S. J. G., Jochum, K. P., Williams, E., Horbe, A. M. C., Rosa, J. W. C., Balsam, W., Adams, D., Mezger, K., and Andreae, M. O., Geochemical and isotopic characterization of the Bod  l   Depression dust source and implications for transatlantic dust transport to the Amazon Basin, *Earth Planet. Sci. Lett.*, 380, 112-123, doi:10.1016/j.epsl.2013.08.028, 2013.
- Andreae, M. O.: Aerosols before pollution, *Science*, 315, 50–51, doi:10.1126/science.1136529, 2007.
- Andreae, M. O. and Gelencs  r, A.: Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols, *Atmos. Chem. Phys.*, 6, 3131–3148, doi:10.5194/acp-6-3131-2006, 2006.
- Andreae, M. O., Berresheim, H., Bingemer, H., Jacob, D. J., Lewis, B. L., Li, S.-M., and Talbot, R. W.: The atmospheric sulfur cycle over the Amazon Basin, 2. Wet Season, *J. Geophys. Res.*, 95, 16813–16824, doi:10.1029/JD095iD10p16813, 1990.
- Andreae, M. O., Acevedo, O. C., Ara  jo, A., Artaxo, P., Barbosa, C. G. G., Barbosa, H. M. J., Brito, J., Carbone, S., Chi, X., Cintra, B. B. L., da Silva, N. F., Dias, N. L., Dias-J  nior, C. Q., Ditas, F., Ditz, R., Godoi, A. F. L., Godoi, R. H. M., Heimann, M., Hoffmann, T., Kesselmeier, J., K  nemann, T., Kr  ger, M. L., Lavric, J. V., Manzi, A. O., Lopes, A. P., Martins, D. L., Mikhailov, E. F., Moran-Zuloaga, D., Nelson, B. W., N  lscher, A. C., Santos Nogueira, D., T. F. Piedade, M., P  hlker, C., P  schl, U., Quesada, C. A., Rizzo, L. V., Ro, C.-U., Ruckteschler, N., S  , L. D. A., de Oliveira S  , M., Sales, C. B., dos Santos, R. M. N., Saturno, J., Sch  ngart, J., S  rgel, M., de Souza, C. M., de Souza, R. A. F., Su, H., Targhetta, N., T  ta, J., Trebs, I., Trumbore, S., van Eijck, A., Walter, D., Wang, Z., Weber, B., Williams, J., Winderlich, J., Wittmann, F., Wolff, S., and Y    ez-Serrano, A. M.: The Amazon Tall Tower Observatory (ATTO): overview of pilot measurements on ecosystem ecology, meteorology, trace gases, and aerosols, *Atmos. Chem. Phys.*, 15, 10723–10776, doi:10.5194/acp-15-10723-2015, 2015.
- Andreae, M. O., Afchine, A., Albrecht, R., Holanda, B. A., Artaxo, P., Barbosa, H. M. J., Borrmann, S., Cecchini, M. A., Costa, A., Dollner, M., F  tterer, D., J  rvinen, E., Jurkat, T., Klimach, T.,

- Konemann, T., Knote, C., Krämer, M., Krisna, T., Machado, L. A. T., Mertes, S., Minikin, A., Pöhlker, C., Pöhlker, M. L., Pöschl, U., Rosenfeld, D., Sauer, D., Schlager, H., Schnaiter, M., Schneider, J., Schulz, C., Spanu, A., Sperling, V. B., Voigt, C., Walser, A., Wang, J., Weinzierl, B., Wendisch, M., and Ziereis, H.: Aerosol characteristics and particle production in the upper troposphere over the Amazon Basin, *Atmos. Chem. Phys.*, 18, 921-961, doi:10.5194/acp-18-921-2018, 2018.
- Artaxo, P., Fernandes, E. T., Martins, J. V., Yamasoe, M. A., Hobbs, P. V., Maenhaut, W., Longo, K. M., and Castanho, A.: Large-scale aerosol source apportionment in Amazonia, *J. Geophys. Res.*, 103(D24), 31837–31847, doi:10.1029/98JD02346, 1998.
- Artaxo, P., Rizzo, L. V., Brito, J. F., Barbosa, H. M., Arana, A., Sena, E. T., Cirino, G. G., Bastos, W., Martin, S. T., Andreae, M., O.: Atmospheric aerosols in Amazonia and land use change: from natural biogenic to biomass burning conditions, *Faraday Discuss.*, 165, 203-235, doi:10.1039/C3FD00052D, 2013.
- Baars, H., Ansmann, A., Althausen, D., Engelmann, R., Artaxo, P., Pauliquevis, T. and Souza, R.: Further evidence for significant smoke transport from Africa to Amazonia, *Geophys. Res. Lett.*, 38, L20802, doi:10.1029/2011GL049200, 2011.
- Bateman, A. P., Gong, Z., Liu, P., Sato, B., Cirino, G., Zhang, Y., Artaxo, P., Bertram, A. K., Manzi, A. O., Rizzo, L. V., Souza, R. A. F., Zaveri, R. A., and Martin, S. T.: Sub-micrometre particulate matter is primarily in liquid form over Amazon rainforest, *Nat. Geosci.*, 9, 34–37, doi:10.1038/ngeo2599, 2016.
- Bateman, A. P., Gong, Z., Harder, T. H., de Sá, S. S., Wang, B., Castillo, P., China, S., Liu, Y., O'Brien, R. E., Palm, B. B., Shiu, H.-W., Cirino, G. G., Thalman, R., Adachi, K., Alexander, M. L., Artaxo, P., Bertram, A. K., Buseck, P. P., Gilles, M. K., Jimenez, J. L., Laskin, A., Manzi, A. O., Sedlacek, A., Souza, R. A. F., Wang, J., Zaveri, R., and Martin, S. T.: Anthropogenic influences on the physical state of submicron particulate matter over a tropical forest, *Atmos. Chem. Phys.*, 17, 1759–1773, doi:10.5194/acp-17-1759-2017, 2017.
- Bazzaz, F. A.: Tropical Forests in a Future Climate: Changes in Biological Diversity and Impact on the Global Carbon Cycle, *Climatic Change*, 39, 317-336, doi:10.1023/A:1005359605003, 1998.
- Buseck, P. R. and Pósfai, M.: Airborne minerals and related aerosol particles: Effects on climate and the environment, *Proc. Natl. Acad. Sci. U.S.A.* 96, 3372-3379,

- doi:10.1073/pnas.96.7.3372, 1999.
- Chen, Y., Shah, N., Huggins, F. E., and Huffman, G. P.: Characterization of ambient airborne particles by energy-filtered transmission electron microscopy, *Aerosol Sci. Technol.*, 39, 509–518, doi:10.1080/027868291001402, 2005.
- Chen, Y., Shah, N., Huggins, F.E., and Huffman, G.P.: Microanalysis of ambient particles from Lexington, KY, by electron microscopy, *Atmos. Environ.*, 40, 651–663, doi:10.1016/j.atmosenv.2005.09.036, 2006.
- Chen, H. H., Navea, J. G., Young, M. A., and Grassian, V. H.: Heterogeneous Photochemistry of Trace Atmospheric Gases with Components of Mineral Dust Aerosol, *J. Phys. Chem. A*, 115, 490-499, doi:10.1021/jp110164j, 2011.
- Chen, Q., Farmer, D. K., Rizzo, L. V., Pauliquevis, T., Kuwata, M., Karl, T. G., Guenther, A., Allan, J. D., Coe, H., Andreae, M. O., Pöschl, U., Jimenez, J. L., Artaxo, P., and Martin, S. T.: Submicron particle mass concentrations and sources in the Amazonian wet season (AMAZE-08), *Atmos. Chem. Phys.*, 15, 3687–3701, doi:10.5194/acp-15-3687-2015, 2015.
- China, S., Wang, B., Weis, J., Rizzo, L., Brito, J., Cirino, G. G., Kovarik, L., Artaxo, P., Gilles, M. K., Laskin, A.: Rupturing of Biological Spores As a Source of Secondary Particles in Amazonia, *Environ. Sci. Technol.*, 50, 12179-12186, doi: 10.1021/acs.est.6b02896, 2016.
- de Sá, S. S., Palm, B. B., Campuzano-Jost, P., Day, D. A., Hu, W., Isaacman-VanWertz, G., Yee, L. D., Brito, J., Carbone, S., Ribeiro, I. O., Cirino, G. G., Liu, Y. J., Thalman, R., Sedlacek, A., Funk, A., Schumacher, C., Shilling, J. E., Schneider, J., Artaxo, P., Goldstein, A. H., Souza, R. A. F., Wang, J., McKinney, K. A., Barbosa, H., Alexander, M. L., Jimenez, J. L., and Martin, S. T.: Urban influence on the concentration and composition of submicron particulate matter in central Amazonia, *Atmos. Chem. Phys.*, 18, 12185-12206, doi:10.5194/acp-18-12185-2018, 2018.
- Eom, H.-J., Gupta, D., Li, X., Jung, H.-J., Kim, H., and Ro, C.-U.: Influence of Collecting Substrates on the Characterization of Hygroscopic Properties of Inorganic Aerosol Particles, *Anal. Chem.*, 86, 2648–2656, doi:10.1021/ac4042075, 2014.
- Eom, H.-J., Gupta, D., Cho, H.-R., Hwang, H. J., Hur, S. D., Gim, Y., and Ro, C.-U.: Single-particle investigation of summertime and wintertime Antarctic sea spray aerosols using low-Z particle EPMA, Raman microspectrometry, and ATR-FTIR imaging techniques, *Atmos. Chem. Phys.*, 16, 13823-13836, 10.5194/acp-16-13823-2016, 2016.

- Fan, J., Rosenfeld, D., Zhang, Y., Giangran, S. E., Li, Z., Machado, L. A. T., Martin, S. T., Yang, Y., Wang, J., Artaxo, P., Barbosa, H. M. J., Braga, R. C., Comstock, J. M., Feng, Z., Gao, W., Gomes, H. B., Mei, F., Pöhlker, C., Pöhlker, M. L., Pöschl, U., and de Souza, R. A. F.: Substantial convection and precipitation enhancements by ultrafine aerosol particles, *Science*, 359, 411-418, 2018.
- Fraund, M., Pham, D. Q., Bonanno, D., Harder, T. H., Wang, B., Brito, J., Sá, S. S. de, Carbone, S., China, S., Artaxo, P., Martin, S. T., Pöhlker, C., Andreae, M. O., Laskin, A., Gilles, M., K., and Moffet, R. C.: Elemental Mixing State of Aerosol Particles Collected in Central Amazonia during GoAmazon2014/15, *Atmosphere*, 8, 173, doi:10.3390/atmos8090173, 2017.
- Fung, K. H. and Tang, I. N.: Raman spectra of singly suspended supersaturated ammonium bisulfate droplets, *Chem. Phys. Lett.*, 147, 509-513, doi:10.1016/0009-2614(88)85017-6, 1988.
- Fuzzi, S., Decesari, S., Facchini, M. C., Cavalli, F., Emblico, L., Mircea, M., Andreae, M. O., Trebs, I., Hoffer, A., Guyon, P., Artaxo, P., Rizzo, L. V., Lara, L. L., Pauliquevis, T., Maenhaut, W., Raes, N., Chi, X., Mayol-Bracero, O. L., Soto-García, L. L., Claeys, M., Kourtchev, I., Rissler, J., Swietlicki, E., Tagliavini, E., Schkolnik, G., Falkovich, A. H., Rudich, Y., Fisch, G., and Gatti, L. V.: Overview of the inorganic and organic composition of size-segregated aerosol in Rondonia, Brazil, from the biomass-burning period to the onset of the wet season, *J. Geophys. Res.* 112, D01201, doi:10.1029/2005JD006741, 2007.
- Geng, H., Jung, H.-J., Park, Y., Hwang, H., Kim, H., Kim, Y.J., Sunwoo, Y., and Ro, C.-U.: Morphological and chemical composition characteristics of summertime atmospheric particles collected at Tokchok Island, Korea, *Atmos. Environ.* 43, 3364–3373, doi:10.1016/j.atmosenv.2009.03.034, 2009.
- Geng, H., Kang, S., Choel, M., Kim, H., and Ro, C.-U.: Characterization of individual submicrometer aerosol particles collected in Incheon, Korea, by quantitative transmission electron microscopy energy-dispersive X-ray spectrometry, *J. Geophys. Res.*, 115, D15306, doi:10.1029/2009JD013486, 2010.
- Geng, H., Ryu, J., Maskey, S., Jung, H.-J., and Ro, C.-U.: Characterisation of individual aerosol particles collected during a haze episode in Incheon, Korea using the quantitative ED-EPMA technique, *Atmos. Chem. Phys.* 11, 1327–1337, doi:10.5194/acp-11-1327-2011, 2011.
- Geng, H., Hwang, H., Liu, X., Dong, S., and Ro, C.-U.: Investigation of aged aerosols in size

- resolved Asian dust storm particles transported from Beijing, China, to Incheon, Korea, using low-Z particle EPMA, *Atmos. Chem. Phys.*, 14, 3307–3323, doi:10.5194/acp-14-3307-2014, 2014.
- Geng, H., Jin, C.-S., Zhang, D.-P., Wang, S.-R., Xu, X.-T., Wang, X.-R., Zhang, Y., Wu, L., and Ro, C.-U.: Characterization of size-resolved urban haze particles collected in summer and winter at Taiyuan City, China using quantitative electron probe X-ray microanalysis, *Atmos. Res.*, 190, 29–42, doi: 10.1016/j.atmosres.2017.02.005, 2017.
- Gilardoni, S., Vignati, E., Marmer, E., Cavalli, F., Belis, C., Gianelle, V., Loureiro, A., and Artaxo, P.: Sources of carbonaceous aerosol in the Amazon basin, *Atmos. Chem. Phys.*, 11, 2747–2764, doi:10.5194/acp-11-2747-2011, 2011.
- Goldstein, J., Newbury, D., Joy, D., Lyman, C., Echlin, P., Lifshin, E., Sawyer, L., and Michael, J.: *Scanning Electron Microscopy and X-ray Microanalysis*, third ed., Kluwer Academic/Plenum Publishers, New York, 2003.
- Gregory, G. L., Harriss, R. C., Talbot, R. W., Rasmussen, R. A., Garstang, M., Andreae, M. O., Hinton, R. R., Browell, E. V., Beck, S. M., Sebacher, D. I., Khalil, M. A. K., Ferek, R. J., and Harriss, S. V.: Air chemistry over the tropical forest of Guyana: *J. Geophys. Res.*, 91, 8603–8612, 1986.
- Haga, D. I., Burrows, S. M., Iannone, R., Wheeler, M. J., Mason, R. H., Chen, J., Polishchuk, E. A., Pöschl, U., and Bertram, A. K.: Ice nucleation by fungal spores from the classes *Agaricomycetes*, *Ustilaginomycetes*, and *Eurotiomycetes*, and the effect on the atmospheric transport of these spores, *Atmos. Chem. Phys.*, 14, 8611–8630, doi:10.5194/acp-14-8611-2014, 2014.
- Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, Th. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155–5236, doi:10.5194/acp-9-5155-2009, 2009.
- Hansen, K., Pryor, S. C., Boegh, E., Hornsby, K. E., Jensen, B., and Sørensen, L. L.: Background concentrations and fluxes of atmospheric ammonia over a deciduous forest, *Agric. For.*

Meteorol., 214-215, 380-392, doi:10.1016/j.agrformet.2015.09.004, 2015.

Hansen, K., Personne, E., Skjøth, C. A., Loubet, B., Ibrom, A., Jensen, R., Sørensen, L. L., and Boegh, E.: Investigating sources of measured forest-atmosphere ammonia fluxes using two-layer bi-directional modelling, *Agric. For. Meteorol.*, 237-238, 80-94, doi:10.1016/j.agrformet.2017.02.008, 2017.

Huang, P. F. and Turpin, B.: Reduction of sampling and analytical errors for electron microscopic analysis of atmospheric aerosols, *Atmos. Environ.*, 30, 4137-4148, 1996.

IBGE: Estimates of the resident population in Brazil and Federative Units, Brazilian Institute of Geography and Statistics, 2017.

Jardine, K., Yañez-Serrano, A. M., Williams, J., Kunert, N., Jardine, A., Taylor, T., Abrell, L., Artaxo, P., Guenther, A., Hewitt, C. N., House, E., Florentino, A. P., Manzi, A., Higuchi, N., Kesselmeir, J., Behrendt, T., Veres, P. R., Derstroff, B., Fuentes, J. D., Martin, S. T., Andreae, M. O.: Dimethyl sulfide in the Amazon rain forest, *Global Biogeochem. Cycles*, 29, 19–32, doi:10.1002/2014GB004969, 2015.

Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, E. J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525–1529, doi:10.1126/science.1180353, 2009.

Jordan, C., Caskey, W., Escalante, G., Herrera, R., Montagnini, F., Todd, R., and Uhl, C.: The nitrogen cycle in a ‘Terra Firme’ rainforest on oxisol in the Amazon territory of Venezuela, *Plant and Soil*, 67, 325-332, doi:10.1007/BF02182779, 1982.

Kellerhals, T., Brüttsch, S., Sigl, M., Knüsel, S., Gäggeler, H. W., and Schwikowski, M.: Ammonium concentration in ice cores: A new proxy for regional temperature reconstruction?,

- J. Geophys. Res., 115, D16123, doi:10.1029/2009JD012603, 2010.
- Kim, J. and Rees, D. C.: Nitrogenase and biological nitrogen fixation, *Biochemistry*, 33, 389–397, 1994.
- Krejci, R., Ström, J., Reus, M. de, Sahle, W.: Single particle analysis of the accumulation mode aerosol over the northeast Amazonian tropical rain forest, Surinam, South America, *Atmos. Chem. Phys.*, 5, 3331–3344, 2005.
- Laskin, A., Gaspar, D. J., Wang, W., Hunt, S. W., Cowin, J. P., Colson, S. D., and Finlayson-Pitts, B. J.: Reactions at Interfaces as a Source of Sulfate Formation in Sea-Salt Particles, *Science*, 301, 340–344, doi:10.1126/science.1085374, 2003.
- Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of Atmospheric Brown Carbon, *Chem. Rev.*, 115, 4335–4382, doi:10.1021/cr5006167, 2015.
- Li, W., Shao, L., Zhang, D., Ro, C.-U., Hu, M., Bi, X., Geng, H., Matsuki, A., Niu, H., and Chen, J.: A review of single aerosol particle studies in the atmosphere of East Asia: morphology, mixing state, source, and heterogeneous reactions, *J. Clean. Prod.*, 112, 1330–1349, doi:10.1016/j.jclepro.2015.04.050, 2016.
- Lin, G., Sillman, S., Penner, J. E., and Ito, A.: Global modeling of SOA: the use of different mechanisms for aqueous-phase formation, *Atmos. Chem. Phys.*, 14, 5451–5475, doi:10.5194/acp-14-5451-2014, 2014.
- Ling, T. Y., and Chan, C. K.: Formation and transformation of metastable double salts from the crystallization of mixed ammonium nitrate and ammonium sulfate particles, *Environ. Sci. & Technol.*, 41, 8077–8083, doi:10.1021/es071419t, 2007.
- Mabrouk, K. B., Kauffmann, T. H., Aroui, H., and Fontana, M. D.: Raman study of cation effect on sulfate vibration modes in solid state and in aqueous solutions, *J. Raman Spectrosc.*, 44, 1603–1608, doi: 10.1002/jrs.4374, 2013.
- Mace, K. A., Artaxo, P., and Duce, R. A.: Water-soluble organic nitrogen in Amazon Basin aerosols during the dry (biomass burning) and wet seasons, *J. Geophys. Res.*, 108(D16), 4512, doi:10.1029/2003JD003557, 2003.
- Martin, S. T., Andreae, M. O., Artaxo, P., Baumgardner, D., Chen, Q., Goldstein, A. H., Guenther, A., Heald, C. L., Mayol-Bracero, O. L., McMurry, P. H., Pauliquevis, T., Pöschl, U., Prather, K. A., Roberts, G. C., Saleska, S. R., Dias, M. A. S., Spracklen, D. V., Swietlicki, E., and Trebs, I.: Sources and properties of Amazonian aerosol particles, *Rev. Geophys.*, 48, RG2002,

doi:10.1029/2008rg000280, 2010a.

Martin, S. T., Andreae, M. O., Althausen, D., Artaxo, P., Baars, H., Borrmann, S., Chen, Q., Farmer, D. K., Guenther, A., Gunthe, S. S., Jimenez, J. L., Karl, T., Longo, K., Manzi, A., Müller, T., Pauliquevis, T., Petters, M. D., Prenni, A. J., Pöschl, U., Rizzo, L. V., Schneider, J., Smith, J. N., Swietlicki, E., Tota, J., Wang, J., Wiedensohler, A., and Zorn, S. R.: An overview of the Amazonian Aerosol Characterization Experiment 2008 (AMAZE-08), *Atmos. Chem. Phys.*, 10, 11415–11438, doi:10.5194/acp-10-11415-2010, 2010b.

Martin, S. T., Artaxo, P., Machado, L. A. T., Manzi, A. O., Souza, R. A. F., Schumacher, C., Wang, J., Andreae, M. O., Barbosa, H. M. J., Fan, J., Fisch, G., Goldstein, A. H., Guenther, A., Jimenez, J. L., Pöschl, U., Silva Dias, M. A., Smith, J. N., and Wendisch, M.: Introduction: Observations and Modeling of the Green Ocean Amazon (GoAmazon2014/5), *Atmos. Chem. Phys.*, 16, 4785–4797, doi:10.5194/acp-16-4785-2016, 2016.

Maskey, S., Choël, M., Kang, S., Hwang, H., Kim, H., and Ro, C.-U.: The influence of collecting substrates on the single-particle characterization of real atmospheric aerosols, *Anal. Chim. Acta*, 658, 120–127, doi:10.1016/j.aca.2009.11.006, 2010.

Medeiros, A. S. S., Calderaro, G., Guimarães, P. C., Magalhaes, M. R., Morais, M. V. B., Rafee, S. A. A., Ribeiro, I. O., Andreoli, R. V., Martins, J. A., Martins, L. D., Martin, S. T., and Souza, R. A. F.: Power plant fuel switching and air quality in a tropical, forested environment, *Atmos. Chem. Phys.*, 17, 8987–8998, doi:10.5194/acp-17-8987-2017, 2017.

Millstein, D. E., Harley, R. A., and Hering, S. V.: Weekly cycles in fine particulate nitrate, *Atmos. Environ.* 42, 632–641, doi:10.1016/j.atmosenv.2007.10.010, 2008.

Möhler, O., Benz, S., Saathoff, H., Schnaiter, M., Wagner, R., Schneider, J., Walter, S., Ebert, V., Wagner, S.: The effect of organic coating on the heterogeneous ice nucleation efficiency of mineral dust aerosols, *Environ. Res. Lett.*, 3, 025007, doi:10.1088/1748-9326/3/2/025007, 2008.

Moran-Zuloaga, D., Ditas, F., Walter, D., Saturno, J., Brito, J., Carbone, S., Chi, X., de Angelis, I. H., Baars, H., Godoi, R. H. M., Heese, B., Holanda, B. A., Lavrič, J. V., Martin, S. T., Ming, J., Pöhlker, M. L., Ruckteschler, N., Su, H., Wang, Y., Wang, Q., Wang, Z., Weber, B., Wolff, S., Artaxo, P., Pöschl, U., Andreae, M. O., and Pöhlker, C.: Long-term study on coarse mode aerosols in the Amazon rain forest with the frequent intrusion of Saharan dust plumes, *Atmos. Chem. Phys.*, 18, 10055–10088, doi:10.5194/acp-18-10055-2018, 2018.

- Neill, C., Piccolo, M. C., Melillo, J. M., Steudler, P. A., Cerri, C. C.: Nitrogen dynamics in Amazon forest and pasture soils measured by ^{15}N pool dilution, *Soil Biol. Biochem.*, 31, 567-572, 1999.
- Palm, B. B., de Sá, S. S., Day, D.A., Campuzano-Jost, P., Hu, W., Seco, R., Sjöstedt, S. J., Park, J.-H., Guenther, A. B., Kim, S., Brito, J., Wurm, F., Artaxo, P., Thalman, R., Wang, J., Yee, L. D., Wernis, R., Isaacman-VanWertz, G., Goldstein, A. H., Liu, Y., Springston, S. R., Souza, R., Newburn, M. K., Alexander, M. L., Martin, S.T., and Jimenez, J. L.: Secondary organic aerosol formation from ambient air in an oxidation flow reactor in central Amazonia, *Atmos. Chem. Phys.*, 18, 467–493, doi:10.5194/acp-18-467-2018, 2018.
- Pöhlker, C., Wiedemann, K. T., Sinha, B., Shiraiwa, M., Gunthe, S. S., Smith, M., Su, H., Artaxo, P., Chen, Q., Cheng, Y. F., Elbert, W., Gilles, M. K., Kilcoyne, A. L. D., Moffet, R. C., Weigand, M., Martin, S. T., Poeschl, U., and Andreae, M. O.: Biogenic potassium salt particles as seeds for secondary organic aerosol in the Amazon, *Science*, 337, 1075–1078, doi:10.1126/science.1223264, 2012.
- Pöhlker, M. L., Ditas, F., Saturno, J., Klimach, T., Hrabě de Angelis, I., Araùjo, A. C., Brito, J., Carbone, S., Cheng, Y., Chi, X., Ditz, R., Gunthe, S. S., Holanda, B. A., Kandler, K., Kesselmeier, J., Könemann, T., Krüger, O. O., Lavrič, J. V., Martin, S. T., Mikhailov, E., Moran-Zuloaga, D., Rizzo, L. V., Rose, D., Su, H., Thalman, R., Walter, D., Wang, J., Wolff, S., Barbosa, H. M. J., Artaxo, P., Andreae, M. O., Pöschl, U., and Pöhlker, C.: Long-term observations of cloud condensation nuclei over the Amazon rain forest – Part 2: Variability and characteristics of biomass burning, long-range transport, and pristine rain forest aerosols, *Atmos. Chem. Phys.*, 18, 10289-10331, doi:10.5194/acp-18-10289-2018, 2018.
- Pöschl, U., Martin, S. T., Sinha, B., Chen, Q., Gunthe, S. S., Huffman, J. A., Borrmann, S., Farmer, D. K., Garland, R. M., Helas, G., Jimenez, J. L., King, S. M., Manzi, A., Mikhailov, E., Pauliquevis, T., Petters, M. D., Prenni, A. J., Roldin, P., Rose, D., Schneider, J., Su, H., Zorn, S. R., Artaxo, P., and Andreae, M. O.: Rainforest Aerosols as Biogenic Nuclei of Clouds and Precipitation in the Amazon, *Science* 329, 1513-1516, doi:10.1126/science.1191056, 2010.
- Pósfai, M., Xu, H., Anderson, and Buseck, P. R.: Wet and dry sizes of atmospheric aerosol particles: An AFM-TEM study, *Geophys. Res. Lett.* 25, 1907–1910, doi:10.1029/98GL01416, 1998.
- Pósfai, M., Anderson, J. R., Buseck, P. R., and Sievering, H.: Soot and sulfate aerosol particles in the remote marine troposphere, *J. Geophys. Res.* 104, 21, 685–21, 693,

- doi:10.1029/1999JD900208, 1999.
- Pósfai, M., Simonics, R., Li, J., Hobbs, P. V., and Buseck, P. R.: Individual aerosol particles from biomass burning in southern Africa: 1. Compositions and size distributions of carbonaceous particles, *J. Geophys. Res.*, 108(D13), 8483, doi:10.1029/2002JD002291, 2003.
- Pósfai, M., Gelencser, A., Simonics, R., Arato, K., Li, J., Hobbs, P. V., and Buseck, P. R.: Atmospheric tarballs: Particles from biomass and biofuel burning, *J. Geophys. Res.*, 109, D06213, doi:10.1029/2003JD004169, 2004.
- Prieto-Taboada, N., Gomez-Laserna, O., Martinez-Arkarazo, I., Olazabal, M. A., and Madariaga, J. M.: Raman Spectra of the Different Phases in the $\text{CaSO}_4\text{--H}_2\text{O}$ System, *Anal. Chem.*, 86, 20, 10131-10137, doi: 10.1021/ac501932f, 2014.
- Qiao, T., Zhao, M., Xiu, G., and Yu, J.: Simultaneous monitoring and compositions analysis of PM1 and PM2.5 in Shanghai: implications for characterization of haze pollution and source apportionment, *Sci. Total Environ.*, 557–558, 386-394, doi:10.1016/j.scitotenv.2016.03.095, 2016.
- Rizzolo, J. A., Barbosa, C. G. G., Borillo, G. C., Godoi, A. F. L., Souza, R. A. F., Andreoli, R. V., Manzi, A. O., Sá, M. O., Alves, E. G., Pöhlker, C., Angelis, I. H., Ditas, F., Saturno, J., Moran-Zuloaga, D., Rizzo, L. V., Rosário, N. E., Pauliquevis, T., Santos, R. M. N., Yamamoto, C. I., Andreae, M. O., Artaxo, P., Taylor, P. E., and Godoi, R. H. M.: Soluble iron nutrients in Saharan dust over the central Amazon rainforest, *Atmos. Chem. Phys.*, 17, 2673-2687, doi:10.5194/acp-17-2673-2017, 2017.
- Saturno, J., Ditas, F., Penning de Vries, M., Holanda, B. A., Pöhlker, M. L., Carbone, S., Walter, D., Bobrowski, N., Brito, J., Chi, X., Gutmann, A., Hrabe de Angelis, I., Machado, L. A. T., Moran-Zuloaga, D., Rüdiger, J., Schneider, J., Schulz, C., Wang, Q., Wendisch, M., Artaxo, P., Wagner, T., Pöschl, U., Andreae, M. O., and Pöhlker, C.: African volcanic emissions influencing atmospheric aerosols over the Amazon rain forest, *Atmos. Chem. Phys.*, 18, 10391–10405, doi:10.5194/acp-18-10391-2018, 2018.
- Sobanska, S., Hwang, H., Choël, M., Jung, H., Eom, H., Kim, H., Barbillat, J., and Ro, C.-U.: Investigation of the chemical mixing state of individual Asian Dust particles by the combined use of Electron Probe X-ray Microanalysis and Raman Microspectrometry, *Anal. Chem.*, 84, 3145-3154, doi:10.1021/ac2029584, 2012.
- Su, L., Yuan, Z., Fung, J. C. H., and Lau, A. K. H.: A comparison of HYSPLIT backward

trajectories generated from two GDAS datasets, *Sci. Total Environ.*, 506-507, 527-537, doi:10.1016/j.scitotenv.2014.11.072, 2015.

Sullivan, R. C., Guazzotti, S. A., Sodeman, D. A., and Prather, K. A.: Direct observations of the atmospheric processing of Asian mineral dust, *Atmos. Chem. Phys.*, 7, 1213-1236, doi:10.5194/acp-7-1213-2007, 2007.

Sutton, M.A., Nemitz, E., Milford, C., Campbell, C., Erisman, J. W., Hensen, A., Cellier, P., David, M., Loubet, B., Personne, E., Schjoerring, J. K., Mattsson, M., Dorsey, J. R., Gallagher, L., Horvath, M. W., Weidinger, T., Meszaros, R., Dämmgen, U., Neftel, A., Herrmann, B., Lehman, B. E., Flechard, C., and Burkhardt, J.: Dynamics of ammonia exchange with cut grassland: synthesis of results and conclusions of the GRAMINA integrated experiment. *Biogeosciences* 6, 2907–2934. doi:10.5194/bg-6-2907-2009, 2009.

Sutton, M.A., Reis, S., Riddick, S.N., Dragosits, U., Nemitz, E., Theobald, M. R., Tang, Y. S., Braban, C. F., Vieno, M., Dore, A. J., Mitchell, R. F., Wanless, S., Daunt, F., Fowler, D., Blackall, T. D., Milford, C., Flechard, C. R., Loubet, B., Massad, R., Cellier, P., Personne, E., Coheur, P. F., Clarisse, L., Van Damme, M., Ngadi, Y., Clerbaux, C., Skjøth, C. A., Geels, C., Hertel, O., Wichink Kruit, R. J., Pinder, R. W., Bash, J. O., Walker, J. T., Simpson, D., Horvath, L., Misselbrook, T. H., Bleeker, A., Dentener, F., and de Vries, W.: Towards a climate-dependent paradigm of ammonia emission and deposition, *Philos. Trans. R. Soc. Lond. Ser. B*, 368, doi:10.1098/rstb.2013.0166, 2013.

Swap, R., Garstang, M., Greco, S., Talbot, R., and Kållberg, P.: Saharan dust in the Amazon Basin, *Tellus*, 44B, 133-149, 1992.

Talbot, R. W., Andreae, M. O., Berresheim, H., Artaxo, P., Garstang, M., Harriss, R. C., Beecher, K. M., and Li, S. M.: Aerosol chemistry during the wet season in Central Amazonia: The influence of long-range transport, *J. Geophys. Res.*, 95, 16,955-16,969, 1990.

ten Brink, H. M.: Reactive uptake of HNO_3 and H_2SO_4 in sea-salt (NaCl) particles, *J. Aerosol Sci.*, 29, 57-64, 10.1016/s0021-8502(97)00460-6, 1998.

Tobo, Y., Zhang, D., Matsuki, A., and Iwasaka, Y.: Asian dust particles converted into aqueous droplets under remote marine atmospheric conditions, *Proc. Natl. Acad. Sci. USA*, 96, 3396–3403, 2010.

Tobo, Y., Prenni, A. J., DeMott, P. J., Huffman, J. A., McCluskey, C. S., Tian, G., Pöhlker, C., Pöschl, U., and Kreidenweis, S. M.: Biological aerosol particles as a key determinant of ice

- nuclei populations in a forest ecosystem, *J. Geophys. Res.*, 118, 10100-10110, doi:10.1002/jgrd.50801, 2013.
- Vekemans, B., Janssens, K., Vincze, L., Adams, F., and Van Espen, P.: Analysis of X-ray spectra by iterative least squares (AXIL): New developments, *X-Ray Spectrom.* 23, 278-285, 1994.
- Wang, A., Freeman, J. J., Jolliff, B. L., and Chou I-M.: Sulfates on Mars: A systematic Raman spectroscopic study of hydration states of magnesium sulfates, *Geochim. Cosmochim. Acta*, 70, 611, doi:10.1016/j.gca.2006.05.022, 2006.
- Wang, B., Harder, T. H., Kelly, S. T., Piens, D. S., China, S., Kovarik, L., Keiluweit, M., Arey, B. W., Gilles, M. K., and Laskin, A.: Airborne soil organic particles generated by precipitation, *Nature Geosci.* 9, 433–437, doi:10.1038/ngeo2705, 2016.
- Worobiec, A., de Hoog, J., Osan, J., Szaloki, I., Ro, C.-U., and Van Grieken, R.: Thermal stability of beam sensitive atmospheric aerosol particles in electron probe microanalysis at liquid nitrogen temperature, *Spectrochim. Acta B* 58, 479–496, 2003.
- Wright, S. J.: Phenological responses to seasonality in tropical forest plants, 440-460, in *Tropical Forest Plant Ecophysiology*, Mulkey S. S., Chazdon, R. L., and Smith A. P. (Eds), Springer, 1996.
- Yeung, M. C., and Chan, C. K.: Water content and phase transitions in particles of inorganic and organic species and their mixtures using Micro-Raman Spectroscopy, *Aerosol Sci. Technol.*, 4, 269-280, doi:10.1080/02786820903583786, 2010.
- Zhang, R., Khalizov, A. F., Khalizov, J., Zhang, D., Xue, H., and McMurry, P. H.: Variability in morphology, hygroscopicity, and optical properties of soot aerosols during atmospheric processing, *Proc. Natl. Acad. Sci. U. S. A.*, 105, 10,291–10,296, doi:10.1073/pnas.0804860105, 2008.

Figure 1. Location of sampling sites at the Brazilian Amazon basin: an urban site in Manaus (S 3°05.753' W 59°59.419') and a rainforest site at ATTO (S 02°647' W 58°59.992'). Map of South America (top left) with the region marked with a red rectangle and a map of the Amazonas state, Brazil (bottom left) also with the region of interest marked in red.





Figure 2. Ten-day (240 h) backward air mass trajectories at 100 m-, 500 m-, and 1000 m-receptor heights; (a)-(d) for the SA1-SA4 samples collected on April 1 and 16-18, 2012 at the ATTO site and (e)-(g) for the SM1-SM3 samples collected on May 1-3, 2012 at the Manaus site. HYbrid Lagrangian Single-Particle Integrated Trajectory (HYSPLIT) model available at the NOAA Air Resources Laboratory's web server (<http://www.arl.noaa.gov/ready/hysplit4.html>) was used



956

Figure 3. Home-made sample holder for TEM grid samples in SEM/EDX measurements and a typical X-ray spectrum of the TEM grids.




Figure 5. Typical SEM images of aerosol particles for (a) stage 1 ($PM_{0.25-0.5}$) of the SM1 sample, (b) stage 2 ($PM_{0.5-1.0}$) of the SM2 sample, (c) stage 3 (PM_{1-2}) of the SM1 sample, and (d) stage 4 ($PM_{2.0-4.0}$) of the SM3 sample, collected at the Manaus site. For convenience, ammonium sulfate, secondary organic aerosol, aluminosilicates, and reacted sea-salt are denoted as “AS”, “SOA”, “AlSi” and “rss”, respectively.




Figure 6. SEIs, X-ray spectra, and element atomic concentrations of SOA, ammonium sulfate (AS), and mixture particles. The inset images in (b), (c), and (f) show the beam damage on the particles after X-ray measurements.




Figure 7. (a) Raman spectra of standard and airborne ammonium sulfate (AS) particles, which were rescaled for clarity. The inset SEI images are for standard and airborne AS particles where the scale bar is 1 μm . The shoulder peak of SO_4^{2-} at 982 cm^{-1} in the airborne AS particles is from K_2SO_4 ; (b) SEI, optical images, X-ray, ATR-FTIR, and Raman spectra of an overloaded $\text{PM}_{0.25-0.5}$ sample collected at ATTO site on June 10, 2014. X-ray, ATR-FTIR, and Raman spectra indicate that AS, organics, and minerals are the major components of the submicron sample. The sloping baseline in the airborne Raman spectrum is due to the fluorescence from organic compounds.




Figure 8. SEIs, X-ray spectra, and element atomic concentrations of aged mineral dust particles.





Figure 9. Number abundances of reacted sea-salt and aged mineral dust particles containing sulfates (■), nitrates (■), and both (■). The size ranges of each sample is given in the x-axis.

(a) aged mineral dust



(b) reacted sea-salts




Figure 10. SEIs, X-ray spectra, and element atomic concentrations of (a) reacted sea-salt, (b) reacted sea-salt with K-salt, and (c) elongated CaSO_4




Figure 11. SEIs, X-ray spectra, and element atomic concentrations of (a) PBA, (b) soot, (c) tarball, (d) fly ash, (e) Ni-containing, and (f) Fe-containing particles.




Figure 12. Typical SEIs of PBA particles from stage 4 of the (a) SA4 and (b) SM3 samples. PBA and PBA/SOA mixture particles are marked with (\rightarrow) and (+), respectively.




Figure 13. Relative **number** abundance of nine different particle types for the SA1-SA4 and SM1-SM3 samples collected at the ATTO and Manaus sites, respectively.




Table 1. Sampling dates, sampling times, and metrological conditions during the samplings.

Sampling site	Sample	Sampling date	Sampling time	T (°C)	RH (%)	Rainfall (mm)	Sampling conditions
ATTO	SA1	April 1, 2012	12:25 - 14:25	28	72	9	Precipitation during the TEM grid exposure
	SA2	April 16, 2012	12:12 - 13:52	28	74	0	Precipitation (1 mm) in the last 24 hours before the TEM grid exposure
	SA3	April 17, 2012	12:13 - 13:53	30	60	0	No precipitation
	SA4	April 18, 2012	12:04 - 13:45	27	75	0	No precipitation
Manaus	SM1	May 1, 2012	12:17 - 13:57	32	57	0	Precipitation (1 mm) in the last 24 hours before the TEM grid exposure
	SM2	May 2, 2012	12:00 - 13:40	27	73	0	Precipitation (2 mm) in the last 24 hours before the TEM grid exposure
	SM3	May 3, 2012	12:06 - 13:46	24	88	8	Precipitation during the TEM grid exposure