

Single-particle characterization of aerosols collected at a remote site in the Amazonian rainforest and an urban site in Manaus, Brazil

Li Wu¹, Xue Li¹, HyeKyeong Kim¹, Hong Geng², Ricardo H. M. Godoi³, Cybelli G. G. Barbosa³, Ana F. L. Godoi³, Carlos I. Yamamoto⁴, Rodrigo A. F. de Souza⁵, Christopher Pöhlker⁶, Meinrat O. Andreae^{6,7}, and Chul-Un Ro^{1,*}

¹ Department of Chemistry, Inha University, Incheon, 402-751, Korea;

² Institute of Environmental Science, Shanxi University, Taiyuan 030006, China;

³ Environmental Engineering Department, Federal University of Parana-UFPR, Curitiba, PR, Brazil;

⁴ Chemical Engineering Department, Federal University of Paraná-UFPR, Curitiba, PR, Brazil;

⁵ Amazonas State University, Superior School of Technology, Manaus, Amazonas, Brazil;

⁶ Multiphase Chemistry & Biogeochemistry Departments, Max Planck Institute for Chemistry, 55020 Mainz, Germany;

⁷ Department of Geology and Geophysics, King Saud University, Riyadh, Saudi Arabia.

*Corresponding author. Tel.: +82 32 860 7676; fax: +82 32 874 9207
E-mail address: curo@inha.ac.kr (C.-U. Ro)

Figure S1. Typical SEM images of aerosol particles for stage 1 (PM_{0.25-0.5}), stage 2 (PM_{0.5-1.0}), and stage 3 (PM_{1.0-2.0}) samples of SA1 collected on Apr. 1, 2012 at the ATTO site. For convenience, ammonium sulfate, secondary organic aerosol, aluminosilicates, and reacted sea-salt are denoted as “AS”, “SOA”, “AlSi” and “rss”, respectively.




Figure S2. Typical SEM images of aerosol particles for stage 1 (PM_{0.25-0.5}), stage 2 (PM_{0.5-1.0}), and stage 3 (PM_{1.0-2.0}) samples of SA2 collected on Apr. 16, 2012 at the ATTO site.




Figure S3. Typical SEM images of aerosol particles for stage 1 (PM_{0.25-0.5}), stage 2 (PM_{0.5-1.0}), and stage 3 (PM_{1.0-2.0}) samples of SA3 collected on Apr. 17, 2012 at the ATTO site.




Figure S4. Typical SEM images of aerosol particles for stage 1 ($PM_{0.25-0.5}$), stage 2 ($PM_{0.5-1.0}$), stage 3 ($PM_{1.0-2.0}$) and stage 4 ($PM_{2.0-4.0}$) samples of SA4 collected on Apr. 18, 2012 at the ATTO site.




Figure S5. Typical SEM images of aerosol particles for stage 1 (PM_{0.25-0.5}), stage 2 (PM_{0.5-1.0}), and stage 3 (PM_{1.0-2.0}) samples of SM1 collected on May. 1, 2012 at the Manaus site.




Figure S7. Typical SEM images of aerosol particles for the stage 1 (PM_{0.25-0.5}), stage 2 (PM_{0.5-1.0}), and stage 3 (PM_{1.0-2.0}) samples of SM3 collected on May. 3, 2012 at the Manaus site.

