

Response to Referees' Comments

Peter Rayner, Anna M. Michalak and Frédéric Chevallier

June 10, 2019

We thank the three anonymous referees for their comments which have allowed us to clarify various points in the paper. There are some common points made by several referees. We will deal with these first as general comments then with particular comments from each referee. We place referees' comments in typewriter font and our replies in Roman

General Comments

There is a general concern from all three reviewers about the purpose and target audience of the paper. The critiques diverge e.g. (paraphrasing) "The paper is not rigorous enough" (R1) or "Section 2.2 is too general" (R3). Unfortunately, as authors, we cannot assume from this divergence that we have the balance about right. It could well be that the "sweet spot" we seek does not exist. Obviously we do not think so and our experience in teaching, both at the summer school that spawned this special issue and more generally supports this view. The question of mathematical rigour is particularly difficult. The usual route to a well-grounded description of probability and its manipulation is via an undergraduate course in measure theory. Very few likely readers of this paper will have undertaken such a course. Reviews or even text on data assimilation do not generally dedicate the space to provide this level of background. We have instead started from a few precepts:

1. The basic concepts are rather intuitive if demonstrated carefully;
2. The best starting point is the axioms of probability, Bayes Theorem is rather a consequence of these. This approach mirrors both Jaynes and Bretthorst (2003) and Tarantola (2005). We currently note this in Section 3 and have added a comment to the introduction.
3. there really is no distinction in principle between problems given names like data assimilation or inversion. As such we very much thank the reviewers for pointing out inconsistencies in language. We have corrected these and expanded on the problem description in the introduction to clarify these relationships.

The other general critique touches the role of the paper as a review, tutorial or perspective. Reviewers have suggested we should survey the current state of the art and describe future directions. This does not fulfil the paper's role within the special issue.

The other papers in the issue do precisely this for their fields, usually by presenting a state-of-the-art example. There seems little value in summarising those results here. We do, however, accept that more forward-looking material on the general problem is warranted and have expanded the relevant section.

Specific Comments

Reviewer I

First we have dealt with the general comments above although the reviewer makes some points as part of these to which we respond directly.

For example, the term "target variable" is used on page 5, line 13 without any prior definition. We have added text defining this and strengthening the link between the definition of events and PDFs.

Furthermore, the manuscript makes references to "the inversion (or inverse) problem" many times without explaining what the inverse problem is (it is only mentioned briefly in Introduction on page 1, line 16). We have switched to the term "data assimilation" or "assimilation" throughout and added text in the introduction on the equivalence of the various names used in the literature.

For example, a schematic of the data assimilation process and the different components would be helpful and could be referred to when describing the different parts of the system. This is an idea we have thought about during the preparation of the paper. Unfortunately there seems so little commonality in data flow among the various implementations of the theory. For example, highly efficient methods for calculating MAP estimates might use the adjoint of a cost function derived from some exponential PDF. They may never calculate a likelihood or probability. Other techniques that map or sample the posterior PDF may only calculate likelihoods or probabilities. Any form of the diagram would need so many variants that it would add confusion rather than clarity.

I also think some concrete examples when describing the different implementations would help readers to follow the whole process better. We have already noted that the article forms part of a special issue. The task of producing a new textbook in this area (essentially updating the 2000 AGU Monograph) is well beyond our scope here.

2. I think the manuscript would be clearer if the authors choose one perspective from the outset (e.g. data assimilation) and present all material from this perspective in a consistent manner. Currently the manuscript seems to borrow many terms from data assimilation, but frequently, especially in later parts, the language switches to that of inverse modelling (e.g. page 10, line 13: "We need therefore to incorporate these extra variables into the inversion process"; there are many other parts where "inverse problem" and "inversion" are mentioned).

Section 6 is even named "Solving the Inverse Problem". I recommend to replace all occurrences of "inversion", "inverse problem" etc. with terms that are more common in data assimilation. **see general comments above.**

3. Data assimilation is probably best known for its applications in numerical weather prediction, where the technique is used mainly to improve meteorological initial conditions to produce better weather forecasts. In biogeochemistry, on the other hand, data assimilation (and related techniques) are more commonly used to constrain parameters. This difference is alluded to in the manuscript (e.g. page 15, lines 3-10). However, I think it would be better if this distinction is explicitly stated in the beginning of the manuscript. This difference explains e.g. why the manuscript does not focus on the dynamical model (the dynamical model for the target variables in biogeochemical applications is often unknown or assumed to be persistence, while the forecast model is an essential component in atmospheric data assimilation). A broader discussion about the choice of assimilation time window would also be helpful. **We do not accept the distinction drawn here between data assimilation as used to estimate different target variables, indeed one purpose of the paper is to demonstrate that the distinction is superficial. We now argue in an expanded section on future developments that so-called "dual state" approaches that seek to constrain the state and parameters of the model should be more widely used in biogeochemical problems. We have added further commentary on assimilation windows at the end of Section 7.**

4. The authors write that "we will not be using mathematically precise language" (page 2, line 17). I can see where the authors are coming from, but I think this does a disservice to the readers. I recommend the authors to remove this sentence and to be mathematically rigorous to the extent feasible. I understand a mathematical precise language will take away some of the simplicity, but I think the benefits outweigh the added complexity; currently the reader may be left wondering where the language is imprecise and be less likely to refer back to the text when e.g. implementing a data assimilation system. **See general comments above.**

5. The manuscript does not talk much about the issues of ill-posed problems, which are common in inverse problems, and the need for regularization (except for under "Historical Overview", page 19). I consider the use of prior error covariances, e.g. errors with a specified correlation length scale, to be a form of regularization. Even if the "true" error correlation length scales are smaller, a larger correlation length scale may be necessary for the data assimilation system to converge to a solution. This constraint will on the other hand lead to larger aggregation errors. It may also be worth to add a discussion

about how the number of observations influences the design of the data assimilation system, e.g. the choice of regularization. The term "ill-posed" has several meanings in mathematics, here we think the author means problems that are not numerically well-behaved, e.g. with poor convergence. We have dealt with the use of other regularisation methods in the historical overview. We do not know of cases where deliberately incorrect choices were made for prior uncertainties in order to facilitate convergence.

Specific comments

1. Page 3, lines 23-25: "As a practical example a frequentist may estimate a mean by averaging his sample while a Bayesian may calculate an integral over her probability density." I do not think this is a good example of the difference between Bayesian and frequentist statisticians. A better example could illustrate how Bayesian and frequentists interpret probabilities (e.g., a frequentist may only consider the long-term frequency of occurrence of a random event, while a Bayesian may draw from other prior information to assign probabilities, even for non-repeatable events). The reviewer's example is correct, this *is* a difference between the two approaches but we don't think it the most important one. We think the point important enough to defend. It is well-stated in the preface to Jaynes and Bretthorst (2003) commencing on p.xxii. Jaynes points out here that the machinery we use allows the use of prior information but its most important distinction is more fundamental, working with the PDFs themselves rather than estimators pre-derived from assumed distributions.

2. Page 4, line 2: "we have followed it [the notation of Ide et al., 1997] here". It would be helpful to the reader to highlight exactly what is new with the notation introduced in this manuscript. Is it simply an extension of the Ide et al. (1997) notation for biogeochemistry data assimilation? Is it a generalization? More about this in the next point. We have added "as closely as possible" to indicate we changed nothing substantial from Ide et al. (1997).

3.1. Consider adding a "Remarks" column and note when e.g. a notation differs from the notation in Ide et al. (1997). We have done this by separating the mathematical definition and textual description and expanding the latter where necessary.

3.2. For the definition of G , should μ and U be bold to show that they are a vector and a matrix, respectively? In that case also change "mean μ and covariance U " to "means μ and covariances U ". We have changed the typeface. The pluralisation is not correct, the mean of a multivariate distribution is still a mean (singular).

3.3. Descriptions of superscripts "a" and "b": "Posterior or analysis" and "Background or prior". Change to "Analysis or posterior" (to be consistent with "Background or prior").

Done.

3.4 Symbol Q: From my understanding Q is often used to denote model uncertainty (for the dynamical model). "Forecast uncertainty" here seems to include uncertainties due to initial conditions and boundary conditions. This is a good point. Q should refer only to forecast uncertainty. This is too subtle a point to make in the table so we have referred to the relevant section on the Kalman Filter and spelled this out more clearly there.

3.5. Description of R: Add "(Observation uncertainty)" or something similar. Added.

3.6. Symbol A: I do not think I have seen "A" used for "posterior uncertainty covariance" before. Maybe use \mathbf{P}^b for background uncertainty covariance and \mathbf{P}^a for analysis uncertainty covariance. We considered this but it generates other inconsistencies, we should, for example, use the same superscripts for the MAP as well as the uncertainty etc. We believe adding one new symbol to Ide et al. (1997) notation is a better solution. This notation is common throughout NWP (e.g. Desroziers et al., 2005).

4. Page 6, Figure 1: It took some time for me to interpret this schematic. It may be helpful to mention that the observation operator in this case is a simple 1:1 mapping to the system state. An excellent idea, done.

5. Page 10, lines 6-7: "then the dynamical model forms part of the mapping between the unknowns and the observations so is properly considered part of the observation operator". I know that this notation is common in e.g. atmospheric inversion, but I personally think it is unfortunate and easily leads to confusion (as the authors also note on page 9, lines 18-19, data uncertainty or data error may refer to the uncertainty due to errors in both observations and the observation operator, which is misleading). In e.g. 4D-Var for atmospheric data assimilation, I believe the observation operator and dynamical model are usually kept separate in the formulation of the cost function. Given that this manuscript focuses on clarity and fundamentals of data assimilation, I think it would wise to adopt the notation of atmospheric data assimilation and not conflate the observation operator and dynamical model. The reviewer is addressing two different problems here. The first is the role of dynamical models and observation operators. We disagree that the dynamical model error is treated explicitly in conventional 4dvar for numerical weather prediction. It is usually embedded in the background error. We wish to emphasise that the distinction is one of use rather than inherent and we have added a sentence to make this point explicit. The other question about observation and model errors is a different problem which we treat elsewhere. The dynamical model applies to the model state variable (see Eq (1) of Ide et al.). In the case of atmospheric inversion, the dynamical model is sandwiched between operations that are either upstream (interpolation, multiplication with a flux pattern, injection, etc.) or downstream (unit changes, spatial sampling) of it because the target variables \mathbf{x} are not the model state variables \mathbf{z} . For this application, it is

then appropriate to call "observation operator" the full operation chain, including the dynamical model. It is also important to account for the uncertainty in the dynamical model in the data error budget.

6. Page 10, lines 12-13: "Frequently we regard these [parameters] as fixed, which is likely to underestimate the uncertainty of the estimates. We need therefore to incorporate these extra variables into the inversion process." I think "need to" is a bit too strong; maybe say instead that some methods incorporate these extra variables. **Agreed, changed.**

7. Page 20, lines 16-19: The authors mention that the time delay in EnKF may be problematic for tracers that live longer than the assimilation window. Is this not a common problem for all implementations? **Yes, but it's particularly severe for the KF which often uses short, rolling windows. We have clarified.**

Technical Corrections

1. Page 1, footnote 1: I find the footnote unnecessary and suggest to put this information directly in the text. **We disagree. Footnotes are for text which a reader might use but which would disturb the flow if included in the body. This seems to fit that description.**

2. Page 5, line 11: "by an observation operator". May be worth mentioning that this operator is sometimes also referred to as the forward operator., **agreed, done.**

3. Page 5, line 13: "target variable". Define the term. **Now done in the introductory section.**

4. Page 7, line 18: "to find variables to which it is sensitive". Replace "it" with "the quantity of interest" or something similar. **done.**

5. Page 8, line 10: Capitalize "This". **done.**

6. Page 8, line 10: Fix citation format (no parentheses). **done.**

7. Page 13, line 14-15: "These estimates will generally yield larger variability than that from our most likely flux". I do not believe "flux" is defined in this context. Maybe change to "realisation". **done.**

8. Page 14, line 23: "superscript f indicates the application of the forward model". I believe the authors are referring to the dynamical or forecast model here. The forward model is, from my understanding, often synonymous with the observation operator. This is indeed potentially confusing, we have replaced "forward" with "dynamical" and introduced the mnemonic "forecast" which is the meaning we intend for the superscript

9. Page 15, line 4: "For data assimilation our motivation is to hindcast the state of the system". Consider changing to "For data assimilation applied to biogeochemistry our motivation

is often to hindcast the state of the system", or something similar. **done as suggested.**

10. Page 16, line 30: "Another advantage is that, ...". It is not clear from the previous sentences that the previous statement (need to run a dynamical model for each realization of the ensemble) is an advantage. Maybe change to "An advantage of EnKF is that, ..." **The real problem was with the previous sentence where we had not clarified the advantage of the EnKF, now done.**

11. Page 19, line 7: Capitalize "possibilities". **done.**

12. Page 19, line 26: "(references (e.g. Gloor et al., 2000;". Remove "(references" or "(e.g.". **removed "references".**

13. Page 20, line 16: Change format of citations (no parentheses). **done.**

Reviewer II

We have responded to most of the general comments in our overall response. Some specific comments are addressed below:

The reviewer remarked that most of the literature cited was from the authors' own work. We don't agree with that critique but also note that, in a small research community, the authors are three active and longstanding researchers with broad collaborations. A strong representation of work which includes them is unsurprising.

General Comments

1. I was not convinced that that all of the methods in widespread use within the field are special cases of the underlying Bayesian formalism. The manuscript often switches between Bayesian methods not found within biogeochemistry but found within atmospheric sciences more generally (e.g. particle filtering), hybrid Bayesian methods (e.g. Michalak et al. 2005) and non-Bayesian methods (e.g. Manning et al. 2011) with little distinction between them. **We acknowledge that there is a trade-off between comprehensiveness and methodological purity but think some of the reviewer's comments here are misplaced. Particle filters and hierarchical methods are firstly Bayesian (even if not all implementations are complete) and in growing use within biogeochemistry. To ignore studies like Manning et al. (2011) risks a false impression of complete methodological unity which is not the case. We would argue that "all of the methods in widespread use ..." is an accurate description at the moment.**

2. Section 3 has huge potential but does not deliver as one of the main contributions listed in the abstract. The notation needs improving rather than simply reiterating. For example, I do not agree that the notation is sufficient for most practical cases as it is neither followed throughout or sufficient for a tutorial. An example is the discussion of hyperparameters there is no notation available in Table 1

to represent a vector of hyperparameters (I would suggest a bold θ). I think that this is a good opportunity for an explicit notation for the MAP estimate vs the mean. The notation in Table 1 is not precise enough. For example, some of the notations are specific to Gaussian distributions. This would be fine if Table 1 were being used to only discuss a special case of Gaussian problems, but later in the manuscript, non-Gaussian formulations are also discussed, which would require a new notation. The reviewer makes two points here, first that the notation isn't good enough and second that we don't follow it throughout. We reiterate the general point that inventing our own notation which we thought perfect would be at best wasted effort and (more likely) add further confusion. The reviewer raises two examples of missing notation. The vector of hyperparameters and the maximum a posteriori (MAP) estimate. We have added θ to Table 1 and a brief mathematical development in the relevant section. We resist the idea of a special notation for the MAP. It is a useful quantity to know but it is often neither a parameter of the PDF and, for sampling approaches, not generated by the method. We believe referring to it as the mode of the posterior PDF is both accurate and parsimonious of notation. The reviewer has not cited a case where we do not follow our own notation. For example, we have a notation for Gaussian distributions and go on to use it being careful to state it is a special though widely-used case. The rest of the methodology we develop using the notation for probabilities we define.

3. Section 4. As this section is fundamental Bayesian theory and if this is a tutorial, there should be a rudimentary explanation of the notation of $p(A \text{ given } B)$ and how (in basic terms) this forms Bayes theorem. It then worth reserving the notation $p(x)$ for the prior probability of x . Following Jaynes and Bretthorst (2003) and Tarantola (2005) we do not believe that Bayes Theorem is fundamental so do not introduce the topic this way. We do introduce the likelihood later.

4. Section 5.6. Here is a good place to introduce the new notation suggested in Section 3 and to expand the definition of Bayes theorem from Section 4. The references used as examples are not conducive to the narrative thus far. The paper Michalak et al. (2005) does not integrate out the hyper-parameters and is therefore this is not a hierarchical method but an empirical hierarchical method (see e.g. Statistics for Spatio-Temporal Data by Cressie & Wikle). We have now added a brief mathematical development using θ as suggested. We already noted that Michalak et al. (2005) was a stepping stone and have added the classification suggested.

5. Section 6.2. Gibbs samplers always have higher acceptance rates in that by design, the Gibbs sampler has an acceptance rate of 1. The real gain is not due to sampling from a univariate distribution but sampling from distributions where there is a closed form expression for the conditionals that can be sampled from directly. The disadvantage is that for many situations,

this is not known a priori. It is not clear whether the work adaptive on page 12, line 3, is meant to refer to Adaptive MCMC. Adaptive MCMC methods have a particular meaning, which is not the same as that described here. It would be better to simply state that improved strategies use gradient information while sampling. It is important to note that these methods maintain ergodicity while sampling. **We think this discussion is taking us too far into implementation details. Our solution is rather to cut back on discussion of different sampling strategies so we have removed the detailed discussion of the Metropolis-Hastings method and do not discuss other sampling strategies in any detail.**

6. Section 7.2. This needs to discuss non-Gaussian problems, especially given the focus on Bayesian inference. For example, should this posterior uncertainty represent the highest posterior density region around the mode or an equal weighted probability region around the median? This brings to light the difficulties when quoting the posterior mean for non-Gaussian posteriors. **We have added a sentence to the end of this section.**

7. Section 8. The final sentence on future methods could have much more discussion. At the moment, only one study is referenced but not really discussed, so it reads more like an afterthought. **We have added a new section based on the drivers identified earlier. It is necessarily speculative.**

8. Section 8 reads like a review paper, but it isn't comprehensive enough. I would suggest that if this is meant to be a tutorial, to not try to be a review also, because section 8 could be a whole paper in and of itself. **This section was added at the request of a previous reviewer who thought that historical context would support the rest of the paper. We have added a sentence to the opening paragraph explaining this.**

Specific Comments

P1 Title: Data and Assimilation should be lower case **done**.

P1 Line 4: . . .for automating part of the process. i.e. the choice of prior distribution is not automated. **This is probably line 14. Replaced with "building and using algorithms for ..."**

Page 1 Line 12-14: An idea is introduced here but the readers are left hanging. Why introduce this example of improving a model but not testing it, but not provide any explanation of what is meant here. **Explanation added.**

Page 2 Line 5: awkward wording demonstration how these many methods are its implementations **changed to "these methods are implementations of that theory."**

Page 2 Lines 21-23: Why are only these few papers referenced? There is a wealth of literature on applications of the theory to different fields and it is unclear why the authors select only four to represent their fields. **We wanted presentations that**

were extensive and in closely-related fields. This critique risks being open-ended; there is always one more paper.

P3 Line 7 (Eq. 1): Define ξ . ξ plays the role of an integration variable which are customarily not defined.

Section 2.2 capitalize non-Bayesian done.

Page 3 Line 19: Explain what is meant by the Replicate Earth Paradigm We believe the explanation would be too long and take us too far afield, we have deleted the reference.

Section 2.2 should come before Section 2.1? This narrows the remainder of the discussion to Bayesian. We think the introduction of events and probabilities should precede the bifurcation between Bayesian and non-Bayesian approaches so have not made this change.

P4 Table 1: This isnt referred to anywhere in the text. Delete extra brackets in description of R. Have added sentence pointing out that symbols used throughout are defined in Table 1, also removed extra brackets.

P5 Line 20 (Eq 20): Make clear that the earlier reference to one variable, one observation has now been expanded into vectors. done

Figure 1. Put all text into the caption below. Why is the x axis labelled unknown in the bottom panel? Label panels (a) and (b). The numbers 1.2 and 0.8, 0.2, -0.2 are not immediately clear what they are showing. We have repositioned the captions and explained the values further.

Page 5 Line 26-27: describe what Equation 2 not well applied means This referred to poor approximations in the solution but this is not the best place to discuss that so we removed the reference.

Page 7 Line 5: Need to describe what you mean what are the common misunderstandings We explained these below so have combined the paragraphs and explained the misunderstandings explicitly.

Page 7 Line 10: Reason 1: Why does limiting target variables underestimate the uncertainty? One might remove an important variable. We have explained this.

Reason 2: This is just repeating what was said above and is not a reason. Agree with the repetition, we have reorganised. We have also added that it is not always easy to guess in advance what will be constrained, especially in complex models.

Page 7 Line 15: Walking through an example setting up this ideal world would be much more helpful for a novice reader than only providing the instructions. We have used an example to illustrate the points below, this is an excellent suggestion. We also deleted the last element in the list since that concerns activities after the assimilation.

Page 7 Line 15: Point 1: Explain what this means. How do you make this decision? The choice is motivated by the science question we are addressing. We have added examples.

Point 3: An example would help. have added an example.

Point 4: What cut-off? **We have lengthened the explanation.**

Page 8 Line 9: K can be either sign so why would s_K be more likely to increase rather than decrease? i.e. a larger s means a more negative s_K if K were negative. I may be misunderstanding what is being said here, but isn't it simpler to say that positive scaling has a minimum of 0 so skewed in one direction while $\log(s)$ can be both positive and negative? **Two points here. First on k the reviewer is correct, we have replaced "value" with "magnitude". We also adopted the second suggestion.**

P8 Line 24: The text does not describe what a uniform prior is (i.e. a uniform distribution). If this is for a novice, needs to be explicit. **We have introduced this on page 5.**

Page 8 Line 19: Discussion about aggregation errors missing. No reference to methods that try to diagnose these (e.g., Turner et al., 2015, Lunt et al., 2016) **We don't think this is the right place to discuss aggregation errors which usually occur with the choice of target variables rather than prior PDF. We have added some more discussion in Section 5.1 including these references.**

Page 8 Line 26: Another disadvantage is that it is not fully hierarchical and that hyper-parameters are not integrated out. **This is true, we discuss it explicitly later.**

Section 5.2. Need a discussion about the uncertainties assumed in the prior as this makes a very large impact. In practice, this is not well known. This could be a practical note on the application to biogeochemistry. **A good point, we have added a paragraph.**

Page 9 Line 11: It is not only resolution that affects H but model structures such as parameterizations. **A good point, we had assumed it but now added an explicit comment.**

Section 5.4. Need much more detail about how uncertainties are treated in H . **We have added a paragraph. We also note this in Section 8 since it is a weakness in current work.**

Page 10 Line 25: Need to state that this is not a fully hierarchical Bayesian method if hyperparameters are not integrated out, and state that the impact is likely an underestimation of uncertainties **Added.**

Page 10 Line 29: This study did integrate hyper-parameters in a hierarchical sense and propagated these uncertainties through to fluxes, but requires MCMC calculations with potentially higher computational cost. **comment added.**

Page 10 line 32: What is a well-known atmospheric inversion? **TRANSCOM, now described in detail.**

Page 11, line 1: If this is a tutorial, then the description and notation should be included in this paper and not just referenced to another paper. It's not clear what the reviewer is referring to here. If it is Wikle and Berliner (2007) then we have essentially done this in the

expanded section with the solution for θ sketched. We would also argue that the general formalism we have used already embodies the hierarchical method. One simply augments the target variables with θ , solves for the joint PDF of the augmented target variables and integrates out whatever one doesn't want.

Page 11 Line 12: A new altered Figure 2 would help show this point We think the textual description is sufficient and that the space for a new figure is not warranted.

Page 11 Line 13: Delete various of done.

Page 12 Line 19 Can talk about the role of thinning chains.

Added description.

Page 12 Line 5- Need references using MCMC We have cut back on the description of various Monte Carlo methods so believe the depth of referencing is sufficient.

Page 12 Line 5 - Need to discuss limitations of MCMC such as convergence issues. Again, we have added a comment that one must watch the sampling properties of whatever method one uses but have avoided technical detail of any particular method.

P13 Line 3: It isn't clear what the word model means here. For a tutorial, an example of what is meant would be helpful. Should have been "observation operator", now changed.

Page 13 Line 5 typo MCMC? changed.

Page 13 Line 16 Needs more explanation such as? The sentence was meant to point to the section immediately below, we have made this explicit.

P13 Line 27: The notation $G(H(x) - y^o, R)$ in this section is different to that defined in Table 1 and does not appear in Ide et al (1997). We have corrected the form to match the definition of G in Table 1.

P13 Line 29 (Eq. 5): I would change $p(x)$ to something else here, e.g., $p(x|y)$, to distinguish it from the prior probability of x done.

P14 Line 16: Specify where this will be described rather than simply later. done.

P15 Line 20 (Eq. 7): Is the meaning of J described before this? What is J? What does it mean? Explain for a beginner. J is actually defined by that equation, we have now added text ahead explaining this.

Section 8: Reserve the term we for subjective choices made specifically by the authors rather than the community in general, as e.g. We can identify..., We now know. . . etc. corrected throughout.

P 19 Line 23: Capital T needed at start of sentence. corrected.

P21 Line 10: Delete apparently done.

Reviewer III

We refer to our general responses to the reviewer's overall comments.

General Comments

1) Section 2.2: Since the paper focuses on Bayesian approach, I don't think the paper needs to discuss non-Bayesian method. We don't discuss it in detail but think readers need to be aware of the difference, especially since Bayesian methods are still regarded as more obscure.

2) Section 3: It is not clear to me what is the difference between x (Target variables for assimilation) and z (model state variable). It is not explained in the paper. Later on, only x is used in the cost function. We have added a paragraph in Section 5.1 to clarify this distinction.

3) Section 5: this section has a lot of useful materials. I would suggest discussing how each element is addressed in specific examples. Section 5.1 gave a recipe to decide target variables. It would be easier to understand if this recipe is discussed within a specific application. Reviewer II also suggested this and we have added an example.

4) Section 6.5: The Kalman filter is discussed in a very general concept here. How Kalman filter has been used in biogeochemistry, and what is the challenge? We have addressed the history in Section 7.

5) Section 7 discussed specific Bayesian methods from computation perspective. Again, I found the description is too general. The applications of these methods in specific problem could be very different. For example, in boundary condition estimation, the prior ensembles may not come from the posterior ensemble of previous step, since there is no dynamical model to propagate information forward. As a result, step 5 described in section 7.4 is not applicable. If there is no dynamical model for the target variables one might question whether the KF or any of its derivatives is a good choice for the assimilation algorithm. As we note later, even algorithms that did not seem to be using a dynamical model were often assuming a persistence model.

References

- Desroziers, G., Berre, L., Chapnik, B., and Poli, P.: Diagnosis of observation, background and analysis-error statistics in observation space, *Quarterly Journal of the Royal Meteorological Society*, 131, 3385–3396, <https://doi.org/10.1256/qj.05.108>, URL <http://dx.doi.org/10.1256/qj.05.108>, 2005.
- Ide, K., Courtier, P., Ghil, M., and Lorenc, A. C.: *Unified Notation for Data Assimilation: Operational, Sequential and Variational*, 1997.
- Jaynes, E. and Bretthorst, G.: *Probability Theory: The Logic of Science*, Cambridge University Press, URL <http://books.google.com.au/books?id=tTN4HuUNXjgC>, 2003.

- Manning, A., O'Doherty, S., Jones, A., Simmonds, P., and Derwent, R.: Estimating UK methane and nitrous oxide emissions from 1990 to 2007 using an inversion modeling approach, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- Michalak, A. M., Hirsch, A., Bruhwiler, L., Gurney, K. R., Peters, W., and Tans, P. P.: Maximum likelihood estimation of covariance parameters for Bayesian atmospheric trace gas surface flux inversions, *J. Geophys. Res.*, 110, D24 107, <https://doi.org/10.1029/2005JD005970>, 2005.
- Tarantola, A.: *Inverse problem theory and methods for model parameter estimation*, siam, 2005.
- Wikle, C. K. and Berliner, L. M.: A Bayesian tutorial for data assimilation, *Physica D: Nonlinear Phenomena*, 230, 1–16, 2007.

Fundamentals of ~~Data Assimilation~~ data assimilation applied to biogeochemistry

Peter J. Rayner¹, Anna M. Michalak², and Frédéric Chevallier³

¹School of Earth Sciences, University of Melbourne, Melbourne, Australia

²Department of Global Ecology, Carnegie Institution for Science, Stanford, USA

³Laboratoire des Sciences du Climat et de l'Environnement, Gif sur Yvette, France

Correspondence: Peter Rayner (prayner@unimelb.edu.au)

Abstract. This article lays out the fundamentals of data assimilation as used in biogeochemistry. It demonstrates that all of the methods in widespread use within the field are special cases of the underlying Bayesian formalism. Methods differ in the assumptions they make and information they provide on the probability distributions used in Bayesian calculations. It thus provides a basis for comparison and choice among these methods. It also provides a standardised notation for the various quantities used in the field.

1 Introduction

The task of improving current knowledge by the consideration of observed phenomena is a fundamental part of the scientific process. The mechanics of the process are a matter for historians of science and a matter of fierce debate, while its fundamental reliability is a matter for philosophers, literally a meta-physical question. Here we are concerned with ~~algorithms for automating~~ building and using algorithms for the process. The step by step construction of such algorithms from agreed upon logical premises is beyond our scope here but is masterfully presented by Jaynes and Bretthorst (2003).

It is important to note at the outset the generality of the machinery we will present, especially because many of the common objections to the methods spring from misunderstandings of this generality. For example, it is a common complaint that data assimilation can improve a model but cannot test it. The methods are, however, capable of considering multiple models and assigning probabilities to them; a test of their relative performance.

At the outset we should settle some questions of terminology. The phrases “data assimilation”, “parameter estimation”, “inverse modelling” and “model-data fusion” are used with overlapping meanings in the literature. ~~Here we will draw no distinction between them.~~ “Inversion” is most commonly used to describe improving knowledge of the inputs of a model given observations of its output. The term arises since the technique reverses the normal direction of causality (Enting, 2002, p.10). “Data assimilation” most commonly refers to the correction of the trajectory of a running model given observations. There are many exceptions to this categorisation and other distinctions could be made. We will use the term data assimilation throughout but readers should interpret the name broadly.

We are concerned with algorithms to improve the performance of models through the use of data. The improvement will come via knowledge of the state (quantities the model calculates), boundary conditions (quantities we insert into the model)

or structure of the model. Performance will be assessed by the model's ability to reproduce observations, especially those not included in the data assimilation algorithm.

The application of data assimilation to biogeochemistry has proceeded from informality to formal methods and from small to larger problems. One can see the study of Fung et al. (1987) as a data assimilation experiment. These authors adjusted
5 respiration parameters in a simple biosphere model so that the seasonal cycle of atmospheric CO₂ concentration better matched the observed seasonal cycle at some stations. Fifteen years later, Kaminski et al. (2002) performed a similar study on a similar model with more stations and a formal algorithm. We see similar developments in the estimation of CO₂ fluxes (e.g. Tans et al. (1990) vs Chevallier et al. (2010)) and ocean productivity (Balkanski et al. (1999) cf Brasseur et al. (2009)).

The flourishing of the general approach has also led to a great diversification of methods, reflecting repeated borrowings
10 from other fields. This has been extremely fruitful, but can be confusing for a novice. Our aim in this paper is to reintroduce the fundamental theory and demonstrate how these ~~many methods are its implementations~~ methods are implementations of that theory. We will aim to tread a careful path between generality and simplicity. If we succeed, a reader should be well-placed to understand the relationships among the methods and applications presented later. We will also introduce a consistent notation that is sufficiently general as to capture the range of possible applications.

15 The structure of the paper is as follows: In Section 2 we introduce the general theory. There is no new material here, but we need a basis for the subsequent discussion. Section 3 introduces a consistent notation. Section 4 presents an overall approach to the solution to the data assimilation problem through the use of a simple example. Section 5 describes the construction of inputs to the problem. Section 6 describes solution methods and how they are special cases of the general theory but take advantage of the circumstances of a particular problem. Section 7 describes some computational aspects related to the solution of data
20 assimilation problems. Finally, Section 8 introduces a range of example applications which illustrate the historical development of the field.

2 Data Assimilation as Statistical Inference

In what follows we will not be using mathematically precise language. We think the trade-off of ambiguity for simplicity suits our introductory task. Some of the key concepts are introduced in Tarantola and Valette (1982) and elaborated further
25 in Tarantola (1987). In addition, Evans and Stark (2002) sought to unify the languages of applied mathematics and statistics used to describe the methods. Jaynes and Bretthorst (2003) gave a thorough exposition on the underlying ideas while Tarantola (2005) provided a comprehensive treatment of both motivations and methods. Other presentations that focus on particular applications include Kalnay (2003) for numerical weather prediction, Rodgers (2000) for atmospheric remote sensing, Enting (2002) and Bocquet et al. (2015) for atmospheric composition and Carrassi et al. (2018) for the geosciences more generally.

30 2.1 Events and Probability

For our purposes, we define an event as a statement about the condition of a system. This is deliberately general; such statements can take many forms. Examples include categorical or discrete statements (e.g. "including isopycnal mixing improves ocean

circulation models”), logical propositions (e.g. “increasing soil temperatures lead to increased soil respiration”) or quantitative statements like “the Amazon forest is a sink of between 1 and 2 PgCyr⁻¹”. The choice of the set of events we consider is the first one we make in setting up any data assimilation problem. We require that any events that are logically incompatible be disjoint (mutually exclusive) and that the set of events be complete.¹

5 The concept of probability is so simple and universal that it is hard to find a definition that is more than a tautology. It is a function mapping the set of events onto the interval [0, 1]. Its simplest properties, often called the Kolmogorov axioms (Jaynes and Bretthorst, 2003, Appendix A) reflect the definition of events, i.e that the probability of the whole domain is 1, and that the probability of the union of two disjoint events is the sum of their individual probabilities. The most common case in natural science is learning about some numerical quantity like a rate constant or a flux. We term the set of these quantities the “target variables” of the problem. They are also called unknowns, parameters or control variables. In this continuous case, events are usually membership of some set (e.g. $x \in (2, 5)$). If our target variables are continuous, we can define a probability density function p (usually abbreviated PDF) such that the probability P that a ~~quantity-target variable~~ x is in some region R is the integral of p over R . In one dimension this simplifies to

$$p(x) = \lim_{\delta x \rightarrow 0} \frac{P(\xi \in (x, x + \delta x))}{\delta x} \quad (1)$$

15 2.2 Bayesian and ~~non-bayesian non-Bayesian~~ inference

At this point, any discussion of statistics bifurcates into two apparently incompatible methodologies, roughly termed Frequentist and Bayesian. Debate between the two schools has been carried on with surprising vigour. See Salsburg (2001) for a general history and Jaynes and Bretthorst (2003) for a partisan view. Characterizing the two schools is far beyond our scope here, especially as we will follow only one of them in much detail, but since even the language of the methods is almost disjoint it is worth introducing the new reader to the major concepts.

Frequentists generally pose problems as the quest for some property of a population. The property must be *estimated* from a sample of the population. *Estimators* are functions that compute these estimates from the sample. The design of estimators with desirable properties (no bias, minimum uncertainty, etc.) is a significant part of the technical apparatus of frequentist inference. Often we seek a physical parameter. This is not a bulk property but a single value that must be deduced from imperfect data. Here we treat each experiment as observing a member of a population so that we can use the same apparatus; ~~the method is summed up in the phrase “the Replicate Earth Paradigm” (Bishop and Abramowitz, 2013).~~

Bayesian statisticians concern themselves with the state of knowledge of a system. They regard the task of inference as improving this state of knowledge. We generate knowledge of some property of the system by applying some mathematical operation to the state of knowledge of the underlying system. Although we may call this estimating some property of the system, the calculations involved are quite different from the estimators of frequentist statistics. As a practical example a frequentist may estimate a mean by averaging his sample while a Bayesian may calculate an integral over her probability density. It is also important to note that for a Bayesian the state of knowledge does not necessarily come only from the sample

¹This definition of event already takes us down one fork of the major distinction in statistics roughly described as Bayesians vs Frequentists. This is a Bayesian definition; a Frequentist will usually refer to an event as the outcome of an experiment.

itself. This use of information external or *prior* to the data is the most important practical difference between the two methods. Throughout this paper we will generally follow a Bayesian rather than frequentist approach. This represents the balance of activity in the field but is not a judgment on their relative merits.

3 Towards a Consistent Notation

5 The notation in use in this field is as varied as the nomenclature. This is unavoidable where similar techniques have been developed independently to answer the needs of many different fields. This diversity complicates the task for novices and experts alike as they compare literature from different fields. Producing a notation at once compact enough to be usable and general enough to cover all cases is a difficult task. Ide et al. (1997) made an attempt focused on numerical weather and ocean prediction. Experience has shown their notation to be sufficient for most practical cases and we have followed it here [as closely](#)
10 [as possible](#). The notation does have an explicitly Bayesian flavour. For example we eschew the use of hats to describe estimates because we regard estimation as an operation on a probability density. [The notation used in this paper and the rest of the issue derives from Table 1 and readers should refer to this table for definitions of symbols.](#)

4 Fundamental Bayesian Theory

In this section we will sketch the fundamental solution of the inverse problem. This will motivate a more detailed description
15 of the various inputs in the following section.

Tarantola (2005) states at the outset of his book that the state of knowledge of a system can be described by a probability distribution or corresponding probability density. Our task is to combine knowledge from measurements with pre-existing knowledge on the system's state to improve knowledge of the system.²

20 The recipe for combining these two sources of knowledge is traditionally introduced via conditional probabilities and expressed as Bayes' Theorem (Laplace, 1774). [Here we follow](#) Tarantola (2005) and Jaynes and Bretthorst (2003) [who](#) derive the machinery directly from the axioms of probability.

We postulate that the true state of the system is consistent with our prior knowledge of it and the measurements we take. The link between the system's state and the observations is created by an observation operator [\(often called a forward operator\)](#).³ Both observations and observation operators are imperfect and we describe the state of knowledge of them both with probability
25 densities.

The case for one target variable and one observation is described in Figure 1, slightly adapted from Rayner (2010). The top panel shows the joint probability distribution for the target variable (blue rectangle) and the measured quantity (red rectangle). Our imperfect knowledge of the mapping between them is shown by the green rectangle. In this simple case the three PDFs are uniform ~~so there is no privileged value for the data or target variable~~ [meaning all values within the interval are equally likely](#).

²Tarantola (2005) uses the term information instead of knowledge. We avoid it here because it has a technical meaning.

³Rayner et al. (2010) refers to this as a model but here we reserve that term for the dynamical evolution of the system.

Table 1. Table of notation used throughout the issue. Borrows heavily from Ide et al. (1997) Appendix 1.

| Basic Notation | | |
|--|---|--|
| Symbol | <u>Definition</u> | Description |
| BOLD | | Matrix |
| bold | | Vector |
| \mathbf{x} | | Target variables for assimilation |
| \mathbf{z} | | Model state variables |
| \mathbf{y} | | Vector of observed variables |
| θ | | <u>hyperparameters (see Section 5.6)</u> |
| J | | Cost function |
| $\mathbf{U}(\mathbf{x}, \mathbf{x}^t)$ | | Uncertainty covariance of \mathbf{x} around some reference point \mathbf{x}^t |
| $\mathbf{C}(\mathbf{x})$ | | Uncertainty correlation of \mathbf{x} |
| $p(x)$ | | Probability density function evaluated at variable x |
| $G(\mathbf{x}, \mu, \mathbf{U})$ | | Multivariate normal (Gaussian) distribution with mean μ and covariance \mathbf{U} |
| H | | Observation operator mapping target or model variables onto observables |
| \mathbf{H} | $\nabla_{\mathbf{x}} \mathbf{y}$ | Jacobian of H , often used in its place, especially for linear problems |
| M | | Model to evolve state vector from one timestep to the next |
| \mathbf{M} | $\nabla_{\mathbf{x}} \mathbf{x}^f$ | Jacobian of M |
| $(\cdot)^a$ | Posterior or analysis | <u>Analysis or posterior</u> |
| $(\cdot)^b$ | | Background or prior |
| $(\cdot)^f$ | | Forecast |
| $(\cdot)^g$ | | (First) guess in iteration |
| $(\cdot)^t$ | | True |
| $(\cdot)^o$ | | Observed |
| δ | | Increment |
| Some Useful Shortcuts | | |
| Symbol | <u>Definition</u> | Description |
| \mathbf{d} | $\mathbf{y} - H(\mathbf{z})$ (Innovation) | <u>Innovation</u> |
| $\mathbf{U}(\mathbf{x})$ | | Uncertainty covariance of \mathbf{x} about its own mean, i.e. the true uncertainty covariance |
| \mathbf{B} | $\mathbf{U}(\mathbf{x}^b)$ (Prior uncertainty covariance) | Prior uncertainty covariance |
| \mathbf{Q} | $\mathbf{U}(\mathbf{x}^f, \mathbf{x}^t)$ (Forecast uncertainty) | <u>Forecast uncertainty (see Section 6.5)</u> |
| \mathbf{R} | $\mathbf{U}((\mathbf{y} - H(\mathbf{z}^t)))$ <u>$\mathbf{U}(\mathbf{y} - H(\mathbf{z}^t))$</u> | <u>Uncertainty of observations around prediction given true state, (often abbreviated observational uncertainty)</u> |
| \mathbf{A} | $\mathbf{U}(\mathbf{x}^a)$ (posterior uncertainty covariance) | posterior uncertainty covariance |

The solution of the problem is the intersection or multiplication of the three PDFs. Adding the requirement that the system state must take a value, we obtain the solution (bottom panel) by projecting the joint PDF onto the x-axis and normalizing it. We write the multiplication as

$$p(\mathbf{x}, \mathbf{y}) \propto p(\mathbf{x}|\mathbf{x}^b) \times p(\mathbf{y}|\mathbf{y}^o) \times p(\mathbf{y}|H(\mathbf{x})) \quad (2)$$

5 where we have extended the case with one variable and one observation to the usual case where both are vectors. We generate the posterior distribution for \mathbf{x} by integrating over possible values of the observed quantity

$$p(\mathbf{x}) \propto \int p(\mathbf{x}, \mathbf{y}) d\mathbf{y} \quad (3)$$

The constant of proportionality follows from normalisation $\int p(\mathbf{x}) d\mathbf{x} = 1$. The last term in Equation 2 is the **uncertainty probability** of a value \mathbf{y} given a value of the target variable \mathbf{x} and is often termed the likelihood. We stress that the generated PDFs are the most complete description of the solution. What we usually think of as the estimated **state-of-the-system-target variables** \mathbf{x}^a is some derived property of $p(x)$, e.g. its mode or mean. Note that if any PDF is mischaracterized **or if Equation 2 is not well-applied**, \mathbf{x}^a can correspond to a very small probability, possibly even smaller than for \mathbf{x}^b . Such a situation may remain unnoticed in the absence of independent direct validation data for \mathbf{x}^a , as is currently the case for global atmospheric inversions. This issue underlines the necessity to keep a clear link with the statistical framework, despite the possible technical complexity of data assimilation systems.

5 The Ingredients


In this section we describe the various inputs to Equation 2 along with some closely related quantities.

5.1 Deciding on Target Variables

20 It is important at the outset to distinguish between target variables \mathbf{x} and state variables \mathbf{z} . Target variables are the set of quantities we wish to learn more about in the assimilation. State variables are the set of quantities calculated by a model. As an example, a chemical transport model (Krol et al., 2005) may have specified inputs of various tracers and will calculate the evolution of those (and other) tracers subject to atmospheric transport and chemical transformation. We will frequently wish to learn about the inputs but not about the concentrations. In that case the inputs are target variables while the concentrations are state variables. We might also choose to learn about the concentrations in the assimilation in which case they become both state and target variables.


25 Although not usually considered part of the problem, the decision on target variables preconditions much of what follows. **It is also the subject of some common misunderstandings and there is some definite guidance for the choice.**

In general the target variables should include anything which is important to the behaviour of the system and is not perfectly known. **The choice should not depend on us guessing which target variables can be constrained by the observations. There are three reasons for this.** A common misunderstanding is to limit the target variables to the smallest possible set. This is risky for three reasons:



(A)

Joint probability distribution for system state (x-axis) and measurement (y-axis). The light-blue rectangle (which extends to infinity in the y direction) represents the prior knowledge of the unknown system state. The light-red rectangle (which extends to infinity in the x direction) represents the knowledge of the true value of the measurement. The green rectangle (extending to infinity along the one-to-one line) represents the state of knowledge of the observation operator. The black triangle (the intersection of the three PDFs) represents the solution as the joint PDF.



(B)

PDF for unknown system state obtained by projecting the black triangle from the left panel onto the x-axis (i.e. integrating over all values of the observation).

Figure 1. Illustration of Bayesian inference for a system with one target variable and one observation. The top panel Part (A) shows the joint probability distribution for target variable (x-axis) and measurement (y-axis). The light-blue rectangle (which extends to infinity in the y direction) represents the prior knowledge of the target variable (uniformly distributed between -0.2 and 0.2). The light-red rectangle (which extends to infinity in the x direction) represents the knowledge of the true value of the measurement (uniformly distributed between 0.8 and 1.2). The green rectangle (extending to infinity along the one-to-one line) represents the state of knowledge of the observation while operator. The observation operator is a simple one-to-one mapping represented by the bottom panel heavy green line. The black triangle (the intersection of the three PDFs) represents the solution as the joint PDF. Part (B) shows the resulting PDF for the target variable variables

1. We often wish to calculate the uncertainty in some predicted quantity. ~~Limiting the number of target variables may underestimate this uncertainty~~; We may inadvertently eliminate a target variable whose uncertainty contributes to that of our quantity of interest.
 2. We are not always capable of guessing in advance which target variables will be well-constrained by the assimilation. We should let the observations determine which variables can be constrained rather than guessing in advance (Wunsch and Minster, 1982);
 3. If a target variable that should be important for explaining an observation is removed, the information from that observation may contaminate other target variables (Tramper and Snieder, 1996; Kaminski et al., 2001). This usually happens when we limit the detail or resolution in our target variables.
- 10 These considerations push us towards complex rather than simple models. A common critique is that we will over-fit the data. This can certainly happen but will be revealed by proper consideration of the posterior uncertainty. We can always post-treat the target variables to seek combinations with lower uncertainty. An example is aggregating a gridded field to lower resolution which is often better constrained by the observations.
- ~~In an ideal world one would~~Point three above has been the subject of much work in the flux inversion community in which
- 15 our target variables are gridded surface fluxes of some atmospheric tracer. Often we group or aggregate grid cells into regions and assume the flux pattern inside these regions is perfectly known. Our target variables comprise multiples or scaling factors for these patterns. Kaminski et al. (2001) showed that incorrect specification of these patterns led to errors in the retrieved multipliers. Following Tramper and Snieder (1996) such errors are known as aggregation errors. Kaminski et al. (2001) proposed an algorithm for deweighting observations most strongly contaminated by aggregation errors. The more common solution is to
- 20 solve at higher resolution (See Section 7.1). Bocquet (2005) pointed out some limitations of this approach with the information from observations being focused more tightly around observing sites as resolution increased. (Rodgers, 2000, Section 3.2.1) also noted an increased impact of the prior as more detail in the target variables meant that more remained unobserved. This has led to various approaches to designing structures for the target variables consistent with the capabilities of the observing system. Wu et al. (2011); Turner and Jacob (2015); Lunt et al. (2016) chose different methods for doing this with Lunt et al. (2016)
- 25 showing how to include the uncertainty in the choice in their assimilation.
- We use an example to illustrate the process. Bodman et al. (2013) used the simple carbon-cycle/climate model MAGICC (Meinshausen et al., 2008) and historical observations of temperature and CO₂ concentration to estimate the global mean temperature and integrated carbon uptake for 2100. The steps involved in choosing which parameters of MAGICC to use as target variables was:
- 30 1. Decide on a quantity of interest. This ~~need~~will be driven by the science question we are addressing, e.g. the global mean temperature in 2100 or the biospheric carbon uptake of North America. Often it will not be a target variable for the assimilation; frequently it is a forecast from the model. It is a function of the target variables for the assimilation target variable but the result of a forecast or diagnostic calculation.

2. Use sensitivity analysis to find variables to which ~~it~~ the quantity of interest is sensitive (e.g. Moore et al., 2012). This may often include forcings that are usually neglected. It is often a brute force calculation performed by changing the variable a small amount though the adjoint analysis discussed in Section 7.1 can also be used.
3. Generate prior PDFs for the most sensitive variables and, with these, calculate the prior PDF in the quantity of interest generated by each potential target variable. ~~These two~~ This and the previous steps should never be separated; the sensitivity to a given target variable is best expressed in uncertainty units for that variable. This provides a common basis for comparing sensitivities of different variables in different units. For example, Bodman et al. (2013) used realisations of the simple climate model MAGICC to calculate the contribution of uncertainties in model parameters (such as climate sensitivity or carbon-cycle feedback parameters) to uncertainty in temperature in 2100.
4. Target the most ~~sensitive variables~~ important variables for our quantity of interest with the cut-off being computationally driven.
5. ~~Calculate the posterior PDF for the quantity of interest. This PDF is generated by propagating the PDFs for the target variable through the function mapping them onto the quantity of interest~~ This led Bodman et al. (2013) to choose a set of eleven target variables.

It is noteworthy how much calculation may be required before we start an assimilation.

5.2 Possible Prior Information

The PDF for the prior $p(\mathbf{x}|\mathbf{x}^b)$ should reflect whatever information we have about the quantity before a measurement is made. This is not always easy given that science is in continual dialogue between existing knowledge and new measurements and it is likely that measurements quite similar to the ones we are about to use have already informed the prior. We should also use any information we have, even if this is only obvious constraints such as positivity. Some of the reluctance to use Bayesian methods comes from the possible impact of the prior PDF on the posterior. This influence can be subtle. For example, if we wish to constrain some physical value K and we approach this by solving for a scaling factor s so that sK appears in our physical problem. We can reasonably expect that s is positive. Choosing a random positive s is far more likely to increase rather than decrease the value magnitude of sK compared to K . ~~Choosing a uniform distribution for $\log(s)$ makes~~ Solving for $\log(s)$ as a target variable means we can make increase or decrease equally likely. ~~this~~ This was first noted by Jeffreys (1939) and is explained in more detail in ~~(Jaynes and Bretthorst, 2003, Section 6.15)~~ Jaynes and Bretthorst (2003) Section 6.15.

In practice, prior information is most often taken from a model providing a mechanistic or process-based representation of the system (e.g. Gurney et al., 2002). The advantage of this approach is that it explicitly incorporates scientific understanding of the functioning of the system into the data assimilation process. Further, model outputs can easily be tailored to the ~~inversion problems, with data being provided for a wide range of temporal or spatial resolutions~~ data assimilation we are performing, e.g. prior information can be provided at whatever time and space resolution we need. The disadvantage is that many alternate representations often exist, and that the choice among them can have a major and scale-dependent influence on the final PDFs

(e.g. Lauvaux et al., 2012; Carouge et al., 2010a), while the relative merits of such alternate priors is difficult to objectively and quantitatively determine when no ground truth is available. The availability of ground truth allows drawing the statistics of model performance, but not necessarily directly at the spatial and temporal scales of the prior information. In this case, one needs to study how the uncertainties of the fluxes change as they are aggregated or disaggregated. These scaling properties are controlled by uncertainty correlations (e.g. Chevallier et al., 2012). The parameters that encapsulate this prior information (such as uncertainty magnitudes or correlation lengths) are also imperfectly known and Section 5.6 describes how they can be incorporated into the formalism. Michalak et al. (2017) describes how they may be diagnosed from the assimilation system.

Alternatives that reduce the reliance on potentially subjective prior information do exist, where less restrictive prior assumptions are made. The case of uniform priors described above is one such example. One practical example in biogeochemical data assimilation is the "geostatistical" approach (e.g. Michalak et al., 2004), where the prior is described as a function of unknown hyperparameters that link ancillary datasets to the state, and where the prior uncertainty covariance is formally described through spatial and temporal correlation analysis that is based as strongly as possible on available observations. The advantage here is the replacement of possibly subjective prior with empirical information (e.g. Gourdji et al., 2012; Miller et al., 2013), while the disadvantage is a more limited formal link to scientific process-based prior understanding of the system.

It is important to note that the various approaches used to construct the prior in data assimilation problems are not disjoint options, but rather can be thought of as points along a continuum of options that reflect different choices about the assumptions that a researcher is willing to make about the system (e.g. Fang et al., 2014; Fang and Michalak, 2015). The choice of prior is an expression of a choice of assumptions. With a careful framing of the specific question to be addressed, and with an explicit examination of the assumptions inherent to different possible approaches, one can design a set of priors that optimally blends existing approaches for a given application.

We also note that often much more attention is paid to the parameters that define the location of the prior PDF (such as the mode) while less attention is paid to its spread (e.g. the variance in the case of Gaussian distributions). The two are equally important as they both control the intersection of PDFs in Figure 1.

5.3 Observations

The PDF for the observation $p(\mathbf{y}|\mathbf{y}^o)$ represents our knowledge of the true value of the measured quantity given the measurement. The characterization of this PDF is the task of metrology discussed by Scholze et al. (2017). An important point to make at this stage is that verification of a measurement means, for our purposes, verification of the PDF (including its mean, i.e. studying possible systematic errors in measurements). That is, it is more important that the PDF we use properly **represent** represents the state of knowledge of the true value, than that the value is as precise as possible.

5.4 Observation Operators

The PDF for the observation operator $p(\mathbf{y}|H(\mathbf{x}))$ represents our state of knowledge of the true value of a modelled quantity that arises from a given state of the system. It includes any artifact caused by the resolution of the target variables (Kaminski et al., 2001; Bocquet, 2005). It also includes uncertain choices about model structure and numerical implementation as well

as model parameters that are not included among the target variables. ~~This PDF~~ The PDF for the observation operator is often hard to verify because there are few cases where we know precisely the relevant values of the system state. Without this
5 knowledge it is hard to ascribe a difference between the simulated and true value to an error in the observation operator or an error in the prior. Frequently we must abstract the observation operator and run it under controlled conditions. An example is deliberate tracer release experiments (e.g. van Dop et al., 1998). Absent such direct verification, calculations like sensitivity analyses or ensemble experiments (e.g. Law et al., 1996) give incomplete guidance. It is important to note that if the PDFs are Gaussian then the errors due to observations and observation operators may be added by adding their covariances (Tarantola,
10 2005, Eq. 1.101). We frequently shorthand this as the data uncertainty (or worse data error) when it is usually dominated by the observation operator. The resulting PDF describes the difference we might expect between the simulated result of the observation operator and the measured value. Thus analysis of the residuals (observation – simulated quantity) can help test the assumed errors (e.g. Desroziers et al., 2005; Kuppel et al., 2013). This forms part of the diagnostics of data assimilation treated in Michalak et al. (2017) and touched on in Section 5.6.

15 In some cases it may be possible to calculate uncertainty in the observation operator in advance rather than diagnosing it from the assimilation. If there is an ensemble of observation operators available (e.g. Gurney et al., 2002) we can use the spread among their outputs as a measure of spread, including likely consistent or systematic errors arising from shared history and components (Bishop and Abramowitz, 2013). We may also perturb parameters in the observation operator to calculate their contribution to uncertainty.

20 5.5 Dynamical Models

Although they are not one of the ingredients in Figure 1, dynamical models are so common in this field that we include them here. Many of the systems studied are dynamical, in that they involve the evolution of the system state through time. We reserve the term dynamical model (often shorthanded to model) for the mathematical function or computer system that performs this evolution. We frequently consider the dynamical model to be deterministic, in which case the set of system states as a function
25 of time are a unique function of the boundary and initial conditions. Some formulations relax this condition, in which case we need a PDF for the model. In the most common case of a first-order model, the PDF represents our state of knowledge of the true state of the system at timestep $i + 1$ given its state at timestep i and perfect knowledge of all relevant boundary conditions. As with observation operators this is hard to generate in practice since we rarely know the preexisting state or boundary conditions perfectly. In repeated assimilation cases like numerical weather prediction the statistics of many cases can
30 give guidance. The PDF can also be deduced from the analysis of the model-data misfits when the statistics of the uncertainty of the model inputs and of the measurement errors are known (Kuppel et al., 2013).

The boundary between the observation operator and the dynamical model depends on our choice of target variables. If we consider only the initial state and assimilate observations over a time-window (the classic case of 4-dimensional variational data assimilation, 4D-Var) then the dynamical model forms part of the mapping between the unknowns and the observations so is properly considered part of the observation operator. If we include the system state from the assimilation window (weak-constraint 4D-Var) then the dynamical model enters as a separate object with its own PDF. We will see shortly how this can be

incorporated explicitly into the statistical framework. One confusing outcome of this distinction is that the same model could play the role of an observation operator or a dynamical model depending on how it is used.

5 5.6 Hierarchical Methods and Hyperparameters

The solution to the inverse data assimilation problem depends on many inputs that are themselves uncertain. These include parameters of the input PDFs such as variances or choices of observation operators. Frequently we regard these as fixed, which is likely to underestimate the uncertainty of the estimates. ~~We need therefore~~ One solution is to incorporate these extra variables into the inversion data assimilation process. The problem is to deal with the complex relationships such as the dependence of the likelihood $p(\mathbf{y}|H(\mathbf{x}))$ on H as well as \mathbf{x} . An explanation and evaluation of methods for such problems is given by Cressie et al. (2009), here we sketch the problem and note some applications in the field.

The chain rule of probabilities (Jaynes and Bretthorst, 2003, Eq. 14.5) can be stated as: For two events A and B

$$P(A, B) = P(A|B)P(B) \quad (4)$$

If the events are regions in parameter space we can develop similar relations with PDFs. We can go further if there are more than two parameters involved, building a hierarchy in which we construct the joint PDF from a base probability and a cascading set of conditional probabilities. For this reason the approach is described as hierarchical.

~~We can substitute~~ In our case we extend our target variables \mathbf{x} by a new set of unknowns θ . θ contains quantities that may affect our data assimilation problem but are not themselves target variables. Examples include parameters describing PDFs (variances, correlation lengths etc) choices of dynamical models or observation operators or even the complexity of the target variables themselves. Some elements in θ may not even concern \mathbf{x} such as the observational uncertainties treated in Michalak et al. (2005).

For our case Equation 4 ~~into~~ becomes

$$P(\mathbf{x}, \theta) = P(\mathbf{x}|\theta)P(\theta) \quad (5)$$

We can then solve Equation 2 ~~and proceed as normal. This substitution may occur~~ for the extended set of target variables \mathbf{x}, θ to obtain $P(\mathbf{x}, \theta)$ and then if we are not interested in θ may integrate Equation 5 to remove it. θ may appear with Equation 2. For example if we are unsure of the uncertainty we should attach to the prior, we may include parameters describing the uncertainty in the first term on the right of Equation 2. The same holds for the measurement process and the second term on the right. Finally if we have more than one potential observation operator we can include a collection of these into the third term on the right.

A prototype of this approach was given by Michalak et al. (2005). They included parameters describing \mathbf{R} (often called hyperparameters) in an atmospheric flux inversionassimilation. While they did not solve for the PDF of these hyperparameters they did calculate the mode of their distribution and recalculated the distributions of fluxes accordingly. This calculation did not include the impact of uncertainties in the hyperparameters on uncertainties in the fluxes. Michalak et al. (2005) and Wu et al. (2013) used the same approach to solve for correlation length scales among fluxes in their inversionflux errors in their

assimilation. Ganesan et al. (2014) took account of uncertainties in diagonal elements of \mathbf{B} and \mathbf{R} and temporal autocorrelations in \mathbf{R} . (Ganesan et al., 2014) did consider the impact of uncertainties in θ on estimated fluxes but at considerable computational
5 cost. Finally Rayner (2017a) considered the uncertainty introduced by an ensemble of observation operators in a well-known
atmospheric inversion the Transport Model Comparison (TRANSCOM) described in Gurney et al. (2002).

Finally we should note that the hierarchical approach outlined here subsumes the description in Section 4. For this reason, and as a continual reminder of the assumptions implicit in selecting models and other inputs, Wikle and Berliner (2007) write the whole methodology in hierarchical form.

10 6 Solving the Inverse-Assimilation Problem

6.1 General Principles

We saw in Section 4 that the solution of the assimilation problem consisted of the multiplication of three PDFs to generate the posterior PDF. Before describing solution methods applicable in various circumstances we point out some general consequences of Equation 2. We begin by stressing that the posterior PDF generated in Equation 2 is the most general form of the
15 solution. Derived quantities are generated by operations on this PDF.

Second, we see that the only physical model involved is the observation operator. All the sophisticated machinery of assimilation is not fundamental to the problem although we need it to derive most of the summary statistics.

Third there is no requirement that a solution exists or, if it does, it can have vanishingly low probability. In Figure 1 this would occur if various of the prior the PDFs were made tighter (smaller rectangles for example) so that they did not overlap. This is a
20 critical scientific discovery because it tells us our understanding of the problem is not consistent with the data, notwithstanding all the uncertainties of the inputs. It points out a boundary of "normal science" as defined by Kuhn (1962).⁴ Michalak et al. (2017) focus on diagnosing such inconsistencies.

6.2 Sampling Methods

A general approach for characterizing the posterior probability distribution of the target variables is to devise an approach for
25 generating a large number of samples, or realizations, from this multivariate posterior distribution. We briefly describe a few example approaches that fall under this umbrella.

One approach to solving Equation 2 is direct calculation of the product for a large number of realizations drawn from the constituent PDFs. Most simply, one can treat the PDF like any other scalar function and map it. This is often achieved by graphs or contour plots of a transformation of the PDF such as the least squares cost function (e.g. Kaminski et al., 2002, Fig. 12). This mapping approach requires one computation of the forward model for each parameter set, so is infeasible for even moderately dimensioned problems. In addition, most of these parameter sets will lie in very low probability regions.

⁴We thank Ian Enting for pointing this out.

A range of more sophisticated Monte Carlo sampling approaches exist for multi-dimensional problems. The most straightforward of these is direct sampling of the posterior PDF, feasible for the case where this posterior PDF is sufficiently simple to allow sampling from the distribution. This would be the case, for example, for multivariate Gaussian or lognormal distributions.

In cases where direct sampling is not feasible, the strategy often becomes one of sequentially sampling from a surrogate distribution (often termed the proposal distribution) in such a way that the ensemble of samples ultimately represents a sample from the more complex target distribution. Arguably, the simplest of the approaches falling under this umbrella is rejection sampling. In this approach, a sample is generated from a proposal distribution that is simple enough to be sampled directly, and that sample is accepted or rejected with a probability equal to the ratio between the likelihood of the sampled value based on the target versus the proposal distribution, normalized by a constant (e.g. Robert and Casella, 2004).

A particularly useful class of Monte Carlo sampling strategies are Markov chain Monte Carlo (MCMC) approaches. In these approaches, the samples form a Markov chain, such that each sample is not obtained independently, but rather as a perturbation of the last previously accepted sample. ~~More formally, a Markov Chain is a Stochastic traversal of some manifold in which the current state is dependent only on the immediately preceding state. The various MCMC approaches differ in how the candidate samples are generated based on the last accepted sample. Usually we construct the new state by adding a random perturbation to the previous state. For example, Metropolis et al. (1953) proposed and Hastings (1970) proved that a Markov chain based on the following algorithm samples an underlying PDF:-~~

1. ~~At iteration i we have state \mathbf{x}_i with probability $p(\mathbf{x}_i)$ and a set of accepted samples $S = (\mathbf{x}_1, \mathbf{x}_2 \dots \mathbf{x}_i)$;~~
2. ~~Generate a new candidate point \mathbf{x}_{i+1}^* by adding a random perturbation to \mathbf{x}_i ;~~
3. ~~Choose a uniformly distributed random number $u \in [0, 1]$;~~
4. ~~If $p(\mathbf{x}_{i+1}^*)/p(\mathbf{x}_i) > u$ we accept \mathbf{x}_{i+1}^* , append it to S and return to step 1, otherwise reject it and return to step 2.~~

~~The critical step is step 4 which is known as the Metropolis rule, and the algorithm overall is known as the Metropolis-Hastings algorithm. This algorithm produces samples with high serial correlation because successive points in the Markov Chain are likely to be close to each other in parameter space. The degree of correlation depends on the size and type of the perturbation applied in Step 2.~~

~~Rather than sampling the posterior PDF directly, one can also sample each of three PDFs sequentially (Figure 1). For this we can use, e.g. with the cascaded Metropolis algorithm (see Tarantola, 2005, Section 6.11). Once we have an accepted point from step 4 above we proceed to apply the Metropolis rule to the other probability distributions in the problem. In practice, if we are solving simultaneously for unknowns and data (the case in Figure 1) then we find a point acceptable to the first two PDFs on the right hand side of Equation 2 and only then go on to apply the rule to the third PDF. This is convenient since the third PDF is the only one requiring the observation operator and is hence the most expensive part of the algorithm.~~

Another commonly-used MCMC algorithm, the Gibbs sampler, can be seen as a special case of the Metropolis Hastings algorithm where the target variables are sampled individually and sequentially, rather than sampling the full multi-variate PDF of target variables as a single step (e.g. Casella and George, 1992). An advantage of the Gibbs sampler is that it can lead

to higher rates of acceptance, because defining strategic approaches for generating perturbations is more straightforward for univariate distributions. A disadvantage is that the overall computational cost can still be higher due to the need to perform sampling multiple times for each overall realisation.

The most important implementation detail for applying MCMC approaches is the choice of the perturbation (~~Step 2 in the case of the Metropolis-Hastings algorithm~~) because this determines how the PDF is sampled. Good strategies are adaptive, lengthening the step size to escape from improbable regions of the parameter space and shortening it to sample probable parts (noting the comments on independence above).

Note that it takes at least one forward run for every accepted point, meaning only inexpensive ~~models~~ observation operators can be treated this way. Furthermore the original algorithm cannot be parallelized because each calculation requires the previous one. Some related techniques are amenable to parallel computation (e.g. Jacob et al., 2011). On the other hand ~~MCMH~~ MCMC makes few assumptions about the posterior PDF, can handle a wide variety of input PDFs and requires no modification to the forward model.

6.3 Summary Statistics

We often describe the solution of the inverse assimilation problem using summary statistics of the posterior PDF. This can be because we assume that the posterior PDF has an analytic form with a few parameters that can describe it (e.g. the mean and standard deviation of the Gaussian) or because we need a numerical representation for later processing or comparison. This nearly universal practice is valuable but requires care. For example, consider a target variable in the form of a timeseries. We can represent the mode of the posterior PDF as a line graph and the posterior uncertainty as a band around this line. The true timeseries is a realisation drawn from within this band. Any statements about the temporal standard deviation of the true timeseries must be based on PDFs generated from these realisations. These estimates will generally yield larger variability than that from our most likely flux realisation.

~~There are some cases that are~~ The linear Gaussian case is so common that ~~they deserve it~~ deserves explicit treatment, however, provided the caveat above is borne in mind.

6.4 The Linear Gaussian Case

If H is linear and the PDFs of the prior, data and model are Gaussian then the posterior PDF consists of the multiplication of three Gaussians. We multiply exponentials by adding their exponents so it follows that the exponent of the posterior has a quadratic form and hence the posterior is Gaussian.

The Gaussian form for the inputs allows one important simplification without approximation. We noted in Section 4 that the solution to the inverse assimilation problem consists of finding the joint posterior PDF for unknowns and observed quantities then projecting this into the subspace of the unknowns. As we saw in section 4 we also calculate the posterior PDF for \mathbf{y} the observed quantity. If there are very large numbers of observations this is computationally prohibitive and usually uninteresting. If the PDFs are Gaussian, Tarantola (2005, Section 6.21) showed that the last two terms of Equation 2 can be combined into a single PDF $G(H(\mathbf{x}) - \mathbf{y}^o, \mathbf{R})$ where $\mathbf{R} = \mathbf{U}(H(\mathbf{x}) - \mathbf{y}) + \mathbf{U}(\mathbf{y}^o - \mathbf{y})$, i.e. by adding the two variances. The PDF now includes

only the unknowns \mathbf{x} and the posterior PDF can be written

$$p(\mathbf{x}|\mathbf{y}) \propto G(\mathbf{x}^b, \mathbf{B}) \times G(\mathbf{H}(\mathbf{x}), \mathbf{y}^o, \mathbf{R}) \quad (6)$$

5 this generates a Gaussian for \mathbf{x} whose mean and variance can be calculated by a range of methods (Rodgers, 2000, ch.1). The posterior covariance \mathbf{A} does not depend on \mathbf{x}^b or \mathbf{y}^o so that one can calculate posterior uncertainties before measurements are taken (Hardt and Scherbaum, 1994). This is the basis of the quantitative network design approaches described in Krause et al. (2008); Kaminski and Rayner (2016).

The largest computation in this method is usually the calculation of \mathbf{H} which includes the response of every observation
10 to every unknown. This may involve many runs of the forward model. Once completed, \mathbf{H} is the Green's function for the problem and instantiates the complete knowledge of the resolved dynamics. It enables a range of other statistical apparatus e.g. Michalak et al. (2005) or Rayner (2017a).

Following the introduction of formal Bayesian methods to atmospheric ~~inversions~~ assimilations by Enting et al. (1993, 1995) these analytical methods became the most common method for this field. Rayner et al. (1999); Bousquet et al. (2000); Roy
15 et al. (2003); Patra et al. (2005, 2006) used repeated forward runs of a single transport model while the series of papers from the Transport Model Intercomparison Project (TransCom) (Gurney et al., 2002, 2003; Law et al., 2003; Gurney et al., 2004; Baker et al., 2006) used repeated runs of an ensemble of models. Meanwhile Kaminski et al. (1999a, b); Rödenbeck et al. (2003); Peylin et al. (2005); Gerbig et al. (2009); Lauvaux et al. (2009); Turnbull et al. (2011); Gourdji et al. (2012); Fang et al. (2014); Fang and Michalak (2015); Lauvaux et al. (2012); Schuh et al. (2013) (among many others) used various forms of transport
20 model in reverse mode. These calculate the same Jacobian but require one run per observation rather than per unknown. The advent of more atmospheric data (especially satellite measurements of concentration) and the need for higher resolution of unknowns has seen these methods largely replaced by the ~~matrix-free methods described later~~ matrix-light methods described in Section 7.1.

6.5 Dynamically Informed Priors: the Kalman Filter

25 A common case in physical systems has the state of the system evolving according to some dynamical model M . This is usually (though not necessarily) some form of first-order differential equation. In such a case a natural algorithm for constraining the evolution of the system with data is

1. At any time step n our knowledge of the system is contained in the probability distribution $p(\mathbf{x}^n)$;
2. Calculate a probability distribution for the new state of the system $p(\mathbf{x}^{f,n+1})$ using the dynamical model M where the
30 superscript f (forecast) indicates the application of the ~~forward~~ dynamical model;
3. Use $p(\mathbf{x}^{f,n+1})$ as the prior PDF for the assimilation step described in Section 4;
4. This yields a posterior PDF $p(\mathbf{x}^{n+1})$ which we use as the input for the next iteration of step 1.

For the simplest case of linear M and H with Gaussian PDFs the algorithm was derived by Kalman (1960). The most difficult step is the generation of $p(\mathbf{x}^f, n)$. For the original Gaussian case where $p(\mathbf{x}^n) = G(\mathbf{x}^n, \mathbf{B}^n)$ this is given by

$$5 \quad p(\mathbf{x}^f, n) = G(M(\mathbf{x}^n), \mathbf{M}\mathbf{B}^n\mathbf{M}^T + \mathbf{Q}) \quad (7)$$

where \mathbf{Q} , the forecast error covariance, represents the uncertainty added to the state of the system by the model itself, i.e. not including uncertainties in the state. The matrix product in Equation 7 makes this approach computationally intractable for large numbers of unknowns and various approaches have been tried using a well-chosen subspace. The ensemble methods discussed in Section 7.3 have generally supplanted these. This also holds for nonlinear variants of the system.

10 The original conception of the Kalman Filter was for dynamical control in which the state of the system is continually adjusted in accordance with arriving observations. For data assimilation applied to biogeochemistry our motivation is to hindcast the state of the system and, optionally, infer some underlying constants (e.g. Trudinger et al., 2007, 2008). A consequence of making inferences at each time separately is that the system may be forced to follow an erroneous observation. For a hindcast we can counter this by expanding our set of unknowns to include not only the current state but the state for several timesteps
15 into the past, and therefore to smooth their previous analyses. This technique is known as the Kalman Smoother (Jazwinski, 1970) and also exists in ensemble variants (Evensen, 2009) and in variational ones (see Section 7.1). Evensen (2009) also showed that the Kalman Filter described above is a special case of the Kalman Smoother with only one time-level considered.

Note that by default, the time series of the maximum a posteriori estimates is not physically consistent, eg it does not conserve tracer mass from one assimilation step to another. However, the time series of the full PDFs includes trajectories that
20 are physically consistent.

7 Computational Aspects

Much of the computational machinery of data assimilation aims to calculate various summary statistics of the posterior PDF. The utility of this approach depends on whether these parameters summarize accurately the posterior PDF. It is best to choose the target summary statistic based on knowledge of the posterior PDF; there is no point estimating two parameters if the
25 posterior distribution is described by only one (e.g. the Poisson Distribution). Here we limit ourselves to discussions of the two most common parameters: the maximum a posteriori estimate and the posterior covariance. Solving for these parameters does not itself assume a Gaussian posterior but we often do assume such normality.

7.1 Iterative Solution for the Mode: Variational Methods

The mode of the posterior PDF is the value of the unknowns that maximizes the posterior probability. If it is often summarised as the maximum a posteriori or MAP estimate. If p is a product of exponentials then we can write

$$p(x) \propto e^{-J(x)} \quad (8)$$

then maximizing p is equivalent to minimizing J .⁵ In the linear Gaussian case J is quadratic so minimizing it is termed a least squares solution. It is perhaps unfortunate that many treatments of data assimilation start from the discussion of a least squares problem and thus hide many of the assumptions needed to get there.

The minimization of J takes us into the realm of numerical methods, beyond our scope. From the numerical viewpoint J is a scalar function of the unknowns \mathbf{x} . Minimizing J is a problem in the calculus of variations and so the methods are commonly termed variational.

The quadratic form is sufficiently common to warrant more development. From Equation 6, J takes the form

$$J(\mathbf{x}) = \frac{1}{2} [(\mathbf{x} - \mathbf{x}^b)^T \mathbf{B}^{-1} (\mathbf{x} - \mathbf{x}^b) + (H(\mathbf{x}) - \mathbf{y}^o)^T \mathbf{R}^{-1} (H(\mathbf{x}) - \mathbf{y}^o)] \quad (9)$$

Most efficient minimization algorithms require the gradient of $J(\mathbf{x})$ ($\nabla_{\mathbf{x}} J$). In general these gradients are precisely calculated by the use of adjoint methods (Griewank, 2000), but they involve the analytical derivation of each line of code of H beforehand. [The only matrices left that are potentially large in this approach are \$\mathbf{B}\$ and \$\mathbf{R}\$.](#)

7.2 Calculation of Posterior Uncertainty

Any reasonable summary of a posterior PDF will involve a parameter describing its spread as well as location. This spread almost always has a higher dimension than the mode because it must describe the joint probabilities of parameters. This immediately raises a problem of storage and computation because even moderate assimilation problems may include 100,000 unknowns for which representation of the joint PDF is intractable. In purely Gaussian problems it can be shown that the Hessian or second derivative $\nabla_{\mathbf{x}}^2 J$ evaluated at the minimum is the inverse covariance of the posterior PDF. This is promising given that many gradient-based algorithms for minimization calculate low-rank approximations of the Hessian (Meirink et al., 2008; Bousserez et al., 2015). Unfortunately they usually commence with the largest eigen-values of the Hessian corresponding to the lowest eigen-values (best-constrained parts) of the covariance. When convergence is achieved we are left with a subspace of the unknowns for which we can comment on the posterior uncertainty and ignorance beyond this. Properly we should restrict analysis of the results to this limited subspace but it usually does not map conveniently onto the physical problem we are studying (Chevallier et al., 2005).

An alternative approach is to generate an ensemble of modes using realizations of the prior and data PDFs (Fisher, 2003; Chevallier et al., 2007). The resulting realizations of the posterior PDF can be used as a sample for estimating population statistics. For Gaussian problems it can be shown that the sample covariance so generated is an unbiased estimator of the posterior covariance. [For non-Gaussian problems we can use the ensemble of realisations to estimate a parameter representing the spread of the distribution or calculate an empirical measure \(commonly the ensemble standard deviation\).](#)

7.3 The Ensemble Kalman Filter

The main limitations of the traditional Kalman Filter for data assimilation are its assumption of Gaussian PDFs, the requirement for linear M and H and the difficulty of projecting the state covariance matrix forward in time. The ensemble approach

⁵We cannot always write the PDFs this way, e.g. the uniform PDFs in Figure 1. In such cases the mode may not be defined.

described in Section 7.2 can also be used to circumvent these problems. The hierarchy of approaches is reviewed in Chen (2003). The most prominent (and most closely related to the original) is the Ensemble Kalman Filter (EnKF) (Evensen, 2003, 5 2009). Instead of representing exactly the PDF by its mean and covariance we generate an ensemble of realisations. We replace Equation 7 by advancing each realisation with the dynamical model then perturbing the result with noise generated from \mathbf{Q} . We then update each ensemble member using the Gaussian theory described in Section 6.4. Equation 6 shows that, as well as a prior estimate (for which we use the ensemble member), we need observations, an observational covariance and a prior covariance. The prior covariance we calculate from the ensemble itself. The observational covariance we assume. In the simplest version 10 of the EnKF we do not use the observations directly but rather perturb these with the observational noise. More recent versions (e.g. Tippett et al., 2003) avoid perturbing the observations but produce the same posterior uncertainty.

Computationally we need to run a dynamical model for each realisation of the ensemble. This differs from Equation 7 ~~where the propagation of the uncertainty scales with the number of unknowns~~ for which we need a run of the dynamical model for every unknown. Another advantage is that, while Equation 7 is exact only for linear models, the ensemble method may capture 15 nonlinear impacts on the state covariance. The biggest disadvantage is the sampling problem for the posterior PDF. The limited ensemble size introduces errors in the magnitudes of the variances and spurious correlations among state variables. Often the covariance must be inflated ad hoc to prevent the ensemble from collapsing to an artificially small variance. Similarly the spurious correlations between, for example, spatially separated points, mean that observations can artificially influence target variables at long distances. This is handled by limiting the range of influence of observations (see Houtekamer and Zhang, 2016, 20 for a review of these techniques for numerical weather prediction), but this fix has also unwanted effects at short distances and is questionable for long-lived tracers (e.g. Miyazaki et al., 2011).

7.4 Combining Filtering and Sampling: the Particle Filter

Particle Filters or Sequential Monte Carlo methods relax the assumptions inherent in the EnKF. Like the EnKF, the Particle Filter samples the underlying PDFs with a series of realisations rather than propagating the mean and variance of the PDF. 25 There are steps in the EnKF that assume Gaussian probability distributions and the Particle Filter avoids these, seeking to fit the true underlying PDF. We parallel the description of the Kalman Filter algorithm (Section 6.5) to highlight the differences.

1. At any time step n our knowledge of the system is contained in the probability distribution $p(\mathbf{x}^n)$. Generate a set of realisations (usually called particles) \mathbf{x}_i^n drawn from $p(\mathbf{x}^n)$;
2. Propagate each realisation forward in time using the dynamical model M to generate realisations of the forecast state 30 $\mathbf{x}_i^{M,n+1}$;
3. Account for the uncertainty in the dynamical model by perturbing $\mathbf{x}_i^{M,n+1}$ according to the model PDF to generate realisations $\mathbf{x}_i^{f,n+1}$ consistent with the dynamical model and its uncertainty;
4. For each $\mathbf{x}_i^{f,n+1}$, generate the simulated observations \mathbf{y}_i using the observation operator H then evaluate the PDF for the observations $p(\mathbf{y})$ at the point \mathbf{y}_i , these probabilities act as weights for $\mathbf{x}_i^{f,n+1}$;

5 5. Using some numerical procedure with inputs of $\mathbf{x}_i^{f,n+1}$ and $p(\mathbf{y}_i)$ generate a posterior PDF $p(\mathbf{x}^{n+1})$ which we use as the prior PDF for the next iteration.

We see that the last three steps mirror the multiplication of PDFs described in Equation 2. Also the difference between the EnKF and Particle Filter lies chiefly in how we generate the posterior PDF from the ensemble of forecast states; the EnKF uses the linear Gaussian approximation while the Particle filter uses some kind of sampling. Thus the Particle Filter is a hybrid of the Kalman Filter and sampling approaches.

10 The relaxation of the Gaussian assumption is at once a strength and weakness of the Particle Filter. It can handle more complex PDFs than the EnKF but it also requires many more samples because it samples the posterior PDF rather blindly while the EnKF is guided by the Kalman update step into regions of high probability. This difference has meant that, while the EnKF has been used successfully on some very large problems (such as numerical weather prediction) the Particle Filter has been limited to the same problem sizes as the approaches described in Section 6.2. More recent developments such as the
15 Marginal Particle Filter (Klaas et al., 2005) or the Equivalent Weights Particle Filter (Ades and van Leeuwen, 2015) try to overcome the sampling problem by introducing additional steps for the resampling of the particles.

An overview of the principal ideas of particle filtering and improvements upon the basic filter is given by Doucet et al. (2001) while van Leeuwen (2009) reviews its application in geophysical systems. Stordal et al. (2011) points to some further generalisations which may combine advantages of the EnKF and Particle Filter.

20 7.5 time window

A common feature of biogeochemical systems is a large range of time-scales. For example, terrestrial models may include both photosynthesis (sub-seconds) to forest succession (centuries). Slow processes often couple weakly to instantaneous observations so they must be constrained by long observational series. This can make the problem computationally intractable. A common solution is to break the problem into shorter assimilation windows. This is different from the case of numerical
25 weather prediction where the main consideration is the need for a new forecast every few hours and the advantage that the dynamical model can be considered perfect for short enough times.

The choice of assimilation window for biogeochemical applications is guided by several factors.

1. Computational cost: If our target variables change with time then longer assimilation windows imply higher dimensions for \mathbf{x} with corresponding space requirements. Any scheme that requires multiple runs over the assimilation window
30 (such as the EnKF or iterative methods) will also take longer.
2. Sensitivity of observations to target variables: These often decrease with time, especially for initial conditions. Once they become either too small or too unreliable (because of accumulating model error) there is little added value in assimilating observations further into the future. We may thus break the problem into several parallel assimilations.

(1) above means the assimilation window must also consider the time resolution for which we wish to solve. If we believe the value of some physical parameter only holds for a finite time (e.g. a month) there is little point exposing it to observations

several years later. In practice the assimilation window is usually set by a trade-off between computational constraints and the time-scale of the slowest processes studied.

8 Historical Overview

Throughout this paper we have adopted a thematic structure. Here we switch views to a narrative description. This is partly to situate the various methods we have described but also to give some signposts for the future. We cannot hope to be exhaustive, nor to trace the rise and fall of each method. Rather we will identify drivers for these developments and show examples of how they have guided the field. Our intention is to support the rest of the paper rather than provide a comprehensive review.

We can identify four major drivers for the development of biogeochemical data assimilation:

1. Development of the underlying methods and their percolation into the field;
2. rise in the scale and complexity of models and data sets;
3. rise in computational power;
4. diversifying needs within and beyond science.

These drivers do not always pull in the same direction, e.g. increases in scale may require simplification of techniques while computational power may allow more sophistication. Different practitioners will also make different choices faced with the same trade-offs so our narrative is not a single thread but some large-scale patterns are evident.

One of the first, clearly recognisable applications of data assimilation to biogeochemistry (Bolin and Keeling, 1963) used a set of measurements of CO₂ concentrations to estimate the meridional diffusion of this gas and the distribution of its sources. ~~We now know~~ It is now clear that the assumptions of Bolin and Keeling (1963) were wrong but, to quote Keeling (1998) “Our results could not be seriously challenged, however, until digital computers, atmospheric circulation models, and considerably more CO₂ data all became available two decades later.”

Bolin and Keeling (1963) confronted the problem that information is generally lost in the transformation of the inputs we seek to the outputs we observe. Approaches to this problem have guided the field since. Other fields face the same problem. Atmospheric remote sensing (Rodgers and Walshaw, 1966) and seismic studies (Backus and Gilbert, 1968) addressed it in related fields and their approaches, refracted through later developments, would eventually enter biogeochemistry. Approaches can be roughly divided into two classes: identifying and limiting the solution to those parts constrained by the data or the addition of ancillary information. The former approach is exemplified by the studies of Brown (1993, 1995) who used the singular value decomposition to establish flux patterns most strongly projected onto the data and limited her analysis to those components. ~~We can also regard later~~ Later studies which intentionally limited the number of unknowns (e.g. Fan et al., 1998; Gloor et al., 2000), can be regarded as less formal examples of this approach while bearing in mind the caution of Wunsch and Minster (1982) about allowing the data to choose the subspace of unknowns we constrain. Other approaches considered the observability of target variables by the available data. Target variables could be aggregated where they were not individually

constrained sufficiently. Manning et al. (2011) aggregated pixels in their flux ~~inversion~~ [assimilation](#) until they reached some
5 threshold of observability while Wu et al. (2011) included observability as a design criterion for their choice of target variables.

The addition of ancillary information is frequently termed regularisation since its practical purpose is to limit implausible solutions in poorly-constrained parts of parameter space. For most problems, solving inverse problems meant solving a version of Equation 9 which ignored the first term on the right-hand side. Regularisation consisted of adding some other term which penalised undesirable solutions. ~~possibilities~~ [Possibilities](#) included “shrinkage estimators” which penalised the overall size
10 of the solution (~~Krakauer et al., 2004~~) ([Shaby and Field, 2006](#)) or smoothing operators which penalised rapid variations in the solution (e.g. Enting, 1987; McIntosh and Veronis, 1993).

The importation of the explicit Bayesian methods which we have reviewed here largely supplanted the alternative regularisation methods though some studies (~~e.g. Issartel, 2005; Manning et al., 2011~~) ([e.g. Krakauer et al., 2004; Issartel, 2005; Manning et al., 2011](#))
15 followed other approaches. Also the geostatistical methods introduced into the field by Michalak et al. (2004) remain in use (e.g. Gourdjji et al., 2008; Miller et al., 2013). Bayesian methods entered biogeochemistry through atmospheric inverse studies (Enting et al., 1993, 1995). Most work since has elaborated different methods of solution or forms of prior knowledge. Almost all methods described in Section 6 have found some application. Often a method will be tried and left until an algorithmic advance or computational improvement makes it feasible.

Classical analytical methods (Section 6.4) were the first workhorse for flux ~~inversions~~ [assimilations](#) of inert tracers like CO₂
20 where the problem is linear. They are still widely used (e.g., Fang et al., 2014; Fang and Michalak, 2015; Henne et al., 2016; Saeki and Patra, 2017; Chen et al., 2017; Shiga et al., 2018; Wang et al., 2018). Originally Jacobians were generated with multiple forward runs, each run corresponding to a desired source component. The development of adjoint transport models allowed one run per observation (e.g. Kaminski et al., 1999a, b; Rödenbeck et al., 2003; Carouge et al., 2010a, b; Thompson et al., 2011; Gourdjji et al., 2012; Fang et al., 2014; Fang and Michalak, 2015). This is more efficient in the usual case of more
25 unknowns than observations. ~~the~~ [The](#) task of generating Jacobians is inherently parallel so the ultimate limitation on these methods is the storage and manipulation of large matrices. These limitations recede with increasing computational power and recent algorithmic advances (e.g. Yadav and Michalak, 2013) have also expanded their domain. Analytical methods are prized for exact calculation of posterior covariance and so are widely used in network design (~~references~~ (e.g. Gloor et al., 2000; Patra et al., 2003; Hungerschofer et al., 2010; Kaminski and Rayner, 2016). Matrix methods ~~also~~ are also often used where
30 more sophisticated statistics require more complex calculations such as the hierarchical calculations of Michalak et al. (2004); Ganesan et al. (2014); Rayner (2017b). They may also be used in weakly nonlinear forms where dimensionality is low enough to permit finite difference sensitivity calculations for each unknown (e.g. Kuppel et al., 2012; Bacour et al., 2015; MacBean et al., 2018; Norton et al., 2018).

Once the number of both observations and target variables renders computation of Jacobians unfeasible, variational methods
35 (section 7.1) are a natural extension of the matrix methods. This evolution is particularly evident in atmospheric problems, driven by increases in satellite data and a drive for higher resolution. Several groups have developed similar systems around adjoints of atmospheric transport models: (e.g. Rödenbeck, 2005; Chevallier et al., 2005; Henze et al., 2007; Bergamaschi et al., 2007; Liu et al., 2014; Wilson et al., 2014) for global models and Brioude et al. (2011); Broquet et al. (2011); Guerrette

and Henze (2015); Zheng et al. (2018b) for regional models. Most of the effort has centred on improving ~~our~~ knowledge of
5 sources and sinks (e.g. Basu et al., 2013; Escribano et al., 2016; Lu et al., 2016; Zheng et al., 2018a) but some have focused
on improving state estimation for forecasting (e.g. Elbern and Schmidt, 2001; Wang et al., 2011; Soon-Young et al., 2016).
To achieve high resolution while maintaining global consistency, some authors also combine methods, using one for the large-
scale and, potentially, a different method for a nested domain. This approach was introduced by Rödenbeck et al. (2009) and
has been recently used by Kountouris et al. (2018).

10 Versions of the Kalman Filter are some of the natural choices whenever the dynamical evolution of the model state is
explicitly included. Few atmospheric studies have exploited this capability though Miyazaki et al. (2011) solved for both fluxes
and concentrations. More common is a random walk or persistence model in which state variables evolve without dynamical
guidance. This was implemented for atmospheric flux ~~inversion-assimilation~~ by Mulquiney et al. (1995); Mulquiney and Norton
(1998); Mulquiney et al. (1998). Trudinger et al. (2002a, b) later used the Kalman smoother to estimate global fluxes from ice
15 core concentration and isotopic measurements. Peters et al. (2005) introduced the CarbonTracker system using a simplified
version of the EnKF and the global atmospheric model TM5 (Krol et al., 2005). TM5's global nested capability has allowed
a large suite of global and regional applications for CarbonTracker (e.g. Peters et al., 2010) for Europe or Kim et al. (2014)
for Asia. Other applications of the EnKF to the estimation of surface sources include (~~Zupanski et al., 2007; Feng et al., 2009~~)
Zupanski et al. (2007); Feng et al. (2009). One problem common to all these models was a potential time delay between target
20 variables and observations. This does not fit immediately into the original formalism of Kalman (1960). A solution was to
keep several time ~~periods-steps~~ in a moving assimilation window but ~~is still problematic for tracers that live longer than the
length-of-an-assimilation-window~~ one may need many steps if the time delay is long. This can vitiate some of the computational
advantages of the EnKF.

Kalman Filters were frequently limited by the need to calculate Jacobians for each target variable. This was circumvented by
25 the introduction of the EnKF. Provided one accounts for the problems of ensemble size the EnKF's combination of efficiency
and simplicity has led it to supplant other KF approaches.

The movement towards inclusion of explicit dynamics in biogeochemical inverse problems has increased the use of EnKF
and variational methods since both handle the nonlinear formulations common in biogeochemical models. In terrestrial models,
Williams et al. (2005); Quaife et al. (2008) included state variables of a simple terrestrial model in an ensemble system
30 while Dutta et al. (2018) solved for parameters. Trudinger et al. (2008) considered the joint estimation of parameters and
state variables using an EnKF and pseudo-observations at a site. Barrett (2002) and Braswell et al. (2005) used variational
approaches to optimise parameters of simple biosphere models against various site data. Rayner et al. (2005) and the papers
that followed it (e.g. Scholze et al., 2007; Knorr et al., 2010; Kaminski et al., 2012; Scholze et al., 2016) used an adjoint
of a more comprehensive (though incomplete) biosphere model coupled to various observation operators in their variational
approach.

There have been some attempts to compare the strengths and weaknesses of the various approaches above. Meirink et al.
5 (2008) or Kopacz et al. (2009) illustrated the equivalence of the variational and the analytical approaches for a given definition
of \mathbf{x} , while Liu et al. (2016) showed some convergence of the variational approach and of the ensemble Kalman filter for

increasing observation coverage. Trudinger et al. (2007) used a simplified terrestrial model and a fairly rigorous protocol to test the efficiency and robustness of several estimation methods. They found no clear differences among methods but did note the importance of careful analysis of assimilation performance (e.g. do the differences between posterior simulation and data agree with the assumed PDF). Chatterjee and Michalak (2013) compared the variational and EnKF approaches to estimating surface fluxes and also noticed that each had comparative strengths and weaknesses. ~~We have described how each of these methods is an alternative algorithm for the same underlying problem.~~ Trudinger et al. (2007) suggests that any of these algorithms is capable of performing well if well handled and analysed.

9 Future Developments

15 In previous sections, we have described how many assimilation methods are alternative algorithms for the same underlying problem. Different methods have risen to prominence in response to combinations of the four drivers mentioned above. Discussion of future directions is necessarily speculative but we can suggest how some of the drivers will evolve and their corresponding influence.

20 Computational constraints are less active now and likely to remain so. this is true even when we consider the increases in data size and complexity. The increased availability of large numbers of processors rather than their individual speed will likely privilege naturally parallel algorithms such as ensemble methods.

25 The rising complexity and diversity of models will also promote ensemble approaches. The strongest analogy is with climate prediction and seasonal forecasting where ensembles frequently include different dynamical formulations as well as different parameter settings. As yet there have been few attempts at multi-model ensemble data assimilation in biogeochemistry with the TransCom project the highest profile example. One reason for this is the scarcity of biogeochemical models with data assimilation capability. Another advantage of ensemble methods is a reduced entry barrier.

A more likely brake on the evolution of data assimilation in biogeochemistry is the quality of the models themselves. A precondition for the utility of data assimilation is that information learned in one environment or regime is applicable elsewhere, or that a model can make coherent inferences from different observations of the same system. As examples, Medvigy et al. (2009) found it difficult to transfer inferences made at one ecosystem site to another while Bacour et al. (2015) found it difficult to fit multiple streams of data with a single set of parameters. We would recommend tighter integration of data assimilation with model development so that components can be tested as they are developed.

Finally, the distinctions we have drawn between methods here represent a snapshot in time. Methods are likely to be matched and combined differently in future. Merging the various approaches, for instance by constructing optimal low-rank solutions even for high-dimensional problems in new systems ~~is seems~~ a promising way forward (Bousserez and Henze, 2018).

5 10 Conclusions

There ~~are an apparently~~ is a wide range of techniques available for assimilating data into models such as those used in biogeochemistry. ~~All~~ Most of these, however, inherit the basic formalism of Bayesian inference in which posterior probabilities are constructed as conjunctions or multiplications of probability distributions describing prior knowledge, measured quantities and models relating the two. The methods differ in the information they assume or adopt on the various components. In general the
10 more restrictive the assumptions the more powerful is the statistical apparatus available to analyze the system.

Code and Data Availability

This paper contains some sketches of mathematical development but does not rely on any particular piece of code or data set.

Author contributions. PR led the writing of the manuscript, AM wrote several sections and redacted the whole manuscript, FC wrote several sections, redacted the whole manuscript and was primarily responsible for regularising the notation.

15 *Competing interests.* The authors have no competing interests.

Acknowledgements. We acknowledge the support from the International Space Science Institute (ISSI). This publication is an outcome of the ISSI's Working Group on "Carbon Cycle Data Assimilation: How to consistently assimilate multiple data streams". Rayner is in receipt of an Australian Professorial Fellowship (DP1096309).

References

- 20 Ades, M. and van Leeuwen, P. J.: The equivalent-weights particle filter in a high-dimensional system, *Quarterly Journal of the Royal Meteorological Society*, 141, 484–503, <https://doi.org/10.1002/qj.2370>, <http://dx.doi.org/10.1002/qj.2370>, 2015.
- Backus, G. and Gilbert, F.: The resolving power of gross earth data, *Geophys. J. R. astr. Soc.*, 13, 247–276, 1968.
- Bacour, C., Peylin, P., MacBean, N., Rayner, P. J., Delage, F., Chevallier, F., Weiss, M., Demarty, J., Santaren, D., Baret, F., and Prunet, P.: Joint assimilation of eddy-covariance flux measurements and FAPAR products over temperate forests within a process-oriented biosphere model, *J. Geophys. Res.*, 120, <https://doi.org/10.1002/2015JG002966>, 2015.
- 25 Baker, D. F., Law, R. M., Gurney, K. R., Rayner, P., Peylin, P., Denning, A. S., Bousquet, P., Bruhwiler, L., Chen, Y.-H., Ciais, P., Fung, I. Y., Heimann, M., John, J., Maki, T., Maksyutov, S., Masarie, K., Prather, M., Pak, B., Taguchi, S., and Zhu, Z.: TransCom 3 inversion intercomparison: Impact of transport model errors on the interannual variability of regional CO₂ fluxes, 1988–2003, *Global Biogeochem. Cycles*, 20, GB1002, doi:10.1029/2004GB002439, 2006.
- 30 Balkanski, Y., Monfray, P., Battle, M., and Heimann, M.: Ocean primary production derived from satellite data: An evaluation with atmospheric oxygen measurements, *Global Biogeochem. Cycles*, 13, 257–271, 1999.
- Barrett, D. J.: Steady state turnover time of carbon in the Australian terrestrial biosphere, *Glob. Biogeochem. cyc.*, 16, 1108, <https://doi.org/10.1029/2002GB001860>, 2002.
- Basu, S., Guerlet, S., Butz, A., Houweling, S., Hasekamp, O., Aben, I., Krummel, P., Steele, P., Langenfelds, R., Torn, M., Biraud, S., Stephens, B., Andrews, A., and Worthy, D.: Global CO₂ fluxes estimated from GOSAT retrievals of total column CO₂, *Atmospheric Chemistry and Physics*, 13, 8695–8717, <https://doi.org/10.5194/acp-13-8695-2013>, <http://www.atmos-chem-phys.net/13/8695/2013/>, 2013.
- 35 Bergamaschi, P., Frankenberg, C., Meirink, J. F., Krol, M., Dentener, F., Wagner, T., Platt, U., Kaplan, J. O., Körner, S., Heimann, M., Dlugokencky, E. J., and Goede, A.: Satellite cartography of atmospheric methane from SCIAMACHY on board ENVISAT: 2. Evaluation based on inverse model simulations, *J. Geophys. Res.*, 112, D02304, doi:10.1029/2006JD007268, 2007.
- 5 Bishop, C. H. and Abramowitz, G.: Climate model dependence and the replicate Earth paradigm, *Climate dynamics*, 41, 885–900, 2013.
- Bocquet, M.: Grid resolution dependence in the reconstruction of an atmospheric tracer source, *Nonlinear Processes in Geophysics*, 12, 219–233, 2005.
- Bocquet, M., Elbern, H., Eskes, H., Hirtl, M., Žabkar, R., Carmichael, G., Flemming, J., Inness, A., Pagowski, M., Pérez Camaño, J., et al.: Data assimilation in atmospheric chemistry models: current status and future prospects for coupled chemistry meteorology models, *Atmospheric chemistry and physics*, 15, 5325–5358, 2015.
- 10 Bodman, R. W., Rayner, P. J., and Karoly, D. J.: Uncertainty in temperature projections reduced using carbon cycle and climate observations, *Nature Climate Change*, <https://doi.org/10.1038/NCLIMATE1903>, 2013.
- Bolin, B. and Keeling, C. D.: Large-scale atmospheric mixing as deduced from the seasonal and meridional variations of carbon dioxide, *J. Geophys. Res.*, 68, 3899–3920, 1963.
- 15 Bousquet, P., Peylin, P., Ciais, P., Le Quééré, C., Friedlingstein, P., and Tans, P. P.: Regional changes in carbon dioxide fluxes of land and oceans since 1980, *Science*, 290, 1342–1346, 2000.
- Bousserez, N. and Henze, D. K.: Optimal and scalable methods to approximate the solutions of large-scale Bayesian problems: theory and application to atmospheric inversion and data assimilation, *Quarterly Journal of the Royal Meteorological Society*, 144, 365–390, 2018.

- Bousserez, N., Henze, D., Perkins, A., Bowman, K., Lee, M., Liu, J., Deng, F., and Jones, D.: Improved analysis-error covariance matrix for high-dimensional variational inversions: application to source estimation using a 3D atmospheric transport model, *Quarterly Journal of the Royal Meteorological Society*, 141, 1906–1921, 2015.
- Brasseur, P., Gruber, N., Barciela, R., Brander, K., Doron, M., El Moussaoui, A., Hobday, A. J., Huret, M., Kremer, A.-S., Lehodey, P., Matear, R., Moulin, C., Murtugudde, R., Senina, I., and Svendsen, E.: Integrating biogeochemistry and ecology into ocean data assimilation systems, *Oceanography*, 22, 206–215, 2009.
- Braswell, B. H., Sacks, W. J., Linder, E., and Schimel, D. S.: Estimating diurnal to annual ecosystem parameters by synthesis of a carbon flux model with eddy covariance net ecosystem exchange observations, *glob. Change Biol.*, 11, 335–355, 2005.
- Brioude, J., Kim, S.-W., Angevine, W. M., Frost, G., Lee, S.-H., McKeen, S., Trainer, M., Fehsenfeld, F. C., Holloway, J., Ryerson, T., et al.: Top-down estimate of anthropogenic emission inventories and their interannual variability in Houston using a mesoscale inverse modeling technique, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- Broquet, G., Chevallier, F., Rayner, P., Aulagnier, C., Pison, I., Ramonet, M., Schmidt, M., Vermeulen, A. T., and Ciais, P.: A European summertime CO₂ biogenic flux inversion at mesoscale from continuous in situ mixing ratio measurements, *J. Geophys. Res.*, 116, D23 303, <https://doi.org/10.1029/2011JD016202>, 2011.
- Brown, M.: Deduction of Emissions of Source Gases Using an Objective Inversion Algorithm and a Chemical Transport Model, *J. Geophys. Res.*, 98, 12 639–12 660, 1993.
- Brown, M.: The singular value decomposition method applied to the deduction of the emissions and the isotopic composition of atmospheric methane, *J. Geophys. Res.*, 100, 11 425–11 446, 1995.
- Carouge, C., Bousquet, P., Peylin, P., Rayner, P. J., and Ciais, P.: What can we learn from European continuous atmospheric CO₂ measurements to quantify regional fluxes: Part 1: Potential of the 2001 network, *Atmospheric Chemistry and Physics*, 10, 3107–3117, <http://www.atmos-chem-phys.net/10/3107/2010/>, 2010a.
- Carouge, C., Rayner, P. J., Peylin, P., Bousquet, P., Chevallier, F., and Ciais, P.: What can we learn from European continuous atmospheric CO₂ measurements to quantify regional fluxes: Part 2: Sensitivity of flux accuracy to inverse setup, *Atmospheric Chemistry and Physics*, 10, 3119–3129, <http://www.atmos-chem-phys.net/10/3119/2010/>, 2010b.
- Carrassi, A., Bocquet, M., Bertino, L., and Evensen, G.: Data assimilation in the geosciences: An overview of methods, issues, and perspectives, *Wiley Interdisciplinary Reviews: Climate Change*, 9, e535, 2018.
- Casella, G. and George, E. I.: EXPLAINING THE GIBBS SAMPLER, *American Statistician*, 46, 167–174, <https://doi.org/10.2307/2685208>, <GotoISI>://WOS:A1992JE81900001, 1992.
- Chatterjee, A. and Michalak, A. M.: Technical Note: Comparison of ensemble Kalman filter and variational approaches for CO₂ data assimilation, *Atmospheric Chemistry and Physics*, 13, 11 643–11 660, <https://doi.org/10.5194/acp-13-11643-2013>, <https://www.atmos-chem-phys.net/13/11643/2013/>, 2013.
- Chen, J. M., Mo, G., and Deng, F.: A joint global carbon inversion system using both CO₂ and ¹³CO₂ atmospheric concentration data, *Geoscientific Model Development*, 10, 1131–1156, <https://doi.org/10.5194/gmd-10-1131-2017>, <https://www.geosci-model-dev.net/10/1131/2017/>, 2017.
- Chen, Z.: Bayesian filtering: From Kalman filters to particle filters, and beyond, *Statistics*, 182, 1–69, 2003.
- Chevallier, F., Fisher, M., Peylin, P., Serrar, S., Bousquet, P., Breon, F.-M., Chedin, A., and Ciais, P.: Inferring CO₂ sources and sinks from satellite observations: method and application to TOVS data, *J. Geophys. Res.*, (submitted), 2005.

- Chevallier, F., Bréon, F.-M., and Rayner, P. J.: The Contribution of the Orbiting Carbon Observatory to the Estimation of CO₂ Sources and Sinks: Theoretical Study in a Variational Data Assimilation Framework, *J. Geophys. Res.*, 112, D09 307, doi:10.1029/2006JD007375, 2007.
- Chevallier, F., Ciais, P., Conway, T. J., Aalto, T., Anderson, B. E., Bousquet, P., Brunke, E. G., Ciattaglia, L., Esaki, Y., Frohlich, M., Gomez, A., Gomez-Pelaez, A. J., Haszpra, L., Krummel, P. B., Langenfelds, R. L., Leuenberger, M., Machida, T., Maignan, F., Matsueda, H., Morgu, J. A., Mukai, H., Nakazawa, T., Peylin, P., Ramonet, M., Rivier, L., Sawa, Y., Schmidt, M., Steele, L. P., Vay, S. A., Vermeulen, A. T., Wofsy, S., and Worthy, D.: CO₂ surface fluxes at grid point scale estimated from a global 21 year reanalysis of atmospheric measurements, *Journal of Geophysical Research: Atmospheres*, 115, n/a–n/a, <https://doi.org/10.1029/2010JD013887>, <http://dx.doi.org/10.1029/2010JD013887>, 2010.
- Chevallier, F., Wang, T., Ciais, P., Maignan, F., Bocquet, M., Altaf Arain, M., Cescatti, A., Chen, J., Dolman, A. J., Law, B. E., Margolis, H. A., Montagnani, L., and Moors, E. J.: What eddy-covariance measurements tell us about prior land flux errors in CO₂-flux inversion schemes, *Global Biogeochemical Cycles*, 26, n/a–n/a, <https://doi.org/10.1029/2010GB003974>, <http://dx.doi.org/10.1029/2010GB003974>, 2012.
- Cressie, N., Calder, C. A., Clark, J. S., Hoef, J. M. V., and Wikle, C. K.: Accounting for uncertainty in ecological analysis: the strengths and limitations of hierarchical statistical modeling, *Ecological Applications*, 19, 553–570, <https://doi.org/10.1890/07-0744.1>, <http://dx.doi.org/10.1890/07-0744.1>, 2009.
- Desroziers, G., Berre, L., Chapnik, B., and Poli, P.: Diagnosis of observation, background and analysis-error statistics in observation space, *Quarterly Journal of the Royal Meteorological Society*, 131, 3385–3396, <https://doi.org/10.1256/qj.05.108>, <http://dx.doi.org/10.1256/qj.05.108>, 2005.
- Doucet, A., de Freitas, N., and Gordon, N., eds.: *Sequential Monte Carlo Methods in Practice*, Springer New York, <https://doi.org/10.1007/978-1-4757-3437-9>, 2001.
- Dutta, D., Schimel, D. S., Sun, Y., van der Tol, C., and Frankenberg, C.: Optimal Inverse Estimation of Ecosystem Parameters from Observations of Carbon and Energy Fluxes, *Biogeosciences Discussions*, 2018, 1–41, <https://doi.org/10.5194/bg-2018-303>, <https://www.biogeosciences-discuss.net/bg-2018-303/>, 2018.
- Elbern, H. and Schmidt, H.: Ozone episode analysis by four-dimensional variational chemistry data assimilation, *Journal of Geophysical Research: Atmospheres*, 106, 3569–3590, 2001.
- Enting, I. G.: On the use of smoothing splines to filter CO₂ data, *J. Geophys. Res.*, 92D, 10977–10984, 1987.
- Enting, I. G.: *Inverse Problems in Atmospheric Constituent Transport*, Cambridge University Press, 2002.
- Enting, I. G., Trudinger, C. M., Francey, R. J., and Granek, H.: Synthesis Inversion of Atmospheric CO₂ Using the GISS Tracer Transport Model, Tech. Paper No. 29, CSIRO Div. Atmos. Res., 1993.
- Enting, I. G., Trudinger, C. M., and Francey, R. J.: A synthesis inversion of the concentration and $\delta^{13}\text{C}$ of atmospheric CO₂, *Tellus*, 47B, 35–52, 1995.
- Escribano, J., Boucher, O., Chevallier, F., and Huneus, N.: Subregional inversion of North African dust sources, *Journal of Geophysical Research: Atmospheres*, 121, 8549–8566, 2016.
- Evans, S. N. and Stark, P.: Inverse Problems as Statistics, *Inverse problems*, 18, R55–R97, 2002.
- Evensen, G.: The ensemble Kalman filter: Theoretical formulation and practical implementation, *Ocean Dynamics*, 53, 343–367, 2003.
- Evensen, G.: The ensemble Kalman filter for combined state and parameter estimation, *Control Systems, IEEE*, 29, 83–104, <https://doi.org/10.1109/MCS.2009.932223>, 2009.

- Fan, S., Gloor, M., Mahlman, J., Pacala, S., Sarmiento, J., Takahashi, T., and Tans, P.: A Large Terrestrial Carbon Sink in North America
20 Implied by Atmospheric and Oceanic CO₂ Data and Models, *Science*, 282, 442–446, 1998.
- Fang, Y. and Michalak, A. M.: Atmospheric observations inform CO₂ flux responses to enviroclimatic drivers, *Global Biogeochemical
Cycles*, 29, 555–566, <GotoISI>://CCC:000356383700003, 2015.
- Fang, Y., Michalak, A. M., Shiga, Y. P., and Yadav, V.: Using atmospheric observations to evaluate the spatiotemporal variability of CO₂
fluxes simulated by terrestrial biospheric models, *Biogeosciences*, 11, 6985–6997, <https://doi.org/10.5194/bg-11-6985-2014>, <GotoISI>:
25 //WOS:000346357100032, 2014.
- Feng, L., Palmer, P., Bösch, H., and Dance, S.: Estimating surface CO₂ fluxes from space-borne CO₂ dry air mole fraction observations
using an ensemble Kalman Filter, *Atmospheric Chemistry and Physics*, 9, 2619–2633, 2009.
- Fisher, M.: Estimation of Entropy Reduction and Degrees of Freedom for Signal for Large Variational Analysis Systems, ECMWF technical
memorandum, European Centre for Medium-Range Weather Forecasts, <http://books.google.com.au/books?id=anikQwAACAAJ>, 2003.
- 30 Fung, I. Y., Tucker, C. J., and Prentice, K. C.: Application of Advanced Very High Resolution Radiometer Vegetation Index to Study
Atmosphere-Biosphere Exchange of CO₂, *J. Geophys. Res.*, 92, 2999–3015, 1987.
- Ganesan, A. L., Rigby, M., Zammit-Mangion, A., Manning, A. J., Prinn, R. G., Fraser, P. J., Harth, C. M., Kim, K.-R., Krummel, P. B., Li, S.,
Mühle, J., O’Doherty, S. J., Park, S., Salameh, P. K., Steele, L. P., and Weiss, R. F.: Characterization of uncertainties in atmospheric trace
gas inversions using hierarchical Bayesian methods, *Atmospheric Chemistry and Physics*, 14, 3855–3864, [https://doi.org/10.5194/acp-14-
35 3855-2014](https://doi.org/10.5194/acp-14-3855-2014), <http://www.atmos-chem-phys.net/14/3855/2014/>, 2014.
- Gerbig, C., Dolman, A. J., and Heimann, M.: On observational and modelling strategies targeted at regional carbon exchange over continents,
BIOGEOSCIENCES, 6, 1949–1959, 2009.
- Gloor, M., Fan, S. M., Pacala, S., and Sarmiento, J.: Optimal sampling of the atmosphere for purpose of inverse modeling: A model study,
Global Biogeochem. Cycles, 14, 407–428, 2000.
- Gourdji, S. M., Mueller, K. L., Schaefer, K., and Michalak, A. M.: Global Monthly-Averaged CO₂ Fluxes Recovered Using a Geostatistical
Inverse Modeling Approach: 2. Results Including Auxiliary Environmental Data, *J. Geophys. Res.*, (in press), 2008.
- Gourdji, S. M., Mueller, K. L., Yadav, V., Huntzinger, D. N., Andrews, A. E., Trudeau, M., Petron, G., Nehrkorn, T., Eluszkiewicz, J.,
5 Henderson, J., Wen, D., Lin, J., Fischer, M., Sweeney, C., and Michalak, A. M.: North American CO₂ exchange: inter-comparison of
modeled estimates with results from a fine-scale atmospheric inversion, *Biogeosciences*, 9, 457–475, [https://doi.org/10.5194/bg-9-457-
2012](https://doi.org/10.5194/bg-9-457-2012), <GotoISI>://WOS:000300229000031, 2012.
- Griewank, A.: *Evaluating Derivatives: Principles and Techniques of Automatic Differentiation*, SIAM, Philadelphia, Pa., 2000.
- Guerrette, J. and Henze, D.: Development and application of the WRFPLUS-Chem online chemistry adjoint and WRFDA-Chem assimilation
10 system, *Geoscientific Model Development*, 8, 1857–1876, 2015.
- Gurney, K. R., Law, R. M., Denning, A. S., Rayner, P. J., Baker, D., Bousquet, P., Bruhwiler, L., Chen, Y.-H., Ciais, P., Fan, S., Fung, I. Y.,
Gloor, M., Heimann, M., Higuchi, K., John, J., Maki, T., Maksyutov, S., Masarie, K., Peylin, P., Prather, M., Pak, B. C., Randerson, J.,
Sarmiento, J., Taguchi, S., Takahashi, T., and Yuen, C.-W.: Towards robust regional estimates of CO₂ sources and sinks using atmospheric
transport models, *Nature*, 415, 626–630, 2002.
- 15 Gurney, K. R., Law, R. M., Denning, A. S., Rayner, P. J., Baker, D., Bousquet, P., Bruhwiler, L., Chen, Y.-H., Ciais, P., Fan, S., Fung, I. Y.,
Gloor, M., Heimann, M., Higuchi, K., John, J., Kowalczyk, E., Maki, T., Maksyutov, S., Peylin, P., Prather, M., Pak, B. C., Sarmiento, J.,
Taguchi, S., Takahashi, T., and Yuen, C.-W.: TransCom 3 CO₂ inversion intercomparison: 1. Annual mean control results and sensitivity
to transport and prior flux information, *Tellus*, 55B, 555–579, doi:10.1034/j.1600-0560.2003.00049.x, 2003.

- Gurney, K. R., Law, R. M., Denning, A. S., Rayner, P. J., Pak, B. C., Baker, D., Bousquet, P., Bruhwiler, L., Chen, Y.-H., Ciais, P.,
 20 Fung, I. Y., Heimann, M., John, J., Maki, T., Maksyutov, S., Peylin, P., Prather, M., and Taguchi, S.: Transcom 3 inversion inter-
 comparison: Model mean results for the estimation of seasonal carbon sources and sinks, *Global Biogeochem. Cycles*, 18, GB1010,
 doi:10.1029/2003GB002111, 2004.
- Hardt, M. and Scherbaum, F.: The Design of Optimum Networks for Aftershock Recordings, *Geophys. J. Int.*, 117, 716–726, 1994.
- Hastings, W. K.: Monte Carlo sampling methods using Markov Chains and their applications, *Biometrika*, 57, 97–109, 1970.
- 25 Henne, S., Brunner, D., Oney, B., Leuenberger, M., Eugster, W., Bamberger, I., Meinhardt, F., Steinbacher, M., and Emmenegger, L.: Validat-
 ion of the Swiss methane emission inventory by atmospheric observations and inverse modelling, *Atmospheric Chemistry and Physics*,
 16, 3683–3710, <https://doi.org/10.5194/acp-16-3683-2016>, <https://www.atmos-chem-phys.net/16/3683/2016/>, 2016.
- Henze, D. K., Hakami, A., and Seinfeld, J. H.: Development of the adjoint of GEOS-Chem, *Atmospheric Chemistry and Physics*, 7, 2413–
 2433, 2007.
- 30 Houtekamer, P. and Zhang, F.: Review of the ensemble Kalman filter for atmospheric data assimilation, *Monthly Weather Review*, 144,
 4489–4532, 2016.
- Hungerschofer, K., Bréon, F.-M., Peylin, P., Chevallier, F., Rayner, P., Klonecki, A., and Houweling, S.: Evaluation of various observing
 systems for the global monitoring of CO₂ surface fluxes, *Atmos. Chem. Phys.*, 10, 10 503–10 520, [https://doi.org/10.5194/acp-10-10503-](https://doi.org/10.5194/acp-10-10503-2010)
 2010, 2010.
- 35 Ide, K., Courtier, P., Ghil, M., and Lorenc, A. C.: Unified Notation for Data Assimilation: Operational, Sequential and Variational, 1997.
- Issartel, J.-P.: Emergence of a tracer source from air concentration measurements, a new strategy for linear assimilation, *Atmospheric Chem-
 istry and Physics*, 5, 249–273, 2005.
- Jacob, P., Robert, C. P., and Smith, M. H.: Using parallel computation to improve independent Metropolis–Hastings based estimation, *Journal
 of Computational and Graphical Statistics*, 20, 616–635, 2011.
- Jaynes, E. and Bretthorst, G.: *Probability Theory: The Logic of Science*, Cambridge University Press, [http://books.google.com.au/books?id=](http://books.google.com.au/books?id=tTN4HuUNXjgC)
 5 [tTN4HuUNXjgC](http://books.google.com.au/books?id=tTN4HuUNXjgC), 2003.
- Jazwinski, A. H.: *Stochastic processes and filtering theory*, Academic Press, New York, 1970.
- Jeffreys, H.: *Theory of Probability.*, 1939.
- Kalman, R. E.: A new approach to linear filtering and prediction problems, *J. Basic Eng. (ASME)*, 82D, 35–45, 1960.
- Kalnay, E.: *Atmospheric modelling, data assimilation and predictability*, Cambridge University Press, 2003.
- 10 Kaminski, T. and Rayner, P. J.: Quantitative Network Design in Biogeochemical Data Assimilation, *Atmos. Chem. Phys.*, (in prep.), 2016.
- Kaminski, T., Heimann, M., and Giering, R.: A coarse grid three-dimensional global inverse model of the atmospheric transport, 1, Adjoint
 Model and Jacobian Matrix, *J. Geophys. Res.*, 104, 18,535–18,553, 1999a.
- Kaminski, T., Heimann, M., and Giering, R.: A coarse grid three-dimensional global inverse model of the atmospheric transport, 2, Inversion
 of the transport of CO₂ in the 1980s, *J. Geophys. Res.*, 104, 18 555–18 582, 1999b.
- 15 Kaminski, T., Rayner, P. J., Heimann, M., and Enting, I. G.: On Aggregation Errors in Atmospheric Transport Inversions, *J. Geophys. Res.*,
 106, 4703–4715, 2001.
- Kaminski, T., Knorr, W., Rayner, P., and Heimann, M.: Assimilating Atmospheric data into a Terrestrial Biosphere Model: A case study of
 the seasonal cycle, *Glob. Biogeochem. Cyc.*, 16, 1066, doi:10.1029/2001GB001463, 2002.

- Kaminski, T., Rayner, P. J., Voßbeck, M., Scholze, M., and Koffi, E.: Observing the continental-scale carbon balance: assessment of sampling complementarity and redundancy in a terrestrial assimilation system by means of quantitative network design, *Atmospheric Chemistry and Physics*, 12, 7867–7879, <https://doi.org/10.5194/acp-12-7867-2012>, <http://www.atmos-chem-phys.net/12/7867/2012/>, 2012.
- Keeling, C. D.: Rewards and Penalties of Monitoring the Earth, *Annu. Rev. Energy Environ.*, 23, 25–82, 1998.
- Kim, J., Kim, H. M., and Cho, C.-H.: The effect of optimization and the nesting domain on carbon flux analyses in Asia using a carbon tracking system based on the ensemble Kalman filter, *Asia-Pacific Journal of Atmospheric Sciences*, 50, 327–344, 2014.
- Klaas, M., de Freitas, N., and Doucet, A.: Toward Practical N2 Monte Carlo: the Marginal Particle Filter, in: *Proceedings of the Twenty-First Conference Annual Conference on Uncertainty in Artificial Intelligence (UAI-05)*, pp. 308–315, AUAI Press, Arlington, Virginia, 2005.
- Knorr, W., Kaminski, T., Scholze, M., Gobron, N., Pinty, B., Giering, R., and Mathieu, P.-P.: Carbon cycle data assimilation with a generic phenology model, *Journal of Geophysical Research: Biogeosciences*, 115, n/a–n/a, <https://doi.org/10.1029/2009JG001119>, <http://dx.doi.org/10.1029/2009JG001119>, 2010.
- Kopacz, M., Jacob, D. J., Henze, D. K., Heald, C. L., Streets, D. G., and Zhang, Q.: Comparison of adjoint and analytical Bayesian inversion methods for constraining Asian sources of carbon monoxide using satellite (MOPITT) measurements of CO columns, *Journal of Geophysical Research: Atmospheres*, 114, 2009.
- Kountouris, P., Gerbig, C., Rödenbeck, C., Karstens, U., Koch, T. F., and Heimann, M.: Atmospheric CO₂ inversions on the mesoscale using data-driven prior uncertainties: methodology and system evaluation, *Atmospheric Chemistry and Physics*, 18, 3027–3045, 2018.
- Krakauer, N. Y., Schneider, T., Randerson, J. T., and Olsen, S. C.: Using generalized cross-validation to select parameters in inversions for regional carbon fluxes, *Geophys. Res. Lett.*, 31, 19 108, <https://doi.org/10.1029/2004GL020323>, 2004.
- Krause, A., Singh, A., and Guestrin, C.: Near-optimal sensor placements in Gaussian processes: Theory, efficient algorithms and empirical studies, *Journal of Machine Learning Research*, 9, 235–284, 2008.
- Krol, M., Houweling, S., Bregman, B., and van den Broek, M.: The two-way nested global chemistry-transport zoom model TM5: algorithm and applications, *Atmos. chem. Phys.*, 4, 2005.
- Kuhn, T. S.: *The structure of scientific revolutions*, University of Chicago, 1962.
- Kuppel, S., Peylin, P., Chevallier, F., Bacour, C., Maignan, F., and Richardson, A.: Constraining a global ecosystem model with multi-site eddy-covariance data, *Biogeosciences*, 9, 3757–3776, 2012.
- Kuppel, S., Chevallier, F., and Peylin, P.: Quantifying the model structural error in carbon cycle data assimilation systems, *Geoscientific Model Development*, 6, 45–55, <https://doi.org/10.5194/gmd-6-45-2013>, <http://www.geosci-model-dev.net/6/45/2013/>, 2013.
- Laplace, P. S.: *Mémoire sur la probabilité des causes par les événements*, in: *Mem. Acad. Roy. Sci.*, vol. 6, pp. 621–656, 1774.
- Lauvaux, T., Pannekoucke, O., Sarrat, C., Chevallier, F., Ciais, P., Noilhan, J., and Rayner, P. J.: Structure of the transport uncertainty in mesoscale inversions of CO₂ sources and sinks using ensemble model simulations, *Biogeosciences*, 6, 1089–1102, 2009.
- Lauvaux, T., Schuh, A. E., Uliasz, M., Richardson, S., Miles, N., Andrews, A. E., Sweeney, C., Diaz, L. I., Martins, D., Shepson, P. B., and Davis, K. J.: Constraining the CO₂ budget of the corn belt: exploring uncertainties from the assumptions in a mesoscale inverse system, *Atmospheric Chemistry and Physics*, 12, 337–354, <https://doi.org/10.5194/acp-12-337-2012>, <http://www.atmos-chem-phys.net/12/337/2012/>, 2012.
- Law, R. M., Rayner, P. J., Denning, A. S., Erickson, D., Fung, I. Y., Heimann, M., Piper, S. C., Ramonet, M., Taguchi, S., Taylor, J. A., Trudinger, C. M., and Watterson, I. G.: Variations in modelled atmospheric transport of carbon dioxide and the consequences for CO₂ inversions, *Global Biogeochem. Cycles*, 10, 783–796, 1996.

- Law, R. M., Chen, Y.-H., Gurney, K. R., and TransCom 3 modellers: TransCom 3 CO₂ inversion intercomparison: 2. Sensitivity of annual mean results to data choices, *Tellus*, 55B, 580–595, doi:10.1034/j.1600-0560.2003.00053.x, 2003.
- Liu, J., Bowman, K. W., Lee, M., Henze, D. K., Bousserez, N., Brix, H., James Collatz, G., Menemenlis, D., Ott, L., Pawson, S., et al.: Carbon monitoring system flux estimation and attribution: impact of ACOS-GOSAT XCO₂ sampling on the inference of terrestrial biospheric sources and sinks, *Tellus B: Chemical and Physical Meteorology*, 66, 22 486, 2014.
- Liu, J., Bowman, K. W., and Lee, M.: Comparison between the Local Ensemble Transform Kalman Filter (LETKF) and 4D-Var in atmospheric CO₂ flux inversion with the Goddard Earth Observing System-Chem model and the observation impact diagnostics from the LETKF, *Journal of Geophysical Research: Atmospheres*, 121, 2016.
- Lu, S., Lin, H., Heemink, A., Fu, G., and Segers, A.: Estimation of volcanic ash emissions using trajectory-based 4D-Var data assimilation, *Monthly Weather Review*, 144, 575–589, 2016.
- Lunt, M. F., Rigby, M., Ganesan, A. L., and Manning, A. J.: Estimation of trace gas fluxes with objectively determined basis functions using reversible-jump Markov chain Monte Carlo, *Geoscientific Model Development*, 9, 3213–3229, 2016.
- MacBean, N., Maignan, F., Bacour, C., Lewis, P., Peylin, P., Guanter, L., Köhler, P., Gómez-Dans, J., and Disney, M.: Strong constraint on modelled global carbon uptake using solar-induced chlorophyll fluorescence data, *Scientific reports*, 8, 1973, 2018.
- Manning, A., O’Doherty, S., Jones, A., Simmonds, P., and Derwent, R.: Estimating UK methane and nitrous oxide emissions from 1990 to 2007 using an inversion modeling approach, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- McIntosh, P. C. and Veronis, G.: Solving Underdetermined Tracer Inverse Problems by Spatial Smoothing and Cross Validation, *J. Phys. Oceanogr.*, 23, 716–730, 1993.
- Medvigy, D., Wofsy, S. C., Munger, J. W., Hollinger, D. Y., and Moorcroft, P. R.: Mechanistic scaling of ecosystem function and dynamics in space and time: Ecosystem Demography model version 2, *J. Geophys. Res.*, 114, G01 002, <https://doi.org/10.1029/2008JG000812>, 2009.
- Meinshausen, M., Raper, S. C. B., and Wigley, T. M. L.: Emulating IPCC AR4 atmosphere-ocean and carbon cycle models for projecting global-mean, hemispheric and land/ocean temperatures: MAGICC 6.0, *Atmospheric Chemistry and Physics Discussions*, 8, 6153–6272, <https://doi.org/10.5194/acpd-8-6153-2008>, <http://www.atmos-chem-phys-discuss.net/8/6153/2008/>, 2008.
- Meirink, J. F., Bergamaschi, P., and Krol, M. C.: Four-dimensional variational data assimilation for inverse modelling of atmospheric methane emissions: method and comparison with synthesis inversion, *Atmospheric Chemistry and Physics*, 8, 6341–6353, 2008.
- Metropolis, N., Rosenbluth, A. W., Rosenbluth, M. N., Teller, A. H., and Teller, E.: Equation of state calculations for fast computing machines, *Journal of Chemical Physics*, 21, 1087–1092, 1953.
- Michalak, A. M., Bruhwiler, L., and Tans, P. P.: A geostatistical approach to surface flux estimation of atmospheric trace gases, *J. Geophys. Res.*, 109, D14 109, doi: 10.1029/2003JD004422, 2004.
- Michalak, A. M., Hirsch, A., Bruhwiler, L., Gurney, K. R., Peters, W., and Tans, P. P.: Maximum likelihood estimation of covariance parameters for Bayesian atmospheric trace gas surface flux inversions, *J. Geophys. Res.*, 110, D24 107, <https://doi.org/10.1029/2005JD005970>, 2005.
- Michalak, A. M., Randazzo, N. A., and Chevallier, F.: Diagnostic methods for atmospheric inversions of long-lived greenhouse gases, *Atmospheric Chemistry and Physics*, 17, 7405–7421, <https://doi.org/10.5194/acp-17-7405-2017>, <https://www.atmos-chem-phys.net/17/7405/2017/>, 2017.
- Miller, S. M., Wofsy, S. C., Michalak, A. M., Kort, E. A., Andrews, A. E., Biraud, S. C., Dlugokencky, E. J., Eluszkiewicz, J., Fischer, M. L., Janssens-Maenhout, G., Miller, B. R., Miller, J. B., Montzka, S. A., Nehrkorn, T., and Sweeney, C.: Anthropogenic emissions of

- methane in the United States, *Proceedings of the National Academy of Sciences of the United States of America*, 110, 20018–20022, <https://doi.org/10.1073/pnas.1314392110>, <GotoISI>://WOS:000328061700028, 2013.
- 20 Miyazaki, K., Maki, T., Patra, P., and Nakazawa, T.: Assessing the impact of satellite, aircraft, and surface observations on CO₂ flux estimation using an ensemble-based 4-D data assimilation system, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- Moore, A. M., Arango, H. G., and Broquet, G.: Estimates of analysis and forecast error variances derived from the adjoint of 4D-Var, *Monthly Weather Review*, 140, 3183–3203, 2012.
- 25 Mulquiney, J. E. and Norton, J. P.: A new inverse method for trace gas flux estimation 1. State-space model identification and constraints, *J. Geophys. Res.*, 103D, 1417–1427, 1998.
- Mulquiney, J. E., Norton, J. P., Jakeman, A. J., and Taylor, J. A.: Random walks in the Kalman filter: Implications for greenhouse gas flux deductions, *Environmetrics*, 6, 473–478, 1995.
- Mulquiney, J. E., Taylor, J. A., Jakeman, A. J., Norton, J. P., and Prinn, R. G.: A new inverse method for trace gas flux estimation 2. Application to tropospheric CFC₃ fluxes, *J. Geophys. Res.*, 103D, 1429–1442, 1998.
- 30 Norton, A. J., Rayner, P. J., Koffi, E. N., and Scholze, M.: Assimilating solar-induced chlorophyll fluorescence into the terrestrial biosphere model BETHY-SCOPE v1.0: model description and information content, *Geoscientific Model Development*, 11, 1517–1536, <https://doi.org/10.5194/gmd-11-1517-2018>, <https://www.geosci-model-dev.net/11/1517/2018/>, 2018.
- Patra, P. K., Maksyutov, S., Baker, D., Bousquet, P., Bruhwiler, L., Chen, Y.-H., Ciais, P., Denning, A. S., Fan, S., Fung, I. Y., Gloor, M., Gurney, K. R., Heimann, M., Higuchi, K., John, J., Law, R. M., Maki, T., Peylin, P., Prather, M., Pak, B., Rayner, P. J., Sarmiento, J. L., Taguchi, S., Takahashi, T., and Yuen, C.-W.: Sensitivity of optimal extension of CO₂ observation networks to model transport, *Tellus B*, 55, 498–511, 2003.
- 35 Patra, P. K., Maksyutov, S., Ishizawa, M., Nakazawa, T., Takahashi, T., and Ukita, J.: Interannual and decadal changes in the sea-air CO₂ flux from atmospheric CO₂ inverse modeling, *Global Biogeochem. Cycles*, 19, GB4013, doi:10.1029/2004GB002257, 2005.
- Patra, P. K., Gurney, K. R., Denning, A. S., Maksyutov, S., Nakazawa, T., Baker, D., Bousquet, P., Bruhwiler, L., Chen, Y.-H., Ciais, P., Fan, S., Fung, I., Gloor, M., Heimann, M., Higuchi, K., John, J., Law, R. M., Maki, T., Pak, B. C., Peylin, P., Prather, M., Rayner, P. J., Sarmiento, J., Taguchi, S., Takahashi, T., and Yuen, C.-W.: Sensitivity of inverse estimation of annual mean CO₂ sources and sinks to ocean-only sites versus all-sites observational networks, *Geophys. Res. Lett.*, 33, 05 814, (doi:10.1029/2005GL025403), 2006.
- 5 Peters, W., Miller, J. B., Whitaker, J., Denning, A. S., Hirsch, A., Krol, M. C., Zupanski, D., Bruhwiler, L., and Tans, P. P.: An ensemble data assimilation system to estimate CO₂ surface fluxes from atmospheric trace gas observations, *J. Geophys. Res.*, 110, 24 304, <https://doi.org/10.1029/2005JD006157>, 2005.
- Peters, W., Krol, M., Van Der Werf, G., Houweling, S., Jones, C., Hughes, J., Schaefer, K., Masarie, K., Jacobson, A., Miller, J., et al.: Seven years of recent European net terrestrial carbon dioxide exchange constrained by atmospheric observations, *Global Change Biology*, 16, 1317–1337, 2010.
- 10 Peylin, P., Rayner, P. J., Bousquet, P., Carouge, C., Hourdin, F., Heinrich, P., Ciais, P., and AEROCARB Contributors: Daily CO₂ flux estimates over Europe from continuous atmospheric measurements: 1, inverse methodology, *Atmospheric Chemistry and Physics*, 5, 3173–3186, <https://doi.org/10.5194/acp-5-3173-2005>, <http://www.atmos-chem-phys.net/5/3173/2005/>, 2005.
- 15 Quaife, T., Lewis, P., De Kauwe, M., Williams, M., Law, B. E., Disney, M., and Bowyer, P.: Assimilating canopy reflectance data into an ecosystem model with an Ensemble Kalman Filter, *Remote Sensing of Environment*, 112, 1347–1364, 2008.
- Rayner, P.: Data Assimilation using an Ensemble of Models: A hierarchical approach, *Atmospheric Chemistry and Physics Discussions*, 2017, 1–16, <https://doi.org/10.5194/acp-2017-10>, <https://www.atmos-chem-phys-discuss.net/acp-2017-10/>, 2017a.

- Rayner, P.: Data Assimilation using an Ensemble of Models: A hierarchical approach, *Atmospheric Chemistry and Physics Discussions*, 2017, 1–16, <https://doi.org/10.5194/acp-2017-10>, <http://www.atmos-chem-phys-discuss.net/acp-2017-10/>, 2017b.
- Rayner, P. J.: The Current State of Carbon Cycle Data Assimilation, *Curr. Opin. Environ. Sustain.*, 2, 289–296, <https://doi.org/10.1016/j.cosust.2010.05.005>, 2010.
- Rayner, P. J., Enting, I. G., Francey, R. J., and Langenfelds, R. L.: Reconstructing the recent carbon cycle from atmospheric CO₂, $\delta^{13}\text{C}$ and O₂/N₂ observations, *Tellus*, 51B, 213–232, 1999.
- 25 Rayner, P. J., Scholze, M., Knorr, W., Kaminski, T., Giering, R., and Widmann, H.: Two decades of terrestrial Carbon fluxes from a Carbon Cycle Data Assimilation System (CCDAS), *Glob. Biogeochem. Cyc.*, 19, 2026, <https://doi.org/10.1029/2004GB002254>, 2005.
- Rayner, P. J., Raupach, M. R., Paget, M., Peylin, P., and Koffi, E.: A new global gridded dataset of CO₂ emissions from fossil fuel combustion: 1: Methodology and evaluation, *J. Geophys. Res.*, 115, D19 306, <https://doi.org/10.1029/2009JD013439>, 2010.
- Robert, C. P. and Casella, G.: *Monte Carlo Statistical Methods*, chap. 2.3.2, pp. 51–53, Springer, 2 edn., 2004.
- 30 Rödenbeck, C.: Estimating CO₂ sources and sinks from atmospheric mixing ratio measurements using a global inversion of atmospheric transport, no. 06 in *Max Planck Institute for biogeochemistry Technical Reports*, Max Planck Society, 2005.
- Rödenbeck, C., Houweling, S., Gloor, M., and Heimann, M.: Time-dependent atmospheric CO₂ inversions based on interannually varying tracer transport, *Tellus*, 55B, 488–497, 2003.
- Rödenbeck, C., Gerbig, C., Trusilova, K., and Heimann, M.: A two-step scheme for high-resolution regional atmospheric trace gas inversions based on independent models, *Atmospheric Chemistry and Physics*, 9, 5331–5342, 2009.
- 35 Rodgers, C. D.: *Inverse methods for atmospheric sounding: Theory and practice*, World Scientific Publishing Co. Pte. Ltd. 2000.
- Rodgers, C. D. and Walshaw, C. D.: The computation of infra-red cooling rate in planetary atmosphere, *Quart. J. Roy. Meteor. Soc.*, 92, 67–92, 1966.
- Roy, T., Rayner, P., Mearns, R., and Francey, R.: Southern hemisphere ocean CO₂ uptake: reconciling atmospheric and oceanic estimates, *Tellus*, 55B, 701–710, doi:10.1034/j.1600-0560.2003.00058.x, 2003.
- Saeki, T. and Patra, P. K.: Implications of overestimated anthropogenic CO₂ emissions on East Asian and global land CO₂ flux inversion, *Geoscience Letters*, 4, 9, <https://doi.org/10.1186/s40562-017-0074-7>, <https://doi.org/10.1186/s40562-017-0074-7>, 2017.
- 5 Salsburg, D.: *The lady tasting tea: How statistics revolutionized science in the twentieth century*, Macmillan, 2001.
- Scholze, M., Kaminski, T., Rayner, P., Knorr, W., and Geiring, R.: Propagating uncertainty through prognostic CCDAS simulations, *J. Geophys. res.*, 112, d17 305, doi:10.1029/2007JD008642, 2007.
- Scholze, M., Kaminski, T., Knorr, W., Blessing, S., Vossbeck, M., Grant, J., and Scipal, K.: Simultaneous assimilation of SMOS soil moisture and atmospheric CO₂ in-situ observations to constrain the global terrestrial carbon cycle, *Remote sensing of environment*, 180, 334–345, 10 2016.
- Scholze, M., Buchwitz, M., Dorigo, W., Guanter, L., and Quegan, S.: Reviews and syntheses: Systematic Earth observations for use in terrestrial carbon cycle data assimilation systems, *Biogeosciences*, 14, 3401–3429, <https://doi.org/10.5194/bg-14-3401-2017>, <https://www.biogeosciences.net/14/3401/2017/>, 2017.
- Schuh, A. E., Lauvaux, T., West, T. O., Denning, A. S., Davis, K. J., Miles, N., Richardson, S., Uliasz, M., Lokupitiya, E., Cooley, D., Andrews, A., and Ogle, S.: Evaluating atmospheric CO₂ inversions at multiple scales over a highly inventoried agricultural landscape, *Global Change Biology*, 19, 1424–1439, <https://doi.org/10.1111/gcb.12141>, <http://dx.doi.org/10.1111/gcb.12141>, 2013.
- 15 Shaby, B. A. and Field, C. B.: Regression tools for CO₂ inversions: application of a shrinkage estimator to process attribution, *Tellus B: Chemical and Physical Meteorology*, 58, 279–292, 2006.

- Shiga, Y. P., Tadić, J. M., Qiu, X., Yadav, V., Andrews, A. E., Berry, J. A., and Michalak, A. M.: Atmospheric CO₂ Observations Reveal
20 Strong Correlation Between Regional Net Biospheric Carbon Uptake and Solar-Induced Chlorophyll Fluorescence, *Geophysical Research Letters*, pp. n/a–n/a, <https://doi.org/10.1002/2017GL076630>, <http://dx.doi.org/10.1002/2017GL076630>, 2017GL076630, 2018.
- Soon-Young, P., Dong-Hyeok, K., Soon-Hwan, L., and Lee, H. W.: Variational data assimilation for the optimized ozone initial state and the short-time forecasting, *Atmospheric Chemistry and Physics*, 16, 3631, 2016.
- Stordal, A. S., Karlsen, H. A., Nævdal, G., Skaug, H. J., and Vallès, B.: Bridging the ensemble Kalman filter and particle filters: the adaptive
25 Gaussian mixture filter, *Computational Geosciences*, 15, 293–305, 2011.
- Tans, P. P., Fung, I. Y., and Takahashi, T.: Observational constraints on the global atmospheric CO₂ budget, *Science*, 247, 1431–1438, 1990.
- Tarantola, A.: *Inverse Problem Theory: Methods for Data Fitting and Parameter Estimation*, Elsevier, Amsterdam, 1987.
- Tarantola, A.: *Inverse problem theory and methods for model parameter estimation*, siam, 2005.
- Tarantola, A. and Valette, B.: Generalized nonlinear inverse problem solved using the least squares criterion, *Reviews of Geophysics*, 20,
30 219–232, 1982.
- Thompson, R. L., Gerbig, C., and Rödenbeck, C.: A Bayesian inversion estimate of N₂O emissions for western and central Europe and the assessment of aggregation errors, *Atmospheric Chemistry and Physics*, 11, 3443–3458, <https://doi.org/10.5194/acp-11-3443-2011>, <http://www.atmos-chem-phys.net/11/3443/2011/>, 2011.
- Tippett, M. K., Anderson, J. L., Bishop, C. H., Hamill, T. M., and Whitaker, J. S.: Ensemble square root filters*, *Monthly Weather Review*,
35 131, 1485–1490, 2003.
- Trampert, J. and Snieder, R.: Model Estimations Biased by Truncated Expansions: Possible Artifacts in Seismic Tomography, *Science*, 271, 1257–1260, 1996.
- Trudinger, C., Raupach, M., Rayner, P., Kattge, J., Liu, Q., Pak, B., Reichstein, M., Renzullo, L., Richardson, A. D., Roxburgh, S., Styles, J., Wang, Y. P., Briggs, P., Barrett, D., and Nikolova, S.: The OptIC project: An intercomparison of optimisation techniques for parameter estimation in terrestrial biogeochemical models, *J. Geophys. Res.*, 112, 2027, <https://doi.org/10.1029/2006JG000367>, 2007.
- Trudinger, C. M., Enting, I. G., and Rayner, P. J.: Kalman filter analysis of ice core data: 1. Method development and testing the statistics, *J. Geophys. Res.*, 107(D20), <https://doi.org/10.1029/2001JD001111>, 2002a.
- Trudinger, C. M., Enting, I. G., Rayner, P. J., and Francey, R. J.: Kalman filter analysis of ice core data: 2. Double deconvolution of CO₂ and δ¹³C measurements, *J. Geophys. Res.*, 107(D20), <https://doi.org/10.1029/2001JD001112>, 2002b.
- Trudinger, C. M., Raupach, M. R., Rayner, P. J., and Enting, I. G.: Using the Kalman filter for parameter estimation in biogeochemical models, *Environmetrics*, 19, 849–870, <https://doi.org/10.1002/env.910>, 2008.
- 10 Turnbull, J. C., Karion, A., Fischer, M. L., Faloona, I., Guilderson, T., Lehman, S. J., Miller, B. R., Miller, J. B., Montzka, S., Sherwood, T., Saripalli, S., Sweeney, C., and Tans, P. P.: Assessment of fossil fuel carbon dioxide and other anthropogenic trace gas emissions from airborne measurements over Sacramento, California in spring 2009, *Atmospheric Chemistry and Physics*, 11, 705–721, <https://doi.org/10.5194/acp-11-705-2011>, <http://www.atmos-chem-phys.net/11/705/2011/>, 2011.
- Turner, A. and Jacob, D.: Balancing aggregation and smoothing errors in inverse models, *Atmospheric Chemistry and Physics*, 15, 7039–
15 7048, 2015.
- van Dop, H., Addis, R., Fraser, G., Girardi, F., Graziani, G., Inoue, Y., Kelly, N., Klug, W., Kulmala, A., Nodop, K., and Pretel, J.: ETEX A European tracer experiment observations, dispersion modelling and emergency response, *Atmospheric Environment*, 32, 4089–4094, 1998.

- van Leeuwen, P. J.: Particle Filtering in Geophysical Systems, *Monthly Weather Review*, 137, 4089–4114, <https://doi.org/10.1175/2009MWR2835.1>, 2009.
- 20 Vanoye, A. Y. and Mendoza, A.: Application of direct regularization techniques and bounded–variable least squares for inverse modeling of an urban emissions inventory, *Atmospheric Pollution Research*, 5, 219–225, 2014.
- Wang, J. S., Kawa, S. R., Collatz, G. J., Sasakawa, M., Gatti, L. V., Machida, T., Liu, Y., and Manyin, M. E.: A global synthesis inversion analysis of recent variability in CO₂ fluxes using GOSAT and in situ observations, *Atmospheric Chemistry and Physics*, 18, 11 097–11 124, <https://doi.org/10.5194/acp-18-11097-2018>, <https://www.atmos-chem-phys.net/18/11097/2018/>, 2018.
- 25 Wang, X., Mallet, V., Berroir, J.-P., and Herlin, I.: Assimilation of OMI NO₂ retrievals into a regional chemistry-transport model for improving air quality forecasts over Europe, *Atmospheric environment*, 45, 485–492, 2011.
- Wikle, C. K. and Berliner, L. M.: A Bayesian tutorial for data assimilation, *Physica D: Nonlinear Phenomena*, 230, 1–16, 2007.
- Williams, M., Schwarz, P. A., Law, B. E., Irvine, J., and Kurpius, M. R.: An improved analysis of forest carbon dynamics using data assimilation, *Glob. Change Biol.*, 11, 89–105, 2005.
- Wilson, C., Chipperfield, M., Gloor, M., and Chevallier, F.: Development of a variational flux inversion system (INVICAT v1. 0) using the TOMCAT chemical transport model, *Geoscientific Model Development*, 7, 2485–2500, 2014.
- 1090 Wu, L., Bocquet, M., Lauvaux, T., Chevallier, F., Rayner, P., and Davis, K.: Optimal Representation of Source-Sink Fluxes for Mesoscale Carbon Dioxide Inversion with Synthetic Data, *J. Geophys. Res.*, 116, D21 304, <https://doi.org/10.1029/2011JD016198>, 2011.
- Wu, L., Bocquet, M., Chevallier, F., Lauvaux, T., and Davis, K.: Hyperparameter Estimation for Uncertainty Quantification in Mesoscale Carbon Dioxide Inversions, *Tellus*, 65b, accepted, 2013.
- 1095 Wunsch, C. and Minster, J.-F.: Methods for box models and ocean circulation tracers: Mathematical programming and non-linear inverse theory, *J. Geophys. Res.*, 87, 5647–5662, 1982.
- Yadav, V. and Michalak, A.: Improving computational efficiency in large linear inverse problems: An example from carbon dioxide flux estimation, *Geoscientific Model Development*, 6, 583–590, 2013.
- Zheng, B., Chevallier, F., Ciais, P., Yin, Y., Deeter, M. N., Worden, H. M., Wang, Y., Zhang, Q., and He, K.: Rapid decline in carbon monoxide emissions and export from East Asia between years 2005 and 2016, *Environmental Research Letters*, 13, 044 007, 2018a.
- 1100 Zheng, T., French, N. H., and Baxter, M.: Development of the WRF-CO₂ 4D-Var assimilation system v1. 0, *Geoscientific Model Development*, 11, 1725–1752, 2018b.
- Zupanski, D., Denning, A. S., Uliasz, M., Zupanski, M., Schuh, A. E., Rayner, P. J., Peters, W., and Corbin, K.: Carbon flux bias estimation employing Maximum Likelihood Ensemble Filter (MLEF), *J. Geophys. Res.*, 112, d17 107, [doi:10.1029/2006JD008371](https://doi.org/10.1029/2006JD008371), 2007.