

Interactive comment on “Evaluating the Recent “2+26” Regional Strategy for Air Quality Improvement During Two Orange Air Pollution Alerts in Beijing: variations of PM_{2.5} concentrations, source apportionment, and the relative contribution of local emission and regional transport” by Ziyue Chen et al.

Anonymous Referee #1

Received and published: 21 December 2018

General comments: This manuscript attempts to evaluate the effects of “2+26” regional integrative strategy on air quality improvement in China, by comparing the variation of PM_{2.5} concentrations and source contribution in Beijing during four pollution episodes. The study uses airborne pollutants modeled by the WRF-CAMx model and the observed PM_{2.5} to understand how “2+26” strategy can affect the PM_{2.5} reduc-

C1

tion during the pollution episodes. The topic is interesting because the impact of emission-reduction on air quality has been still unclear, and the results were very helpful to control the air pollution in cities. However, the authors have not made best use of the model results. The model can provide information on meteorological condition, emission and regional distribution of pollutants. These information should be supplemented and analyzed to know the influence of meteorological condition on the implementation of “2+26” strategy. There needs to be significant improvement in results interpretation and more analysis needs to be done to know the control factors of variation of PM_{2.5} and SNA. I would thus recommend a major revision to improve this manuscript.

Specific comments: Q1. In Discussion Section, the authors stated the influence of meteorological factors on PM_{2.5} concentration and explained the different effects of “2+26” strategy on PM_{2.5} reduction for different pollution episodes. Using WRF-CAMx model, author can obtain the detailed meteorological field. However, only average RH and wind speed for four pollution episodes was shown in Table 2. Then how authors determine the airborne pollutants in Beijing was transported from neighboring areas during pollution episodes? There is also no meteorological data under unpolluted weathers (AQI < 50), how can we know the meteorological difference between pollution and clean days? I can't follow “a high-humidity condition” without comparison with a background value. In Line 438-444, the author pointed that “although unified emission-reduction measures were implemented in its neighboring areas, the significantly restricted regional transport did not fully project the effect of the “2+26” strategy to the local PM_{2.5} concentrations in Beijing”, so to what extent can meteorological condition affect the implementation of “2+26” strategy? And under what circumstances the meteorological condition will have important effect on implementation? Q2. There is no detailed information of meteorological parameters and concentration of SO₂, NO_x, and NH₃ during four pollution episodes. However, the SNA formation is related with the precursor. I think the related information should be supplemented and analyzed before comparison of PM_{2.5} reduction. Q3. In Introduction Section, the authors explained the

C2

specific emission-reduction measures in detail. However, there is no emission data for airborne pollutants, such as SO₂, NO_x, NH₃, dust, etc., from 2013 to 2018. Whether the emission of all these precursor gas was really reduced greatly by implementing “2+26” strategy? I suggested that the authors analyze the emission variation in detail to evaluate the “2+26” strategy. Q4. Section 2.2.2. Supplement the sampling duration, sample numbers and the membrane to collected PM_{2.5}. Was the sampling duration 15min for ions and 1 hour for OC/EC? Q5. Revise the Section 2.3.1 to make the description of WRF-CAMx more concise Q6. Section 2.3.1 In this manuscript, the authors simulated several episodes during 2013-2017. During these years, emissions in China changed obviously due to lots of national strategies. Emission inventory is an important factor which would influence model results. So please clarify which years’ emission inventories were used in this study? Did you consider “coal to gas” strategy in your emission inventory? Q7. L207-209. The input and output of CMAx is in binary format. However, output from MCIP is in NETCDF format, please clarify how to use NETCDF meteorological data in CMAx? Q8. Section 2.4 The authors explained that meteorological parameters contributed to the underprediction of simulated PM_{2.5}, could you give out some information about the model performance of meteorological parameters such as T, RH, WS, WD? Q9. The author found that composition of PM_{2.5} changed obviously due to the national strategies, therefore it is important to show the model performance of inorganic components in PM_{2.5} such as SO₄²⁻, NO₃⁻, NH₄⁺ but not only show the result of PM_{2.5}. If the model performance is satisfied, further analysis of PSAT would be reasonable, otherwise, results of PSAT would not be convincing. Q10. Supplement the criteria or error index that can verify the satisfactory simulation for PSAT. Q11. In Section 3.2, only the variation of ions in PM_{2.5} was discussed. Organic compounds are one of major components of PM_{2.5}. Since the OC/EC has been analyzed, I think the OC variation should be discussed here. Q12. From Fig. 4b, very high concentration of NO₂⁻ was observed during the pollution episode in March, 2018. The value is very abnormal, almost two times higher than NO₃⁻. In general, nitrite shows very low concentration in atmospheric aerosols and contributes little to

C3

water soluble inorganic ions. What’s the reason for this abnormal value? I think the authors should check the data and discussed the reason. Q13. L320-321. Which data can support the “The main source for NO₃⁻ is vehicle exhaust” in Beijing? How did you verified the vehicle exhaust was main sources of NO₃⁻ in Beijing just as the cited reference suggested in other cities? I think the source appointment of NO₃⁻ will be helpful to support your suggestion on vehicle exhaust in conclusion. Q14. L320-321. As “The main source for NO₃⁻ is vehicle exhaust” and the vehicles that cannot meet the Environmental Levels I and II was forbidden during orange alerts, why the concentration of NO₃⁻ was much higher during orange alerts in Mar, 2018 than that in March, 2013 without emission-reduction (Table 4)? Increased NO₃⁻ corresponded to decreased concentration of NH₄⁺ during pollution episodes, so what’s possible existing form of NO₃⁻ in PM_{2.5}? Q15. L364-365. Please clarify what changes have been made to the air pollutants emission after “Coal to Gas”. Q16. L378-383. According to Fig.6, the local emission contributed 49.46% - 88.35% to PM_{2.5} during four pollution episode, indicating the local emission had a great effect on PM_{2.5} in Beijing. This is contradicting L92-93. And the different emission-reduction strategies did not lead to a clear pattern for the regional transport. So the PM_{2.5} reduction really was a result of “2+26” strategies or the meteorological condition? Whether the strict regulation on vehicle exhaust will be more effective than that of regional emission control under specific wind direction? The meteorological condition should be analyzed in detail for each pollution episode. Q17. Overall, some of the conclusions on page 20 appear to be speculation with little data or discussion to support it, such as L494-495. Analysis and discussion on regional distribution of PM_{2.5} needs to be supplemented.

Technical corrections: L139. Supplement the link of website PM25.in. It’s difficult to follow. L164. Change “ãÄÄ” to “,”. L183. Change “*” to “iCt”. P12. Fig.4, change “NO2-” to “NO2-”

Please also note the supplement to this comment:

<https://www.atmos-chem-phys-discuss.net/acp-2018-1085/acp-2018-1085-RC1->

C4

supplement.pdf

Interactive comment on Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2018-1085>, 2018.