1 The role of chlorine in global tropospheric chemistry

Xuan Wang¹, Daniel J. Jacob^{1,2}, Sebastian D. Eastham³, Melissa P. Sulprizio¹, Lei Zhu¹, Qianjie
Chen⁴, Becky Alexander⁵, Tomás Sherwen^{6,7}, Mathew J. Evans^{6,7}, Ben H. Lee⁵, Jessica D.
Haskins⁵, Felipe D. Lopez-Hilfiker⁸, Joel A. Thornton⁵, Gregory L. Huey⁹, and Hong Liao¹⁰
¹School of Engineering and Applied Sciences, Harvard University, Cambridge, Massachusetts, USA
²Department of Earth and Planetary Sciences, Harvard University, Cambridge, Massachusetts, USA
³Laboratory for Aviation and the Environment, Massachusetts Institute of Technology, Cambridge, Massachusetts,

- 9 USA
- ⁴ Department of Chemistry, University of Michigan, Ann Arbor, Michigan, USA
- ⁵ Department of Atmospheric Sciences, University of Washington, Seattle, USA
- ⁶ Wolfson Atmospheric Chemistry Laboratories, Department of Chemistry, University of York, York, UK
- ¹³ ⁷National Centre for Atmospheric Science, University of York, York, UK
- ⁸ Paul Scherrer Institute, Villigen, Switzerland
- ⁹ School of Earth and Atmospheric Science, Georgia Institute of Technology, Atlanta, GA, USA
- ¹⁰ School of Environmental Science and Engineering, Nanjing University of Information Science and Technology,
- 17 Nanjing, China
- 18 Correspondence to: Xuan Wang (wangx@seas.harvard.edu)

19 Abstract. We present a comprehensive simulation of tropospheric chlorine within the GEOS-Chem global 3-D model 20 of oxidant-aerosol-halogen atmospheric chemistry. The simulation includes explicit accounting of chloride 21 mobilization from sea-salt aerosol by acid displacement of HCl and by other heterogeneous processes. Additional 22 small sources of tropospheric chlorine (combustion, organochlorines, transport from stratosphere) are also included. 23 Reactive gas-phase chlorine Cl*, including Cl, ClO, Cl₂, BrCl, ICl, HOCl, ClNO₃, ClNO₂, and minor species, is 24 produced by the HCl + OH reaction and by heterogeneous conversion of sea-salt aerosol chloride to BrCl, ClNO₂, Cl₂, 25 and ICl. The model simulates successfully the observed mixing ratios of HCl in marine air (highest at northern mid-26 latitudes) and the associated HNO₃ decrease from acid displacement. It captures the high ClNO₂ mixing ratios 27 observed in continental surface air at night, and attributes the chlorine to HCl volatilized from sea salt aerosol and 28 transported inland following uptake by fine aerosol. The model simulates successfully the vertical profiles of HCl 29 measured from aircraft, where enhancements in the continental boundary layer can again be largely explained by 30 transport inland of the marine source. It does not reproduce the boundary layer Cl₂ mixing ratios measured in the 31 WINTER aircraft campaign (1-5 ppt in the daytime, low at night); the model is too high at night, which could be due 32 to uncertainty in the rate of the $CINO_2 + CI^-$ reaction, but we have no explanation for the high observed Cl_2 in daytime. 33 The global mean tropospheric concentration of Cl atoms in the model is 620 cm⁻³ and contributes 1.0% of the global 34 oxidation of methane, 20% of ethane, 14% of propane, and 4% of methanol. Chlorine chemistry increases global mean 35 tropospheric BrO by 85%, mainly through the HOBr + Cl^{-} reaction, and decreases global burdens of tropospheric 36 ozone by 7% and OH by 3% through the associated bromine radical chemistry. ClNO₂ chemistry drives increases in

37 ozone of up to 8 ppb over polluted continents in winter.

1 1 Introduction

2 Mobilization of chloride (Cl⁻) from sea salt aerosol (SSA) is a large source of chlorine gases to the troposphere

3 (Graedel and Keene, 1995; Finlayson-Pitts, 2003). These gases may generate chlorine radicals with a broad range of

4 implications for tropospheric chemistry including the budgets of ozone, OH (the main tropospheric oxidant), volatile

6 Simpson et al., 2015). Only a few global models have attempted to examine the implications of tropospheric chlorine

organic compounds (VOCs), nitrogen oxides, other halogens, and mercury (Saiz-Lopez and von Glasow, 2012;

7 chemistry on a global scale (Singh and Kasting, 1988; Long et al., 2014; Hossaini et al., 2016). Here we present a

8 more comprehensive analysis of this chemistry within the framework of the GEOS-Chem chemical transport model

9 (CTM).

5

10 Saiz-Lopez and von Glasow (2012) and Simpson et al. (2015) present recent reviews of tropospheric halogen 11 chemistry including chlorine. Sea-salt aerosols represent a large chloride flux to the atmosphere but most of that 12 chloride is removed rapidly by deposition. Only a small fraction is mobilized to the gas phase as HCl or other species. 13 Additional minor sources of tropospheric chlorine include open fires, coal combustion, waste incineration, industry, 14 road salt application, fugitive dust, and ocean emission of organochlorine compounds (Lobert et al., 1999; McCulloch 15 et al., 1999; Sarwar et al., 2012; WMO, 2014; Kolesar et al., 2018). It is useful to define Cly as total gas-phase inorganic 16 chlorine, excluding particle phase Cl⁻. Most of this Cl_v is present as HCl, which is removed rapidly by deposition but 17 also serves as a source of chlorine radicals. Rapid cycling takes place between the chlorine radicals and other chlorine 18 gases, eventually returning HCl. Thus it is useful to define reactive chlorine Cl* as the ensemble of Cly gases other

19 than HCl.

20 Cycling of chlorine affects tropospheric chemistry in a number of ways (Finlayson-Pitts, 2003;Saiz-Lopez and von 21 Glasow, 2012). Acid displacement of Cl⁻ by nitric acid (HNO₃) is a source of NO₃⁻ aerosol (Massucci et al., 1999). Cl 22 atoms provide a sink for methane, other volatile organic compounds (VOCs) (Atkinson, 1997), dimethyl sulfide (DMS) 23 (Hoffmann et al., 2016; Chen et al., 2017), and mercury (Horowitz et al., 2017). Cycling between Cl radicals and their 24 reservoirs drives catalytic ozone loss, and converts nitrogen oxide radicals (NO_x \equiv NO + NO₂) to HNO₃, decreasing 25 both ozone and OH. On the other hand, aqueous-phase reaction of Cl⁻ with N₂O₅ in polluted environments produces 26 ClNO₂ radicals that photolyze in the daytime to return Cl atoms and NO₂, stimulating ozone production (Behnke et 27 al., 1997; Osthoff et al., 2008). Chlorine also interacts with other halogens (bromine, iodine), initiating further radical 28 chemistry that affects ozone, OH, and mercury.

A number of global modeling studies have investigated tropospheric halogen chemistry but most have focused on bromine and iodine, which are more active than chlorine because of the lower chemical stability of HBr and HI (Yang et al., 2005; Saiz-Lopez et al., 2014). Interest in global modeling of tropospheric chlorine has focused principally on quantifying the Cl atom concentration as a sink for methane (Keene et al., 1990; Singh et al., 1996). Previous global 3-D models found mean tropospheric Cl atom concentrations of the order of 10³ cm⁻³, with values up to 10⁴ cm⁻³ in the marine boundary layer (MBL), and contributing 2-3% of atmospheric methane oxidation (Long et al., 2014; Hossaini et al., 2016; Sherwen et al., 2016b; Schmidt et al., 2016). A regional modeling study by Sarwar et al. (2014) 1 included ClNO₂ chemistry in a standard ozone mechanism and found increases in surface ozone mixing ratios over 2 the US of up to 7 ppb ($ppb \equiv nmol mol^{-1}$).

3 Here we present a more comprehensive analysis of global tropospheric chlorine chemistry and its implications, 4 building on previous model development of oxidant-aerosol-halogen chemistry in GEOS-Chem. A first capability for 5 modeling tropospheric bromine in GEOS-Chem was introduced by Parrella et al. (2012). Eastham et al. (2014) 6 extended it to describe stratospheric halogen chemistry including chlorine and bromine cycles. Schmidt et al. (2016) 7 updated the tropospheric bromine simulation to include a broader suite of heterogeneous processes, and extended the 8 Eastham et al. (2014) stratospheric chlorine scheme to the troposphere. Sherwen et al. (2016a;b) added iodine 9 chemistry and made further updates to achieve a consistent representation of tropospheric chlorine-bromine-iodine 10 chemistry in GEOS-Chem. Chen et al. (2017) added the aqueous-phase oxidation of SO₂ by HOBr and found a large 11 effect on the MBL bromine budget. Our work advances the treatment of tropospheric chlorine in GEOS-Chem to 12 include in particular a consistent treatment of SSA chloride and chlorine gases, SSA acid displacement 13 thermodynamics, improved representation of heterogeneous chemistry, and better accounting of chlorine sources. We 14 evaluate the model with a range of global observations for chlorine and related species. From there we quantify the 15 global tropospheric chlorine budgets, describe the principal chemical pathways, and explore the impacts on 16 tropospheric chemistry.

17 2 Model description

18 2.1 GEOS-Chem model with Cl+Br+I halogen chemistry

19 We build our new tropospheric chlorine simulation capability onto the standard version 11-02d of GEOS-Chem 20 (http://www.geos-chem.org). The standard version includes a detailed tropospheric oxidant-aerosol-halogen 21 mechanism as described by Sherwen et al. (2016b) and Chen et al. (2017). It includes 12 gas phase Cl_v species: Cl_v 22 Cl₂, Cl₂O₂, ClNO₂, ClNO₃, ClO, ClOO, OClO, BrCl, ICl, HOCl, and HCl. It allows for heterogeneous chemistry 23 initiated by SSA Cl⁻ but does not actually track the SSA Cl⁻ concentration and its exchange with gas-phase Cl_v. Here 24 we add two new transported reactive species to GEOS-Chem to describe Cl⁻ aerosol, one for the fine mode (< 1µm 25 diameter) and one for the coarse mode (> 1µm diameter). The standard GEOS-Chem wet deposition schemes for 26 water-soluble aerosols (Liu et al., 2001) and gases (Amos et al., 2012) are applied to Cl⁻ aerosol and Cl_y gases 27 respectively, the latter with Henry's law constants from Sander (2015). Dry deposition of Cl⁻ aerosol follows that of 28 SSA (Jaeglé et al., 2011), and dry deposition of Cl_v gases follows the resistance-in-series scheme of Wesely (1989) as 29 implemented in GEOS-Chem by Wang et al. (1998). We also add to the model two SSA alkalinity tracers in the fine 30 and coarse modes, and retain the inert SSA tracer to derive local concentrations of non-volatile SSA cations (Section 31 2.3). SSA debromination by oxidation of Br⁻ as described by Schmidt et al. (2016) was included only as an option in 32 standard version 11-02d of GEOS-Chem because of concern over excessive MBL BrO (Sherwen et al., 2016b). 33 However, Zhu et al. (2018) show that it allows in fact a successful simulation of MBL BrO when one accounts for 34 new losses in the standard model from aqueous-phase oxidation of SO₂ by HOBr (Chen et al., 2017) and oxidation of 35 marine acetaldehyde by Br atoms (Millet et al., 2010). We include SSA debromination in this work.

- 1 We present a 1-year global simulation for 2016 driven by GEOS-FP (forward processing) assimilated meteorological
- 2 fields from the NASA Global Modeling and Assimilation office (GMAO) with native horizontal resolution of 0.25°
- 3 $\times 0.3125^{\circ}$ and 72 vertical levels from the surface to the mesosphere. Our simulation is conducted at 4° \times 5° horizontal
- 4 resolution and meteorological fields are conservatively regridded for that purpose. Stratospheric chemistry is
- 5 represented using 3-D monthly mean production rates and loss rate constants from a fully coupled stratosphere-
- 6 troposphere GEOS-Chem simulation (Murray et al., 2012; Eastham et al., 2014).

7 2.2 Sources of chlorine

- 8 Table 1 lists the global sources and sinks of tropospheric gas-phase inorganic chlorine (Cl_y) and reactive chlorine (Cl*)
- 9 in our model. The main source is mobilization of Cl⁻ from SSA. SSA emission is computed locally in GEOS-Chem
- 10 separately for fine and coarse as the integrals of the size-dependent source function over two size bins, fine (0.2-1 µm
- 11 diameter) and coarse (1-8 µm diameter). The source function depends on wind speed and sea surface temperature
- 12 (Jaeglé et al., 2011). We obtain a global SSA source of 3230 Tg a⁻¹ for 2016, corresponding to 1780 Tg a⁻¹ Cl⁻
- 13 (assuming fresh SSA to be 55.05% Cl⁻ by mass (Lewis and Schwartz, 2013), of which 16% is in the fine mode and
- 14 84% is in the coarse mode. Only 64 Tg Cl⁻ a^{-1} (3.6%) is mobilized to Cl_y by acid displacement and other heterogeneous
- 15 reactions, while the rest is deposited. 42% of the mobilization is from fine SSA and 58% is from coarse SSA. Details
- 16 of this mobilization are in Sections 2.3 and 2.4. 80% of the mobilization is by acid displacement to HCl, which is in
- 17 turn efficiently deposited. Only 19% of HCl is further mobilized to Cl* by reaction with OH to drive chlorine radical
- 18 chemistry. Direct generation of Cl* from SSA through heterogeneous chemistry provides a Cl* source of comparable
- 19 magnitude to HCl + OH, with dominant contributions from HOBr + Cl^{-} and $N_2O_5 + Cl^{-}$ (the latter in polluted high-
- 20 NO_x environments).
- 21 Cl* can also be produced in the model by atmospheric degradation of the organochlorine gases CH₃Cl, CH₂Cl₂, CHCl₃,
- 22 and CH₂ICl. These gases are mainly of biogenic marine origin, with the exception of CH₂Cl₂ which has a large
- 23 industrial solvent source (Simmonds et al., 2006). Mean tropospheric lifetimes are 520 days for CH₃Cl, 280 days for
- 24 CH₂Cl₂, 260 days for CHCl₃, and 0.4 days for CH₂ICl. Emissions of CH₃Cl, CH₂Cl₂, and CHCl₃ are implicitly treated
- in the model by specifying monthly mean surface air boundary conditions in 5 latitude bands (60-90°N, 30-60°N, 0-
- 26 30°N, 0-30°S, and 30-90°S) from AGAGE observations (Prinn et al., 2018). Emission of CH₂ICl is from Ordóñez et
- al. (2012), as described by Sherwen et al. (2016a). Tropospheric oxidation of hydrochlorofluorocarbons (HCFCs) is
- 28 neglected as a source of Cl* because it is small compared to the other organochlorines. The stratospheric source of
- 29 Cl_y from chlorofluorocarbons (CFCs), HCFCs, and CCl₄ is included in the model on the basis of the Eastham et al.
- 30 (2014) GEOS-Chem stratospheric simulation as described in Section 2.1. Tropospheric organochlorines give a global
- 31 Cl* source of 3. 3 Tg Cl a⁻¹ in Table 1, smaller than that from heterogeneous SSA Cl⁻ reactions (11.9 Tg Cl a⁻¹) or
- 32 oxidation of HCl by OH (9.7 Tg Cl a⁻¹). The stratosphere is a minor global source of tropospheric Cl* (0.06 Tg Cl a⁻¹)
- ¹) although it could be important in the upper troposphere (Schmidt et al., 2016).

- 1 We also include primary HCl emissions from open fires. We apply the emission factors of (HCl + Cl⁻) from Lobert et
- 2 al. (1999) for different vegetation types to the GFED4 (Global Fire Emissions Database) biomass burned inventory
- 3 (van der Werf et al., 2010; Giglio et al., 2013), resulting in a global source of 7.6 Tg Cl a⁻¹ emitted as HCl.
- 4 Anthropogenic sources of HCl include coal combustion, waste incineration, and industrial activities. The only global 5 emission inventory is that of McCulloch et al. (1999), which gives a total of 6.7 Tg Cl a⁻¹. As shown in Section 4.2, 6 we find that this greatly overestimates atmospheric observations of HCl over the US. National inventories of HCl from 7 coal combustion available for China (236 Gg Cl a⁻¹ in 2012; (Liu et al., 2018)) and the US (69 Gg Cl a⁻¹ in 2014; (US 8 EPA, 2018)) are respectively six and seven times lower than McCulloch et al. (1999) for those countries. We choose 9 therefore not to include anthropogenic HCl emissions in our standard simulation, as they are small in any case from a 10 global budget perspective. We show in Section 4.2 that we can account for HCl observations in continental air largely 11 on the basis of the SSA Cl⁻ source. We also do not consider Cl* generation from snow/ice surfaces which could be 12 important in the Arctic spring MBL (Liao et al., 2014) but is highly uncertain and would only affect a small
- 13 atmospheric domain.

We do not include the anthropogenic Cl⁻ source from fugitive dust, although it might be important in contributing to chloride levels in continental surface air (Sarwar et al., 2012). The global source of anthropogenic fugitive dust is

- estimated to be less than 13 Tg a^{-1} (Philip et al., 2017), of which 0.3% by mass is estimated to be chloride (Reff et al.,
- 17 2009). This corresponds to a chloride source of less than 0.39 Tg Cl a^{-1} , negligible on a global scale.

18 2.3 HCl/Cl⁻ acid displacement thermodynamics

SSA Cl⁻ can be displaced to HCl by strong acids (H₂SO₄, HNO₃) once the SSA is sufficiently aged that its initial supply of alkalinity (\equiv HCO₃⁻ + 2×CO₃²⁻) has been exhausted. The acid displacement is described by

21	$Cl^- + HNO_3 \rightleftharpoons HCl + NO_3^-$	(R1)
22	$Cl^- + H_2SO_4 \rightleftharpoons HCl + HSO_4^-$	(R2)

with equilibrium constants from Fountoukis and Nenes (2007). (R1) must be treated as an equilibrium because HNO_3 and HCl have comparable effective Henry's law constants. H_2SO_4 has a much lower vapor pressure so that (R2) fully displaces HCl. Additional displacement of HCl by HSO_4^- does not take place because HSO_4^- is a much weaker acid than HCl (Jacob et al., 1985).

- Alkalinity initially prevents any acid displacement in freshly emitted SSA. Alkalinity is emitted as 0.07 mole equivalents per kg of dry SSA (Gurciullo et al., 1999), and is transported in the model as two separate tracers for fine and coarse SSA. It is consumed over time by uptake of acids (SO₂, H₂SO₄, HNO₃, and HCl) as described by Alexander et al. (2005), and once fully consumed it is set to zero (titration). The SSA is then diagnosed as acidified, enabling acid displacement by (R1)-(R2). In our simulation, alkalinity is titrated everywhere shortly after emission except in
- 32 some areas of the Southern Ocean, which is consistent with the model results of Alexander et al. (2005) and Kasibhatla
- 33 et al. (2018).

- 1 Observations in the MBL indicate that fine SSA is usually internally mixed with sulfate-nitrate-ammonium (SNA)
- 2 aerosols while coarse SSA is externally mixed (Fridlind and Jacobson, 2000; Dasgupta et al., 2007). Acid displacement
- 3 for the acidified fine SSA is thus computed by adding HCl/Cl⁻ to the SNA thermodynamics. The local thermodynamic
- 4 gas-aerosol equilibrium for the resulting H₂SO₄-HCl-HNO₃-NH₃-NVC system is calculated with ISORROPIA II
- 5 (Fountoukis and Nenes, 2007). The calculation is done assuming an aqueous aerosol even if relative humidity is below
- 6 the deliquescence point (metastable state). NVC (non-volatile cations) describes the sum of cations emitted as SSA
- 7 and is treated in ISORROPIA II using Na⁺ as proxy. Here NVC is emitted as 16.4 moles equivalent per kg of dry SSA
- 8 to balance the emission of SSA anions including Cl⁻, alkalinity, and sea-salt sulfate. The NVC concentration is
- 9 determined locally from the mass concentration of the inert SSA tracer.
- 10 Acid displacement for acidified coarse SSA is assumed to be driven by uptake of strong acids from the gas phase,
- 11 mainly HNO₃ (Kasibhatla et al., 2018). The ISORROPIA II calculation is conducted with 2 gas species (HNO₃ and
- HCl) and 4 aerosol species (NVC, Cl⁻, SO₄²⁻, and NO₃⁻). Here the sulfate includes only the emitted sea-salt component

13 and that produced by heterogeneous SO_2 oxidation in coarse SSA (Alexander et al., 2005). In the case of coarse

14 aerosols, there may be significant mass transfer limitation to reaching gas-aerosol thermodynamic partitioning (Meng

- 15 and Seinfeld, 1996). To account for this limitation, the concentrations are adjusted after the ISORROPIA II calculation
- 16 following the dynamic method of Pilinis et al. (2000). This 2-step thermodynamics approach has been used in previous
- 17 studies (Koo et al., 2003; Kelly et al., 2010).

18 2.4 Heterogeneous chemistry of Cl⁻

19 Table 2 lists the heterogeneous reactions of Cl⁻ other than acid displacement. The loss rate of a gas species X due to 20 reaction with Cl⁻ is calculated following Jacob (2000):

$$21 \qquad \frac{dn_X}{dt} = -\left(\frac{r}{D_g} + \frac{4}{c\gamma([\text{Cl}^-])}\right)^{-1} An_X \tag{1}$$

Here n_X is the number density of species X (molecules of X per unit volume of air), *A* is the aerosol or cloud surface area concentration per unit volume of air, *r* is the effective particle radius, D_g is the gas-phase molecular diffusion coefficient of X, *c* is the average gas-phase thermal velocity of X, and γ is the reactive uptake coefficient which is a function of the aqueous-phase molar Cl⁻ concentration [Cl⁻] (moles of Cl⁻ per liter of water). Values of γ in Table 2 are mostly from recommendations by the International Union of Pure and Applied Chemistry (IUPAC) (Ammann et al., 2013).

The heterogeneous reactions take place in both clear-air aerosol and clouds. The GEOS-FP input meteorological data include cloud fraction and liquid/ice water content for every grid cell. Concentrations per cm³ of air of aerosol-phase species (including fine and coarse Cl⁻ and Br⁻) within a grid cell are partitioned between clear air and cloud as determined by the cloud fraction. Clear-air aqueous-phase concentrations for use in calculating heterogeneous reaction rates are derived from the RH-dependent liquid water contents of fine and coarse SSA using aerosol hygroscopic growth factors from the Global Aerosol Database (GADS, (Koepke, 1997)) with update by Lewis and 1 Schwartz (2006). In-cloud aqueous-phase concentrations are derived using liquid and ice water content from the 2 GEOS-FP meteorological data. Values of r in equation (1) are specified as RH-dependent effective radii for the

- 3 different clear-air aerosol components (Martin et al., 2003), and are set to 10 µm for cloud droplets and 75 µm for ice
- 4 particles. These effective radii are also used to infer the area concentrations A on the basis of the mass concentrations.
- 5
- Heterogeneous chemistry in ice clouds is restricted to the unfrozen layer coating the ice crystal, which is assumed to
- 6 be 1% of the ice crystal radius (Schmidt et al., 2016).
- 7 The heterogeneous uptake of HOBr, HOCl, and ClNO₂ as well as further aqueous-phase reaction with Cl⁻ in Table 2
- 8 are pH-dependent, with a higher efficiency in acidic solutions (Fickert et al., 1999; Roberts et al., 2008; Abbatt et al.,
- 9 2012). They are considered only when SSA alkalinity has been titrated, and $CINO_2 + CI^-$ further requires pH <2,
- 10 following laboratory results presented in Roberts et al. (2008). The pH of chloride-containing fine aerosol after
- 11 alkalinity has been titrated is calculated by ISORROPIA II. Liquid cloud water pH is calculated in GEOS-Chem
- 12 following Alexander et al. (2012), with update to include Cl⁻ and NVC. Coarse-mode SSA and ice cloud pH are
- 13 assumed to be 5 and 4.5 respectively (Schmidt et al., 2016).

14 **3** Global budget and distribution of tropospheric chlorine

- 15 Figure 1 describes the global budget and cycling of tropospheric inorganic chlorine in GEOS-Chem. The dominant
- 16 source of Cl_y is acid displacement from SSA. The global rate of Cl_y generation from acid displacement is 52 Tg Cl a
- ¹, close to the observationally based estimate of 50 Tg Cl a^{-1} by Graedel and Keene (1995), and lower than the model 17
- 18 estimate of 90 Tg Cl a^{-1} from Hossaini et al. (2016), who treated displacement of Cl⁻ by HNO₃ as an irreversible rather
- 19 than thermodynamic equilibrium process. HCl is the largest reservoir of Cl_v in the troposphere, with a global mean
- 20 tropospheric mixing ratio of 60 ppt (ppt \equiv pmol/mol).
- 21 Acid displacement generates Cl_y as HCl, which is mostly removed by deposition. Broader effects of chlorine on 22 tropospheric chemistry take place through the cycling of radicals originating from production of reactive chlorine Cl* 23 \equiv Cl_v – HCl. HCl contributes 9.7 Tg Cl a⁻¹ to Cl* through the reaction between HCl and OH. Beside this source, Cl⁻ provides a Cl* source of 12 Tg Cl a⁻¹ through heterogeneous reactions with principal contributions from HOBr + Cl⁻ 24 $(8.6 \text{ Tg Cl a}^{-1})$ and $N_2O_5 + \text{Cl}^{-1}(1.8 \text{ Tg Cl a}^{-1})$. This heterogeneous source of 12 Tg Cl a⁻¹ is much higher than previous 25 26 estimates of 5.6 Tg Cl a⁻¹ (Hossaini et al., 2016) and 6.1 Tg Cl a⁻¹ (Schmidt et al., 2016). Schmidt et al. (2016) only 27 considered the HOBr + Cl⁻ reaction. Production of the chlorine radicals Cl and ClO is by the HCl + OH reaction (45%) 28 and by photolysis of BrCl (40%), ClNO₂ (8%), Cl₂ (4%), and ICl (2%). Loss of Cl* is mainly through the reaction of 29 Cl with methane (46%) and other organic compounds (CH₃OH 15%, CH₃OOH 11%, C₂H₆ 8%, higher alkanes 8%,
- 30 and CH₂O 7%).
- 31 Conversion of Cl to ClO* drives some cycling of chlorine radicals, but the associated chain length versus Cl* loss is
- short $(4.1 \times 10^4/2.5 \times 10^4 = 1.6)$. Conversion of Cl to ClO is mainly by reaction with ozone (98%), while conversion of 32

1 ClO back to Cl is mostly by reaction with NO (72%), driving a null cycle as NO₂ photolyzes to regenerate NO and 2 ozone.

3 Figure 2 shows the annual mean global distributions of HCl mixing ratios and Cl atom concentrations. The mixing 4 ratio of HCl decreases from the surface to the middle troposphere, reflecting the SSA source, and then increases again 5 in the upper troposphere where it is supplied by transport from the stratosphere and has a long lifetime due to lack of 6 scavenging. Remarkably, Cl atom concentrations show little decrease with altitude, contrary to the common 7 assumption that tropospheric Cl atoms should be mainly confined to the MBL where the SSA source resides (Singh 8 et al., 1996). We find that the effect of the SSA source is offset by the slower sink of Cl* at higher altitudes due to the 9 strong temperature dependence of the reactions between Cl atom and organic compounds. Transport of HCl and Cl* 10 from the stratosphere also contribute to the source of Cl atoms in the upper troposphere.

HCl mixing ratios in marine surface air are usually highest along polluted coastlines where the large sources of HNO₃ and H₂SO₄ from anthropogenic NO_x and SO₂ emissions drive acid displacement from SSA. By contrast, HCl mixing ratios over the Southern Ocean are low because of the low supply of acid gases. The distribution of Cl atoms in surface air reflects its sources from both HCl + OH and the heterogeneous production of Cl*. The highest concentrations are in northern Europe due to production of ClNO₂ from the N₂O₅ + Cl⁻ reaction (R3). Cl atom concentrations in marine air are shifted poleward relative to HCl because of increasing bromine radical concentrations (Parrella et al., 2012), driving BrCl formation by the HOBr + Cl⁻ reaction (R5).

18 Figure 3 shows the global mean vertical distributions of reactive chlorine species (Cl*) in continental and marine air. 19 Mean boundary layer mixing ratios are higher over land than over the ocean because of the $CINO_2$ source from N_2O_5 20 +Cl⁻ in high-NO_x polluted air (Thornton et al., 2010). ClNO₂ mixing ratios are much higher than in the Sherwen et al. 21 (2016b) model which restricted its production to SSA, reflecting the importance of HCl dissolved in SNA aerosol 22 which allows further transport inland. High mixing ratios of ClNO₃ in the upper troposphere are due to transport from 23 the stratosphere and inefficacy of the sinks from hydrolysis and heterogeneous chemistry. In the marine MBL we find 24 comparable contributions from HOCl (mainly in daytime) and Cl₂ and ClNO₂ (mainly at night). The BrCl mixing ratio 25 is much lower than in the previous model studies of Long et al. (2014) and Sherwen et al. (2016b) which had very 26 large sources from the HOBr + Cl⁻ reaction (R5). Our lower BrCl mixing ratio is due to competition from the HOBr + 27 S(IV) reaction (Chen et al., 2017) and to oceanic VOC emissions (Millet et al., 2010), both of which act to depress 28 bromine radical concentrations in the MBL (Zhu et al., 2018). Further discussion of the BrCl source is presented in 29 Section 5.2.

30 4 Comparison to observations

31 Here we compare the model simulation for 2016 to observations for gas-phase chlorine and related species collected

32 in different years, assuming interannual variability to be a minor factor in model error. Previous evaluation of the

33 GEOS-Chem sea salt source by Jaeglé et al. (2011) showed general skill in simulating SSA observations and we do

not repeat this evaluation here. We also do not consider data affected by local anthropogenic sources because they would not be properly resolved at the $4^{\circ} \times 5^{\circ}$ grid resolution of our model.

3 4.1 Surface air observations

4 Table 3 compares our simulated Cl⁻ SSA deficits to an ensemble of marine air observations compiled by Graedel and 5 Keene (1995). The Cl⁻ deficit is relative to seawater composition and provides an indicator of the mobilization of Cl⁻ 6 through acid displacement and heterogeneous chemistry. The observations show a wide range from -50% to +90%, 7 and Graedel and Keene (1995) emphasize that uncertainties are large. Slight negative deficits in the observations could 8 be caused by titration of alkalinity by HCl but large negative deficits are likely due to error. Mean model deficits 9 sampled for the regions and months of the observations range from +4% to +40%, not inconsistent with the 10 observations. The largest model deficits are in polluted coastal regions because of acid displacement and this is also 11 where the measured deficits are largest.

12 Figure 4 compares simulated HCl and HNO₃ mixing ratios to concurrent observations of both gases at coastal sites 13 and over oceans. The data are arranged from left to right by increasing latitude. Mean HCl mixing ratios average 323 ppt in the model and 347 ppt in the observations for the ensemble of regions. The HCl source in the model is mainly 14 15 acid displacement from SSA. A sensitivity simulation without acid displacement from SSA has less than 7 ppt HCl 16 in all regions. The model captures the spatial variability of the mean HCl mixing ratios across locations (r = 0.88), 17 which largely reflects the HCl enhancement at polluted coastal sites and northern mid-latitudes (Figure 2). Simulated 18 HNO₃ mixing ratios average 190 ppt across locations as compared to 137 ppt in the observations, again with good 19 simulation of spatial variability (r = 0.96) driven by NO_x emissions. HNO₃ mixing ratios are sensitive to acid 20 displacement from SSA, as the sensitivity simulation without acid displacement shows mean values of 441 ppt that 21 are much higher than observed. This could partly explain the general model problem of HNO₃ overestimation in 22 remote air (Bey et al., 2001).

23 Figure 5 shows 2016 annual mean observations of PM_{2.5} Cl⁻ (mass concentration in particles less than 2.5 µm diameter) 24 from the US Interagency Monitoring of Protected Visual Environments (IMPROVE) network (Malm et al., 1994). 25 Corresponding model values are shown as background contours for fine Cl⁻ (< 1µm diameter and internally mixed 26 with SNA aerosol) are for total Cl⁻. One would expect the IMPROVE concentrations to be higher than the model fine 27 Cl⁻ (because of the larger size cut) and lower than total Cl⁻, and this is generally the case. The model is consistent with 28 observations in the continental interior, which we attribute to inland transport of marine HCl incorporated into SNA 29 aerosol. Fine Cl⁻ concentrations can actually be higher over the continent than over the ocean because of HCl 30 displacement from the coarse SSA followed by re-condensation on anthropogenic SNA aerosol. The model 31 underestimates the observations over the Southwest US and this may be due to a missing dust source. We find that 32 IMPROVE Cl⁻ and dust concentrations are positively correlated (R = 0.3-0.9) in this region.

A number of surface air measurements have been made of Cl^* as the water-soluble component of Cl_y after removal of HCl (Keene et al., 1990), although most of these measurements are below the detection limit (Table 4, mostly from 1 Keene et al. (2009)). This Cl* has been commonly assumed to represent the sum of Cl_2 and HOCl (Pszenny et al., 2 1993) but it would also include $ClNO_2$, $ClNO_3$, and minor components of Cl*. Table 4 shows that simulated Cl* mixing 3 ratios are consistent with the measurements to the extent that comparison is possible. Simulated Cl* over remote 4 oceans is dominated by HOCl, but $ClNO_2$ is responsible for the high values over the Atlantic Ocean near Europe.

5 Lawler et al. (2009) measured Cl₂ and BrCl mixing ratios at Cape Verde in the tropical Atlantic for 5 days in May-6 June 2007, and Lawler et al. (2011) measured Cl₂ and HOCl mixing ratios at the same site for 7 days in May-June 7 2009. The observations show a diurnal cycle with mixing ratios of Cl_2 highest at night, and HOCl highest in the day, 8 consistent with the model. Observed mixing ratios in background marine air are in the range 0-30 ppt for Cl_2 and 0-2 9 ppt for BrCl at night, and 0-5 ppt for HOCl in the daytime. Corresponding mean model values are 0.3 ppt for Cl₂, 1.8 10 ppt for BrCl, and 5 ppt for HOCl, with little day-to-day variability. Lawler et al. (2011) also sampled long-range 11 outflow from Europe for 3 days in 2009 with daytime HOCl and nighttime Cl₂ mixing ratio ranges of 40-200 ppt and 12 5-40 ppt respectively but the model does not capture these enhancements. Sommariva and von Glasow (2012) 13 suggested that a lower aerosol pH and/or slower rate for HOCl + Cl⁻ could explain the high HOCl in European outflow 14 but this would also cause Cl_2 to be lower. We have no explanation for the high Cl_2 values observed by Lawler et al. 15 (2009, 2011) in marine air or for the joint observed enhancements of HOCl and Cl₂ in European outflow.

16 Many surface observations of ClNO₂ have been made in nighttime urban environments. These are difficult to compare 17 to the model because of the $4^{\circ} \times 5^{\circ}$ grid resolution and because of nighttime stratification of the surface layer (the 18 lowest model grid level extends up to 130 m above the surface). In addition, the publications usually report maxima 19 instead of means. Table 5 shows a comparison for representative sites, indicating that the model offers a credible 20 simulation within the above caveats. The previous GEOS-Chem simulation of Sherwen et al. (2016b) only considered 21 $CINO_2$ production in SSA and as a result their $CINO_2$ mixing ratios were consistently below a few ppt at continental 22 sites. Our simulation can reproduce the observed >100 ppt concentrations at these sites because it accounts for HCl 23 dissolved in SNA aerosol, allowing marine influence to extend further inland as also shown in the comparison to the

24 IMPROVE Cl⁻ data (Figure 5).

25 **4.2** Comparison to aircraft measurements

26 The WINTER aircraft campaign over the eastern US and offshore in February-March 2015 provides a unique data set 27 for evaluating our model. Measurements included HCl, ClNO₂, HOCl, Cl₂, and ClNO₃ by Iodide, Time of Flight, 28 Chemical Ionization Mass Spectrometry (I-TOF-CIMS) (Lee et al., 2018). We focus on the first four measurements 29 because calibration for ClNO₃ needs further examination. The mean 1-s detection limits for HCl, ClNO₂, HOCl, and 30 Cl_2 were 100, 2, 2, and 1 ppt respectively (Lee et al., 2018). The estimated calibration uncertainty is $\pm 30\%$ for all 31 chlorine species. As discussed in Lee et al. (2018), labeled 15-N₂O₅ was added to the inlet tip during WINTER flights 32 to quantify inlet production of ClNO₂, which was found to be negligible (<<10% of measured ClNO₂), but inlet 33 production of Cl₂, for example from surface reactions of HOCl with adsorbed HCl, was not evaluated.

Figure 6 compares the observed median vertical profiles of HCl, ClNO₂, HOCl, and Cl₂ in WINTER to the model sampled along the flight tracks for the corresponding period. Figure 7 compares the median diurnal variations below 1 km altitude, separately over ocean and land. We exclude daytime (10:00-16:00 local) data for ClNO₂ in Figure 6 because its mixing ratios are near-zero (Figure 7).

5 The WINTER observations of HCl show median values of 380 ppt near the surface, dropping to a background of 100-6 200 ppt in the free troposphere (Figure 6). The model is lower than the observations in the lowest 2 km but within the 7 calibration uncertainty. The free tropospheric background in the model is much lower than observed but the 8 observations are near the 100 ppt detection limit. HCl mixing ratios in the lowest km average 60% higher over ocean 9 than over land in both the observations and the model, reflecting the marine source.

10 Also shown in Figure 6 is a sensitivity simulation including anthropogenic HCl emissions from McCulloch et al. (1999) 11 as described in Section 2.2. The resulting model mixing ratios are too high though still within the calibration 12 uncertainty. Based on sampling of power plant plumes during WINTER campaign, Lee et al. (2018) inferred a 13 HCl:SO₂ emission mass ratio of 0.033 from power plants. Adding this emission to the standard simulation, scaled to 14 the SO₂ emissions in GEOS-Chem (from the EPA National Emission Inventory over the US) increases modeled 15 mixing ratios of HCl in the continental boundary layer by 18% along the WINTER flight tracks, improving agreement 16 with observations relative to the standard simulation but still representing a relatively minor source. ClNO₂, HOCl, 17 and Cl₂ mixing ratios increase by12%, 8%, and 4% respectively.

18 Figure 8 compares the model HCl vertical profiles to measurements by the Georgia Tech CIMS instrument during the 19 SEAC⁴RS campaign over the Southeast US in August-September 2013 (Toon et al., 2016), the KORUS-AQ campaign 20 over and around the Korea peninsula in May-June 2015, and the INTEX-B campaign over the North Pacific in May 21 2006 (Kim et al., 2008). The standard model simulation without anthropogenic chlorine successfully simulates the 22 boundary layer HCl observations during SEAC⁴RS and KORUS-AQ, but adding the McCulloch et al. (1999) 23 anthropogenic inventory results in large overestimates. Boundary layer HCl mixing ratios over land are much lower 24 in SEAC⁴RS than in WINTER and this is well reproduced by the model, where the difference is due to seasonal 25 contrast in the SSA source and in the inflow of marine air. The free tropospheric background observed in SEAC⁴RS, 26 KORUS-AQ, and INTEX-B data is only ~25 ppt, much lower than in WINTER (100-200 ppt), whereas the model 27 free tropospheric background is consistently 20-50 ppt in all four campaigns. The WINTER observations are near 28 their 100 ppt detection limit as pointed out above. Modeled vertical profile of HCl during INTEX-B is very similar to 29 that of KORUS-AQ, but overestimates observed boundary layer HCl by a factor of 5. The median value of observed 30 surface HCl mixing ratio is 24 ppt during INTEX-B, which is much lower than all the surface air observations in 31 Section 4.1.

Mixing ratios of $CINO_2$ observed in WINTER are above the detection limit only in the lowest km of atmosphere at night, and are much higher over the ocean than over land. This is well simulated by the model (Figures 6 and 7), and

reflects the nighttime source from the $N_2O_5 + Cl^2$ heterogeneous reaction from the combined with the fast loss by

35 photolysis in daytime. Previous studies have suggested that the Bertram and Thornton (2009) representation of the

- 2 et al., 2013; McDuffie et al., 2018a). By using a box model applied to the WINTER observations, McDuffie et al.
- 3 (2018a;b) found that both the N_2O_5 uptake rate and the ClNO₂ production yield were overestimated by the Bertram
- 4 and Thornton (2009) parameterization. One important difference with these previous studies is the assumption of
- 5 aerosol mixing state. GEOS-Chem assumes that Cl⁻ is present only in SNA and SSA when doing the calculation of
- 6 N₂O₅ reactive uptake rates, assuming an external mixture of aerosol types (Martin et al., 2003; Evans and Jacob, 2005;
- 7 Jaeglé et al., 2018). This decreases both the N₂O₅ uptake rate and the ClNO₂ yield as compared to using the
- 8 parameterization of Bertram and Thornton (2009) on internally mixed aerosols. Jaeglé et al. (2018) reported that using
- 9 this external mixing assumption leads to good agreement between modeled and observed N_2O_5 during WINTER
- 10 campaign. The mean ClNO₂ yield (φ in R3) below 1 km in our model is 0.20 during the WINTER campaign, very
- 11 close to that calculated from observation (0.218) by McDuffie et al., (2018a).
- 12
- Nighttime Cl_2 mixing ratios in WINTER are greatly overestimated by the model. Under polluted wintertime conditions such as in WINTER the $ClNO_2 + Cl^-$ reaction greatly enhances Cl_2 production in the model:

15
$$\text{ClNO}_2 + \text{Cl}^- \rightarrow \text{NO}_2^- + \text{Cl}_2$$
 (R7)

The reactive uptake coefficient for (R7) in Table 2 is based on a single laboratory study (Roberts et al., 2008). It 16 17 requires an aerosol pH < 2 and this condition is generally met for our model simulation of the WINTER environment, 18 consistent with the observation-based analysis of aerosol pH by Guo et al. (2016) for the eastern US in winter. A 19 sensitivity simulation without (R7) is shown as dashed red lines in Figure 7 and can reproduce the low Cl₂ mixing 20 ratios observed over the ocean at night. The analysis of WINTER data by McDuffie et al. (2018b) finds that the 21 correlation between particle acidity and Cl_2 observations is opposite of the trend expected from (R7), though there 22 may be no trend on sufficiently acidic (i.e. pH < 2) aerosol. Further study of that reaction is needed. Similar reactions 23 such as $CINO_2 + Br^2$ (R8) and $CINO_2 + I^2$ are also very uncertain as their reaction rate coefficients are very different in different studies (Fickert et al., 1998; Frenzel et al., 1998; Schweitzer et al., 1998). 24

The model underestimates the WINTER observations of HOCl and Cl_2 in daytime, over the ocean as well as over land. These species have short lifetimes against photolysis (less than a few minutes). Direct anthropogenic emission from coal combustion has been proposed (Chang et al., 2002) but would only be observed in plumes and not over the oceans. Matching the >1 ppt Cl_2 observed during daytime is particularly problematic since it would require a large photochemical source absent from the model. Lawler et al. (2011) suggested a fast daytime HOCl source from a hypothetical light-dependent Cl⁻ oxidation. The measurements of Cl_2 are also possibly subject to positive artifact from rapid heterogeneous conversion of chlorine species on the surface of the TOF-CIMS inlet (Lee et al., 2018).

1 5 Global implications of tropospheric chlorine chemistry

2 5.1 Cl atom and its impact on VOCs

3 The global mean pressure-weighted tropospheric Cl atom concentration in our simulation is 620 cm⁻³, while the MBL 4 concentration averages 1200 cm⁻³ (Figure 2). Our global mean is lower than the previous global model studies of 5 Hossaini et al. (2016) (1300 cm⁻³) and Long et al. (2014) (3000 cm⁻³), which had excessive Cl* generation as discussed 6 above. It is consistent with the upper limit of 1000 cm⁻³ inferred by Singh et al. (1996) from global modeling of C_2Cl_4 7 observations (C_2Cl_4 is highly reactive with Cl atoms). Isotopic observations of methane have been used to infer a Cl 8 atom concentration in the MBL higher than 9000 cm⁻³ in the extra-tropical Southern Hemisphere (Platt et al., 2004; 9 Allan et al., 2007), much higher than our estimate of 800 cm⁻³ over this region. More recently, Gromov et al. (2018) 10 revisited these data together with added constraints from CO isotope measurements and concluded that extra-tropical 11 Southern Hemisphere concentrations of Cl atoms in the MBL should be lower than 900 cm⁻³, consistent with our 12 estimate.

13 Tropospheric oxidation by Cl atoms drives a present-day methane loss rate of 5.3 Tg a⁻¹ in our model, contributing

14 only 1.0% of total methane chemical loss. It has more significant impact on the oxidation of some other VOCs,

15 contributing 20% of the global loss for ethane, 14% for propane, 10% for higher alkanes, and 4% for methanol.

16 5.2 Impact on bromine and iodine chemistry

- 17 Bromine radicals (BrO_x \equiv Br + BrO) and iodine radicals (IO_x \equiv I + IO) affect global tropospheric chemistry by 18 depleting ozone and OH (Parrella et al., 2012; Sherwen et al., 2016b). Br atoms are also thought to drive the oxidation 19 of elemental mercury (Holmes et al., 2006). Chlorine chemistry increases IO_x mixing ratios by 16% due to the 20 reactions of HOI, INO₂, and INO₃ with Cl⁻(R11-R13), producing ICl which photolyzes rapidly to I atoms (Figure 1). 21 The effect on bromine is more complicated. Bromine radicals originate from photolysis and oxidation of 22 organobromines emitted by the ocean, as well as from SSA debromination (Yang et al., 2005). They are lost by 23 conversion to HBr which is efficiently deposited. Parrella et al. (2012) pointed out that heterogeneous chemistry of 24 HBr (dissolved as Br⁻) is critical for recycling bromine radicals and explaining observed tropospheric BrO mixing 25 ratios in the background troposphere:
- 26 $HOBr(aq) + Br^- + H^+ \rightarrow Br_2 + H_2O$ (R14) 27 $Br_2 + hv \rightarrow 2Br$ (R15)

Chloride ions and dissolved SO₂ (S(IV) \equiv HSO₃⁻ + SO₃²⁻) can however compete with Br⁻ for the available HOBr (Chen et al., 2017):

$30 \text{HOBr}(aq) + \text{Cl}^- + \text{H}^+ \rightarrow \text{BrCl} + \text{H}_2\text{O}$	(R5)
---	------

31 $HOBr(aq) + HSO_3^{-}/SO_3^{2-} \rightarrow HBr + HSO_4^{-}/SO_4^{2-}$ (R16)

32 Chen et al. (2017) pointed out that reaction (R16) effectively decreases BrO mixing ratios by producing HBr which is 33 rapidly deposited instead of contributing to BrO_x cycling. They found in a GEOS-Chem simulation that global 1 tropospheric BrO mixing ratios decreased by a factor of 2 as a result. Reaction (R5) may however have a compensating 2 or opposite effect. It propagates the cycling of BrO_x if BrCl volatilizes:

3 $\operatorname{BrCl} + hv \to \operatorname{Br} + \operatorname{Cl}$ (R17)

4 but it may also generate new BrO_x if BrCl reacts with Br^- in the aqueous phase to produce Br_2 (Wang et al., 1994):

5	$BrCl(aq) + Br^{-} \rightleftharpoons Br_2Cl^{-}$	(R18)
6	$Br_2Cl^- \rightleftharpoons Br_2(aq) + Cl^-$	(R19)

The sequence (R5) + (R18) + (R19) with Cl⁻ as a catalyst has the same stoichiometry as (R14) and thus contributes to HBr recycling in the same way. We find in the model that it is globally 30 times faster than (R14) and therefore much more effective at regenerating bromine radicals. In GEOS-Chem, the rate of reaction (R5) computed from Table 2 is applied to the following stoichiometry reflecting the ensemble of reactions (R5, R14, R18, and R19):

11
$$HOBr(aq) + YBr^{-} + (1 - Y)Cl^{-} + H^{+} \rightarrow YBr_{2} + (1 - Y)BrCl + H_{2}O$$
 (R5+R14+R18+R19)

where *Y* is the yield of Br_2 and 1-*Y* is the yield of BrCl. *Y* is calculated following the laboratory study of Fickert et al. (1999):

14
$$Y = 0.41 \log_{10}([Br^-]/[Cl^-]) + 2.25$$
 for $[Br^-]/[Cl^-] < 5 \times 10^{-4}$ (2)
15 $Y = 0.90$ for $[Br^-]/[Cl^-] > 5 \times 10^{-4}$ (3)

This mechanism was first included in GEOS-Chem version 11-02d by Chen et al. (2017), who did not however have an explicit SSA Cl⁻ simulation (they instead assumed a fixed SSA [Cl⁻] = 0.5 M, and considered only dissolved HCl in cloud).

19 Chen et al. (2017) found in their GEOS-Chem simulation that the global tropospheric BrO burden was 8.7 Gg without 20 the HOBr + S(IV) reaction (R16), and dropped to 3.6 Gg when that reaction was included. Previous GEOS-Chem 21 model estimates of the global tropospheric BrO burden were 3.8 Gg (Parrella et al., 2012), 5.7 Gg (Schmidt et al., 22 2016), and 6.4 Gg (Sherwen et al., 2016b). Our simulation features many updates relative to Chen et al. (2017) 23 including not only explicit SSA Cl⁻ but also explicit calculation of aerosol pH with ISORROPIA II for the rates of 24 reactions in Table 2. By including explicit SSA Cl⁻, the cloudwater [Cl⁻] in our model is much higher than that in Chen 25 et al. (2017) and more comparable to measurements ($\sim 10^{-4}$ M in typical cloud; (Straub et al., 2007)). We find in our standard simulation a global tropospheric BrO burden of 4.2 Gg, 17% higher than Chen et al. (2017). 26

Figure 9 shows the change of surface BrO mixing ratios due specifically to tropospheric chlorine chemistry, as obtained by difference with a sensitivity simulation including none of the Cl_y chemistry shown in Figure 1. The inclusion of chlorine chemistry increases the global tropospheric BrO burden by 85%. More than 80% of this change

30 is caused by the HOBr + Cl⁻ reaction as discussed above. Other significant contributions include $ClNO_3 + Br^-$ and

1 $CINO_2 + Br^{-}$. The largest BrO increases (1-2 ppt) are in surface air over the high northern latitudes oceans where SSA 2 emissions are high and acidic conditions promote HOBr + Cl⁻ chemistry.

3 5.3 Impact on tropospheric ozone and OH

Figure 9 also shows the effects of chlorine chemistry on NO_x, OH, and ozone concentrations. The global tropospheric burdens decrease by 5% for NO_x, 3% for OH, and 7% for ozone. The inter-hemispheric (N/S) ratio of tropospheric mean OH decreases from 1.14 to 1.12. Models tend to overestimate global mean tropospheric OH and its inter-hemispheric ratio relative to the constraint from methylchloroform which suggests a ratio of 0.85-0.98 (Naik et al., 2013;Voulgarakis et al., 2013). The effect of chlorine chemistry on the N/S ratio is slight but in the right direction.

9 The chlorine-induced decreases in Figure 9 are mainly through bromine chemistry initiated by chlorine (Section 5.2),

and have spatial distributions characteristic of bromine chemistry with maxima at high latitudes as discussed by

11 Schmidt et al. (2016). There are specific chlorine mechanisms including catalytic ozone loss through HOCI formation

12 and photolysis:

4

5 6

7

8

$13 \qquad \text{Cl} + 0_3 \rightarrow \text{Cl}0 + 0_2 \tag{F}$
--

14	$ClO + HO_2 \rightarrow HOCl + O_2$	(R21)
----	-------------------------------------	-------

- 15 $HOCI \xrightarrow{hv} CI + OH$ (R22)
- 16 Net: $O_3 + HO_2 \rightarrow OH + 2O_2$
- 17 and also loss of NO_x :
- 18 $\text{ClO} + \text{NO}_2 + \text{M} \rightarrow \text{ClNO}_3 + \text{M}$ (R23)
- 19 $\text{ClNO}_3 + \text{H}_2\text{O} \rightarrow \text{HOCl} + \text{HNO}_3$ (R24)

However, we find that the rates are very small compared to similar mechanisms involving bromine and iodine because
the stability of HCl quenches Cl* radical cycling.

22

23 A particular situation arises over polluted continents due to ClNO₂ chemistry. Production of ClNO₂ at night from the 24 $N_2O_5 + Cl^-$ heterogeneous reaction, followed by photolysis in the morning to release Cl and NO₂, provides a source of 25 radicals and ozone. This explains the increases of OH over North America and Europe in Figure 9. The effect is most 26 important at high northern latitudes in winter due to the longer night. To isolate the impact on ozone we conducted a 27 sensitivity simulation with no ClNO₂ production, setting $\varphi = 0$ for reaction (R3) in Table 2. The surface air ozone 28 enhancement due to $CINO_2$ chemistry is found to be the largest (~8 ppb) in European winter, due to the large supply 29 of Cl⁻ from the North Atlantic combined with high NO_x emissions. Other polluted continents see ozone increases of 30 1-5 ppb in winter. The effect in summer is less than 1 ppb. These results are similar to previous regional modeling 31 studies by Sarwar et al. (2014) and Sherwen et al. (2017).

2 GEOS-Chem model version 11-02d including SSA debromination. Our explicit treatment of chlorine chemistry and 3 thermodynamic representation of aerosol pH increases the global tropospheric BrO burden by 40%. Most of this 4 change is caused by faster HOBr + Cl⁻ reaction at high latitudes, resulting from higher Cl- concentration in our model 5 particularly in cloud. The decrease of BrO in the tropical MBL is caused by an increase in aerosol pH (pH was 6 previously assumed to be 0 for computation of bromine chemistry), which slows down the acid-catalyzed recycling 7 of bromine by reactions (R5) and (R14). Our computed global tropospheric burdens decrease by 4% for NO_x, 2% for 8 OH, and 4% for ozone relative to version 11-02d, again due to the more active bromine chemistry. The increase of 9 OH over continental regions is due to our accounting of HCl dissolved in SNA aerosol, allowing marine influence to 10 extend further inland to drive ClNO₂ chemistry.

Figure S1 shows the differences of BrO, NO_x, OH, and ozone concentrations between our model and the standard

11 6 Conclusions

1

We have added to the GEOS-Chem model a comprehensive and consistent representation of tropospheric chlorine chemistry. This includes in particular explicit accounting of the mobilization of sea salt aerosol (SSA) chloride (Cl⁻), by acid displacement of HCl as well as by other heterogeneous processes. Cycling of inorganic gas-phase chlorine species (Cl_y) generated from SSA and other sources is simulated and coupled to the model aerosol-oxidant-bromineiodine chemistry. With our work, GEOS-Chem now has a complete simulation of halogen (Cl+Br+I) chemistry in both the troposphere and stratosphere.

18 Emission of chlorine in the model is mainly as sea-salt aerosol (1780 Tg Cl a⁻¹). Other sources (combustion, 19 organochlorines, stratospheric input) are also included but are small in comparison. Most of the sea-salt aerosol 20 chloride is removed by deposition, but 3.6% is mobilized to inorganic gas-phase chlorine (Cl_v) through acid 21 displacement to HCl (52 Tg a⁻¹) and through other heterogeneous chemistry producing more reactive chlorine species 22 (12 Tg a^{-1}) . We define reactive chlorine (Cl*) as the ensemble of Cl_y species excluding HCl and including Cl, ClO, Cl₂, BrCl, HOCl, ClNO₂, ClNO₃, plus other minor species. Oxidation of HCl by OH provides a Cl* source of 9.7 Tg 23 24 a^{-1} , comparable to the heterogeneous source from HOBr + Cl⁻ (8.6 Tg a^{-1}). N₂O₅ + Cl⁻ (1.8 Tg a^{-1}) is also important 25 in polluted environments. Cycling between Cl* species drives radical chlorine (Cl/ClO) chemistry but chain lengths 26 are limited by fast conversion to HCl and subsequent deposition.

HCl mixing ratios in the model are highest over the oceans downwind of polluted continents due to effective acid displacement from sea-salt aerosol by HNO₃ and H₂SO₄. Mixing ratios are much lower over the Southern Ocean where the supply of acids is low. The dominant daytime Cl* species is generally HOCl while BrCl, Cl₂, and ClNO₂ dominate at night. ClNO₃ dominates in the upper troposphere due to stratospheric input. Chlorine atom concentrations are highest over Europe in winter due to ClNO₂ chemistry, and are otherwise high over the northern mid-latitudes oceans where the supply of acidity promotes Cl formation both through HCl and through acid-catalyzed heterogeneous processes.

1 Comparison of model results to observations in marine surface air show that the model is usually able to reproduce

2 the range and distributions of observed sea-salt aerosol chloride deficits, HCl mixing ratios, and Cl* mixing ratios. In

- 3 particular, concurrent observations of HCl and HNO₃ in coastal/marine air worldwide show high correlation with the
- 4 model including high HCl mixing ratios at northern mid-latitudes combined with depressed HNO₃. Consideration of
- 5 acid displacement greatly improves model agreement with HNO₃ observations in marine air. The model can also
- 6 successfully simulate observations of high ClNO₂ at night including in continental air. The chlorine in that case
- 7 originates from sea-salt aerosol transported far inland following uptake of volatilized HCl by sulfate-nitrate-
- 8 ammonium (SNA) aerosol. The model cannot reproduce the very high HOCl and Cl₂ concentrations observed by
- 9 Lawler et al. (2009;2011) at Cape Verde in the tropical Atlantic.
- 10 Comparisons of model results to aircraft campaign observations from WINTER (eastern US and offshore, Feb-Mar

11 2015), SEAC4RS (Southeast US, Aug-Sep 2013), and KORUS-AQ (Korean Peninsula, April-Jun 2016) show general

12 consistency for HCl vertical profiles. Continental boundary layer HCl mixing ratios in these campaigns can be mostly

13 accounted for by the marine source transported inland, though power plants could make a minor contribution.

14 WINTER observations also include ClNO₂ and Cl₂, and HOCl. The observed ClNO₂ is mainly confined to the

15 nighttime marine boundary layer and is consistent with the model. Observed Cl₂ concentrations at night are much

16 lower than the model, which has a large source under the WINTER conditions from the $CINO_2 + CI^2$ heterogeneous

17 reaction. The rate coefficient for this reaction is from only one laboratory study.

18 The model simulates a global mean Cl atom concentration of 620 cm⁻³ in the troposphere and 1200 cm⁻³ in the marine 19 boundary layer (MBL), lower than previous global model studies that had excessive generation of Cl* but consistent 20 with independent proxy constraints. We find that oxidation by Cl atoms accounts for only 1.0% of the global loss of 21 atmospheric methane but has larger effects on the global losses of ethane (20%), propane (14%), and methanol (4%). 22 Chlorine chemistry increases global tropospheric BrO by 85%, and decreases ozone and OH by 7% and 3% 23 respectively, relative to a sensitivity simulation with no chlorine chemistry. The large effect on BrO is due to 24 production of bromine radicals by the HOBr + Cl^{-} heterogeneous reaction, and the decreases of ozone and OH are 25 mainly through the induced bromine chemistry. An exception is winter conditions over polluted regions, where ClNO₂ 26 chemistry increases ozone mixing ratios by up to 8 ppb.

Author contributions. XW, DJJ, and HL designed the study. XW developed the chlorine model code, performed the simulations and analyses. SDE, MPS, LZ, QC, BA, TS, and MJE contributed to the GEOS-Chem halogen model development. BHL, JDH, FDL, and JAT conducted and processed the measurement during WINTER campaign. GLH conducted and processed the measurement during SEAC⁴RS and KORUS-AQ campaigns. XW and DJJ prepared the manuscript with contributions from all co-authors.

Data availability. The model code is available from the corresponding author upon request, and will be made available to the community through the standard GEOS-Chem (http://www.geos-chem.org) in the future. Data of WINTER campaign are available to the general public at https://www.eol.ucar.edu/field_projects/winter. Data of NASA SEAC⁴RS AND KORUS-AQ missions and are available to the general public through the NASA data archive

- 1 (https://www-air.larc.nasa.gov/cgi-bin/ArcView/seac4rs and https://www-air.larc.nasa.gov/cgi-
- 2 bin/ArcView/korusaq). IMPROVE data are available through the Federal Land Manager Environmental database
- 3 (http://views.cira.colostate.edu/fed).

4 Acknowledgments. This work was supported by the Atmospheric Chemistry Program of the US National Science 5 Foundation and by the Joint Laboratory for Air Quality and Climate (JLAQC) between Harvard and the Nanjing 6 University for Information Science and Technology (NUIST). QC and BA were supported by National Science 7 Foundation (AGS 1343077). We thank Prasad S. Kasibhatla for insightful discussion. WINTER data are provided by 8 NCAR/EOL under sponsorship of the National Science Foundation (https://www.eol.ucar.edu/field projects/winter). 9 SEAC⁴RS, KORUS-AQ, and INTEX-B data are provided by NASA LaRC Airborne Science Data for Atmospheric 10 Composition (https://www-air.larc.nasa.gov). IMPROVE is a collaborative association of state, tribal, and federal 11 agencies, and international partners. US Environmental Protection Agency is the primary funding source, with contracting and research support from the National Park Service. The Air Quality Group at the University of 12 13 California, Davis is the central analytical laboratory, with ion analysis provided by Research Triangle Institute, and 14 carbon analysis provided by Desert Research Institute.

1 References

Abbatt, J. P. D., Lee, A. K. Y., and Thornton, J. A.: Quantifying trace gas uptake to tropospheric aerosol: recent advances and remaining challenges, Chemical Society Reviews, 41, 6555-6581, 10.1039/C2CS35052A, 2012.

Alexander, B., Park, R. J., Jacob, D. J., Li, Q. B., Yantosca, R. M., Savarino, J., Lee, C. C. W., and Thiemens, M. H.:
Sulfate formation in sea-salt aerosols: Constraints from oxygen isotopes, Journal of Geophysical Research:
Atmospheres, 110, doi:10.1029/2004JD005659, 2005.

- Alexander, B., Allman, D. J., Amos, H. M., Fairlie, T. D., Dachs, J., Hegg, D. A., and Sletten, R. S.: Isotopic
 constraints on the formation pathways of sulfate aerosol in the marine boundary layer of the subtropical northeast
 Atlantic Ocean, Journal of Geophysical Research: Atmospheres, 117, doi:10.1029/2011JD016773, 2012.
- Attantic Ocean, Journal of Geophysical Research. Atmospheres, 117, doi:10.1029/2011j1
- Allan, W., Struthers, H., and C. Lowe, D.: Methane carbon isotope effects caused by atomic chlorine in the marine boundary layer: Global model results compared with Southern Hemisphere measurements, 2007.
- 12 Ammann, M., Cox, R. A., Crowley, J. N., Jenkin, M. E., Mellouki, A., Rossi, M. J., Troe, J., and Wallington, T. J.:
- 13 Evaluated kinetic and photochemical data for atmospheric chemistry: Volume VI heterogeneous reactions with
- 14 liquid substrates, Atmospheric Chemistry and Physics, 13, 8045-8228, 10.5194/acp-13-8045-2013, 2013.
- 15 Amos, H. M., Jacob, D. J., Holmes, C. D., Fisher, J. A., Wang, Q., Yantosca, R. M., Corbitt, E. S., Galarneau, E.,
- 16 Rutter, A. P., Gustin, M. S., Steffen, A., Schauer, J. J., Graydon, J. A., Louis, V. L. S., Talbot, R. W., Edgerton, E. S.,
- 17 Zhang, Y., and Sunderland, E. M.: Gas-particle partitioning of atmospheric Hg(II) and its effect on global mercury
- 18 deposition, Atmos. Chem. Phys., 12, 591-603, 10.5194/acp-12-591-2012, 2012.
- Atkinson, R.: Gas-Phase Tropospheric Chemistry of Volatile Organic Compounds: 1. Alkanes and Alkenes, J. Phys.
 Chems. Ref. Data, 26(2), 215-290, 1997.
- 21 Bannan, T. J., Booth, A. M., Bacak, A., Muller, J. B. A., Leather, K. E., Le Breton, M., Jones, B., Young, D., Coe, H.,
- Allan, J., Visser, S., Slowik, J. G., Furger, M., Prévôt, A. S. H., Lee, J., Dunmore, R. E., Hopkins, J. R., Hamilton, J.
 F., Lewis, A. C., Whalley, L. K., Sharp, T., Stone, D., Heard, D. E., Fleming, Z. L., Leigh, R., Shallcross, D. E., and
- Percival, C. J.: The first UK measurements of nitryl chloride using a chemical ionization mass spectrometer in central
- London in the summer of 2012, and an investigation of the role of Cl atom oxidation, Journal of Geophysical Research:
- 26 Atmospheres, 120, 5638-5657, doi:10.1002/2014JD022629, 2015.
- 27 Bannan, T. J., Bacak, A., Le Breton, M., Flynn, M., Ouyang, B., McLeod, M., Jones, R., Malkin, T. L., Whalley, L.
- 28 K., Heard, D. E., Bandy, B., Khan, M. A. H., Shallcross, D. E., and Percival, C. J.: Ground and Airborne U.K.
- 29 Measurements of Nitryl Chloride: An Investigation of the Role of Cl Atom Oxidation at Weybourne Atmospheric
- 30 Observatory, Journal of Geophysical Research: Atmospheres, 122, 11,154-111,165, doi:10.1002/2017JD026624,
- 31 2017.
- Bari, A., Ferraro, V., Wilson, L. R., Luttinger, D., and Husain, L.: Measurements of gaseous HONO, HNO3, SO2,
 HCl, NH3, particulate sulfate and PM2.5 in New York, NY, Atmospheric Environment, 37, 2825-2835,
 10.1016/s1352-2310(03)00199-7, 2003.
- Behnke, W., George, C., Scheer, V., and Zetzsch, C.: Production and decay of CINO2 from the reaction of gaseous
 N2O5 with NaCl solution: Bulk and aerosol experiments, Journal of Geophysical Research: Atmospheres, 102, 3795 3804, doi:10.1029/96JD03057, 1997.
- Bertram, T. H., and Thornton, J. A.: Toward a general parameterization of N2O5 reactivity on aqueous particles: the
 competing effects of particle liquid water, nitrate and chloride, Atmos. Chem. Phys., 9, 8351-8363, 10.5194/acp-9 8351-2009, 2009.
 - 19

- 1 Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q., Liu, H. Y., Mickley, L. J., and
- 2 Schultz, M. G.: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and 3
- evaluation, Journal of Geophysical Research: Atmospheres, 106, 23073-23095, doi:10.1029/2001JD000807, 2001.

4 Chang, S., McDonald-Buller, E., Kimura, Y., Yarwood, G., Neece, J., Russell, M., Tanaka, P., and Allen, D.: 5 Sensitivity of urban ozone formation to chlorine emission estimates, Atmospheric Environment, 36, 4991-5003, 6 https://doi.org/10.1016/S1352-2310(02)00573-3, 2002.

7 Chen, Q., Schmidt, J. A., Shah, V., Jaeglé, L., Sherwen, T., and Alexander, B.: Sulfate production by reactive bromine: 8 Implications for the global sulfur and reactive bromine budgets, Geophysical Research Letters, 44, 7069-7078, 9 doi:10.1002/2017GL073812, 2017.

10 Crisp, T. A., Lerner, B. M., Williams, E. J., Quinn, P. K., Bates, T. S., and Bertram, T. H.: Observations of gas phase hydrochloric acid in the polluted marine boundary layer, Journal of Geophysical Research: Atmospheres, 119, 6897-11 6915, doi:10.1002/2013JD020992, 2014. 12

13 Dasgupta, P. K., Campbell, S. W., Al-Horr, R. S., Ullah, S. M. R., Li, J., Amalfitano, C., and Poor, N. D.: Conversion 14 of sea salt aerosol to NaNO3 and the production of HCl: Analysis of temporal behavior of aerosol chloride/nitrate and 15 gaseous HCl/HNO3 concentrations with Atmospheric Environment, 41, 4242-4257, AIM, 16 10.1016/j.atmosenv.2006.09.054, 2007.

- 17 Eastham, S. D., Weisenstein, D. K., and Barrett, S. R. H.: Development and evaluation of the unified tropospheric-
- 18 stratospheric chemistry extension (UCX) for the global chemistry-transport model GEOS-Chem, Atmospheric
- 19 Environment, 89, 52-63, https://doi.org/10.1016/j.atmosenv.2014.02.001, 2014.
- 20 Evans, M. J., and Jacob, D. J.: Impact of new laboratory studies of N2O5 hydrolysis on global model budgets of 21 tropospheric nitrogen oxides, ozone, and OH, Geophysical Research Letters, 32, doi:10.1029/2005GL022469, 2005.
- 22 Faxon, C., Bean, J., and Ruiz, L.: Inland Concentrations of Cl2 and ClNO2 in Southeast Texas Suggest Chlorine 23 Chemistry Significantly Contributes to Atmospheric Reactivity, Atmosphere, 6, 1487-1506, 10.3390/atmos6101487, 24 2015.
- 25 Fickert, S., Helleis, F., Adams, J. W., Moortgat, G. K., and Crowley, J. N.: Reactive uptake of CINO2 on aqueous 26 bromide solutions. Journal of Physical Chemistry A, 102(52), 10689-10696. https://doi.org/10.1021/jp983004n, 1998.
- 27 Fickert, S., Adams, J. W., and Crowley, J. N.: Activation of Br2 and BrCl via uptake of HOBr onto aqueous salt 28 solutions, Journal of Geophysical Research: Atmospheres, 104, 23719-23727, doi:10.1029/1999JD900359, 1999.
- 29 Finlayson-Pitts, B. J.: The Tropospheric Chemistry of Sea Salt: A Molecular-Level View of the Chemistry of NaCl 30 and NaBr, Chemical Reviews, 103, 4801-4822, 10.1021/cr020653t, 2003.
- 31 Fountoukis, C., and Nenes, A.: ISORROPIA II: a computationally efficient thermodynamic equilibrium model for 32 K⁺-Ca²⁺-Mg²⁺-NH₄⁺-Na⁺-SO₄²⁻-NO₃⁻-Cl⁻-H₂O aerosols,, Atmos. Chem. Phys., 7, 4639-4659, 10.5194/acp-7-33 4639-2007, 2007.
- 34 Frenzel, A., Scheer, V., Sikorski, R., George, C., Behnke, W., & Zetzsch, C.: Heterogeneous Interconversion Reactions of BrNO₂, ClNO₂, Br₂, and Cl₂. The Journal of Physical Chemistry A, 102(8), 1329-1337. 35 https://doi.org/10.1021/jp973044b, 1998. 36
- 37 Fridlind, A. M., and Jacobson, M. Z.: A study of gas-aerosol equilibrium and aerosol pH in the remote marine
- boundary layer during the First Aerosol Characterization Experiment (ACE 1), Journal of Geophysical Research: 38
- 39 Atmospheres, 105, 17325-17340, doi:10.1029/2000JD900209, 2000.

- 1 Giglio, L., Randerson, J. T., and van der Werf, G. R.: Analysis of daily, monthly, and annual burned area using the
- 2 fourth-generation global fire emissions database (GFED4), Journal of Geophysical Research: Biogeosciences, 118,
- 3 317-328, doi:10.1002/jgrg.20042, 2013.
- Graedel, T. E., and Keene, W. C.: Tropospheric budget of reactive chlorine, Global Biogeochemical Cycles, 9, 47-77,
 doi:10.1029/94GB03103, 1995.
- 6 Gromov, S., Brenninkmeijer, C. A. M., and Jöckel, P.: A very limited role of tropospheric chlorine as a sink of the 7 greenhouse gas methane, Atmospheric Chemistry and Physics, 18, 9831-9843, 10.5194/acp-18-9831-2018, 2018.
- Gurciullo, C., Lerner, B., Sievering, H., and Pandis, S. N.: Heterogeneous sulfate production in the remote marine
 environment: Cloud processing and sea-salt particle contributions, J. Geophys. Res., 104(D17), 21,719 21,731, 1999.
- 10 Guo, H., Sullivan, A. P., Campuzano-Jost, P., Schroder, J. C., Lopez-Hilfiker, F. D., Dibb, J. E., Jimenez, J. L.,
- 11 Thornton, J. A., Brown, S. S., Nenes, A., and Weber, R. J.: Fine particle pH and the partitioning of nitric acid during
- 12 winter in the northeastern United States, Journal of Geophysical Research: Atmospheres, 121, 10,355-310,376,
- 13 doi:10.1002/2016JD025311, 2016.
- 14 Haskins, J. D., Jaeglé, L., Shah, V., Lee, B. H., Lopez-Hilfiker, F. D., Campuzano-Jost, P., Schroder, J. C., Day, D.
- 15 A., Guo, H., Sullivan, A. P., Weber, R., Dibb, J., Campos, T., Jimenez, J. L., Brown, S. S., and Thornton, J. A.:
- 16 Wintertime Gas-Particle Partitioning and Speciation of Inorganic Chlorine in the Lower Troposphere Over the
- 17 Northeast United States and Coastal Ocean, Journal of Geophysical Research: Atmospheres, 123, 12,897-812,916,
- 18 doi:10.1029/2018JD028786, 2018.
- 19 Hoffmann, E. H., Tilgner, A., Schrödner, R., Bräuer, P., Wolke, R., and Herrmann, H.: An advanced modeling study
- 20 on the impacts and atmospheric implications of multiphase dimethyl sulfide chemistry, Proceedings of the National
- 21 Academy of Sciences, 113, 11776-11781, 10.1073/pnas.1606320113, 2016.
- Holmes, C. D., Jacob, D. J., and Yang, X.: Global lifetime of elemental mercury against oxidation by atomic bromine
 in the free troposphere, Geophysical Research Letters, 33, doi:10.1029/2006GL027176, 2006.
- 24 Horowitz, H. M., Jacob, D. J., Zhang, Y., Dibble, T. S., Slemr, F., Amos, H. M., Schmidt, J. A., Corbitt, E. S., Marais,
- 25 E. A., and Sunderland, E. M.: A new mechanism for atmospheric mercury redox chemistry: implications for the global
- 26 mercury budget, Atmos. Chem. Phys., 17, 6353-6371, 10.5194/acp-17-6353-2017, 2017.
- Hossaini, R., Chipperfield, M. P., Saiz-Lopez, A., Fernandez, R., Monks, S., Feng, W., Brauer, P., and von Glasow,
 R.: A global model of tropospheric chlorine chemistry: Organic versus inorganic sources and impact on methane
 oxidation, Journal of Geophysical Research: Atmospheres, 121, 14,271-214,297, doi:10.1002/2016JD025756, 2016.
- 30 Impey, G. A., Mihele, C. M., Anlauf, K. G., Barrie, L. A., Hastie, D. R., and Shepson, P. B.: Measurements of
- 31 Photolyzable Halogen Compounds and Bromine Radicals During the Polar Sunrise Experiment 1997, Journal of
- 32 Atmospheric Chemistry, 34, 21-37, 10.1023/a:1006264912394, 1999.
- Jacob, D. J., Waldman, J. M., Munger, J. W., and Hoffmann, M. R.: Chemical composition of fogwater collected
 along the California coast, Environmental Science & Technology, 19, 730-736, 10.1021/es00138a013, 1985.
- Jacob, D. J.: Heterogeneous chemistry and tropospheric ozone, Atmospheric Environment, 34, 2131-2159, https://doi.org/10.1016/S1352-2310(99)00462-8, 2000.
- Jaeglé, L., Quinn, P. K., Bates, T. S., Alexander, B., and Lin, J. T.: Global distribution of sea salt aerosols: new
 constraints from in situ and remote sensing observations, Atmospheric Chemistry and Physics, 11, 3137-3157,
 10.5194/acp-11-3137-2011, 2011.
- Jaeglé, L., Shah, V., Thornton, J. A., Lopez-Hilfiker, F. D., Lee, B. H., McDuffie, E. E., Fibiger, D., Brown, S. S.,
 Veres, P., Sparks, T. L., Ebben, C. J., Wooldridge, P. J., Kenagy, H. S., Cohen, R. C., Weinheimer, A. J., Campos, T.

- 1 L., Montzka, D. D., Digangi, J. P., Wolfe, G. M., Hanisco, T., Schroder, J. C., Campuzano-Jost, P., Day, D. A.,
- Jimenez, J. L., Sullivan, A. P., Guo, H., and Weber, R. J.: Nitrogen Oxides Emissions, Chemistry, Deposition, and
 Export Over the Northeast United States During the WINTER Aircraft Campaign, Journal of Geophysical Research:
- 4 Atmospheres, 123, 12,368-312,393, doi:10.1029/2018JD029133, 2018.
- 5 Jeong, D., Seco, R., Gu, D., Lee, Y., Nault, B. A., Knote, C. J., McGee, T., Sullivan, J. T., Jimenez, J. L., Campuzano-6 Jost, P., Blake, D. R., Sanchez, D., Guenther, A. B., Tanner, D., Huey, L. G., Long, R., Anderson, B. E., Hall, S. R.,
- Jost, F., Blace, D. R., Saltenez, D., Guenner, A. D., Famier, D., Huey, E. G., Eolig, R., Anderson, D. E., Han, S. R.,
 Ullmann, K., Shin, H. J., Herndon, S. C., Lee, Y., Kim, D., Ahn, J., and Kim, S.: Integration of Airborne and Ground
- 8 Observations of Nitryl Chloride in the Seoul Metropolitan Area and the Implications on Regional Oxidation Capacity
- 9 During KORUS-AQ 2016, Atmos. Chem. Phys. Discuss., 2018, 1-25, 10.5194/acp-2018-1216, 2018.
- 10 Kasibhatla, P., Sherwen, T., Evans, M. J., Carpenter, L. J., Reed, C., Alexander, B., Chen, Q., Sulprizio, M. P., Lee,
- J. D., Read, K. A., Bloss, W., Crilley, L. R., Keene, W. C., Pszenny, A. A. P., and Hodzic, A.: Global impact of nitrate
- photolysis in sea-salt aerosol on NOx, OH, and O3 in the marine boundary layer, Atmos. Chem. Phys., 18, 11185-11203, 10.5194/acp-18-11185-2018, 2018.
- 14 Keene, W. C., Pszenny, A. A. P., Jacob, D. J., Duce, R. A., Galloway, J. N., Schultz-Tokos, J. J., Sievering, H., and
- 15 Boatman, J. F.: The geochemical cycling of reactive chlorine through the marine troposphere, Global Biogeochemical
- 16 Cycles, 4, 407-430, doi:10.1029/GB004i004p00407, 1990.
- 17 Keene, W. C., Stutz, J., Pszenny, A. A. P., Maben, J. R., Fischer, E. V., Smith, A. M., von Glasow, R., Pechtl, S.,
- 18 Sive, B. C., and Varner, R. K.: Inorganic chlorine and bromine in coastal New England air during summer, Journal of
- 19 Geophysical Research: Atmospheres, 112, 10.1029/2006jd007689, 2007.
- 20 Keene, W. C., Long, M. S., Pszenny, A. A. P., Sander, R., Maben, J. R., Wall, A. J., O'Halloran, T. L., Kerkweg, A.,
- 21 Fischer, E. V., and Schrems, O.: Latitudinal variation in the multiphase chemical processing of inorganic halogens
- 22 and related species over the eastern North and South Atlantic Oceans, Atmos. Chem. Phys., 9, 7361-7385,
 - 23 10.5194/acp-9-7361-2009, 2009.
 - Kelly, J. T., Bhave, P. V., Nolte, C. G., Shankar, U., and Foley, K. M.: Simulating emission and chemical evolution
 of coarse sea-salt particles in the Community Multiscale Air Quality (CMAQ) model, Geosci. Model Dev., 3, 257 273, 10.5194/gmd-3-257-2010, 2010.
- 27 Kercher, J. P., Riedel, T. P., and Thornton, J. A.: Chlorine activation by N2O5: simultaneous, in situ detection of
- CINO2 and N2O5 by chemical ionization mass spectrometry, Atmos. Meas. Tech., 2, 193-204, 10.5194/amt-2-193 2009, 2009.
- 30 Kim, S., Huey, L. G., Stickel, R. E., Pierce, R. B., Chen, G., Avery, M. A., Dibb, J. E., Diskin, G. S., Sachse, G. W.,
- McNaughton, C. S., Clarke, A. D., Anderson, B. E., and Blake, D. R.: Airborne measurements of HCl from the marine
 boundary layer to the lower stratosphere over the North Pacific Ocean during INTEX-B, Atmos. Chem. Phys. Discuss.,
 8, 3563, 3595, https://doi.org/10.5104/acpd.8, 3563, 2008, 2008
- 33 8, 3563-3595, https://doi.org/10.5194/acpd-8-3563-2008, 2008.
- Kim, M. J., Farmer, D. K., and Bertram, T. H.: A controlling role for the air-sea interface in the chemical processing
 of reactive nitrogen in the coastal marine boundary layer, Proceedings of the National Academy of Sciences, 111,
 3943-3948, 10.1073/pnas.1318694111, 2014.
- Kim, P. S., Jacob, D. J., Fisher, J. A., Travis, K., Yu, K., Zhu, L., Yantosca, R. M., Sulprizio, M. P., Jimenez, J. L.,
 Campuzano-Jost, P., Froyd, K. D., Liao, J., Hair, J. W., Fenn, M. A., Butler, C. F., Wagner, N. L., Gordon, T. D.,
 Welti, A., Wennberg, P. O., Crounse, J. D., St. Clair, J. M., Teng, A. P., Millet, D. B., Schwarz, J. P., Markovic, M.
 Z., and Perring, A. E.: Sources, seasonality, and trends of southeast US aerosol: an integrated analysis of surface,
- 41 aircraft, and satellite observations with the GEOS-Chem chemical transport model, Atmos. Chem. Phys., 15, 10411-
- 42 10433, 10.5194/acp-15-10411-2015, 2015.

- 1 Knipping, E. M., and Dabdub, D.: Modeling Cl2 formation from aqueous NaCl particles: Evidence for interfacial
- 2 reactions and importance of Cl2 decomposition in alkaline solution, Journal of Geophysical Research: Atmospheres,
- 3 107(D18), 4360, doi:10.1029/2001JD000867, 2002.
- Koepke, P., M. Hess, I. Schult, and E.P. Shettle Global Aerosol Data Set, Report No. 243, Max-Planck-Institut für
 Meteorologie, Hamburg, 1997.
- Kolesar, K. R., Mattson, C. N., Peterson, P. K., May, N. W., Prendergast, R. K., and Pratt, K. A.: Increases in
 wintertime PM2.5 sodium and chloride linked to snowfall and road salt application, Atmospheric Environment, 177,
- 8 195-202, https://doi.org/10.1016/j.atmosenv.2018.01.008, 2018.
- Koo, B., Gaydos, T. M., and Pandis, S. N.: Evaluation of the Equilibrium, Dynamic, and Hybrid Aerosol Modeling
 Approaches, Aerosol Science and Technology, 37, 53-64, 10.1080/02786820300893, 2003.
- Lawler, M. J., Finley, B. D., Keene, W. C., Pszenny, A. A. P., Read, K. A., von Glasow, R., and Saltzman, E. S.:
 Pollution-enhanced reactive chlorine chemistry in the eastern tropical Atlantic boundary layer, Geophysical Research
- 13 Letters, 36, 10.1029/2008gl036666, 2009.
- Lawler, M. J., Sander, R., Carpenter, L. J., Lee, J. D., von Glasow, R., Sommariva, R., and Saltzman, E. S.: HOCl and
 Cl<sub>2</sub> observations in marine air, Atmospheric Chemistry and Physics, 11, 7617-7628,
 10.5194/acp-11-7617-2011, 2011.
- 17 Lee, B. H., Lopez-Hilfiker, F. D., Schroder, J. C., Campuzano-Jost, P., Jimenez, J. L., McDuffie, E. E., Fibiger, D. L.,
- 18 Veres, P. R., Brown, S. S., Campos, T. L., Weinheimer, A. J., Flocke, F. F., Norris, G., O'Mara, K., Green, J. R.,
- 19 Fiddler, M. N., Bililign, S., Shah, V., Jaeglé, L., and Thornton, J. A.: Airborne Observations of Reactive Inorganic
- 20 Chlorine and Bromine Species in the Exhaust of Coal-Fired Power Plants, Journal of Geophysical Research:
- 21 Atmospheres, 123, 11,225-211,237, 10.1029/2018jd029284, 2018.
- Lewis, E., and Schwartz, S.: Comment on "size distribution of sea-salt emissions as a function of relative humidity",
 Atmos. Env., 40 (3): 588-590., 2006.
- Lewis, E., and Schwartz, S.: Sea Salt Aerosol Production: Mechanisms, Methods, Measurements and Models, 2013.
- 25 Liao, J., Huey, L. G., Liu, Z., Tanner, D. J., Cantrell, C. A., Orlando, J. J., Flocke, F. M., Shepson, P. B., Weinheimer,
- A. J., Hall, S. R., Ullmann, K., Beine, H. J., Wang, Y., Ingall, E. D., Stephens, C. R., Hornbrook, R. S., Apel, E. C.,
- 27 Riemer, D., Fried, A., Mauldin, R. L., Smith, J. N., Staebler, R. M., Neuman, J. A., and Nowak, J. B.: High levels of
- 28 molecular chlorine in the Arctic atmosphere, Nature Geoscience, 7, 91-94, 10.1038/ngeo2046, 2014.
- Liu, H., Jacob, D. J., Bey, I., and Yantosca, R. M.: Constraints from 210Pb and 7Be on wet deposition and transport in a global three-dimensional chemical tracer model driven by assimilated meteorological fields, Journal of
- 31 Geophysical Research: Atmospheres, 106, 12109-12128, doi:10.1029/2000JD900839, 2001.
- Liu, Q., and Margerum, D. W.: Equilibrium and Kinetics of Bromine Chloride Hydrolysis, Environmental Science &
 Technology, 35, 1127-1133, 10.1021/es001380r, 2001.
- Liu, Y., Fan, Q., Chen, X., Zhao, J., Ling, Z., Hong, Y., Li, W., Chen, X., Wang, M., and Wei, X.: Modeling the impact of chlorine emissions from coal combustion and prescribed waste incineration on tropospheric ozone formation in China, Atmospheric Chemistry and Physics, 18, 2709-2724, 10.5194/acp-18-2709-2018, 2018.
- Lobert, J. M., Keene, W. C., Logan, J. A., and Yevich, R.: Global chlorine emissions from biomass burning: Reactive
 Chlorine Emissions Inventory, Journal of Geophysical Research: Atmospheres, 104, 8373-8389,
 doi:10.1029/1998JD100077, 1999.
- 40 Long, M. S., Keene, W. C., Easter, R. C., Sander, R., Liu, X., Kerkweg, A., and Erickson, D.: Sensitivity of 41 tropospheric chemical composition to halogen-radical chemistry using a fully coupled size-resolved multiphase

chemistry–global climate system: halogen distributions, aerosol composition, and sensitivity of climate-relevant
 gases, Atmospheric Chemistry and Physics, 14, 3397-3425, 10.5194/acp-14-3397-2014, 2014.

Malm, W. C., Sisler, J. F., Huffman, D., Eldred, R. A., and Cahill, T. A.: Spatial and seasonal trends in particle
 concentration and optical extinction in the United States, Journal of Geophysical Research: Atmospheres, 99, 1347 1370, doi:10.1029/93JD02916, 1994.

- Martin, R. V., Jacob, D. J., Yantosca, R. M., Chin, M., and Ginoux, P.: Global and regional decreases in tropospheric
 oxidants from photochemical effects of aerosols, Journal of Geophysical Research: Atmospheres, 108,
 doi:10.1029/2002JD002622, 2003.
- 9 Massucci, M., Clegg, S. L., and Brimblecombe, P.: Equilibrium Partial Pressures, Thermodynamic Properties of 10 Aqueous and Solid Phases, and Cl2 Production from Aqueous HCl and HNO3 and Their Mixtures, The Journal of
- 11 Physical Chemistry A, 103, 4209-4226, 10.1021/jp9847179, 1999.

12 McCulloch, A., Aucott, M. L., Benkovitz, C. M., Graedel, T. E., Kleiman, G., Midgley, P. M., and Li, Y.-F.: Global

13 emissions of hydrogen chloride and chloromethane from coal combustion, incineration and industrial activities:

- 14 Reactive Chlorine Emissions Inventory, Journal of Geophysical Research: Atmospheres, 104, 8391-8403,
- 15 10.1029/1999jd900025, 1999.
- 16 McDuffie, E. E., Fibiger, D. L., Dubé, W. P., Lopez Hilfiker, F., Lee, B. H., Jaeglé, L., Guo, H., Weber, R. J., Reeves,
- 17 J. M., Weinheimer, A. J., Schroder, J. C., Campuzano-Jost, P., Jimenez, J. L., Dibb, J. E., Veres, P., Ebben, C., Sparks,
- 18 T. L., Wooldridge, P. J., Cohen, R. C., Campos, T., Hall, S. R., Ullmann, K., Roberts, J. M., Thornton, J. A., and
- Brown, S. S.: ClNO2 Yields From Aircraft Measurements During the 2015 WINTER Campaign and Critical
- 20 Evaluation of the Current Parameterization, Journal of Geophysical Research: Atmospheres, 123, 12,994-913,015,
- 21 10.1029/2018jd029358, 2018a.
- 22 McDuffie, E. E., Fibiger, D. L., Dubé, W. P., Lopez-Hilfiker, F., Lee, B. H., Thornton, J. A., Shah, V., Jaeglé, L.,
- 23 Guo, H., Weber, R. J., Michael Reeves, J., Weinheimer, A. J., Schroder, J. C., Campuzano-Jost, P., Jimenez, J. L.,
- 24 Dibb, J. E., Veres, P., Ebben, C., Sparks, T. L., Wooldridge, P. J., Cohen, R. C., Hornbrook, R. S., Apel, E. C., Campos,
- 25 T., Hall, S. R., Ullmann, K., and Brown, S. S.: Heterogeneous N2O5 Uptake During Winter: Aircraft Measurements
- 26 During the 2015 WINTER Campaign and Critical Evaluation of Current Parameterizations, Journal of Geophysical
- 27 Research: Atmospheres, 123, 4345-4372, doi:10.1002/2018JD028336, 2018b.
- Meng, Z., and Seinfeld, J. H.: Time scales to achieve atmospheric gas-aerosol equilibrium for volatile species,
 Atmospheric Environment, 30, 2889-2900, https://doi.org/10.1016/1352-2310(95)00493-9, 1996.
- Mielke, L. H., Furgeson, A., and Osthoff, H. D.: Observation of ClNO2 in a Mid-Continental Urban Environment,
 Environmental Science & Technology, 45, 8889-8896, 10.1021/es201955u, 2011.
- Mielke, L. H., Stutz, J., Tsai, C., Hurlock, S. C., Roberts, J. M., Veres, P. R., Froyd, K. D., Hayes, P. L., Cubison, M.
 J., Jimenez, J. L., Washenfelder, R. A., Young, C. J., Gilman, J. B., de Gouw, J. A., Flynn, J. H., Grossberg, N., Lefer,
 B. L., Liu, J., Weber, R. J., and Osthoff, H. D.: Heterogeneous formation of nitryl chloride and its role as a nocturnal
 NOx reservoir species during CalNex-LA 2010, Journal of Geophysical Research: Atmospheres, 118, 10,638-610,652,
 doi:10.1002/jgrd.50783, 2013.
- Mielke, L. H., Furgeson, A., Odame-Ankrah, C. A., and Osthoff, H. D.: Ubiquity of ClNO2 in the urban boundary
 layer of Calgary, Alberta, Canada, Canadian Journal of Chemistry, 94, 414-423, 10.1139/cjc-2015-0426, 2015.
- 39 Millet, D. B., Guenther, A., Siegel, D. A., Nelson, N. B., Singh, H. B., de Gouw, J. A., Warneke, C., Williams, J.,
- 40 Eerdekens, G., Sinha, V., Karl, T., Flocke, F., Apel, E., Riemer, D. D., Palmer, P. I., and Barkley, M.: Global
- 41 atmospheric budget of acetaldehyde: 3-D model analysis and constraints from in-situ and satellite observations,
- 42 Atmos. Chem. Phys., 10, 3405-3425, 10.5194/acp-10-3405-2010, 2010.

- 1 Murray, L. T., Jacob, D. J., Logan, J. A., Hudman, R. C., and Koshak, W. J.: Optimized regional and interannual variability of lightning in a global chemical transport model constrained by LIS/OTD satellite data, Journal of
- 2
- 3 Geophysical Research: Atmospheres, 117, doi:10.1029/2012JD017934, 2012.

4 Naik, V., Voulgarakis, A., Fiore, A. M., Horowitz, L. W., Lamarque, J. F., Lin, M., Prather, M. J., Young, P. J.,

5 Bergmann, D., Cameron-Smith, P. J., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R., Eyring, V., Faluvegi, G.,

Folberth, G. A., Josse, B., Lee, Y. H., MacKenzie, I. A., Nagashima, T., van Noije, T. P. C., Plummer, D. A., Righi, 6

- 7 M., Rumbold, S. T., Skeie, R., Shindell, D. T., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., and Zeng, G.:
- 8 Preindustrial to present-day changes in tropospheric hydroxyl radical and methane lifetime from the Atmospheric 9 Chemistry and Climate Model Intercomparison Project (ACCMIP), Atmospheric Chemistry and Physics, 13, 5277-
- 10 5298, 10.5194/acp-13-5277-2013, 2013.
- 11 Ordóñez, C., Lamarque, J. F., Tilmes, S., Kinnison, D. E., Atlas, E. L., Blake, D. R., Sousa Santos, G., Brasseur, G.,
- 12 and Saiz-Lopez, A.: Bromine and iodine chemistry in a global chemistry-climate model: description and evaluation
- 13 of very short-lived oceanic sources, Atmos. Chem. Phys., 12, 1423-1447, 10.5194/acp-12-1423-2012, 2012.
- 14 Osthoff, H. D., Roberts, J. M., Ravishankara, A. R., Williams, E. J., Lerner, B. M., Sommariva, R., Bates, T. S.,
- 15 Coffman, D., Quinn, P. K., Dibb, J. E., Stark, H., Burkholder, J. B., Talukdar, R. K., Meagher, J., Fehsenfeld, F. C.,
- 16 and Brown, S. S.: High levels of nitryl chloride in the polluted subtropical marine boundary layer, Nature Geoscience,
- 17 1, 324-328, 10.1038/ngeo177, 2008.
- 18 Parrella, J. P., Jacob, D. J., Liang, Q., Zhang, Y., Mickley, L. J., Miller, B., Evans, M. J., Yang, X., Pyle, J. A., Theys,
- 19 N., and Van Roozendael, M.: Tropospheric bromine chemistry: implications for present and pre-industrial ozone and
- 20 mercury, Atmospheric Chemistry and Physics, 12, 6723-6740, 10.5194/acp-12-6723-2012, 2012.
- 21 Philip, S., Martin, R. V., Snider, G., Weagle, C. L., van Donkelaar, A., Brauer, M., Henze, D. K., Klimont, Z.,
- 22 Venkataraman, C., Guttikunda, S. K., and Zhang, Q.: Anthropogenic fugitive, combustion and industrial dust is a
- 23 significant, underrepresented fine particulate matter source in global atmospheric models, Environmental Research
- 24 Letters, 12, 044018, 10.1088/1748-9326/aa65a4, 2017.
- 25 Phillips, G. J., Tang, M. J., Thieser, J., Brickwedde, B., Schuster, G., Bohn, B., Lelieveld, J., and Crowley, J. N.:
- Significant concentrations of nitryl chloride observed in rural continental Europe associated with the influence of sea 26
- 27 salt chloride and anthropogenic emissions, Geophysical Research Letters, 39, n/a-n/a, 10.1029/2012gl051912, 2012.
- 28 Pilinis, C., Capaldo, K. P., Nenes, A., and Pandis, S. N.: MADM-A New Multicomponent Aerosol Dynamics Model, 29 Aerosol Science and Technology, 32, 482-502, 10.1080/027868200303597, 2000.
- 30 Platt, U., Allan, W., and Lowe, D.: Hemispheric average Cl atom concentration from 13C/12C ratios in atmospheric 31 methane, Atmos. Chem. Phys., 4, 2393-2399, 10.5194/acp-4-2393-2004, 2004.
- 32 Priestley, M., le Breton, M., Bannan, T. J., Worrall, S. D., Bacak, A., Smedley, A. R. D., Reyes-Villegas, E., Mehra, A., Allan, J., Webb, A. R., Shallcross, D. E., Coe, H., and Percival, C. J.: Observations of organic and inorganic 33
- 34 chlorinated compounds and their contribution to chlorine radical concentrations in an urban environment in northern
- 35 Europe during the wintertime, Atmos. Chem. Phys., 18, 13481-13493, 10.5194/acp-18-13481-2018, 2018.
- 36 Prinn, R. G., Weiss, R. F., Arduini, J., Arnold, T., DeWitt, H. L., Fraser, P. J., Ganesan, A. L., Gasore, J., Harth, C. 37 M., Hermansen, O., Kim, J., Krummel, P. B., Li, S., Loh, Z. M., Lunder, C. R., Maione, M., Manning, A. J., Miller, B. R., Mitrevski, B., Mühle, J., O'Doherty, S., Park, S., Reimann, S., Rigby, M., Saito, T., Salameh, P. K., Schmidt, 38 39 R., Simmonds, P. G., Steele, L. P., Vollmer, M. K., Wang, R. H., Yao, B., Yokouchi, Y., Young, D., and Zhou, L.: 40 History of chemically and radiatively important atmospheric gases from the Advanced Global Atmospheric Gases
- 41 Experiment (AGAGE), Earth Syst. Sci. Data, 10, 985-1018, 10.5194/essd-10-985-2018, 2018.
- Pszenny, A. A. P., Keene, W. C., Jacob, D. J., Fan, S., Maben, J. R., Zetwo, M. P., Springer-Young, M., and Galloway, 42
- 43 J. N.: Evidence of inorganic chlorine gases other than hydrogen chloride in marine surface air, Geophysical Research
- 44 Letters, 20, 699-702, doi:10.1029/93GL00047, 1993.

- 1 Pszenny, A. A. P., Moldanová, J., Keene, W. C., Sander, R., Maben, J. R., Martinez, M., Crutzen, P. J., Perner, D.,
- 2 and Prinn, R. G.: Halogen cycling and aerosol pH in the Hawaiian marine boundary layer, Atmos. Chem. Phys., 4,
- 3 147-168, 10.5194/acp-4-147-2004, 2004.
- Reff, A., Bhave, P. V., Simon, H., Pace, T. G., Pouliot, G. A., Mobley, J. D., and Houyoux, M.: Emissions Inventory
 of PM2.5 Trace Elements across the United States, Environmental Science & Technology, 43, 5790-5796,
 10.1021/es802930x, 2009.
- Riedel, T. P., Wagner, N. L., Dubé, W. P., Middlebrook, A. M., Young, C. J., Öztürk, F., Bahreini, R., VandenBoer,
 T. C., Wolfe, D. E., Williams, E. J., Roberts, J. M., Brown, S. S., and Thornton, J. A.: Chlorine activation within urban
 or power plant plumes: Vertically resolved ClNO2and Cl2measurements from a tall tower in a polluted continental
- 10 setting, Journal of Geophysical Research: Atmospheres, 118, 8702-8715, 10.1002/jgrd.50637, 2013.
- Roberts, J. M., Osthoff, H. D., Brown, S. S., and Ravishankara, A. R.: N2O5 Oxidizes Chloride to Cl2 in Acidic
 Atmospheric Aerosol, Science, 321, 1059-1059, 10.1126/science.1158777, 2008.
- Roberts, J. M., Osthoff, H. D., Brown, S. S., Ravishankara, A. R., Coffman, D., Quinn, P., and Bates, T.: Laboratory
 studies of products of N2O5 uptake on Cl– containing substrates, Geophysical Research Letters, 36,
 doi:10.1029/2009GL040448, 2009.
- Saiz-Lopez, A., and von Glasow, R.: Reactive halogen chemistry in the troposphere, Chem Soc Rev, 41, 6448-6472,
 10.1039/c2cs35208g, 2012.
- 18 Saiz-Lopez, A., Fernandez, R. P., Ordóñez, C., Kinnison, D. E., Gómez Martín, J. C., Lamarque, J.-F., and Tilmes,
- S.: Iodine chemistry in the troposphere and its effect on ozone, Atmos. Chem. Phys., 14, 13119-13143,
- 20 https://doi.org/10.5194/acp-14-13119-2014, 2014.
- 21 Sander, R., Pszenny, A. A. P., Keene, W. C., Crete, E., Deegan, B., Long, M. S., Maben, J. R., and Young, A. H.: Gas
- 22 phase acid, ammonia and aerosol ionic and trace element concentrations at Cape Verde during the Reactive Halogens
- in the Marine Boundary Layer (RHaMBLe) 2007 intensive sampling period, Earth System Science Data, 5, 385-392,
 - 24 10.5194/essd-5-385-2013, 2013.
 - Sander, R.: Compilation of Henry's law constants (version 4.0) for water as solvent, Atmospheric Chemistry and
 Physics, 15, 4399-4981, 10.5194/acp-15-4399-2015, 2015.
 - Sanhueza, E., and Garaboto, A.: Gaseous HCl at a remote tropical continental site, Tellus B: Chemical and Physical
 Meteorology, 54, 412-415, 10.3402/tellusb.v54i4.16675, 2002.
 - Sarwar, G., Simon, H., Bhave, P., and Yarwood, G.: Examining the impact of heterogeneous nitryl chloride production
 on air quality across the United States, Atmospheric Chemistry and Physics, 12, 6455-6473, 10.5194/acp-12-6455-
 - 31 2012, 2012.
 - Sarwar, G., Simon, H., Xing, J., and Mathur, R.: Importance of tropospheric ClNO2 chemistry across the Northern
 Hemisphere, Geophysical Research Letters, 41, 4050-4058, doi:10.1002/2014GL059962, 2014.
 - 34 Schmidt, J. A., Jacob, D. J., Horowitz, H. M., Hu, L., Sherwen, T., Evans, M. J., Liang, Q., Suleiman, R. M., Oram,
 - 35 D. E., Le Breton, M., Percival, C. J., Wang, S., Dix, B., and Volkamer, R.: Modeling the observed tropospheric BrO
 - 36 background: Importance of multiphase chemistry and implications for ozone, OH, and mercury, Journal of
 - 37 Geophysical Research: Atmospheres, 121, 11,819-811,835, 10.1002/2015jd024229, 2016.
 - Schweitzer, F., Mirabel, P., & George, C.: Multiphase Chemistry of N2O5, ClNO2, and BrNO2. The Journal of Physical Chemistry A, 102(22), 3942-3952. https://doi.org/10.1021/jp980748s, 1998.
 - Sherwen, T., Evans, M. J., Carpenter, L. J., Andrews, S. J., Lidster, R. T., Dix, B., Koenig, T. K., Sinreich, R., Ortega,
 I., Volkamer, R., Saiz-Lopez, A., Prados-Roman, C., Mahajan, A. S., and Ordóñez, C.: Iodine's impact on tropospheric

- 1 oxidants: a global model study in GEOS-Chem, Atmospheric Chemistry and Physics, 16, 1161-1186, 10.5194/acp-2 16-1161-2016, 2016a.
- 3 Sherwen, T., Schmidt, J. A., Evans, M. J., Carpenter, L. J., Großmann, K., Eastham, S. D., Jacob, D. J., Dix, B.,
- 4 Koenig, T. K., Sinreich, R., Ortega, I., Volkamer, R., Saiz-Lopez, A., Prados-Roman, C., Mahajan, A. S., and Ordóñez,
- 5 C.: Global impacts of tropospheric halogens (Cl, Br, I) on oxidants and composition in GEOS-Chem, Atmospheric
- 6 Chemistry and Physics, 16, 12239-12271, 10.5194/acp-16-12239-2016, 2016b.
- Sherwen, T., Evans, M. J., Sommariva, R., Hollis, L. D. J., Ball, S. M., Monks, P. S., Reed, C., Carpenter, L. J., Lee,
 J. D., Forster, G., Bandy, B., Reeves, C. E., and Bloss, W. J.: Effects of halogens on European air-quality, Faraday
 Discuss, 200, 75-100, 10.1039/c7fd00026j, 2017.
- Simmonds, P. G., Manning, A. J., Cunnold, D. M., McCulloch, A., O'Doherty, S., Derwent, R. G., Krummel, P. B.,
 Fraser, P. J., Dunse, B., Porter, L. W., Wang, R. H. J., Greally, B. R., Miller, B. R., Salameh, P., Weiss, R. F., and
 Prinn, R. G.: Global trends, seasonal cycles, and European emissions of dichloromethane, trichloroethene, and
 tetrachloroethene from the AGAGE observations at Mace Head, Ireland, and Cape Grim, Tasmania, Journal of
 Geophysical Research: Atmospheres, 111, doi:10.1029/2006JD007082, 2006.
- Simpson, W. R., Brown, S. S., Saiz-Lopez, A., Thornton, J. A., and Glasow, R.: Tropospheric halogen chemistry:
 sources, cycling, and impacts, Chem Rev, 115, 4035-4062, 10.1021/cr5006638, 2015.
- 17 Singh, H. B., and Kasting, J. F.: Chlorine-hydrocarbon photochemistry in the marine troposphere and lower 18 stratosphere, Journal of Atmospheric Chemistry, 7, 261-285, 10.1007/BF00130933, 1988.
- Singh, H. B., Thakur, A. N., Chen, Y. E., and Kanakidou, M.: Tetrachloroethylene as an indicator of low Cl atom
 concentrations in the troposphere, Geophysical Research Letters, 23, 1529-1532, 10.1029/96gl01368, 1996.
- Sommariva, R., and von Glasow, R.: Multiphase halogen chemistry in the tropical Atlantic Ocean, Environ Sci
 Technol, 46, 10429-10437, 10.1021/es300209f, 2012.
- 23 Sommariva, R., Hollis, L. D. J., Sherwen, T., Baker, A. R., Ball, S. M., Bandy, B. J., Bell, T. G., Chowdhury, M. N.,
- Cordell, R. L., Evans, M. J., Lee, J. D., Reed, C., Reeves, C. E., Roberts, J. M., Yang, M., and Monks, P. S.: Seasonal
 and geographical variability of nitryl chloride and its precursors in Northern Europe, Atmospheric Science Letters,
 19, e844, doi:10.1002/asl.844, 2018.
- Straub, D. J., Lee, T., and Collett Jr., J. L.: Chemical composition of marine stratocumulus clouds over the eastern
 Pacific Ocean, Journal of Geophysical Research: Atmospheres, 112, doi:10.1029/2006JD007439, 2007.
- Tham, Y. J., Yan, C., Xue, L., Zha, Q., Wang, X., and Wang, T.: Presence of high nitryl chloride in Asian coastal
 environment and its impact on atmospheric photochemistry, Chinese Science Bulletin, 59, 356-359, 10.1007/s11434013-0063-y, 2014.
- Thornton, J. A., Kercher, J. P., Riedel, T. P., Wagner, N. L., Cozic, J., Holloway, J. S., Dubé, W. P., Wolfe, G. M.,
 Quinn, P. K., Middlebrook, A. M., Alexander, B., and Brown, S. S.: A large atomic chlorine source inferred from
 mid-continental reactive nitrogen chemistry, Nature, 464, 271, 10.1038/nature08905, 2010.
- 35
- Toon, O. B., Maring, H., Dibb, J., Ferrare, R., Jacob, D. J., Jensen, E. J., Luo, Z. J., Mace, G. G., Pan, L. L., Pfister,
 L., Rosenlof, K. H., Redemann, J., Reid, J. S., Singh, H. B., Thompson, A. M., Yokelson, R., Minnis, P., Chen, G.,
 Jucks, K. W., and Pszenny, A.: Planning, implementation, and scientific goals of the Studies of Emissions and
 Atmospheric Composition, Clouds and Climate Coupling by Regional Surveys (SEAC4RS) field mission, Journal of
- 40 Geophysical Research: Atmospheres, 121, 4967-5009, doi:10.1002/2015JD024297, 2016.
- 41 US EPA: 2014 National Emissions Inventory, https://www.epa.gov/air-emissions-inventories/2014-national-42 emissions-inventory-nei-data, last access: 28 August 2018, 2018.

- 1 van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R.
- 2 S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, 3 agricultural, and peat fires (1997–2009), Atmos. Chem. Phys., 10, 11707-11735, 10.5194/acp-10-11707-2010, 2010.
- agricultural, and peat fires (1997-2009), Atmos. Chem. Phys., 10, 11707-11735, 10.5194/acp-10-11707-2010, 2010

4 Voulgarakis, A., Naik, V., Lamarque, J. F., Shindell, D. T., Young, P. J., Prather, M. J., Wild, O., Field, R. D.,

5 Bergmann, D., Cameron-Smith, P., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R. M., Eyring, V., Faluvegi,

6 G., Folberth, G. A., Horowitz, L. W., Josse, B., MacKenzie, I. A., Nagashima, T., Plummer, D. A., Righi, M.,

- Rumbold, S. T., Stevenson, D. S., Strode, S. A., Sudo, K., Szopa, S., and Zeng, G.: Analysis of present day and future
 OH and methane lifetime in the ACCMIP simulations, Atmos. Chem. Phys., 13, 2563-2587, 10.5194/acp-13-2563-
- OH and methane methane in the ACCMIP simulations, Atmos. Chem. Phys., 15, 2505-2587, 10.5194/acp-15-2505 2013, 2013.
- 9 2015, 2015.
- 10 Wagner, N. L., Riedel, T. P., Young, C. J., Bahreini, R., Brock, C. A., Dubé, W. P., Kim, S., Middlebrook, A. M.,
- 11 Öztürk, F., Roberts, J. M., Russo, R., Sive, B., Swarthout, R., Thornton, J. A., VandenBoer, T. C., Zhou, Y., and
- Brown, S. S.: N2O5 uptake coefficients and nocturnal NO2 removal rates determined from ambient wintertime
- 13 measurements, Journal of Geophysical Research: Atmospheres, 118, 9331-9350, doi:10.1002/jgrd.50653, 2013.
- 14
- 15 Wang, T., Tham, Y. J., Xue, L., Li, Q., Zha, Q., Wang, Z., Poon, S. C. N., Dubé, W. P., Blake, D. R., Louie, P. K. K.,
- 16 Luk, C. W. Y., Tsui, W., and Brown, S. S.: Observations of nitryl chloride and modeling its source and effect on ozone
- in the planetary boundary layer of southern China, Journal of Geophysical Research: Atmospheres, 121, 2476-2489,
- 18 doi:10.1002/2015JD024556, 2016.
- 19 Wang, T. X., Kelley, M. D., Cooper, J. N., Beckwith, R. C., and Margerum, D. W.: Equilibrium, Kinetic, and UV-
- 20 Spectral Characteristics of Aqueous Bromine Chloride, Bromine, and Chlorine Species, Inorganic Chemistry, 33,
- 21 5872-5878, 10.1021/ic00103a040, 1994.
- Wang, Y., Jacob, D. J., and Logan, J. A.: Global simulation of tropospheric O3-NO x -hydrocarbon chemistry: 1.
 Model formulation, Journal of Geophysical Research: Atmospheres, 103, 10713-10725, doi:10.1029/98JD00158, 1998.
- Wesely, M. L.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models,
 Atmospheric Environment (1967), 23, 1293-1304, https://doi.org/10.1016/0004-6981(89)90153-4, 1989.
- WMO: Scientific Assessment of Ozone Depletion: 2014, World Meteorological Organization, Global Ozone Research
 and Monitoring Project—Report No. 55, 416 pp., World Meteorological Organization, Geneva, Switzerland, 2014.
- Yang, X., Cox, R. A., Warwick, N. J., Pyle, J. A., Carver, G. D., O'Connor, F. M., and Savage, N. H.: Tropospheric
 bromine chemistry and its impacts on ozone: A model study, Journal of Geophysical Research, 110,
 10.1029/2005jd006244, 2005.
- 32 Zhu, L., Jacob, D. J., Eastham, S. D., Sulprizio, M. P., Wang, X., Sherwen, T., Evans, M. J., Chen, Q., Alexander, B.,
- 33 Koenig, T. K., Volkamer, R., Huey, L. G., Le Breton, M., Bannan, T. J., and Percival, C. J.: Effect of sea-salt aerosol
- on tropospheric bromine chemistry, Atmos. Chem. Phys. Discuss., 2018, 1-17, 10.5194/acp-2018-1239, 2018.
- 35 36

1 Ta	ble 1: Global sources a	nd sinks of gas-phas	e inorganic (Cl _v) and	reactive (Cl*) tropo	spheric chlorine ^a .
-------------	-------------------------	----------------------	------------------------------------	----------------------	---------------------------------

	Cly (Gg Cl a ⁻¹)	Cl* (Gg Cl a ⁻¹)
Total source	75200	25000
Sea Salt	63900	11900
Acid displacement ^b	52000	-
$HOBr + Cl^{-}$	8590	8590
$N_2O_5 + Cl^2$	1810	1810
HOI, $IONO_x^c + Cl^-$	641	641
$CINO_2 + Cl^-$	327	327
$OH + Cl^{-}$	403	403
$ClNO_3 + Cl^-$	64	64
$HOCl + Cl^{-}$	61	61
HCl + OH	-	9720
Organochlorines	3320	3300
$CH_3Cl + OH^{d}$	2200	2180
$CH_2Cl_2 + OH$	780	780
$CHCl_3 + OH$	298	298
$CH_2ICl + OH$	46	46
Stratosphere ^e	380	64
Anthropogenic HCl ^f	(6660)	-
Open fires	7640	-
Total sink	75200	25000
Deposition	71400	346
Dry	35200	170
Wet	36200	176
Uptake by alkaline SSA	3800	-
Conversion to HCl ^g	-	24600
Tropospheric mass (Gg)	316	12
Lifetime (hours)	37	3.8

2 3 ^a Annual totals for 2016 computed from GEOS-Chem. Gas-phase inorganic chlorine is defined as $Cl_y \equiv Cl + 2 \times Cl_2 + 2 \times Cl_2O_2 + Cl$ $CINO_2 + CINO_3 + CIO + CIOO + OCIO + BrCl + ICl + HOCl + HCl.$ Reactive chlorine is defined as $Cl^* \equiv Cl_y - HCl$. Thus the

source of HCl can be inferred from the Table entries as $Cl_y - Cl^*$. The definition of Cl_y excludes aerosol Cl⁻, which has a very large 4 5 6

sea salt source of 1780 Tg Cl a⁻¹ but is mainly removed by deposition, HCl is the dominant component of Cl_y but is also mostly removed by deposition. Reactive chlorine Cl* is the chemical family principally involved in radical cycling.

7 ^b Net production minus loss of HCl from acid aerosol displacement by HNO₃ and H₂SO₄ computed as thermodynamic equilibrium. 8 ^c $IONO_x \equiv IONO + IONO$

9 ^d The source from CH₃Cl + Cl is not shown since it contributes < 1% of CH₃Cl oxidation. Same for other organochlorines.

10 ^e Net stratospheric input to the troposphere.

11 ^f Coal combustion, waste incineration, and industrial activities. These emissions are only included in a sensitivity simulation (see

12 Section 2.2 and 4.2 for details) and are therefore listed here in parentheses. Emissions of anthropogenic fugitive dust are estimated 13 as less than 390 Gg a⁻¹ (Section 2,2) and are not included in the model.

^g From reactions of Cl atoms (see Figure 1). 14

1 Table 2: Heterogeneous reactions of Cl⁻and reactive uptake coefficients (y) ^a.

	Reaction	Reactive uptake coefficient (y)	Footnote
R3	$N_2O_5 + \phi Cl^- + (1 - \phi)H_2O \rightarrow \phi ClNO_2 + (2 - \phi)NO_3^- + 2(1 - \phi)H^+$	$\begin{split} \gamma &= Bk_{2f}' \left(1 - \frac{1}{\left(\frac{k_3[\text{H}_2\text{O}]}{k_{2b}[\text{NO}_3^-]}\right) + 1 + \left(\frac{k_4[\text{Cl}^-]}{k_{2b}[\text{NO}_3^-]}\right)} \right) \\ k_{2f}' &= \beta \left(1 - e^{-\delta[\text{H}_2\text{O}]} \right); \ \varphi = \left(\frac{k_2[\text{H}_2\text{O}]}{k_3[\text{Cl}^-]} + 1\right)^{-1} \\ B &= 3.2 \times 10^{-8}s; \ k_3/k_2 = 450 \end{split}$	b
		$\beta = 1.15 \times 10^{6} s^{-1}; \ \delta = 0.13 M^{-1}$	
		$k_3/k_{2b} = 0.06$; $k_4/k_{2b} = 29$	
R4	$OH + Cl^- \rightarrow 0.5Cl_2 + OH^-$	$\gamma=0.04[Cl^-]$	с
		$\gamma = \left(\frac{1}{\Gamma_{\rm b}} + \frac{1}{\alpha_{\rm b}}\right)^{-1}$	
R5	$HOBr + Cl^- + H^+ \rightarrow BrCl + H_0$	$\Gamma_{\rm b} = 4H_{HOBr}RTI_rk_b[{\rm Cl}^-][{\rm H}^+]f(r,I_r)/c$	d
K5	$HOBT + CI + H^+ \rightarrow BrCI + H_2O$	$I_r = \sqrt{D_l / (k_b [\text{Cl}^-][\text{H}^+])}$; $\alpha_b = 0.6$	
		$k_b = 2.3 \times 10^{10} \mathrm{M}^{-2} \mathrm{s}^{-1}$; $D_l = 1.4 \times 10^{-5} \mathrm{cm}^2 \mathrm{s}^{-1}$	
R6	$\text{ClNO}_3 + \text{Cl}^- \rightarrow \text{Cl}_2 + \text{NO}_3^-$	$\gamma = 0.0244$	e
R7	$CINO_2 + Cl^- \rightarrow NO_2^- + Cl_2$	$\gamma = \left(\frac{1}{\Gamma_{\rm b}} + \frac{1}{\alpha_{\rm b}}\right)^{-1} (\text{pH} < 2), \gamma = 0 (\text{pH} > 2)$ $\Gamma_{\rm b} = 4H_{CINO_2}RTI_r k^{II} [\text{CI}^-]f(r, I_r)/c$	
		$I_r = \sqrt{D_l / (k^{II} [\text{Cl}^{-1}])}; \alpha_b = 0.01$ $k^{II} = 10^7 \text{M}^{-2} \text{s}^{-1}; D_l = 1 \times 10^{-5} \text{cm}^2 \text{s}^{-1}$	
R8	$CINO_2 + Br^- \rightarrow NO_2^- + BrCl$	$\gamma = \left(\frac{1}{\Gamma_{\rm b}} + \frac{1}{\alpha_{\rm b}}\right)^{-1}$ $\Gamma_{\rm b} = 4H_{CINO_2}RTI_rk^{II}[{\rm Br}^-]f(r, I_r)/c$ $I_r = \sqrt{D_l/(k^{II}[{\rm Br}^-])}; \alpha_b = 0.01$ $H_{CINO_2}{}^2D_lk^{II} = 0.101 {\rm Mcm}^2 {\rm s}^{-2}$	
R9	$HOCl + Cl^- + H^+ \rightarrow Cl_2 + H_2O$	$\begin{split} \gamma &= \min(\left(\frac{1}{\Gamma_{\rm b}} + \frac{1}{\alpha_{\rm b}}\right)^{-1}, 2 \times 10^{-4}) \\ \Gamma_{\rm b} &= 4H_{HOCl}RTI_{r}k_{t}[{\rm Cl}^{-}][{\rm H}^{+}]f(r, I_{r})/c \\ I_{r} &= \sqrt{D_{l}/(k_{t}[{\rm Cl}^{-}][{\rm H}^{+}])}; \alpha_{\rm b} = 0.8 \\ k_{t} &= 1.5 \times 10^{4}{\rm M}^{-2}{\rm s}^{-1}; D_{l} = 2 \times 10^{-5}{\rm cm}^{2}{\rm s}^{-1} \end{split}$	
R10	$\mathrm{NO}_3 + \mathrm{Cl}^- \rightarrow \mathrm{NO}_3^- + \mathrm{Cl}^-$	$\gamma = \left(\frac{1}{\Gamma_{\rm b}} + \frac{1}{\alpha_{\rm b}}\right)^{-1}$ $\Gamma_{\rm b} = 4H_{NO_3}RTI_rk'[{\rm Cl}^-]f(r, I_r)/c$ $I_{\rm r} = \sqrt{D_l/(k'[{\rm Cl}^-])}; \alpha_{\rm b} = 0.013$ $k' = 2.76 \times 10^6 {\rm M}^{-2} {\rm s}^{-1}; D_l = 1 \times 10^{-5} {\rm cm}^2 {\rm s}^{-1}$	
R11	$IONO_2 + Cl^- \rightarrow ICl + NO_3^-$	$\gamma=8.5{\times}10^{-3}$	f
R12	$IONO + Cl^- \rightarrow ICl + NO_2^-$	$\gamma = 0.017$	f
R13	$HOI + Cl^- \rightarrow ICl + OH^-$	$\gamma=8.5{\times}10^{-3}$	f

- ^a Formulations for the reactive uptake coefficient γ are from IUPAC (Ammann et al., 2013) unless stated otherwise in the footnote
- column. Brackets denote aqueous-phase concentrations in unit of M (moles per liter of water). R is the ideal gas constant. c is the
- average gas-phase thermal velocity for the reactant with Cl⁻. The reactive uptake coefficient is used to calculate the reaction rate
- following equation (1). $f(r, I_r) = \operatorname{coth}(r/I_r) (I_r/r)$ is a spherical correction to mass transfer where I_r is a reacto-diffusive length
- scale and r is the radius of the aerosol particle or cloud droplet.
- 1 2 3 4 5 6 7 ^b Bertram and Thornton (2009); Roberts et al. (2009). This parametrization is only applied for SNA and SSA aerosols. For N₂O₅
- uptake on other aerosols, we assume $\varphi = 0$ and follow Jaeglé et al., (2018).
- 8 ^c Knipping and Dabdub (2002)
- 9 ^d k_b is based on Liu and Margerum (2001). R5 competes with the heterogeneous reactions HOBr+Br⁻ and HOBr+S(IV) as given by
- 10 Chen et al. (2017). The BrCl product may either volatilize or react with Br to produce Br₂ and return Cl following Fickert et al. 11 (1999), as described in Section 5.2.
- 12 ^e Assumes that Cl⁻ is present in excess so that γ does not depend on [Cl⁻]. However, R6 competes with the heterogeneous
- 13 reaction ClNO₃+Br⁻ as given by Schmidt et al. (2016), with the branching ratio determined by the relative rates.
- 14 ^f These reactions are based on Sherwen et al. (2016a) and only take place in SSA.
- 15

1 Table 3: Chloride deficits in sea salt aerosol.^a

Location	Modeled Cl ⁻ deficit (%)	Measured Cl ⁻ deficit (%)
North Carolina coast	+40	-1 to +90
Townsville coast, Australia	+23	+33
California coast	+21	+2 to +75
Greenland Sea	+18	+6 to +22
North Atlantic Ocean	+14	-24 to +54
Equatorial Atlantic	+12	+11 to +64
Puerto Rico coast	+9	+7 to +25
Pacific Ocean	+6	-22 ~ +40
Cape Grim, Australia	+4	-50~+15

^a Deficits relative to seawater composition. Observations compiled by Graedel and Keene (1995) are reported there as ranges for individual regions and months, with the ranges likely reflecting measurement uncertainty rather than physical variability. Model

values are means for the regions and months of observations.

2

3

6 7

, 8 9

Table 4: Surface air mixing ratios of reactive chlorine (Cl*)^a

Location	Modeled Cl* (ppt)	Measured mean Cl^* (ppt)	Reference
Atlantic cruise near Europe	43	27 ^b	Keene et al. (2009)
Appledore Island (US east coast)	17	< 20	Keene et al. (2007)
Atlantic cruise near North Africa	5	< 24	Keene et al. (2009)
Southern Ocean cruise	4	< 24	Keene et al. (2009)
Hawaii	4	6	Pszenny et al. (2004)
Tropical Atlantic cruise	2	< 24	Keene et al. (2009)
Alert (Canada)	0.2	< 14	Impey et al. (1999)

10 ^a Reactive chlorine Cl* is the ensemble of gas-phase inorganic chlorine species excluding HCl. Measurements are 24- hour

11 averages. Model values are monthly means in 2016 taken for the same month and location as the observations.

12 ^b Median value

1 Table 5: Comparison of modeled maximum CINO ₂ mixing ratios to surface observation	1	Table 5: Comparison of m	nodeled maximum ClNO ₂	2 mixing ratios to surface	observations ^a
---	---	--------------------------	-----------------------------------	----------------------------	---------------------------

Location	Date	Observed (ppt)	Simulated (ppt)	References
Manchester, UK	Oct-Nov 2014	510	400	Priestley et al. (2018)
Weybourne, UK	Jun-Aug 2015	1100	1200	Sommariva et al. (2018)
East Anglia Coast, UK	Jan 2014	100	400	Bannan et al. (2017)
Leicester, UK	Feb 2016	730	760	Sommariva et al. (2018)
London, UK	July-Aug 2012	730	510	Bannan et al. (2015)
Calgary, Canada	Apr 2010	240	130	Mielke et al. (2011)
Calgary, Canada	Sep 2010-Mar 2011	340	170	Mielke et al. (2015)
Penlee Point, UK	Apr-May 2015	920	870	Sommariva et al. (2018)
Kleiner Feldberg, Germany	Aug-Sep 2011	850	400	Phillips et al. (2012)
Long Island Sound	Mar 2008	200	210	Kercher et al. (2009)
Olympic Park, South Korea	May-Jun 2016	780	520	Jeong et al. (2018)
Taehwa Research Forest, South Korea	May-Jun 2016	2600	220	Jeong et al. (2018)
Boulder, Colorado	Feb 2009	440	130	Thornton et al. (2010)
Pasadena, California	May-Jun 2010	3500	360	Mielke et al. (2013)
Offshore of Los Angeles, California	May-Jun 2010	1800	500	Riedel et al. (2013)
La Jolla, California	Feb 2013	65	65	Kim et al. (2014)
Houston, Texas	Aug-Sep 2006	1200	150	Osthoff et al. (2008)
Houston, Texas	Sep 2013	140	18	Faxon et al. (2015)
Hong Kong, China	Aug, 2012	1900	200	Tham et al. (2014)
Hong Kong, China	Nov-Dec, 2013	4700	410	Wang et al. (2016)

^a Observed and modeled values are maxima for the reporting period. Model maxima are based on hourly values sampled at the same location and time period as the observations. The sites are listed in order of decreasing latitude.



1

Figure 1: Global budget and cycling of tropospheric inorganic chlorine (Cl_y) in GEOS-Chem. The figure shows global annual mean rates (Gg Cl a⁻¹), masses (Gg), and mixing ratios (ppt, in brackets) for simulation year 2016. Read 2.5(4) as 2.5×10⁴. ClO* stands for ClO + OClO + ClO₂ + 2×Cl₂O₂; 84% is present as ClO. Reactions producing Cl atoms and related to Cl heterogeneous chemistry are shown in red and green, respectively. The dotted box indicates the Cl_y family, and arrows into and out of that box represent general sources and sinks of Cl_y. Reactions with rate < 100 Gg Cl a⁻¹ are not shown. Anthropogenic emissions of HCl as indicated by a dashed line are only included in a sensitivity simulation (see Section 2.2

8 and 4.2 for details).



Surface and zonal mean HCI mixing ratios and CI atom concentrations



Figure 2: Global distributions of annual mean HCl mixing ratios and Cl atom concentrations in GEOS-Chem. The top panels show surface air mixing ratios/concentrations. The bottom panels show zonal mean mixing ratios/concentrations as a function of latitude and altitude. Dashed lines indicate the tropopause.





marine air. Stratospheric conditions are excluded.





Figure 4: HCl and HNO₃ surface air mixing ratios at coastal/island sites and from ocean cruises, arranged from left to right in order of increasing latitude. Observations are means (black circles) or medians (black triangles) depending on availability from the publications. Model values are monthly means for the sampling locations. Also shown are results from a sensitivity simulation with no mobilization of Cl⁻ from sea salt aerosol (SSA). References: (1, 2, 5, 10) Keene et al. (2009); (3) Sanhueza and Garaboto (2002); (4) Sander et al. (2013); (6) Dasgupta et al. (2007); (7) Crisp et al. (2014); (8) Bari et al. (2003); (9) Keene et al. (2007).

8

9





Figure 5: Aerosol Cl⁻ concentrations in surface air over the contiguous US. Values are annual means for 2016. GEOS-Chem model values are shown as contours separately for fine Cl⁻ (<1 µm diameter) and total Cl⁻. Observations from the IMPROVE

5 network (<2.5 μm diameter) are shown as circles and are the same in both panels; one would expect them to be higher than 6 the model fine Cl⁻ but lower than total Cl⁻.





Figure 6: Vertical profiles of HCl, nighttime ClNO₂, HOCl, and Cl₂ mixing ratios during the WINTER campaign over the eastern US and offshore in February-March 2015. Observations from Haskins et al. (2018) are shown as individual 1-minute data points, with medians and 25th-75th percentiles in 500-m vertical bins. Measurements below the detection limit are treated as the median of 0 and detection limit. ClNO₂ data exclude daytime (10:00-16:00 local) when mixing ratios are near zero both in the observations and in the model (Figure 7). Model values are shown as medians sampled along the flight tracks. Also shown are results from a sensitivity simulation including the anthropogenic chlorine inventory of McCulloch et al. (1999).



Diurnal variations of chlorine species during WINTER

1

Figure 7: Median diurnal variations of HCl, ClNO₂, and Cl₂ mixing ratios below 1 km altitude during the WINTER aircraft campaign over the eastern US and offshore in February-March 2015. The data are separated between ocean (left panels) and land (right panels). Model values are compared to observations from Haskins et al. (2018). Vertical bars show the 25th-75th percentiles in the observations. Measurements below the detection limit are treated as the median of 0 and detection limit. Also shown are results from a sensitivity simulation excluding ClNO₂+Cl⁻, which has negligible effect on HCl, ClNO₂, and HOCl, but brings the Cl₂ simulation in much better agreement with observations at night.





Figure 8: Vertical profiles of HCl mixing ratios during the SEAC⁴RS aircraft campaign over the Southeast US (95°-81.5°W, 30.5°-39°N) in August-September 2013, during the KORUS-AQ aircraft campaign over and around the Korean peninsula (120°-132°E, 32°-38°N) in May-June 2015, and during the INTEX-B aircraft campaign over North Pacific (175°E-50°W, 40°-61°N) in May 2006. Observations from the Georgia Tech CIMS instrument are shown as gray points (1-minute averages), with medians and 25th-75th percentiles in 1-km vertical bins. Model values are sampled along the flight tracks and for the measurement period. Measurements below the detection limit are treated as the median of 0 and detection limit. Note the difference in scales between panels.



Chlorine driven changes in BrO, NO_x, OH, and ozone

2

Figure 9: Effects of tropospheric chlorine chemistry on BrO, NO_x, OH, and ozone concentrations. The top panels show the annual mean surface concentrations of BrO, NO_x, OH, and ozone simulated in our standard model including tropospheric chlorine chemistry. The lower panels show the changes in annual mean mixing ratios/concentrations due to tropospheric chlorine chemistry, as determined by difference with a sensitivity simulation including no Cl_y production and cycling. The middle panels show the changes in surface air concentrations and the bottom panels show the changes in zonal mean mixing concentrations as a function of latitude and altitude. Black dashed lines indicate the tropopause. Numbers in bottom panels show the global tropospheric mean differences.

- 10
- 11
- 12