

## ***Interactive comment on “The unintended consequence of SO<sub>2</sub> and NO<sub>2</sub> regulations over China: increase of ammonia levels and impact on PM<sub>2.5</sub> concentrations” by Mathieu Lachatre et al.***

**Mathieu Lachatre et al.**

mathieu.lachatre@lisa.u-pec.fr

Received and published: 3 April 2019

We wish to thank the referee for his/her helpful comments. The comments of the referee are in bold and our answers in normal black.

Similar topic and conclusions have been shown in at least two recent studies (Fu et al., 2017, Liu et al., 2018). It is important to highlight the difference and new insights in the present work.

Fu et al., (2017), Liu et al., (2018) were already mentioned in manuscript initial version and more details about new insights from our study have been added during revision.

C1

The present study brings new insights concerning NO<sub>x</sub> emissions reduction impact on ammonia, described P12L9. In the study Liu et al., (2018) do not process to a SO<sub>2</sub> emissions changes only simulation, which has shown in our study a large increase of nitrate production and helped us to figure out that change of SO<sub>2</sub> and NO<sub>x</sub> emissions combine have produce more NH<sub>3</sub> released in the gas phase than SO<sub>2</sub> emissions changes alone. Fu et al., 2017 conclusion are considered and compared P15L29-L30. If our results agree with those presented in Fu et al., 2017, it brings a more precise view of Inorganic PM system with the insight brought by the cation / anion ratio and altitude analysis. This work on PM helped us to understand nitrates conservation (mentioned in Liu et al., 2018 from ground measures) between 2011 and 2015. Also, we have used information from IASI instrument to evaluate modelled NH<sub>3</sub> evolution.

We added a sentence P3L5: “A very recent study by Liu et al. (2018) suggests that ammonia increase mainly comes from SO<sub>2</sub> emission policies. They found that the changes in NO<sub>x</sub> emissions decreased the NH<sub>3</sub> column concentrations in their study period. On the contrary, Fu et al. (2017) have shown that SO<sub>2</sub> and NO<sub>2</sub> emissions control was an important factor affecting the significant enhancement of NH<sub>3</sub> column concentrations over China during the period 2011–2014. In addition, our study also presents a comparison to NH<sub>3</sub> IASI satellite observations.”

We added a sentence P12L9: “This statement on NO<sub>x</sub> emission evolution impacts is different from that in Liu et al. (2018), in which NO<sub>x</sub> emission reduction is considered as not responsible for the NH<sub>3</sub> increase between 2011 and 2015.” This additional NO<sub>x</sub> emission dependence is an important and original point of our study.

We modified a sentence P19L12 “Liu et al. (2018) estimated a +35% NH<sub>3</sub> columns increase over the North China Plain, between 2011 and 2015, taking account of SO<sub>2</sub> emissions decrease, a value close to our result for this case (+27% between 2011A and 2015B).”

We added a sentence P15L29-L30 “In the future, emissions reductions for NH<sub>3</sub> and

C2

anions precursors should lead to less  $\text{NH}_4\text{NO}_3(\text{p})$  and  $(\text{NH}_4)_2\text{SO}_4(\text{p})$  formation, reducing observed PM levels, which was already suggested in Fu et al. (2017).”

In addition, the paper requires extensive English editing. English editing has been performed with the help of a native English speaking colleague.

Specific comments:

1. Page 4, Line 6: Meteorology predictions need to be validated before exploring its impacts on  $\text{NH}_3$  concentrations. In this case we use a meteorological fields provided by the Integrated Forecasting System of ECMWF which is an operational product extensively validated by the center (Owens and Hewson, 2018). As an example, ECMWF Zenith Tropospheric Delay (ZTD) has been evaluated from GPS ZTD (Chen et al., 2010) the bias ranged from 11.5 to -28.6 mm with a corresponding average of -10.5 mm. Jingjing et al., 2015 evaluated Planetary Boundary Layer Height with CALYPSO. Moreover this product is based on meteorological analysis, which means that observations (in situ, satellite) are used to correct the initial state of the model every 6 hours which is, for temperature, humidity, a guarantee of the good quality of the fields.

2. It's better to put the model validation part (section 3.4 and 3.5) to the first part of section 3, because it's the foundation of the following analysis. In our study, we assume that our main result on ammonia increase (section 3.5) should be kept as the final part of our paper, just before the conclusion, as the IASI/CHIMERE comparison and evaluation. It is the final point of our paper, which validate the consistency of hypothesis made on emissions and meteorological changes, investigated separately in section 3.1 and 3.2.

Validation of  $\text{SO}_2$  and  $\text{NO}_x$  predictions need to be added.  $\text{SO}_2$  and  $\text{NO}_x$  columns predictions from emissions update have indeed been compared to the OMI satellite evolution in Part 2.2. It should be recalled that our emission estimations for  $\text{SO}_2$  and  $\text{NO}_x$  have also been compared and are consistent with the new MEIC inventory. (See below, answer to 4.)

C3

We added a sentence (P7L5): ““Emission update allowed to reproduce correctly  $\text{SO}_2$  and  $\text{NO}_2$  column evolutions, with for  $\text{SO}_2$  -44% (CHIMERE) and -53% (OMI) between 2011 and 2015, and for  $\text{NO}_2$  -31% (CHIMERE) and -23% (OMI) between 2013 and 2015.”

3. Page 3, Line 6-7: Why the operationally provided IASI level 2 data cannot be used to analyze the inter-annual  $\text{NH}_3$  variability? This is fully explain in Van Damme et al., 2017 : “The analysis of ANNI- $\text{NH}_3$ -v2.1 time series revealed several sharp discontinuities which seemed to coincide with IASI L2 version changes (see Fig. 3). In particular, a noticeable overall increase in the  $\text{NH}_3$  columns was found to correspond with the change from v5 to v6, and a smaller decrease was observed with the introduction of v6.2. As we will show below, these are a direct consequence of algorithmic changes to the retrieved temperature of the surface and lower troposphere. Following these findings, the need arose for a self-consistent IASI  $\text{NH}_3$  dataset, which uses stable and uniform input data. The ECMWF ERA-Interim reanalysis (Dee et al., 2011) is very suitable for this purpose, as it provides all the necessary meteorological parameters and covers the whole IASI time period.”

We added the following sentences in the text (P4L7): “For this study we used the dataset ANNI- $\text{NH}_3$ -v2.2R-I, relying on ERA-Interim ECMWF (European Centre for Medium-Range Weather Forecasts) meteorological input data rather than the operationally provided Eumetsat IASI Level 2 (L2) data used for the standard near-real-time version. The analysis of ANNI- $\text{NH}_3$ -v2.1 time series indeed revealed sharp discontinuities coinciding with IASI L2 version changes (Van Damme et al., 2017). With the ECMWF ERA-Interim reanalysis, the time series is now coherent in time (excepted for the cloud coverage flag) and can therefore be used to study interannual  $\text{NH}_3(\text{g})$  variability over East China between 2011 and 2015 (Figure 1).”

4. Page 4: In the EDGAR-HTAP-v2.2 inventory you used for 2010, Chinese emissions are derived from the MEIC inventory. The MEIC inventories for 2012, 2014 and 2016 are available in its website (<http://www.meicmodel.org/>). Why not use the MEIC inven-

C4

tory directly for 2013 and 2015? What is difference between your updated emissions for 2013 and 2015 and those in MEIC? We have initiated our work on ammonia since mid 2016, and at this time, updated emissions inventories were not available..

We now compare latest MEIC inventory (Zheng et al., 2018, Figure 1) to our updated emissions, which brings similar evolution of the emissions between 2011 and 2015. We added the following sentences: P6L12: "A recent study from Zheng et al. (2018) evaluated NOX emissions evolution of -17.4% between 2011 and 2015, similar to our -24% evolution."

P7L4: "Zheng et al. (2018) evaluated SO2 emissions evolution of -41.9% between 2011 and 2015, again similar to our -37.5% evolution."

Minor comments:

Page 1, line 4: The full name for "IASI" need to be given. Included Page 2, line 5: "NH3(g) Chinese emissions " should be "NH3(g) emissions in China" Modified

Page 2, line 23: "observed" should be deleted Modified

Page 2, line 25: "ran" should be "conducted" Modified

Page 10, line 2: "reaction" should be deleted. Modified

Page 12, line 16 to Page 13, line 2: The English grammar for the last sentence need to be checked. Sentence has been reformulated P13L9:

"However, for conditions of weak atmospheric dispersion or high humidity, as in the Sichuan province and the Chongqing municipality (located in an orographic depression), sulphates can be formed formed closer to sources. In this area, sulphates largely contribute to the PM column, as much as 32% as compared to 23% over East China, SO2-4(p), see Figure S8 in supplement file"

Page 16, line 4-7: It's difficult to understand these sentences, and the statement need to be improved. Sentence has been reformulated P15L20: "A probable explanation is

C5

the following: first July and August correspond to the monsoon season, with higher water vapour content and solar radiation over the study area. This leads to enhanced OH radical concentrations (up to twice the annual mean) to form H2SO4(g) and HNO3(g). Second, higher water content inducts more SO2(g) dissolution in aqueous phase. Both factors, then induce more SO42-(p) formation (Stockwell and Calvert,2016) decreasing by this way the C/A ratio."

Page 18: It's difficult to read Table 3. Better presentation and interpretation are needed. Table has remained identical but we tried to be more explicit in Table caption to help reader to quickly understand what "Changes" are indicating in Table 3 P18. "Table 3. Daily PM2.5 comparison between model and measurements for 2013C and 2015C simulations. "Changes" corresponds to differences between 2013C and 2015C comparisons on one hand and 2013A and 2015A ones on the other (i.e. BiasChanges = BiasC - BiasA). Bias and NRMSE are normalized using the measurement mean. R corresponds to the Pearson correlation coefficient and n represents the number of available daily means."

References:


Chen, Q., Song, S., Stefan, H., Yuei-An, L., Zhu, W., and Jingyang, Z.: Assessment of ZTD derived from ECMWF/NCEP data with GPS ZTD over China, GPS Solut, <https://doi.org/10.1007/s10291-010-020>, 2010. Fu, X., Wang, S., Xing, J., Zhang, X., Wang, T., and Hao, J.: Increasing Ammonia Concentrations Reduce the Effectiveness of Particle Pollution Control Achieved via SO2 and NOX Emissions Reduction in East China, Environmental Science and Technology Letters, 4, 221–227, <https://doi.org/10.1021/acs.estlett.7b00143>, <https://doi.org/10.1021/acs.estlett.7b00143>, 2017. Jingjing, L., Jianping, H., Bin, C., Tian, Z., Hongru, Y., Hongchun, J., Zhongwei, H., and Beidou, Z.: Comparisons of PBL heights derived from CALIPSO and ECMWF reanalysis data over China, Journal of Quantitative Spectroscopy and Radiative Transfer, 153, 102 – 112, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2014.10.011>, <http://www.sciencedirect.com/s>

C6

topical issue on optical particle characterization and remote sensing of the atmosphere: Part II, 2015. Liu, M., Huang, X., Song, Y., Xu, T., Wang, S., Wu, Z., Hu, M., Zhang, L., Zhang, Q., Pan, Y., and Zhu, T.: Rapid SO<sub>2</sub> emission reductions significantly increase tropospheric ammonia concentrations over the North China Plain, *Atmospheric Chemistry and Physics Discussions*, 2018, 1–19, <https://doi.org/10.5194/acp-2018-880>, <https://www.atmos-chem-phys-discuss.net/acp-2018-880/>, 2018. Owens, R. G. and Hewson, T.: ECMWF Forecast User Guide, Reading, <https://doi.org/10.21957/m1cs7h>, <https://software.ecmwf.int/wiki/display/FUG/Forecast+User+Guide>, <p> Replaces previous editions that were available as PDF documents.</p>, 2018. Van Damme, M., Whitburn, S., Clarisse, L., Clerbaux, C., Hurtmans, D., and Coheur, P.-F.: Version 2 of the IASI NH<sub>3</sub> neural network retrieval algorithm: near-real-time and reanalysed datasets, *Atmos. Meas. Tech.*, 10, 4905-4914, <https://doi.org/10.5194/amt-10-4905-2017>, 2017. Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L., Zhang, Y., Zhao, H., Zheng, Y., He, K., and 35 Zhang, Q.: Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions, *Atmospheric Chemistry and Physics*, 18, 14 095–14 111, <https://doi.org/10.5194/acp-18-14095-2018>, <https://www.atmos-chem-phys.net/18/14095/2018/>, 2018.

Interactive comment on *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2018-1092>, 2018.

C7



**Fig. 1.** Emissions evolution in China, from 2010 to 2017 (in Tg.yr<sup>-1</sup>) from Zheng et al, 2018

C8