Atmos. Chem. Phys. Discuss., https://doi.org/10.5194/acp-2018-111-SC1, 2018 © Author(s) 2018. This work is distributed under the Creative Commons Attribution 4.0 License.





Interactive comment

## Interactive comment on "Modelling black carbon absorption of solar radiation: combining external and internal mixing assumptions" by Gabriele Curci et al.

## C. He

cenlinhe@ucar.edu

Received and published: 31 May 2018

This manuscript is generally well written, but there seems to be one important piece missing, which is related to BC particle structures. This study assumed spheres for externally mixed BC and core-shell coating structure for internally mixed BC. However, recent observations (e.g., China et al., 2015; Wang et al., 2017) have shown irregular fractal aggregates for externally mixed BC and various non-core-shell coating structures for internally mixed BC. Further modeling studies (e.g., Scarnato et al., 2013; He et al., 2015, 2016) have indicated a large variation in BC optical properties due to the observed complex fractal and coating structures. Thus, the assumptions in the current



Discussion paper



manuscript may lead to uncertainty in the calculations of BC optical properties. Thus, I suggest that the authors include these recent studies and add some discussions on this important issue.

References

China, S., et al.: Morphology and mixing state of aged soot particles at a remote marine free troposphere site: Implications for optical properties, Geophys. Res. Lett., 42, 1243–1250, doi:10.1002/2014gl062404, 2015.

He, C., et al.: Variation of the radiative properties during black carbon aging: theoretical and experimental intercomparison, Atmos. Chem. Phys., 15, 11967-11980, doi:10.5194/acp-15-11967-2015, 2015.

He, C., et al.: Intercomparison of the GOS approach, superposition T-matrix method, and laboratory measurements for black carbon optical properties during aging, J. Quant. Spectrosc. Radiat. Transf., 184, 287–296, doi:10.1016/j.jqsrt.2016.08.004, 2016.

Scarnato, B. V., et al.: Effects of internal mixing and aggregate morphology on optical properties of black carbon using a discrete dipole approximation model, Atmos. Chem. Phys., 13, 5089–5101, doi:10.5194/acp-13-5089-2013, 2013.

Wang, Y., et al.: Fractal dimensions and mixing structures of soot particles during atmospheric processing, Environ. Sci. Technol. Lett., 4, 487-493, doi:10.1021/acs.estlett.7b00418, 2017.

**ACPD** 

Interactive comment

Printer-friendly version

Discussion paper



Interactive comment on Atmos. Chem. Phys. Discuss., https://doi.org/10.5194/acp-2018-111, 2018.