Solubility and Solution-phase Chemistry of Isocyanic Acid, Methyl Isocyanate, and Cyanogen Halides

James M. Roberts¹, and Yong Liu²

1. NOAA/ESRL Chemical Sciences Division, Boulder, Colorado, 80305

2. Department of Chemistry, University of Colorado, Denver, Denver, Colorado, 80217

for submission to Atmospheric Chemistry and Physics November, 2018

52 Abstract53

54 Condensed phase uptake and reaction are important atmospheric removal processes for reduced nitrogen 55 species, isocyanic acid (HNCO), methyl isocyanate (CH₃NCO) and cyanogen halides (XCN, X =Cl, Br, I), yet many 56 of the fundamental quantities that govern this chemistry have not been measured or are understudied. Solubilities 57 and first-order reaction rates of these species were measured for a variety of solutions using a bubble flow reactor 58 method with total reactive nitrogen (N_r) detection. The aqueous solubility of HNCO was measured as a function of 59 pH, and exhibited the classic behavior of a weak acid, with an intrinsic Henry's law solubility of 20 (±2) M/atm, and 60 a K_a of 2.0 (± 0.28) $\times 10^{-4}$ M (which corresponds to pK_a = 3.7 ± 0.06) at 298K. The temperature dependence of HNCO 61 solubility was very similar to other small nitrogen-containing compounds and the dependence on salt concentration 62 exhibited the "salting out" phenomenon that was also similar to small polar molecules. The rate constant of reaction 63 of HNCO with 0.45 M NH₄⁺ was measured at pH=3, and found to be 1.2 (\pm 0.1) \times 10⁻³ M⁻¹sec⁻¹, which is much faster 64 than the rate that would be estimated from rate measurements at much higher pHs, and the assumption that the 65 mechanism is solely by reaction of the un-dissociated acid with NH₃. The solubilities of HNCO in the non-polar 66 solvents n-octanol (n-C₈H₁₇OH) and tridecane (C₁₃H₂₈) were found to be higher than aqueous solution for n-octanol 67 $(87 \pm 9 \text{ M/atm at } 298\text{K})$ and much lower than aqueous solution for tridecane $(1.7 \pm 0.17 \text{ M/atm at } 298\text{K})$, but the 68 first-order loss rate of HNCO in n-octanol was determined to be relatively slow 5.7 (± 1.4) ×10⁻⁵sec⁻¹. The aqueous 69 solubility of CH₃NCO was measured at several pHs and found to be 1.3 (±0.13) M/atm independent of pH, and 70 CH₃NCO solubility in n-octanol was also determined at several temperatures and ranged from 4.0 (±0.5) to 2.8 71 (± 0.3) M/atm. The aqueous hydrolysis of CH₃NCO was observed to be slightly acid-catalyzed, in agreement with 72 literature values, and reactions with n-octanol ranged from 2.5 (± 0.5) to 5.3 (± 0.7) $\times 10^{-3}$ sec⁻¹ from 298 to 310K. 73 The aqueous solubilities of XCN, determined at room temperature and neutral pH, were found to increase with 74 halogen atom polarizability from 1.4 (±0.2) M/atm for CICN, 8.2 (±0.8) M/atm for BrCN, to 270 (±54) M/atm for 75 ICN. Hydrolysis rates, where measurable, were in agreement with literature values. The atmospheric loss rates of 76 HNCO, CH₃NCO, and XCN due to heterogeneous processes are estimated from solubilities and reaction rates. 77 Lifetimes of HNCO range from about 1 day against deposition to neutral pH surfaces in the boundary layer, but 78 otherwise can be as long as several months in the mid troposphere. The loss of CH₃NCO due to aqueous phase 79 processes is estimated to be slower than, or comparable to, the lifetime against OH reaction (3 months). The loss of 80 XCNs due to aqueous uptake are estimated to range from quite slow, lifetime of 2-6 months or more for ClCN, 1 81 week to 6 months for BrCN, to 1 to 10 days for ICN. These characteristic times are shorter than photolysis lifetimes 82 for CICN, and BrCN, implying that heterogeneous chemistry will be the controlling factor in their atmospheric 83 removal. In contrast, the photolysis of ICN is estimated to be faster than heterogeneous loss for average mid-latitude 84 conditions.

- 86 I. Introduction
- 87

88 The earth's atmosphere is a highly oxidizing environment in which chemical compounds are typically 89 destroyed through radical pathways. The reduced nitrogen species, isocyanic acid (HNCO) and hydrogen cyanide 90 (HCN), are an exception to this, as they have slow reactions with atmospheric radicals and have primarily 91 condensed-phase sources and sinks. Cyanogen halides (XCN, where X = CI, Br, I) are compounds that are present in 92 the environment, and whose atmospheric chemistry is of emerging interest. XCN compounds likewise have very 93 slow reaction rates with radical species and, with the exception of ICN, very slow photolysis rates in the 94 troposphere. These general classes of reduced nitrogen species, isocyanates (R-NCO), cvanides (RCN), and 95 cyanogen halides (XCN) have potential health impacts that are related to their condensed phase chemistry. 96 Therefore, information on solubility and reaction rates are needed to understand the atmospheric fate of such 97 compounds and define their impact on human and ecosystem health. Five reduced nitrogen species will be focused 98 on here: isocyanic acid, HNCO, methyl isocyanate, CH₃NCO, cyanogen chloride, ClCN, cyanogen bromide, BrCN, 99 and cyanogen iodide, ICN. 100 The isocyanate compounds are products of the pyrolysis or combustion of N-containing materials (biomass, 101 polyurethanes) (Blomqvist et al., 2003; Koss et al., 2018) and the two simplest ones, HNCO and CH₃NCO, have 102 also been observed in interstellar and cometary media (Goesmann et al., 2015; Halfen et al., 2015). The atmospheric 103 chemistry of HNCO has received considerable attention in the past few years as it has become clear that it is present 104 in ambient air, and could be related to health impacts through specific biochemical pathways (Roberts et al., 2011) 105 involving the reaction of cyanate ion with proteins. There are relatively few observations of HNCO in ambient air, 106 and those show that there are "background" mixing ratios that range from 10pptv to over several ppbv depending on 107 the nature of regional sources, and that peak mixing ratios approaching a few ppbv are observed in areas impacted 108 by local biomass burning (Chandra and Sinha, 2016; Kumar et al., 2018; Mattila et al., 2018; Roberts et al., 2014; 109 Sarkar et al., 2016; Wentzell et al., 2013; Woodward-Massey et al., 2014; Zhao et al., 2014). The aqueous phase 110 solubility of HNCO was examined by Roberts et al., (Roberts et al., 2011) and Borduas et al., (2016), wherein it was 111 found that HNCO is only slightly soluble at pHs characteristic of atmospheric aerosol (pH= 2-4) and as a weak acid 112 (pKa=3.9), it is quite soluble at physiologic conditions (pH=7.4). Attempts to model the global distribution of 113 HNCO (Young et al., 2012) and the cloud water uptake of HNCO (Barth et al., 2013) used the limited solubility and 114 hydrolysis data available at that time, (Jensen, 1958; Roberts et al., 2011). Several aspects of HNCO solubility 115 remain unknown, such salt effects on aqueous solubility, and solubility in non-aqueous solvents, a property 116 important for predicting HNCO behavior in biological systems. The pH dependent hydrolysis of HNCO had been 117 studied some time ago (Jensen, 1958), the mechanism for this process involves three separate reactions; 118 $HNCO + H_3O^+ \implies NH_4^+ + CO_2$ (R1) 119 $HNCO + H_2O \implies NH_3 + CO_2$ (R2) 120 $NCO^{-} + 2H_2O \implies NH_3 + HCO_3^{-}$ (R3) 121 and Borduas et al. (2016), recently re-measured these rates under a wider range of conditions and found their

122 measurements to be essentially consistent with the previous work at pHs of interest in the atmosphere. Rates of

reaction of HNCO with other compounds in aqueous solution are not as well studied, especially under atmospheric conditions, e.g. low pH, relatively high ionic strength. Rates of reaction of HNCO/NCO⁻ with nitrogen bases have been measured but only at the pK_as of the BH⁺, which are typically pH 9-10 (Jensen, 1959; Williams and Jencks,

126 1974a, b).

127 Methyl isocyanate is most notable for its part in the one of the largest industrial disasters in history, when a 128 large quantity of CH₃NCO was released from a chemical plant and fumigated the city of Bhopal, India. There are 129 other, more common sources of CH₃NCO to the atmosphere including combustion of biomass (Koss et al., 2018) 130 and N-containing polymers such as polyurethanes and isocyanate foams (Bengtstrom et al., 2016; Garrido et al., 131 2017), and cooking (Reves-Villegas et al., 2018). Recent measurements of CH₃NCO in laboratory wildfire studies 132 have observed mixing ratios up to 10 ppby or so in fuels characteristic of western North America (Koss et al., 2018). 133 CH₃NCO is also produced in photochemical reactions of methylisothiocyanate (CH₃NCS), which is the main 134 degradation product of the agricultural fungicide metam-sodium (CH₃NHCS₂Na) (Geddes et al., 1995). In addition, 135 CH₃NCO has been observed in studies of the photooxidation of amides (Barnes et al., 2010; Borduas et al., 2015; 136 Bunkan et al., 2015) and by extension will be formed in dimethyl amine oxidation. To our knowledge there is only 137 one reported set of ambient measurements of CH₃NCO, conducted near a field where metam-sodium was being used 138 as a soil fumigant (Woodrow et al., 2014), and the resulting CH₃NCO mixing ratios were as high as 1.7 ppby. The 139 California Office of Environmental Health Hazard Assessment has placed an inhalation reference exposure level of 140 0.5 ppby (1 µg/m³) on CH₃NCO due to its propensity to cause respiratory health effects (State of California, 2008). 141 There have been only a few studies of the gas phase loss rates of CH₃NCO including reaction with OH 142 radical (Lu et al., 2014), which appears to be slow based on the mostly recent measurements (Papanastasiou et al., in 143 preparation, 2018), reaction with chlorine atoms (Cl) which might be as much as 20% of OH under some 144 atmospheric conditions (Papanastasiou et al., in preparation, 2018), and UV photolysis which has a negligible 145 contribution to atmospheric loss (Papanastasiou et al., in preparation, 2018). Thus, heterogeneous uptake might 146 compete with these gas phase loss processes. The solubility of CH₃NCO has not been previously determined 147 experimentally, but is probably low, <2 M/atm, by analogy to CH₃NCS (3.7 M/atm) (Geddes et al., 1995). In 148 addition, there are no data on the solubility of CH₃NCO in non-aqueous solvents. The hydrolysis of CH₃NCO is acid 149 catalyzed; 150 $CH_3NCO + H_2O + H^+ => CH_3NH_3^+ + CO_2$ (R4)

151 $CH_3NCO + H_2O => CH_3NH_2 + CO_2$

producing methyl amine and carbon dioxide. The rate constants for these reactions are fairly well established (Al-Rawi and Williams, 1977; Castro et al., 1985).

(R5)

Cyanogen halides are less well studied as atmospheric species, but have potentially important environmental sources. Cyanogen chloride was once produced as a chemical warfare agent, however its importance to the atmosphere is more related to its possible formation in the reaction of active chlorine species (HOCl/OCl⁻, chloramines) with N-containing substrates such as amino acids and humic substances (Na and Olson, 2006; Shang et al., 2000; Yang and Shang, 2004). These reactions are known to be important in systems where chlorination is used for disinfection such as swimming pools and water treatment (see for example (Afifi and Blatchley III, 2015), and

- 160 perhaps indoor surfaces (J. Abbatt, personal communication). We are not aware of any measurements of CICN in
- ambient air. Cyanogen bromide can likewise be formed through reactions of HOBr/OBr- with reduced nitrogen
- 162 species, and there are observations of BrCN in bromide-containing waters that have been received chlorine
- 163 treatment (see for example (Heller-Grossman et al., 1999). The formation results from the facile reaction of
- 164 HOCl/OCl- with bromide to make HOBr/OBr-, which then reacts with nitrogen species in the water. In addition,
- there is a natural source of BrCN from at least one strain of marine algae (Vanelslander et al., 2012) that is thought
- 166 to be related to allelopathic activity, i.e. secreted to control the growth of competing organisms. This marine algae
- source may be responsible for BrCN levels observed in remote atmospheres (J.A. Neuman and P.R. Veres, personal
- 168 communication). Cyanogen iodide can also potentially be formed from the chlorination of water or wastewater
- because iodide is easily oxidized by HOCl/OCl⁻, however iodide is usually quite small in concentration, so the
- 170 several studies that report total cyanogen halides report CICN and BrCN but not ICN (Diehl et al., 2000; Yang and
- 171 Shang, 2004). There are also biochemical pathways for ICN formation involving several enzymes that are part of the
- immune defense system (see for example (Schlorke et al., 2016)), but the extent to which ICN might be volatilized
- 173 from those systems is not clear. There are some observations of ICN in the remote marine troposphere (J.A. Neuman
- 174 and P.R. Veres, personal communication), but their origin is currently unclear.

The possible gas phase loss processes of cyanogen halides include reaction with radicals or ozone, and photolysis. Radical reaction rates (OH, Cl) have not been measured at room temperatures, but are likely to be slow due the electronegativity of each group (X-CN). The UV-visible absorption spectra of all three of these compounds have been measured (Barts and Halpern, 1989; Felps et al., 1991; Hess and Leone, 1987; Russell et al., 1987), and indicate a range of photolysis behavior ranging from no tropospheric photolysis of ClCN, to slight photolysis of BrCN, and faster photolysis of ICN. The rates of photolysis need to be balanced against condensed phase losses of XCN compounds to obtain a full picture of their atmospheric losses.

182The aqueous phase solution chemistry of cyanogen halides is not as well studied as the isocyanates. The183aqueous solubilities of XCN compounds are not known with the exception of ClCN whose solubility is thought to be184fairly low, 0.6 - 0.52 M/atm at 293-298K (Weng et al., 2011; Yaws and Yang, 1992) as reported by (Hilal et al.,1852008). The hydrolysis of XCN compounds are known to be base-catalyzed and so involve the following reactions;

(R7)

186

187 $XCN + 2H_2O => HOCN + H_3O^+ + X^-$ (R6)

188
$$XCN + H_2O + OH^- => HOCN + X^- + H_2O$$

189

with R6 being fairly slow at medium to low pH (Bailey and Bishop, 1973; Gerritsen et al., 1993). The product,
cyanic acid, HOCN, is unstable with respect to HNCO in aqueous solution (Belson and Strachan, 1982);

192

$$HOCN + H^+ \Longrightarrow HNCO + H^+$$
(R8)

194

Thus, XCN compounds represent potential intermediates in the condensed-phase formation of HNCO, for whichthere is some observational evidence (Zhao et al., 2014). So, in addition to being active halogen species, XCN

- 197 compounds represent potential condensed phase source of HNCO in systems where there is halogen activation and 198 there are reduced nitrogen species present, e.g. wildfire plumes, bio-aerosols and indoor surfaces.
- 199 Measurements of solubility and reaction rates will be presented here for HNCO, CH₃NCO, and the XCN
- 200 species: CICN, BrCN, and ICN. The aqueous solubility of HNCO was measured as a function of pH, temperature
- and salt concentration. The rate of reaction of HNCO with NH4⁺ was measured at pH3, to examine the importance of
- this reaction to atmospheric uptake of HNCO. The solubilities of HNCO in the non-polar solvents n-octanol and
- tridecane were also measured as a function of temperature, and the first-order loss rate of HNCO in n-octanol was
- also determined. The aqueous solubility of CH₃NCO was measured at several at several pHs, and the solubility in n-
- 205 octanol was also determined at several temperatures. Finally, the aqueous solubility of CICN, BrCN, and ICN were
- determined at room temperature, and at 273.15 K (CICN, BrCN) and neutral pH, and the solubility and first loss of
- these compounds in n-octanol was also determined. These data will be used to estimate atmospheric lifetimes
- against aqueous uptake and to assess the relative bioavailability of these compounds.
- 209

210 II. Methods

211 Most of the techniques used for the work presented here have largely been presented elsewhere (Borduas et 212 al., 2016; Kames and Schurath, 1995; Kish et al., 2013; Roberts, 2005) and will only be briefly summarized here. 213 The basic principle is that the compound of interest is equilibrated with solution in a bubble flow reactor, and then 214 removed from the gas-phase and the exponential decay of the signal due to loss of the compound is measured with a 215 sensitive and selective method. The dependence of decay rates on flow rate-to-liquid volume ratio can then be 216 related to solubility and first-order loss rate due to reaction in solution. This technique relies on being able to 217 produce a consistent gas stream of the compound of interest, and being able to selectively detect the compound 218 exiting the reactor. This method has limitations in that the solubility must be within a certain range, and the first-219 order loss rate slow enough that there are measurable amounts of compound exiting the reactor.

220

- 221 A. Preparation of Gas-Phase Standards
- 223 The general system used for preparation of gas phase streams of HNCO, CH₃NCO, BrCN, and ICN was the 224 capillary diffusion system described by (Williams et al., 2000) and (Roberts et al., 2010). Isocyanic acid was 225 produced in a steady stream by heating the trimer, cyanuric acid (Sigma-Aldrich, USA) to 250°C under N₂ and 226 establishing a constant diffusion rate through a short length of capillary tubing (1mm ID x 5cm length). Care was 227 taken to condition the system for several days before use, by keeping the system under flow and at a minimum of 228 125°C even when not in active use, to prevent the build-up of unwanted impurities, particularly NH₃. Standards in 229 the range of several ppmv in 40 SCCM could easily be prepared in this way. 230 The same capillary diffusion cells were used for CH₃NCO preparation, starting with a sample of the pure
- liquid (Alinda Chemicals, UK). FTIR analysis of samples of this material were found to contain small amounts (3%)
 of siloxanes, which probably came from a chloro-silane added as a stabilizer, but no measurable presence of any
- 233 other nitrogen compounds. The high volatility of CH₃NCO (BP 38 °C) required that low concentration solution (1%

vol/vol) of CH₃NCO in n-tridecane (C₁₃H₂₈) solvent at a temperature of 0°C be used in the diffusion cell. Under

these conditions a 40 SCCM stream resulted in a mixing ratio of 10ppmv. The output of the source was stable for

- 236 long periods of time (days) and could be used for the solubility study and calibration of other instruments. The
- source was also analyzed by an H₃O⁺ chemical ionization mass spectrometric system, which showed that it had no
- impurities detectable above the 1% level.
- The preparation of a gas phase standard of CICN is described by Stockwell et al., (2018) and is based on chemical conversion of an HCN calibration mixture. It has been known for some time that HCN reacts readily with active chlorine compounds to yield CICN (Epstein, 1947), for example:
- $HCN + HOC1 \Longrightarrow ClCN + H_2O$

(R9)

- 243 In fact, this reaction has been used as the basis for measuring HCN in the gas phase by conversion to ClCN with 244 detection by gas chromatography with electron capture (Valentour et al., 1974). In those systems, Chloramine-T 245 (Sigma-Aldrich), a non-volatile sulfonyl N-chloro compound, has proven useful. The method used in this work 246 consisted of passing a small stream (5-10 SCCM) of a 10ppmv gas-phase standard of HCN in N_2 , combined with 247 humidified Zero Air (ZA, 80% RH, 30-50 SCCM) over a bed packed with glass beads coated with a solution of 248 Chloramine-T. The glass beads were prepared by coating glass 3 mm OD beads with a 2 g/100mL solution and 249 packing ~20cc of them in a 12.7mm OD PFA tube and flowing ZA over them until they appeared dry. The reaction 250 was shown to be essentially 100% ($\pm 10\%$) when conducted in a humidified atmosphere (RH $\geq 60\%$), PTR-MS, and 251 FTIR analysis of the gas stream before and after passing through the chlorination bed. The ClCN source was also 252 checked by measuring the total nitrogen content of the gas stream before and after the chlorination step, and the 253 resulting signal was found to be $98\pm1\%$ of the original HCN standard. This means that the combination of the 254 chlorination reaction and Nr conversion (see below) were at least 98% efficient. 255 Preparation of BrCN and ICN gas streams was accomplished with the diffusion cell apparatus using
- commercially available samples of BrCN (98% purity, Sigma-Aldrich) and ICN (97% purity ACROS Organics),
 that were used without further purification. BrCN is a volatile solid, so was kept in a diffusion cell at 0°C while in
- use. ICN is a relatively non-volatile solid and so was placed in a diffusion cell and heated to 80°C while in use.
- These resulted in sample streams that had mixing ratios on the order of 250-350 ppbv in 1 SLPM. Analysis by
- 260 iodide ion chemical ionization mass spectrometry indicated traces of the molecular halogen species (Br₂, I₂), but no
- 261 other significant N-containing species.
- 262
- 263 B. Detection of Nitrogen Compounds
- 264

The method for detection of the compounds studied in this work relies on high temperature conversion of any N-containing species, except for N_2 or N_2O , to nitric oxide (NO) and detection of the resulting NO by O₃ chemiluminescence (Williams et al., 1998). This technique, which we will refer to as Total Reactive Nitrogen, N_r, has been shown to measure a wide range of reduced nitrogen species as well as the more familiar oxides of nitrogen (Hardy and Knarr, 1982; Saylor et al., 2010; Stockwell et al., 2018), provided care is taken to convert any nitrogen distribution formed in the Pt converter hash to NO prime to detection (Calcurch et al., 2007). In this work this work

dioxide that is formed in the Pt converter back to NO prior to detection (Schwab et al., 2007). In this work, this was

- accomplished with a solid molybdenum tube operated at between 350 and 450 °C, with the addition of a small
- amount of pure H₂ resulting in a 0.8% mixing ratio in the catalyst flow. The detection system was routinely
- 273 calibrated with a NO standard (Scott-Marrin, Riverside, CA) and the conversion efficiency was confirmed with a
- low concentration (10ppmv) HCN standard (GASCO, Oldsmar, FL). The high conversion efficiencies (≥98%) for
- HNCO and CICN were confirmed by other methods as described by Stockwell et al. (2018). The conversion
- efficiencies for BrCN, and ICN are assumed to be equally high due to the fact the X-CN bond strengths of these
- 277 compounds are lower than for H-CN and Cl-CN (Davis and Okabe, 1968) and the CH₃-NCO bond is weaker than
- the H-NCO bond (Woo and Liu, 1935). Although readily measured here, a solubility measurement of this kind does
- 279 not require the determination of the absolute concentration of the analytes, it only requires that the measurement be
- 280 linear (i.e. constant sensitivity) throughout the range of signals measured. The NO instrument is linear from the low
- 281 pptv into the low ppmv range, the chief limitation being the ability to count photon rates above 5MHz. The
- 282 magnitude of the gas phase sources used and the flow rate of the instrument (1 SLPM) insured that instrument
- signals did not reach the non-linear range.

284 The requirement for the detection method to be selective could be an issue with a general method such as 285 Nr. In practice, the reactions of the nitrogen species studied here form products that are not volatile under the 286 conditions used in this work, and so do not interfere with the measurement. In aqueous-phase reactions. HNCO 287 produces NH₃/NH₄⁺, CH₃NCO produces CH₃NH₂/CH₃NH₃⁺, and XCN compounds produce HOCN/NCO⁻ all of 288 which are non-volatile in the pH ranges at which those experiments were conducted. The products of the organic-289 phase reactions are not as well known: tridecane should not react with HNCO, n-octanol will form carbamyl or 290 methyl carbamyl groups with n-octyl substituents which should be non-volatile. Possible reactions of XCN 291 compounds with n-octanol are less well known, particularly in the absence of water in the solution, so those 292 experiments will need to be interpreted with care.

The bubble flow reactor has been described in a number of publications (Borduas et al., 2016; Kames and Schurath, 1995; Kish et al., 2013; Roberts, 2005), so will be only briefly summarized here. The reactor used for the most of the experiments is a modification of the one described by Roberts (2005), the main modification being a reduction in volume to 125 mL. Liquid volumes used in the experiments ranged from 20 to 50 mL, and the volumetric flow rates used ranged from 170 to 1070 ambient mL/min. Temperatures were measured using a calibrated mercury thermometer, and in temperatures different than room temperatures were controlled using a water bath with either ice/water, or a temperature control system. The uncertainties in the temperatures were ±0.5 °C.

300 The bubble flow reactor method relies on the rapid equilibration of a gas stream that contains the analyte of 301 interest, with solution by means of the creation of small, finely divided bubbles. In the system used here, these 302 bubbles are created by passing the gas stream through a fine glass frit, situated at the bottom of the glass vessel. The 303 main sample flow is passed through the bubbler and into the detector stream to establish a baseline. A small flow of 304 the analyte is added upstream of the reactor by means of a PFA solenoid valve to start the measurement and the 305 effluent is monitored until the measured concentration attains equilibrium. At this point, the analyte entering the 306 reactor is switch off, and the concentration exiting the reactor begins to decay. This decay is due to a combination of 307 loss of the analyte as it re-equilibrates with the gas stream, and first-order loss in the solution due to reaction. Under

308conditions of rapid equilibration, this decay takes the form of a single exponential equation, dependent on the ratio309of flow rate (ϕ , cm³/sec) to liquid volume (V, cm³), the Henry's Law solubility (M/atm), and the first-order loss rate310(k):311 $ln(C_0/C_t) = [\phi/(HRTV) + k]t$ 313Eq. (1)

Measurements performed at a series of ϕ/V should be linear with a slope of the decay rate (d $ln(C_0/C_t)/dt$) vs phi/V of 1/HRT, where R is the ideal gas constant, and T the temperature (K), and an x-intercept of k, the first-order loss rate (sec⁻¹). In practice, the linearity of this relationship and the performance of the measurement at different liquid volumes and flow rates that result in the same ϕ/V provide a check on the assumption of rapid equilibration within the reactor.

319 Attempts to measure the solubility of ICN with the glass bubbler system described above were 320 unsuccessful, because ICN did not equilibrate at the levels and timescales typical of the other compounds measured 321 in this work, and the decay profiles were not reproducible nor exponential. The possibility that this was due to 322 higher solubility, faster reaction, or decomposition of ICN on glass surfaces was explored by using a smaller reactor 323 fabricated from 12.7 mm O.D. PFA tubing and PFA compression fittings (see supplemental Figure S1). In these 324 experiments, liquid volumes of between 1.0 and 2.0 mL and flow rates of 100 to 600 ambient cc/min were used. 325 This resulted in equilibration and decay profiles more similar to the other experiments, when the solubility of ICN in 326 water was measured at room temperature.

327 Solution for the aqueous solubility/reaction experiments were prepared from reagent-grade materials. The 328 pH 2-4 buffer solutions were commercial preparations, made from citric acid monohydrate with differing amounts 329 of hydrochloric acid, sodium chloride, and sodium hydroxide (Fixanal, Fluka Analytical), having anion 330 concentrations ranging from 0.08 to approximately 0.2 M. The manufacturer specifications (Fluka, Sigma-Aldrich) 331 of the pH=3 buffer showed a slight temperature dependence, with the pH ranging from 3.03 at 0 °C to 2.97 at 90 °C. 332 An ammonium chloride solution of 0.45 M was prepared through addition of a measured amount of the solid to the 333 pH=3 buffer. Sodium chloride solutions ranging up to 2.5 M were prepared gravimetrically in the pH buffer 334 solution. The pHs of NH₄Cl and NaCl were measured at room temperature with a pH meter and found to be within 335 0.1 pH unit of the nominal buffer pH value.

337 III. Results and Discussion

338

336

A typical experiment consisted of a series of exposures of the solution of interest to the analyte in a gas stream at a series of known flow rates. The analyte was introduced by switching the small flow from the source into the bubbler gas stream using an all-Teflon PFA 3-way valve (Roberts, 2005). The total reactive nitrogen content of the gas exiting the bubbler was measured continuously and approached a plateau as the analyte equilibrated with the solution. At this point, the analyte was switched out of the bubbler stream, using the 3-way valve so as not to otherwise perturb the flow through the system. The Nr concentration exiting the bubbler decayed exponentially due 345 to a combination of re-equilibration and first-order reactive loss of dissolved analyte (due to hydrolysis for 346 example). An example of the data generated by these experiments is shown in Figures 1 and 2, which show the 347 exponential decays for a series of gas flow rates (Figure 1) and the correlation of the decay rates versus the ratio of 348 volumetric flow rate to solution volume (Figure 2). The uncertainties in the Henry's coefficients are derived from a 349 combination of the reproducibility of the decay rates, the agreement between decay rates at the same ϕ/V (but 350 different flows and liquid volumes) and the fits to the slope of relationships like those shown in Figure 2, and were 351 generally $\pm 10\%$ or better. The uncertainties in first-order lass rate are the corresponding uncertainties in the 352 intercepts. The results of the experiments with HNCO, CH₃NCO, ClCN, BrCN, and ICN with the variety of solvents 353 and conditions employed are summarized in Table 1 and described below. 354 355 A. Results for Aqueous Solution 356 357 1. Solubility and Reactions of HNCO 358 359 Here we report results for pHs between 2 and 4, and for the temperature range 279.5 to 310.0 K at pH=3. In 360 addition, we report data for the effect of salt concentrations on the solubility at pH=3, and the effect of ammonium 361 concentrations on solubility and apparent first-order loss rate in solution. The dependence of aqueous solubility of 362 HNCO on pH is expected given it is a weak acid; 363 HNCOg <=> HNCOaq Eq. (2) Hhnco 364 $HNCO_{aq} + H_2O_{aq} \iff H_3O^+ + NCO^-$ Ka Eq. (3) 365 so that the effective Henry's coefficient has a term that accounts for the acid-base equilibrium and that equilibrium 366 constant: 367 $H_{eff} = H_{HNCO}(1 + K_a/[H_+])$ Eq. (4) 368 The plot of H_{eff} vs 1/[H+] is shown in Figure 3, the slope of which is $H_{HNCO}*K_a$ and the intercept is the intrinsic 369 Henry's Law constant, H_{HNCO}. The resulting fit ($R^2 = 0.99$) gave a H_{HNCO} = 20 (±2) M/atm, and a K_a of 2.0 (±0.28) 370 $x10^{-4}$ M (which corresponds to pK_a = 3.7 ± 0.06). The uncertainties in these numbers were derived from the standard 371 deviations of the fitted parameters, where the value for K_a is the propagated uncertainty in both H and the slope. 372 Figure 4 shows the comparison of the H measurements from this work with those of Borduas et al., (2016) plotted 373 according to Eq. 4 equation. There are approximately 20% differences in the two data sets, which is just at the limits 374 of the quoted uncertainties, when both the uncertainties in the intrinsic H and pKa are taken into account. 375 The temperature dependence of the solubility measured at pH=3, obeys the simple Van't Hoff relationship; 376 $d\ln H_{\rm eff}/d(1/T) = -\Delta H_{\rm soln}/R$ Eq. (5) 377 shown in Figure 5 as a linear relationship of log H vs. 1/T. These data were not corrected for the slight dependence 378 of the buffer pH on temperature (3.02-2.99 pH units over this range). The slope of the correlation yields a ΔH_{soln} of 379 -37.2 ± 3 kJ/mole, calculated using dimensionless Henry's coefficients (H_{eff}RT), (Sander, 2015). This enthalpy of 380 solution agrees with that measured by Borduas et al., (2016) (-34 \pm 2 kJ/mole) within the stated uncertainties. 381 Moreover, this enthalpy is similar to those of other small N-containing molecules: HCN (-36.6 kJ/mole), CH₃CN (-

- 382 34.1 kJ/mole), and nitromethane (-33.3 kJ/mole) (Sander, 2015), but different than that of formic acid (-47.4
- 383 kJ/mole) which was used by the cloud uptake modeling study (Barth et al., 2013).
- 384 Often the Henry's Law solubility can depend on salt concentration of the solution, usually resulting in a 385 lower solubility (salting out), but occasionally resulting in a higher solubility (salting in), with higher salt 386 concentrations. These effects are most applicable to aerosol chemistry, where ionic strengths can be quite high. This 387 effect on HNCO solubility was measured at pH=3 and 298 K for NaCl solutions between 0 and 2.5 M concentration. 388 The results, shown in Figure 6, exhibit the classic "salting out" effect where HNCO was only about 60% as soluble 389 at 2.5 M compared to the standard pH=3 buffer. The Setschenow constant, k_s , can be determined by the relationship: 390 $-Log(H/H_0) = k_s x[I]$ Eq. (6) 391 where H is the Henry's coefficient at a given ionic strength, I, and H₀ is the Henry's coefficient in pure water. For a 392 salt with two singly charged ions, I is equal to the salt concentration. In this experiment, k_s was found to be 0.097 ± 393 0.011 M^{-1} . The magnitude of the salting out effect on HNCO is similar or slightly smaller than those found for other
- small organic compounds in NaCl (Clever, 1983; Schumpe, 1993).

The net hydrolysis reaction rates observed in this study are listed in Table 1, and range from 0.22 to 4.15 $\times 10^{-3}$ s⁻¹ and are both pH and temperature dependent. The main reactions of HNCO/NCO⁻ in aqueous solution are hydrolysis reactions that involve the acid or its conjugate anion, as detailed in Reactions 1-3 noted above. The expression for the net hydrolysis reaction is;

399
$$k_{hydr} = \frac{k_1 [H^+]^2 + k_2 [H^+] + k_3 K_a}{K_a + [H^+]}$$
Eq. (7)

as derived by Borduas et al., 2016. The rates of these reactions that were determined in several previous studies
(Borduas et al., 2016; Jensen, 1958) and are in reasonable agreement except for Reaction 3, which is not
atmospherically relevant. Equation 7 was used to calculate the values from those two studies that would correspond
to the rates at pH=3 measured in our work, and are also listed in Table 1. The rate constants reported in this work
agree within the range observed in the two previous studies, except for one temperature, and the relative standard
deviations of mean values calculated from all three observations ranged from 5 to 30%.

The above hydrolysis reactions represent a lower limit on the condensed phase loss of HNCO, so reaction with other species present in the condensed phase might result in faster loss, and produce unique chemical species. HNCO/NCO⁻ are known to react with a variety of organic compounds having an "active hydrogen" (Belson and Strachan, 1982) through simple addition across the N=C bond. For example, alcohols react to yield carbamates, i.e. esters of carbamic acid:

$$411 \qquad HNCO + ROH => H_2NC(O)OR \qquad (R10)$$

412 In the same fashion, HNCO/NCO⁻ can react with ammonia in solution to yield urea

- 413 $HNCO + NH_3 <= > H_2NC(O)NH_2$ (R11) 414 And in a more general sense, react with amines to yield substituted ureas:
- 415 $HNCO + RNH_2 \ll H_2NC(O)NHR$
 - 416 Reaction 11 is known to be an equilibrium that lies far to the product side under all conditions pertinent to this work
 - 417 (Hagel et al., 1971). While the forward reaction rate for R11 has been measured under neutral to slightly basic
 - 418 conditions (Jensen, 1959; Williams and Jencks, 1974b), it has not been measured at pHs applicable to atmospheric

(R12)

419 aerosol or cloud droplets, i.e. pH=2-4. These previous studies have assumed that the mechanism involves the 420 reaction of the un-ionized species, e.g. NH₃ and HNCO, although there is some evidence that Reaction 12 for some 421 amines (RNH₂) has a more complicated reaction mechanism (Williams and Jencks, 1974a). As a consequence of this 422 assumption, the previous studies reported their reaction rates corrected for the acid-base equilibria of each species. 423 The solubility/reaction experiment in this work was performed at pH=3 and $[NH_4^+]$ of 0.45 M, so a substantial 424 correction of the literature values for the acid-base equilibria in the case of NH_4^+ and a minor correction for the 425 dissociation of HNCO was required in order to compare with our result. The results of our study (Table 1) show that 426 the solubility of HNCO in NH₄Cl solution at 292 K is essentially the same as that of the pH=3 buffer alone (31.5 ± 3) 427 vs 32.6 ± 3 M/atm). This implies that R11 does not impact the aqueous solubility. However, the measured first-order 428 loss rate, 1.2 (± 0.03) ×10⁻³ sec⁻¹ is faster than the hydrolysis at pH3, 0.66 (± 0.06) ×10⁻³ sec⁻¹. The reaction can be 429 expressed as the sum of hydrolysis and reactions of HNCO and NCO⁻ with NH₄⁺ (the predominant form at pH3). $\frac{d[HNCO+NCO]}{[HNCO+NCO]} = -(k_{hydr} + k_{11}[NH_4^+])dt$ 430 Eq. 8 431 We calculate a value of 1.2 (± 0.1) $\times 10^{-3}$ M⁻¹sec⁻¹ for k₁₁ from our measurements which is much faster than the rate 432 constants reported by previous studies, 5×10^{-6} M⁻¹sec⁻¹ (Jensen, 1959) and 1.5×10^{-5} M⁻¹sec⁻¹ (Williams and Jencks, 433 1974b), when corrected for acid-base equilibria. 434 435 Solubility and Reactions of CH₃NCO 436 437 The solubility and first-order loss rate of CH₃NCO were measured at pH=2 and pH=7 at 298 K, and the 438 results are listed in Table 1. The Henry's coefficients, 1.3 (±0.13) and 1.4 (±0.14) M/atm, were lower than those 439 measured for HNCO, and independent of pH, within the uncertainties of the measurements. This is consistent with 440 CH₃NCO being a less polar compound, with no dissociation reactions that might be pH dependent. In addition, these 441 results imply that solution complexation due to the presence of anions does not affect CH₃NCO solubility, at least at 442 the concentrations and anions present in the pH=2 buffer solution, 0.2 M for the sum of citrate and chloride. 443 The first-order loss rates of CH₃NCO, presumably due to hydrolysis, did show a pH dependence that 444 implies acid catalysis. These hydrolysis rates were faster than the rates for HNCO at the same temperatures and pHs. 445 Solution-based studies of CH₃NCO hydrolysis in the presence of strong acid anions (Al-Rawi and Williams, 1977; 446 Castro et al., 1985) have shown that a complex mechanism takes place, involving a reversible complexation, (shown 447 here for HSO₄⁻): 448 449 $CH_3NCO + HSO_4^- \iff CH_3NH-C(O)-OSO_3^-$ (R13) 450 451 happening in parallel with catalyzed and un-catalyzed hydrolysis: 452 453 $CH_3NCO + H_3O^+ \Longrightarrow CH_3NH_3^+ + CO_2$ (R4) 454 $CH_3NCO + H_2O => CH_3NH_2 + CO_2$ (R5) 455

456	Rate constants for reactions R4 and R5 were reported by Castro et al., (1985) but the prec	cision of these were				
457	somewhat compromised by the presence of the R13 equilibrium. In this study, the Henry	's coefficient results imply				
458	a negligible role for complexation, so the following simplified expression for the pH dep	endence is used:				
459						
460	$k_{\text{MIC}} = k_5 + k_4 [\text{H}_3\text{O}^+]$	(Eq9)				
461						
462	to derive the following values for $k_5 = 1.9 (\pm 0.6) \times 10^{-3} \text{ sec}^{-1}$, and $k_4 = 0.13 (\pm 0.07) \text{ M}^{-1}\text{sec}^{-1}$	⁻¹ . These values are in				
463	reasonable agreement with the value for k ₅ given by Al-Rawi and Williams, (1977), 1.47 $\times 10^{-3}$ sec ⁻¹ considering					
464	those measurements were at 1M KCl, and the value for $k_4 = 0.16 \text{ M}^{-1}\text{sec}^{-1}$ given by Castro et al., (1985) for reaction					
465	with HCl in the absence of a buffer.					
466						
467	3. Solubility and Reactions of XCN Compounds.					
468						
469	The solubilities and first-order loss rates of XCN compounds were measured at	room temperature and				
470	neutral pH in pure DI water, and at ice/water temperature for ClCN and BrCN. The result	ting Henry's coefficients				
471	are listed in Table 1. The CICN solubility was essentially the same as that measured for CH ₃ NCO at room					
472	temperature, and is in reasonable agreement with the value of 0.52 M/atm at 298 K based on a model estimate (Hilal					
473	et al., 2008), and one reported measurement, 0.6 M/atm at 293 K (Weng et al., 2011). In contrast, BrCN was more					
474	soluble than ClCN, 8.2 ±0.8 M/atm at 296°K, but fairly insoluble in an absolute sense. The temperature dependences					
475	of HCICN and HBrCN were as expected, showing higher solubility at lower temperatures, however, they had very					
476	different heats of solution, -27.8 kJ/mole for ClCN, and -38.3 kJ/ mole for BrCN, although there are only two data					
477	points for each compound. Both the higher solubilities and larger ΔH_{soln} , could be a result of the higher dipole					
478	moment and polarizability of BrCN relative to CICN (Maroulis and Pouchan, 1997).					
479	The solubility of ICN was measured a room temperature using a combination of	Edifferent flow rates (208 –				
480	760 amb cm ³ /min) and liquid volumes (1.95 and 0.95mL). A plot of the decay rates versu	is ϕ/V for those runs is				
481	shown in Figure 7. Where those data sets overlap there is agreement to within about 15%	, implying that the				
482	equilibration could be fast enough to meet the criteria for these types of flow experiments. The resulting Henry's					
483	coefficient, 270 (±54) M/atm. Is significantly larger than the other two XCN compounds, but is consist with the					
484	trend of increasing solubility with dipole moment and polarizability. Attempts to use the small reactor to measure					
485	the solubility of ICN at ice/water temperatures was not successful, e.g. did not yield simple single exponential					
486	decays with time, under the same range of flow conditions as used in the room temperature experiment.					
487	The hydrolysis of XCN compounds is known to be base-catalyzed, and can be susceptible to anion					
488	complexation (Bailey and Bishop, 1973; Gerritsen et al., 1993) in a manner similar to CH	I3NCO:				
489						
490	$XCN + A^{-} < = > [XCN \cdot A]^{-}$	(R14)				
491	$XCN + 2H_2O \Longrightarrow HOCN + H_3O^+ + X^-$	(R6)				
492	$XCN + OH^- + H_2O => HOCN + X^- + H_2O$	(R7)				

493	
494	This complexation can be ignored in our study for CICN and BrCN since the experiment was performed in
495	DI water. Accordingly, the expression for the CICN and BrCN hydrolysis rate constant is;
496	
497	$k_{\rm XCN} = k_{\rm w} + k_{\rm OH} [\rm OH^{-}] $ (Eq10)
498	
499	Bailey and Bishop (1973) found $k_w = 2.58 \times 10^{-6} \text{ sec}^{-1}$ and $k_{OH} = 4.53 \text{ M}^{-1} \text{sec}^{-1}$ at 299.7 K, for ClCN, which
500	corresponds to 3.03 x10 ⁻⁶ at pH7. This is consistent with the results of this study which found that the first-order loss
501	rate was zero, within the error of the linear fit ($\pm 4.2 \times 10^{-4} \text{ sec}^{-1}$). The study of BrCN hydrolysis of Gerritsen et al.,
502	1993 did not derive k_w nor did it present sufficient data for k_w to be estimated. However, there are two other studies
503	that presented data from which k_w can be estimated, and those range from $1.9 - 9.2 \times 10^{-5}$ sec ⁻¹ , (Heller-Grossman et al. 1997) and 1997 are secret as the secret secre
504	al., 1999; Vanelslander et al., 2012).
505	The hydrolysis of ICN is slightly more complicated since there is some evidence that ICN might complex
506	with iodide (Gerritsen et al., 1993). The room temperature hydrolysis rate observed in our experiment was not
507	significantly different than zero, 4.4 (\pm 7.6) ×10 ⁻⁵ sec ⁻¹ , but is in the same range of the rate constant estimated from
508	the data given by Gerritsen, et al., (1993), by extrapolating their rate constant vs. [OH ⁻] data to zero [OH ⁻], assuming
509	no complexation reactions.
510	
511	B. Non-aqueous Solution
512	
513	Solubility in non-aqueous solvents is a standard indicator of how compounds will be distributed between
514	different compartments in the environment, i.e. lipids in the body, organic aerosols in the atmosphere. In addition,
515	the ratio of organic to aqueous solubility (Kow) is used to estimate membrane transport of a chemical species, a key
516	factor in estimating physiologic effects of a pollutant. Several non-aqueous solvents were used in this study,
517	tridecane to represent a completely non-polar solvent and n-octanol, which is used as a standard material for such
518	studies.
519	
520	1. Solubility and Reactions of HNCO
521	
522	The experiments performed on HNCO were conducted with tridecane, 10% (V/V) n-octanol/tridecane, and
523	pure n-octanol, and the results are summarized in Table 1. HNCO is the least soluble in tridecane, 1.7 (±0.17) M/atm
524	and increasingly soluble as the proportion of n-octanol is increased, to pure n-octanol, 87 (±9) M/atm at 298 K.
525	Experiments at two other temperatures were performed to confirm that these solubilities follow the expected
526	temperature dependence, and to obtain the solubility in pure n-octanol at body temperature (310 K) to match data for
527	the aqueous solubility. The lower solubility of HNCO in tridecane is expected since tridecane is completely non-
528	polar and has no tendency to hydrogen bond or interact with the polarizable end of the HNCO molecule. In contrast,

529 the increase in solubility of HNCO with increasing proportion of n-octanol is due to the polar –OH group at the end 530 of the molecule.

The rate of reactions of HNCO with the non-aqueous solvents were below the limit of detection by this method for all combinations except for pure n-octanol at 310 K. Even still, the measured rate was quite a bit lower than the corresponding hydrolysis rate in aqueous solution at pH=3. The manner in which these two factors (solubility and reaction) affect the net uptake and loss of HNCO will be discussed below.

- 535
- 536 537

2. Solubility and Reactions of CH₃NCO

The solubility of CH₃NCO in n-octanol was measured at several temperatures, as summarized in Table 1. The value for 298 K is approximately 3 times higher than that of aqueous solubility, and has the expected temperature dependence. In addition, the first-order reaction rates for CH₃NCO in n-octanol were in the same range or slightly higher than the aqueous reactions. The reaction with n-octanol is expected to go via the carbamylation reaction (R9), although there is some evidence that this reaction has as more complex mechanism possibly involving multiple alcohol molecules (Raspoet et al., 1998). These rates are much faster than the corresponding rates for HNCO, and may provide some guidance concerning the loss rates of CH₃NCO to heterogeneous processes.

545 546

547

3. CICN and BrCN

548 The solubilities of CICN and BrCN in n-octanol were measured at room temperature. Cyanogen chloride has an 549 only a slightly higher Henry's coefficient in n-octanol than in water, in contrast to BrCN, which is relatively more 550 soluble in n-octanol. The first-order loss rates of CICN and BrCN could be determined from the flow reactor 551 experiments and were 1.3 (± 0.4) $\times 10^{-3}$ sec⁻¹ and 9 (± 2) $\times 10^{-5}$ sec⁻¹, respectively. Reactions of ClCN with alcohols are 552 known (see for example (Fuks and Hartemink, 1973)), and form carbamates, in a mechanism that appears to be second-553 order in the alcohol, and acid catalyzed, but rate constants for ClCN-alcohol reactions have not been reported to our 554 knowledge. There are studies of rates of reactions of CICN with nucleophiles, e.g. nitrogen bases, and those reactions 555 appear to result in -CN substitution and formation of a Cl⁻ ion (Edwards et al., 1986). In addition, BrCN has been used 556 by protein chemists to selectively cleave disulfide bonds and has been used for some time by synthetic chemists to 557 selectively convert tertiary amines to secondary amines (Siddiqui and Siddiqui, 1980; von Braun and Schwarz, 1902) 558 and can carbamylate amino acids (Schreiber and Witkop, 1964). The importance of these reactions to the atmospheric 559 fate of XCN compounds remains an open question, but it is important to note that they constitute losses of active 560 halogen, i.e. conversion of the halogen to a halide ion.

- 561
- 562

4. Octanol/Water Partition Coefficients

563

The ratio of solubilities between a non-polar solvent and water is a fundamental quantity that is useful in predicting the fate of a compound in the environment and biological systems (Leo et al., 1971). This parameter is 566 used to predict lipid solubility, membrane transport, and the potential of uptake of a particular compound by organic

aerosol. n-Octanol is a standard non-polar solvent that is commonly for this purpose, as it has an overall non-polar

- 568 character with a substituent that is capable of hydrogen bonding. The data from this study can be used to calculate
- the octanol/water partition coefficients for HNCO, CH₃NCO, ClCN, and BrCN as the ratio of the respective Henry's

570 coefficients;

580

596

571 $K_{ow} = H_{oct}/H_{H2O}$

(Eq.11)

572 The results are listed in Table 3 along with Kows for some related small molecules. Both CH₃NCO, and BrCN are 573 fundamentally more soluble in n-octanol than in water, while CICN has nearly the same solubility in both materials. 574 The weak acid equilibrium of HNCO makes it more soluble in n-octanol at pH 3, but much more soluble in water at 575 neutral pH. Formic acid is a similarly weak acid (pKa = 3.77) and so is a good point of comparison to HNCO. The 576 n-octanol partition coefficient of HNCO is a factor of 15 larger than that of formic acid. Similarly, the n-octanol 577 partition coefficient of CH₃NCO is 6.8 times larger than that of acetonitrile. The two cyanogen halides measured 578 here had differing behavior, with CICN showing almost no difference in solubility, and BrCN having about the same 579 increase in solubility in n-octanol as HNCO and CH₃NCO.

581 IV. Atmospheric and Environmental Chemistry Implications582

583 The atmospheric loss of the compounds studied here are either: solely or predominantly through 584 heterogeneous uptake and reaction for HNCO, CH₃NCO, CICN, BrCN, or in the case of ICN due to both 585 heterogeneous chemistry and photolysis. The aqueous solubility and reaction data from this study allow some 586 prediction of uptake parameters and loss rates in some important systems, e.g. cloud water and natural water 587 surfaces like oceans. In addition, some indications can be gained about the uptake of HNCO, CH₃NCO, ClCN, and 588 BrCN to organic aerosol, using n-octanol as a model. Finally, the n-octanol/water partition coefficient is often used 589 as a key parameter in modeling cross-membrane transport, and the data from this study can be used to predict the 590 behavior of these reduced-N compounds relative to other well-studied compounds.

591 The reactive uptake of HNCO, CH₃NCO and XCN on environmental surfaces, small particles and aqueous 592 droplets can be parameterized using the uptake coefficient, γ , defined as the fraction of collisions of a molecule with 593 a surface that lead to incorporation of that molecule in the condensed phase. If solubility and reaction are the 594 limiting processes, a good assumption for the species in this work, then γ_{fxn} can be estimated from the following 595 equation (Kolb et al., 1995):

$$\gamma_{rxn} = \frac{4HRT\sqrt{kD_a}}{\langle c \rangle} \tag{Eq.12}$$

where H and k are the Henry's coefficient and first-order loss rate in solution measured in this work, R is the gas constant, T is temperature, D_a is the diffusion coefficient in aqueous solution (assumed here to be 1.9×10^{-5} cm²sec⁻¹ for HNCO and 1.6×10^{-5} cm²sec⁻¹ for CH₃NCO and ClCN, 1.2×10^{-5} cm²sec⁻¹ for BrCN, and 1.0×10^{-5} cm²sec⁻¹ for ICN at 298 K), and <c> is the mean molecular velocity. The results of these calculations are shown in Figure 8 for the H measurements reported here, k_{hydr} for HNCO reported by Borduas et al., (2016), k_{hdyr} or CH₃NCO from this work and k_{hydr} for ClCN from Bailey and Bishop (1973). 603 Deposition of a compound to the surface can be parameterized as essentially two processes taking place in 604 series, physical transport within the planetary boundary layer to the surface and then chemical uptake on the surface 605 (see for example (Cano-Ruiz et al., 1993)). In this formulation, the deposition velocity, v_d (the inverse of the total 606 resistance) is expressed as follows:

607 608

$$v_d = \frac{1}{\frac{1}{v_t} + \frac{1}{\frac{\zeta C}{4}}}$$
(Eq.13)

609

623

610 where $1/v_t$ is the resistance due to transport, and $\frac{1}{\frac{1}{\gamma_{\frac{2}{4}}}}$ is the resistance due to chemical uptake. For a species for 611 which uptake is rapid, e.g. a highly soluble acid, the chemical resistance becomes small and $v_d \approx v_t$. This is the case 612 for HNCO deposition to land or natural water surfaces (pHs ~7-8). Typical v_t s are on the order of 0.5 to 1 cm/sec for 613 a reasonably mixed boundary layer (Wesely and Hicks, 2000). For compounds for which γ is quite small, the 614 chemical term predominates.

615
$$v_d \cong \gamma \frac{\langle c \rangle}{4}$$
 (Eq. 14)

616 The lifetime of a species within the PBL then can be estimated as h/v_d , where h is the boundary layer height. The 617 lifetime estimates for HNCO, CH₃NCO, and XCN compounds are given in Table 2, and range from the short 618 lifetime noted for HNCO, to quite long lifetimes for the least soluble species, for example ClCN.

The loss rates due to uptake of species to atmospheric aerosol particles can be estimated from the pH
dependent uptake coefficients in Figure 8, using parameterizations described in the literature (Davidovits et al.,
2006; Sander, 1999). In the limited case of surface-controlled uptake, i.e. neglecting gas phase diffusion, the loss of
a species is;

$$k = \frac{A\gamma < c>}{4}$$
(Eq. 15)

where A is the aerosol surface area. If we take the γ s from Figure 8, and assume highly polluted conditions, A = 1000 μ m²/cm³ and pHs between 1 and 2, then the lifetimes listed in Table 2 are arrived at. The values for HNCO and CH₃NCO show a range because the uptake is pH dependent, and it should be noted that the values for CH₃NCO, ClCN, and BrCN are over estimated by this method, as their chemistry is slow enough that a volume-based estimate may be more appropriate. The more important effect here is that the γ values are based on hydrolysis losses, which are undoubtedly much slower than many of the solution-phase reactions that these species can undergo, hence the lifetimes against aerosol deposition are upper limits.

631The loss of HNCO to cloudwater is the subject of extensive work discussed by Barth et al, 2013, and no632attempt will be made here to update that analysis. The fastest loss rates for HNCO were observed in warm dense633clouds into which SO2 was also dissolving and adding considerable acidity, so that value for HNCO was included in634Table 2. For the other compounds we use a simple parameterization of cloudwater reaction to estimate the in-cloud635loss rates for CH₃NCO and the XCN compounds. In the estimate of reaction rate:636 $k = k_1 L_c HRT$ (Eq. 16)

- 637 k₁ is the liquid phase rate constant, L_c is the cloud liquid water content, and H is the Henry's coefficient. If we
- 638 assume a L_c of 2 x10⁻⁶, and T \cong 298 K, and we use the H and k values measured in this work (the exception is that
- the literature value for CH_3NCO at pH=2 was used), then the values for lifetimes of CH_3NCO , and XCN compounds
- 640 listed in Table 2 were obtained. Below we discuss the characteristic times obtained for each compound in the
- 641 context of what else is known about their sources and atmospheric chemistry.

643 A. HNCO

642

- 644 645 The loss of HNCO via heterogeneous processes occurs in two separate regimes: in aerosols and cloud 646 droplets at relatively low pH, and in surface waters and on terrestrial surfaces that are neutral or slightly basic in pH. 647 In the former case, HNCO solubility is relatively low but hydrolysis is acid catalyzed. In the latter case, solubility is 648 high enough that uptake will be limited by the transport of HNCO to the surface, much like other strong acids such 649 as HNO₃. Ambient measurements of HNCO at surface sites are consistent with deposition of HNCO to the ground, 650 exhibiting diurnal profiles similar to those of O₃ or HNO₃ (Kumar et al., 2018; Roberts et al., 2014; Mattila et al.,
- 651 2018; Zhao et al., 2014).
- 652 Several aspects of the aqueous solubility and hydrolysis, and heterogeneous removal of HNCO have been 653 examined in modeling studies. A global modeling study by Young et al. (2012) was a first attempt to model global 654 HNCO by scaling the source to fire emissions of HCN. Loss of HNCO was assumed to be due to wet and dry 655 deposition with efficiencies similar to HNO₃ and HC(O)OH, and that HNCO was lost once it was taken up by 656 clouds. Young et al., concluded that HNCO had an average lifetime of about 37 days. Barth et al., (2013) addressed 657 part of this analysis by modeling the cloud removal of HNCO using actual solubility and reaction data in a cloud 658 parcel model, albeit, the hydrolysis rates used were from Jensen, (1957) which were approximately 50% higher than 659 the Borduas results, and the temperature dependence of H was assumed equal to that of formic acid, and resulted in 660 higher solubilities at low temperature. This cloud model showed that cloud water uptake was reversible in that most 661 cases hydrolysis was slow enough that some HNCO returned to the gas phase after cloud evaporation. The Barth et 662 al., study estimated HNCO lifetimes as short as 1 hour in warm polluted clouds (i.e. high $SO_2 => H_2SO_4$ formation). 663 The results of our study and those of Borduas et al., (2016) add to these analyses in that now the measured 664 temperature dependence of H can be used, and the hydrolysis rate constants can be updated. 665 The results in this paper allow for further refinement of HNCO loss estimates. For example, the salting-out 666 effect may be important for aerosol with high inorganic content, and high ammonium concentrations will result in 667 reactive loss rates that are faster than hydrolysis. The solubility of HNCO in aerosol particles with substantial
- organic character can be higher or low depending on the nature of substituent groups, e.g. degree of -OH
- 669 functionalization.

670 In studies of the condensed phase oxidation of dissolved N species, as well as biological processes produce 671 cyanate ion, there is a growing recognition that cyanate is part of the natural N cycle in the ocean (see (Widner et al., 672 2013) and references there-in). Observed near-surface cyanate levels often reached a few 10s of nM in near shore 673 productive areas. The observations of cloud/aerosol source of HNCO presented in (Zhao et al., 2014) on the coast 674 of California might be explained by a combination of this NCO⁻ seawater source and aerosol/cloud water 675 acidification by local sources of strong acids, particularly HNO₃. In specific, acidification of sea spray containing

- 676 about 10 nM NCO⁻ to pH=4 or so, would correspond to H_{eff} of around 50 M/atm, and result in an equilibrium
- 677 HNCO concentration of several hundred pptv. Such a source would most likely be limited by the concentration of
- 678 sea salt-derived aerosol, but could easily account for the source implied by the measurements of (Zhao et al., 2014).

680 B. CH₃NCO

679

690

681 The atmospheric chemistry of CH₃NCO is less well studied than HNCO. There is a single reported 682 measurement of the reaction rate of CH₃NCO with OH by relative rates which gave $k = 3.6 \times 10^{-12}$ molec cm⁻³ sec⁻¹ 683 (Lu et al., 2014). The uptake coefficients estimated for CH₃NCO in Figure 8 are relatively low with only a slight 684 increase at the lowest pHs in atmospheric media. As a consequence, atmospheric lifetimes of CH₃NCO towards 685 surface deposition are estimated to be quite long, 6 months or more if hydrolysis is the sole loss process. The loss 686 due to aerosol of cloudwater uptake is estimated to be slightly faster, due primarily to the slight acid-catalysis of the 687 CH₃NCO hydrolysis rate. The lifetime estimates should be considered upper limits since there are number of 688 condensed-phase reactions that might be faster than hydrolysis, and would need to be the subject of further research. 689

691 C. CICN, BrCN, and ICN 692

693 To date, we know of no observations of CICN in the ambient atmosphere, but its formation in the 694 chlorination of water, waste water, and swimming pools (Afifi and Blatchley III, 2015; Daiber et al., 2016; Lee et 695 al., 2006) indicates that there could be sources from human activities, including the use of chlorine bleach for 696 cleaning indoor surfaces. In addition, there might also be a source from aerosol systems were chlorine is being 697 activated, i.e. oxidized from Cl⁻ to ClNO₂, Cl₂, or HOCl (see for example (Roberts et al., 2008)) in the presence of 698 reduced nitrogen. The results of our solubility measurements indicate that CICN will volatilize from the condensed 699 phase fairly readily, so its atmospheric removal should be considered. BrCN has been observed in systems where 700 bromide-containing water or wastewater were treated with halogens (Heller-Grossman et al., 1999), and there are 701 biological mechanisms that make BrCN and ICN as well (Schlorke et al., 2016; Vanelslander et al., 2012). The 702 potential for remote atmospheric sources of these compounds is currently being investigated, but BrCN could be the 703 result of the same bromine activation chemistry that depletes ground level ozone in that environment (Simpson et 704 al., 2007).

Gas phase radical reactions of XCN compounds have not been studied under atmospheric conditions. A few studies at higher temperatures and the studies of HCN and CH₃CN can be used to roughly predict how fast the relevant reactions are. For example, the reactions of ClCN and BrCN with O atoms at 518-635 K are very slow (<3 $\times 10^{-15}$ molec cm⁻³ sec⁻¹, (Davies and Thrush, 1968)) and the reaction of Cl atom with ClCN at high temperature is also quite slow (<1.0 ×10⁻¹⁴ molec cm⁻³ sec⁻¹, (Schofield et al., 1965)). However, these observations do not preclude the presence of another reaction channel at low temperature, e.g. a mechanism involving addition to the CN group.

- 711 The reactions of HCN and CH₃CN with OH, Cl atom and O atom at atmospherically relevant temperatures are all
- 712 quite slow, implying such addition channels are not likely to be substantially faster for these XCN compounds. We
- conclude that rate constants for the reactions of OH or Cl with X-CN compounds are likely quite low ($<2 \times 10^{-14}$

714 molec cm⁻³ sec⁻¹), making the lifetimes of these compounds against these reactions on the order of a year or longer. 715 The UV-visible absorption spectra of all three of these compounds have been measured (Barts and Halpern, 1989; 716 Felps et al., 1991; Hess and Leone, 1987; Russell et al., 1987), have maxima that range from <200nm for ClCN, 717 202nm for BrCN, and 250nm for ICN, with absorption that tails into near-UV and visible wavelengths, (see Figure 718 S2 in the Supplemental Material). Extrapolation of the spectra, combined with photo fluxes estimated from the 719 NCAR TUV model for mid-summer 40° North at the surface, result in a range of photolysis behavior ranging from 720 no tropospheric photolysis of ClCN, to slight photolysis of BrCN ($\tau \approx 135$ days), and faster photolysis of ICN ($\tau \approx 9$ 721 hours). The above gas phase processes provide the context in which to assess the importance of condensed phase 722 loss processes of ClCN, BrCN, and ICN. Rates of loss of XCN compounds due to surface deposition, cloudwater or 723 aerosol uptake would need to be faster than the gas phase processes to be important in the atmosphere. In addition, 724 condensed phase reactions convert XCN to halide ions either by hydrolysis to cyanate, or creation of a carbamyl 725 functionalities. Only photolysis reforms the halogen atom, and therefore maintains active halogen reaction chain. 726 Estimated atmospheric lifetimes of XCN compounds against loss due to condensed phase reactions listed in Table 2 727 shows a general trend. The lifetimes become shorter as the halogen atom goes from Cl to Br to I, primarily due to 728 higher solubilities. The actual condensed phase losses are likely much shorter than those estimated here because of 729 faster condensed phase reactions that are not taken into account by the brief analysis presented here.

730

732

731 D. Solubility in non-polar media, uptake to organic aerosol, and membrane transport.

733 The solubilities of HNCO, CH₃NCO, and BrCN in n-octanol were roughly a factor 4 larger than water, 734 while that of CICN was virtually the same. Reaction rates with n-octanol were the same or slower than for aqueous 735 solutions, except for CICN which was faster than hydrolysis at pH=7. As a result, loss due to uptake to organic 736 aerosol will be only slightly faster for all of these species. Membrane transport is a key process in determining the 737 extent to which a chemical species will impact biological systems. Simple membrane transport models parameterize 738 this process as diffusion through a lipid bi-layer according to a partition coefficient, K_p, which the ratio of 739 solubilities in lipid versus aqueous media (Missner and Pohl, 2009), and Kow is often used for this partition 740 coefficient. The results of our work indicate that both HNCO and CH3NCO are more soluble in n-octanol than 741 water, in contrast to other similar small organic acids and N-containing compounds (Table 3). These features will 742 need to be accounted for in assessing the connection by between environmental exposure to HNCO, CH₃NCO, 743 CICN and BrCN and resulting biochemical effects. 744 745 V. Data availability. The data are available on request. 746

VI. Author contributions. YL and JR performed the laboratory experiments and JR and YL wrote the paper.

- 747 748
- 749 VII. Competing interests. The author declare no competing interests.
- 750

752	not be construed as an endorsement.
753	
754	IX. Acknowledgements
755	
755	
756 757 758 759 760	We thank Drs. James B. Burkholder, Françios Bernard, for help with the FTIR analysis and Drs. Patrick Veres and Bin Yuan for the CIMS analyses of XCN samples. We thank Dr. Dimitris Papanastasiou, Dr. J. Andrew Neuman, and Dr. Patrick Veres for communicating the results of their work prior to publication. This work was supported in part by NOAA's Climate and Health of the Atmosphere Initiatives.
761 762 763	X. References
764 765 766	Afifi, M. Z. and Blatchley III, E. R.: Seasonal dynamics of water and air chemistry in an indoor chlorinated swimming pool, Water Res., 68, 771-783, 2015.
767 768 769 770	Al-Rawi, H. and Williams, A.: Elimination-addition mechanisms of acyl group transfer: the hydrolysis and synthesis of carbamates, J. Am Chem. Soc., 99, 2671-2678, 1977.
771 772 773	Bailey, P. L. and Bishop, E.: Hydrolysis of Cyanogen Chloride, Journal of the Chemical Society, Dalton Transactions, doi: 10.1039/dt9730000912, 1973. 912-916, 1973.
774 775 776	Barnes, I., Solignac, G., Mellouki, A., and Becker, K. H.: Aspects of the atmospheric chemistry of amides, Chem Phys Chem, 11, 3844-3857, 2010.
777 778 779	Barth, M. C., Cochran, A. C., Fiddler, M. N., Roberts, J. M., and Bililign, S.: Numerical modeling of cloud chemistry effects on isocyanic acid (HNCO), J. Geophys. Res., 118, 2013.
780 781 782	Barts, S. A. and Halpern, J. B.: Photodissociation of CICN between 190 and 213 nm, J. Phys. Chem., 93, 7346-7351, 1989.
783 784 785	Belson, D. J. and Strachan, A. N.: Preparation and properties of isocyanic acid, Chem. Soc. Rev., 11, 41-56, 1982. Bengtstrom, L., Salden, M., and Stec, A. A.: The role of isocyanates in fire toxicity, Fire Sci. Rev., 5, 1-23, 2016.
786 787 788	Blomqvist, P., Hertzberg, T., Dalene, M., and Skarping, G.: Isocyanates, aminoisocyanates, and amines from fires - a screening of common materials found in buildings, Fire Mater., 27, 275-294, 2003.
789 790 791 792	Borduas, N., da Silva, G., Murphy, J. G., and Abbatt, J. P. D.: Experimental and theoretical understanding of the gas phase oxidation of atmospheric amides with OH radicals: Kinetics, products, and mechanisms, J. Phys. Chem. A, 119, 4298-4308, 2015.
793 794 795	Borduas, N., Place, B., Wentworth, G. R., Abbatt, J. P. D., and Murphy, J. G.: Solubility and reactivity of HNCO in water: insights into HNCO's fate in the atmosphere, Atmos. Chem. Phys., 16, 703-714, 2016.
796 797 798 799	Bunkan, A. J. C., Hetzler, J., Mikoviny, T., Wisthaler, A., Nielsen, C. J., and Olzmann, M.: The reactions of N- methylformamide and N,N-dimethylformamide with OH and their photo-oxidation under atmospheric conditions: experimental and theoretical studies, Phys. Chem. Chem. Phys., 17, 7046-7059, 2015.
800 801 802	Cano-Ruiz, J. A., Kong, D., Balas, R. B., and Nazaroff, W. W.: Removal of reactive gases at indoor surfaces: Combining maas transport and surface kinetics Atmos. Environ., 27A, 2039-2050, 1993.

752

VIII. Disclaimer. Any mention of commercial products or brands were solely for identifying purposes and should

- Castro, E. A., Moodie, R. B., and Sansom, P. J.: The Kinetics of Hydrolysis of Methyl and Phenyl Isocyanates, J.
 Chem. Soc. Perkin Trans II, 2, 737-742, 1985.
- 805

- 806 Chandra, B. P. and Sinha, V.: Contribution of post-harvest agricultural paddy residue fires in the N.W.
- 807 Indo-Gangetic Plain to ambient carcinogenic benzenoids, toxic isocyanic
- 808 acid and carbon monoxide, Environ. Intl., 88, 187-197, 2016. 809
- 810 Clever, H. L.: Sechenov Salt-Effect Parameter, J. Chem. Eng. Data, 28, 340-343, 1983. 811

Baiber, E. J., DeMarini, D. M., Ravuri, S. A., Liberatore, H. K., Cuthbertson, A. A., Thompson-Klemish, A., Byer,
J. D., Schmid, J. E., Afifi, M. Z., Blatchley III, E. R., and Richardson, S. D.: Progressive Increase in Disinfection
Byproducts and Mutagenicity from Source to Tap to Swimming Pool and Spa Water: Impact of Human Inputs,
Environ. Sci. Technol., 50, 6652-6662, 2016.

- 816
 817 Davidovits, P., Kolb, C. E., Williams, L. R., Jayne, J. T., and Worsnop, D. R.: Mass accommodation and chemical
 818 reactions at gas-liquid interfaces, Chemical Reviews, 106, 1323-1354, 2006.
- Bavies, P. B. and Thrush, B. A.: Reactions of oxygen atoms with hydrogen cyanide, cyanogen chloride and cyanogen bromide, Trans. Faraday Soc., 64, 1836-1843, 1968.
- Bavis, D. D. and Okabe, H.: Determination of bond dissociation energies in hydrogen cyanide, cyanogen and
 caynogen halides by the photodissociation mehtod., J. Chem. Phys., 49, 5526-5531, 1968.
- Diehl, A. C., Speitel Jr., G. E., Symons, J. M., Krasner, S. W., Hwang, C. J., and Barrett, S. E.: DBP formation
 during chloramination, J. Am. Water Works Assoc., 92, 76-90, 2000.
- Edwards, J. O., Erstfeld, T. E., Ibne-Rasa, K. M., Levey, G., and Moyer, M.: Reaction rates for nucleophiles with
 cyanogen chloride: Comparison with two other digonal carbon compounds, Int. J. Chem. Kinet., 18, 165-180, 1986.
- Epstein, J.: Estimation of microquantities of cyanide, Anal. Chem., 19, 272-274, 1947.
- Felps, W. S., Rupnik, K., and McGlynn, S. P.: Electronic spectroscopy of the cyanogen halides. , J. Phys. Chem., 95,
 639-656, 1991.
- Fuks, R. and Hartemink, M. A.: The acid-catalyzed reaction of cyanogen chloride with aliphatic alcohols. A general
 synthesis of aliphatic carbamates, Bulletin des Societes Chimiques Belges, 82, 23-30, 1973.
- Garrido, M. A., Gerecke, A. C., Heeb, N., Font, R., and Conesa, J. A.: Isocyanate emissions from pyrolysis of
 mattresses containing polyurethane foam, Chemosphere, 168, 667-675, 2017.
- Geddes, J. D., Miller, G. C., and Taylor Jr., G. E.: Gas phase photolysis of methyl isothiocyanate, Environ. Sci.
 Technol., 29, 2590-2594, 1995.
- Gerritsen, C. M., Gazda, M., and Margerum, D. W.: Non-metal redox kinetics: Hypobromite and hypoiodite
 reactions with cyanide and the hydrolysis of cyanogen halides, Inorg. Chem., 32, 5739-5748, 1993.
- Goesmann, F., Rosenbauer, H., Bredehoft, J. H., Cabane, M., Ehrenfreund, P., Gautier, T., Giri, C., Kruger, H., Le
 Roy, L., MacDermott, A. J., McKenna-Lawlor, S., Meierhenrich, U. J., Munoz Caro, G. M., Raulin, F., Roll, R.,
 Steele, A., Steininger, H., Sternberg, R., Szopa, C., Thiemann, W., and Ulamec, S.: Organic compounds on comet
- 67P/Churyumov-Gerasimenko revealed by COSAC mass spectrometry, Science, 349, aab0689-0681,0683, 2015.
- Hagel, P., Gerding, J. J. T., Fieggen, W., and Bloemendal, H.: Cyanate formation in solutions of urea .1. Calculation
 of cyanate concentrations at different temperature and pH, Biochimica et Biophysica Acta, 243, 366, 1971.
- Halfen, D. T., Ilyushin, V. V., and Ziurys, L. M.: Interstellar detection of methyl isocyanate CH₃NCO in Sgr B2(N):
 A link from molecular clouds to comets, Astrophysical Journal Letters, 812, L5, 2015.

862

Hardy, J. E. and Knarr, J. J.: Technique for measuring the total concentration of gaseous fixed nirogen species, J. Air
 Pollut. Contr. Assoc., 32, 376-379, 1982.

Heller-Grossman, L., Idin, A., Limoni-Relis, B., and Rebhun, M.: Formation of cyanogen bromide and other volatile
DBPs in the disinfection of bromide-rich lake water., Environ. Sci. Technol., 33, 932-937, 1999.

Hess, W. P. and Leone, S. R.: Absolute I* quantum yields for the ICN Ã state by diode laser gain-vs-absorption
spectroscopy, J. Chem. Phys., 86, 3773-3780, 1987.

Hilal, S. H., Ayyampalayam, S. N., and Carreira, L. A.: Air-Liquid Partition Coefficient for a Diverse Set of Organic
Compounds: Henry's Law Constant in Water and Hexadecane, Environ. Sci. Technol., 42, 9231-9236, 2008.

Jensen, M. B.: On the kinetics of the decomposition of cyanic acid, Acta Chem. Scand., 12, 1657-1670, 1958.

Jensen, M. B.: On the kinetics of the reaction of cyanic acid with amines, Acta Chem. Scand., 13, 289-300, 1959.

Kames, J. and Schurath, U.: Henry's law and hydrolysis-rate constants for peroxyacetyl nitrates (PANs) using a
homogeneous gas-phase source, J. Atmos. Chem., 21, 151-164, 1995.

Kish, J. D., Leng, C., Kelley, J., Hiltner, J., Zhang, Y., and Liu, Y.: An improved approach for measuring Henry's
law coefficients of atmospheric organics, Atmos Environ., 79, 561-565, 2013.

Kolb, C. E., Worsnop, D. R., Zahniser, M. S., Davidovits, P., Keyser, L. F., Leu, M.-T., Molina, M. J., Hanson, D.
R., Ravishankara, A. R., Williams, L. R., and Tolbert, M. A.: Laboratory studies of atmospheric heterogeneous
chemistry. In: Progress and Problems in Atmospheric Chemistry, Barker, J. R. (Ed.), Advanced Series in Physical
Chemistry, 3, World Scientific, Singapore, 1995.

Koss, A. R., K., S., Gilman, J. B., Selimovic, V., Coggon, M. M., Zarzana, K. J., Yuan, B., Lerner, B. M., Brown, S.
S., Jimenez, J. L., J., K., Roberts, J. M., Warneke, C., Yokelson, R. J., and de Gouw, J.: Non-methane organic gas
emissions from biomass burning: identification, quantification, and emission factors from PTR-ToF during the
FIREX 2016 laboratory experiment, Atmos. Chem. Phys., 18, 3299-3319, 2018.

Kumar, V., Chandra, B. P., and Sinha, V.: Large unexplained suite of chemically reactive compounds present in
ambient air due to biomass fires., Scientific Reports, 8, 2018.

- Lee, J. H., Na, C., Ramirez, R. L., and Olsen, T. M.: Cyanogen chloride precursor analysis in chlorinated river
 water, Environ. Sci. Technol., 40, 1478-1484, 2006.
- Leo, A., Hansch, C., and Elkins, D.: Partition coefficients and their uses, Chem. Rev., 71, 525-616, 1971.

Lu, Z., Hebert, V. R., and Miller, G. C.: Gas-Phase Reaction of Methyl Isothiocyanate and Methyl Isocyanate
with Hydroxyl Radicals under Static Relative Rate Conditions, J. Agric. Food Chem., 62, 1792-1795, 2014.

Maroulis, G. and Pouchan, C.: Dipole polarizability and hyperpolarizability of FCN, ClCN, BrCN and ICN, Chem.
Phys., 216, 67-76, 1997.

Mattila, J. M., Brophy, P., Kirkland, J., Hall, S., Ullman, K., E.V., F., Brown, S., McDuffie, E., Tevlin, A., and
Farmer, D. K.: Tropospheric sources and sinks of gas-phase acids in the Colorado Front Range, Atmos. Chem.
Phys., 18, 12315-12327, 2018.

910 Missner, A. and Pohl, P.: 110 Years of the Meyer–Overton rule: Predicting membrane permeability of gases and 911 other small compounds, Chem Phys Chem, 10, 1405-1414, 2009.

912

913 Na, C. and Olson, T. M.: Mechanism and kinetics of cyanogen chloride formation from the chlorination of glycine,

914 Environ. Sci. Technol., 40, 1469-1477, 2006.

- 915
- 816 Raspoet, G., Nguyen, M. T., McGarraghy, M., and Hegarty, A. F.: The alcoholysis reaction of isocyanates giving
- 917 urethanes: Evidence for a multimolecular mechanism, J. Org. Chem., 63, 6878-6885, 1998.
- 918

919 Reyes-Villegas, E., Bannan, T., Le Breton, M., Mehra, A., Priestley, M., Percival, C., Coe, H., and Allan, J. D.:

- Online chemical characterization of food-cooking organic aerosols: Implications for Source Apportionment,
 Environ. Sci. Technol., 52, 5308-5318, 2018.
- Roberts, J. M.: Measurement of the Henry's law coefficient and the first order loss rate of PAN in n-octanol,
 Geophys. Res. Lett., 31, L08803, 2005.
- 924 Geophys. Res. Lett., 31, L08803, 2005. 925
- Roberts, J. M., Osthoff, H. D., Brown, S. S., and Ravishankara, A. R.: N₂O₅ Oxidizes Chloride to Cl₂ in Acidic
 Atmospheric Aerosol, Science, 321, 1059., 2008.
- 929 Roberts, J. M., Veres, P., Warneke, C., Neuman, J. A., Washenfelder, R. A., Brown, S. S., Baasandorj, M.,
- Burkholder, J. B., Burling, I. R., Johnson, T. J., Yokelson, R. J., and de Gouw, J.: Measurement of HONO, HNCO,
 and other inorganic acids by negative-ion proton-transfer chemical-ionization mass spectrometry (NI-PT-CIMS):
 Application to biomass burning emissions., Atmos. Meas. Tech., 3, 981-990, 2010.
- 933
 934 Roberts, J. M., Veres, P. R., Cochran, A. K., Warneke, C., Burling, I. R., Yokelson, R. J., Lerner, B. M., Gilman, J.
- B., Kuster, W. C., Fall, R., and de Gouw, J.: Isocyanic acid in the atmosphere and its possible link to smoke-related
- 936 health effects, PNAS, 108, 8966-8971, 2011.
- Roberts, J. M., Veres, P. R., VandenBoer, T. C., Warneke, C., Graus, M., Williams, E. J., Holloway, J. S., Lefer, B.,
 Brock, C. A., Bahreini, R., Ozturk, F., Middlebrook, A. M., Wagner, N. L., Dube, W. P., and de Gouw, J. A.: New
 insights into atmospheric sources and sinks of isocyanic acid, HNCO, from recent urban and regional observations,
 J. Geophys. Res., 119, 2014.
- Russell, J. A., McLaren, I. A., Jackson, W. M., and Halpern, J. B.: Photolysis of BrCN between 193 and 266 nm, J.
 Phys. Chem., 91, 3248-3253, 1987.
- Sander, R.: Compilation of Henry's law constants (version 4.0) for water as solvent, Atmos. Chem. Phys., 15, 43994981, 2015.
- 948
- Sander, R.: Modeling atmospheric chemistry: Interactions between gas-phase species and liquid cloud/aerosol
 particles, Surv. Geophys., 20, 1-31, 1999.
- Sarkar, C., Sinha, V., Kumar, V., Rupakheti, M., Panday, A., Mahata, K. S., Rupakeheti, D., Kathayat, B., and
 Lawrence, M. G.: Overview of VOC emissions and chemistry from PTR-TOF-MS measurements during the
 SusKat-ABC campaign: high acetaldehyde, isoprene and isocyanic acid in wintertime air of the Kathmandu Valley,
 Atmos. Chem. Phys., 16, 3979-4003, 2016.
- Saylor, R. D., Edgerton, E. S., Hartsell, B. E., Baumann, K., and Hansen, D. A.: Continuous gaseous and total ammonia measurements from the southeastern aerosol research and characterization (SEARCH) study, Atmos.
- 959 Environ., 44, 4994-5004, 2010. 960
- Schlorke, D., Flemming, J., Birkemeyer, C., and Arnhold, J.: Formation of cyanogen iodide by lactoperoxidase., J.
 Inorg. Biochem., 154, 35-41, 2016.
- Schofield, D., Tsang, W., and Bauer, S. H.: Thermal decomposition of ClCN, J. Chem. Phys., 42, 2132 2137,
 1965.
- 967 Schreiber, J. and Witkop, B.: The reaction of cyanogen bromide with mono- and diamino Acids, J. Am. Chem. Soc.,
 86, 2441-2445, 1964.
- 970 Schumpe, A.: The estimation of gas solubilities in salt solutions, Chem. Eng. Sci., 48, 153-158, 1993.

994

- 972 Schwab, J. J., Li, Y., Bae, M.-S., Demerjian, K. L., Hou, J., Zhou, X., Jensen, B., and Pryor, S.: A laboratory
- 973 intercomparision of real-time gaseous ammonia measurement methods, Environ. Sci. Technol., 41, 8412-8419,
 974 2007.
 975
- Shang, C., Gong, W.-l., and Blatchley III, E. R.: Breakpoint chemistry and volatile byproduct formation resulting
 from chlorination of maodel organic-N compounds, Environ. Sci. Technol., 34, 1721-1728, 2000.
- 979 Siddiqui, S. and Siddiqui, B. S.: Some extensions of von braun (BrCN) reaction on organic bases, Z. Naturforsch.,
 980 35b, 1049-1052, 1980.
 981
- Simpson, W. R., von Glasow, R., Riedel, K., Anderson, P., Ariya, P., Burrows, J., Carpenter, L. J., FrieB, U.,
 Goodsite, M. E., Heard, D., Hutterli, M., Jacobi, H.-W., Kaleschke, L., Neff, B., Plane, J., Platt, U., Richter, A.,
- Roscoe, H., Sander, R., Shepson, P., Sodeau, J., Steffen, A., Wagner, T., and Wolff, E.: Halogens and their role in
 polar boundary-layer ozone depletion, Atmospheric Chemistry And Physics, 7, 4375-4418, 2007.
- 987
 988
 989
 989
 State of California, OEHHA Chemical Database Air: https://oehha.ca.gov/air/chemicals/methyl-isocyanate, accessed August 23, 2017.
- Stockwell, C. E., Kupc, A., Witkowski, B., Talukdar, R. K., Liu, Y., Selimovic, V., Zarzana, K. J., Sekimoto, K.,
 Warneke, C., Washenfelder, R. A., Yokelson, R. J., Middlebrook, A. M., and Roberts, J. M.: Characterization of a
 catalyst-based conversion technique to measure total particle nitrogen and organic carbon and comparison to a
 particle mass measurement instrument, Atmos. Meas. Tech., 11, 2749-2768, 2018.
- Valentour, J. C., Aggarwal, V., and Sunshine, I.: Sensitive gas chromatographic determination of cyanide, Anal.
 Chem., 46, 924-925, 1974.
- Vanelslander, B., Paul, C., Grueneberg, J., Prince, E. K., Gillard, J., Sabbe, K., Pohnert, G., and Vyeverman, W.:
 Daily bursts of biogenic cyanogen bromide (BrCN) control biofilm formation around a marine benthic diatom.,
 Proc. Natl Acad. Sci., 109, 2412-2417, 2012.
- 1002 von Braun, J. and Schwarz, R.: Die Einwirkung von Bromcyan auf tertiäre Amine, Chem. Ber., 35, 1279-1285, 1902.
 1004
- Weng, S.-C., Weaver, W. A., Blatchley, T. N., Cramer, J. S., Chen, J., and Blatchley III, E. R.: Dynamics of gasphase trichloramine (NCl₃) in chlorinated, indoor swimming pool facilities, Indoor Air, 21, 391-399, 2011.
- Wentzell, J. B., Liggio, J., Li, S.-M., Vlasenko, A., Staebler, R., Lu, G., Poitras, M.-J., Chan, T., and Brook, J. R.:
 Measurements of gas phase acids in Diesel exhaust: A relevant source of HNCO?, Environ. Sci. Technol., 47, 76637671, 2013.
- Wesely, M. L. and Hicks, B. B.: A review of the current status of knowledge on dry deposition, Atmospheric
 Environment, 34, 2261-2282, 2000.
- Widner, B., Mulholland, M. R., and Mopper, K.: Chromatographic determination of nanomolar cyanate
 concentrations in estuarine and sea waters by precolumn fluorescence derivatization. , Anal. Chem., 85, 6661-6666,
 2013.
- Williams, A. and Jencks, W. P.: Acid and base catalysis of urea synthesis: Nonlinear Bronsted plots consistent with
 a diffusion-controlled proton-transfer mechanism and reactions of imaidazole and N-methylimidazole with cyanic
 acids, J. Chem. Soc., Perkins II, 1974a. 1760-17868, 1974a.
- Williams, A. and Jencks, W. P.: Urea Synthesis from amines and cyanic acid: kinetic evidence for a zwitterionic
 intermediate, J. Chem. Soc., Perkins II, 1974b. 1753-1759, 1974b.
- 1025

- 1026 Williams, E. J., Baumann, K., Roberts, J. M., Bertman, S. B., Norton, R. B., Fehsenfeld, F. C., Springston, S. R.,
- Nunnermacker, L. J., Newman, L., Olszyna, K., Meagher, J., Hartsell, B., Edgerton, E., Pearson, J. R., and Rodgers,
 M. O.: Intercomparison of ground-based NO_y measurement techniques, Journal of Geophysical Research Atmentaneous 102, 222(1, 22200, doi:10.2020/20200174, 1000
- 1029 Atmospheres, 103, 22261-22280, doi: 22210.21029/22298JD00074, 1998. 1030
- Williams, J., Roberts, J. M., Bertman, S. B., Stroud, C. A., Fehsenfeld, F. C., Baumann, K., Buhr, M. P., Knapp, K.,
 Murphy, P. C., Nowick, M., and Williams, E. J.: A method for the airborne measurement of PAN, PPN, and MPAN,
 J. Geophys. Res.-Atmos., 105, 28,943-928,960, 2000.
- Woo, S.-C. and Liu, T.-K.: The absorption spectra and dissociation energies of cyanic acid and some isocyanates, J.
 Chem. Phys., 3, 544-546, 1935.
- Woodrow, J. E., LePage, J. T., Miller, G. C., and Herbert, V. R.: Determination of methyl isocyanate in outdoor
 residential air near metam-sodium soil fumigations, J. Agric. Food Chem., 62, 8921-8927, 2014.
- Woodward-Massey, R., Taha, Y. M., Moussa, S. G., and Osthoff, H. D.: Comparison of negative-ion proton-transfer
 with iodide ion chemical ionization mass spectrometry for quantification of isocyanic acid in ambient air, Atmos
 Environ., 98, 693-703, 2014.
- Yang, X. and Shang, C.: Chlorination byproduct formation in the presence of humic acid, model nitrogenous orgainc
 compounds, ammonia, and bromide, Environ. Sci. Technol., 38, 4995-5001, 2004.
- Yaws, C. L. and Yang, H.-C.: Henry's law constant for compounds in water. . In: Thermodynamic and physical
 properties, Yaws, C. L. (Ed.), Gulf Publishing Company, Houston, TX, 1992.
- Young, P. J., Emmons, L. K., Roberts, J. M., Lamarque, J.-F., Wiedinmyer, C., Veres, P., and VandenBoer, T. C.:
 Isocyanic acid in a global chemistry transport model: Tropospheric distribution, budget, and identification of regions with potential health impacts, J. Geophys. Res.-Atmos., 117, 2012.
- 1055 Zhao, R., Lee, A. K. Y., Wentzell, J. J. B., McDonald, A. M., Toom-Sauntry, D., Leaitch, W. R., Modini, R. L.,
- 1056 Corrigan, A. L., Russell, L. M., Noone, K. J., Schroder, J. C., Bertram, A. K., Hawkins, L. N., Abbatt, J. P. D., and
- Liggio, J.: Cloud partitioning of isocyanic acid (HNCO) and evidence of secondary source of HNCO in ambient air,
 Geophys. Res. Lett., 41, 2014.
- 1059



Figure 1. Plots of the ratio of HNCO concentration at time t, C_i, to the initial concentration, C_o, versus time for a series of flow rates. The solvent was tridecane (C₁₃H₂₈) and 299 K.



 $\begin{array}{c} 1066 \\ 1067 \end{array}$

1068Figure 2. Plots of HNCO loss rate versus the ratio of volumetric flow rate, ϕ , to solution volume, V, for the1069experiment shown in Figure 1, (circles), and the experiment with 10% w/w n-octanol in tridecane at 299 K1070(triangles).



Figure 3. Plot of effective Henry's coefficient of HNCO vs 1/[H+] for the measurements at pH=1, pH=2 and pH=3, and 298 K.



Figure 4. Comparison of effective Henry's coefficients of HNCO measured in this work (blue) with those reported by Borduas et al., 2016, plotted versus pH, according to Equation 4.





Figure 6. Dependence of the Henry's coefficient (H) at a given salt concentration, relative to that with no added salt (H₀) versus NaCl molarity. $R^2 = 0.960$





Figure 7., The Plot of ICN loss rate versus the ratio of volumetric flow rate, ϕ , to solution volume, V, for the 1103 experiment involving the solubility of ICN in water with the small reactor. The line is the least-square fit to the data 1104 $(R^2 = 0.968)$ 1105

- 1106
- 1107



1108 1109 1110 Figure 8. The uptake coefficients of HNCO, CH₃NCO, CICN, BrCN, and ICN as a function of pH for aqueous solution at approximately 298 K. The shaded areas show the range of pHs characteristic of: aerosols (light brown), 1111 cloud/fog water (green), human physiology (red), and ocean surface water (light blue).

Solute	Solvent	Temp. (°K)	pН	Salt, Reactant	H _{eff} , M/atm	Literature H	k^{I} , (x10 ³), sec ⁻¹	Literature k, (x10 ³), sec ⁻¹
HNCO	H ₂ O	279	3.0	-	68 ±7	73ª	0.22	0.24 ^a 0.17 ^b
		286.5	3.0	-	42 ±4	51ª	0.383	0.43 ^a 0.41 ^b
		291	3.0	-	33 ±3	40 ^a	0.657	0.63^{a} 0.72^{b}
		296.5	3.0	-	26 ±2.6	31 ^a	1.02	0.96 ^a 1.32 ^b
		310	3.0	-	12 ±1.2	17ª	4.15	2.6 ^a 5.6 ^b
		298	2.0	-	18 ±1.8		22.2 ± 0.08	
		298	3.0	-	26 ±2.6		1.02 ± 0.13	
		298	4.0	-	59 ±5.9		0.72 ± 0.11	
		298	3.0	0M NaCl	26 ±2.6			
		298	3.0	0.25M NaCl	24.6 ±2.5			
		298	3.0	0.5M NaCl	20.9 ±2.1			
		298	3.0	1.0M NaCl	19.4 ±2.0			
		298	3.0	2.5M NaCl	14.5 ± 1.5			
		292	3.0	0.45M NH ₄ Cl	31.5 ±3.2		1.2	0.005 - 0.015 ^c
	Tridecane (TD)	298	-		1.7 ±0.17		<0.043	
	TD + 10% n- Octanol	283	-		13.2±1.6		0.16 ±0.18	
	TD + 10% n- Octanol	298	-		8.3 ±0.8		< 0.03	
	n-Octanol	298	-		87 ±9		< 0.015	
	n-Octanol	310	-		51 ±5		0.057 ±0.014	
CH ₃ NCO	H ₂ O	298	2.0		1.3 ±0.13		3.2 ±0.3	2.5 ^d , 3.1 ^e
		298	7.0		1.4 ± 0.14		1.9 ±0.6	1.34 ^d , 1.47 ^e
	n-Octanol	298	-		4.0 ± 0.5		2.5 ± 0.5	
	n-Octanol	310	-		2.8 ± 0.3		5.3 ±0.7	
CICN	H ₂ O	299.5	7.0		1.4 ± 0.14	$0.6^{\rm f}, 0.52^{\rm g}$	0.0 ± 0.42	3.03 x10 ^{-3h}
	H ₂ O	273.15	7.0		4.5 ±0.4		0.015 ± 0.016	
	n-Octanol	299.5			1.9 ±0.2		1.3 ±0.4	
BrCN	H ₂ O	296	7.0		8.2 ±0.8		$6.2 \pm 3.7 \text{ x} 10^{-2}$	$1.9 - 9.2 \text{ x} 10^{-2i}$
	H ₂ O	273.15	7.0		32.7 ±3		2.4 ±0.5 x10 ⁻²	
	n-Octanol	297			31 ±3		$9 \pm 2 \times 10^{-2}$	
ICN	H ₂ O	296	7.0		270 ±54		$4.4 \pm 7.6 \text{ x} 10^{-2}$	$\sim 3.4 \text{ x} 10^{-2j}$

Table 1. Summary of solubility and loss rate measurements of HNCO, CH₃NCO and XCN compounds.

a. Calculated from the temperature and pH dependent data reported by Borduas et al., (2016).

- b. Calculated from the temperature and pH dependent data reported by Jensen, (1958).
- c. These were calculated from rates measured at higher pHs, assuming the mechanism is $HNCO + NH_3 => H_2NC(O)NH_2$
- d. From k_{H+} and k_w reported by Williams and Jencks (1974a)
- e. From k_{H^+} for HCl and k_w reported by Castro et al., (1985)
- f. Measured value at 293K, reported by Weng et al., (2011).
- g. Modelled value at 298K, reported by Hilal et al., (2008).
- h. From k_w and k_{OH} reported by Bailey and Bishop (1973).
- i. Estimated from Heller-Grossman et al., 1999, and Vanelslander, et al., 2012.
- j. Estimated from Gerritsen et al., (1993).

Process	HNCO	CH ₃ NCO	CICN	BrCN	ICN
BL deposition	1-2 days	0.5 yrs	yrs	0.5 yrs	5-10 days
Aerosol dep.	6-12 days	2 -4 months	yrs	0.6 yrs.	8 days
In-cloud rxn	2-6 hrs	2 months	10-20 wks.	1-3 wks	1-3 days

Table 2. Estimates of HNCO, CH₃NCO, and XCN compounds against loss due to heterogeneous processes.

Compound	Temperature	LogKow	
HNCO	298	0.64	
	310	0.63	
CH ₃ NCO	298	0.49	
CICN	299.5	0.13	
BrCN	297	0.61 ^a	
HC(O)OH	298	-0.54 ^b	
CH ₃ NO ₂	293	-0.33 ^b	
HCN	?	0.66 ^c	
CH ₃ CN	298	-0.34 ^b	

Table 3. Octanol/Water partition coefficients for HNCO, CH₃NCO, ClCN, BrCN and related compounds.

a. based on extrapolated H_{H2O} at 297K b. (Sangster, 1989) c. (EPA, 1989)