

Author’s Response to interactive comments by Anonymous Referees #1 and #2 on “Estimating background contributions and U.S. anthropogenic enhancements to maximum ozone concentrations in the northern U.S.” by David D. Parrish

The author is grateful for the referees’ thoughtful comments regarding this paper. Responses to all comments are included here, and where appropriate the manuscript has been revised as described herein, and indicated in the “tracked changes” manuscript copy at the end of this response. Many of the referees’ comments are incorrect in some respects, or are not germane to the issues that this manuscript investigates; in such cases detailed responses are included below, and some additions have been made to the manuscript to more fully discuss these issues.

I believe that this manuscript is very important and deserves publication. U.S. policy makers must set national ambient air quality standards (NAAQS) for criteria air pollutants including ozone. A major uncertainty they face is the contribution to measured ambient concentrations made by transported background ozone. It is possible that a standard could be set at an ozone concentration below that background contribution, at least in some regions of the country, and thus be impossible to meet through control of U.S. precursor emissions. If it is also possible, indeed likely, that the background contribution varies substantially over the country, so any standard would be more difficult to meet in some regions of the country than others. Currently, policy makers must rely on estimates of the background ozone contribution calculated by models of atmospheric transport and chemistry. However, these estimates vary widely among different models, and are recognized to be so uncertain that their utility to policy makers is limited. The value of this paper is that it presents an observationally based estimate of the background ozone contribution that policy makers can compare and contrast with model estimates. As noted by the referees (and acknowledged and discussed in the paper) this observationally based approach is not perfect, but all analyses suggest that the results are more accurate than the currently available model results, and that the paper deserves publication. The comments of each referee are reproduced below (*in blue italics*) followed by my response (in plain text).

Referee #1 Comments

For reference, I have numbered the paragraphs from the first referee’s comments; when a paragraph included multiple comments, those are addressed separately with numbers and letters (e.g., paragraph 3 is divided into 3a and 3b).

1) This paper fits long-term trends of ozone design values (ODVs) in the northeastern and rural western US to exponential decay forms with a pre-derived decay scale (21 years) from prior work for Los Angeles, and infers US background ODVs from the asymptote. It concludes that the ODV in the northeastern US is 45.8 ppb, points out that it represents a large fraction (65%) of the current NAAQS over which the US has no regulatory authority, and that it is much larger than models implying that models have large errors.

This summary is mostly accurate, if perhaps overly concise. However, the last phrase is not accurate; the response to comment 5 below discusses the issues that imply large model errors.

2) I have a number of problems with this paper, and not sure that they can be fixed, so ACP may need to do arbitration or seek another reviewer. As I see it, there is no reason that the 21-year ODV decay time scale from LA would apply to other regions. The rural western US is mostly flat, and the northeastern US has a very different trend history initiated with the early 2000s NOx SIP Call. In fact, it seems from Figure 6 that the trend in the Northeast since 2000 could be fit to a linear decrease just as well as to an exponential decrease, and the linear decrease would imply a zero background ODV – which does not make sense of course

but makes the point that there is no robustness to the estimate of background ODV presented here from the asymptote to the exponential decay curve. As the paper points out, a 10% change in the decay time scale would lead to a 5 ppb change in background ODV – but there is much more than 10% leeway to the fit in Figure 6.

The referee correctly identifies one of the more difficult aspects of the analysis presented in this paper. In the northeastern U.S., the long-term decreases of ozone design values have not continued in a consistent manner long enough for all three parameters of Equation 1 to be precisely extracted from the available data sets. Thus, applying the exponential decay time from the entire southern California region (not just LA) is a convenient, and as we demonstrate below, accurate approximation. Section 4.2 of the original manuscript did discuss reasons that the long-term ozone changes may be similar to those in southern California; this discussion is retained in the revised manuscript. The northeastern U.S. does have a mixed trend history before 2000, but ODVs over much of this region have decreased similarly to those in California. Figure 12a illustrates the great value of comparing the two most populous U.S. urban areas in this manner.

The referee suggests that a linear fit to the data would be appropriate. Indeed, the estimation of background ODVs in the northeastern U.S. can be based on linear fits to the long-term changes as illustrated in Figure 1. Importantly, a linear decrease by itself does not imply a zero background ODV, or any other specific value; rather it implies that the ODVs can decrease




Figure 1: Comparison of linear fits (solid black lines) and exponential fits (colored lines) to the maximum observed ODVs in the New York City urban area (from Figure 12 of the original paper), and all of the ODVs observed in Vermont (from Figure S7 of the original Supporting Information). The dotted lines are extrapolations of the linear fits, with the red circle indicating their intersection at an ODV value of 48.5 ppb.

indefinitely to increasingly negative ODVs – which indeed does not make sense. However, two linear fits to different time series of ODVs in a region with a uniform background ODVs, do imply a particular background ODV value, which is given by the intersection of the extrapolations of the two linear fits. This follows because the background ODV does not depend upon the magnitude of the anthropogenic enhancement of ODVs, so both time series should meet at that intersection when U.S. emissions of anthropogenic ozone

precursors are reduced to zero. For two extreme northeastern U.S. data sets (the maximum ODVs observed in the New York City urban area, and all of the ODVs observed in the much more rural state of Vermont) the background ODV from the intersection of the linear fits (48.5 ppb) is in reasonable accord with the asymptotes approached by the two exponential curves for New York City and Vermont (45.8 and 45.6 ppb, respectively). The important point of this figure and discussion is not that the linear fits are realistic (they are not), but rather that the estimation of the background ODV is not strongly dependent on the validity of the exponential fits, because they can be estimated nearly as well from linear fits. Thus, contrary to the referee’s assertion, the estimates of background ODVs presented in the manuscript are robust.

Importantly, there is an independent analysis that can also estimate the U.S. background ODVs without the assumption of any specific functional form for the time dependence of the ODV enhancements. Section 2.3 of Parrish et al. (2017) discusses this different, somewhat more general approach, which is based upon correlations between ODVs recorded in different regions with substantially different anthropogenic enhancements. The time series of ODVs from one region is selected as a reference, and other time series are linearly correlated with this reference. A different assumption does underlie this approach, namely that the

U.S. background ODVs vary in a similar manner, but not necessarily as an exponential decrease, in all regions under consideration. Given this assumption, the U.S. background ODV for a region is taken as the ODV where the line derived from a linear correlation of that region's ODVs with those of the reference region intercepts the 1:1 line; at that point the extrapolated ODVs from the two time series would be equal, which is necessarily at that regionally uniform U.S. background ODV. Section 2.3 of Parrish et al. (2017) show that the results of this approach for seven southern California air basins are nearly identical to the results from the fits to Equation 1.

The results from applying this second approach to the northeastern U.S. are shown in Figures S11-S18 of the revised Supplement for the 13 regional data sets given by the black lines in Figure 8 of the revised manuscript. Here the time series of maximum observed ODVs in the New York City urban area (included in Figure 12a of the revised manuscript) is selected as the reference, because they are some of the largest ODVs recorded in the northeastern U.S., and because after 2000 this time series closely follows an exponential decrease with little interannual variability. Figure S18 of the Supplement illustrates all of the linear regressions for the 13 regional data sets, and Table S3 summarizes the results. These results have significant variability (25 to 62 ppb), because the correlation slopes are near unity, so it is difficult to determine the intercept of the linear correlation with the 1:1 line. However, the average of the derived background ODVs (49.2 ± 3.9 ppb for standard linear regressions and 42.5 ± 5.7 ppb for reduced major axis regressions, where 95% confidence limits are indicated) bracket the result (45.8 ± 1.7 ppb) reported in the manuscript for the exponential fits, and neither average is statistically significantly different from that result. The analysis presented in this comment demonstrates that the fitting of long-term trends of ODVs to Equation 1 of the text does not compromise the accuracy of the results presented in the discussion paper.

Changes incorporated in revised manuscript: Section 3.2.3 “Alternative approach for estimating U.S. background ODV in northeastern states”, has been added to the manuscript; it contains much of the above discussion describing this alternate approach. The results are included in the Supplement - Figures S11-S18 and Table S3. The agreement between the two analysis approaches is now briefly discussed in Section 4.2 “Possible shortcomings of the analysis”.

3a) There is also no physical rationale for a single time scale in the exponential decay of ODV, and in the absence of such a rationale any interpretation or extrapolation can be very foolish. The decrease of ODV in the Northeast is thought to be driven mainly by US NO_x emissions, which have decreased linearly since 2000 according to EPA although Jiang et al. (PNAS 2018) suggest that they have been flat since 2009 – in any case, I don't see how either scenario would drive an exponential decay of ozone. Even if the response of ozone to NO_x emissions was exponential, we would need a sum of exponentials to describe the ozone decrease because different anthropogenic NO_x sources have decreased at different rates, and NO_x emissions from fertilizer use and small industries have not decreased at all according to EPA. The effect of anthropogenic emissions on ozone is thus much more complicated than can be explained with a single exponential, and even if one can achieve such a fit to the data there is no rationale for extrapolation without understanding.

There is a strong rationale for a single time scale in the exponential decay of ODVs, but that rationale is more mathematical than physical. In the entire southern California region considered by Parrish et al. (2017), a single time scale in the exponential decay of ODVs captured 98.4% of the total variability in the maximum ODVs in 7 air basins, as indicated by the $r^2=0.984$ for the correlation between the observed ODVs and the values predicted from the single time scale in the exponential decay of Equation 1 (see Figure 5 of Parrish et al., 2017). Such strong correlations are only rarely encountered in geophysical research. In 4 of the 7 air basins, this exponential decay covered the entire 36-year 1980-2015 time series of ODVs. In the present manuscript, a single time scale in the exponential decay of Equation 1 captures 89% of the total variability in the maximum ODVs in the 8 northeastern U.S. states over the 18-year 2000-2017 period. The shorter time period northeastern U.S. accounts for the smaller r^2

found in the present work. This strong agreement between the fits to Equation 1 and the observations in these two diverse U.S. regions is the primary rationale for using a single time scale in the exponential decay, and this strong agreement clearly demonstrates that the effect of anthropogenic emissions on ozone is not more complicated than can be explained with a single exponential. It would be of interest to understand the origin of this strong agreement, but until robust model calculation can reproduce the observed long-term trends in ODVs such an understanding is not available.

Changes incorporated in revised manuscript: The strong agreement between the fits to the exponential decrease and the observations was already discussed in Section 4.2 “Possible shortcomings of the analysis”; no further discussion has been added.

3b) Considering that NO_x emissions from fertilizer use and small industries have not decreased, and that some VOC emissions have not decreased (reference in the paper to McDonald), one must conclude that the background ODV derived in this paper is biased high, possibly by a large amount.

The referee is correct that the background ODVs derived in this paper can be biased high. This is because the fits to Equation 1 separate the ODV time trend into two contributions, one time independent (y_0) and one time dependent ($A \exp\{-(\text{year}-2000)/t\}$). Any contribution to ODVs from emissions that have not decreased would be counted in the time independent term, and thus bias its estimate high compared to the true U.S. background ODV. However, generally speaking there are no indications that this bias is significant, and we have some checks on the magnitude of the potential bias. For example, the same y_0 value is found throughout the northeastern U.S. in regions that are highly industrialized with no significant agriculture (i.e., the major metropolitan areas), rural areas with significant agriculture (e.g., the Hudson River Valley), and forested regions with little agriculture. If NO_x emissions from fertilizer use or small industries were important, their importance is expected to be reflected in spatial variability of the derived y_0 values. A similar argument was made by Parrish et al. (2017), where the same y_0 value was found for 6 of the 7 California air basins, where the degree of industrialization and land use varied from the Los Angeles urban area, to the rich agricultural areas of the San Joaquin Valley, and to the sparsely populated Mojave desert. Some significant examples of derived y_0 values that are biased high have been identified and discussed. In the present paper, the y_0 values derived for the most densely populated areas (see discussion of Figure 8) are higher than expected from comparison with nearby areas; this is discussed as possibly caused by emissions of the volatile chemical products discussed by McDonald et al. (2018). Parrish et al. (2017) discuss the source of the high bias of the derived y_0 value for the Salton Sea air basin (NO_x emissions from the highly fertilized agriculture of California’s Imperial Valley). Beyond these limited examples, there is no indication of a significant, general high bias in the results.

Changes incorporated in revised manuscript: This possible bias was already discussed in Sections 4.1 “Implications of the results” and 4.2 “Possible shortcomings of the analysis”; no further discussion has been added.

4) Indeed, a punch line of the paper is that the background ODVs inferred from the exponential fit are 65-90% of the NAAQS. That seems like a big fraction, but the background ODV estimates are biased high (see above). In addition this is misleading, considering that the ODV is depleted under polluted conditions. In the northeastern US in particular, the ozone background is highest in subsiding northerly flow, whereas the ODV exceedances are under stagnant conditions with southerly flow where background ozone is much lower.

It is difficult to respond to this comment, as several issues are combined. First, there is no evidence that the background ODV estimates are generally biased high by a significant amount, although there are some specific instances of biases that are discussed, and their influence avoided (see response to comment 3b). It is not clear what the referee means by the statement “that

the ODV is depleted under polluted conditions”; the ODV is defined as the 3-year average of the annual fourth-highest MDA8 ozone concentration, and these highest MDA8 ozone concentrations do occur under polluted conditions. If they are the highest concentrations, how can they be depleted? The analysis presented in the paper makes no assumption about, and does not depend upon, any particular flow regime, whether subsiding northerly flow or stagnant conditions with southerly flow. I can find no relevance in this comment.

Changes incorporated in revised manuscript: No further discussion has been added.

5) As the paper points out, model estimates of background ozone are much lower than what is presented in this paper. The paper attributes this to error in the models. It is fair to say that there is a ± 10 ppb uncertainty in model estimates, as quoted in the paper. But that uncertainty is not a bias, whereas the background estimate in this paper is unarguably biased high. The paper does point out to some extent the uncertainties in its estimate of background ODVs, but that disappears in the abstract where the message is about the high contribution of the background to the NAAQS and how the models need to be corrected.

Model estimates of background ozone are both higher and lower than the result presented in this paper. The 45.8 ± 1.7 ppb estimate of the background ODV derived in our analysis for the northeastern US is smaller than the model results illustrated in a recent assessment of background ozone over the U.S. (Jaffe et al., 2018); the color scale of their figure 3 indicates a U.S. background ODV in the northeastern U.S. in the range of 50-60 ppb. Our paper does cite other model calculations that give results that average 4 to 12 ppb lower than the 45.8 ppb estimate.

The uncertainty in model estimates of ODVs is actually significantly larger than ± 10 ppb. The paper does have the sentence: “Jaffe et al. (2018) estimate an uncertainty in modeled seasonal mean U.S. background ozone of about ± 10 ppb, with greater uncertainty for individual days (such as those that define the ODV), and Guo et al. (2018) find biases as high as 19 ppb in modeled seasonal mean MDA8 ozone.” Much of the uncertainty in model results is indeed due to biases. The referee’s reference to a high bias in our analysis is discussed in the response to comment 3b) above.

It is clear from the large systematic, average differences among the results from different models that models have large errors, and that to be more useful to policy makers, the models do need to be improved.

The abstract does focus on the high contribution of the background to the NAAQS and the need for further model improvement, since those are among the primary conclusions of the paper. The abstract also states “The approach used in this work has some unquantified uncertainties that are discussed.”

Changes incorporated in revised manuscript: The last sentence of the abstract has been changed to read “Further model improvement is required until their output can accurately reproduce the time series and variability of observed ODVs. Ideally, the uncertainties in the modeling and observational based approaches can then be reduced through additional comparisons.”

6) Aside from these basic issues of scientific content, I found the paper to be much longer than it needs to be. Figures 1 and 2 show the Pacific Northwest and the Midwest but these then drop from the radar screen, why even bother? Descriptive discussions of population, topography, etc. don’t seem necessary. There’s a lot of chattiness and repetition.

The description of the Pacific Northwest and the Midwest are necessary for two reasons: first, to show that the ozone design values measured throughout the entire tier of northern U.S. states follow a common pattern (i.e., the magnitude of ODV enhancements correlate with population density, and have decreased in at least a qualitatively similar manner), and second, to justify the focus on two of the four regions, namely on the two regions presenting the greatest extremes. This paper is intended

for a journal with an international readership, so I believe that descriptive discussions of population and topography are needed for readers that are not familiar with these U.S. details.

Changes incorporated in revised manuscript: The paper has been revised to remove chattiness and repetition where they could be found.

Referee #2 Comments

The analysis is designed to separately quantify the U.S. background ozone design values (ODVs) and the enhancements of the ODVs above that background contribution due to U.S. anthropogenic precursor emissions. The U.S. background ozone design value is assumed to be the maximum ozone DV that would exist in these regions in the absence of U.S. anthropogenic precursor emissions. The US background and US anthropogenic increment are derived from a simple exponential function, analogous to the function derived for California subregions in Parrish et al. (2017).

Although the idea of a simple model to describe design value behavior is appealing, there are several problems with this approach.

(1) The simple exponential function has been applied to separate the contributions of anthropogenic US precursor emissions to the ozone design values from the contributions by US background ozone (i.e., ozone that would be present in the absence of US anthropogenic precursor emissions). This formulation of the ozone problem is based upon a chemical transport modeling definition of US background ozone; the US background ozone can be estimated by “zeroing-out” US anthropogenic emissions in a chemical transport model. In areas far from the Pacific Coast, where US background concentrations enter the country, it is difficult to see how this simple observational model can untangle the interactions between US biogenic emissions (part of the background) and US anthropogenic emissions (part of the anthropogenic component). It is not at all clear that the asymptotic value approached by the exponential equation in this manuscript represents US background, or some mixture of US background (e.g., biogenic VOCs) combined with an especially persistent US anthropogenic (NO_x) component that hasn't yet been substantially reduced by control strategies.

The referee speculates that there may be “an especially persistent US anthropogenic (NO_x) component that hasn't yet been substantially reduced by control strategies” that can confound the analysis presented in the manuscript. As discussed in the paper, this is a concern. However, there is no evidence for such a component with a NO_x emission magnitude that is comparable to the well-known emission sectors (mobile, industrial and power plant sources) that have been effectively reduced by control strategies. The possible influence of such an emission component must be kept under consideration, but in that absence of any evidence of such a component, there is no justification for rejecting the results of the analysis presented in the manuscript.

Notably, the emission inventories typically used for regional photochemical modeling do not include any such “especially persistent US anthropogenic (NO_x) component”. If such a component did exist, photochemical modeling results also would be biased. Such a possibility further emphasizes the value of the present observationally based estimate of U.S. background ozone, so that biases in either the model results or the observationally based estimate can be understood through comparison of results, and the results obtained from each approach improved.

Changes incorporated in revised manuscript: The manuscript has already discussed biases due to anthropogenic precursors that have not been reduced by emission controls. No further discussion of this issue has been added.

(2) Interannual variation of ozone data is smoothed because three years of data are averaged together to get a design value. The rationale for using the three-year average of the fourth high is that attainment of the ozone standard is linked to a three-year average, and hence, it is important to study the behavior of this somewhat unwieldy metric in order to reach policy-relevant conclusions.

But a design value is defined for a specific metropolitan area. The highest three-year average of the fourth high at any monitoring site within the metro area is the design value. Since the analysis does not examine ODVs for individual metro areas, or select the fourth high for each metro area for each year, the ODVs described in the paper are not actually the ODVs used in regulatory applications. It can be argued that this distinction is scientifically trivial, but in this case we are discussing policy, not science, so the distinction is important. The author could redefine the regions according to the EPA's definition of nonattainment areas to match the policy definition.

The referee's description of ozone design values is not entirely complete. As discussed in the manuscript, an ODV is defined for each monitor in the U.S. The ODVs analyzed in this paper are indeed the ODVs that are considered in regulatory applications. EPA's process for defining nonattainment areas (rural as well as metropolitan areas) involves many considerations, including the ODV for the area; the area ODV is simply the highest ODV of those recorded at any of the sites within the area during each year. Since the highest site ODVs are included in our analysis, we are analyzing the ODVs actually used in regulatory applications. To assuage the referee's concern, Figure 2 illustrates the analysis for nonattainment area ODVs, and Table 1 gives the derived parameters. (The four nonattainment areas that lie entirely within the northeastern states are included; the fifth nonattainment area is only partially in the area under consideration, so it is not included.) These nonattainment areas are described in the manuscript. As can be seen from comparing Table 1 below with Table 2 of the manuscript (and properly considering the




Figure 2: Time series of nonattainment area ODVs (solid symbols) and the ODVs reported from all monitoring sites (open symbols) in the four nonattainment areas included within the northeastern U.S. states examined in our analysis. The curves are fits of Eq. 1 to respective colored symbols, or to all ODVs (black curves) in the areas.

Table 1. Results of least-squares fits to Eq. 1 illustrated in Figure 1; RMSD indicates the root-mean-square deviation between the observed ODVs and the derived fit.

Nonattainment area/sites	y_0 (ppb)	A (ppb)	RMSD (ppb)	years fit
NY-NJ-LI-CT area/ODVs	63 ± 12	40 ± 16	3.0	2000-2017
NY-NJ-LI-CT/all sites	53 ± 4	42 ± 6	5.8	2000-2017
Greater CT area/ODVs	59 ± 12	36 ± 17	3.2	2000-2017
Greater CT/all sites	54 ± 7	39 ± 10	4.1	2000-2017
Jamestown, NY area/ODVs	45 ± 14	51 ± 19	3.7	2000-2017
Jamestown, NY/all rural upwind sites	42 ± 7	50 ± 10	5.1	2000-2017
Dukes Co, MA/all sites	44 ± 9	52 ± 13	3.2	2000-2017

parameter confidence limits included in the tables), the analysis presented in the manuscript is directly relevant to the nonattainment area ODVs. The results for the NY-NJ-LI-CT and Greater CT nonattainment areas in Table 1 are in close accord with the Connecticut and New York/maximum results, respectively, given in Table 2 of the manuscript. Similarly, the results for the Jamestown, NY and Dukes Co, MA nonattainment areas in Table 1 are in close accord with the New York/rural upwind and Massachusetts/coastal results, respectively, given in Table 2 of the manuscript. The analysis for the nonattainment areas is entirely consistent with the analysis and discussion included in the manuscript; nothing new would be gained by including this additional analysis in the manuscript.

Changes incorporated in revised manuscript: No further discussion of this issue has been added to the manuscript.

But the statistical analysis would still be somewhat clumsy; the three-year averages smooth out much of the interannual variability, and can cause autocorrelation issues, as the author notes. There are, however, other metrics just as relevant to policy as the ODVs used in this study. A better metric for individual monitoring sites would be the 4th high (98th percentile) maximum daily eight-hour ozone concentration for each year at each site. If the fourth-high/98th percentile metric for each year at each site were analyzed, there would be no overlap among years, eliminating problems with autocorrelation and excessive smoothing of interannual variations among years, yet the analysis would be at least as relevant to regulatory status as the current analysis. The physical interpretation of the data would be simplified as well, because the metric itself would be more closely tied to the observations of a single site and single year, instead of being smeared over three years.

Using a different metric would help resolve an issue related to the smoothing of interannual variation. The author asserts that the simple exponential model of ODV trends has achieved a degree of success in describing the variation, based upon the confidence intervals. These confidence intervals have been modified to account for covariance due to lack of independence among ODVs. But the interannual variation would be larger if the analysis had been performed on annual 98th percentiles rather than ODVs for each site, and it is unclear whether the modification of confidence intervals to account for covariance also accounts for the reduction of interannual variation. The results would be more compelling if the interannual variation had not been shaved down by using three-year running averages.

The preceding two paragraphs criticize the statistical fitting technique utilized in the manuscript; however, the referee's discussion is not entirely correct. The goal of the statistical analysis is to extract the systematic, long-term change in a set of observed ODVs (or fourth-high/98th percentile) as accurately as possible, given the interannual variability about those long-term changes. Importantly, no statistically significant information regarding the long-term change is lost by working with 3-year means (i.e., the ODVs) rather than annual mean data (i.e., fourth-high/98th percentile). A qualitative explanation for this can be

given. Deriving 3-year means from annual data involves an averaging process, which minimizes the sum of the squares of the deviations of the annual data from the derived 3-year means. The fitting procedure employed in the analysis to derive the long-term change minimizes the sum of the squares of the deviations of the 3-year ODVs from the derived long-term change. The final result is independent of whether the sum of the squares of the deviations is minimized in two steps (3-year mean calculation followed by the fit to the long-term change), or in one step (extracting the long-term change directly from the higher frequency annual data). We work with 3-year mean ODVs because they are of most policy relevance, and we have properly dealt with the autocorrelation issues. And the confidence limits derived from the procedure for fitting the long-term changes properly account for the uncertainty arising from interannual variations, whether that fit is to 3-year or annual means.

Additionally, working with annual mean data (i.e. fourth-high/98th percentile) would worsen a separate autocorrelation concern. Interannual meteorological and climate variations can drive differences in ozone photochemical production and transport, differences that can persist over multiple years, so that annual mean data may be autocorrelated. Working with 3-year mean ODVs reduces the influence of this additional autocorrelation.

Changes incorporated in revised manuscript: No further discussion of these issues has been added to the manuscript.

(3) Increasing the interannual variation, however, would probably worsen another issue: the inability of the model to converge on a solution for the three model parameters. As the author notes on page 7, lines 10-16, the shorter data record for the northern regions appears to be preventing estimation of the three parameters of the exponential function. To resolve this issue, the author has assumed that one of the parameters can be set at the same value as derived for California. As the author notes, the value of $\tau = 21.9$ years derived from California implicitly assumes that control strategies have produced approximately equal relative reductions in anthropogenic ozone enhancements throughout the country. This assumption is questionable, and the results for the northeastern states seems to show that it is unwarranted.

As discussed in my response immediately preceding this comment, no statistically significant information regarding the long-term change is lost working with 3-year means. Increasing the interannual variation by working with annual means would not worsen the uncertainty in deriving the three model parameters; the greater number of independent data available for the fitting procedure would closely compensate for the larger interannual variation.

The referee is correct that the assumption of $\tau = 21.9$ years is questionable, and that question is addressed in detail in my response to Comment 2 of referee #1. Notably, the referee does not state in what way “the results for the northeastern states seems to show that it (the assumption) is unwarranted”; thus I am unable to respond to the last part of this comment.

Changes incorporated in revised manuscript: No further discussion of this issue has been added to the manuscript.

(4) Table 2 shows the derived values of y_0 (US background ozone) and A (US anthropogenic component) for subsets of monitoring sites. The values of US background ozone for low altitude sites vary from 41 ± 10 ppb in suburban Massachusetts to 61 ± 6 ppb in coastal Connecticut. This is a large variation over a short distance for a value that is supposed to reflect relatively unvarying background ozone.

The referee correctly points out that the derived y_0 values in Table 2 of the manuscript for the low altitude sites vary from 41 ± 10 ppb in suburban Massachusetts to 61 ± 6 ppb in coastal Connecticut and suggests that this variation is large. This large variation arises from two sources. First, as reflected in the 95% confidence limits given in the table, there is statistical uncertainty in deriving the parameter values from the fits of Equation 1 to the ODVs; the confidence limits for the two extreme results that the referee quotes indicate that the statistically significant portion of the variation is likely not as large as the range in the extreme

values might indicate. Second, as discussed in the paper, some of the variation does arise from real departures of the derived y_0 values from the true U.S. background ozone. Figure 2 below illustrates an analysis of the distribution of the derived y_0 values that allows these two sources of variation to be examined separately. The ordinate scale of this figure is designed so that a normal distribution will lie on a straight line. In this figure, 13 of the 17 derived y_0 values do define a normal distribution with a median of 47.7 ppb and a standard deviation of 4.5 ppb. We believe that the median of the 13 values accurately reflects U.S. background ozone and the standard deviation reflects the uncertainty in determining each y_0 value. This result is consistent with the 45.8 ± 1.7 ppb estimate of the U.S. background ODV given in the manuscript. The variation of the results from these 13 groups of sites is consistent with the statistically derived confidence limits given in the manuscript. The highest 4 of 17 derived y_0 values in Figure 2 define a high value tail; these are the 4 y_0 values that are discussed separately in the manuscript. Figure 2 nicely illustrates the value of our analysis based upon the simple model encapsulated in Equation 1 – the large majority of the results provide a robust estimate of the regional U.S. background ozone, and deviations from a uniform result point toward issues that deserve further investigation.




Figure 3: Cumulative probability plot of the y_0 determinations listed in Table 2 of the manuscript. The line shows a linear regression fit to the open points, which defines a normal distribution with the median and standard deviation annotated in the figure. The four points at the higher y_0 values correspond to the colored curves in Figure 8 of the manuscript.

Changes incorporated in revised manuscript: Figure 2 is now included as Figure S10 in the supplement, and a short description of this analysis has been added to Section 3.2.1 of the manuscript.

One interpretation is that the Connecticut “background O3” includes a lot of ozone generated in the New York City area, and that therefore the simple exponential model cannot determine US background. The author chose a different interpretation, and re-set the US background to a lower value, which increased the US anthropogenic component to more acceptable levels. This portion of the analysis is not convincing, and seems to be an attempt to compensate for the simple model’s shortcomings. In fact, it is essentially an admission that the original simple model cannot be used to distinguish between US background ozone and US anthropogenic ozone.

This interpretation suggested by the referee cannot be correct. Connecticut ODVs certainly do reflect a lot of ozone generated in the New York City area. However, this New York City generated ozone has decreased over the years; thus, our analysis would properly include it in Connecticut’s time varying term ($A \cdot \exp\{-\text{(year-2000)}/t\}$), not in Connecticut’s constant y_0 term. The simple exponential model can indeed determine U.S. background ozone, within the caveats thoroughly discussed in the manuscript.

The referee does not correctly describe the reasoning in our manuscript. We followed two clearly distinct steps. The first step is the purely mathematical fitting of Equation 1 to the observed ODVs. The success of this step is judged by the good agreement between the fits and the observations. The second step is the physical interpretation of the A and y_0 parameters. As discussed in the manuscript and illustrated in Figure 3 above, the derived y_0 values provide substantial information regarding the U.S. background ozone, but we do not argue that the two quantities are identical in all situations. Hence, the four values in the high value tail, which includes the Connecticut y_0 value discussed by the referee, require separate discussion. In the physical interpretation of the A and y_0 parameters we do not re-set the U.S. background ozone to a lower value; rather we discuss why the derived Connecticut y_0 value is not equal to the U.S. background ozone. This is not an attempt to compensate for the shortcomings of the simple model; rather it is an identification of an issue identified by the simple model that deserves further investigation. It is not an admission that the original simple model cannot be used to distinguish between US background ozone and US anthropogenic ozone; in the majority of the ODV time series investigated the model does provide a robust distinction, and the few exceptions identify interesting issues.

Changes incorporated in revised manuscript: No further discussion of this issue has been added to the manuscript.

Ultimately, I have concluded that the assertions claiming that US background and anthropogenic increment can be derived from the simple exponential function are not compelling, especially for the northeastern states. It is possible that the analysis could be re-worked, by changing the ozone metric to the annual 98th percentile, but the failure of the method to derive the three parameters of the model even with three-year running averages, and its inability to distinguish between US background and US anthropogenic in the northeastern states suggests that this simple approach is flawed for the regions to which it has been applied in this study. I thought that the study was interesting, and the author did a commendable job in explaining the uncertainties and possible shortcomings of the approach. This admirable transparency in describing the methods is worthy of emulation, but I do not think the study should be published in its present form.

The author appreciates the referee's positive comments. However, as discussed in detail above and in the response to comment 2 of referee #1, the referee's comments do not raise valid concerns regarding the analysis presented in the manuscript. The referee's overall recommendations stated here are not justified.

The critical comments provided by the referees have sharpened the analysis presented, and formulating the above responses has (hopefully) clarified the discussion. All of the objections raised by the referees have been answered. This is an important paper; please consider this resubmitted manuscript for publication.

Estimating background contributions and U.S. anthropogenic enhancements to maximum ozone concentrations in the northern U.S.

David D. Parrish^{1,2,3}

¹ Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, USA

² NOAA/ESRL Chemical Sciences Division, Boulder, Colorado, USA

³ David.D.Parrish, LLC, Boulder, Colorado, USA

Correspondence to: David D. Parrish (David.D.Parrish@noaa.gov)

10 **Abstract.** U.S. ambient ozone concentrations have two components: U.S. background ozone and enhancements produced
from the country's anthropogenic precursor emissions; only the enhancements effectively respond to national emission
controls. We investigate the temporal evolution and spatial variability of the largest ozone concentrations, i.e., those that
define the ozone design value (ODV) upon which the National Ambient Air Quality Standard (NAAQS) is based, within the
northern tier of U.S. states. We focus on two regions: rural western states, with only small anthropogenic precursor
15 emissions, and the urbanized northeastern states, which include the New York City urban area, the nation's most populated.
The U.S. background ODV (i.e., the ODV remaining if U.S. anthropogenic precursor emissions were reduced to zero) is
estimated to vary from 54 to 63 ppb in the rural western states, and to be smaller and nearly constant (45.8 ± 1.7 ppb)
throughout the northeastern states. These U.S. background ODVs correspond to 65 to 90% of the 2015 NAAQS of 70 ppb.
Over the past two to three decades U.S. emission control efforts have decreased the anthropogenic ODV enhancements at an
20 approximately exponential rate with an e-folding time constant of ~ 22 years. These ODV enhancements are small in the rural
western states (2.4 ± 1.2 ppb in 2000), with much larger state maximum ODV enhancements (~ 35 -64 ppb in 2000) in the
northeastern states. The U.S. background ODV contribution is significantly larger than the present-day ODV enhancements
due to photochemical production from U.S. anthropogenic precursor emissions in the urban as well as the rural regions
investigated. Forward projections of past trends suggest that average maximum ODVs in northeastern U.S. will drop below
25 the NAAQS of 70 ppb by about 2021, assuming that the exponential decrease of the ODV enhancements can be maintained
and the U.S. background ODV remains constant. This estimate is much more optimistic than in the Los Angeles urban area,
where a similar approach estimates ~ 2050 for the maximum ODV to reach 70 ppb (Parrish et al., 2017). The primary reason
for this large difference is the significantly higher U.S. ODV background (62.0 ± 2.0 ppb) estimated for the Los Angeles
urban area. The approach used in this work has some unquantified uncertainties that are discussed. Models can also estimate
30 U.S. background ODVs; some of those results are shown to correlate with the observational estimates derived here (r^2 values
for different models are ~ 0.31 to 0.85), but are on average systematically lower by 4 to 12 ppb. Further model improvement

is required until their output can accurately reproduce the time series and spatial variability of observed ODVs. Ideally, the uncertainties in the model and observational based approaches can then be reduced through additional comparisons.

1 Introduction

The U.S. has a long-standing air quality problem associated with elevated ozone concentrations (e.g., NRC, 1991).

5 | Fortunately, this problem has been greatly improved over the past 3 to 5 decades, particularly in urban areas. For example, through the 1960s and 1970s the Los Angeles urban area (i.e., California's South Coast Air Basin – SoCAB) endured maximum 1-hr average and maximum daily 8-hr average (MDA8) ozone mixing ratios that exceeded 500 and 300 ppb, respectively (ppb = nmole ozone per mole air) (Parrish and Stockwell, 2015). The National Ambient Air Quality Standard (NAAQS) is based on the ozone design value (ODV), which is defined as the 3-year average of the annual fourth-highest
10 | daily maximum 8-hour average (MDA8) ozone concentration; in 2015 the NAAQS was lowered, now requiring that ODVs not exceed 70 ppb. A fit to the long-term trend of the maximum ODVs recorded in the SoCAB indicates that these highest ozone concentrations decreased from 289 to 102 ppb over the 36-year, 1980 to 2015 period (Parrish et al., 2017). This decrease demonstrates that controls on U.S. ozone precursor emissions have been remarkably effective in reducing maximum ambient ozone concentrations. However, much additional emission reduction efforts are required to reach the
15 | NAAQS of 70 ppb. A critical question has relevance to policy development for managing U.S. ozone concentrations: What is the limit to which ODVs can be reduced by controlling U.S. anthropogenic emissions? One goal of this work is to provide an approximate answer to this question.

Both natural and anthropogenic processes interact to determine the temporal and spatial distribution of surface ozone concentrations in both urban and rural areas. Thus, even if U.S. anthropogenic emissions of ozone precursors were
20 | completely eliminated, ambient ozone concentrations throughout the U.S. would be non-zero due to contributions from natural sources of ozone, enhanced by anthropogenic contributions from other countries. Parrish et al. (2017) estimate that this remaining ODV (denoted as U.S. background ODV) would be 62.0 ± 1.9 ppb in the Los Angeles urban area. This contribution is the limit to which the ODVs can be reduced by U.S. emission controls alone; it is so large that there is little margin for enhancement of ambient ozone concentrations by photochemical production from U.S. anthropogenic precursor
25 | emissions before the NAAQS of 70 ppb is exceeded. In this paper we extend the analysis of Parrish et al. (2017) to the northern tier of U.S. states, which includes the most populated U.S. urban area (the New York City metropolitan area), as well as vast, sparsely populated regions.

The northern U.S. states can be conveniently divided into four regions (Figure 1): three Pacific Northwest states, three rural western states, three Midwest states, and eight northeastern states. The northeastern states are the most heavily populated,
30 | with the New York City; Boston, Massachusetts; and Providence, Rhode Island urban areas. Two regions are moderately

David Parrish 1/13/2019 6:26 PM

Deleted: , and

David Parrish 1/13/2019 6:28 PM

Deleted:

David Parrish 1/13/2019 6:24 PM

Deleted: two

David Parrish 1/13/2019 6:25 PM

Deleted: can

David Parrish 1/14/2019 4:13 PM

Deleted: situation

David Parrish 1/14/2019 4:14 PM

Deleted: mandating

David Parrish 1/14/2019 4:16 PM

Deleted: further

populated: the Midwest states (with Detroit, Michigan; Minneapolis-St. Paul, Minnesota; and Milwaukee, Wisconsin) and the Pacific Northwest states (with Seattle, Washington and Portland, Oregon). These urban areas all have estimated 2017 populations > 1.5 million. In contrast, the rural western states (Montana, North Dakota and South Dakota) have no cities with populations >260,000. Wyoming could potentially be included with these three rural western states; however,

5 Wyoming's maximum ODVs occur in winter due to photochemical ozone production within a basin containing intense oil and gas production activities (Oltmans et al., 2014). Since this is a very different situation from that in the other rural states, only the Yellowstone National Park site from this state is considered; it is adjacent to Montana and distant from the oil and gas activities. This site is included because it has a measurement record that has been examined in previous analyses of long-term trends of U.S. background ozone concentrations (e.g., Lin et al., 2017).

10 The temporal histories of the ambient ozone concentrations measured in these four regions approximately correlate with the magnitude of the urban populations (Figure 2). The largest ODVs have been recorded in the northeastern states, the lowest in the rural western states, with intermediate ODVs in the other two regions. As expected, the ODVs in none of these regions have approached the maximum ODVs recorded in the SoCAB (indicated by blue lines in Figure 2). The analysis that follows
15 in this paper considers only the two more extreme regions – the rural western states, where the ODVs have remained approximately constant at relatively small values over the 39 years of measurements, and the northeastern states, where the largest ODVs in the northern U.S. have been recorded.

The northeastern U.S. and the rural western states present dramatic contrasts. Although the eight northeastern states total only ~40% of the land area of the three western states, their population is a factor of ~16 larger. These differences imply a factor of ~40 greater population density in the northeast, despite large rural areas remaining within that region (e.g., parts of
20 Maine, Vermont and New Hampshire). This comparison emphasizes the concentration of population in urban areas in the northeast, with the 2017 New York City urban area population of > 20 million compared to no cities significantly exceeding 1% of that population in the rural western states. Consistent with this population distribution, the Northeast U.S. has suffered from ozone air pollution for decades (e.g., Wolff and Lioy, 1980). There are no designated ozone nonattainment areas in the

25 western rural states, while U.S. EPA's "Green Book" (<https://www.epa.gov/green-book>) currently lists five ozone nonattainment areas in the eight northeastern states. The two nonattainment areas designated as "Moderate" - New York-Northern New Jersey-Long Island, NY-NJ-CT and Greater Connecticut, CT - include the New York City urban area and the regions directly downwind. The other three, designated as "Marginal", include southern New Jersey in the Philadelphia-Wilmington-Atlantic City, PA-NJ-MD-DE exceedance area, plus two sparsely populated areas - Dukes County, MA with a population of ~17,000 and Jamestown, NY with a population of ~95,000. Transport of ozone into these latter two areas is
30 accepted as the dominant cause of the observed high ozone concentrations that led to their designations (Wilcox, 1996).

David Parrish 1/14/2019 4:18 PM
Deleted: average

David Parrish 1/14/2019 4:19 PM
Deleted: ,

David Parrish 1/14/2019 4:19 PM
Deleted: despite substantial improvements in ozone air quality (e.g., Lin et al., 2000), the

In this paper we examine the temporal and spatial variability of the highest ozone concentrations (i.e., the ODVs) observed over the past three to four decades in these two contrasting regions of the northern U.S. The analysis is designed to separately estimate the U.S. background ODVs and the enhancements of the ODVs above that background contribution due to U.S. anthropogenic precursor emissions. The U.S. background ODV estimates quantify the maximum ozone concentrations that would exist in these regions in the absence of U.S anthropogenic precursor emissions. We also aim to quantify the temporal evolution and spatial variability of the US anthropogenic enhancements of the ODVs, and based on past trends, project the expected time required for the maximum ozone concentrations to decrease to the 70 ppb NAAQS in the northeastern U.S. Photochemical modeling systems are generally utilized for such quantifications and projections (e.g., Dolwick et al., 2015; Emery et al., 2012; and Fiore et al., 2014). However, present model quantifications of U.S. ozone concentrations have quite large uncertainties (Jaffe et al., 2018; Guo et al., 2018). Parrish et al. (2017) present an observational based analysis that utilized the temporal record of measured maximum ODVs in eight air basins in Southern California for these purposes; here we apply this latter approach in the northern U.S. Comparison of these observational based results with those from modeling efforts identify needs for further research.

The analysis approach in this paper relies on differences in the temporal behavior of the U.S. background ODV (assumed approximately constant) and ODV enhancements due to US anthropogenic precursor emissions (assumed to have decreased exponentially over the past decades in response to U.S emission controls.) One complication arises from previously published studies that identified a multitude of additional processes that potentially can make systematic contributions on a variety of time scales to the variability of ozone concentrations at U.S. surface sites, including: stratospheric intrusions, which can bring particularly high ozone concentrations to the surface (Langford et al., 2009, 2014; Lin et al., 2012a, 2015); increasing Asian anthropogenic emissions, which are believed to raise ozone concentrations over the U.S. (Jacob et al., 1999; Lin et al., 2012b); increasing frequency of wildfires, which can produce episodic ozone enhancements (McKeen et al., 2002; Jaffe, 2008, 2013; Pfister et al., 2016); variable meteorological conditions, which can lead to changes in transport patterns (Wang et al., 2016) or changes in the conditions conducive to photochemical ozone production (Shen and Mickley, 2017; Shen et al., 2017), increasing methane, which is argued to increase global ozone concentrations (Fiore et al., 2008, and references therein); and a warming climate, which has been argued may partially offset air quality improvement from regional emission controls by (Fiore et al., 2015). However, there has been little in the way of systematic, quantitative analysis of the effects of these additional processes on ODVs across the U.S. Figure 2 shows clear overall reductions of ODVs across four decades, consistent with control of U.S. ozone precursor emissions, but there is also significant variability of ODVs about the long-term trends. In this work, we first quantify the U.S. background ODVs and the temporal effects of US anthropogenic precursor emission reductions, and then discuss the influence of other processes through analysis of the fraction of the ODV variability that cannot be accounted for by the first two factors.

David Parrish 1/14/2019 4:21 PM

Deleted: quantify

David Parrish 1/14/2019 4:22 PM

Deleted: s will be taken as estimates of

David Parrish 1/14/2019 4:23 PM

Deleted: In contrast,

David Parrish 1/14/2019 4:24 PM

Deleted: will help to

David Parrish 1/14/2019 4:25 PM

Deleted: have

Many papers have investigated U.S. surface ozone trends (see Lin et al., 2017 and references therein). These studies have treated a variety of statistics (medians, means, and various percentiles) to characterize the ozone concentration distribution. It is important to note that all trends investigated in this paper are based on the ODV statistic. The reason for this choice is that the NAAQS is based on this statistic, and so is most relevant for policy considerations. The ODV corresponds to ~98th percentile of the MDA8 concentrations during the ozone season. As a consequence, the U.S. background ODVs that we discuss are significantly larger than average or median background ozone concentrations that are often discussed in the literature. Given these different choices, care must be taken in comparing trends derived in this work with those from other analyses.

The sources of data and the analysis methods are discussed in the next section, followed by the applications of those methods to quantify the U.S. background ODVs and the enhancements from U.S. anthropogenic ozone precursor emissions in the western rural region (Section 3.1) and the northeastern U.S. (Section 3.2). The larger temporal ODV trends and the greater spatial variation of those trends in the northeastern U.S. provide the basis for the elucidation of several features of regional ozone trends, which are briefly discussed in the Appendix. Section 4 gives a summary of the approach and the results, discusses implications of those results, and identifies needs for further research.

2 Data and Methodology

This work considers Ozone Design Values (ODVs) reported from the beginning of U.S. ozone monitoring in the mid-1970s through 2017 in seventeen northern U.S. states; in-depth analyses focus on three, sparsely populated, western rural states and on eight states in the much more heavily populated northeast U.S. The ODV, the statistic upon which the U.S. NAAQS is based, is calculated each year for each ozone monitoring station in the U.S. if the measurements achieve the specified completeness criteria. Each year the ODVs recorded at all monitoring stations in the U.S. are added to EPA's AQS data archive (<https://www.epa.gov/aqs>). The ODVs reported for all years at all sites in all seventeen northern states were downloaded from this archive; only the ODVs marked as valid were retained for analysis. Exceptional events that have concurrence from the U.S. EPA were excluded. Table S1 summarizes these archived ODVs for each of the states, including the number of monitoring sites, the years spanned by the reported ODVs, and their maximum and minimum values. The number of monitoring sites in the states varies from 83 in Wisconsin to 4 in Rhode Island, with a total of 457 in the seventeen states. The reported ODVs span the range from 169 ppb to 41 ppb. It should be noted that very few sites have continuous measurements over the complete time spans indicated, and that many sites operated for only short periods. All reported ODVs are included in this analysis, even if only a single ODV was reported for a particular site. It is implicitly assumed that the temporal discontinuities associated with initiation or termination of individual sites does not prevent an accurate quantification of temporal trends of ODVs within the regions selected for analysis.

David Parrish 1/14/2019 4:27 PM

Deleted: the

David Parrish 1/14/2019 4:27 PM

Deleted: ODV

David Parrish 1/14/2019 4:27 PM

Deleted: Assuming that the four highest MDA8 ozone concentrations occur in the warmer half of the year when the highest ozone concentrations generally occur, t

David Parrish 1/14/2019 4:30 PM

Deleted: Most notably the Connecticut Department of Energy and Environmental Protection submitted an exceptional event demonstration to the U.S. EPA showing that emissions from the 2016 Fort McMurray wildfire caused elevated ozone levels throughout Connecticut on May 25 and 26, 2016, to which the U.S. EPA concurred. (http://www.ct.gov/deep/lib/deep/air/ozone/2016_exceptional_event_request/ct_deep_ft_mcmurray_ee_demo_concurrence_letter_and_tsd_7_31_2017.pdf).

David Parrish 1/14/2019 4:31 PM

Deleted: The earliest ODV was reported in 1973 in New York.

An examination of long-term trends of ODVs has been reported for southern California air basins (Parrish et al., 2017) that utilized an exponential function with a constant positive offset (Eq. 1) to quantify the temporal evolution of the ODVs in each air basin:

$$\text{ODV} = y_0 + A \exp\{-(\text{year}-2000)/\tau\}. \quad (1)$$

5 Mathematically, the first term of Eq. 1, y_0 , is the asymptotic value toward which the ODVs approach as the year of the ODV increases, and the second term is the enhancement of the ODVs above y_0 , which decreases exponentially with an e-folding time constant of τ years. Thus, A is the enhancement of the ODVs above y_0 in a reference year, defined in this work as the year 2000. Parrish et al. (2017) show that a single value of $\tau = 21.9 \pm 1.2$ years, a single value of $y_0 = 62.0 \pm 1.9$ ppb, and a different value of A in each air basin provided an excellent fit ($r^2 = 0.984$) to the maximum ODVs recorded in six southern California air basins over the 1980-2015 period (shorter periods ending in 2015 in two of those air basins). In this work Eq. 1 with τ set to the same value (21.9 years) is used to derive fits to the evolution of ODVs in the northern U.S. states.

A well-established conceptual model (e.g., Parrish et al., 1986) provides qualitative support for the application of Eq. 1. At any given location in the U.S., we can consider ambient ozone concentrations to be composed of two contributions: 1) background ozone and 2) enhancements resulting from ozone produced from photochemical processing of U.S. anthropogenic emissions of ozone precursors. The first contribution is defined as the ozone that would be present in the absence of U.S. emissions of ozone precursors from anthropogenic sources; this is ozone transported into the U.S., plus that produced over the U.S. from naturally emitted precursors, modified by loss processes. The U.S. Environmental Protection Agency (EPA) has defined this contribution as U.S. background ozone (e.g., Dolwick et al., 2015). The second contribution accounts for ozone produced from U.S. anthropogenic precursor emissions, ~~includes~~ that transported into a region from upwind U.S. sources, and that produced locally and regionally. We identify the first term of Eq. 1, y_0 , as an estimate of the ODV that would result from U.S. background ozone alone (i.e., U.S. background ODV), and the second term as an estimate of the enhancement of observed ODVs above y_0 due to ~~contributions~~ from U.S. anthropogenic precursor emissions. This interpretation implies that the average of the annual fourth-highest MDA8 ozone concentration in the absence of U.S. anthropogenic emissions would equal y_0 , and that in the year 2000 the long-term trend in ODVs is best fit by the sum of y_0 plus A ; hence A is identified as the ODV enhancement above y_0 in the year 2000. It should be noted that in the absence of anthropogenic emissions, the four days of highest MDA8 ozone concentrations that would determine y_0 are likely not the same days that determined the actual ODV in any particular year. Indeed, Parrish et al. (2017) show that there has been a systematic shift of the days with maximum ozone concentrations to earlier in the year as the U.S. anthropogenic ozone contribution, with a summertime maximum due to its photochemical source, has decreased, and the U.S. background ozone contribution, which has a summertime maximum, has come to dominate to a progressively greater extent.

David Parrish 1/14/2019 4:35 PM
Deleted: which can be further divided into two separate contributions -

David Parrish 1/14/2019 4:35 PM
Deleted: the sum of the two

Equation 1 implies that the U.S. anthropogenic contribution has decreased exponentially as emission control efforts reduced ozone produced from all U.S. anthropogenic sources. Associating U.S. background ozone with the parameter y_0 is equivalent to extrapolating long-term trends of observed ODVs to the limit of zero U.S. anthropogenic emissions, when the exponential term in Eq. 1 becomes zero. Parrish et al. (2017) used 36 year ODV records (1980-2015) in southern California to derive a value of $\tau = 21.9$ years for that region of the country. As will be shown in the following discussion, the ODV records in the northern U.S. do not show consistent decreasing trends over such long time periods. As a consequence it is not possible to precisely extract 3 parameters from the regression fits of the measurement records to Eq. 1. Here we take the value of $\tau = 21.9$ years derived from California to apply to all of the northern U.S., which implicitly assumes that control strategies have produced approximately equal relative reductions in anthropogenic ozone enhancements throughout the country. Fixing the value of τ allows precise determinations of the other two parameters (y_0 and A) from the fits to Eq. 1.

Two differences between the application here and that of Parrish et al. (2017) should be noted. First, the former work chose 1980 as the reference year, while here we choose the year 2000. The curves derived from the fits to Eq. 1 and the values derived for the y_0 parameter do not depend on the choice of reference year, while the values derived for the A parameter do. Consequently, the A parameters derived here cannot be directly compared with those given for California by Parrish et al. (2017); however, the second term of Eq. 1 with the parameters derived in this work and year set to 1980 does give values that can be directly compared to the A parameter values from that earlier work. Second, the former work considered only the maximum ODV recorded in each year at any of the sites within a given air basin, while here we primarily analyze the ODVs from all sites recorded in a given year in selected regions, although we do also consider the maximum ODVs recorded in each of the northeastern U.S. states. Since y_0 is an estimate of the U.S. background ODV, the value derived for this parameter is expected to be independent of whether the fit of Eq. 1 is made to all ODVs or to only the maximum ODVs in a particular region.

As discussed in the Introduction, the conceptual model utilized for interpretation of Eq. 1 is simplified; it assumes that decreasing U.S. anthropogenic emissions is the only cause of ODV variability at a particular location. Other factors such as rising anthropogenic emissions in Asia, variable occurrence of wild fires, and interannual meteorological and climate variability can also potentially affect observed ODVs, but they cannot be simply included in Eq. 1. The approach taken here is to interpret the observed ODVs solely on the basis of Eq. 1, and then to examine the fraction of the ODV variability captured by that interpretation. The remaining fraction of the variability is then attributed to other factors, including those listed above. Here we use two statistics to quantify the fraction of the variability not captured by the fits to Eq. 1. The root-mean-square deviation (RMSD) between the derived fit and the observed ODVs gives an absolute measure (in ppb) of the ODV variability not captured by Eq. 1. The square of the correlation coefficient (r^2) between the observed ODVs and the values derived from the fit to Eq. 1 gives a measure of the fraction of the total variability of the ODVs that is captured by that fit; the difference between unity and that r^2 value is then a relative measure (as a fraction) of the ODV variability not

David Parrish 1/14/2019 4:36 PM
Deleted: two
David Parrish 1/14/2019 4:36 PM
Deleted: s
David Parrish 1/14/2019 4:36 PM
Deleted: have

captured by Eq. 1. Other factors must then account for the variability quantified by those statistics. In the earlier study of ODVs in southern California air basins (Parrish et al., 2017), the derived $r^2 = 0.984$ and the RMSD ≈ 4 ppb indicate that all factors not included in Eq. 1 account for no more than 1.6% of the total variability in the basin maximum ODVs analyzed in that work, and contribute a RMSD to those ODVs of no more than ~ 4 ppb.

5 A potential complication in the interpretation of the two terms of Eq. 1 arises if there is a significant fraction of U.S. anthropogenic ozone precursor emissions that has not been reduced by emission controls. Ozone produced from such emissions would not have decreased in the same manner as that produced from most U.S. anthropogenic emissions, which could raise the derived value of y_0 above the actual U.S. background ODV. Parrish et al. (2017) have discussed this issue with regard to the emissions associated with the intense agricultural activity in the Imperial Valley of the Salton Sea air
10 basin, where the derived y_0 is higher than in other southern California air basins. The final section of this paper briefly considers the possible impact of this complication in the northeastern U.S. states.

In this work we consistently give 95% confidence limits for derived parameters, unless indicated otherwise. Most of the analysis in this work is based on non-linear, least-squares regression fits of the archived ODVs to Eq. 1, and interpretation of the derived values for the parameters y_0 and A . In this interpretation it is important to properly consider the uncertainty of the
15 derived parameter values. We derive the 95% confidence limits from the least-squares fitting routine that have been adjusted to account for the known covariance between the recorded ODVs. Each ODV is a three-year running mean; therefore only every third ODV is independent from the others determined at a given site. Consequently, the number of independent ODVs in each fit is approximately a factor of three smaller than the number of reported ODVs. Thus, the fitting routine underestimates the true confidence limits of the derived parameters; all derived confidence limits have been increased by a
20 factor of $3^{1/2}$ to account for this covariance, giving more realistic confidence limits. There are additional sources of covariance between the ODVs included in any particular fit. The ODVs from different sites within a region can co-vary due to regionally coherent interannual variability. Also temporal interannual variability may possibly lead to covariance between ozone concentrations measured in successive years at a site. We are not able to account for the effect of this additional covariance; the derived confidence limits are thus lower limits for the true confidence limits of the derived parameters.

25 **3 Results**

Here we fit the time series of ODVs from the western rural states (Section 3.1) and the northeastern states (Section 3.2) to Eq. 1 and discuss the results in the context of the conceptual model introduced above. This model considers the recorded ODVs to comprise two contributions: 1) an approximately constant U.S. background ODV (i.e., the ODV that would exist if U.S. anthropogenic ozone precursor emissions were reduced to zero), which is identified with y_0 in Eq. 1, and 2)
30 enhancements due to local and regional photochemical production of ozone from the actual U.S. anthropogenic ozone precursor emissions, which are approximated by the second term in Eq. 1. These photochemical enhancements have been

progressively reduced over past decades by U.S. emission controls. Section 3.2 and the Appendix discuss further details of the spatial and temporal variability of ODVs in the northeastern U.S.

3.1 ODVs in western rural states

The sparsely populated, three-state, western rural region generally lies on the Northern U.S. Great Plains downwind of more mountainous terrain to the west. Figure 3 shows a topographical map of the region, with the locations of the ozone monitoring sites indicated. The region gradually slopes to the east and north. With the exception of Yellowstone NP at 2.43 km, all of the monitoring sites lie below 1.55 km elevation.

The histories of the ODVs recorded in the region (Figure 4 with data statistics and fitted curves summarized in Table 1) indicate that throughout the measurement period, there has been little enhancement of ozone concentrations due to anthropogenic influences and little variability due to any cause. The 283 tabulated ODVs recorded over 39 years at 35 sites in the three states average 59.3 ppb with a standard deviation of 3.7 ppb – strong support for the assumption of an approximately constant U.S. background ODV within this region. At the individual sites and within each state the entire measurement records are all well described by averages with small standard deviations (Table 1): < 3 ppb in Montana and North Dakota, and < 4 ppb in South Dakota, the state whose sampling sites span the largest elevation range (0.34 to 1.55 km). U.S. background ODVs generally increase with the elevation of the sampling site (e.g., see discussion in Jaffe et al., 2018), so larger variability is expected when the monitoring sites within a state span a larger range of elevations. Fits to Eq. 1 included in Figure 4 show small, statistically significant anthropogenic enhancements of ODVs in Montana and North Dakota, with A values of 2.1 ± 2.1 ppb and 2.5 ± 1.5 ppb, respectively. These fits slightly reduce the variability – the RMSD of all of the data about the fitted curves in Figure 4 is 2.7 ppb, compared to the standard deviation of 3.7 ppb for the average of all ODVs in the region. The derived values for the y_0 parameter all lie within a range of 9 ppb, but there are some statistically significant differences: a maximum in South Dakota (62.7 ± 5.1 ppb), a minimum in Montana (53.7 ± 1.9 ppb) with North Dakota in between (57.0 ± 1.5 ppb). Consistent with the site elevation differences, the average ODV at Yellowstone NP (elevation 2.43 km) is significantly larger than that at Glacier NP (elevation 0.96 km): 64.0 ± 2.1 and 54.5 ± 1.3 ppb, respectively (where the standard deviation of each data set is indicated); however statistical uncertainty prevents an attribution of the cause of this difference to either larger y_0 values at Yellowstone NP (59.1 ± 6.1 vs. 54.8 ± 2.9 ppb) or larger A values (6.8 ± 8.3 vs. -0.3 ± 3.1 ppb).

3.2 ODVs in northeastern states

A topographical map showing the network of ozone monitoring sites in the eight northeastern U.S. states is given in Figure 5. All of the ODVs recorded in four of the eight states are plotted in Figures 6 and 7, along with curves showing fits of Eq. 1 to the ODVs from selected groups of sites over selected time periods. These groups of sites, and similarly selected groups in

David Parrish 1/14/2019 4:41 PM
Deleted: the

David Parrish 1/17/2019 11:18 AM
Deleted: slightly

David Parrish 1/14/2019 4:44 PM
Deleted: , with

David Parrish 1/14/2019 4:44 PM
Deleted: and

the other four states, represent different environments within each state. Figures S1-S8 of the Supplement show detailed ODV temporal plots for all eight states with fits of Eq. 1 to the ODVs for all selected groups of sites; the locations of these groups are indicated on maps included in these figures. In all, seventeen groups of sites within the eight states were selected; they are listed in Table 2. Superimposed curves in Figures 6, 7 and S1-S8 indicate the fits of Eq. 1 to the ODVs from all groups of sites; the parameters derived from those fits are included in Table 2. The fits to the ODVs in the rural western states include the full time span of all available results. However in the northeastern U.S., for most of the groups of selected sites, a consistent decrease in ODVs is not defined by the available data over the full measurement record (e.g., the four groups of selected sites in Figures 6 and 7). The strategy utilized here is to fit the ODVs over the time period beginning when a clear, consistent decrease in ODVs is first established, and continuing through 2017, the most recent ODV available. In all cases these fits begin by 2000, with some beginning earlier - either at the start of measurement record, 1990 or 1995 determined by the best, consistent fit to the functional form of Eq. 1.

There are some consistent general features of the ODV time series and the corresponding fits that inform the following analysis:

- Throughout the measurement record, the largest ODVs are found in the states that contain the New York City metropolitan area (New York, New Jersey and southwestern Connecticut), or that lie directly downwind (coastal Connecticut and Long Island, New York). Such sites compose two of the selected groups of sites in New York and Connecticut (see highlighted points in that area in Figure 5), whose ODVs and fits of Eq. 1 are highlighted in Figure 6.
- In several states, the largest ODVs are recorded at coastal sites (i.e., Connecticut, Massachusetts, New Hampshire and Maine in Figures 6, 7, S3, S5, S6 and S8). Two Massachusetts coastal sites are relatively isolated, located on the offshore island of Martha's Vineyard and near the tip of Cape Cod; however, these sites record some of the highest ODVs within that state (see Figure S5). Dukes County, which includes only Martha's Vineyard and nearby smaller islands, with a total population of ~17,000 has been designated as a marginal non-attainment area for ozone. The high ODVs at coastal sites emphasize the important, widely-discussed (e.g., Wolff and Lioy, 1980; Wilcox, 1996) role of transport in bringing high ozone concentrations from the major East Coast urban areas far downwind, particularly when that transport occurs over the waters of Long Island Sound and the Coastal Atlantic Ocean.
- In the past, ODVs at rural, generally upwind sites on the western border of New York (green symbols in on the left in Figure 5) were significantly smaller than in the northeastern U.S. urban areas, although in recent years that difference has diminished (Figure 6). These upwind rural areas in New York, and similar sites in Vermont (Figure S7), experienced ozone concentrations exceeding 80 ppb throughout the measurement record until about 2005. These high concentrations caused Chautauqua County, N.Y., with a population of ~95,000, to be designated as a marginal non-attainment area, again emphasizing the importance of ozone transport in the northeastern U.S., although in this case the source of the transported ozone is not as clearly established.

Additional systematic features of the ODV time series in the northeastern U.S. are discussed in the Appendix.

All of the curves derived from the fits of Eq. 1 to the long-term trends of the ODVs shown in Figures 6, 7 and S1-S8 are compared in Figure 8, with the corresponding parameters included in Table 2. Except for the four fits denoted by the colored dotted and dashed curves, all fits are similar in the sense that they exhibit the same relative long-term decrease and are asymptotically approaching approximately the same value of y_0 . The same relative long-term decrease is necessarily forced by the use of the same value of $\tau = 21.9$ years in all fits. However, the derived A and y_0 values do provide information regarding the spatial and temporal variation of ODVs over the past two to three decades. Three of the four curves with noticeably different behavior are from fits to the groups of sites with the highest recently reported ODVs (Connecticut, especially the coastal sites, and the New York sites highlighted in Figures 5 and 6); these are discussed further in the Appendix. The fourth exception is the one high elevation site (Mt. Washington in New Hampshire at an elevation of 1.9 km), which is also discussed separately in the Appendix. The parameters in Table 2 provide the basis for quantitatively comparing the fits throughout the northeastern U.S. in the next two sections.

3.2.1 Estimation of U.S. background ODV in northeastern states

The arithmetic mean of all y_0 values in Table 2 (excluding the four exceptions indicated in Figure 8) is 45.9 ppb with a standard deviation of 3.2 ppb. The average of these y_0 values weighted with the inverse square of the respective confidence limits is 45.8 ± 1.7 ppb, where the 95% confidence limit of this average is indicated. All of the y_0 values in Table 2 agree (excluding the four exceptions noted above) with these average values within their indicated confidence limits. Figure S10 of the Supplement shows an investigation of the distribution of the y_0 determinations; 13 of the 17 derived y_0 values approximately define a normal distribution with a median of 47.7 ppb and a standard deviation of 4.5 ppb. The median is interpreted as representing a common regional y_0 value, and the standard deviation as reflecting the uncertainty in determining each y_0 value. This median is consistent with the above averages. The highest 4 of 17 derived y_0 values define a high value tail; these are the 4 four exceptions indicated in Figure 8.

Recalling earlier discussion, we identify the average $y_0 = 45.8 \pm 1.7$ ppb as the best estimate of the U.S. background ODV throughout the northeastern U.S.; there is no discernable spatial variability within this region. This value is significantly smaller than the value of 62.0 ± 1.9 ppb derived for southern California (Parrish et al., 2017); however even in northeastern U.S., the U.S. background ODV amounts to 65% of the 70 ppb NAAQS.

3.2.2 Estimation of U.S. anthropogenic ODV enhancements in northeastern states

The fits to Eq. 1 with $\tau = 21.9$ years provide estimates of A , the enhancements of the ODVs above y_0 in the reference year 2000; Table 2 lists these values for the 17 selected groups of sites from two-parameter fits, i.e., fits with y_0 and A as independent parameters determined from the least-squares fits themselves. The discussion above estimated a constant value of $y_0 = 45.8 \pm 1.7$ ppb for the entire northeastern U.S. region. Based on this result, we can derive one-parameter fits to Eq. 1

David Parrish 1/14/2019 4:48 PM

Deleted: The fits of Eq. 1 to the long-term trends of the ODVs from the seventeen selected groups of sites shown in Figures 6, 7 and S1-S8 elucidate the temporal and spatial variability of ODVs in the northeastern U.S.

David Parrish 1/14/2019 4:48 PM

Deleted: these fits

David Parrish 1/14/2019 4:49 PM

Deleted: Considerable similarity is apparent between the fits to the ODVs from the different selections of sites in the eight states.

David Parrish 1/14/2019 4:50 PM

Deleted: in this region

David Parrish 1/15/2019 1:28 PM

Deleted: y_0 as an estimate of U.S. background ODV, i.e., the ODV expected if U.S. anthropogenic emissions were reduced to zero. We take the latter value of

David Parrish 1/15/2019 1:28 PM

Deleted: this

David Parrish 1/15/2019 1:04 PM

Deleted: U.S. background ODV

David Parrish 1/15/2019 1:04 PM

Deleted: in the U.S. background ODV

with y_0 held constant at this value of 45.8 ppb; the results of these fits are included in Table 1 as the A^* values. (Such a fit is not included for the Mt. Washington results, since U.S. Background ODV is evidently greater than 45.8 ppb as discussed in the Appendix). The A^* values generally agree with the A values from the two-parameter fits within their confidence limits, which are smaller, since only one parameter need be derived. The exceptions to the agreement between A and A^* are the fits to the exceptions discussed earlier - the two groups of Connecticut sites and the New York maximum ozone sites, which are the upper three colored curves in Figure 8. In Table 2 the A values for these three groups of sites are anomalously small compared to the results from neighboring groups of sites (i.e., New Jersey, Rhode Island, Massachusetts/coastal); the A^* values for all of these neighboring groups of sites agree more closely. In the following discussion we take these A^* values as the best estimate for the U.S. anthropogenic ODV enhancements in the northeastern states.

An overview of the spatial variation of the ODV enhancements across the northeastern U.S. can be obtained from a contour plot (Figure 9) derived from the A^* values in Table 2. The groups of selected sites fit to Eq. 1 give only coarse spatial resolution across the northeastern U.S.; thus the contour plot has uncertainties not apparent from the smooth spatial variability of this figure. This uncertainty has been mitigated in the contour plot calculation by inclusion of duplicate values added at the site locations in each selected group of sites; these additions ensure that the contouring program reproduces a more nearly constant value over the sometimes large regions covered by the selected groups of sites from which the individual A^* values were derived. Despite the uncertainties, the contour plot does give a useful, semi-quantitative representation of the magnitude and regional variation of the ODV enhancements in the northeastern U.S. Note that the contour plot and the A and A^* values of Table 1 describe the ODV enhancements in the year 2000. As is apparent from Eq. 1 and the illustrated temporal trends in the figures, the ODVs have decreased throughout the last two to three decades. The e-folding time of $\tau = 21.9$ years implies that between the reference year of 2000 and 2017, the ODV enhancements decreased by a factor of 2.2. Hence, dividing the year 2000 ODVs in the contour plot by that factor gives an approximation of the 2017 ODV enhancements.

The ability of Eq. 1 to accurately reproduce observed ODVs can be judged by comparing those observed ODVs with the values predicted from the fits derived with $y_0 = 45.8$ ppb and $\tau = 21.9$ years. Figure 10a shows this comparison as a correlation plot. The fits for ODVs recorded at all sites in the eight northeastern states over the entire measurement period are calculated from the A^* values at each site interpolated from the contour plot of Figure 9. The correlation is high ($r^2 = 0.71$) for the 1719 separate ODV values recorded at the 148 sites over the 2000-2017 period, but significantly lower for earlier years as expected from the figures illustrating the derived fits. A general decrease in ODVs throughout the region did not begin until 2000, which is about the time that the U.S. EPA "NOx SIP Call" began reducing power plant NOx emissions across much of the eastern U.S. (Aleksic et al., 2013). There is significant scatter about the 1:1 line in the comparison in Figure 10a; the RMSD between observed and calculated ODVs is 5.6 ppb for the 2000-2017 period. Much of this scatter is due to variability in ODVs recorded at different sites within a given region, which arises from differences in local

David Parrish 1/17/2019 12:26 PM
Deleted: y_0
David Parrish 1/17/2019 12:27 PM
Formatted: Font:Not Italic
David Parrish 1/17/2019 12:27 PM
Formatted: Font:Not Italic

David Parrish 1/14/2019 4:55 PM
Deleted: as they existed
David Parrish 1/14/2019 4:55 PM
Deleted: We have used an
David Parrish 1/14/2019 4:55 PM
Deleted:
David Parrish 1/14/2019 4:55 PM
Deleted: to represent this decrease. This
David Parrish 1/14/2019 4:56 PM
Deleted: (the latest year of data included in this analysis)
David Parrish 1/17/2019 12:29 PM
Deleted: that

photochemical ozone production and transport patterns. This variability can be reduced by comparing state maximum ODVs, rather than individual site ODVs. Figure 11 plots the time series of the maximum ODVs recorded in each year for the eight northeastern states with respective fits over the 2010-2017 period. The derived A^* values (given in Table 3) are somewhat larger than would be expected from the contour plot in Figure 9, consistent with consideration of only the maximum ODVs recorded in each state. Stronger correlation ($r^2 = 0.89$) is found for the fits to the state maximum ODVs as expected, since considering only the largest of the states' ODVs in a given year removes much of the regional variability across the state.

3.2.3 Alternative approach for estimating U.S. background ODVs in the northeastern U.S. states

An independent analysis approach discussed in Section 2.3 of Parrish et al. (2017) can also estimate U.S. background ODVs without assuming any specific functional form for the time dependence of the ODV enhancements. Different assumptions underlie this analysis - namely that all of the ODV time series under consideration follow the same functional form, but not necessarily an exponential decrease, and that all time series are approaching a common U.S. background ODV. These assumptions imply that all of the time series will converge to a common ODV as anthropogenic precursor emissions are reduced to zero; this common ODV is necessarily the regional U.S. background ODV. In practice this analysis uses correlations between time series of ODVs that have different anthropogenic ODV enhancements. One time series is selected as a reference; in the examples discussed here the time series with the largest ODVs is selected. Other time series are then linearly correlated with this reference. The intercept of each linear correlation with the 1:1 line then provides an estimate of the U.S. background ODV; at that point the ODVs from the two time series are equal. Parrish et al. (2017) show that the results of this approach for seven southern California air basins are nearly identical to the results from fits to Equation 1.

This correlation approach applied to the northeastern U.S. is shown in Figures S11-S18 of the Supplement. All 13 regional data sets that give the black lines in Figure 8 are included in this analysis. The time series of maximum observed ODVs in the New York City urban area (discussed in Section 4 and shown in Figure 12a) is selected as the reference, because they are some of the largest ODVs recorded in the northeastern U.S., and after 2000 this time series closely follows an exponential decrease with little interannual variability. Figure S18 of the Supplement shows all of the linear regressions for the 13 regional data sets, and Table S3 summarizes the results. These results have significant variability (25 to 62 ppb) due to the relatively short 2000-2017 data records and because the slopes are not widely different from unity, preventing a precise determination of the intercepts of the correlations with the 1:1 line. However, the average of the derived background ODVs (49.2 ± 3.9 ppb for ordinary linear regressions and 42.5 ± 5.7 ppb for reduced major axis regressions, where 95% confidence limits are indicated) bracket the result (45.8 ± 1.7 ppb) derived from the exponential fits, and neither average is statistically significantly different from that earlier result. The agreement between these two approaches for estimating U.S. background ODVs shows that the assumption of an exponential decrease in the ODV enhancements is not essential for estimating the background ODV (although that approach does give more precise results), and increases our confidence in the results of each approach.

4 Discussion and Conclusions

The analysis presented in this paper has two complementary parts. First, time series of the highest ozone concentrations (i.e., the ODVs, the statistic upon which the NAAQS is based) observed in selected regions in the northern U.S. are fit to the simple mathematical relationship given by Eq. 1. This equation has two terms - one constant and one that exponentially decreases with an e-folding time of 21.9 years. Equation 1 provides fits to the most recent two to four decades of all time series of ODVs considered here through two variable parameters: y_0 , the magnitude of the constant term, and A , the magnitude of the exponentially decreasing term in the year 2000. The success of this mathematical process is judged through standard statistical tests that (1) quantify how well Eq. 1 captures the variability of the ODV time series to which the equation is fit, and (2) quantify the uncertainty of the derived parameter values. The second part of the analysis is the physical interpretation of the parameters derived from the fits to Eq. 1; y_0 is taken as an estimate of the U.S. background ODV (i.e., the ODV that would exist if U.S. anthropogenic emissions of ozone precursors were reduced to zero), and the second term of Eq. 1 is interpreted as an estimate of the regional enhancement of the observed ODVs above that U.S. background ODV due to photochemical production of ozone from U.S. anthropogenic precursor emissions. These estimates apply to the region of the ODV determinations, and vary between selected regions. Thus, the derived values of y_0 and A with their calculated uncertainties quantify the estimates of the two contributions to observed ODVs. Parrish et al. (2017) applied this same analysis approach to large areas of southern California, including the Los Angeles urban area.

In this paper the above-described analysis is applied to the northern tier of U.S. states, with a focus on two contrasting regions – three sparsely populated rural western states (Montana, North Dakota, and South Dakota, with map in Figure 3) and eight northeastern states (New York, New Jersey, and the six New England states, with map in Figure 5). This selection is made in order to use the marked regional contrasts to judge the value of the physical interpretation of Eq. 1. The northeastern states contain the New York City urban area – the most populous U.S. urban area – while the western rural states contain no large cities. The time series of ODVs recorded in these two regions differ markedly. The rural western states span a distance of ~1700 km and cover a land area of >0.75 million km², but the ODVs recorded at 35 different sites throughout this area over a 39-year period show remarkably little variability (Figures 2 and 4) with an overall standard deviation of 3.7 ppb. In contrast, the ODVs recorded in the northeastern states vary from >160 to <50 ppb (Figures 2, 6, 7 and S1-S8).

There are significant spatial variations in the derived U.S. background ODVs. Although ODVs have varied little, either spatially or temporally, within the rural western states through the entire measurement record, small but statistically significant regional variability in the U.S. background ODV and the small anthropogenic enhancements are quantified (Table 1); accounting for their associated variability reduces the standard deviation of the residuals to 2.7 ppb from the overall standard deviation of 3.7 ppb for all ODVs recorded over 39 years in the western rural states. Within the northern U.S., the largest y_0 value is found for South Dakota (62.7 ± 5.1 ppb), which agrees within statistical confidence limits with that

David Parrish 1/18/2019 12:04 PM

Deleted: The fits of Eq. 1 to time series of ODVs in selected groups of sites in these two regions allow the estimation of both the U.S. background ODV and the enhancements of the ODVs due to U.S. anthropogenic precursor emissions, and elucidate the spatial variability of both.

derived for southern California (62.0 ± 1.9 ppb). Within the western rural states, Montana has the smallest U.S. background ODV (53.7 ± 1.9 ppb) with North Dakota (57.0 ± 1.5 ppb) between those two values. The U.S. background ODV in the northeastern U.S. states (45.8 ± 1.7 ppb) is significantly smaller than in any of the western U.S. regions. For context, these U.S. background ODVs account for 65 to 90% of the 2015 NAAQS of 70 ppb. No statistically significant differences are discernable over the northeastern U.S. states, except at the one relatively high elevation site (Mt. Washington in New Hampshire at 1.9 km), which has a larger $y_0 = 66 \pm 7$ ppb that is believed to be characteristic of the lower free troposphere (see Appendix for further discussion).

In the western rural states, the anthropogenic ODV enhancements, as quantified by the derived A parameters, are quite small with a weighted average of 2.4 ± 1.2 ppb calculated for the three states plus Yellowstone NP (see Table 1 for derived parameters). In the northeastern U.S. the A parameters have a systematic spatial dependence as shown by the contour plot in Figure 9, with the largest values (>54 ppb) immediately downwind of New York City decreasing to <22 ppb over northeastern Maine. Importantly, these derived A parameters quantify the anthropogenic ODV enhancements in the year 2000. By 2017 these enhancements had decreased by a factor of 2.2; thus the largest ODV enhancements immediately downwind of New York City have decreased to ~ 25 ppb.

15 4.1 Implications of the results

This analysis presented here and the results of Parrish et al. (2107) demonstrate that throughout diverse regions of the country (i.e., northern rural states, the northeastern U.S., and California) the U.S. background ODV contribution is significantly larger than the present-day ODV enhancements due to photochemical production from U.S. anthropogenic precursor emissions. This comparison is true not only in rural areas, but also in the two most populous U.S. urban areas, New York City and Los Angeles. Since these ODVs are the largest observed ozone concentrations upon which the NAAQS is based, degraded air quality due to elevated ozone concentrations can be attributed primarily to the hemisphere-wide source that is responsible for the U.S. background ODV, with local and regional photochemical production enhancing that background to a significant, but smaller amount.

Forward projections of the fits to the maximum ODVs (shown in Figures 11 and 12a) allow an estimate of future trends of ODVs in the northeastern U.S. under the assumptions that 1) the U.S. background ODV (i.e., y_0) remains constant at 45.8 ppb throughout the region, and 2) the exponential decrease in ODV enhancements can be maintained with an e-folding time, τ , of 21.9 years, by means of continued emission reduction efforts. These projections suggest that the maximum ODVs throughout the northeastern U.S. will drop below the 2015 NAAQS of 70 ppb by about 2021. However, these projections do not account for the variability of observed maximum ODVs (i.e., RMSD of 3.9 ppb in the northeastern U.S.) about the fitted curves, so that even after 2020 this variability will likely result in the continued occasional recording of ODVs above 70 ppb.

5 These forward projections cannot account for any systematic deviations of the ODVs from the behavior given by Eq. 1. The recent temporal evolution of ODVs in Connecticut appears to differ significantly from the general regional behavior (see Figures 6-8 and 11). In the discussion of the fit to Eq. 1 of the Connecticut ODVs, as well as the maximum New York City ODVs this difference was noted (see dashed colored curves in Figure 8), but nevertheless the temporal evolution was forced with $y_0 = 45.8$ ppb in deriving the A^* values given in Table 2 and in deriving the contour plot of Figure 9. The different behavior and fits for Connecticut are due to the most recent five years of ODV values lying above the expected trend, as most clearly shown in Figure 11. The cause of this difference is not understood. Whether this difference is simply a statistical fluctuation cannot be determined at this time; however, random fluctuations of similar magnitude are only rarely apparent in the temporal records of ODVs in the states discussed. McDonald et al. (2018) have recently discussed a class of ozone precursor emissions, i.e., volatile chemical products - including pesticides, coatings, printing inks, adhesives, cleaning agents, and personal care products - that have not been addressed by emission controls to the same extent as other emission sectors. The impact of this emission sector on ODVs has not been quantified, but is expected to be most significant in areas of largest population density, exactly the regions where the significant differences in temporal evolution of ODVs are noted.

15 The higher U.S. background ODV (y_0) in southern California of 62.0 ± 1.9 ppb (Parrish et al., 2017) compared to the value of 45.8 ± 1.7 ppb derived here for the northeastern U.S. implies much less difficulty in achieving the 2015 ozone NAAQS of 70 ppb in the New York City (NYC) urban area compared to Los Angeles (LA). This situation arises because the northeastern U.S. has a much larger margin for anthropogenic enhancement of ODVs while still attaining the NAAQS. Figure 12 compares these U.S. background ODVs and the maximum ODVs in these two urban areas. In 2015 these curves indicated maximum ODVs of 78 and 102 ppb in NYC and LA, respectively. To lower the maximum ODVs to 70 ppb would require respective decreases in total ODVs of 10% in NYC and 31% in LA. However, only the ODV enhancements due to U.S. anthropogenic emissions can be addressed by local and regional controls of ozone precursor emissions. In 2015 these enhancements were about 25% larger in LA than in NYC (32 and 40 ppb, respectively). To reach a maximum ODV of 70 ppb requires ODV enhancement reductions of 25% in NYC and 80% (i.e. a factor of 5 reduction) in LA. The exponential term of Eq. 1 projects that such reductions of the 2015 ODV enhancements will require 5 years in NYC and 35 years in LA; hence the projected years of 2021 and 2050 in NYC (discussed above) and LA (Parrish et al., 2017), respectively. From the perspective of lowering maximum ODVs to the ozone NAAQS, the most important difference between NYC and LA urban areas is the higher U.S. background ODV in LA, although the 25% larger anthropogenic ODV enhancements in LA play a secondary role. This comparison provides an insightful context for the consideration of relative anthropogenic enhancements of ozone concentrations across the country.

30 It should be noted that from the perspective of human health, continuing efforts to reduce ambient ozone concentrations are beneficial, despite the difficulty of achieving the NAAQS. Recent studies establish human health impacts from long-term ozone exposure over several years (Turner et al., 2016; Di et al., 2017; Berger et al., 2017). Therefore, any reduction in

David Parrish 1/15/2019 1:36 PM
Deleted: by

ozone concentrations below present levels, regardless of whether or not ODVs remain above 70 ppb, will benefit U.S. human health.

4.2 Possible shortcomings of the analysis

In this work we have used Eq. 1 to quantitatively describe the temporal evolution of ODVs throughout the northern U.S.; this equation accounts for the U.S. background ODV and enhancements of the ODVs due to U.S. anthropogenic emissions. As noted in the Introduction, a multitude of additional processes (i.e., systematic departures of the temporal evolution of the ODVs from a purely exponential decrease, possible variation in U.S. background ODVs from a single y_0 value associated with stratospheric intrusions, rising anthropogenic emissions in Asia, variable or increasing occurrence of wild fires, meteorological and climate variations driving differences in ozone photochemical production between years, and transport differences associated with interannual meteorological and climate variability) have been identified as possibly affecting ozone concentrations on a variety of time scales at U.S. surface sites. However, there has been little in the way of systematic, quantitative analysis of the effects of these additional processes on ODVs across the U.S. Our approach here is to quantify the fraction of the variation of the ODVs captured by Eq. 1 in the rural western states and in the northeastern states, and then to attribute the remaining, unexplained variability to these additional influences. Section 2 discusses the use of the r^2 and RMSD statistics derived from the fits of Eq. 1 to the observed ODVs to quantify the variability contributions.

The analysis of the ODVs in the western rural states (Section 3.1) demonstrates that in this vast region, the U.S. background ODV contribution overwhelmingly dominates the observed ODVs; the 283 ODVs reported from 35 sites over 39 years of measurements in the three states have a standard deviation of only 3.7 ppb. At the individual sites and within each state the entire measurement records are all well described by averages with even smaller standard deviations (Table 1). For example, Glacier NP is a single site with a 27-year measurement record that is often utilized for characterizing background ozone concentrations (see Lin et al., 2017 and references therein); the ODVs at this site have a standard deviation of only 1.4 ppb over the entire record. Fits to Eq. 1 included in Figure 4 show indications of only small anthropogenic enhancements of ODVs above y_0 (see Table 1). These fits reduce the remaining variability in the ODVs to a limited extent – the RMSD of all of ODVs about the fitted curves in Figure 4 is 2.7 ppb. Thus, contributions of all factors to ODV variability, except for changing anthropogenic precursor emissions, including the multitude of effects discussed above plus any spatial variability between sites within the individual states, are small in these western rural states - no larger than the overall 2.7 ppb RMSD about the fitted curves.

The northeastern U.S. states present a strong contrast to the western rural states. In this region variation in the anthropogenic ODV enhancements dominates the total variability, and Eq. 1 captures the large majority of this variability. In Figure 10 the r^2 values for 18 years (2000-2017) indicate that Eq. 1 captures more than two-thirds of the variability of the 1719 ODVs reported from all 147 sites in the northeastern U.S., and 89% of the variability of the maximum ODVs in the eight states. The

David Parrish 1/15/2019 1:41 PM

Deleted: only small

5 difference between these percentages is attributed to spatial variability in the ODV enhancements not accurately represented by the contour plot of Figure 9, plus interannual variability in the spatial distribution of ODVs within the states. The RMSD between observed and calculated values is 5.6 and 3.5 ppb for all ODVs and the state maximum ODVs, respectively; these values provide estimates of the total influence of all other factors affecting ODVs over the region. Together, the root mean square contribution from these factors to the variability in the state maximum ODVs is no more than 3.5 ppb, which accounts for no more than 11% of their total variability over the 2000-2016 period. There are indications that the influence of these additional factors is even smaller than the 3.5 ppb RMSD of the state maximum ODVs, as several of the RMSD values for groups of sites in Table 2 are smaller than that value. In summary, Eq. 1 is remarkably successful at capturing a large fraction of the ODV variability in the northeastern U.S. states from 2000-2017; the combined influences from all other factors make only relatively minor contributions to the ODV variability across this entire region. Guo et al. (2018) discuss a contrasting result; they suggest that monthly regional mean U.S. background MDA8 ozone concentrations vary by up to 15 ppb from year to year, and that a 3-year averaging period (as is used to define the ODV) is not long enough to eliminate interannual variability in background ozone on the days of highest observed ozone. This is not a direct comparison, but it is clear that Guo et al. (2018) overestimate the actual variability of the observed ODVs in the two northern U.S. regions examined in this work and in southern California examined by Parrish et al. (2017).

David Parrish 1/15/2019 1:46 PM
Deleted: about

An uncertainty in the fits of the ODV time series to the exponential decay of the ODV enhancement term in Eq. 1 is the determination of the time constant, τ . The clear decrease in ODVs across the entire northeastern U.S. did not begin until about 2000; the 18 year period that decrease has continued is not long enough for fits of Eq. 1 to accurately derive all three parameters. The approach we have taken is to use $\tau = 21.9$ years, the value determined for southern California (Parrish et al., 2017), in these two northern U.S. regions. It is not clear how the time scales of ODV enhancement reductions compare between California and the northeastern U.S. In California, precursor emission reductions may have been faster, because that state may have had more aggressive emission control measures, but they may also have been slower because controls on eastern coal-fired power plants dramatically reduced NOx emissions. This latter reduction would not have occurred in California where such power plants are located downwind, out-of-state. On the other hand, emission reduction rates could be roughly the same, as most northeastern U.S. states have adopted the California on-road light-duty motor vehicle emission control program, and this is a large source sector both in California and the Northeast. The y_0 values derived from the fits are sensitive to the selected τ value, with a larger value of τ giving a smaller value for y_0 , and vice versa. For example, using the Boston data from Figure S5, increasing the assumed value of τ by 10% (from 21.9 to 24.1 years) decreases the derived y_0 estimate by 6% (from 45.8 to 43.3 ppb). Setting $\tau = 21.9$ years means that the confidence limits derived in this analysis are necessarily lower limits. Importantly, the agreement between the results from these exponential fits and the results from the alternative approach discussed in Section 3.2.3 suggests that this additional uncertainty is not large.

David Parrish 1/13/2019 5:05 PM
Deleted: this

David Parrish 1/13/2019 5:06 PM
Deleted: analysis is the definition of the time constant, τ , for

Finally, Eq. 1 implicitly assumes that all sectors of anthropogenic U.S. ozone precursor emissions have been reduced by emission controls at approximately the same rate. However, in some respects this is a poor approximation in that some emission sectors have received lesser efforts than others. Any emissions that have not been reduced would tend to lead to an overestimate in the U.S. background ODV, since ozone produced from those emissions would not have decreased. Parrish et al. (2017) note that continuing agricultural emissions in the Salton Sea Air Basin may account for the anomalously high y_0 value derived for that region, and the possible influence volatile chemical products (McDonald et al., 2018) in the northeastern U.S. is mentioned above. It is not possible to account for uncertainties in the results that may arise from this issue.

4.3 Needs for further research efforts

10 Accurately quantifying the U.S. background contribution to ODVs (i.e., the limit to which ODVs can be reduced through U.S. anthropogenic emission reductions alone) is important from the perspective of determining the extent of emission reductions required to attain the ozone NAAQS. In this work we have determined the value of the parameter y_0 of Eq. 1 within relatively small uncertainties (95% confidence limits of ~2ppb). These uncertainties are derived from the scatter in the observed ODVs about the fits to Eq. 1. However, identifying the value of y_0 with U.S. background ODVs brings in
15 additional possible uncertainties (see discussion in the preceding section) that have not been quantified. Traditionally, models have been used to estimate U.S. background ozone (see Jaffe et al., 2018 and references therein), but the models utilized in these efforts have significant shortcomings (e.g., see discussion in Parrish et al., 2017), that lead to large uncertainties in the results. Jaffe et al. (2018) estimate an uncertainty in modeled seasonal mean U.S. background ozone of about ± 10 ppb, with greater uncertainty for individual days (such as those that define the ODV), and Guo et al. (2018) find
20 biases as high as 19 ppb in modeled seasonal mean MDA8 ozone. Thus, ~~modeling and the observational based approach discussed in this paper~~ are both available for estimating U.S. background ODVs, but each has significant, poorly quantified uncertainties.

The estimates for the U.S. background ODV derived in this work can be compared with model results. Fiore et al. (2014) compare calculations of the fourth highest MDA8 North American background (NAB) ozone (also called policy-relevant
25 background (PRB) ozone) from two global models. The NAB concentration is that which would be present if anthropogenic emissions were reduced to zero throughout North America, not just in the U.S. NAB ozone concentrations are therefore somewhat smaller than U.S. background ozone concentrations, but for the purposes of this comparison, we can ignore this difference. The color scales in their Figures 2 and 10 allow estimates of the U.S. background ODV from the GEOS-Chem and AM3 models, respectively. Similarly, the color scale in Figure 6 of Emery et al. (2012) allows estimates of results from
30 a different version of the GEOS-Chem model for the fourth highest MDA8 PRB. Figure S9 and Table S2 compare the model results with the observationally based estimates of U.S. background ODV derived in this work. These model results do have

David Parrish 1/18/2019 12:23 PM

Deleted: two

David Parrish 1/18/2019 12:24 PM

Deleted: approaches

some skill in calculating the U.S. background ODVs. For five regions (three western rural states, the northeastern U.S. region, and the South Coast Air Basin) the model-observation correlations give r^2 values varying from 0.31 to 0.85, but the model results are on average systematically lower by 4 to 12 ppb. Importantly, the model results disagree with each other, as well as with the observationally based results.

5 In summary, effective air quality management can be usefully informed by quantification of U.S. background ODVs. However, given the relatively small differences between estimated U.S. background ODVs and the 2015 ozone NAAQS of 70 ppb, these quantifications will be of more utility if they are accurate to within a couple of ppb (see Figure 12 and associated discussion). Currently, two general approaches are available for estimating U.S. background ODVs (the observational based method discussed here and in Parrish et al. (2017) and a variety of modelling approaches), but the
10 | limited comparisons of results from these two approaches [and between the different model results](#) indicate differences much larger than ideal. However, the magnitudes of these disagreements are within the uncertainty of the model estimates as discussed by Jaffe et al. (2018) and Guo et al. (2018). Further improvement is required in modeling systems until their output can accurately reproduce the magnitude and variability of the time series of observed ODVs discussed here; these model calculations [could](#) then provide accurate determination of the U.S. background ODVs, the ODV enhancements from U.S.
15 | anthropogenic emissions, and robust interpretations of the parameters y_0 and A derived in this work. Until that model improvement is accomplished, the observationally based approach utilized in this work can provide useful estimates for air quality management guidance, as well as for comparison with evolving model calculations.

David Parrish 1/18/2019 12:28 PM

Deleted: can

David Parrish 1/18/2019 12:29 PM

Deleted: from the fits to Eq. 1 to observed ODVs

Appendix A. Additional features of ODV time series in the northeastern states

The text of the paper briefly described some consistent general features of the ODV time series and the corresponding fits to
20 Eq. 1 for selected groups of sites in the northeastern U.S. that guided the analysis. Here some additional features of interest are briefly discussed:

- New York has two non-attainment areas. In addition to the New York-N. New Jersey-Long Island, NY-NJ-CT moderate non-attainment area with a population of more than 20 million, there is the Chautauqua County (Jamestown), NY marginal non-attainment area with a population of less than 100,000. In Figure S1 the two sites in this latter non-
25 | attainment area are highlighted in purple; the ODVs from these two sites do not differ markedly from the other upwind sites on the western border of the state. In this analysis ODVs from all of the upwind sites are considered together.
- Sites in the New York urban area and regions downwind with over-water transport paths from that urban area have recorded the largest observed ODVs. Consistent with this identification, Vermont, the only state with neither major urban areas nor an over-ocean transport path from the New York City area, records the smallest maximum ODVs (see
30 | Table 3 and Figure S7).
- Although some sites the New York urban area record high ODVs, some other sites in central urban areas in the northeast U.S. are record the lowest ODVs (e.g., New Haven, Connecticut; Providence, Rhode Island, particularly before 2000;

and Boston Massachusetts in; see Figures S3, S4 and S7, respectively). This behavior is consistent with fresh NO_x emissions in urban areas reducing the ozone concentrations in air masses transported into those areas. This is evidently a very localized phenomenon, as the suburban sites adjacent to Boston (Figure S5) exhibit ODVs similar to other coastal sites in the state.

- 5 • The farthest downwind coastal monitoring site in northeast Maine (Figure S8) records significantly lower ODVs than other coastal sites, suggesting that ozone concentrations may decrease during transport due to dilution and/or ozone loss to surface deposition.
- Interestingly, Connecticut had much higher maximum ODVs than any other state before 1985 (all points above 140 ppb in Figure 2); their cause is unknown. Since 1985 Connecticut ODVs have been similar to those of neighboring states.
- 10 • Through the measurement record, the differences between maximum and minimum ODVs have decreased, both within individual states and throughout the entire region.
- There is one monitoring site at a relatively elevated location in the northeastern U.S. - Mt. Washington in New Hampshire at 1.9 km above sea level (asl). Although the ODV record at this site (Figure S6) is generally not higher than others recorded in New Hampshire, the fit to Eq. 1 shows a much smaller decrease than seen at any other site in the entire region. These ODVs followed a temporal evolution different from any of the other sites in the region (see curves in Figure 8 and parameters in Table 2). The A value (8 ± 8 ppb) is much smaller than that of any other selected set of sites, and the U.S. background ODV ($y_0 = 66 \pm 7$ ppb) is significantly higher than the common y_0 value of 45.8 ± 1.7 ppb derived for the entire northeastern U.S. This difference is attributed to the vertical gradient of ozone over the northeastern U.S. Ozone concentrations in the free troposphere increase with altitude (e.g., see Figure 2 of Fehsenfeld et al., 2006), and it is these higher altitude air parcels that impact Mt. Washington. The value of y_0 derived at Mt. Washington is in reasonable accord with the average ozone concentrations measured over the eastern U.S. by the MOZAIC program in the years near 2000 (Fehsenfeld et al., 2006). The enhancement of the ODVs (i.e., the in A value) in the free troposphere observations at Mt. Washington is much smaller than the enhancements seen at the other sites, which are all located within the planetary boundary layer. Note that the temporal evolution described by the parameters in Table 2 and illustrated in Figures 9 and S6 implies that the Mt. Washington summit site will soon record the highest ODVs in New Hampshire and higher than other sites in the northeastern U.S. outside of and immediately downwind from the New York City urban area; in 2017 Mt. Washington did report the largest ODV in New Hampshire.
- 20 • The Cadillac Mountain site at in Maine is at a somewhat elevated location (0.47 km asl). In contrast to Mt. Washington, the Cadillac Mountain ODVs (Figure S8) are generally similar to, although slightly higher than, others recorded at the southwest Maine coastal sites. Evidently Cadillac Mountain receives primarily boundary layer air masses. One coastal site is at a relatively high elevation (0.47 km) on Cadillac Mountain.
- 25
- 30

Competing interests. The author declares that he has no conflict of interest. He is the sole proprietor of David.D.Parrish, LLC, which has had contracts funded by several state and federal agencies and private companies. One of those contracts funded some of this work.

Acknowledgements. Some of the content of this paper was originally developed as a report submitted in fulfillment of the
5 Technical Services Agreement between the Northeast States for Coordinated Air Use Management (NESCAUM) and
David.D.Parrish, LLC funded under Agreement No. 101132 from the New York State Energy Research and Development
Authority (NYSERDA). NYSEDA has not reviewed the information contained herein, and the opinions expressed in this
report do not necessarily reflect those of NYSEDA or the State of New York. The author appreciates the comments and
discussion provided by Paul Miller of NESCAUM and Tom Ryerson, Fred Fehsenfeld, Owen Cooper and Andrew Langford
10 of NOAA ESRL CSD. The scientific results and conclusions, as well as any views or opinions expressed herein, are those of
the author, and do not necessarily reflect the views of NESCAUM, NOAA, or the Department of Commerce.

References

- Aleksic, N., Ku, J.-Y., Sedefian, L.: Effects of the NO_x SIP Call program on ozone levels in New York. *J. Air Waste Manage. Assoc.* 63, 1335–1342, 2013.
- Berger, R.E., Ramaswami, R., Solomon, C.G., Drazen, J.M.: Air pollution still kills, *N. Engl. J. Med.*, 376, 2591-2592, 2017.
- 5 Di, Q., Wang, Y., Zanobetti, A., Wang Y., Koutrakis P., Choirat C., Dominici F., Schwartz, J.D., Air Pollution and Mortality in the Medicare Population, *N. Engl. J. Med.*, 386, 2513-2522. 2017.
- Dolwick, P., Akhtar, F., Baker, K.R., Possiel, N., Simon, H., Tonnesen, G.: Comparison of background ozone estimates over the western United States based on two separate model methodologies, *Atmos. Environ.*, 109, 282-296, doi.org/10.1016/j.atmosenv.2015.01.005, 2015.
- 10 Emery, C., Jung, J., Downey, N., Johnson, J., Jimenez, M., Yarwood, G., Morris, R.: Regional and global modeling estimates of policy relevant background ozone over the United States, *Atmos. Environ.*, 47, 206-217, doi:10.1016/j.atmosenv.2011.11.012, 2012.
- Fehsenfeld, F. C., et al.: International Consortium for Atmospheric Research on Transport and Transformation (ICARTT): North America to Europe—Overview of the 2004 summer field study, *J. Geophys. Res.* 2006, 111, D23S01, doi:10.1029/2006JD007829.
- 15 Fiore, A.M., West, J.J., Horowitz, L.W., Naik, V., and Schwarzkopf, M.D.: Characterizing the tropospheric ozone response to methane emission controls and the benefits to climate and air quality, *J. Geophys. Res.*, 113, D08307, doi:10.1029/2007JD009162, 2008.
- Fiore, A.M., et al.: Estimating North American background ozone in U.S. surface air with two independent global models: Variability, uncertainties, and recommendations, *Atmos. Environ.*, 96, 284-300, doi.org/10.1016/j.atmosenv.2014.07.045, 2014.
- 20 Fiore, A.M., Naik, V., and Leibensperger, E.M.: Air Quality and Climate Connections, *J. Air Waste Manage.*, 65, 645–685, doi:10.1080/10962247.2015.1040526, 2015.
- Guo, J.J., Fiore, A.M., Murray, L.T., Jaffe, D.A., Schnell, J.L., Moore, C.T., and Milly, G.P.: Average versus high surface ozone levels over the continental USA: model bias, background influences, and interannual variability. *Atmos. Chem. Phys.*, 18, 12123–12140, doi.org/10.5194/acp-18-12123-2018, 2018.
- 25 Jacob, D.J., Logan, J.A., Murti, P.P.: Effect of rising Asian emissions on surface ozone in the United States, *Geophys. Res. Lett.*, 26, 2175–2178, doi:10.1029/1999gl900450, 1999.
- Jaffe, D., Chand, D., Hafner, W., Westerling, A., and Spracklen, D.: Influence of fires on ozone concentrations in the western US, *Environ. Sci. Technol.*, 42, 5885–5891, doi:10.1021/es800084k, 2008.
- 30 Jaffe, D., Wigder, N., Downey, N., Pfister, G., Boynard, A., and Reid, S. B.: Impact of Wildfires on Ozone Exceptional Events in the Western US, *Environ. Sci. Technol.*, 47, 11065–11072, doi:10.1021/es402164f, 2013.

- Jaffe, D.A., Cooper, O.R., Fiore, A.M., Henderson, B.H., Tonneson, G.S., Russell, A.G., Henze, D.K., Langford, A.O., Lin, M., Moore, T.: Scientific assessment of background ozone over the U.S.: Implications for air quality management. *Elem. Sci. Anth.* 6 56 doi.org/10.1525/elementa.309, 2018.
- Langford, A.O., Aikin, K.C., Eubank, C.S., Williams, E.J.: Stratospheric contribution to high surface ozone in Colorado during springtime, *Geophys. Res. Lett.*, 36, L12801, doi:10.1029/2009gl038367, 2009.
- Langford, A.O., Senff, C., Alvarez II, R., Brioude, J., Cooper, O., Holloway, J., Lin, M., Marchbanks, R., Pierce, R., Sandberg, S., Weickmann, A., Williams, E.: An overview of the 2013 Las Vegas Ozone Study (LVOS): Impact of stratospheric intrusions and long-range transport on surface air quality, *Atmos. Environ.*, 109, 305–322, doi:10.1016/j.atmosenv.2014.08.040, 2014.
- 10 Lin, C.-Y. C., Jacob, D. J., Fiore, A. M.: Trends in violations of the ozone air quality standard in the continental United States, 1980-1998, *Atmos. Environ.*, 35, 3217–3228, 2000.
- Lin, M., Fiore, A.M., Cooper, O.R., Horowitz, L.W., Langford, A.O., Levy, H., Johnson, B.J., Naik, V., Oltmans, S.J., Senff, C.J.: Springtime high surface ozone events over the western United States: Quantifying the role of stratospheric intrusions, *J. Geophys. Res.*, 117, D00V22, doi:10.1029/2012jd018151, 2012a.
- 15 Lin, M., Fiore, A.M., Horowitz, L.W., Cooper, O.R., Naik, V., Holloway, J., Johnson, B.J., Middlebrook, A.M., Oltmans, S.J., Pollack, I.B., Ryerson, T.B., Warner, J.X., Wiedinmyer, C., Wilson, J., Wyman, B.: Transport of Asian ozone pollution into surface air over the western United States in spring, *J. Geophys. Res.*, 117, D00V07, doi:10.1029/2011jd016961, 2012b.
- Lin, M.; Fiore, A.M., Horowitz, L.W., Langford, A.O., Oltmans, S.J., Tarasick, D., Rieder, H.E.: Climate variability modulates western US ozone air quality in spring via deep stratospheric intrusions, *Nat. Commun.*, 6, 7105, doi:10.1038/ncomms8105, 2015.
- Lin, M., Horowitz, L.W., Payton, R., Fiore, A.M., Tonneson, G.: US surface ozone trends and extremes from 1980 to 2014: quantifying the roles of rising Asian emissions, domestic controls, wildfires, and climate. *Atmos. Chem. Phys.*, 17, 2943-2970, doi.org/10.5194/acp-17-2943-2017, 2017.
- 25 McDonald, B.C., et al.: Volatile chemical products emerging as largest petrochemical source of urban organic emissions, *Science*, 359, 760-764, 2018.
- McKeen, S.A., Wotawa, G., Parrish, D.D., Holloway, J.S., Buhr, M.P., Hübler, G., Fehsenfeld, F.C. Meagher, J.F.: Ozone production from Canadian wildfires during June and July of 1995, *J. Geophys. Res.*, 107 (D14), doi:10.1029/2001JD000697, 2002.
- 30 National Research Council (NRC), 1991. Rethinking the Ozone Problem in Urban and Regional Air Pollution. National Academy Press, Washington, DC.
- Oltmans, S.; Schnell, R.; Johnson, B.; Pétron, G.; Mefford, T.; Neely III, R.: Anatomy of wintertime ozone associated with oil and natural gas extraction activity in Wyoming and Utah, *Elem. Sci. Anth.*, 2, 24, doi.org/10.12952/journal.elementa.000024, 2014.

- Parrish, D.D., Fahey, D.W., Williams, E.J., Liu, S.C., Trainer, M., Murphy, P.C., Albritton, D.L., Fehsenfeld, F.C.: Background Ozone and Anthropogenic Ozone Enhancement at Niwot Ridge, Colorado, *J. Atmos. Chem.*, 4, 63-80, 1986.
- 5 Parrish, D.D., and Stockwell, W.R.: Urbanization and air pollution: Then and now, *Eos: Earth & Space Science News*, 96, 10-15, 2015.
- Parrish, D.D., Young, L.M., Newman, M.H., Aikin, K.C., and Ryerson; T.B.: Ozone Design Values in Southern California's Air Basins: Temporal Evolution and U.S. Background Contribution, *J. Geophys. Res. Atmos.*, 122, 11,166–11,182, doi.org/10.1002/2016JD026329, 2017
- 10 Pfister, G.G., Wiedinmyer, C., and Emmons, L.K.: Impacts of the fall 2007 California wildfires on surface ozone: Integrating local observations with global model simulations, *Geophys. Res. Lett.*, 35, L19814, doi:10.1029/2008GL034747, 2008.
- Shen, L., and Mickley, L.J.: Effects of El Niño on summertime ozone air quality in the eastern United States, *Geophys. Res. Lett.*, 44, 12,543-12,550, doi.org/10.1002/2017GL076150, 2017.
- Shen, L., Mickley, L.J., Leibensperger, E.M., and Li, M.: Strong dependence of U.S. summertime air quality on the decadal variability of Atlantic sea surface temperatures, *Geophys. Res. Lett.*, 44, 12,527–12,535, doi.org/10.1002/2017GL075905, 2017.
- 15 Turner, M.C., Jerrett, M., Pope III, C.A., Krewski, D., Gapstur, S.M., Diver, W.R., Beckerman, B.S., Marshall, J.D., Su, J., Crouse, D.L., Burnett, R.T.: Long-Term Ozone Exposure and Mortality in a Large Prospective Study, *Am. J. Respir. Crit. Care Med.*, 193(10), 1134-1142, 2016.
- 20 Wang, Y., Jia, B., Wang, S.-C., Estes, M., Shen, L., and Xie, Y.: Influence of the Bermuda High on interannual variability of summertime ozone in the Houston–Galveston–Brazoria region, *Atmos. Chem. Phys.*, 16, 15265-15276, https://doi.org/10.5194/acp-16-15265-2016, (2016).
- Wilcox, G.L.: New England and the Challenge of Interstate Ozone Pollution Under the Clean Air Act of 1990. *B.C. Envtl. Aff. L. Rev.*, 24, 1-101, http://lawdigitalcommons.bc.edu/ealr/vol24/iss1/2, 1996.
- 25 Wolff, G. T., Lioy, P. J.: Development of an ozone river associated with synoptic scale episodes in the eastern United States. *Environ. Sci. Technol.*, 14, 1257–1260, 1980.

Table 1. Results of least-squares fits to Eq. 1 illustrated in Figure 1; RMSD indicates the root-mean-square deviation between the observed ODV's and the derived fit.

State/Site	Avg. \pm Std. Dev. (ppb)	y_0 (ppb)	A (ppb)	RMSD (ppb)	years fit
Montana	55.4 \pm 2.2	53.7 \pm 1.9	2.1 \pm 2.1	2.0	1979-2017
Glacier NP	54.5 \pm 1.3	54.8 \pm 2.9	-0.3 \pm 3.1	1.4	1991-2017
Yellowstone NP	64.0 \pm 2.1	59.1 \pm 6.1	6.8 \pm 8.3	1.7	1999-2017
North Dakota	59.3 \pm 2.7	57.0 \pm 1.5	2.5 \pm 1.5	2.5	1982-2017
South Dakota	61.5 \pm 3.8	62.7 \pm 5.1	-1.9 \pm 7.8	3.8	1990-2017

5 **Table 2. Results of least-squares fits to Eq. 1 illustrated in Figures 6-8 and S1-S8; RMSD indicates the root-mean-square deviation between the observed ODV's and the derived fit.**


State/sites	y_0 (ppb)	A (ppb)	RMSD (ppb)	A^* (ppb)	years fit
New York/maximum	53 \pm 6	43 \pm 9	3.9	53 \pm 2	2000-2017
New York/rural upwind	42 \pm 7	50 \pm 10	5.1	44 \pm 2	2000-2017
New Jersey/all sites	43 \pm 4	57 \pm 6	4.6	54 \pm 2	2000-2017
Connecticut/all sites	56 \pm 5	40 \pm 7	5.0	55 \pm 2	2000-2017
Connecticut/coastal	61 \pm 6	36 \pm 8	4.1	57 \pm 3	2000-2017
Rhode Island/all sites	49 \pm 8	44 \pm 12	4.0	49 \pm 3	2000-2017
Massachusetts/Boston	46 \pm 6	27 \pm 6	3.1	27 \pm 2	1990-2017
Massachusetts/suburban	41 \pm 10	52 \pm 14	3.3	45 \pm 3	2000-2017
Massachusetts/coastal	44 \pm 9	52 \pm 13	3.2	49 \pm 3	2000-2017
New Hampshire/coastal	49 \pm 6	35 \pm 8	3.7	38 \pm 2	1995-2017
New Hampshire/northwest	45 \pm 6	29 \pm 9	3.7	28 \pm 2	2000-2017
New Hampshire/Mt. Washington	66 \pm 7	8 \pm 8	2.9	---	1993-2017
Vermont /all sites	46 \pm 7	34 \pm 10	2.7	33 \pm 2	2000-2017
Maine/interior	44 \pm 8	23 \pm 10	5.8	21 \pm 3	1990-2017
Maine/NE coast	47 \pm 5	22 \pm 5	2.0	23 \pm 2	1991-2017
Maine/SW coast	49 \pm 5	36 \pm 5	4.1	39 \pm 2	1990-2017
Maine/Cadillac Mtn.	52 \pm 16	36 \pm 20	5.2	44 \pm 5	1997-2017

David Parrish 1/4/2019 4:00 PM
Deleted: 57

David Parrish 1/4/2019 4:02 PM
Deleted: 45

10 **Table 3. Results of least-squares fits of Eq. 1 to the state maximum ODV's illustrated Figure 11; y_0 and τ were held constant at 45.8 ppb and 21.9 years, respectively. The absolute root-mean-square deviations between the observed ODV's and the derived fits are indicated. Year_{NAAQS} indicates the projected year that the fit to the state maximum ODV dropped to, or will drop to the NAAQS of 70 ppb.**

State	A^* (ppb)	RMSD (ppb)	Year _{NAAQS}
Connecticut	61 \pm 7	5.8	2021
Maine	48 \pm 4	3.2	2015
Massachusetts	53 \pm 5	3.9	2017
New Hampshire	43 \pm 4	3.0	2013
New Jersey	64 \pm 5	3.7	2021
New York	58 \pm 4	3.0	2019
Rhode Island	52 \pm 4	3.4	2017
Vermont	35 \pm 3	2.1	2008



5 **Figure 1:** Maps of the northern U.S. with all ozone monitoring sites indicated by grey circles. The colored lines indicate the four regions considered: three Pacific Northwest states (blue dashed), three rural western states (green solid), three midwestern states (dotted orange), and eight northeastern states (purple dashed).




Figure 2: Time series of all ODVs (grey symbols) reported from all monitoring sites in the four northern U.S. regions shown in Figure 1. The numbers of monitoring sites and reported ODVs are annotated for each region. The red symbols give the averages and 2- σ confidence limits for all ODVs reported in each year. For comparison, the blue curve in each panel indicates a fit to the time history of the maximum ODVs recorded in the Los Angeles urban area (Parrish et al., 2017). The dotted line indicates the 2015 NAAQS of 70 ppb.




Figure 3: Topographical map of the three rural western states with symbols indicating the locations of the monitoring sites. The two colored symbols indicate two long-term sites in national parks that are discussed in detail. (Note that Yellowstone National Park is located in Wyoming, but is nevertheless considered here.)

5




Figure 4: Time series of all ODVs (grey symbols) reported from all monitoring sites in three rural western states, plus Yellowstone NP located in Wyoming. The two sets of colored symbols are results from two long-term sites in national parks. The curves are fits of Eq. 1 to all data from the three states (black curves), and to the national park data in the respective colors.

5




Figure 5: Topographical map of the eight northeastern states with symbols indicating the locations of the ozone monitoring sites. Three groups of colored symbols indicate groups of sites that are discussed in detail. The inset gives the abbreviations for each of the eight states.




Figure 6: Time series of all ODVs (grey symbols) reported from all monitoring sites in New York and Connecticut. The three sets of colored symbols indicate the results from groups of sites that are discussed in detail. The curves are fits of Eq. 1 to respective colored symbols, and to all data points for Connecticut.

5




Figure 7: Time series of all ODVs (grey symbols) reported from all monitoring sites in Massachusetts and Maine. The four sets of colored symbols indicate the results from groups of sites that are discussed in detail. The curves are fits of Eq. 1 to respective colored symbols.




Figure 8: Comparison of fits of the ODVs shown in Figures 6, 7 and S1-S8 to Eq. 1. The parameters of these fits are included in Table 2.

5





Figure 9: Approximate contour plot of the enhancement of ozone design values due to photochemical production from U.S. anthropogenic emissions in the year 2000, estimated from the A^* values given in Table 2.



5 Figure 10. Comparison of observed ODVs color-coded by year with those calculated from Eq. 1 for a) all monitoring sites and b) for the maximum observed in each state. The dashed lines indicate the 1:1 relationships with y_0 indicated by the larger circle, and the dotted lines the NAAQS. The number of data, square of the correlation coefficient, and the root-mean-square difference between the observed and calculated ODVs for 2000-2017 are annotated.





Figure 11: Time series of maximum ODVs reported from any site within each of the eight northeastern states. The solid curves colors are fits of Eq. 1 to respective colored symbols for the 2010-2017 period. The derived A^* values from these latter fits are given in Table 3. The dashed lines are projections of the solid curves.

5



10 Figure 12. Comparison of maximum observed ODVs in the New York City and Los Angeles urban areas. a) Temporal trend of observations (symbols) and fit to Eq. 1, including extrapolation to infinite time; annotations indicate year that extrapolations decrease to 70 ppb. The New York City results are the maxima from either the states of New York or New Jersey, and the Los Angeles results are those for the South Coast Air Basin (Figure 8 of Parrish et al., 2017). b) Bar graph indicating maximum ODVs in 2000 and 2015 (hatched bars) and the estimated U.S. background ODV (solid bars); the maximum ODVs are derived from the fits to Eq. 1 included in a).
15