

Integration of Airborne and Ground Observations of Nitryl Chloride in the Seoul Metropolitan Area and the Implications on Regional Oxidation Capacity During KORUS-AQ 2016

Daun Jeong¹, Roger Seco¹, Dasa Gu^{1,a}, Youngro Lee², Benjamin A. Nault^{3,4}, Christoph J. Knote⁵, Tom Mcgee⁶, John T. Sullivan⁶, Jose L. Jimenez^{3,4}, Pedro Campuzano-Jost^{3,4}, Donald R. Blake¹, Dianne Sanchez¹, Alex B. Guenther¹, David Tanner², L. Gregory Huey², Russell Long⁷, Bruce E. Anderson⁸, Samuel R. Hall⁹, Kirk Ullmann⁹, Hye-jung Shin¹⁰, Scott C. Herndon¹¹, YoungJae Lee¹⁰, Danbi Kim¹⁰, Joonyoung Ahn¹⁰, and Saewung Kim¹

¹Department of Earth System Science, University of California, Irvine, Irvine, CA, USA

²School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA

³Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO, USA

⁴Department of Chemistry, University of Colorado, Boulder, CO, USA

⁵Meteorologisches Institut, Ludwig-Maximilians-Universität München, München, Germany

⁶Atmospheric Chemistry and Dynamics Laboratory, NASA GSFC, Greenbelt, MD, USA

⁷Office of Research and Development, U.S. EPA, Research Triangle Park, NC, USA

⁸NASA Langley Research Center, Hampton, VA, USA

⁹National Center for Atmospheric Research, Boulder, CO, USA

¹⁰National Institute of Environmental Research, Incheon, South Korea

¹¹Aerodyne Research Inc., Billerica, MA, United States

^acurrently at: Division of Environment and Sustainability, Hong Kong University of Science and Technology, Hong Kong, China

Correspondence: Saewung Kim (saewung.kim@uci.edu)



Abstract. Nitryl chloride (CINO₂) is a radical reservoir species that releases chlorine radicals upon photolysis. An integrated analysis of the impact of CINO₂ on regional photochemistry in the Seoul Metropolitan Area (SMA) during the Korean-United States-Air Quality (KORUS-AQ) 2016 field campaign is presented. Comprehensive multiplatform observations were conducted aboard the NASA DC-8 and at two ground sites (Olympic Park, OP; Taehwa Research Forest, TRF), representing an urbanized area and a forested suburban region, respectively. Positive correlations between daytime Cl₂ and CINO₂ were observed at both sites, the slope of which were dependent on O₃ levels. The possible mechanisms are explored through box model simulations constrained with observations. The overall diurnal variations of CINO₂ at both sites appeared similar but the night time variations were systematically different. For about half of the observation days at the OP site the level of CINO₂ increased at sunset but rapidly decreased at around midnight. On the other hand, high levels were observed throughout the night at the TRF site. Significant levels of CINO₂ were observed at both sites for 4-5 hours after sunrise. Airborne observations, box model calculations, and back trajectory analysis consistently show that these high levels of CINO₂ in the morning are likely from vertical or horizontal transport of air masses from the west. Box model results show that chlorine radical initiated chemistry can impact the regional photochemistry by elevating net chemical production rates of ozone by ~ 25 % in the morning.

1 Introduction

Nitryl chloride (CINO₂) is a night time radical reservoir that generates chlorine radicals (Cl[•]) upon sunrise (R1), with a lifetime (τ_{CINO_2}) of around 30 minutes at midday in the northern hemisphere mid-latitude summer, under clear sky conditions ($J_{CINO_2} \approx 5.47 \times 10^{-4} \text{ s}^{-1}$, (Madronich and Flocke, 1998)). It is produced through heterogeneous reaction of chloride (Cl⁻) containing aerosols and dinitrogen pentoxide (N₂O₅(g)), which is generated from an equilibrium reaction with gas-phase nitrate radical (NO₃) and nitrogen dioxide (NO₂) (R 2-4, (Finlayson-Pitts et al., 1989)). In acidic aerosols (\approx pH 1.8), uptake of N₂O₅ (g) can also produce gas-phase chlorine (Cl₂, R5), resulting from enhanced CINO₂ uptake coefficient of up to 3 - 4 orders of magnitude higher than neutral pH (Roberts et al., 2008). However, this reaction has yet to be proven in ambient conditions.

During the day, N₂O₅ exists at low levels due to its thermal instability (Malko and Troe, 1982) and the short lifetime of NO₃ ($\tau_{NO_3} < 5$ s) from photolysis and reaction with NO (Wayne et al., 1991). Particulate Cl⁻ and chlorine containing gas species can come from both natural sources such as sea salt and biomass burning (Blanchard, 1985; Woodcock, 1953), and anthropogenic sources such as steel making, incineration, bleaching processes, and coal-fired power plants (Hov, 1985; Reff et al., 2009; Tanaka et al., 2000; Lee et al., 2018; Fu et al., 2018). The efficiency of CINO₂ production depends on heterogeneous loss of N₂O₅, which is a function of the N₂O₅ aerosol uptake coefficient ($\gamma_{N_2O_5}$), aerosol surface area, and N₂O₅ mean molecular speed, as well as the yield of CINO₂ (ϕ_{CINO_2}) (e.g., Thornton et al. 2003; Schweitzer et al. 1998; Behnke et al. 1997; Hu and Abbatt 1997; Bertram and Thornton 2009). Many recent studies have reported discrepancies between field derived and laboratory parameterized $\gamma_{N_2O_5}$ (e.g., Brown et al. 2009; Chang et al. 2016; Morgan et al. 2015; Phillips et al. 2016; McDuffie et al. 2018b; Tham et al. 2016; Wang et al. 2017a, b) and ϕ_{CINO_2} (e.g., McDuffie et al. 2018a; Riedel et al. 2013; Ryder et al. 2015; Tham et al. 2018; Thornton et al. 2010; Wagner et al. 2013; Wang et al. 2017a, b). In a nocturnal boundary layer, CINO₂ can accumulate to significant levels due to its long lifetime ($\tau_{CINO_2} > 30$ h) with slow

loss mechanisms through heterogeneous uptake (Behnke et al., 1997; Frenzel et al., 1998; George et al., 1995). At sunrise, CINO₂ rapidly photolyzes to generate chlorine radicals (Cl[·]), which can react with most volatile organic compounds (VOCs). For alkanes, Cl[·] has up to 1 - 2 orders of magnitude larger rate constants than hydroxyl radicals (e.g., $k_{OH+n-C_4H_{10}} = 2.4 \times 10^{-12}$, $k_{Cl+n-C_4H_{10}} = 2.2 \times 10^{-10}$ at 298 K) (Atkinson, 1997; Atkinson and Arey, 2003). Therefore, Cl[·] can potentially influence the radical pool (HO_x-RO_x) and ozone (O₃) level, which can also affect the formation of secondary aerosols. This influence can be most prominent in the morning when concentrations of other oxidants are low (i.e., NO₃ and ·OH) (Finlayson-Pitts, 1993; Hov, 1985; Young et al., 2014).



The first ambient measurements of CINO₂ were carried out by Osthoff et al. (2008), from a ship sampling along the southeastern U.S. coast in 2006. In that study, CINO₂ was observed up to ~1 ppbv at night time, particularly during the time period influenced by urban pollution and ship plumes of the Houston ship channel. Since then, a growing number of measurements reported significant levels of CINO₂, especially in polluted coastal regions with sources from natural and anthropogenic chloride and nitrogen oxides. Riedel et al. (2012) measured up to ~2 ppbv of CINO₂ off the coast of Santa Monica Bay, on board the research vessel Atlantis. Recent studies show that high levels of CINO₂ are also present in mid-continental regions. Thornton et al. (2010) measured up to ~ 400 pptv in Boulder, Colorado, which is ~ 1,400 km away from the coastline. Mielke et al. (2011) reported up to ~ 250 pptv in Calgary, Alberta, Canada, during spring, which is ~ 800 km from the coastline. Back trajectory analysis results showed that the observations were most likely not influenced by marine air masses. More recently (in the past 5 years), increasing number of CINO₂ observations have been conducted in Asia consistently showing significant levels of CINO₂ present in the boundary layer (e.g., Tham et al. 2018, 2016; Wang et al. 2016, 2017b, 2014; Yun et al. 2018; Liu et al. 2017). CINO₂ observations at semi-rural (Wangdu of Hebei province) and urban (Hong Kong, Jinan) regions in China have measured up to 2 ppbv and 776 pptv respectively (Tham et al., 2016; Wang et al., 2017a). At the mountain top (957 m above sea level) in Hong Kong, up to 4.7 ppbv of CINO₂ was reported (Wang et al., 2016). The high levels of CINO₂ in these studies were mostly correlated with continental pollution in vicinity (e.g., power plant plumes, biomass burning). A recent study by Yun et al. (2018) reported the highest-recorded mixing ratio of CINO₂ (8.3 ppb), during a severe haze event in a semi-rural site downwind of the Pearl River Delta in the winter. Overall, observations have shown that CINO₂ is ubiquitous in the tropospheric boundary layer.

However, measurements are still limited, as discrepancies remain between global chemical transport models and observations. Uncertainties in model simulated CINO₂ can arise from limited emission inventories, low resolution of the grid,

uncertainties in $\gamma_{N_2O_5}$ and ϕ_{ClNO_2} parameterization, complexity of the terrain, and meteorological conditions and these have been dealt in previous studies (e.g., Zhang et al. 2017; McDuffie et al. 2018b, a; Lowe et al. 2015; Sarwar et al. 2012, 2014; Sherwen et al. 2017). For instance, smoothing out local $ClNO_2$ peaks by diluting local NO_x emissions, will result in limited NO_3 and N_2O_5 production. According to Sarwar et al. (2012, 2014), the Community Multiscale Air Quality (CMAQ) model with a finer grid (i.e. 12 km) simulated $ClNO_2$ that corresponded better to the observations, compared to the model runs with coarser grid size (i.e., 108 km), embedded with similar chemistry. Another modeling study by Sherwen et al. (2017) compared the $ClNO_2$ levels between the GEOS-Chem simulations and observations in inland areas (i.e., London, UK and a mountain top near Frankfurt, Germany) during the summer of 2015. Compared to observations, the simulations underestimated the $ClNO_2$ maxima levels by ~ 7 times in inland areas (Sherwen et al., 2017). Modeling studies have consistently suggested the significance of Cl^- initiated reactions in regional and global O_3 production and in the lifetime of VOCs in the troposphere (Knipping and Dabdub, 2003; Tanaka et al., 2000, 2003; Sarwar et al., 2014; Sherwen et al., 2016; Simon et al., 2009). Sarwar et al. (2014) explored the production of $ClNO_2$ from sea salt and biomass burning and its impact in the Northern Hemisphere by including $ClNO_2$ formation chemistry in the CMAQ model. The results showed that, compared to the simulations without $ClNO_2$ formation, monthly 8 h wintertime maximum O_3 and $\cdot OH$ increased up to 15 % and 20 %, respectively. The impact was the largest in China and Western Europe. In the Hong Kong-Pearl River Delta (HK-PRD) region, Li et al. (2016) simulated up to ~ 1 ppbv of $ClNO_2$ originating from sea salt, biomass burning, and anthropogenic emissions (e.g., coal combustion) with the Weather Research and Forecasting coupled with Chemistry (WRF-CHEM) model. This resulted in ~ 16 % O_3 increase in the planetary boundary. Another modeling study of WRF-CHEM embedded with an updated chlorine chemistry, simulated 3-6 % of surface O_3 increase in the North China Plain and Yangtze River Delta during the summer (Zhang et al., 2017). A recent study by Wang et al. (2019) updated the standard version of the GEOS-Chem (Chen et al., 2017; Sherwen et al., 2016) to better track partitioning between aerosol chloride and gas-phase chlorine species. Comparison between their model simulations with and without $ClNO_2$ production showed enhanced O_3 up to 8 ppb during the winter season in Europe.

East Asian countries are of particular interest due to the rapid economic growth in the past decades with high anthropogenic emissions from densely populated megacities (e.g., Shanghai, Guangzhou, Beijing, Tokyo, Seoul). The majority of the world's megacities are situated in coastal regions (Neumann et al., 2015) with high NO_x emissions and abundant sources of chloride from both anthropogenic and natural origin. These regional characteristics likely promote $ClNO_2$ production. Moreover, considering that nearly half the population in the world lives near the coast, defined as < 100 km from coastline (Hinrichsen, 1998), a careful evaluation of the impact of $ClNO_2$ on local tropospheric chemistry is crucial. In this study, we present $ClNO_2$ observation results from the Korean - United States Air Quality (KORUS-AQ) study conducted in the Seoul Metropolitan Area (SMA), South Korea during late Spring (May 2 to June 12, 2016). The field campaign was an international collaboration between the National Institute of Environmental Research (NIER) of South Korea and the National Aeronautics and Space Administration (NASA) of the United States with the aim to better understand the impact of a megacity on regional air quality. A comprehensive suite of measurements were deployed at two super sites (Olympic Park site, OP; Taewha Research Forest, TRF) and aboard the NASA DC-8 to make airborne observations over the South Korean peninsula and the Yellow Sea. We present observational and box model results to evaluate the impact of $ClNO_2$ towards regional air quality in SMA.

2 Methods

2.1 KORUS-AQ 2016 Field Campaign and Observation Sites

We present observations carried out at Olympic Park, (OP; lat:37° 30' 32.904" N, lon:127° 7' 20.136" E), Taehwa Research Forest (TRF; lat: 37° 19' 14.484" N, lon:127° 18' 32.58" E) and on the NASA DC-8. The two ground sites were within the SMA region, which is the second largest metropolitan area in the world with a population of ~ 24 million (Park et al., 2017). As shown in Figure 1(a), the OP site is located in the southern part of Seoul, surrounded by high rise residence buildings and close to major freeways. The TRF site is in the middle of a forested area, ~ 26 km southeast of the OP site. Previous studies have shown that the TRF site is affected by both aged anthropogenic air masses from the city and fresh biogenic emissions from the forest (Kim et al., 2016, 2015). Both sites were ~ 50 km to the east of the nearest coastline. Figure 1(b) shows the flight tracks of the NASA DC-8, during the KORUS-AQ campaign. Spiral patterns were conducted near the TRF site to measure a vertical profile of the troposphere. Airborne observations were carried out during the daytime, between 8:00 and 17:00 local time. A summary of the analytical techniques of the measurements presented in this study are shown in Table 1. Meteorology during the observation period can be classified into dynamic (May 4th - 16th), stagnation (May 17th - 22nd), transport (May 25th - 31st), and blocking period. During the stagnant period, high pressure system was persistent in the Korean peninsula resulting in local air masses to be more dominant within the SMA compared to the dynamic and transport (May 25nd - 31st) periods. Rex block patterns were observed during the blocking period (June 1st - 6th). During this period, a high pressure system was adjacent to a low pressure over the Korean peninsula resulting in more local influence with occasional stagnation.

2.2 Chemical Ionization Mass Spectrometry and Calibration

A THS Instruments LLC Chemical Ionization Mass Spectrometer (CIMS), using iodide (I^-) as the reagent ion was used for measuring Cl_2 and $CINO_2$ at the two ground supersites and on the NASA DC-8. The system was similar to what is described in Slusher et al. (2004) and Liao et al. (2011), and the inlet configuration during the campaign is shown in Figure S1. Ambient air was sampled through a stainless steel donut shaped inlet at TRF and a Polytetrafluoroethylene (PTFE) tube inlet at OP. The stainless steel donut inlet has been shown to effectively avoid wall loss of reactive halogens during previous campaigns (Liao et al., 2011). The lengths of the inlet lines of the three CIMS systems were 20 - 30 cm. The PTFE inlet line at the TRF site was washed on a weekly basis and the ones at the OP and DC-8 were not washed routinely during the campaign due to difficulties on detaching the inlet. The potential bias of interactions of Cl_2 and $CINO_2$ inside the inlet were not tested but the artifacts have been shown to be negligible in various field conditions (Riedel et al., 2012; Thornton et al., 2010; Liao et al., 2014). Therefore, the use of different types of inlets (e.g., the use of the donut), described above, at the two ground sites and on the DC-8 is not expected to be an issue for the quantitative comparisons in this study. The sampled air went through the first 3-way valves to be delivered to an ambient or charcoal scrubber mode for background, alternating every 5 minutes. The second 3-way valve was for heated (150 °C) and unheated cycles. $CINO_2$ and Cl_2 were only quantified during the unheated cycles to avoid any potential artifacts as described in Liu et al. (2017). A total of 3000 standard liters per minute (slpm) was drawn in with a blower with an additional flow of 4 slpm drawn at the end of the inlet to reduce the residence time and 1 slpm was sampled into the CIMS. All

the inlet parts, after the blower, including the fittings and tubings, were made of PTFE. In the flow tube, the target compounds
135 form clusters with I^- (R 5-6, (Huey, 2007; Huey et al., 1995; McNeill et al., 2006)), which were generated by flowing 1 slpm
 N_2 through a methyl iodide (CH_3I) permittube oven maintained at 50 °C. Polonium (NRD LLC, Static Master, Model: 2U500,
Activity: 20 mCi) was used as the radioactive source for ionization. Clusters of Cl_2 isotopes were detected at the mass to charge
ratio (m/z) of 197 and 199, and $ClNO_2$ was measured at 208 and 210. The natural abundance of Cl_2 and $ClNO_2$ isotopes are
approximately 9:6:1 ($^{35}Cl^{35}Cl : ^{35}Cl^{37}Cl : ^{37}Cl^{37}Cl$) and 3:1 ($^{35}ClNO_2 : ^{37}ClNO_2$) respectively. Mass 201 ($^{37}Cl^{37}Cl$) was not
140 considered in the data processing due to artifacts.



Calibrations of Cl_2 and $ClNO_2$ were carried out during and after the campaign. Cl_2 in a cylinder (Airgas, 10 ppm in N_2)
was diluted with zero air to be sampled in either ambient or scrubber (charcoal) mode (Figure S1). The Cl_2 in the cylinder
145 was quantified through the method described by Liao et al. (2012) and was 8.84 ± 0.43 ppm. $ClNO_2$ was synthesized, based
on Thaler et al. (2011). Briefly, Cl_2 gas in N_2 was passed through a pyrex reservoir (diameter = 1.3 cm, length = 5.5 cm)
containing a bed of NaCl (MACRON) and $NaNO_2$ (Sigma Aldrich) with a molar ratio of 10 to 1. This slurry mixture contains
 NO_2^- that reacts with the flowing Cl_2 to generate $ClNO_2$. The output flow was further diluted with 4 L min^{-1} of zero air
in order to sufficiently provide gas flow. The flow containing synthesized $ClNO_2$ was then analyzed at m/z of 208 and 210
150 with the CIMS. NO_2 and $NO-NO_y$ were simultaneously measured with a Cavity Ring Down Spectroscopy (CRDS, Los Gatos
Research, detection limit: 10 pptv, precision: 50 pptv at 1σ , model: 907-0009-0002) and chemiluminescence (CL, Thermo
Scientific, detection limit: 50 pptv, model: 42 i) respectively. $ClNO_2$ is detected as NO_y in the CL through conversion to NO
on the heated (325 °C) molybdenum catalytic converter (Williams et al., 1998). The efficiency of the conversion was assumed
to be unity. Therefore, $ClNO_2$ could be determined by comparing the three instruments and subtracting the byproducts (HONO
155 and NO_2) from the total NO_y . The averaged sensitivity of Cl_2 was 31.5 ± 11.2 Hz/ppt and $ClNO_2$ was 19.7 ± 1.5 Hz/ppt. The
2 sigma detection limits of Cl_2 and $ClNO_2$ were 2.9 and 1.5 ppt, respectively, over 30 min.

2.3 Modeling

We used Framework for 0-D Atmospheric Modeling (FOAM v3.1) for simulating three types of simulations: 1) daytime Cl_2
production (Figure 5), 2) in-situ $ClNO_2$ production in the morning (Figure 8), and 3) testing the impact of measured $ClNO_2$ on
160 the regional tropospheric chemistry (Figure 10). FOAM is a MATLAB based open-source box model. Detailed descriptions of
the model can be found in Wolfe et al. (2016). Each step of the model was constrained with the averaged meteorology param-
eters (e.g., pressure, temperature, relative humidity) and trace gases observed in the two ground sites during the campaign. The
constrained trace gases include $ClNO_2$, Cl_2 , O_3 , NO, NO_2 , CO, CH_4 , and 20 non-methane hydrocarbons including 8 alkanes
(i.e., ethane, propane, iso-butane, n-butane, iso-pentane, n-pentane, n-hexane, and n-heptane), that have relatively high reaction
165 rate constants with Cl. A constant meteorology and trace gas observation set, collected at the corresponding time period, were

constrained throughout the 72-hour model simulation presented in Figure 5. Then, the Cl_2 concentrations at the end of the 72-hour simulation are compared to simultaneously observed mixing ratios of ClONO_2 in Figure 5. Simulations in Figure 8 were similarly constrained as those in Figure 5 but allow ClONO_2 concentrations to vary with time in order to assess ClONO_2 production predicted by the model. The model simulation presented in Figure 10 was constrained with a diurnal variation of the parameters. A full diurnal cycle of the model was for 24 hours consisting a total of 864 steps and each step was integrated for 100 seconds. Each step of the model was constrained with observations measured at that time of day. To assess the impact of ClONO_2 chemistry on net O_3 production, all species were constrained except for NO_2 and O_3 , which were initialized with observed values and allowed to vary in time. Photolysis rate constants were derived through the hybrid method (Wolfe et al., 2016) in the F0AM box model. This method uses clear sky solar spectra from the tropospheric ultraviolet and visible radiation model (TUV v 5.2) and cross sections and quantum yields suggested by IUPAC. To capture the effects of pollution on photolysis rates, the ratio of the measured J_{NO_2} to the F0AM modeled J_{NO_2} was calculated. This ratio was then applied to other photolysis rate constants calculated in the model. Measured J_{NO_2} was taken from the DC-8 actinic flux measurements (Charged-coupled device Actinic Flux Spectroradiometer; CAFS) when flying near SMA at altitudes under 1 km. A diurnal cycle was applied to the DC-8 measurement to determine j -values at other times of day. Photolysis rate constants of ClONO_2 , Cl_2 , and ClONO_2 were not present in the F0AM model and therefore taken directly from the DC-8 measurements to be used in the model runs in this study. The Master Chemical Mechanism v3.3.1 (MCM) was taken from <http://mcm.leeds.ac.uk/MCM> and embedded in the box model. MCM v3.3.1 has a detailed gas photochemistry (i.e., 5832 species and 17224 reactions), including the oxidation of CH_4 and 142 non-methane primary emitted VOCs (Jenkin et al., 2015). Since MCM v3.3.1 only includes $\text{Cl}\cdot$ reactions with alkane species, additional chlorine chemistry was embedded in the model, similar to what Riedel et al. (2014) reported. This was done by including multiple $\text{Cl}\cdot$ precursors (e.g., Cl_2 , ClONO_2 , HCl , ClONO_2 , HOCl) and $\text{Cl}\cdot$ reactions with non-alkane VOCs, such as alkene, alcohol, aromatics, alkynes, ketones, organic acids and nitrates. All the reactions embedded in the model can be found in the supplementary of Riedel et al. (2014) and Wolfe et al. (2016). Boundary layer height, emissions, and depositions were not considered in the model. More details on the setup of the box model are in the supplement material (S3). The FLEXible PARTi-cle dispersion model (FLEXPART v9.1, <https://www.flexpart.eu>) was used for the air mass source contribution (Figure 3) and backward trajectory analysis (Figure 9). The backward trajectories reported in our study were initialized 9:00 LST at TRF, following it 24 hours back in time. Only the center of the mass-weighted particles are shown in Figure 9 and clusters are included in the supporting information. These clusters represent fractional contributions of air masses (Figure S10). The trajectories were driven by the National Centers for Environmental Prediction (NCEP) Global Forecast System (GFS) with a 0.25 degree resolution. Influence of air mass originating from the ocean at TRF and OP was calculated every 6 hours following an air mass 5 days back in time. Meteorology was driven by WRF with a 5 km horizontal resolution. Since emissions of CO are very low in the ocean, and assumed to be inert in the model, it was used as a tracer for contribution of air originating from the ocean within a given air mass at each ground site.

3 Results and Discussions

3.1 ClNO₂ Observations

200 Figure 2 shows the temporal variation of trace gases measured during the campaign at (a) the OP site (May 17th - June 11th) and (b) the TRF site (May 5th - June 11th). The OP site, which was located near heavy traffic, showed high levels of NO_x throughout the campaign. During most nights (except for May 24th - 26th, 30th - 31st, and June 6th - 7th), O₃ was completely titrated by NO. On the other hand, at the TRF site, which is a forested region downwind of the urban area, O₃ remained at ~ 30 ppbv throughout the night. During the measurement period, measurable amounts of ClNO₂ were observed at both ground
205 sites (Figure 3). The maximum observed ClNO₂ was ~800 pptv (5 min averaged) and ~2.5 ppbv (5 min averaged) at the OP and TRF sites, respectively. At both sites, ClNO₂ started accumulating at sunset and rapidly photolyzed upon sunrise, which was ~5:30 local standard time (LST) during the campaign. Nighttime relationship between ClNO₂ and Cl₂ varied day by day and did not show a clear correlation. This implies that the sources or loss processes of Cl₂ and ClNO₂ were not consistent at
210 coast of LA.

Daytime (11:00 - 18:00, LST) ClNO₂ was up to ~100 pptv at OP and ~250 pptv at TRF (Figure 4). The level showed a positive correlation with Cl₂, especially in relatively high O₃ conditions (> 50 ppbv). When O₃ was relatively low (< 50 ppbv), Cl₂ production was suppressed, while ClNO₂ was not necessarily limited. Excluding the days with low O₃ (i.e., May 26th and 29th for OP and May 6th, 29th, and June 4th for TRF), the relationship between daytime ClNO₂ and Cl₂ showed
215 positive correlation with R² of 0.49 and 0.80 for OP and TRF, respectively. This positive correlation is consistent with the results reported by Liu et al. (2017) in the North China Plain. In their study, up to ~ 450 pptv of both Cl₂ and ClNO₂ was measured during the daytime (10:00 - 20:00, LST), with strong correlation of R² = 0.83. Cl₂ levels were also suppressed in low O₃ and OH conditions during low solar radiation periods. Therefore, the authors suggested that daytime Cl₂ levels could be positively related to photochemical activities. Considering the short lifetime of Cl₂ and ClNO₂ during the day (i.e., 11:00 -
220 18:00 LST in our study), the levels we observed are likely affected through local production. According to Liu et al. (2017), the air mass showed moderate correlation to SO₂ with possible influences from power plants. However, in this study, the ClNO₂ measured at both the OP and TRF sites was weakly correlated with SO₂ (R² = 0.02), which implies that the air masses that we sampled are not fresh emissions from coal combustion activities such as power plants.

The first possibility we explored is the direct generation of Cl₂ from reactions in acidic particles. ClNO₂ is very insoluble
225 ($\gamma_{ClNO_2} \approx 10^{-6}$ (Rossi, 2003)) in near-neutral pH. However, according to Roberts et al. (2008), γ_{ClNO_2} can increase up to 3 orders of magnitude in acidic surfaces (~ pH 1.8) leading to direct production of gas-phase Cl₂. Aerosol acidity was mostly below pH 2 during the campaign, based on thermodynamic calculations, constrained with airborne observations (Figure S3). Therefore, the efficiency of this reaction in ambient conditions requires further investigation. Another possibility is the autocatalytic production of Cl₂ from heterogeneous reactions of gas-phase ClONO₂ (i.e., $ClONO_{2(g)} + Cl^{-}_{(aq)} + H^{+}_{(aq)}$
230 $\rightarrow Cl_{2(g)} + HNO_3$, (Gebel and Finlayson-Pitts, 2001; Deiber et al., 2004)) and HOCl (i.e., $HOCl_{(g)} + Cl^{-}_{(aq)} + H^{+}_{(aq)} \rightarrow Cl_{2(g)} + H_2O$, (Vogt et al., 1996)) on particles. These reactions are also favored as particle acidity increases. In order to further

investigate its possibility, daytime Cl_2 was simulated by constraining the box model with measurements of ClONO_2 and other trace gases corresponding to each data point in Figure 4. Based on the availability of parameters, we were able to simulate 1680 and 1229 runs for the OP and TRF, respectively. This corresponds to more than 96 % of the daytime data points shown in Figure 4. γ_{ClONO_2} and γ_{HOCl} were set to 0.06 (Deiber et al., 2004; Hanson et al., 1994; Hanson and Ravishankara, 1994), which is an upper-limit of previous laboratory studies, and the yields were assumed to be unity. HCl generation from hydrogen abstraction of VOCs by Cl^\cdot were included in the mechanisms used in the model runs. The end points of the 72 hour simulation results are presented in Figure 5. As shown in the figure, the box model simulations were able to reproduce the positive correlation between Cl_2 and ClONO_2 . Moreover, modeled Cl_2 was suppressed in low O_3 conditions, which corresponds to the observations. This can be explained by Cl^\cdot reacting with O_3 , producing ClO^\cdot , leading to gas-phase ClONO_2 and HOCl production. These can react on acidic aerosols to generate Cl_2 . Sources of Cl^\cdot could be from photo-labile gas-phase chlorine compounds (e.g., Cl_2 , ClONO_2 , ClONO , HOCl) or oxidation of gas-phase HCl by OH . Although the reaction between HCl and OH is relatively slow ($k = 7.86 \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ at 298K, (Atkinson et al., 2007)), it has been reported to be a significant source of Cl^\cdot in the daytime (Riedel et al., 2012). A sensitivity test was carried out by comparing modeled Cl_2 between runs with and without HCl production from oxidation of VOCs by Cl^\cdot (Figure S4 c,d). The results show that production of Cl_2 was suppressed by 40 - 70 % when HCl was not generated in the model. This significant contribution of gas-phase HCl as a Cl^\cdot source, should be an upper-limit as the deposition of HCl was not considered in the model. Nonetheless, our analysis leads us to conclude that the mechanisms we have explored could be the main contributors of the daytime Cl_2 production during KORUS-AQ.

3.2 Sources of ClONO_2

FLEXPART source contribution analysis shows that the level of ClONO_2 at the ground sites was highly correlated with the origin of the air mass (Figure 3). During the nights shaded in red in Figure 3 (OP: May 20th, 22nd, June 2nd, and 7th; TRF: May 11th, 19th-22nd, June 2nd, and June 6th-7th), there was limited production of ClONO_2 at the surface. These periods mostly corresponded to meteorological conditions of stagnation or blocking events, which both resulted in localized air masses to be more dominant with limited influence from the west coast. Stagnation events can be characterized by low wind speeds and increased atmospheric stability, possibly leading to enhanced levels of pollutants like NO_x . Previous studies have shown that stagnant conditions can result in enhanced levels of N_2O_5 driven by high ozone and NO_2 . However, ClONO_2 production was limited during stagnation events in this study. This is likely due to limited availability of chloride as shown in submicron particle measurements of aerosol mass spectrometer (AMS) at the ground site for OP and airborne over TRF (Figure 3). Whether the chloride is from the ocean or anthropogenic emissions is uncertain since large point sources, such as power plants or petrochemical facilities, are also present along the west coast of the SMA. On the nights of May 20th and May 22nd, rapid changes in air quality were observed with fast shifts in O_3 , SO_2 , and CO . This corresponded with changes in ClONO_2 and Cl_2 (Figure S7). These events suggest the importance of boundary layer advection in controlling the ClONO_2 levels in the region.

Different diurnal variations of ClONO_2 were observed between OP and TRF (Figure 6). The measurements were averaged over selected days (OP: May 18th-20th, 22nd, 23rd, 29th, June 4th; TRF: May 5th, 8th, 9th, 12th, 17th, 18th, 30th, June 8th, 10th) that showed these two distinct profiles at each site. The description on these profiles are further explained in the following

sentences. At the TRF site (Figure 6b), far from direct NO emissions, significant levels of ClNO₂ were sustained throughout the night during most of the observation period with rapid photolysis upon sunrise. On the other hand, at OP (Figure 6 (a)), ClNO₂ started to increase upon sunset, followed by a rapid drop at around 22:00 LST. The trend was consistent with slower nitrate radical production rate ($d[\text{NO}_3]/dt = [\text{NO}_2][\text{O}_3]k$, where $k = 3.52 \times 10^{-17}$ at 298 K, (Atkinson et al., 2004)) as O₃ was titrated to zero by NO close to midnight. The wind direction, SO₂, and CO did not correlate. This suppressed ClNO₂ production in urbanized regions with high NO levels, have also been reported by Osthoff et al. (2018). However, significant levels of N₂O₅ and ClNO₂ could have been present in the upper part of the surface layer as shown in previous studies (Baasandorj et al., 2017; Young et al., 2012; Yun et al., 2018). According to Baasandorj et al. (2017), O₃ was completely titrated at the surface in Salt Lake Valley, Utah, while elevated mixing ratios of N₂O₅ were observed at 155 meters above ground level, at a site along the valley wall. On the other hand, airborne measurements at the LA basin (Young et al., 2012) showed a relatively uniform ClNO₂ profile throughout the nocturnal boundary layer as O₃ did not change significantly within the observed altitude range (< 600 m). During the 2015 Megacity Air Pollution Study (MAPS, Seoul, 2015), a Cavity Ringdown Spectrometer (CRDS) was installed on top of the Seoul tower in May - June that measured N₂O₅, NO_x, and O₃ (Brown et al., 2017). The elevation of the measurement site was 360 m above sea level (ASL), allowing for sampling further away from direct NO emissions. In their study, the average nighttime O₃ mixing ratio was around 50 ppbv and N₂O₅ was observed most nights, with mixing ratios reaching up to 5 ppbv. Therefore, it is very likely that ClNO₂ levels higher than the surface measurements could have been present at higher elevation during the observation period.

At both sites, ClNO₂ levels started to increase or sustained after the first 2-3 hours of rapid net loss upon sunrise. In the morning, ClNO₂ positively correlated to Cl₂ levels, but did not follow the nitrate production rate at the site (Figure S8). Box model simulations, initially constrained with observed ClNO₂ level, showed rapid photolysis upon sunrise (Figure S5, red dashed line). At TRF, this corresponded to the measurements until 7 - 8 am LST, when a second ClNO₂ peak was observed (Figure 6b). This ClNO₂ peak in the morning was observed about half the observation days during the campaign. With the net ClNO₂ production rate from the observation, and the loss rate from the simulated ClNO₂ from photolysis, a production rate of 400 pptv h⁻¹ would be required to reconcile the observation. In the case of ClNO₂ observed on May 5th at TRF (an insert of Figure S6b), a maximum of 2.5 ppbv h⁻¹ of ClNO₂ production rate was required in the morning to reconcile the observations. At OP, 18 pptv h⁻¹ was required for the 7 averaged days. The ClNO₂ production rate required in the morning at TRF was much higher than the previous studies that have also reported high sustained levels of ClNO₂ in the morning (i.e., 20 - 200 pptv h⁻¹) (Faxon et al., 2015; Bannan et al., 2015; Tham et al., 2016). In these previous studies, three possibilities have been suggested that could explain the high sustained levels of ClNO₂ in the early morning : 1) in-situ generation of ClNO₂, 2) transport of ClNO₂ within the boundary layer, and 3) entrainment of ClNO₂ from residual layer. Each possibility is explored below.

In order to explore the possibility of in-situ formation, box model simulations of ClNO₂ production from heterogeneous reaction of N₂O₅ and chloride containing aerosols were conducted. N₂O₅ was calculated assuming a photo-stationary state of NO₃ (Brown et al., 2005). Aerosol surface area was taken from airborne observations over TRF. Based on the box model results in Figure 8, even with an assumption of 100 % yield, ClNO₂ from heterogeneous reaction was not able to reconcile the observed

level. Using the dry surface area for the first order loss of N_2O_5 on aerosols certainly could result in an underestimation of ClNO_2 production in the model. Kim et al. (2017, 2018) observed hygroscopic growth factor of less than 1.5 in the SMA region for particles below 150 nm during the KORUS campaign period. In other words, the discrepancy between observed and modeled ClNO_2 of more than 50-fold cannot be reconciled by this underestimation. The box model simulation on gas phase
305 production of ClNO_2 (i.e., $\text{Cl}\cdot_{(g)} + \text{NO}_{2(g)} + \text{M} \rightarrow \text{ClONO}_{(g)} + \text{M}$, $\text{Cl}\cdot_{(g)} + \text{NO}_{2(g)} + \text{M} \rightarrow \text{ClNO}_{2(g)} + \text{M}$) showed at most 2-10 pptv of ClNO_2 and ClONO (Figure S6).

Therefore, horizontal or vertical transport from local sources would be the most likely explanation for the high ClNO_2 in the morning. Although ClNO_2 readily photolyzes during the day ($\tau_{\text{ClNO}_2} \approx 30$ min at midday), the lifetime could be significantly long enough in the early morning to allow for transport of ClNO_2 to the ground sites. Based on the NCAR TUV v5.2 model,
310 the lifetime of ClNO_2 , averaged between 5:30 and 8:30 LST was ~ 2 hours under clear sky conditions. Figure 9 shows back trajectory analysis initiated at 9 am local time at TRF. At high ClNO_2 days with the morning peaks, most of the air masses were from the west. During KORUS, the DC-8 did not fly to the west of the SMA in the early morning. However, there are large point sources, such as petrochemical facilities and industries, and vehicular emissions to the west and south west of the SMA region. Sullivan et al. (2019) reported that this resulted in enhanced levels of O_3 in receptor regions (i.e., Taehwa Research
315 Forest) downwind when westerlies were prevalent. Therefore, favorable conditions such as high chloride content in aerosols from both anthropogenic and natural sources and high levels of NO_x - O_3 could have led to significant levels of ClNO_2 to build up and transported to TRF before being completely photolyzed. During the campaign, influence of large biomass burning was negligible as reported in Tang et al. (2018, 2019).

At night time, the nocturnal boundary layer is decoupled from the residual layer (Stull, 1988), where the pollution from
320 the previous day resides. Being removed from direct NO emissions near the surface, N_2O_5 can effectively accumulate in the residual layer, with the major loss process being heterogeneous reaction on aerosols. Therefore, high levels of NO_2 and O_3 formed during the day can be trapped in the residual layer resulting in significant levels of ClNO_2 persisting throughout the night. Figure 7, shows (a) regional and (b) vertical distribution of airborne ClNO_2 throughout the campaign in the morning (8:00 - 8:30 LST) over the SMA region (lat: $37^\circ 12' 0'' \text{ N} - 37^\circ 38' 60'' \text{ N}$, lon: $126^\circ 54' 0'' \text{ E} - 127^\circ 47' 60'' \text{ E}$). During 3 flights
325 (i.e., May 25th, May 31st, and June 10th), ClNO_2 was observed in the residual layer with a max of ~ 230 pptv. However, the remaining flights observed an average of 17 ± 56 pptv of ClNO_2 (black circles). Even the three days (i.e., May 25th, 31st, and June 10th), that ClNO_2 was observed in the residual layer, the level (max 230 pptv) could not reconcile the observed levels at the TRF site, which was 342 ± 330 pptv when averaged over the corresponding 3 days at 8:00 - 8:30 LST. However, it is possible that the air mass that was measured by the DC-8 was not representative of the air mass aloft at the west side of the
330 ground observation sites. Backtrajectory analysis initialized at 9:00 local time showed that the TRF site was affected by both the residual layer and below (Figure S10). The enhancement of O_3 and SO_2 concurrent to elevation of ClNO_2 could be due to the transport from the residual layer where pollution from high point sources from the other day was trapped within. From the current dataset, it would be difficult to derive a clear conclusion on whether the cause of the significant ClNO_2 in the morning was dominantly of transport from horizontal, vertical, or both.

335 3.3 Impacts of ClNO₂ on O₃

Cl· produced from ClNO₂ photolysis can influence the local air quality through reactions with VOCs followed by enhanced production of O₃. The possible impact of Cl· initiated reactions on the local chemistry were investigated by running box model simulations constrained with measured ClNO₂. A 24 hour diurnal variation of ClNO₂ was averaged over the same selected days as in Figure 6, and these were constrained throughout the model simulations. The results illustrate that when the model was constrained with ClNO₂ and Cl· initiated chemistry, higher levels of O₃ were simulated (Figure 10) compared to the base runs without ClNO₂. The averaged net O₃ production rate was enhanced by up to 2 % and 25 % at OP and TRF in the morning and by 1 % and 2 % when averaged during the day. The OP had 7 times lower Cl· than the TRF site due to low ClNO₂ levels (~60 pptv) in the morning. Since the box model simulations in our study did not take into consideration boundary layer height dynamics, emission, and deposition, this net production rate is the result of just chemical production and loss. For ·OH, the net production rate at TRF increased by 2 % in the morning. The results particularly from TRF are comparable with the previous study in the mountaintop site in Hong Kong, China (Wang et al., 2016). The enhancement of O₃ (max - min) was higher than their moderate ClNO₂ case (11 %) but lower than the high ClNO₂ plume case (41 %).

4 Conclusions

Comprehensive measurements of ClNO₂, Cl₂, other trace gases, and aerosol concentrations and properties have been conducted on the NASA DC-8 and at two ground sites during the KORUS-AQ 2016 field campaign. The observed averaged diurnal variations are largely consistent with the previous observations and our understanding on the photochemistry of ClNO₂. The presence of ClNO₂ was substantially suppressed during strong stagnation events, which could have prevented the transport of chloride near the coast. During the night, Cl₂ and ClNO₂ levels were not correlated while moderate to strong positive relationships were observed at daytime. Through box model simulations, we presented a quantitative analysis of the daytime observations. The results showed that heterogeneous reactions of ClONO₂ and HOCl in acidic aerosols may be responsible for the positive correlation between Cl₂ and ClNO₂, as well as its dependency on O₃. The second ClNO₂ peak in the morning, observed 4-5 hours after sunrise, required a significant source of ClNO₂ (up to 2.5 ppbv h⁻¹). Previous studies have attributed high sustained ClNO₂ in the morning to transport from the residual layer (Tham et al., 2016; Wang et al., 2016). In this study, box model runs of heterogeneous and gas-phase production of ClNO₂ could not reconcile the observed levels. Airborne observations near the ground sites in the early morning showed negligible ClNO₂ levels in the residual layer in most of the days. However, there is still a possibility of the contribution of vertical transport from the residual layer. Although the current data set is limited for us to pinpoint on the vertical locations (i.e., boundary layer v.s. residual layer), back trajectories illustrate that ClNO₂ rich air masses were mostly transported from the west, where there are significant sources of precursors. This shows that different meteorological or chemical conditions of the sites can lead to various causes of high ClNO₂ levels in the early morning. Finally, box model simulations constrained with observations suggest that Cl· initiated chemistry can lead up to ~25 % increase of net chemical O₃ production rate in the morning.

Data availability. Dataset used in this study is open to public and can be downloaded at <https://www-air.larc.nasa.gov/missions/korus-aq/>

Author contributions. DJ, RS, DG, YL, DT, SK, and GH designed and executed field measurements for collecting ClNO₂ and Cl₂; BAN, JLJ, and PCJ provided the airborne AMS data; CK ran the FLEXPART analysis; TM and JS provided O₃ measurements; DRB provided the WAS data; DS and AG provided the PTR-ToF-MS data; RL provided NO_x and O₃ data; BA provided the data from LARGE; SRH and KU provided data from CAFS; HS provided AMS data at OP; SH provided HCHO measurements at TRF; YL, DK, and JA provided CO measurements at TRF. DJ and SK prepared the original manuscript, and all other authors contributed in editing the manuscript.

Acknowledgements. This study is supported by NIER and NASA (NNX15AT90G). Also supported by Ministry of Science and ICT of South Korea. CU HR-AMS measurements and pH and aerosol liquid water calculations (BAN, PCJ, and JLJ) were supported by NASA grant NNX15AT96G and 80NSSC18K0630. We thank the Wisthaler research group (University of Oslo, University of Innsbruck) for providing airborne VOC data, John Crouse and Paul Wennberg (CALTECH) for the HNO₃ data, Hwajin Kim (Korea Institute of Science and Technology) for the discussions on aerosol composition, and Siyuan Wang (NCAR) for discussions on box model simulations. The authors appreciate logistical support from the research and supporting staff at Taehwa research forest operated by Seoul National University.

References

- 380 Atkinson, R.: Gas-Phase Tropospheric Chemistry of Volatile Organic Compounds: 1. Alkanes and Alkenes, *J. Phys.Chem. Ref. Data*, 26, 215–290, <https://doi.org/10.1063/1.556012>, 1997.
- Atkinson, R. and Arey, J.: Gas-phase tropospheric chemistry of biogenic volatile organic compounds: A review, *Atmos. Environ.*, 37, S197–S219, [https://doi.org/https://doi.org/10.1016/S1352-2310\(03\)00391-1](https://doi.org/https://doi.org/10.1016/S1352-2310(03)00391-1), 2003.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I - gas phase reactions of O_x, HO_x, NO_x, and SO_x species, *Atmos. Chem. Phys.*, 4, 1461–1738, <https://doi.org/10.5194/acp-4-1461-2004>, 2004.
- 385 Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume III - Gas phase reactions of inorganic halogens, *Atmos. Chem. Phys.*, 7, 981–1191, <https://doi.org/10.5194/acp-7-981-2007>, 2007.
- 390 Baasandorj, M., Hoch, S. W., Bares, R., Lin, J. C., Brown, S. S., Millet, D. B., Martin, R., Kelly, K., Zarzana, K. J., Whiteman, C. D., Dube, W. P., Tonnesen, G., Jaramillo, I. C., and Sohl, J.: Coupling between Chemical and Meteorological Processes under Persistent Cold-Air Pool Conditions: Evolution of Wintertime PM_{2.5} Pollution Events and N₂O₅ Observations in Utah’s Salt Lake Valley, *Environ. Sci. Technol.*, 51, 5941–5950, <https://doi.org/10.1021/acs.est.6b06603>, 2017.
- Bannan, T. J., Booth, A. M., Bacak, A., Muller, J. B. A., Leather, K. E., Breton, M. L., Jones, B., Young, D., Coe, H., Allan, J., Visser, S., Slowik, J. G., Furger, M., Prévôt, A. S. H., Lee, J., Dunmore, R. E., Hopkins, J. R., Hamilton, J. F., Lewis, A. C., Whalley, L. K., Sharp, T., Stone, D., Heard, D. E., Fleming, Z. L., Leigh, R., Shallcross, D. E., and Percival, C. J.: The first UK measurements of nitryl chloride using a chemical ionization mass spectrometer in central London in the summer of 2012, and an investigation of the role of Cl atom oxidation, *J. Geophys. Res. Atmos.*, 120, 5638–5657, <https://doi.org/10.1002/2014JD022629>, 2015.
- 395 Behnke, W., George, C., Scheer, V., and Zetzsch, C.: Production and decay of ClNO₂ from the reaction of gaseous N₂O₅ with NaCl solution: Bulk and aerosol experiments, *J. Geophys. Res. Atmos.*, 102, 3795–3804, <https://doi.org/10.1029/96JD03057>, 1997.
- 400 Bertram, T. H. and Thornton, J. A.: Toward a general parameterization of N₂O₅ reactivity on aqueous particles: the competing effects of particle liquid water, nitrate and chloride, *Atmos. Chem. Phys.*, 9, 8351–8363, <https://doi.org/10.5194/acp-9-8351-2009>, 2009.
- Blanchard, D.: The oceanic production of atmospheric sea salt, *J. Geophys. Res.*, 90, 961–963, <http://onlinelibrary.wiley.com/doi/10.1029/JC090iC01p00961/full>, 1985.
- 405 Brown, S. S., Osthoff, H. D., Stark, H., Dubé, W. P., Ryerson, T. B., Warneke, C., de Gouw, J. A., Wollny, A. G., Parrish, D. D., Fehsenfeld, F. C., and Ravishankara, A. R.: Aircraft observations of daytime NO₃ and N₂O₅ and their implications for tropospheric chemistry, *J. Photochem. Photobiol. A Chem.*, 176, 270–278, <https://doi.org/10.1016/j.jphotochem.2005.10.004>, 2005.
- Brown, S. S., Dubé, W. P., Fuchs, H., Ryerson, T. B., Wollny, A. G., Brock, C. A., Bahreini, R., Middlebrook, A. M., Neuman, T. A., Atlas, E., Roberts, J. M., Osthoff, H. D., Trainer, M., Fehsenfeld, F. C., and Ravishankara, A. R.: Reactive uptake coefficients for N₂O₅ determined from aircraft measurements during the Second Texas Air Quality Study: Comparison to current model parameterizations, *J. Geophys. Res.*, 114, 1–16, <https://doi.org/10.1029/2008JD011679>, 2009.
- 410 Brown, S. S., An, H. J., Lee, M., Park, J. H., Lee, S. D., Fibiger, D. L., McDuffie, E. E., Dubé, W. P., Wagner, N. L., and Min, K. E.: Cavity enhanced spectroscopy for measurement of nitrogen oxides in the Anthropocene: results from the Seoul tower during MAPS 2015, *Faraday Discuss.*, 200, 529–557, <https://doi.org/10.1039/C7FD00001D>, 2017.

- 415 Chang, W. L., Brown, S. S., Stutz, J., Middlebrook, A. M., Bahreini, R., Wagner, N. L., Dubé, W. P., Pollack, I. B., Ryerson, T. B., and Riemer, N.: Evaluating N_2O_5 heterogeneous hydrolysis parameterizations for CalNex 2010, *J. Geophys. Res.*, 121, 5051–5070, <https://doi.org/10.1002/2015JD024737>, 2016.
- Chen, Q., Schmidt, J. A., Shah, V., Jaeglé, L., Sherwen, T., and Alexander, B.: Sulfate production by reactive bromine: Implications for the global sulfur and reactive bromine budgets, *Geophys. Res. Lett.*, 44, 7069–7078, <https://doi.org/10.1002/2017GL073812>, 2017.
- 420 Colman, J. J., Swanson, A. L., Meinardi, S., Sive, B. C., Blake, D. R., and Rowland, F. S.: Description of the analysis of a wide range of volatile organic compounds in whole air samples collected during PEM-Tropics A and B, *Anal. Chem.*, 73, 3723–3731, <https://doi.org/10.1021/ac010027g>, 2001.
- Deiber, G., George, C., Le Calvé, S., Schweitzer, F., and Mirabel, P.: Uptake study of ClONO_2 and BrONO_2 by Halide containing droplets, *Atmos. Chem. Phys.*, 4, 1291–1299, <https://doi.org/10.5194/acp-4-1291-2004>, 2004.
- 425 Faxon, C. B., Bean, J. K., and Ruiz, L. H.: Inland Concentrations of Cl_2 and ClNO_2 in Southeast Texas Suggest Chlorine Chemistry Significantly Contributes to Atmospheric Reactivity, *Atmosphere*, 6, 1487–1506, <https://doi.org/10.3390/atmos6101487>, 2015.
- Finlayson-Pitts, B. J.: Chlorine Atoms as a Potential Tropospheric Oxidant in the Marine Boundary Layer, *Res. Chem. Intermediat.*, 19, 235–249, <https://doi.org/10.1163/156856793X00091>, 1993.
- Finlayson-Pitts, B. J., Ezell, M. J., and Pitts Jr, J. N.: Formation of chemically active chlorine compounds by reactions of atmospheric NaCl particles with gaseous N_2O_5 and ClONO_2 , *Nature*, 337, 241–244, <https://doi.org/https://doi.org/10.1038/337241a0>, 1989.
- 430 Frenzel, A., Scheer, V., Sikorski, R., George, C., Behnke, W., and Zetzsch, C.: Heterogeneous Interconversion Reactions of BrNO_2 , ClNO_2 , BrO_2 , and ClO_2 , *J. Phys. Chem. A*, 102, 1329–1337, <https://doi.org/10.1021/jp973044b>, <http://dx.doi.org/10.1021/jp973044b>, 1998.
- Fu, X., Wang, T., Wang, S., Zhang, L., Cai, S., Xing, J., and Hao, J.: Anthropogenic emissions of hydrogen chloride and fine particulate chloride in China, *Environ. Sci. Technol.*, 52, 1644–1654, <https://doi.org/10.1021/acs.est.7b05030>, 2018.
- 435 Gebel, M. E. and Finlayson-Pitts, B. J.: Uptake and reaction of ClONO_2 on NaCl and synthetic sea salt, *J. Phys. Chem. A*, 105, 5178–5187, <https://doi.org/10.1021/jp0046290>, 2001.
- George, C., Behnke, W., Scheer, V., Zetzsch, C., Magi, L., Ponche, J. L., and Mirabel, P.: Fate of ClNO_2 over aqueous solutions containing iodide, *Geophys. Res. Lett.*, 22, 1505–1508, <https://doi.org/10.1029/95GL01417>, 1995.
- Hanson, D. R. and Ravishankara, A. R.: Reactive Uptake of ClONO_2 onto Sulfuric Acid Due to Reaction with HCl and H_2O , *J. Phy. Chem.*, 440 98, 5728–5735, <https://doi.org/10.1021/j100073a026>, 1994.
- Hanson, D. R., Ravishankara, a. R., and Solomon, S.: Heterogeneous reactions in sulfuric acid aerosols: A framework for model calculations, *J. Geophys. Res.*, 99, 3615, <https://doi.org/10.1029/93JD02932>, 1994.
- Hinrichsen, D.: *Coastal Waters of the World: Trends, Threats, and Strategies*, Island Press, Washington, DC, <http://arp.sagepub.com/content/30/4/473>, 1998.
- 445 Hov, O.: The effect of chlorine on the formation of photochemical oxidants in Southern Telemark, Norway, *Atmos. Environ.*, 19, 471–485, [https://doi.org/10.1016/0004-6981\(85\)90168-4](https://doi.org/10.1016/0004-6981(85)90168-4), 1985.
- Hu, J. H. and Abbatt, J. P. D.: Reaction probabilities for N_2O_5 hydrolysis on sulfuric acid and ammonium sulfate aerosols at room temperature, *J. Phys. Chem. A*, 101, 871–878, <https://doi.org/10.1021/jp9627436>, 1997.
- Huey, L. G.: Measurement of Trace Atmospheric Species by Chemical Ionization Mass Spectrometry: Speciation of Reactive Nitrogen and Future Directions, *Mass Spectrom. Rev.*, 26, 166–184, <https://doi.org/10.1002/mas.20118>, 2007.
- 450 Huey, L. G., Hanson, D. R., and Howard, C. J.: Reactions of SF_6^- and I^- with atmospheric trace gases, *J. Geophys. Res.*, 99, 5001–5008, <https://doi.org/10.1021/j100014a021>, 1995.

- Jenkin, M. E., Young, J. C., and Rickard, A. R.: The MCM v3.3.1 degradation scheme for isoprene, *Atmos. Chem. Phys.*, 15, 11 433–11 459, <https://doi.org/10.5194/acp-15-11433-2015>, 2015.
- 455 Kim, N., Park, M., Yum, S. S., Park, J. S., Song, I. H., Shin, H. J., Ahn, J. Y., Kwak, K. H., Kim, H., Bae, G. N., and Lee, G.: Hygroscopic properties of urban aerosols and their cloud condensation nuclei activities measured in Seoul during the MAPS-Seoul campaign, *Atmos. Environ.*, 153, 217–232, <https://doi.org/10.1016/j.atmosenv.2017.01.034>, <http://dx.doi.org/10.1016/j.atmosenv.2017.01.034>, 2017.
- Kim, N., Park, M., Yum, S. S., Park, J. S., Shin, H. J., and Ahn, J. Y.: Impact of urban aerosol properties on cloud condensation nuclei (CCN) activity during the KORUS-AQ field campaign, *Atmos. Environ.*, 185, 221–236, <https://doi.org/10.1016/j.atmosenv.2018.05.019>,
460 <https://doi.org/10.1016/j.atmosenv.2018.05.019>, 2018.
- Kim, S., Kim, S. Y., Lee, M., Shim, H., Wolfe, G. M., Guenther, A. B., He, A., Hong, Y., and Han, J.: Impact of isoprene and HONO chemistry on ozone and OVOC formation in a semirural South Korean forest, *Atmos. Chem. Phys.*, 15, 4357–4371, <https://doi.org/10.5194/acp-15-4357-2015>, 2015.
- Kim, S., Sanchez, D., Wang, M., Seco, R., Jeong, D., Hughes, S., Barletta, B., Blake, D. R., Jung, J., Kim, D., Lee, G., Lee, M., Ahn, J.,
465 Lee, S. D., Cho, G., Sung, M. Y., Lee, Y. H., Kim, D. B., Kim, Y., Woo, J. H., Jo, D., Park, R., Park, J. H., Hong, Y. D., and Hong, J. H.: OH reactivity in urban and suburban regions in Seoul, South Korea—an East Asian megacity in a rapid transition, *Faraday Discuss.*, 189, 231–251, <https://doi.org/10.1039/c5fd00230c>, 2016.
- Knipping, E. M. and Dabdub, D.: Impact of chlorine emissions from sea-salt aerosol on coastal urban ozone, *Environ. Sci. Technol.*, 37, 275–284, <https://doi.org/10.1021/es025793z>, 2003.
- 470 Lee, B. H., Lopez-Hilfiker, F. D., Schroder, J. C., Campuzano-Jost, P., Jimenez, J. L., McDuffie, E. E., Fibiger, D. L., Veres, P. R., Brown, S. S., Campos, T. L., Weinheimer, A. J., Flocke, F. F., Norris, G., O’Mara, K., Green, J. R., Fiddler, M. N., Bililign, S., Shah, V., Jaegle, L., and Thornton, J. A.: Airborne observations of reactive inorganic chlorine and bromine species in the exhaust of coal-fired power plants, *J. Geophys. Res. Atmos.*, 123, 11 225–11 237, <https://doi.org/10.1029/2018JD029284>, <http://doi.wiley.com/10.1029/2018JD029284>, 2018.
- Li, Q., Zhang, L., Wang, T., Tham, Y. J., Ahmadov, R., Xue, L., Zhang, Q., and Zheng, J.: Impacts of heterogeneous uptake of dinitrogen
475 pentoxide and chlorine activation on ozone and reactive nitrogen partitioning: Improvement and application of the WRF-Chem model in southern China, *Atmos. Chem. Phys.*, 16, 14 875–14 890, <https://doi.org/10.5194/acp-16-14875-2016>, 2016.
- Liao, J., Sihler, H., Huey, L. G., Neuman, J. A., Tanner, D. J., Friess, U., Platt, U., Flocke, F. M., Orlando, J. J., Shepson, P. B., Beine, H. J., Weinheimer, A. J., Sjostedt, S. J., Nowak, J. B., Knapp, D. J., Staebler, R. M., Zheng, W., Sander, R., Hall, S. R., and Ullmann, K.: A comparison of Arctic BrO measurements by chemical ionization mass spectrometry and long path-differential optical absorption
480 spectroscopy, *J. Geophys. Res. Atmos.*, 116, D00R02, <https://doi.org/10.1029/2010JD014788>, 2011.
- Liao, J., Huey, L. G., Tanner, D. J., Flocke, F. M., Orlando, J. J., Neuman, J. A., Nowak, J. B., Weinheimer, A. J., Hall, S. R., Smith, J. N., Fried, A., Staebler, R. M., Wang, Y., Koo, J. H., Cantrell, C. A., Weibring, P., Walega, J., Knapp, D. J., Shepson, P. B., and Stephens, C. R.: Observations of inorganic bromine (HOBr, BrO, and Br₂) speciation at Barrow, Alaska, in spring 2009, *J. Geophys. Res. Atmos.*, 117, D00R16, <https://doi.org/10.1029/2011JD016641>, 2012.
- 485 Liao, J., Huey, L. G., Liu, Z., Tanner, D. J., Cantrell, C. A., Orlando, J. J., Flocke, F. M., Shepson, P. B., Weinheimer, A. J., Hall, S. R., Ullmann, K., Beine, H. J., Wang, Y., Ingall, E. D., Stephens, C. R., Hornbrook, R. S., Apel, E. C., Riemer, D., Fried, A., Mauldin, R. L., Smith, J. N., Staebler, R. M., Neuman, J. A., and Nowak, J. B.: High levels of molecular chlorine in the Arctic atmosphere, *Nat. Geosci.*, 7, 91–94, <https://doi.org/10.1038/ngeo2046>, 2014.

- 490 Liu, X., Qu, H., Huey, L. G., Wang, Y., Sjostedt, S., Zeng, L., Lu, K., Wu, Y., Hu, M., Shao, M., Zhu, T., and Zhang, Y.: High Levels
of Daytime Molecular Chlorine and Nitryl Chloride at a Rural Site on the North China Plain, *Environ. Sci. Technol.*, 51, 9588–9595,
<https://doi.org/10.1021/acs.est.7b03039>, 2017.
- 495 Lowe, D., Archer-Nicholls, S., Morgan, W., Allan, J., Utembe, S., Ouyang, B., Aruffo, E., Le Breton, M., Zaveri, R. A., Di Carlo, P., Percival,
C., Coe, H., Jones, R., and McFiggans, G.: WRF-Chem model predictions of the regional impacts of N₂O₅ heterogeneous processes on
night-time chemistry over north-western Europe, *Atmos. Chem. Phys.*, 15, 1385–1409, <https://doi.org/10.5194/acp-15-1385-2015>, issn =
16807324, 2015.
- Madronich, S. and Flocke, S.: Handbook of Environmental Chemistry, in: Handbook of Environmental Chemistry, chap. The role of solar
radiation in atmospheric chemistry, pp. 1–26, Springer_Verlag, Heidelberg, p. boule edn., 1998.
- Malko, M. W. and Troe, J.: Analysis of the unimolecular reaction N₂O₅ + M ⇌ NO₂ + NO₃ + M, *Int. J. Chem. Kinet.*, 14, 399–416,
<https://doi.org/10.1002/kin.550140407>, 1982.
- 500 McDuffie, E. E., Fibiger, D. L., Dubé, W. P., Lopez Hilfiker, F., Lee, B. H., Jaeglé, L., Guo, H., Weber, R. J., Reeves, J. M., Weinheimer,
A. J., Schroder, J. C., Campuzano-Jost, P., Jimenez, J. L., Dibb, J. E., Veres, P., Ebben, C., Sparks, T. L., Wooldridge, P. J., Cohen, R. C.,
Campos, T., Hall, S. R., Ullmann, K., Roberts, J. M., Thornton, J. A., and Brown, S. S.: ClNO₂ Yields From Aircraft Measurements
During the 2015 WINTER Campaign and Critical Evaluation of the Current Parameterization, *J. Geophys. Res. Atmos.*, 123, 12,994–
13,015, <https://doi.org/10.1029/2018JD029358>, 2018a.
- 505 McDuffie, E. E., Fibiger, D. L., Dubé, W. P., Lopez-Hilfiker, F., Lee, B. H., Thornton, J. A., Shah, V., Jaeglé, L., Guo, H., Weber, R. J.,
Michael Reeves, J., Weinheimer, A. J., Schroder, J. C., Campuzano-Jost, P., Jimenez, J. L., Dibb, J. E., Veres, P., Ebben, C., Sparks, T. L.,
Wooldridge, P. J., Cohen, R. C., Hornbrook, R. S., Apel, E. C., Campos, T., Hall, S. R., Ullmann, K., and Brown, S. S.: Heterogeneous
N₂O₅ uptake during winter: Aircraft measurements during the 2015 WINTER campaign and critical evaluation of current parameteriza-
tions, *J. Geophys. Res. Atmos.*, 123, 4345–4372, <https://doi.org/10.1002/2018JD028336>, 2018b.
- 510 McNeill, V. F., Patterson, J., Wolfe, G. M., and Thornton, J. A.: The effect of varying levels of surfactant on the reactive uptake of N₂O₅ to
aqueous aerosol, *Atmos. Chem. Phys.*, 6, 1635–1644, <https://doi.org/10.5194/acp-6-1635-2006>, 2006.
- Mielke, L. H., Furgeson, A., and Osthoff, H. D.: Observation of ClNO₂ in a mid-continental urban environment, *Environ. Sci. Technol.*, 45,
8889–96, <https://doi.org/10.1021/es201955u>, <http://www.ncbi.nlm.nih.gov/pubmed/21877701>, 2011.
- Morgan, W. T., Ouyang, B., Allan, J. D., Aruffo, E., Di Carlo, P., Kennedy, O. J., Lowe, D., Flynn, M. J., Rosenberg, P. D., Williams, P. I.,
515 Jones, R., McFiggans, G. B., and Coe, H.: Influence of aerosol chemical composition on N₂O₅ uptake: airborne regional measurements
in northwestern Europe, *Atmos. Chem. Phys.*, 15, 973–990, <https://doi.org/10.5194/acp-15-973-2015>, 2015.
- Müller, M., Mikoviny, T., Feil, S., Haidacher, S., Hanel, G., Hartungen, E., Jordan, A., Märk, L., Mutschlechner, P., Schottkowsky, R., Sulzer,
P., Crawford, J. H., and Wisthaler, A.: A compact PTR-ToF-MS instrument for airborne measurements of volatile organic compounds at
high spatiotemporal resolution, *Atmos. Meas. Tech.*, 7, 3763–3772, <https://doi.org/10.5194/amt-7-3763-2014>, 2014.
- 520 Nault, B. A., Campuzano-Jost, P., Day, D. A., Schroder, J. C., Anderson, B., Beyersdorf, A. J., Blake, D. R., Brune, W. H., Choi, Y., Corr,
C. A., de Gouw, J. A., Dibb, J., DiGangi, J. P., Diskin, G. S., Fried, A., Huey, L. G., Kim, M. J., Knote, C. J., Lamb, K. D., Lee, T.,
Park, T., Pusede, S. E., Scheuer, E., Thornhill, K. L., Woo, J. H., and Jimenez, J. L.: Secondary organic aerosol production from local
emissions dominates the organic aerosol budget over Seoul, South Korea, during KORUS-AQ, *Atmos. Chem. Phys.*, 18, 17 769–17 800,
<https://doi.org/10.5194/acp-18-17769-2018>, 2018.
- 525 Neumann, B., Vafeidis, A. T., Zimmermann, J., and Nicholls, R. J.: Future coastal population growth and exposure to sea-level rise and
coastal flooding - A global assessment, *PLoS ONE*, 10, <https://doi.org/10.1371/journal.pone.0118571>, 2015.

- Osthoff, H. D., Roberts, J. M., Ravishankara, A. R., Williams, E. J., Lerner, B. M., Sommariva, R., Bates, T. S., Coffman, D., Quinn, P. K., Dibb, J. E., Stark, H., Burkholder, J. B., Talukdar, R. K., Meagher, J., Fehsenfeld, F. C., and Brown, S. S.: High levels of nitryl chloride in the polluted subtropical marine boundary layer, *Nat. Geosci.*, 1, 324–328, <https://doi.org/10.1038/ngeo177>, <http://www.nature.com/doi/10.1038/ngeo177>, 2008.
- Osthoff, H. D., Odame-ankrah, C. A., Taha, Y. M., Tokarek, T. W., Schiller, C. L., Haga, D., Jones, K., and Vingarzan, R.: Low Levels of Nitryl Chloride in the Lower Fraser Valley of British Columbia, *Atmos. Chem. Phys.*, 18, 6293–6315, <https://doi.org/http://doi.org/10.5194/acp-18-6293-2018>, 2018.
- Park, M. S., Park, S. H., Chae, J. H., Choi, M. H., Song, Y., Kang, M., and Roh, J. W.: High-resolution urban observation network for user-specific meteorological information service in the Seoul Metropolitan Area, South Korea, *Atmos. Meas. Tech.*, 10, 1575–1594, <https://doi.org/10.5194/amt-10-1575-2017>, 2017.
- Phillips, G. J., Thieser, J., Tang, M., Sobanski, N., Schuster, G., Fachinger, J., Drewnick, F., Borrmann, S., Bingemer, H., Lelieveld, J., and Crowley, J. N.: Estimating N_2O_5 uptake coefficients using ambient measurements of NO_3 , N_2O_5 , ClNO_2 and particle-phase nitrate, *Atmos. Chem. Phys.*, 16, 13 231–13 249, <https://doi.org/10.5194/acp-16-13231-2016>, 2016.
- Reff, A., Bhave, P. V., Simon, H., Pace, T. G., Pouliot, G. A., Mobley, J. D., and Houyoux, M.: Emissions Inventory of $\text{PM}_{2.5}$ Trace Elements across the United States, *Environ. Sci. Technol.*, 43, 5790–5796, <https://doi.org/10.1021/es802930x>, <http://pubs.acs.org/doi/abs/10.1021/es802930x>, 2009.
- Riedel, T. P., Bertram, T. H., Crisp, T. A., Williams, E. J., Lerner, B. M., Vlasenko, A., Li, S. M., Gilman, J., De Gouw, J., Bon, D. M., Wagner, N. L., Brown, S. S., and Thornton, J. A.: Nitryl chloride and molecular chlorine in the coastal marine boundary layer, *Environ. Sci. Technol.*, 46, 10 463–10 470, <https://doi.org/10.1021/es204632r>, 2012.
- Riedel, T. P., Wagner, N. L., Dubé, W. P., Middlebrook, A. M., Young, C. J., Öztürk, F., Bahreini, R., Vandenboer, T. C., Wolfe, D. E., Williams, E. J., Roberts, J. M., Brown, S. S., and Thornton, J. A.: Chlorine activation within urban or power plant plumes: Vertically resolved ClNO_2 and Cl_2 measurements from a tall tower in a polluted continental setting, *J. Geophys. Res. Atmos.*, 118, 8702–8715, <https://doi.org/10.1002/jgrd.50637>, 2013.
- Riedel, T. P., Wolfe, G. M., Danas, K. T., Gilman, J. B., Kuster, W. C., Bon, D. M., Vlasenko, A., Li, S. M., Williams, E. J., Lerner, B. M., Veres, P. R., Roberts, J. M., Holloway, J. S., Lefer, B., Brown, S. S., and Thornton, J. A.: An mcm modeling study of nitryl chloride (ClNO_2) impacts on oxidation, ozone production and nitrogen oxide partitioning in polluted continental outflow, *Atmos. Chem. Phys.*, 14, 3789–3800, <https://doi.org/10.5194/acp-14-3789-2014>, 2014.
- Roberts, J. M., Osthoff, H. D., Brown, S. S., and Ravishankara, A. R.: N_2O_5 oxidizes chloride to Cl_2 in acidic atmospheric aerosol, *Science*, 321, 1059, <https://doi.org/10.1126/science.1158777>, 2008.
- Rossi, M. J.: Heterogeneous Reactions on Salts, *Chem. Rev.*, 103, 4823–4882, <https://doi.org/10.1021/cr020507n>, 2003.
- Ryder, O. S., Campbell, N. R., Shalowski, M., Al-Mashat, H., Nathanson, G. M., and Bertram, T. H.: Role of Organics in Regulating ClNO_2 Production at the Air–Sea Interface, *J. Phys. Chem. A*, 119, 8519–8526, <https://doi.org/10.1021/jp5129673>, 2015.
- Sarwar, G., Simon, H., Bhave, P., and Yarwood, G.: Examining the impact of heterogeneous nitryl chloride production on air quality across the United States, *Atmos. Chem. Phys.*, 12, 6455–6473, <https://doi.org/10.5194/acp-12-6455-2012>, 2012.
- Sarwar, G., Simon, H., Xing, J., and Mathur, R.: Importance of tropospheric ClNO_2 chemistry across the Northern Hemisphere, *Geophys. Res. Lett.*, 41, 4050–4058, <https://doi.org/10.1002/2014GL059962>, 2014.
- Schweitzer, F., Mirabel, P., and George, C.: Multiphase Chemistry of N_2O_5 , ClNO_2 , and BrNO_2 , *J. Phys. Chem. A*, 102, 3942 – 3952, <https://doi.org/10.1021/jp980748s>, 1998.

- 565 Sherwen, T., Schmidt, J. A., Evans, M. J., Carpenter, L. J., Großmann, K., Eastham, S. D., Jacob, D. J., Dix, B., Koenig, T. K., Sinreich, R., Ortega, I., Volkamer, R., Saiz-Lopez, A., Prados-Roman, C., Mahajan, A. S., and Ordóñez, C.: Global impacts of tropospheric halogens (Cl, Br, I) on oxidants and composition in GEOS-Chem, *Atmos. Chem. Phys.*, 16, 12 239–12 271, <https://doi.org/10.5194/acp-16-12239-2016>, 2016.
- Sherwen, T., Evans, M. J., Sommariva, R., Hollis, L. D., Ball, S. M., Monks, P. S., Reed, C., Carpenter, L. J., Lee, J. D., Forster, 570 G., Bandy, B., Reeves, C. E., and Bloss, W. J.: Effects of halogens on European air-quality, *Faraday Discuss.*, 200, 75–100, <https://doi.org/10.1039/c7fd00026j>, 2017.
- Shetter, R. E. and Müller, M.: Photolysis frequency measurements using actinic flux spectroradiometry during the PEM-Tropics mission: Instrumentation description and some results, *J. Geophys. Res. Atmos.*, 104, 5647–5661, <https://doi.org/10.1029/98JD01381>, 1999.
- Simon, H., Kimura, Y., McGaughey, G., Allen, D. T., Brown, S. S., Osthoff, H. D., Roberts, J. M., Byun, D., and Lee, D.: Modeling 575 the impact of ClNO₂ on ozone formation in the Houston area, *J. Geophys. Res.*, 114, D00F03, <https://doi.org/10.1029/2008JD010732>, <http://doi.wiley.com/10.1029/2008JD010732>, 2009.
- Slusher, D. L., Huey, L. G., Tanner, D. J., Flocke, F. M., and Roberts, J. M.: A thermal dissociation - Chemical ionization mass spectrometry (TD-CIMS) technique for the simultaneous measurement of peroxyacyl nitrates and dinitrogen pentoxide, *J. Geophys. Res.*, 109, D19 315, <https://doi.org/10.1029/2004JD004670>, 2004.
- 580 Stull, R. B.: An introduction to boundary layer meteorology, vol. 13, Dordrecht; Boston : Kluwer Academic Publishers, 1988.
- Sullivan, J. T., McGee, T. J., Sumnicht, G. K., Twigg, L. W., and Hoff, R. M.: A mobile differential absorption lidar to measure sub-hourly fluctuation of tropospheric ozone profiles in the Baltimore-Washington, D.C. region, *Atmos. Meas. Tech.*, 7, 3529–3548, <https://doi.org/10.5194/amt-7-3529-2014>, 2014.
- Sullivan, J. T., McGee, T. J., Stauffer, R. M., Thompson, A. M., Weinheimer, A., Knote, C., Janz, S., Wisthaler, A., Long, R., Szykman, J., 585 Park, J., Lee, Y., Kim, S., Jeong, D., Sanchez, D., Twigg, L., Sumnicht, G., Knepp, T., and Schroeder, J. R.: Taehwa Research Forest: A receptor site for severe pollution events in Korea during 2016, *Atmos. Chem. Phys. Discuss.*, pp. 1–30, <https://doi.org/10.5194/acp-2018-1328>, 2019.
- Tanaka, P. L., Oldfield, S., Neece, J. D., Mullins, C. B., and Allen, D. T.: Anthropogenic sources of chlorine and ozone formation in urban atmospheres, *Environ. Sci. Technol.*, 34, 4470–4473, <https://doi.org/10.1021/es991380v>, 2000.
- 590 Tanaka, P. L., Riemer, D. D., Chang, S., Yarwood, G., McDonald-Buller, E. C., Apel, E. C., Orlando, J. J., Silva, P. J., Jimenez, J. L., Canagaratna, M. R., Neece, J. D., Mullins, C. B., and Allen, D. T.: Direct evidence for chlorine-enhanced urban ozone formation in Houston, Texas, *Atmos. Environ.*, 37, 1393–1400, [https://doi.org/10.1016/S1352-2310\(02\)01007-5](https://doi.org/10.1016/S1352-2310(02)01007-5), 2003.
- Tang, W., Arellano, A. F., DiGangi, J. P., Choi, Y., Diskin, G. S., Agustí-Panareda, A., Parrington, M., Massart, S., Gaubert, B., Lee, Y., Kim, D., Jung, J., Hong, J., Hong, J.-W., Kanaya, Y., Lee, M., Stauffer, R. M., Thompson, A. M., Flynn, J. H., and Woo, J.-H.: Evaluating 595 high-resolution forecasts of atmospheric CO and CO₂ from a global prediction system during KORUS-AQ field campaign, *Atmos. Chem. Phys.*, 18, 11 007–11 030, <https://doi.org/10.5194/acp-18-11007-2018>, 2018.
- Tang, W., Emmons, L. K., Arellano Jr., A. F., Gaubert, B., Knote, C., Tilmes, S., Buchholz, R. R., Pfister, G. G., Diskin, G. S., Blake, D. R., Blake, N. J., Meinardi, S., DiGangi, J. P., Choi, Y., Woo, J.-H., He, C., Schroeder, J. R., Suh, I., Lee, H.-J., Jo, H.-Y., Kanaya, Y., Jung, J., Lee, Y., and Kim, D.: Source contributions to carbon monoxide concentrations during KORUS-AQ based on CAM-chem model 600 applications, *J. Geophys. Res. Atmos.*, pp. 1–27, <https://doi.org/10.1029/2018jd029151>, 2019.
- Thaler, R. D., Mielke, L. H., and Osthoff, H. D.: Quantification of nitryl chloride at part per trillion mixing ratios by thermal dissociation cavity ring-down spectroscopy, *Anal. Chem.*, 83, 2761–2766, <https://doi.org/10.1021/ac200055z>, 2011.

- 605 Tham, Y. J., Wang, Z., Li, Q., Yun, H., Wang, W., Wang, X., Xue, L., Lu, K., Ma, N., Bohn, B., Li, X., Kecorius, S., Größ, J., Shao, M., Wiedensohler, A., Zhang, Y., and Wang, T.: Significant concentrations of nitryl chloride sustained in the morning: Investigations of the causes and impacts on ozone production in a polluted region of northern China, *Atmos. Chem. Phys.*, 16, 14 959–14 977, <https://doi.org/10.5194/acp-16-14959-2016>, 2016.
- Tham, Y. J., Wang, Z., Li, Q., Wang, W., Wang, X., Lu, K., Ma, N., Yan, C., Kecorius, S., Wiedensohler, A., Zhang, Y., and Wang, T.: Heterogeneous N_2O_5 uptake coefficient and production yield of ClNO_2 in polluted northern China: roles of aerosol water content and chemical composition, *Atmos. Chem. Phys.*, 18, 13 155–13 171, <https://doi.org/10.5194/acp-18-13155-2018>, 2018.
- 610 Thornton, J. A., Braban, C. F., and Abbatt, J. P. D.: N_2O_5 hydrolysis on sub-micron organic aerosols: the effect of relative humidity, particle phase, and particle size, *Phys. Chem. Chem. Phys.*, 5, 4593–4603, <https://doi.org/10.1039/b307498f>, <http://xlink.rsc.org/?DOI=b307498f>, 2003.
- Thornton, J. A., Kercher, J. P., Riedel, T. P., Wagner, N. L., Cozic, J., Holloway, J. S., Dubé, W. P., Wolfe, G. M., Quinn, P. K., Middlebrook, A. M., Alexander, B., and Brown, S. S.: A large atomic chlorine source inferred from mid-continental reactive nitrogen chemistry, *Nature*, 615 464, 271–274, <https://doi.org/10.1038/nature08905>, <http://www.nature.com/doifinder/10.1038/nature08905>, 2010.
- Vogt, R., Crutzen, P., and Sander, R.: A mechanism for halogen release from sea-salt, *Nature*, 383, 327–331, <https://doi.org/10.1038/383327a0>, 1996.
- Wagner, N. L., Riedel, T. P., Young, C. J., Bahreini, R., Brock, C. A., Dubé, W. P., Kim, S., Middlebrook, A. M., Öztürk, F., Roberts, J. M., Russo, R., Sive, B., Swarthout, R., Thornton, J. A., VandenBoer, T. C., Zhou, Y., and Brown, S. S.: N_2O_5 uptake coefficients and nocturnal NO_2 removal rates determined from ambient wintertime measurements, *J. Geophys. Res. Atmos.*, 118, 9331–9350, <https://doi.org/10.1002/jgrd.50653>, 2013.
- Wang, T., Tham, Y. J., Xue, L., Li, Q., Zha, Q., Wang, Z., Poon, S. C. N., Dubé, W. P., Blake, D. R., Louie, P. K. K., Luk, C. W. Y., Tsui, W., and Brown, S. S.: Observations of nitryl chloride and modeling its source and effect on ozone in the planetary boundary layer of southern China, *J. Geophys. Res. Atmos.*, 121, 2476–2489, <https://doi.org/10.1002/2015JD024556>. Received, 2016.
- 625 Wang, X., Wang, T., Yan, C., Tham, Y. J., Xue, L., Xu, Z., and Zha, Q.: Large daytime signals of N_2O_5 and NO_3 inferred at 62 amu in a TD-CIMS: Chemical interference or a real atmospheric phenomenon, *Atmos. Meas. Tech.*, 7, 1–12, <https://doi.org/10.5194/amt-7-1-2014>, 2014.
- Wang, X., Wang, H., Xue, L., Wang, T., Wang, L., Gu, R., Wang, W., Tham, Y. J., Wang, Z., Yang, L., Chen, J., and Wang, W.: Observations of N_2O_5 and ClNO_2 at a polluted urban surface site in North China: High N_2O_5 uptake coefficients and low ClNO_2 product yields, *Atmos. Environ.*, 156, 125–134, <https://doi.org/10.1016/j.atmosenv.2017.02.035>, 2017a.
- 630 Wang, X., Jacob, D. J., Eastham, S. D., Sulprizio, M. P., Zhu, L., Chen, Q., Alexander, B., Sherwen, T., Evans, M. J., Lee, B. H., Haskins, J. D., Lopez-Hilfiker, F. D., Thornton, J. A., Huey, G. L., and Liao, H.: The role of chlorine in global tropospheric chemistry, *Atmos. Chem. Phys.*, 19, 3981–4003, <https://doi.org/10.5194/acp-2018-1088>, 2019.
- Wang, Z., Wang, W., Tham, Y. J., Li, Q., Wang, H., Wen, L., Wang, X., and Wang, T.: Fast heterogeneous N_2O_5 uptake and ClNO_2 production in power plant and industrial plumes observed in the nocturnal residual layer over the North China Plain, *Atmos. Chem. Phys.*, 17, 175194, 12 361–12 378, <https://doi.org/10.5194/acp-17-12361-2017>, 2017b.
- 635 Wayne, R. P., Barnes, I., Biggs, P., Burrows, J. P., Canosa-Mas, C. E., Hjorth, J., Le Bras, G., Moortgat, G. K., Perner, D., Poulet, G., Restelli, G., and Sidebottom, H.: The nitrate radical: Physics, chemistry, and the atmosphere, *Atmos. Environ.*, 25, 1–203, [https://doi.org/10.1016/0960-1686\(91\)90192-A](https://doi.org/10.1016/0960-1686(91)90192-A), 1991.

- 640 Weinheimer, A. J., Walega, J. G., Ridley, B. A., Gary, B. L., Blake, D. R., Blake, N. J., Rowland, F. S., Sachse, G. W., Anderson, B. E., and Collins, J. E.: Meridional distributions of NO_x , NO_y , and other species in the lower stratosphere and upper troposphere during AASE II, *Geophys. Res. Lett.*, 21, 2583–2586, <https://doi.org/10.1029/94GL01897>, 1994.
- Williams, E., Baumann, K., Roberts, J. M., Bertman, S. B., Norton, R. B., Fehsenfeld, C., Springston, S. R., Nunnermacker, L. J., Newman, L., Olszyna, K., Meagher, J., Hartsell, B., Edgerton, E., Pearson, J. R., and Rodgers, M. O.: Intercomparison of ground-based NO_y measurement techniques, *J. Geophys. Res.*, 103, 22 261–22 280, 1998.
- 645 Wolfe, G. M., Marvin, M. R., Roberts, S. J., Travis, K. R., and Liao, J.: The framework for 0-D atmospheric modeling (F0AM) v3.1, *Geosci. Model Dev.*, 9, 3309–3319, <https://doi.org/10.5194/gmd-9-3309-2016>, 2016.
- Woodcock, A. H.: Salt Nuclei in Marine Air As a Function of Altitude and Wind Force, *J. Meteorol.*, 10, 362–371, [https://doi.org/10.1175/1520-0469\(1953\)010<0366:SNIMAA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1953)010<0366:SNIMAA>2.0.CO;2), <http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%281953%29010%3C0366%3ASNIMAA%3E2.0.CO%3B2>, 1953.
- 650 Young, C. J., Washenfelder, R. A., Roberts, J. M., Mielke, L. H., Osthoff, H. D., Tsai, C., Pikelnaya, O., Stutz, J., Veres, P. R., Cochran, A. K., Vandenboer, T. C., Flynn, J., Grossberg, N., Haman, C. L., Lefer, B., Stark, H., Graus, M., De Gouw, J., Gilman, J. B., Kuster, W. C., and Brown, S. S.: Vertically resolved measurements of nighttime radical reservoirs in Los Angeles and their contribution to the urban radical budget, *Environ. Sci. Technol.*, 46, 10 965–10 973, <https://doi.org/10.1021/es302206a>, 2012.
- 655 Young, C. J., Washenfelder, R. A., Edwards, P. M., Parrish, D. D., Gilman, J. B., Kuster, W. C., Mielke, L. H., Osthoff, H. D., Tsai, C., Pikelnaya, O., Stutz, J., Veres, P. R., Roberts, J. M., Griffith, S., Dusanter, S., Stevens, P. S., Flynn, J., Grossberg, N., Lefer, B., Holloway, J. S., Peischl, J., Ryerson, T. B., Atlas, E. L., Blake, D. R., and Brown, S. S.: Chlorine as a primary radical: Evaluation of methods to understand its role in initiation of oxidative cycles, *Atmos. Chem. Phys.*, 14, 3427–3440, <https://doi.org/10.5194/acp-14-3427-2014>, 2014.
- Yun, H., Wang, W., Wang, T., Xia, M., Yu, C., Wang, Z., Poon, S. C. N., Yue, D., and Zhou, Y.: Nitrate formation from heterogeneous uptake of dinitrogen pentoxide during a severe winter haze in southern China, *Atmos. Chem. Phys.*, 18, 17 515–17 527, <https://doi.org/10.5194/acp-18-17515-2018>, 2018.
- 660 Zhang, L., Li, Q., Wang, T., Ahmadov, R., Zhang, Q., Li, M., and Lv, M.: Combined Impacts of Nitrous Acid and Nitryl Chloride on Lower Tropospheric Ozone: New Module Development in WRF-Chem and Application to China, *Atmos. Chem. Phys.*, 17, 1–31, <https://doi.org/https://doi.org/10.5194/acp-17-9733-2017>, 2017.

Table 1. Summary of the measurements carried out during the KORUS-AQ 2016 field campaign, used in this study.

| Compound | Method | | |
|------------------------------------|-------------------------|-----------------------------|-------------------------|
| | TRF | OP | DC-8 |
| NO | ^a CL | ^b FRM | ^c CL |
| NO ₂ | ^d CRDS | ^e CAPS | ^c CL |
| O ₃ | ^f DIAL | ^g SL-UV | ^c CL |
| SO ₂ | ^h PF | ⁱ UV florescence | x |
| CO | ^j IR | ^k NDIR | x |
| CINO ₂ ,Cl ₂ | ^l CIMS | ^l CIMS | ^l CIMS |
| VOCs | ^m PTR-ToF-MS | ^o QCL | ^p PTR-ToF-MS |
| | ⁿ TILDAS | | ^q WAS |
| chloride (< 1 μm) | x | | |
| nitrate (< 1 μm) | x | ^r ToF-AMS | ^s HR-ToF-AMS |
| sulfate (< 1 μm) | x | | |
| surface area (< 200 nm) | x | x | ^t SMPS |
| surface area (200 nm - 5 μm) | x | x | ^u LAS |
| Jvalues | x | x | ^v CAFS |

^aChemiluminescence with a molybdenum converter (Thermo Scientific 42i - TL), ^bChemiluminescence detector (Federal Reference Method, Teledyne T200U), ^cNCAR 4-channel chemiluminescence (Weinheimer et al., 1994), ^dCavity Ring Down Spectroscopy (Los Gatos Research NO₂ analyzer), ^eCavity Attenuated Phase Shift spectroscopy (Teledyne T500U CAPS analyzer), ^fNASA TROPospheric OZone Differential Absorption Lidar (Sullivan et al., 2014), ^gUV photometric method (2B 211), ^hPulsed fluorescence method (Thermo Scientific 43i-HL), ⁱUV florescence method (KENTEK), ^jInfrared CO analyzer (Thermo Scientific 48i-HL), ^kNon-Dispersive Infrared CO analyzer (KENTEK), ^lChemical Ionization Mass Spectrometer (Slusher et al., 2004), ^mProton-Transfer-Reaction Time-of-Flight Mass Spectrometer (IONICON), ⁿTunable Infrared Laser Direct Absorption Spectroscopy (Aerodyne), ^oQuantum Cascade Laser spectrometer (Aerodyne), ^pUniversity of Oslo/Innsbruck Proton-Transfer-Reaction Time-Of-Flight Mass Spectrometer (Müller et al., 2014), ^qWhole Air Sampler (Colman et al., 2001), ^rAerosol Mass Spectrometer (Aerodyne), ^sUniversity of Colorado, Boulder,Aerosol Mass Spectrometer (Nault et al., 2018), ^tNASA, Scanning Mobility Particle Sizer, ^uNASA, Laser Aerosol Spectrometer, ^vNCAR, Charged-coupled device Actinic Flux Spectroradiometer (Shetter and Müller, 1999)





Figure 1. (a) Location of two ground sites (Taehwa Research Forest and Olympic Park) where the chemical ionization mass spectrometer (CIMS) was installed during the KORUS-AQ 2016 field campaign (b) Airborne measurements of ClNO_2 and DC-8 flight tracks during the whole campaign. The ClNO_2 data points are 60 sec averaged and color coded by time of day of the measurement. The marker size is proportional to the mixing ratio of ClNO_2 .




Figure 2. Temporal variation of trace gases measured at the (a) Olympic Park site (OP) and (b) Taehwa Research Forest (TRF). For both OP and TRF, the frequency of the averaged data is 10 min for NO_x and 1 min for O₃.




Figure 3. CINO₂ and Cl₂ observation results at (a) OP and (b) TRF averaged over 5 min. FLEXPART back trajectory analysis were made for source contribution of CO-like substance originating from the ocean, assuming inert CO. Aerosol chloride mass concentration (ambient $\mu\text{g m}^{-3}$) was measured at the ground for the OP site and on the NASA DC-8 for TRF. For the airborne chloride, measurements below 1 km over the TRF site is shown. Red shades are the time frames with limited CINO₂ production. The time frames for each meteorological condition that dominated during the observation period are classified in black arrows at the bottom of the Figures 3a and 3b




Figure 4. Scatter plot of daytime (11:00 - 18:00 local time) CINO₂ and Cl₂ at (a) OP and (b)TRF, color coded with measured O₃. 5 min averaged data for the whole campaign were used for both sites. Data points of Cl₂ below detection limit (2.9 ppt, 2 σ , over 30 min) are shown for the purpose of comparison to observed CINO₂ levels.




Figure 5. Correlation between box model simulated daytime (11:00 - 18:00 local time) Cl_2 and measured ClNO_2 at (a) OP and (b)TRF, color coded with measured O_3 .




Figure 6. Diurnal variation of ClNO_2 and other trace gases measured during the campaign and averaged over selected days at (a) OP (7 days) and (b) TRF (9 days). Night time is shown as grey shades.




Figure 7. (a) Regional and (b) vertical distribution of airborne ClONO_2 measured over the Seoul Metropolitan Region (SMA) in the morning (8:00 - 8:30 local time).




Figure 8. Steady state CINO₂, simulated from a box model constrained with airborne measurements (blue) and ground site data from TRF (red), when there was a morning CINO₂ peak. Averaged values of the model runs are shown here with standard deviations.




Figure 9. FLEXPART back trajectories from Taehwa Research Forest. Trajectories were initialized at 9 am local time and went 24 h backwards. Only the center trajectories with the highest percentage of airmasses are presented. Trajectories for days with high levels of CINO₂ (> 500 pptv) at night are in red and the remaining days are shown in sky blue.




Figure 10. Box model simulations of chlorine radicals and O₃ at (a) OP and (b) TRF, constrained with ClNO₂ and other trace gases observed during the field campaign. Percent difference of net O₃ production rate (NetP(O₃), blue line) was calculated from the difference of the NetP(O₃) between simulations with and without ClNO₂ constrained in the model (i.e., $100 \cdot (w_{\text{ClNO}_2} - w_{\text{noClNO}_2}) / w_{\text{ClNO}_2}$).