

Dear Authors

**Dear Editor,**

5 ***Thank you for those comments we used to improve the manuscript,***

10 Congratulations on a much-improved draft. In general, my sense is that you have adequately addressed all the reviewer concerns. However, I think a "little bit" got through the gaps, so please take another read through the manuscript to make sure that the whole is harmonized for usage, corrections, etc. For one example, the correction of "Mongu Inn" was not extended throughout the manuscript. To me, this is not a significant issue, but since the Reviewer pointed it out, please either correct throughout (also the usage of "Mongu Station"), or add a sentence at the first use indicating that a variety of terms are used to reference this site.

15 ***We have read the article to harmonize it. In parallel, the Mongu AERONET site is now mentioned as « Mongu Inn » throughout the manuscript,***

20 I also wanted to point out a more significant issue: in addressing Reviewer #1 on the difference between DRE and DRF, it appears that you have mixed them up by writing "(DRF; with respect to noaerosols)". This is of course a point that could be critical to the reader's understanding of everything going on, so please make certain that this is corrected throughout the manuscript. I refer you to Heald et al.'s paper on the topic: <https://www.atmos-chem-phys.net/14/5513/2014/acp-14-5513-2014.html>

25 ***This is right and we have now replaced the term DRF by DRE over the whole manuscript following the Heald et al. (2014) convention as we have effectively investigated the « instantaneous radiative impact » of smoke aerosols in this study. In order to be precise, we have also included in the introduction the Heald et al. (2014) reference.***

30

35

40

# **Simulation of the transport, vertical distribution, optical properties and radiative impact of smoke aerosols with the ALADIN regional climate model during the ORACLES-2016 and LASIC experiments.**

45

Marc Mallet<sup>1</sup>, Pierre Nabat<sup>1</sup>, Paquita Zuidema<sup>2</sup>, Jens Redemann<sup>3</sup>, Andrew Mark Sayer<sup>4,5</sup>, Martin Stengel<sup>6</sup>, Sebastian Schmidt<sup>7</sup>, Sabrina Cochrane<sup>7</sup>, Sharon Burton<sup>8</sup>, Richard Ferrare<sup>8</sup>, Kerry Meyer<sup>5</sup>, Pablo Saide<sup>9</sup>, Hiren Jethva<sup>4,5</sup>, Omar Torres<sup>5</sup>, Robert Wood<sup>10</sup>, David Saint Martin<sup>1</sup>, Romain Roehrig<sup>1</sup>, Christina Hsu<sup>5</sup> and Paola Formenti<sup>11</sup>

50

## Affiliation

<sup>1</sup> Centre National de Recherches Météorologiques, UMR3589, Météo-France-CNRS, Toulouse, France

<sup>2</sup> Rosenstiel School of Marine and Atmospheric Sciences, University of Miami, Miami, FL, USA

<sup>3</sup> University of Oklahoma, USA

55 <sup>4</sup> Universities Space Research Association, Columbia, MD, USA

<sup>5</sup> NASA Goddard Space Flight Center, Greenbelt, MD, USA

<sup>6</sup> Deutscher Wetterdienst (DWD), Offenbach, Germany

<sup>7</sup> University of Colorado, Boulder, USA

<sup>8</sup> NASA Langley Research Center, Hampton, Virginia, USA

60 <sup>9</sup> University of California, Los Angeles (UCLA), USA

<sup>10</sup> University of Washington, Seattle, USA

<sup>11</sup> Laboratoire Interuniversitaire des Systèmes Atmosphériques, UMR CNRS 7583, Université Paris Est Créteil et Université Paris Diderot, Institut Pierre Simon Laplace, France

65 *Correspondence to:* M. Mallet (marc.mallet@meteo.fr)

70

## 75 Abstract

Estimates of the direct radiative **effect forcing** (DRE) from absorbing smoke aerosols over the Southeast Atlantic Ocean (SAO) require simulation of the microphysical and optical properties of stratocumulus clouds as well as of the altitude and shortwave (SW) optical properties of biomass burning aerosols (BBA). In this study, we take advantage of the large number of observations  
80 acquired during the ORACLES-2016 and LASIC projects during September 2016 and compare them with datasets from the ALADIN-Climate regional model. The model provides a good representation of the liquid water path but the low cloud fraction is underestimated compared to satellite data. The modeled total column smoke aerosol optical depth (AOD) and Above Cloud AOD are consistent ( $\sim 0.7$  over continental sources and  $\sim 0.3$  over SAO at 550 nm) with MERRA2,  
85 OMI or MODIS data. The simulations indicate smoke transport over SAO occurs mainly between 2 and 4 km, consistent with surface and aircraft lidar observations. The BBA single scattering albedo is slightly overestimated compared to AERONET, and more significantly when compared to Ascension Island surface observations. The difference could be due to the absence of internal mixing treatment in the ALADIN-Climate model. The SSA overestimate leads to underestimate the  
90 simulated SW radiative heating compared to ORACLES data. ALADIN-Climate simulates a positive (monthly mean) SW **DRE** of about  $+6 \text{ W}\cdot\text{m}^{-2}$  over SAO ( $20^{\circ}\text{S}$ – $10^{\circ}\text{N}$  and  $10^{\circ}\text{W}$ – $20^{\circ}\text{E}$ ) at the top of the atmosphere and in all-sky conditions. Over the continent, the presence of BBA is shown to significantly decrease the net surface SW flux, through direct and semi-direct effects, which is compensated by a decrease (monthly mean) in sensible heat fluxes ( $-25 \text{ W}\cdot\text{m}^{-2}$ ) and surface  
95 land temperature ( $-1.5 \text{ }^{\circ}\text{C}$ ) over Angola, Zambia and Congo notably. The surface cooling and the lower tropospheric heating decreases the continental planetary boundary layer height by about  $\sim 200$  m.

## 1. Introduction

100 Southern Africa is one of the main sources of biomass burning aerosols (BBA) at the global scale. When the intense smoke plumes are transported over the Southeast Atlantic Ocean (SAO), they are able to produce a significant positive (warming) direct radiative **effect forcing** (DRE; **which represents the instantaneous radiative impact of aerosols on the energy balance, Heald et al., 2014**) **with respect to no-aerosols**) at the top of the atmosphere (TOA) in the SW spectral range and in all-  
105 sky conditions (De Graaf et al., 2012, 2014, Feng and Christopher, 2015, Zuidema et al., 2016). Over this specific region, the sign of the **DRE** is found to be opposite to the cooling effect generally exerted by scattering aerosols at TOA. Based on the combination of satellite observations from A-Train data sets (MODIS, CERES and OMI), Feng and Christopher (2015) indicate a regional-

110 averaged instantaneous (i.e., time of observations) **DRE** of about  $+37 \text{ W.m}^{-2}$  (regional mean;  $20^{\circ}\text{S}$ – $10^{\circ}\text{N}$  and  $10^{\circ}\text{W}$ – $20^{\circ}\text{E}$ ) for August 2006, with the highest magnitude of the forcing reaching  $+138 \text{ W.m}^{-2}$  at TOA. Significant positive values are also underlined by De Graaf et al. (2012, 2014), who estimate an averaged **DRE** (August 2006) of about  $+23 \text{ W.m}^{-2}$  near the Southern African coast. In parallel, Meyer et al. (2013) report an instantaneous (near local noon for Aqua overpass) regional mean above-cloud radiative forcing efficiency from  $50 \text{ W.m}^{-2}\text{AOD}^{-1}$  to  $65 \text{ W.m}^{-2}\text{AOD}^{-1}$  by using  
115 their bias-adjusted MODIS cloud retrievals. By using SCIAMACHY observations and radiative transfer model calculations, De Graaf et al. (2014) further estimate a SW **DRE** of about  $\sim +30/35 \text{ W.m}^{-2}$  over the same domain ( $4$ – $18^{\circ}\text{S}$ ,  $5^{\circ}\text{W}$ – $14^{\circ}\text{E}$ ) in August and September (2006-2009).

This positive sign (warming) of the **DRE** is mainly due to the presence of highly reflective stratocumulus clouds (Sc) over the SAO. Although such a positive **DRE** is occasionally observed  
120 over other regions, such as the northeast Pacific during extreme summertime biomass burning events in continental North America (Mallet et al., 2017), the SAO clearly represents the main region at the global scale where such positive forcings can be observed every years at a seasonal-time scale. Indeed, this significant radiative forcing is due to the persistent biomass burning emissions over Central Africa during the July-August-September-October (JASO) period. Smoke  
125 emissions over the central Africa are also related to a significant inter-annual variability, associated with an important increase over the 1979-2015 period (Hodnebrog et al., 2016).

All studies clearly underline the importance of both the aerosol radiative properties of smoke plumes (e.g., Aerosol Optical Depth, AOD, single scattering albedo, SSA), their vertical structures (notably, the localization of smoke vs. Sc; Johnson et al., 2004) as well as the underlying cloud  
130 properties (e.g., cloud optical depth (COD), liquid water path (LWP)) on the produced positive SW **DRE** at TOA. As an example, Feng and Christopher (2015) report a critical COD of  $\sim 12$ - $20$  capable of changing the sign of the **DRE** from negative to positive (at TOA) for BBA characterized by SSA  $\sim 0.91$  and AOD  $\sim 1.0$  (at 550 nm). In the case of more absorbing smoke (SSA  $\sim 0.85$ ), the ranges for critical COD are strongly reduced and reach  $\sim 2$ - $4$ . Chand et al. (2009) also underline the importance  
135 of cloud coverage on the **DRE** exerted at TOA by smoke over SAO. In addition, Sakaeda et al. (2011) provided model estimates of regional radiative forcing from direct and semi-direct effects, with important implications on cloud properties (cloud fraction notably). These complex processes, involving both microphysical and optical properties of BBA and Sc explain, at least partially, the large difficulty of recent global climate models (GCM) in reproducing the **DRE** of smoke over this  
140 specific region (Stier et al., 2013).

In that context, it appears crucial to evaluate carefully and constrain both smoke aerosols and Sc properties in GCM or in their regional configurations (regional climate models, RCM) before

running them over a long-time period, for radiative budget and climatic considerations. The main objectives of this study is to investigate the transport of BBA over SOA, the vertical layering as well as optical properties using the ALADIN-Climate model. In addition, the induced SW **DRE** at TOA and the possible impact of BBA on the regional (continental) climate are also analysed. This work has been conducted in the context of several international field campaigns over the SAO region, including the ObseRvations of Aerosols above Clouds and their intEractionS (ORACLES) (Zuidema et al., 2016), the Layered Atlantic Smoke Interactions with Clouds (LASIC, Zuidema et al., 2018), the AErosol RAdiation and CLOuds in Southern Africa (AEROCLO-sA) and the Cloud-Aerosol-Radiation Interactions and Forcing : Year 2017 (CLARIFY) projects. More specifically, this study takes advantage of the large number of in situ observations acquired from aircraft and surface measurements during September 2016 for the ORACLES-1 and LASIC projects. This unique dataset is combined to satellite aerosol and cloud retrievals (MODIS, OMI and SEVIRI) and re-analysis (MERRA2, CAMS and MACC) data.

We focus our analyses on specific properties which are important to study the radiative effect of BBA over SAO. For Sc clouds, these are low cloud fraction (LCF), liquid water path (LWP) and COD. For BBA, special attention is paid to AOD, Above Cloud AOD (ACAOD), extinction vertical profiles, SSA, and SW radiative heating induced by ~~due to~~ smoke. The **DRE** at the surface and TOA are estimated and analysed in addition to climatic implications, especially those exerted by BBA over the Central Africa continent. The regional modeling model used in this work is the ALADIN-Climate model (Nabat et al., 2015a, 2015b; Daniel et al., 2018), which has been modified in a recent configuration to better represent smoke aerosols and notably their SW optical properties.

This article is organized as follows. First, details on these recent developments are provided in section 2 as well as the design of the ALADIN-Climate simulations. Section 3 reports the complete dataset (satellites, reanalysis, in situ surface and aircraft observations). The analyses of the comparisons between simulated and observed Sc and aerosol properties are presented in section 4. Based on the comparisons, we analyse more specifically the concentration of smoke aerosols over biomass burning sources and during the transport, the altitude of BBA, as well as absorbing properties and induced SW heating rate due to smoke. In addition, the impact of the elevated relative humidity within smoke plumes on optical properties is also investigated in section 4. Finally, section 5 focuses on the analyses of the SW **DRE** exerted by smoke aerosols at TOA during September 2016, as well as their impact on the continental climate in terms of the surface energy budget (temperature, sensible heat fluxes) and lower troposphere dynamics (planetary boundary layer (PBL), notably).

## **2. The regional ALADIN-Climate model**

## 2.1. Aerosol Scheme

The recent aerosol scheme (TACTIC, Tropospheric Aerosols for Climate in CNRM-CM) included in the ALADIN-Climate model accounts for sulfate, organic (OC) and black (BC) carbon, dust and primary sea-salt particles (Nabat et al., 2015b; Michou et al., 2015). The biomass-burning emissions from the CMIP6 inventory have been used for BC, OC and sulfur gaseous SO<sub>2</sub>. In **the model ALADIN-Climate**, mineral dust and sea-salt emissions are interactively connected with surface meteorological fields and soil properties (Nabat et al., 2015a). The emission of mineral dust is taken into account following ~~the~~ Marticorena and Bergametti (1995) **parameterization** and the current formulation for primary sea-spray is based on Schulz et al. (2004). This model includes advection by atmospheric winds, diffusion by turbulence, surface emissions as well as dry and wet (in-cloud and below cloud) removal processes.

For the primary BC and OC species and secondary sulfates, a bulk approach is applied whereby a fixed aerosol size distribution is assumed for calculating aerosol properties, while for mineral dust and sea salt particles, a more explicit size representation is used based on 3 bins for dust and sea-salt. The TACTIC scheme assumes an external mixture of the different aerosol species. For specific situations, this could potentially represent a limitation, especially with regard to possible BC mixing (internal/external) state, which can significantly affect SW absorption (Fierce et al., 2016) **by aerosols**. Knowing that, specific attention is being paid in this study to the simulated absorbing properties (SSA) of BBA, as well as the associated SW heating ~~rate~~.

The radiative properties (mass extinction efficiency (MEE), SSA, and asymmetry parameter (ASY)) of each aerosol species are calculated for the different spectral bands of the Fouquart and Morcrette radiation scheme (FMR, Morcrette, 1989) and the rapid radiative transfer model (RRTM, Mlawer et al., 1997), for the SW and Longwave (LW) radiations, respectively. Aerosol **DRE** at the surface and at TOA (in SW and LW spectral range and for both clear-sky and all-sky conditions) is diagnosed using a double call (with and without aerosols) to the radiation schemes during the model integration. In addition, the semi-direct radiative forcing, which represents the modifications of the cloud properties and atmospheric dynamics due to absorption of SW radiations by smoke, is derived from the direct effect. In its current version, BBA are represented by two different tracers (primary BC and OC) with fixed microphysical and radiative properties without any consideration of possible differences between fossil fuel and biomass-burning emissions. This hypothesis implies that the radiative, hygroscopic properties and e-folding time (aging) of carbonaceous species are similar for both anthropogenic and smoke emissions.

## 2.2 Smoke Radiative properties

210 Two tracers have been recently implemented in ALADIN-Climate describing, respectively, the mass concentration of fresh (less hygroscopic) and aged (more hygroscopic) smoke aerosols, following the methodology presented in Bellouin et al. (2011). This allows to distinguish aerosols from biomass burning and anthropogenic emissions and to monitor specific properties, such as e-folding time, hygroscopic and optical properties. In the ALADIN-Climate model, aging from the fresh mode to hygroscopic aged is quantified using an e-folding time of 6 hours according to Abel et al. (2003). This value is two times higher than the one (~3 h) recently proposed by Vakkari et al. (2018) for southern African savannah. The smoke over the SAO is expected to have aged by 5-7 days (Adebiyi and Zuidema, 2017; Diamond et al., 2018). While studies of the BBA chemical composition and attribution for the smoke's optical and hygroscopic properties are still ongoing, preliminary results indicate smoke aging increases its ability to function both as a cloud condensation nucleus and to absorb SW shortwave radiation (Zuidema et al., 2018).

For each tracer, dry-state aerosol size distributions are assumed based on lognormal function (Table 1) similar to those implemented in the Hadley Centre global climate model, HadGEM2-ES (Bellouin et al., 2011). The smoke dry-state refractive indices used to calculate radiative properties are also reported in the Table 1 (at 550 nm). The values of the real and imaginary refractive indices have been updated using the AERONET observations obtained by Eck et al. (2013) in Zambia (Mongu Inn station). Although they indicate a pronounced seasonal cycle in the real and imaginary parts of the refractive index from AERONET data, we have used a mean value of 0.03 (at 550 nm) for the imaginary component in our Mie calculations (Table 1). This represents an important limitation and the seasonal cycle of smoke absorbing properties (average SSA at 440 nm from 0.83 in early July to 0.92 in mid-October, as noted by Eck et al., 2013) is not well represented in ALADIN-Climate. Due to period investigated in this study (August 2016), the implications are expected to be moderate as smoke are highly absorbing for this month over the continent and SAO (Eck et al., 2013, Zuidema et al., 2018). However, this would be a more severe limitation for simulations encompassing the full biomass burning season.

SW radiative properties have been calculated for the specific wavelength bands of the FMR radiation scheme. The values in the SW spectral ranges are reported in the Table 1. At 550 nm and in dry state, the calculated radiative properties are 4.05 m<sup>2</sup>.g<sup>-1</sup>, 0.84 and 0.51 for the MEE, SSA, and ASY for the « fresh » smoke tracer (Table 1). The values for « aged » smoke are, respectively, 5.05 m<sup>2</sup>.g<sup>-1</sup>, 0.90 and 0.58 (Table 1). The MEE used in the model for « aged » smoke is found to be consistent with those reported by Reid et al. (2005). As BBA are known to be hydrophilic (Rissler et al., 2006), the dependence of the radiative properties to relative humidity (RH) has been included for both tracers. This dependence is formulated as described by Solmon et al. (2006) :

$$MEE_{\text{wet}} = MEE_{\text{dry}}(1-RH)^{-\alpha} \quad (1)$$

245 where  $MEE_{\text{wet}}$  and  $MEE_{\text{dry}}$  are for wet and dry conditions. We have selected a value of 0.26 and 0.15 for the parameter  $\alpha$  in order to reproduce the changes of MEE with RH for aged and fresh smoke, respectively. At very high humidity ( $RH > 99\%$ ) maximum thresholds of 8.5 and 16.9  $\text{m}^2 \cdot \text{g}^{-1}$  are considered for fresh and aged smoke, in order to avoid unrealistic values of MEE. In a similar way, we have also implemented a dependence of smoke SSA on RH using the same relationship as  
250 (1) (Mallet et al., 2017). The values of  $\alpha$  have been fixed to 0.015 (0.02) for aged (fresh) smoke to represent the variations of SSA with RH, as reported in Bellouin et al. (2011).

### 2.3 Aerosols and Cloud interactions

“Aerosol-cloud” interactions were represented using a simple parametrization, ~~similar to most GCMs,~~ thereby maintaining the low numerical costs necessary for climate and ensemble  
255 simulations. The activation of hydrophilic particles to cloud droplets is not explicitly resolved ~~in ALADIN-Climate~~ and the first indirect radiative effect is implemented for hydrophilic sulfates, organic carbonaceous and sea-spray aerosols. This first indirect effect is represented by a simple relationship in ALADIN-Climate relating the mass of hydrophilic aerosols to the cloud droplet number concentration (CDNC) based on the work of Martin et al. (1994). The radiative properties  
260 (COD, SSA and ASY) of liquid clouds are calculated in the SW spectral region by the parameterizations proposed by Slingo and Schrecker (1982). In the present work, we do not discuss possible first indirect effects between BBA and Sc which will be addressed and analyzed in a specific future companion study. The impact of aerosols on liquid clouds via the second indirect effect (precipitation modulation due to the hygroscopic aerosols) is currently under development. In  
265 these simulations, the autoconversion rate from water cloud to rain is not sensitive to the aerosol loading and the value of  $8 \cdot 10^{-4} \text{ kg} \cdot \text{kg}^{-1}$  (Smith et al., 1990) is used for the critical cloud water mixing ratio.

## 3. Model Configuration and Data Used

### 3.1. Simulation Design and Important Physics Options

270 The ALADIN-Climate simulations cover the period from 01 August to 31 October 2016. The lateral boundary conditions are provided by ERA-INTERIM (Dee et al., 2011). The possible long-range transport of BBA is not forced at the lateral boundary conditions but rather a large domain (Latitude : -37.1°S to 09.4°N ; Longitude :-33.4°W to 45.4°E) is defined encompassing the main biomass-burning sources. The horizontal resolution of the model is 12 km with 91 vertical levels  
275 (from 1015 to 0.01 hPa). The land surface is treated using the SURFEX model (Masson et al., 2013). As detailed latter, we also use a spectral nudging method described in Radu et al. (2008). The FMR (RRTM) radiative transfer scheme is used to calculate the SW (LW) radiation. Finally, it



should be mentioned that the possible impact of BBA reducing the sea-surface temperature (SST) is not treated here and the ALADIN-Climate model is used in a forced mode configuration (with fixed  
280 SST). This possible impact of BBA is outside the scope of the present study.

As mentioned previously, the biomass-burning emissions from the CMIP6 inventory have been used ~~for BC, OC and sulfur gaseous SO<sub>2</sub>~~. These emissions are monthly-averaged, reconstructed for the 1997–2015 period from the Global Fire Emissions Database version 4 with small fires (GFED4s). The methodology is described by van Marleet al. (2017). In this work, the ALADIN-  
285 Climate simulation uses one of the latest historical year (2014) for BC and OC emissions, which cannot allow the model to reproduce precisely the daily aerosol variations especially when they are controlled more by emissions than dynamical aspects. However, it should be noted that the AOD anomaly for September 2014 compared to the 2008-2015 CAMS reanalyses period (Figure S1 in Appendix), indicates moderate differences of about ~0.05-0.1. Additionally, Sayer et al. (2019)  
290 found similar above-cloud and total-column AOD over the southern Atlantic Ocean in 2014 and 2016. These suggest that the effect of potential differences between 2014 and 2016 over the biomass burning region is likely to be small. One of the interest of using such methodology is to evaluate the ALADIN-Climate model in its climate configuration, which will be exactly the same as used to address the radiative and climatic impact of BBA at climatic scale.

Following the study of Petrenko et al. (2017), an adjustment factor of 2.5 is applied to the biomass burning emissions. BBA are emitted into the first vertical level of the model, without any considerations of pyroconvective processes, as no clear consensus on such processes exists over this region. For example, Labonne et al. (2007) showed that smoke plumes are generally confined in the ~~planetary boundary layer (PBL)~~ close to the main biomass-burning source regions. In the  
300 simulation, fire emissions from the savannah are emitted at the lowest model level, allowing subgrid-scale turbulence mixing through the boundary layer. The diurnal cycle of smoke emission is not taken into account, which could impact the temporal variations of the aerosol loadings (Xu et al., 2016). We assume that the main smoke emissions transported over SAO are included in the domain defined in Figure 1. Finally, a climatology is used ~~in the model~~ for organic aerosols  
305 produced from vegetation biogenic emission and water vapor released from biomass combustion is not treated in the model.

The BBA mass is known to increase during aging due to the condensation of volatile organic compounds. In the absence of a explicit representation of secondary organic aerosols (SOA) production in ALADIN-Climate, a ratio of particulate organic matter (POM) to primary OC has  
310 been used for artificially representing SOA formation within the smoke plume, when the mass is transferred from the fresh to aged mode in the model. The lack of a complete representation of SOA

in current climate models obviously represents an important source of uncertainties in the estimation of BBA concentration (Johnson et al., 2016). For this ALADIN-Climate simulation, an average POM/OC ratio of 2.3 is applied (Formenti et al., 2003) based on SAFARI-2000 data. This value is consistent with the recent results obtained by Vakkari et al. (2018) and higher than the one (1.6) retained in the HadGem global model (Bellouin et al., 2011, Johnson et al., 2016).

Regarding the large uncertainties related to the POM to OC ratio, two sensitivity tests using different POM to OC ratios (2 and 3) have been performed, showing an important impact of  $\pm 0.15$  on BBA AOD over the continent (Figure S2 in the supplement material). An additional simulation tested the sensitivity of BBA AOD to the e-folding time using the recent value proposed by Vakkari et al. (2018). The results (Figure S2 in Supplement Material) indicate a slight AOD decrease of about -0.05 when averaged over the box\_S (15-25°E / 5-15°S, see Figure 1), suggesting in this case a higher sensitivity of AOD to the choice of the POM to OC ratio.

Four ALADIN-Climate simulations (excluding the sensitivity tests only shown in Figure S2, Supplement Material) are performed. The first one (control run, CTL) does not take BBA into account, while the second simulation (named SMK) includes the direct and semi-direct radiative effect of BBA. As mentioned previously and as absorbing properties of smoke are fixed, the seasonal variations of smoke SSA during the biomass burning season as described in Eck et al. (2013) is not represented in the model. To address this limitation, a simulation (named SMK\_SSA) has been performed using less absorbing smoke (SSA of 0.92 at 550 nm). Finally, a nudged (~~on wind and relative humidity~~) simulation (named SMK\_SN) investigates more specifically the impact of the water vapor transported within the smoke plume on BBA optical properties and the associated SW radiative heating. For the latter, the nudging does not affect PBL, which can be independently influenced by the radiative effects of smoke.

### 3.2 Surface, Aircraft, Satellite and Reanalysis dataset

Different datasets of aerosol and cloud properties from surface, remote-sensing, and reanalysis, have been used for evaluating the ALADIN-Climate simulations. Satellite and reanalysis data are summarized in the Table 2.

#### 3.2.1. LASIC Surface Observations (Ascension Island)

SSA (at 529 nm) at Ascension Island was estimated from the in-situ measurement of the scattering coefficient estimated by a nephelometer and the absorption coefficient deduced from a Particle Soot Absorption Photometer (PSAP). The PSAP measurements incorporate an average of the Virkkula (2010) and Ogren (2010) wavelength-averaged corrections, and are collected at standard temperature and pressure, with dilution corrections applied. The RH of the air entering the PSAP is estimated to be 25% or less, while the air entering the nephelometer is measured, with values

ranging between 45%-60%. Differences in the RH are speculated to bias the SSA higher rather than lower, because drying will reduce the coating thickness on the refractory black carbon, reducing lens-induced enhancement of shortwave absorption. The original nephelometer scattering measurements at 550 nm are converted to estimated values at 529 nm using the scattering-derived angstrom exponent. These measurements are also reported in Zuidema et al. (2018). An independent evaluation of the SSA in August-September 2017 using an Aerodyne Cavity Attenuated Phase Shift-SSA instrument is consistent with the values reported here (Tim Onasch, personal communication).

### 3.2.2. AERONET retrievals

Two continental (Mongu Inn and Lubango) and one maritime (Ascension Island) AERONET sites extend local comparisons to the atmospheric column and for different aerosol variables. As described by Dubovik and King (2000), the AERONET network allows retrieval of microphysical (volume size distribution) and optical (refractive indexes, SSA, ASY and scattering/absorption optical depth) properties of aerosols, as well as their spectral dependence in SW spectral range. The uncertainty of retrieved SSA is  $\pm 0.03$  for AOD (440 nm)  $> 0.2$  for water soluble aerosols and for AOD (440 nm)  $> 0.5$  (zenith angle larger than 50 degrees) for desert dust and BBA. For AOD (440 nm)  $< 0.2$ , the SSA accuracy is  $\pm 0.05$ – $0.07$  (Dubovik et al., 2000). In this study, we focus our analyses on Level 2 AOD and SSA AERONET products from AERONET version 2.

### 3.2.3 Aircraft Observations

#### 3.2.3.1 Aerosol Extinction profiles

The NASA Langley 2<sup>nd</sup> generation High Spectral Resolution Lidar (HSRL-2) has been in operation during ORACLES-1 aboard the NASA ER2. HSRL-2 measures particulate backscatter and extinction at 355 nm and 532 nm using the HSRL technique (Shipley et al., 1983) and aerosol backscatter at 1064 nm. All three wavelengths also measure depolarization. The HSRL technique uses a separate filtered channel at each HSRL wavelength that is sensitive to molecular scattering, but not aerosol scattering. This channel therefore provides a direct observation of the attenuation of the signal and allows for the direct calculation of aerosol extinction without external constraints or assumptions on the aerosol optical depth or lidar ratio. From this, the vertically-resolved particulate extinction is derived by comparison to a molecular density profile from direct measurement or a model. For ORACLES-1, the HSRL-2 retrieval uses molecular density profiles from MERRA2 (Gelaro et al. 2017). The filtering is accomplished with an iodine gas filter at 532 nm (Hair et al. 2008) and a density-tuned field-widened Michelson interferometer at 355 nm (Burton et al. 2018). More information about the instrument, calibrations, and algorithms is given by Hair et al. (2008) and Burton et al. (2015, 2018). The vertical resolution for extinction is 315 m and for backscatter and depolarization is 15 m. The horizontal resolution is 60 seconds for extinction and 10 seconds

380 for backscatter and depolarization, or approximately 10 km (extinction) and 1.8 km (backscatter).  
The ORACLES HSRL-2 extinction product can be found at  
<https://espoarchive.nasa.gov/archive/browse/oracles/id8/ER2>.

#### 3.2.3.2 SW Heating Rate estimates

385 Heating rate profiles segregated by absorber (aerosols, water vapor, oxygen) are determined using  
the spectral information from the Solar Spectral Flux Radiometer (SSFR). SSFR measures  
upwelling (nadir) and downwelling (zenith) irradiance from 350-2100 nm. The zenith light collector  
is actively leveled, which allows SSFR to obtain spectral irradiance measurements throughout spiral  
profiles that extend from the top of the aerosol layer to the bottom of the cloud layer. These  
measurements lend themselves to a new algorithm for retrieving aerosol SSA and ASY from 350-  
390 860 nm. It uses measurements made during the spiral aircraft descents to separate changes in  
upwelling, downwelling and net irradiance due to the aerosol layer from those due to the underlying  
cloud field (Cochrane et al., 2018).

The 4STAR spectral AOD, HSRL-2 extinction profiles, and the spectral SSA and ASY retrievals,  
provide the inputs for the heating rate profiles calculated with the LibRadtran radiative transfer tool  
395 (Mayer and Kylling, 2005). The retrieved intensive aerosol properties SSA and ASY are vertically  
homogeneous, whereas the spectral extinction coefficient from the merged 4STAR HSRL-2  
measurements varies with altitude. The cloud albedo below the aerosol layer is directly measured  
by SSFR (the reflectance has been estimated around  $\sim 800$  m above cloud top), and the atmospheric  
water vapor profile is determined from in situ measurements. Using these inputs, two calculations  
400 of the heating rate profile are done: one with, and one without aerosols, and the difference is  
reported as aerosol heating rate. The accuracy of the calculation is ensured by comparing the  
calculated irradiance spectrum above and below the aerosol layer with the SSFR measurements.  
After this step, the heating rate is spectrally integrated over the solar wavelength range (380-2125  
nm).

#### 405 **3.2.4 Satellite Retrievals**

##### 3.2.4.1 MODIS and MISR dataset

The Deep Blue ACAOD retrieved from the MODIS instruments is described in Sayer et al. (2019),  
which is a slightly updated version of the demonstration algorithm presented in Sayer et al. (2016).  
In brief, this algorithm performs a multispectral weighted least-squares fit of measured reflectance  
410 in four bands across the visible spectral region (centered near 470, 550, 650, and 870 nm) to  
simultaneously retrieve ACAOD and the COD. The minimization is performed using optimal  
estimation. This provides estimates of the uncertainty on retrieved parameters, as well as an  
indicator of how well the retrieval solution is able to fit the measurements. Retrievals where the

forward model is expected to be inappropriate, or where the measurements do not constrain the  
415 retrieved quantities, are filtered out.

The main updates since Sayer et al (2016) are twofold. First, the radiative transfer lookup tables  
have been updated to include dimensions for surface pressure and surface albedo, which improves  
the realism of the forward model over land. Surface pressure is estimated using terrain altitude,  
while the surface albedo is taken from a climatology based on the MODIS gap-filled snow-free land  
420 albedo data set (Sun et al., 2017). The second is that, rather than applying the retrieval to each  
individual pixel, the pixels are aggregated into 10 km x10 km (effective nadir resolution) boxes,  
chosen to match the resolution of the level 2 MODIS aerosol products. Then, the median reflectance  
of water cloud pixels within each box is used for the retrieval. Use of median reflectance decreases  
sensitivity to factors such as 3D effects or cloud detection errors. Additionally, if a 10 km x 10 km  
425 pixel has a water cloud fraction under 0.75, it is excluded, for the same reason.

The MOD06ACAERO (Meyer et al., 2015) products are also used. These use reflectance  
observations at 6 MODIS spectral channels (0.46, 0.55, 0.66, 0.86, 1.24, and 2.1  $\mu\text{m}$ ) to  
simultaneously retrieve ACAOD, and the COD and the cloud effective radius (CER) of the  
underlying marine boundary layer clouds. Retrievals are performed at the pixel level (here, every  
430 fifth native 1 km pixel) on both Terra (morning) and Aqua (afternoon) MODIS data. Output  
includes pixel-level estimates of retrieval uncertainty that accounts for known and quantifiable error  
sources (e.g., radiometry, atmospheric profiles, cloud and aerosol radiative models). Assumptions  
regarding the cloud forward model and ancillary data usage are consistent with those of the  
operational MODIS cloud products (MOD/MYD06) (Platnick et al., 2017). Note that both these  
435 data sets represent only the partial column AOD, i.e. the AOD above the liquid cloud top, and that  
ice phase clouds are not processed. In addition, we have also used MODIS-Terra and Aqua  
combined Deep Blue/Dark Target data set  
(AOD\_550\_Dark\_Target\_Deep\_Blue\_Combined\_Mean\_Mean) from the latest Collection 6.1  
(Sayer et al., 2014) and MISR (MIL3MAE monthly mean data at  $0.5^\circ$  of resolution; Kahn et al.,  
440 2015) (see Table 2).

#### 3.2.4.2 OMI dataset

The Ozone Monitoring Instrument (OMI) sensor, operating since October 2004 onboard of the EOS  
Aura satellite, is a spectrometer with a high spectral resolution (Levelt et al., 2006). OMI offers  
nearly the daily global coverage with a spatial resolution for the UV-2 and VIS (UV-1) channels  
445 ranging from  $13 \times 24 \text{ km}^2$  at nadir. The OMAERUV\_v003 product contains retrievals from the  
OMI near-UV algorithm (Torres et al., 2007). This algorithm derives a variety of aerosol radiative  
properties, such as an aerosol index (AI), AOD, AAOD (uncertainty of  $\pm (0.05 + 30\%)$ ) for clear-

sky conditions. For this study, we have used ACAOD (Jethva et al., 2018) retrieved at 500 nm (Table 2). An above-cloud aerosol retrieval technique was also applied to the multi-year record of OMI observations to deduce a global product of ACAOD on a daily scale (Jethva et al., 2018).

#### 3.2.4.3. SEVIRI dataset

Spatiotemporally highly resolved geostationary satellite observations are taken here from the CCloud property dAtAset based on SEVIRI edition 2 (CLAAS-2, Benas et al., 2017). The CLAAS-2 dataset is based on measurements of the Spinning Enhanced Visible and Infrared Imager (SEVIRI) and was generated and released by the EUMETSAT Satellite Application Facility on Climate Monitoring (CM SAF) as the successor of CLAAS (Stengel et al., 2014). CLAAS-2 includes a variety of cloud properties, of which LWP, COD and CER were used in this study. CLAAS-2 COD and CER are retrieved, similarly to the widely used cloud retrieval method described in Nakajima and King (1990), under the assumption of plane-parallel cloud layers. Lookup tables are pre-calculated and used to map SEVIRI reflectance at 0.6  $\mu\text{m}$  and 1.6  $\mu\text{m}$  wavelengths to COD and CER as function of satellite-sun geometries and cloud phase. For liquid clouds, COD and CER are used to calculate LWP following Stephens (1978). The algorithm is initially described in Roebeling et al. (2006) with more details and updates given in Benas et al. (2017), which also report validation exercises. The CLAAS-2 Level-2 data are instantaneous data on native SEVIRI resolution with a temporal resolution of 15 minutes. For this study, the data are projected onto a regular latitude-longitude grid using nearest neighbor approach. It should be noted that Sc cloud retrievals could be affected by the presence of BBA over SAO. Recently, Seethala et al. (2018) indicate that, in the aerosol-affected months of July, August and September, SEVIRI LWP (based on the 1.6 $\mu\text{m}$  CER) is biased by  $\sim 16\%$ .

#### 3.2.5. Reanalyses of atmospheric composition

Two different reanalyses are used for aerosols and clouds (Table 2). The European Center for Medium-Range Weather Forecasts (ECMWF) reanalysis of global atmospheric composition since 2003 includes five main aerosol species. The first generation of ECMWF reanalysis (Morcrette et al., 2009), issued by the GEMS (Global and regional Earth-System (atmosphere) Monitoring using Satellite and in situ data) project, covers the period 2003-2008. The Monitoring Atmospheric Composition and Climate (MACC) is the second-generation product and provides improvements in sulfate distributions and has extended to the 2003-2011 period (Benedetti et al., 2009). Here, we use MACC NRT daily data sets at 1.125 $^\circ$  resolution of the anthropogenic SW direct effect forcing at TOA in all-sky conditions (Table 2). In addition, we use the Modern-Era Retrospective Analysis for Research and Applications (MERRA2) reanalysis, generated with version 5.2.0 of the Goddard Earth Observing System (GEOS) atmospheric model and data assimilation system (DAS). The

system, the input data streams and their sources, and the observation and background error statistics are fully documented in Rienecker et al. (2011). We rely on the AOD for the different species at  $0.5^\circ \times 0.625^\circ$  spatial resolution (Table 2).

## 485 **4. Microphysical and Optical properties of Sc and BBA**

### **4.1. Sc properties**

The different properties of Sc are analysed over the box  $10\text{-}20^\circ\text{S} / 0\text{-}10^\circ\text{E}$ , defined by Klein and Hartmann (1993), referenced in the following as box\_O.

#### **4.1.1. Macrophysical and Microphysical Sc properties**

490 Figure 2a reports the daily mean LWP ( $\text{g}\cdot\text{m}^{-2}$ ), LCF as well as liquid COD, averaged over the box\_O for September 2016. A reasonable agreement is evident in the LWP between ALADIN-Climate (SMK simulation), SEVIRI and ERA-INT data. The simulated LWP is in the range of ERA-INT and SEVIRI data, if slightly less, by  $-8 \text{ g}\cdot\text{m}^{-2}$  in the mean compared to ERA-INT. The LWP maxima are also well simulated, especially for the 25-28 September, with LWP  $\sim 90 \text{ g}\cdot\text{m}^{-2}$ . The temporal  
495 correlation is about  $\sim 0.45$  between ERA-INT and the SEVIRI LWP.

In contrast, an important negative bias of approximately  $\sim -20\%$  is detected in the simulated low cloud fraction compared to ERA-INT values (Fig. 2b). This result is related to a well-known bias (underestimates of low cloud fraction) detected in most GCMs over the Sc regions (Nam et al., 2012). The mean modeled LCF is 57%, compared to 80% in the ERA-INT dataset. The poor  
500 representation of LCF over Sc regions remains an open issue outside of the scope of this work. Nevertheless, the analysis and discussions of the following results take into account this underestimates, notably for SW heating rate and the **DRE** exerted by BBA at TOA.

#### **4.1.2. Optical Sc properties**

Figure 2c represents the daily COD estimated by ALADIN-Climate using the Slingo et al. (1998)  
505 and Nielsen et al. (2015) parameterizations, which are used to calculate liquid cloud optical properties at different wavelengths from LWP and CER and different spectral coefficients. This figure also includes an ALADIN-Climate simulation with a fixed CER of  $10 \mu\text{m}$ , close to the SEVIRI values (Figure S3 in Supplemental Material). Retrieved values from MODIS and SEVIRI are also reported in Figure 2c. The SEVIRI and MODIS-Aqua values are consistent with each other,  
510 for a mean COD of  $\sim 8$  for both instruments. The COD is overestimated by the model in both configurations but especially when the Slingo parameterization is used (average COD value of 11.5). The Nielsen parametrization slightly reduces the bias for a mean COD of 9.5. As the LWP is realistically simulated by the model, this negative bias could be due to errors (underestimates) in the simulated CER by ALADIN-Climate. Our sensitivity test conducted using a fixed CER of  $10 \mu\text{m}$   
515 indicates a reduced bias (+0.25). As mentioned previously, the BBA indirect radiative effect is not

addressed in this study but is known to be an important issue over SAO (Costantino and Bréon, 2013, Lu et al., 2017). The indirect radiative effect is also relevant to **DRE** as it strongly depends on the albedo of the underlying Sc. The impact of BBA on cloud microphysical properties will be studied in a future work.

## 520 **4.2. Biomass Burning Aerosols**

### **4.2.1. AOD over biomass burning source**

Figure 3 compares the (monthly mean) total (clear-sky, not above-cloud) AOD simulated by ALADIN-Climate to those derived from MODIS-Terra and Aqua combined Deep Blue/Dark Target data set (AOD\_550\_Dark\_Target\_Deep\_Blue\_Combined\_Mean\_Mean) from the latest Collection 6.1 (Sayer et al., 2014) and MISR. The total AOD obtained from the MERRA2 and CAMS reanalyses are also indicated. All AODs are at 550 nm. Over Angola and Zambia, the model is able to simulate a regional pattern of AOD consistent with the MODIS and MISR retrievals, with AODs of ~0.7-0.9, even if BBA are located too far south compared to satellite data. A general good agreement is also observed compared to the CAMS reanalyses, even if the model underestimates AOD over eastern Gabon. Over the continent, significant differences (overestimates) appear clearly 530 compared to MERRA2. This difference with reanalyses product can be due to the scaling factors applied for biomass burning emissions in the different models. Over the ocean, ALADIN-Climate is found to be very consistent with MERRA2 (AOD ~0.7 near the Angola coast) but the differences are large compared to the satellite retrievals. The latter AODs are higher (~0.7-1.0), especially for 535 MODIS. It should be mentioned that 50% of MODIS AOD over the SAO is due to coarse mode according to the level 2 retrievals.

In addition, the land-ocean contrast in AOD detected by MODIS and MISR, with lower AODs over the continent and higher AODs over the ocean, is not observed in ALADIN-Climate. This contrast in AOD is still detected in MERRA2 data, but it is not as large as in the satellite retrievals. It is 540 likely that some of the land/ocean contrast in the satellite data comes from different factors. The first is that the over-land and over-water algorithms are different and may have different biases. The second is that cloud fraction is also significantly higher over the water than over the land, meaning that typically more days of data contribute to the monthly mean over land than over water. Both of these effects could suppress or enhance any real land-ocean contrast in the AOD. Finally, part of the 545 contrast could be due to the presence of sea-spray aerosols, certainly about ~0.1. As well as continued refinement of AOD retrieval algorithms, it is recommended that future work attempts to quantify the potential magnitude of these sampling effects on land/ocean contrast, which have received comparatively little attention to date. Finally, the satellite data also indicate a larger spatial extent to the aerosol loading over the ocean compared to ALADIN-Climate. This difference could



550 be due to possible overactive aerosol deposition in our simulations. A specific section (4.2.2) is dedicated to using ACAOD products for evaluating the model over the ocean.

In order to make comparisons over the continent, a second box (box\_S, 15-25°E / 5-15°S, see Figure 1) is defined over biomass burning sources. Figure 4a indicates the daily-mean AOD at 550 nm averaged over box\_S from MODIS, MERRA2 and ALADIN-Climate. There is a good agreement, with monthly means of 0.59, 0.62 and 0.49 for MODIS, ALADIN-Climate and MERRA2, respectively. The model is very consistent with the MODIS retrievals and slightly higher than MERRA2. Differences between MERRA2 and ALADIN-Climate may reflect the different model biomass burning emissions. Compared to MODIS, a small mean bias of +0.04 is found in the model over smoke sources. In terms of temporal correlation, the score is worse compared to MODIS. As mentioned previously, this could be due to the time frequency of biomass burning emission (monthly-mean emission) imposed in the ALADIN-Climate model. The absence of spectral nudging in the ALADIN-Climate simulation can also explain part of the low temporal correlation, which is clearly higher (0.73) for MERRA2 data.

Data from three AERONET stations (Mongu Inn (Zambia), Lubango (Angola) and Ascension Island) provide additional AOD evaluation. Figure 5 indicates the daily-mean AERONET and ALADIN-Climate (only daytime values) AOD at each station. At Lubango, the model is able to correctly simulate the AOD, except in the beginning of September when it is overestimated. The maxima (AOD of 0.6) of 20 and 27 September are also well represented by the model. The total monthly-mean AOD simulated by ALADIN-Climate (0.41) is consistent with AERONET (0.43), if with a small mean negative bias. Over the Mongu Inn station, the comparisons indicate a more pronounced negative bias (mean value of -0.14). This is due to a nearly constant underestimate of total AOD throughout the period of the simulation, leading to a simulated monthly-mean of 0.47, lower than the one observed (0.61). In parallel, the two maxima detected in AERONET data are well captured by the model. The first one, occurring between the 18 and the 21 September, is simulated too early and its magnitude is overestimated by about ~0.2. The second (23-29 September) is better reproduced in terms of magnitude (~ 1.0) but a significant underestimate in its duration is observed. Indeed, the BBA event starts around the 22 September in the observations, while it is simulated between 26 and 29 September by the model. Finally, the model is able to correctly simulate the magnitude of AOD along the transport over SAO to the remote location of Ascension Island (Figure 5). The simulated monthly-mean value (0.26) is comparable to AERONET (0.21) with a small positive bias, primarily because of overestimates during the 06-10 September period. This suggests that winds and aerosol deposition are also well represented. To summarize, the analysis of AOD comparisons demonstrate that the ALADIN-Climate model

reasonably simulates the magnitude of AOD during September 2016, even if some biases are  
585 detected (possibly due to the monthly-mean emissions imposed in the model that do not allow the  
model to represent some daily variations precisely), especially over the ocean (a negative bias,  
primarily near the coast) when compared to satellite data.

#### 4.2.2 ACAOD over SAO

Due to the significant presence of Sc over SAO, the use of satellite clear-sky AOD products as a  
590 model evaluation tool is limited. This limitation is overcome with new retrievals of ACAOD from  
MODIS and OMI, summarized in Table 2. Figure 6 indicates consistent estimates of monthly-mean  
ACAOD between ALADIN-Climate, MODIS-DB, MOD06ACAERO and OMI. It should be noted  
that the ACAOD is calculated as the integration of the aerosol extinction from the cloud top to the  
model top in the ALADIN-Climate model. For all independent estimates, Figure 6 indicates values  
595 of about  $\sim 0.4-0.5$  (550 nm) near the Angola coast. ACAOD then decreases to  $\sim 0.2$  over SAO.  
ACAOD is underestimated by ALADIN-Climate over the ocean, especially when compared to the  
two different MODIS products. Indeed, MODIS-DB and MOD06ACAERO data reveal a larger  
regional extent over the ocean compared to the model. The ACAOD extent is less pronounced in the  
OMI data, which is thereby more consistent with ALADIN-Climate.

600 Additional comparisons were performed over box\_O using MODIS-DB products. Figure 4b  
indicates daily-mean model ACOAD obtained for the SMK and SMK\_SN (not discussed in this  
part) simulations and MODIS-DB. The monthly-mean ACAOD is underestimated by the model  
(SMK configuration, red dotted line) with a mean value of 0.20 (at 550 nm), 0.06 less than MODIS-  
DB (0.26). The ALADIN-Climate simulation underestimates two maxima observed by MODIS-DB,  
605 around 03-05 and 20-24 of September, explaining part of the negative bias (-0.06) in the SMK  
simulation. For the rest of the time period, we observe a realistic estimation of ACAOD by the  
model. It should be mentioned that additional analyses of the simulated ACAOD is also discussed in  
Shinozuka et al. (2018) for different boxes defined in the ORACLES-1 program, indicating similar  
results with an underestimate of ACAOD. As for AOD, ALADIN-Climate is shown here to simulate  
610 realistically the concentration of aerosols transported above clouds over SAO. This allows to  
address the SW **DRE** exerted by BBA at TOA during ORACLES, after investigating the vertical  
structure (part 4.2.3) and SW absorbing properties (part 4.2.4).

#### 4.2.3 BBA Vertical Structure

##### 4.2.3.1 ALADIN-Climate Extinction vertical profiles

615 The vertical distribution of the modeled BBA extinction is analyzed over the continent and SAO in  
Figure 7 as the monthly-mean extinction vertical profiles (at 550 nm) for two different transects at  
latitudes of 8 and 15°S. For both transects, the highest extinctions are identified over the continent,

close to biomass burning sources, with extinctions  $\sim 0.2 \text{ km}^{-1}$ . The amplitude of BBA extinction coefficient (top and bottom right panels) decreases during the transport, reaching values of  $\sim 0.05$ -  
620  $0.10 \text{ km}^{-1}$  (at 8 and 15°S, respectively) for longitudes near 0°. For both profiles, the top of the smoke plume is around  $\sim 5000 \text{ m}$  over the continent in the simulations, **very consistently** with the altitude of the top plume ~~over the continent~~ reported by Das et al. (2017) from CALIOP ~~lidar~~ observations. This analysis would suggest that injection heights are not that important in this region and aerosols are mainly lofted by convection over the biomass burning sources.

625 Over the SAO, two different well-distinguished aerosol layers are simulated; a first one mainly located in the MBL and mostly due to primary sea spray aerosols and a second BBA layer located above, between 2000 and 4000 m. The top of the marine aerosol plume is simulated around  $\sim 1000 \text{ m}$  ~~in the model~~ and is separated from the smoky layer by a clean atmospheric layer, especially at 15°S, characterized by extinction near  $\sim 0.05 \text{ km}^{-1}$ . For both transects, the top of the smoke plume  
630 decreases from 10°E to 10°W, starting around  $\sim 5000 \text{ m}$  near the coast to reach  $\sim 4000 \text{ m}$  at 10°W. This BBA stratification over SAO is consistent with the vertical structure reported by Das et al. (2017) for latitudes comprised between 0 and 10°S. Indeed, they report a transport of smoke that mainly occurs between 2000 and 4000 m over SAO, contrary to the different models used in this study, which both indicate a more pronounced decline of the altitude of BBA during the transport.

635 This result is also consistent with a previous study from Haywood et al. (2003) over this region. The elevated plume is mainly composed by BBA characterized by a decrease in extinction (mainly due to a decrease of BBA concentration) during the transport from  $0.15 \text{ km}^{-1}$  (at 15°E) to  $0.08 \text{ km}^{-1}$  (near 0°) for the transect at 8°S. Such extinction values are consistent with those reported by Das et al. (2017), who indicated CALIOP extinction around  $\sim 0.1$ - $0.15 \text{ km}^{-1}$  over SAO (for latitudes  
640 between 0 and 10°S). The extinction due to BBA is negligible in the simulation for longitudes highest than  $\sim 10^\circ\text{W}$ , especially for the transect at 15°S. For both transects reported in the Figure 7 (left panels), significant extinctions are simulated within the MBL, with values of about  $\sim 0.2$ - $0.25 \text{ km}^{-1}$  mostly due to the presence of primary sea-spray aerosols. Figure 7 indicates the highest values at 8°S compared to 15°S.

645 Based on the transects, no favorable conditions are identified allowing an efficient mixing of BBA within the MBL during the transport of aerosols over SAO. Such results are found to be different to the schematic view of Gordon et al. (2018), who proposed that an efficient mixing of smoke only occurs around 0° E - 10° W within the MBL. This could limit the possible impact of BBA on cloud droplet concentrations and Sc properties. In the ALADIN-Climate simulations, smoke aerosols  
650 primarily remain above the MBL during transport, with little vertical mixing. For this reason, in the

model, the impact of BBA on the Sc microphysical/optical properties will be primarily through the semi-direct radiative effect.

#### 4.2.3.2 Comparison with HSRL-2 Extinction

Aerosol extinction coefficients derived at three different wavelengths (355, 532 and 1064 nm) by the HSRL-2 instrument permit local 1-D comparisons with ALADIN-Climate simulations. In addition to evaluating the simulated extinction vertical profiles, the spectral dependence of the model calculated extinction can also be evaluated. Figure 8 reports the extinction vertical profiles for three different days (12, 22 and 24 September), as well as RH profiles obtained from MERRA2 and ALADIN-Climate (black dotted and solid lines, respectively). Those specific days have been chosen to represent different locations within Box\_O (Figure 1). The vertical profile of CF simulated by ALADIN-Climate (yellow dotted line) is also included in Figure 8. It should be mentioned that the wavelengths of ALADIN-Climate are not exactly the same as those from HSRL-2, especially for the UV spectral band (355 and 440 nm, respectively). In addition and due to the significant CF, HSRL-2 data are not necessarily available near the surface and remain above cloud top (~2000 m) in most cases.

For September 12, our simulations indicate that the vertical structure of the BBA plume (dashed red, purple and blue lines) is not well represented in the model even though both the model and HSRL-2 place most of aerosol above the MBL. At both times (11:00 and 13:00 UTC), the aerosol extinction coefficients simulated by ALADIN-Climate are overestimated (underestimated) for altitudes between 1500-3000 m (3000-6000 m). This compensation of errors leads to a consistent averaged (between 1500 and 6000 m) integrated extinction in the simulation ( $0.07 \text{ km}^{-1}$  at 550 nm) compared to HSRL-2 observations ( $0.06 \text{ km}^{-1}$ ) at 13:00 UTC but significant underestimates at 11:00 UTC ( $0.05$  and  $0.13 \text{ km}^{-1}$ , for ALADIN-Climate and HSRL-2, respectively). In addition, we observe important biases in the simulated RH, especially at 11:00 UTC between the surface and 3000 m (positive bias) and (negative bias) above 3000 m, which could partly explain the extinction biases.

For the 22 and 24 September, Figure 8 indicates that the altitude of the smoke plume over SAO is realistically represented by ALADIN-Climate, especially for the plume located between 3000 and 6000 m. However, a second aerosol layer, which is observed from HSRL-2 between 2000 and 3000 m for September 24 at 11:00 UTC (Figure 8e) is absent in ALADIN-Climate and simulated below (between 1000 and 2000 m). For the same day at 12:00 UTC (Figure 8f), similar conclusions are obtained with a maxima at ~3500 m which is well simulated by ALADIN-Climate but the second plume (observed around ~2000-2500 m from HSRL-2) is totally absent in the model. For both days, Figure 8 reveals that the magnitude of the simulated extinction is generally underestimated compared to HSRL-2, true at each wavelength. As an example, for 24/09 at 11:00 UTC (Figure 8a),

685 the local maxima ( $\sim 0.30 \text{ km}^{-1}$ ) derived by the HSRL-2 instrument at 5000 m is significantly lower  
( $\sim 0.15 \text{ km}^{-1}$ ) in the model. This is also observed for the second aerosol plume at  $\sim 2500$  m for that  
day. Such conclusions can be drawn for all cases (12, 24 and 26 of September), with a negative  
(mean) bias (indicated only at 550 nm and for the whole atmospheric column) of between -0.01 and  
-0.08  $\text{km}^{-1}$ . This could be attributed to incorrect smoke emissions, e-folding time, OC to POM ratio,  
690 optical properties (especially the mass extinction efficiencies) of BBA, as well as the different  
parameterizations used for representing hygroscopic properties of aged smoke. In section 4.3,  
specific attention is paid to the impact of RH transported within the smoke plume on BBA  
extinctions.

#### 4.2.4 BBA (SW) Absorbing properties and Heating rate

##### 695 4.2.4.1 Absorbing properties at the biomass burning source

The magnitude and the sign of the **DRE** of BBA exerted over SAO is highly sensitive to the smoke  
SSA. The monthly-mean (whole-column integrated) SSA (for the fine aerosols) simulated by  
ALADIN-Climate for September 2016 (Figure S4 in Supplement) indicates values of about  $\sim 0.85$   
(at 550 nm) over a large part of the subcontinent. SSA increases near the coast ( $\sim 0.89$ - $0.90$ ) and  
700 during transport over the SAO ( $\sim 0.92$  to  $0.95$ ). Local comparisons at the two continental  
AERONET stations (Mongu **Inn** and Lubango, Figure S4 in Supplement) reveal a good agreement  
between the simulated and observed SSA, characterized by low bias of about  $+0.01$ / $-0.02$ . A larger  
negative bias is observed and documented at the Lubango site. However, the day to day variability  
is not represented in the model and SSA is nearly constant ( $\sim 0.83$ - $0.84$ ) in the simulation. As an  
705 example, the lowest values ( $\sim 0.79$ - $0.80 \pm 0.04$ ) detected by AERONET are absent in the model.  
The same conclusion is obtained for the highest values derived from observations, especially at  
Lubango (Figure S4 in Supplement). However, such results indicate the ability of the model at  
reproducing absorbing properties of BBA close to biomass burning emissions with limited bias ( $-$   
 $0.02$ / $+0.01$ ).

##### 710 4.2.4.2 Absorbing properties over SAO

The model comparison to in-situ surface-based SSA values at Ascension Island reveals more  
discrepancy. Figure 9 shows the daily-mean SSA obtained at the surface from in-situ observations  
and calculated with ALADIN-Climate at two different altitudes (0.2 and 3 km). The model is not  
able to reproduce the low values (mean of 0.87) observed at the surface. Indeed, near the surface,  
715 the simulation indicates a simulated SSA of nearly 1 for all of the September 2016 period. The  
model MBL optical properties are mainly controlled by primary marine aerosols (see Figure 7)  
leading to SSA close to unity. This highlights also that the mixing of BBA within the MBL is  
possibly underestimated in the model, although LASIC observations also show little smoke is

720 present at the surface in September (Zuidema et al., 2018). The LASIC site is also located on the remote windward side of the island and is not affected by local sources, of which there are few to begin with (no trash burning on the island).

It should be noted that the low values of SSA obtained at Ascension island could reflect long-term ageing processes for the BBA, ~~which that~~ are not currently included in ALADIN-Climate. One indication that the aerosol sampled at the LASIC is aged is through the parameters f44 and f60 (the 725 fraction of the organic aerosol mass spectrum signal at m/z 44 and 60 respectively), in the data from the Aerosol Chemical Species Monitor (Alison Aiken, personal communication). The LASIC f44 and f60 values of approximately 0.2 and 0.002 respectively are characteristic of highly aged aerosols (Cubison et al., 2011). This chemical process could increase the absorbing efficiencies of BBA (Fierce et al., 2016) due to the « lensing » effect (increase of SW radiations reflected to the 730 absorbing core) during the transport (decrease of SSA) and could explain the opposite results obtained in the model, which simulates an increase of SSA (not shown, Figure S4) from biomass burning sources to SAO.

#### 4.2.4.3 SW Heating Rate

735 Figure 10 indicates the SW heating rates only due to BBA for two transects defined at latitudes of 8 and 15°S, similar to Fig. 8. The effect of BBA is isolated by subtracting the heating rates in the simulations without BBA from those with BBA. Significant additional SW heating is simulated over the continent and between 2 and 4 km over SAO due to the presence of absorbing smoke. Over the continent, the additional heating is about ~1 °K by day with maxima near ~1.5 °K by day for altitude of ~4000m at 8°S. The simulated heating is approximately 1°K/day near the coast, 740 decreasing to 0.5°K/day during transport. For both transects, SW heating occurs mainly between 2 and 4 km over SAO. At 15°S, the SW heating is less pronounced than at 8°S, in agreement with the difference in the extinction profiles (see Fig. 7). The SW heating due to BBA absorption is clearly visible only above the MBL and there is no clear additional SW heating within it.

Such values of SW heating due to smoke appear to fall well within the range of values reported by 745 different modeling studies by Tummon et al. (2010), Gordon et al. (2018), Adebisi et al., (2015) or Wilcox et al. (2010), who reported, respectively, additional SW heating due to smoke of 1 (JJAS period), 0.34 (5 days of simulations), 1.2 (for fine AOD > 0.2) and 1.5 °K by day. In addition, Keil and Haywood (2003) estimated a SW heating rate of 1.8°K/day near the coast using a radiative transfer model and observations during SAFARI-2000. The temperature change (estimated through 750 two parallel simulations including or not smoke) due to BBA is about +0.5-0.8 °K between 2 and 4 km (Figure S5 in Supplement material) in a good agreement with the value (+0.5 °K) of Sakaeda et al. (2011) or more recently proposed by Gordon et al. (2018). It should be noted the large impact of

less absorbing BBA (mostly present at the end of the biomass burning season, Eck et al., 2013) on SW heating rate as reported in Table 3. Results obtained with the SMK\_SSA simulations indicate a change from 1.15°K/day (SMK) to 0.58°K/day (SMK\_SSA) over the box\_S.

ORACLES SW heating rates retrieved from the SSFR retrievals of SSA and ASY (see Part 3.2.3.2) in conjunction with HSRL2 extinction profiles (Part 3.2.3.1) are also used to assess our simulations. Figure 11 indicates the instantaneous (12:00 UTC) SW heating (only due to aerosols) vertical profiles obtained for 20 September from SSFR and ALADIN-Climate. Two ALADIN-Climate heating profiles indicate clear-sky and all-sky conditions. Figure 11 indicates that the location of the additional SW heating due to BBA is well represented by the model, with a notable increase between 3 and 4 km, in agreement with the SSFR retrievals. SW Heating between 2 and 2.5 °K/day are simulated at these altitudes with the highest values obtained under all-sky conditions (black dashed lines). However, significant underestimates are observed within the smoke layer, where SSFR observations indicate SW heating of about ~3 to 3.5 °K/ day. As the Sc COD is found to be consistent between simulations and the SSFR cloud retrievals (COD ~9), we hypothesize that the difference in SW heating is due to local underestimates of aerosol extinction as well as of BBA absorption in the model. A second aspect concerns the large underestimate of SW heating around ~1.5 km in the ALADIN-Climate simulation. Indeed, the local maxima of ~2.5 °K/ day obtained from SSFR observations is totally absent in the model, but can be traced back to another layer detected by HSRL.

#### 4.3 Impact of the RH transported within the smoke plume on optical properties

As mentioned previously, a specific simulation (SMK\_SN) that includes the method of spectral nudging (Radu et al. 2008) was also performed. The nudging is applied to wind vorticity and divergence, surface pressure, temperature and specific humidity, using a constant rate above 700 hPa, a relaxation zone between 700 and 850 hPa, while the levels below 850 hPa are free. This simulation was motivated by different studies (Haywood et al., 2003, Adebisi et al., 2015) that indicated a correlation between BBA and specific humidity. In these studies, biomass burning plumes are associated with specific humidities greater than 2 g.kg<sup>-1</sup>, while outside the smoke plumes the values are less than 1 g.kg<sup>-1</sup>. To date, few regional climate modeling studies have investigated the potential role of the relative humidity on smoke optical properties within this specific atmospheric layer, that could, in turn, impact the DRE exerted by BBA. This ALADIN-Climate simulation (SMK\_SN) addresses this specific point.

Figure 4b showing the daily-mean ACAOD (averaged over the box\_O) from MODIS-DB along with the simulations through comparing the ACAOD with and without the nudging of RH, indicates an impact. For SMK\_SN, the negative bias is reduced compared to MODIS-DB and equals to -0.02

(bias of -0.06 for the SMK run). The maxima in ACAOD observed between 19 and 25 September is better reproduced in the SMK\_SN simulation, consistent with a slightly improved temporal correlation (0.42).

790 In addition to the satellite observations, we also used HSRL-2 vertical profiles of extinction (already presented in Figure 8) to investigate the impact of RH on BBA optical properties. Figure 12 shows, for 24 September only, the vertical profiles of RH by MERRA2 and ALADIN-Climate (SMK and SMK\_SN simulations), as well as aerosol extinction (at 550 nm) from HSRL-2 and the model (SMK and SMK\_SN). A significant improvement is evident in the SMK\_SN RH vertical  
795 profiles, reducing the bias with MERRA2, especially at 09:00 and 12:00 UTC. For each case, Figure 12 indicates that RH is better represented in SMK\_SN especially at altitudes where the transport of smoke occurs, i.e., between 2000 and 5000 m (Figure 7).

These changes in RH profiles in SMK\_SN run impact the BBA optical properties for the different cases, notably by increasing extinction within the smoke plume and a remarkable agreement in  
800 extinction is observed between HSRL-2 and the SMK\_SN simulation. As an example, at 09:00 UTC, the improvement of the simulated RH between 3500 and 6000 m in SMK\_SN significantly reduces the bias in the simulated extinction at those altitudes. At 4500 m, the simulated extinction is very consistent with HSRL-2, with maxima around  $\sim 0.2 \text{ km}^{-1}$ . Similar conclusions are also observed for the other cases presented in Figure 12. At 11:00 UTC, important improvements are  
805 found for altitudes between 2000 and 5000 m in the SMK\_SN simulation compared to SMK. At this time, a negative bias persists between 2000 and 3500 m, marked by a bias in RH even in the nudged simulation. These results are consistent with the study of Adebisi et al. (2015) who used CALIPSO smoke extinction profiles to show that the largest extinction coefficients co-occur with high RH ( $\sim 80\%$ ) at the top of the BBA layer. As discussed recently in Kar et al. (2018), this increase could be  
810 due to enhancement of the size of aged smoke during the transport over SAO.

A second important aspect of these results concerns the possible overestimates of the increase of extinction with RH as parameterized in the present version of ALADIN-Climate. As indicated for both cases, excellent agreement is generally observed in the extinction profiles even if some slight negative bias in RH remain. This can be clearly detected at 4500 m (09:00 UTC) or 4000 m (11:00  
815 UTC). At 12:00 UTC and for altitudes between 1000 and 2000 m, RH simulated in SMK\_SN is consistent with MERRA2 data, while the simulated extinction is overestimated.

These original results, using for the first time coincident in-situ observations and nudged simulations (allowing the capture of the elevated humidity transported within smoke plume) of aerosol extinction within BBA plume, clearly indicate the significant impact of RH on BBA optical  
820 properties. This underlines the importance of including in models the fire processes related to the



presence of humidity in the smoke plume over SAO. A second important aspect concerns the presence of possible errors in the actual parameterization used in ALADIN-Climate to calculate the evolution of BBA extinction with RH. In that sense, nudged simulations, associated with in-situ data obtained during ORACLES would certainly provide a unique opportunity to test and constrain the hydrophylic properties of BBA over SAO. Future work will extend significantly the number of cases studied to test the robustness of these first results.

## 5. Direct Radiative Forcing and impact of BBA

### 5.1. DRE exerted at TOA

The monthly-mean DRE exerted at TOA (in the visible spectral range) is indicated in the Figure 13 in clear-sky (left) and all-sky (right) conditions. The ALADIN-Climate estimates do not include possible SST adjustments due to BBA radiative effects, even if this could be important over SAO (Sakaeda et al., 2011). Figure 13 indicates an important regional gradient in the sign of DRE over the domain in all-sky conditions, with a rather negative forcing over the continent (net cooling) and positive (net heating) over SAO. Over the continent, the mean DRE is found to be mostly negative (~-5/-15 W.m<sup>-2</sup>) over Angola, with local maxima up to -20 W.m<sup>-2</sup>. An interesting result concerns the presence of significant positive forcings along the coast from Gabon to Namibia, with values of ~+10 to +20 W.m<sup>-2</sup>. Such significant positive forcing at TOA are correlated with both the presence of Sc along the coast of Angola, Republic of Congo and Gabon (see Figure S6 in Supplement material) and the high surface albedo over Namibia (Figure 1).

On the contrary, over SAO, DRE exerted at TOA is found to be mainly positive in all-sky conditions in agreement with a large literature focused on this region (Meyer et al., 2013; Feng and Christopher, 2015; De Graaf et al., 2012, 2014; Zuidema et al., 2016). The impact of the presence of Sc on the sign of DRE at TOA is clearly shown when comparing the ALADIN-Climate simulations in clear-sky and all-sky conditions. The large cooling effect at TOA is replaced by a significant heating over a large part of SAO. However and when averaged over the same region (20°S–10°N and 10°W–20°E) as defined in Feng and Christopher (2015), an important underestimate is detected compared to satellite observations. Indeed, the instantaneous (at satellite time overpassing) monthly-averaged DRE is found to be about +6 W.m<sup>-2</sup> in ALADIN-Climate and ~+35 W.m<sup>-2</sup> in the study of Feng and Christopher (2015). A better agreement is obtained with Oikawa et al. (2013), who reported an annual mean of +3 W.m<sup>-2</sup> over Southern Africa using CALIPSO and GCM simulation. More recently, Gordon et al. (2018) indicate a regional DRE of +11 W.m<sup>-2</sup> at TOA close to the one obtained in this study, but for five smoky days. We suspect the underestimate of LCF (Figure 2b) to be mostly responsible for this large difference with the Feng and Christopher (2015) estimates. Interestingly, comparisons with the climatological estimates based on MACC NRT data

855 for the period 2010-2015 (Figure S7 of Supplement material) indicate important differences. Figure S7 indicates that the positive **DRE** simulated by ALADIN-Climate is absent in the MACC NRT data, except locally over the continent.

Finally, the SMK\_SSA simulations indicate a significant changes in the monthly-mean (September 2016) **DRE** exerted at TOA, passing from a positive (+4.2 W.m<sup>-2</sup>) to negative (-0.54 W.m<sup>-2</sup>) **DRE** 860 (Figure 13 and Table 3) over the Box\_O. This means that the positive **DRE** at TOA could be weaker at the end of BBA season (late october). Figure 13 indicates a more intense negative **DRE** at TOA over smoke sources due to more scattering BBA and over the box\_S, the monthly-mean value (Table 3) is increasing from -3.9 W.m<sup>-2</sup> (SMK) to -7.3 W.m<sup>-2</sup> (SMK\_SSA). In addition, a weaker positive **DRE** is observed at TOA along the Southern African coast and Gabon due to more 865 scattering smoke.

## **5.2. Impact on the continental surface energy budget and dynamics**

The potential impact of BBA on the « continental climate » has been investigated by using the differences between the CTL and SMK simulations. Figure 14 shows the monthly-mean difference (September 2016) of the following variables; surface net SW radiations (upper left), 2-meter 870 temperature (T2m, upper right), sensible heat fluxes (SHF, bottom left) and the PBL height (bottom right). The potential effect of BBA on the continental precipitation is not studied as little or no precipitation occurs south of approximately 8°S during the austral winter season.

Smoke aerosols are responsible of an important dimming of about -30 to -50 W.m<sup>-2</sup> (monthly mean) over the continent and -10 to -40 W.m<sup>-2</sup> over SAO during September 2016, with the highest impact 875 logically located over smoke sources. Such estimates are consistent with those reported by Sakaeda et al. (2011) or Tummon et al. (2010). This impact of BBA results in an important decrease in the T2m over Congo, Angola and Zambia, as well as certain regions of Southern Africa. The impact is approximately ~-1 to -3 °C over the continent, in good agreement with the values reported by Sakaeda et al. (2011). When averaged over box\_S (5-15° S / 15-25° E), the impact of BBA on T2m 880 is about -1.7°C during September 2016 (Figure 15a). The daily-mean impact of BBA remains constant during this period, except for the end of September when the effect is negligible. For the 26 to 31 September, we hypothesize that compensations should occur, the « dynamical » effect of BBA being more important than the « dimming » effect. As mentioned previously and contrary to Sakaeda et al. (2011), the impact of BBA on SST is not quantified as the ALADIN-Climate 885 simulations have been performed with prescribed SST. Finally, it should be noted that the changes of the surface temperature per unit of AOD (averaged for all the period of simulation) is about ~ -2.5°. This value is found to be higher than the one (-1.5° per unit AOD) reported by Zhang et al. (2016) for an extreme biomass burning event occurring over Central Canada during June 2015. The

890 difference could be due to the absorbing properties of BBA, which are more pronounced in the present study compared to Zhang et al. (2016) with SSA of 0.94 (550 nm). This could favor higher dimming effect and related impact on the surface temperature over the Angola region.

In parallel with the changes in surface temperature, the sensible heat fluxes (SHF, Figure 14, bottom left) significantly decrease, meaning weaker fluxes, over almost the entire subcontinent, with maxima in the main biomass burning sources. The decrease is about -20 to -30  $\text{W.m}^{-2}$  over the 895 continent, with a mean value of -25  $\text{W.m}^{-2}$  when averaged over the box\_S (Figure 15b). The impact of smoke is important throughout the whole time period, with SHF changing from a (monthly-mean) value of 85  $\text{W.m}^{-2}$  to 60  $\text{W.m}^{-2}$  for the CTL and SMK runs, respectively. This is consistent with the findings of Sakaeda et al. (2011) or Tummon et al. (2010). Indeed, the latter report a decrease over almost the entire subcontinents, with a maximum (decrease of ~50%) in the main 900 smoke region. As a result of the significant decrease in T2m and SHF over much of the subcontinent, the PBL height (defined as the top of the PBL) also decreases in the SMK run. This decrease is significant over much of the subcontinent, in accordance with the results of Tummon et al. (2010), with regional maxima up to ~ -400 m (Figure 14, bottom right). The lowest changes are observed along the west coast between 10 and 15°S and small regions of increased PBL height 905 occur over southern Namibia and Northern Angola consistent with increases in the surface temperature. When averaged over the continent (box\_S), the decrease in the PBL height is also significant, changing from ~1200 to ~1000 m (monthly-averaged) for the CTL and SMK simulations (Figure 15c), respectively. The decrease is nearly constant during September 2016, except for 25 to 30 September (decrease), and not necessarily correlated with AOD. The difference 910 in the PBL height can reach a maximum decrease of 300 m (Figure 15c). In parallel, the impact of BBA on the near-surface (10m) wind speed (not shown) indicates a general decrease (of about -0.5  $\text{m.s}^{-1}$ ) over most of the continent. Over the ocean, the impact is more complex with the presence of a contrast, characterized by an increase (decrease) of the surface wind around 0-10°W/15-30°S (latitudes higher than 15°S).

915 Finally, the comparisons with the SMK\_SSA simulations (not shown, Figure S8) indicate a decrease of the surface radiative forcing both over continent and ocean. As reported in Table 3, the monthly-mean **DRE** at BOA is about -39  $\text{W.m}^{-2}$  and -25  $\text{W.m}^{-2}$  over Box\_S for the SMK and SMK\_SSA simulations, respectively (similar results are obtained over SAO). This is due to the decrease of SW radiations absorbed by smoke in the SMK\_SSA simulation, increasing the SW radiations reaching 920 the surface. This could be also due, to a lesser extent, to changes in the aerosol loading due to modifications of the dynamics and precipitations between the two simulations. This induces a less pronounced impact of BBA on the surface temperature and sensible heat fluxes in SMK\_SSA. The

increase of SW surface radiations, associated to lower absorption by BBA, decrease the impact on the PBL development (Figure S8). As mentioned previously, these results suggest that the impact of BBA on the surface fluxes and dynamics are certainly weaker at the end of the biomass burning season.

## 6. Conclusion

The transport, vertical structure, SW radiative heating, SW direct radiative **effect forcing** and climatic impact exerted by absorbing BBA in the SAO have been estimated for September 2016 using the ALADIN-Climate model in the context of the ORACLES and LASIC projects. The model is able to represent LWP and COD well, although with a large underestimate in LCF. The simulated BBA AOD is consistent over the continent ( $\sim 0.7$  at 550 nm) compared to MERRA2 or MODIS data and also locally against AERONET data. We have also used new recent retrievals of ACAOD (OMI or MODIS) to demonstrate the ability of the model to reproduce reasonable values of smoke concentrations above Sc during the transport over SAO.

The simulations indicate the transport of BBA over SAO mainly occurs between 2 and 4 km, consistent with aircraft lidar observations. There is some indication that the entrainment of BBA in the MBL could be underestimated by the model contrary to the recent literature (Zuidema et al., 2018). This possible bias could lead to underestimate the BBA indirect forcing in this ALADIN-Climate configuration. In parallel, the absorbing properties (SSA) of BBA are consistent over biomass-burning sources compared to AERONET but significantly higher when compared to Ascension Island (LASIC) surface observations. The significant difference could be due to the absence of internal mixing treatment in the model, a lack of representation of the long-range aging processes, and/or the absence of mixing of BBA in the MBL. In addition, the important SW absorption by BBA produces an additional SW heating of  $\sim 1$  °K/day.

The ALADIN-Climate simulations reveal a significant regional gradient in the sign of the SW **DRE** at TOA (all-sky conditions and fixed SST conditions), with mostly negative (continent) and positive (SAO) forcing, mainly due to changes in the underlying albedo associated with highly absorbing BBA. Over the continent, an intense monthly-mean positive forcing ( $+10/+15$  W.m<sup>-2</sup>) is simulated over the Gabon, part of the Congo and Angola, mainly due to the presence of low Sc. Over SAO, a **DRE** of  $+6$  W.m<sup>-2</sup> (20°S–10°N and 10°W–20°E) is simulated at TOA during all the ORACLES-1 period.

One of the main original results concerns the use of coincident in-situ observations and nudged simulations (allowing to capture the elevated humidity transported within smoke plume) of aerosol extinction within BBA plume. Results highlight the significant effect of enhanced moisture on BBA extinction that considerably reduces the negative bias (in the simulated extinction) in the nudged

simulation (SMK\_SN) with ORACLES-1 data, compared to the SMK (no nudging) run. A second important aspect concerns the possible errors in the actual parameterization used to estimate the changes in BBA extinction with RH in the model. Indeed, our results indicate a possible overestimate of the increase in smoke extinction due to RH when compared to ORACLES-1 observations. Nudged simulations, associated to in-situ observations, would certainly provide a unique data-set to test and constrain the hygroscopic properties of BBA over SAO. All of these points have possible implications for **DRE** considerations and future works will extend significantly the number of cases studied to test the robustness of the results.

For September 2016, the important negative surface dimming due to BBA (around -5 to -15 W.m<sup>-2</sup>) over the subcontinent significantly modifies the surface energy budget over much of Southern Africa. Indeed, the decrease in the net surface SW radiations is compensated by a decrease in sensible heat fluxes (-25 W.m<sup>-2</sup>, monthly mean) and surface land temperature (-1.5 °C) over Angola, Zambia and Congo notably. The association of the surface cooling and the lower tropospheric heating tends to decrease the continental PBL height over the continent by about ~200 m.

Finally, the indirect radiative effect exerted by BBA remains to be investigated with the ALADIN-Climate model using ORACLES/CLARIFY/AEROCLO-sA data in a manner similar to that presented here for **DRE** considerations. Once evaluated for all forcings over this region, long-term simulations (ERA-INT period) will be done to assess the possible feedbacks of BBA on Sc properties and the regional radiative budget at a climatic scale.

980

985

990

### **Author contribution**

PN and MM designed the ALADIN-Climat configuration and MM performed the simulations. RR, DSM, PN developed the ALADIN-Climat model. AMS, MS, KM, HJ, OT and CH have participated to provide the different aerosol satellite products. PZ provided in-situ LASIC (Ascension Island) observations; SS and SC data from the SSFR instrument. SB and RF participated to provide the HSRL-2 data. JR, RW, PS, PF and all co-authors have participated to the analysis of all observations and simulations. MM finalised the manuscript with contributions from all co-authors.

1000

1005

1010

1015

1020

1025

## Acknowledgment

This work has been conducted in the frame of the ORACLES investigation under the National Aeronautics and Space Administration's Earth Venture program. The LASIC extinction and SSA values are available through doi:10.5439/1369240, obtained from the Atmospheric Radiation Measurement (ARM) User Facility, a U.S. Department of Energy (DOE) Office of Science user facility managed by the Office of Biological and Environmental Research. PZ acknowledges support from DOE ASR grant DE-SC0018272. AERONET data are available from <https://aeronet.gsfc.nasa.gov>. MODIS data are available from the NASA Level 1 and Atmospheres Data Center at <https://ladsweb.modaps.eosdis.nasa.gov/>. MODIS above-cloud data products are available from the authors on request. MISR data are available from the NASA Atmospheric Sciences Data Center at <https://eosweb.larc.nasa.gov/>. M. Mallet, P. Nabat and P. Formenti acknowledge the AErosols, Radiation and CLOUDs in southern Africa (Aeroclo-sA) project, supported by the French National Research Agency under grant agreement n° ANR-15-CE01-0014-01, the French national programme LEFE/INSU, the Programme national de Télédetection Spatiale (PNTS, <http://www.insu.cnrs.fr/pnts>), grants n° PNTS-2016-02 and PNTS-2016-14, the French National Agency for Space Studies (CNES), the Centre National de la Recherche Scientifique (CNRS), the South African National Research Foundation (NRF) under grant UID 105958, and the European Union's 7th Framework Programme (FP7/2014-2018) under EUFAR2 contract n°312609. We thank the CNRM Climate model development team.

1050

1055

1060

1065

1070

## References

- Adebisi, A. A., Zuidema, P., and S. J. Abel, S.J.: The convolution of dynamics and moisture with the presence of shortwave absorbing aerosols over the southeast Atlantic. *J. Climate*, 28,1997–2024, doi: 10.1175/JCLI-D-14-00352.1, 2015
- 1075 Adebisi, A. and Zuidema, P.: The role of the southern African easterly jet in modifying the southeast Atlantic aerosol and cloud environments. *Q. J. R. Meteorol. Soc.*, 142 (April), p. 1574-1589 doi: [10.1002/qj.2765](https://doi.org/10.1002/qj.2765), 2016.
- 1080 Bellouin, N., J. Rae, A. Jones, C. Johnson, J. Haywood, and O. Boucher, O.: Aerosol forcing in the Climate Model Intercomparison Project (CMIP5) simulations by HadGEM2-ES and the role of ammonium nitrate, *J. Geophys. Res.*, 116,D20206, doi:10.1029/2011JD016074, 2011.
- 1085 Benedetti, A., Morcrette, J.-J., Boucher, O., Dethof, A., Engelen, R. J., Fisher, M., Flentjes, H., Huneeus, N., Jones, L., Kaiser, J. W., Kinne, S., Mangold, A., Razinger, M., Simmons, A. J., Suttie, M., and the GEMS-AER team: Aerosol analysis and forecast in the ECMWF Integrated Forecast System: 2. Data assimilation, *J. Geophys. Res.*, 114, D13205, <https://doi.org/10.1029/2008JD011115>, 2009.
- 1090 Benas, N., Finkensieper, S., Stengel, M., van Zadelhoff, G.-J., Hanschmann, T., Hollmann, R., and Meirink, J. F.: The MSG-SEVIRI-based cloud property data record CLAAS-2, *Earth Syst. Sci. Data*, 9, 415-434, <https://doi.org/10.5194/essd-9-415-2017>, 2017.
- 1095 Burton, S. P., Hair, J. W., Kahnert, M., Ferrare, R. A., Hostetler, C. A., Cook, A. L., Harper, D. B., Berkoff, T. A., Seaman, S. T., Collins, J. E., Fenn, M. A., and Rogers, R. R.: Observations of the spectral dependence of linear particle depolarization ratio of aerosols using NASA Langley airborne High Spectral Resolution Lidar, *Atmos. Chem. Phys.*, 15, 13453-13473, 10.5194/acp-15-13453-2015, 2015.
- 1100 Burton, S. P., Hostetler, C., Cook, A., Hair, J., Seaman, S. T., Scola, S. J., Harper, D., Smith, J. A., Fenn, M. A., Ferrare, R., Saide, P. E., Chemyakin, E., and Müller, D.: Calibration of a high spectral resolution lidar using a Michelson interferometer, with data examples from ORACLES, *Appl Opt*, 2018, in press.
- Chand, D., R. Wood, T. L. Anderson, S. K. Satheesh, and R. J. Charlson. "Satellite-Derived Direct Radiative Effect of Aerosols Dependent on Cloud Cover." *Nature Geoscience* 2, no. 3 (March 2009): 181–84. <https://doi.org/10.1038/ngeo437>, 2009.
- 1105 Cochrane, S., K., S. Schmidt, H. Chen, P. Pilewskie, S. Kittelman, W. Gore, J. Redemann, S. LeBlanc, K. Pistone, S. Platnick, K. Meyer, R. Ferrare, S. Burton, C. Hostetler, and Dobracki, A.: Aircraft-Derived Above-Cloud Aerosol Radiative Effects Based on ORACLES 2016 and ORACLES 2017 Experiments, in preparation in ACP, 2018.
- 1110 Cubison, M. J., Ortega, A. M., Hayes, P. L., Farmer, D. K., Day, D., Lechner, M. J., Brune, W. H., Apel, E., Diskin, G. S., Fisher, J. A., Fuelberg, H. E., Hecobian, A., Knapp, D. J., Mikoviny, T., Riemer, D., Sachse, G. W., Sessions, W., Weber, R. J., Weinheimer, A. J., Wisthaler, A., and Jimenez, J. L.: Effects of aging on organic aerosol from open biomass burning smoke in aircraft and laboratory studies, *Atmos. Chem. Phys.*, 11, 12049-12064, <https://doi.org/10.5194/acp-11-12049-2011>, 2011.
- 1115 Das, S., H. Harshvardhan, H. Bian, M. Chin, G. Curci, A. P. Protonotariou, T. Mielonen, K. Zhang, H. Wang, and Liu, X.: Biomass burning aerosol transport and vertical distribution over the South African-Atlantic region, *J. Geophys. Res. Atmos.*, 122, 6391–6415, doi:10.1002/2016JD026421, 2017.
- 1120 De Graaf, M., L. G. Tilstra, P. Wang, and Stammes, P.: Retrieval of the aerosol direct radiative effect over clouds from spaceborne spectrometry, *J. Geophys. Res.*, 117, D07207, doi:10.1029/2011JD017160, 2012.
- De Graaf, M., N. Bellouin, L. G. Tilstra, J. Haywood, and Stammes, P.: Aerosol direct radiative effect of smoke over clouds over the southeast Atlantic Ocean from 2006 to 2009, *Geophys. Res. Lett.*, 41, 7723–7730, doi:10.1002/2014GL061103, 2014.
- 1125 Diamond, M. S., Dobracki, A., Freitag, S., Small Griswold, J. D., Heikkila, A., Howell, S. G., Kacarab, M. E., Podolske, J. R., Saide, P. E., and Wood, R.: Time-dependent entrainment of smoke presents an observational challenge for assessing aerosol–cloud interactions over the southeast Atlantic Ocean, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2018-461>, in review, 2018.



- 1130 Dubovik, O., and King, M.D.: A flexible inversion algorithm for retrieval of aerosol optical properties from Sun and sky radiance measurements, *J. Geophys. Res.*, 105(D16), 20673–20696, doi: 10.1029/2000JD900282, 2000.
- Dubovik, O., A. Smirnov, B. N. Holben, M. D. King, Y. J. Kaufman, T. F. Eck, and Slutsker I.: Accuracy assessments of aerosol optical properties retrieved from Aerosol Robotic Network (AERONET) Sun and sky radiance measurements, *J. Geophys. Res.*, 105(D8), 9791–9806, doi: 10.1029/2000JD900040, 2000.
- 1135 Eck, T. F., et al. : A seasonal trend of single scattering albedo in southern African biomass-burning particles: Implications for satellite products and estimates of emissions for the world’s largest biomass-burning source, *J. Geophys. Res. Atmos.*, 118, 6414–6432, doi:10.1002/jgrd.50500, 2013.
- 1140 Fierce, L., T.C. Bond, S.E. Bauer, F. Mena, and Riemer, N.: Black carbon absorption at the global scale is affected by particle-scale diversity in composition. *Nature Commun.*, 7, 12361, doi:10.1038/ncomms12361, 2016.
- Feng, N. and Christopher, S.A. : Measurement-based estimates of direct radiative effects of absorbing aerosols above clouds, *J. Geophys. Res. Atmos.*, vol. 120, no. 14, pp. 6908–6921, 2015.
- 1145 Fierce, L. et al., Black carbon absorption at the global scale is affected by particle-scale diversity in composition. *Nat. Commun.* 7:12361 doi: 10.1038/ ncomms12361, 2016.
- 1150 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., and Reichle, R.: The modern-era retrospective analysis for research and applications, version 2 (MERRA-2), *J Climate*, 30, 5419-5454, 10.1175/JCLI-D-16-0758.1, 2017.
- Gordon, H., Field, P. R., Abel, S. J., Johnson, B. T., Dalvi, M., Grosvenor, D. P., Hill, A. A., Miltenberger, A. K., Yoshioka, M., and Carslaw, K. S.: Large simulated radiative effects of smoke in the south-east Atlantic, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2018-305>, in review, 2018.
- 1155 Hair, J. W., Hostetler, C. A., Cook, A. L., Harper, D. B., Ferrare, R. A., Mack, T. L., Welch, W., Izquierdo, L. R., and Hovis, F. E.: Airborne High Spectral Resolution Lidar for profiling aerosol optical properties, *Appl Optics*, 47, 6734-6752, 10.1364/AO.47.006734, 2008.
- 1160 Haywood, J., S. R. Osborne, P. N. Francis, A. Keil, P. Formenti, M. O. Andreae, and P. H. Kaye: The mean physical and optical properties of regional haze dominated by biomass burning aerosol measured from the C-130 aircraft during SAFARI 2000, *J. Geophys. Res.*, 108, 8473, doi:10.1029/2002JD002226, 2003.
- 1165 Heald, C.L., et al., **Contrasting the direct radiative effect and direct radiative forcing of aerosols**, *Atmos. Chem. Phys.*, 14, 5513-5527, 2014.
- Hodnebrog, O., Myhre, G., Forster, P. M., Sillmann, J., and Samset, B. H.: Local biomass burning is a dominant cause of the observed precipitation reduction in southern Africa, *Nat. Commun.*, 7, 11236, 2016.
- 1170 Jethva, H., Torres, O., and Ahn, C.: A 12-Year Long Global Record of Optical Depth of Absorbing Aerosols above the Clouds Derived from OMI/OMACA Algorithm, [11, 5837-5864](https://doi.org/10.5194/amt-11-5837-2018), <https://doi.org/10.5194/amt-11-5837-2018>, 2018.
- 1175 Johnson, B. T., Shine, K. P., & Forster, P. M.: The semi-direct aerosol effect: Impact of absorbing aerosols on marine stratocumulus. *Quarterly Journal of the Royal Meteorological Society*, 130, 1407–1422, 2004.
- Johnson, B. T., Haywood, J. M., Langridge, J. M., Darbyshire, E., Morgan, W. T., Szpek, K., Brooke, J. K., Marengo, F., Coe, H., Artaxo, P., Longo, K. M., Mulcahy, J. P., Mann, G. W., Dalvi, M., and Bellouin, N.: Evaluation of biomass burning aerosols in the HadGEM3 climate model with observations from the SAMBBA field campaign, *Atmos. Chem. Phys.*, 16, 14657-14685, <https://doi.org/10.5194/acp-16-14657-2016>, 2016.
- 1180 Kahn, R. A., and Gaitley, B.J.: An analysis of global aerosol type as retrieved by MISR, *J. Geophys. Res. Atmos.*, 120, 4248–4281, doi:10.1002/2015JD023322, 2015.
- 1185 Kar, J., Vaughan, M., Tackett, J., Liu, Z., Omar, A., Rodier, S., Trepte, C., and Lucker, P.: Swelling of transported smoke from savanna fires over the Southeast Atlantic Ocean, *Rem. Sen. Env.*, 211, 105-111, <https://doi.org/10.1016/j.rse.2018.03.043>, 2018.

- 1190 Labonne, M., Breon, F.-M., Chevallier, F.: Injection height of biomass burning aerosols as seen from a spaceborne lidar. *Geophysical Research Letters*, 34, L11806. <https://doi.org/10.10292007GL029311>, 2007.
- Mallet, M., Solmon, F., Roblou, L., Peers, F., Turquety, S., Waquet, F., Jethva, H, Torres, O.: Simulation of Optical Properties and Direct and Indirect Radiative Effects of Smoke Aerosols Over Marine Stratocumulus Clouds During Summer 2008 in California With the Regional Climate Model RegCM, *Journal of Geophys. Res.-Atmosphere*, 122, 10288-10313, 2017.
- 1195 Marticorena, B. and Bergametti, G.: Modeling the atmosphere dustcycle: 1. Design of a soil-derived dust emission scheme, *J. Geo-phys. Res.*, 100, 16415–16430, 1995.
- 1200 Martin, G. M., Johnson, D. W., and Spice, A.: The measurement and parameterization of effective radius of droplets in warm stratiform clouds, *J. Atmos. Sci.*, 51, 1823–1842, 1994.
- 1205 Masson, V., Le Moigne, P., Martin, E., Faroux, S., Alias, A., Alkama, R., Belamari, S., Barbu, A., Boone, A., Bouysse, F., Brousseau, P., Brun, E., Calvet, J.-C., Carrer, D., Decharme, B., Delire, C., Donier, S., Essaouini, K., Gibelin, A.-L., Giordani, H., Habets, F., Jidane, M., Kerdraon, G., Kourzeneva, E., Lafaysse, M., Lafont, S., Lebeaupin Brossier, C., Lemonsu, A., Mahfouf, J.-F., Marguinaud, P., Mokhtari, M., Morin, S., Pigeon, G., Salgado, R., Seity, Y., Taillefer, F., Tanguy, G., Tulet, P., Vincendon, B., Vionnet, V., and Voldoire, A.: The SURFEXv7.2 land and ocean surface platform for coupled or offline simulation of earth surface variables and fluxes, *Geosci. Model Dev.*, 6, 929-960, <https://doi.org/10.5194/gmd-6-929-2013>, 2013.
- 1210 Mayer, B. and Kylling, A.: Technical note: The libRadtran software package for radiative transfer calculations - description and examples of use, *Atmos. Chem. Phys.*, 5, 1855-1877, <https://doi.org/10.5194/acp-5-1855-2005>, 2005.
- 1215 Meyer, K., S. Platnick, L. Oreopoulos, and Lee D.: Estimating the direct radiative effect of absorbing aerosols overlying marine boundary layer clouds in the southeast Atlantic using MODIS and CALIOP, *J. Geophys. Res. Atmos.*, 118, 4801–4815, doi:10.1002/jgrd.50449, 2013.
- 1220 Meyer K., Platnick S., and Zhang Z.: Simultaneously inferring above-cloud absorbing aerosol optical thickness and underlying liquid phase cloud optical and microphysical properties using MODIS, *J Geophys Res. Atmos.*, 119, 5104–5114, doi:10.1002/2013JD021053, 2015.
- 1225 Mlawer, E.J., Taubman, S.J., Brown, P.D., Iacono, M.J. and Clough, S.A. : Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, *Journal of Geophysical Research* 102: doi: 10.1029/97JD00237. Issn: 0148-0227, 1997.
- Morcrette JJ : Description of the radiation scheme in the ecmwf model. Tech. rep, ECMWF, 1989.
- 1230 Michou, M., Nabat, P., and Saint-Martin, D.: Development and basic evaluation of a prognostic aerosol scheme (v1) in the CNRM Climate Model CNRM-CM6, *Geosci. Model Dev.*, 8, 501-531, [DOI:10.5194/gmd-8-501-2015](https://doi.org/10.5194/gmd-8-501-2015), 2015.
- Ogren, J.: Comment on “Calibration and Intercomparison of Filter-Based Measurements of Visible Light Absorption by Aerosols.”. *Aerosol Sci. Technol.*, 44: 589–591, 2010.
- 1235 Oikawa, E., T. Nakajima, T. Inoue, and Winker D.: A study of the shortwave direct aerosol forcing using ESSP/CALIPSO observation and GCM simulation, *J. Geophys. Res. Atmos.*, 118, 3687–3708, doi:10.1002/jgrd.50227, 2013.
- 1240 Nabat, P., Somot, S., Mallet, M., Sevault, F., Chiacchio, M., and Wild, M.: Direct and semi-direct aerosol radiative effect on the Mediterranean climate variability using a coupled regional climate system model, *Clim. Dynam.*, 44, 1127–1155, doi:10.1007/s00382-014-2205-6, 2015a.
- 1245 Nabat, P., Somot, S., Mallet, M., Michou, M., Sevault, F., Driouech, F., Meloni, D., di Sarra, A., Di Biagio, C., Formenti, P., Sicard, M., Léon, J.-F., and Bouin, M.-N.: Dust aerosol radiative effects during summer 2012 simulated with a coupled regional aerosol– atmosphere–ocean model over the Mediterranean, *Atmos. Chem. Phys.*, 15, 3303–3326, doi:10.5194/acp-15-3303-2015, 2015b.
- Nakajima, T., and King, M.D.: Determination of the Optical Thickness and Effective Particle Radius of Clouds from Reflected Solar Radiation Measurements. Part 1: Theory. *J. Atmos. Sci.*, 47, 1878-1893, 1990.

- 1250 Nam, C., Bony, S., Dufresne, J.-L., and Chepfer, H.: The 'too few, too bright' tropical low-cloud problem in CMIP5 models. *Geophys. Res. Letters*, 39 (21), 2012.
- 1255 Petrenko, M., Kahn, R., Chin, M., and Limbacher, J.: Refined use of satellite aerosol optical depth snapshots to constrain biomass burning emissions in the GOCART model. *Journal of Geophysical Research: Atmospheres*, 122, 10,983–11,004. <https://doi.org/10.1002/2017JD026693>, 2017.
- 1260 Radu, R., M. Déqué, and S. Somot, S.: Spectral nudging in a spectral regional climate model. *Tellus*, 60A, 898–910, doi:10.1111/j.1600-0870.2008.00341.x, 2008.
- Reid, J. S., Eck, T. F., Christopher, S. A., Koppmann, R., Dubovik, O., Eleuterio, D. P., Holben, B. N., Reid, E. A., and Zhang, J.: A review of biomass burning emissions part III: intensive optical properties of biomass burning particles, *Atmos. Chem. Phys.*, 5, 827-849, <https://doi.org/10.5194/acp-5-827-2005>, 2005.
- 1265 Roebeling, R.A., A.J. Feijt and Stammes, P.: Cloud property retrievals for climate monitoring: implications of differences between SEVIRI on METEOSAT-8 and AVHRR on NOAA-17, *J. Geophys. Res.*, 111, D20210, doi:10.1029/2005JD006990, 2006.
- 1270 Sakaeda, N., R. Wood, and Rasch, P.J.: Direct and semidirect aerosol effects of southern African biomass burning aerosol, *J. Geophys. Res.*, 116, D12205, doi:10.1029/2010JD015540, 2011.
- Sayer, A. M., L. A. Munchak, N. C. Hsu, R. C. Levy, C. Bettenhausen, and Jeong, M.-J.: MODIS Collection 6 aerosol products: Comparison between Aqua's e-Deep Blue, Dark Target, and "merged" data sets, and usage recommendations, *J. Geophys. Res. Atmos.*, 119, 13,965–13,989, doi: 10.1002/2014JD022453, 2014.
- 1275 Sayer A. M., N. C. Hsu, C. Bettenhausen, J. Lee, J. Redemann, B. Schmid, and Shinozuka, Y.: Extending "Deep Blue" aerosol retrieval coverage to cases of absorbing aerosols above clouds: Sensitivity analysis and first case studies, *J. Geophys. Res. Atmos.*, 121, 4830–4854, doi: 10.1002/2015JD024729, 2016.
- 1280 Sayer, A. M., Hsu, N. C., Lee, J., Kim, W. V., Burton, S., Fenn, M. A., Ferrare, R. A., Kacenelenbogen, M., LeBlanc, S., Pistone, K., Redemann, J., Segal-Rozenhaimer, M., Shinozuka, Y., and Tsay, S.-C.: Two decades observing smoke above clouds in the south-eastern Atlantic Ocean: Deep Blue algorithm updates and validation with ORACLES field campaign data, *Atmos. Meas. Tech. Discuss.*, <https://doi.org/10.5194/amt-2019-58>, in review, 2019.
- 1285 Schulz, M., de Leeuw, G., and Balkanski, Y.: Sea-salt aerosol source functions and emissions, in: *Emission of Atmospheric Trace Compounds*, edited by: Granier, C., Artaxo, P., and Reeves, C. E., Kluwer Acad., Norwell, Mass., Springer Netherlands, 333-359, 2004.
- 1290 Seethala, C., Meirink, J. F., Horváth, Á., Bennartz, R., and Roebeling, R.: Evaluating the diurnal cycle of South Atlantic stratocumulus clouds as observed by MSG SEVIRI, *Atmos. Chem. Phys.*, 18, 13283-13304, <https://doi.org/10.5194/acp-18-13283-2018>, 2018.
- 1295 Shinozuka, Y., P. E. Saide, S. P. Burton, P. Zuidema, G. A. Ferrada, M. Mallet, Y. Zhang, R. Leung, L. Pfister, J.-M. Ryoo, S. G. Howell, S. Freitag, A. Dobracki, J. R. Podolske, S. LeBlanc, R. Ferrare, D. Henze, D. Noone, M. Kacarab, A. Nenes, E. J. Stith, J. R. Bennett, G. R. Carmichael, S. J. Doherty, R. Wood, and Redemann, J.: Global and regional modeling of absorbing aerosols in the southeast Atlantic during ORACLES 2016: comparisons to new observations, in preparation in ACPD, 2019.
- 1300 Shipley, S. T., Tracy, D. H., Eloranta, E. W., Trauger, J. T., Sroga, J. T., Roesler, F. L., and Weinman, J. A.: High Spectral Resolution Lidar to Measure Optical-Scattering Properties of Atmospheric Aerosols .1. Theory and Instrumentation, *Appl Optics*, 22, 3716-3724, 10.1364/AO.22.003716, 1983.
- 1305 Slingo, A. and Schrecker, H.M. : On the shortwave radiative properties of stratiform water clouds. *Q. J. R. Meteorol. Soc.*,108: 407-426, 1982.
- Stephens, G. : Radiation profiles in extended water clouds, II : Parametrization schemes, *J. Atmos. Sci.*, 35, 2123-2132, 1978.

- 1310 Stengel, M., Kniffka, A., Meirink, J. F., Lockhoff, M., Tan, J., and Hollmann, R.: CLAAS: the CM SAF cloud property data set using SEVIRI, *Atmos. Chem. Phys.*, 14, 4297-4311, <https://doi.org/10.5194/acp-14-4297-2014>, 2014.
- 1315 Stier, P., Schutgens, N. A. J., Bellouin, N., Bian, H., Boucher, O., Chin, M., Ghan, S., Huneeus, N., Kinne, S., Lin, G., Ma, X., Myhre, G., Penner, J. E., Randles, C. A., Samset, B., Schulz, M., Takemura, T., Yu, F., Yu, H., and Zhou, C.: Host model uncertainties in aerosol radiative forcing estimates: results from the AeroCom Prescribed intercomparison study, *Atmos. Chem. Phys.*, 13, 3245-3270, <https://doi.org/10.5194/acp-13-3245-2013>, 2013.
- 1320 Sun, Q., Wang, Z., Li, Z., Erb, A., & Schaaf, C. B.: Evaluation of the Global MODIS 30 Arc-Second Spatially and Temporally Complete Snow-Free Land Surface Albedo and Reflectance Anisotropy Dataset. *International Journal of Applied Earth Observation and Geoinformation*, 58, 36–49, 2017.
- Tummon, F., F. Solmon, C. Liousse, and Tadross, M. : Simulation of the direct and semidirect aerosol effects on the southern Africa regional climate during the biomass burning season, *J. Geophys. Res.*, 115, D19206, doi:10.1029/2009JD013738, 2010.
- 1325 Vakkari, V., Beukes, J.P., Dal Maso, M., Aurela, M., Josipovic, M., and Van Zyl, P.G.: Major secondary aerosol formation in southern African open biomass burning plumes, *Nature Geoscience*, 1-4, 2018.
- 1330 van Marle, M. J. E., Kloster, S., Magi, B. I., Marlon, J. R., Daniau, A.-L., Field, R. D., Arneth, A., Forrest, M., Hantson, S., Kehrwald, N. M., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Yue, C., Kaiser, J. W., and van der Werf, G. R.: Historic global biomass burning emissions for CMIP6 (BB4CMIP) based on merging satellite observations with proxies and fire models (1750–2015), *Geosci. Model Dev.*, 10, 3329-3357, <https://doi.org/10.5194/gmd-10-3329-2017>, 2017.
- 1335 Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 $\lambda$  PSAP), *Aerosol Science and Technology*, 44:8, 706-712, DOI: [10.1080/02786826.2010.482110](https://doi.org/10.1080/02786826.2010.482110), 2010.
- Xu, H., Guo, J. P., Ceamanos, X., Roujean, J. L., Min, M., Carrer, D.: On the influence of the diurnal variations of aerosol content to estimate direct aerosol radiative forcing using MODIS data. *Atmospheric Environment*, 141, 186–196. <https://doi.org/10.1016/j.atmosenv.2016.06.067>, 2016.
- 1340 Wilcox, E. M.: Stratocumulus cloud thickening beneath layers of absorbing smoke aerosol, *Atmos. Chem. Phys.*, 10, 11769-11777, <https://doi.org/10.5194/acp-10-11769-2010>, 2010.
- 1345 Zhang, J., Reid, J. S., Christensen, M., and Benedetti, A.: An evaluation of the impact of aerosol particles on weather forecasts from a biomass burning aerosol event over the Midwestern United States: observational-based analysis of surface temperature, *Atmos. Chem. Phys.*, 16, 6475-6494, <https://doi.org/10.5194/acp-16-6475-2016>, 2016.
- Zuidema P, et al.: Smoke and clouds above the southeast Atlantic: Upcoming field campaigns probe absorbing aerosols impact on climate. *Bull Am Meteorol Soc* 97:1131 – 1135, 2016.
- 1350 Zuidema, P., A. Sedlacek, C. Flynn, S. Springston, R. Delgado, J. Zhang, A. Aiken and Muradyan, P. The Ascension Island boundary layer in the remote southeast Atlantic is often smoky. *Geophys. Res. Lett.*, **45**, pp. 4456-4465 doi:[10.1002/2017GL076926](https://doi.org/10.1002/2017GL076926), 2018.

1355

1360

1365

Aerosol Species	$r_0$	$\sigma$	Density	n / k	MEE	SSA	ASY
<i>Fresh smoke</i>	0.10	1.30	1.350	1.50 / 0.03	4.05	0.84	0.51
<i>Aged smoke</i>	0.12	1.30	1.350	1.50 / 0.03	5.05	0.90	0.58

*Note. Here  $r_0$  and  $\sigma$  are the median radius (in  $\mu\text{m}$ ) and geometric standard deviation of the lognormal distribution. Mass density is reported in  $\text{g}\cdot\text{cm}^{-3}$ ,  $m$  is the complex refractive index, and MEE and SSA the mass extinction efficiency ( $\text{m}^2\text{g}^{-1}$ ) and single scattering albedo in dry state and reported at 550 nm.*

1370 **Table 1.** Parameters describing aerosol components used in the ALADIN-Climate model for the two smoke tracers and the resulting optical properties.

1375

1380

1385

1390

1395

	<b>Aerosols</b>	<b>Clouds</b>	<b>Spatial / Temporal Resolutions</b>	<b>Available Period</b>
MOD06ACAERO	ACAOD	#	0.1 × 0.1 ° / daily	June-Oct. 2000(T)/03(A) - 2019
MODIS DB	ACAOD	#	0.5 × 0.5 °/daily	2000(T)/02(A) - 2017
MODIS DT/DB	AOD	LWP, COD	1 × 1 ° / daily	2000 - 2019
OMI	ACAOD	#	0.5 × 0.5 ° / daily	2004 - 2019
MISR	AOD	#	0.5 × 0.5 ° / monthly	2000 - 2017
SEVIRI	#	LWP, COD	0.5 × 0.5° / daily	2004 - 2015
ERA-Interim	#	LWP, LCF	0.7 × 0.7 ° / daily	1979 - 2018
MERRA2	AOD by species	#	0.5 x 0.625 ° / hourly	1980 - 2018
CAMS	AOD by species	#	0.5 x 0.625 ° / hourly	2008 - 2015
MACC	DRE TOA (all-sky)	#	1.125 × 1.125 ° / daily	2003 - 2011

1400

**Table 2.** Satellite (T:Terra, A:Aqua) and reanalysis data used in this study to analyse aerosols and Sc clouds microphysical/optical properties.

1405

1410

1415

1420

	DRE_BOA		DRE_TOA		SW Heating (3 km)	
	Box_S	Box_O	Box_S	Box_O	Box_S	Box_O
ALD_SMK	-38.93	-15.73	-3.92	+4.22	1.15	0.75
ALD_SMK_SSA	-24.81	-8.17	-7.37	-0.54	0.58	0.30

1425 **Table 3.** Monthly mean (September 2016) all-sky direct radiative **effect forcing** at TOA and BOA ( $\text{W.m}^{-2}$ ) and SW heating rate at 3 km ( $^{\circ}\text{K/day}$ ) for the two boxes (Box\_S and Box\_O), obtained from the SMK and SMK\_SSA simulations.

1430


1435

1440

1445

1450

1455



1460

1465

1470

1475 **Figure 1.** Domain defined for the ALADIN-Climate model simulations (here, the surface albedo is represented). The two different boxes (box\_O and box\_S) and the Ascension Island are indicated. Red stars represent the localisation of different profiles studied in the Part 4.2.3.2.

1480

1485

1490



1495

1500


1505

1510


1515

1520

1525




1525 **Figure 2.** Daily-mean Sc properties (LWP, LCF and COD) simulated by ALADIN-Climate and from ERA-INT and SEVERI data. Values have been averaged over the box\_O.



**Figure 3.** Total monthly-averaged Total AOD (at 550 nm) for september 2016 simulated by ALADIN-Climate (all times used), CAMS, MERRA2 and derived from the MODIS Terra, Aqua 1530 (combined DB/DT products) and MISR sensors.

1535






**Figure 4.** Total daily-mean AOD (at 550 nm) averaged over the box\_S (5-15S / 15-25E) from the ALADIN-Climate model (10:30 and 13:30 UTC outputs are used), MERRA2 and derived from 1540 MODIS-Aqua DT data (left). Daily-mean ACAOD (at 550 nm) estimated from the MODIS-DB satellite and two different configurations of the ALADIN-Climate model (SMK and SMK\_SN) averaged over the box\_O (0-10N / 10-20E) (right). Uncertainties related to DT AOD and DB ACAOD are also indicated.

1545

1550


1555



1560 **Figure 5.** Comparisons of daily-mean total AOD obtained at the Lubango, Mongu Inn and Ascension Island AERONET stations (daytime model outputs only) for September 2016.

1565


1570



**Figure 6** Total monthly-averaged ACAOD (at 550 nm) for September 2016 simulated by the 1575 ALADIN-Climate (10:30 and 13:30 UTC outputs only are used) model and derived from MODIS (Deep-Blue and Meyer retrievals) and OMI instruments.

1580


1585



**Figure 7.** Monthly averaged vertical profiles of the total aerosol (left) and smoke (right) extinction coefficient (at 550 nm) simulated by the ALADIN-Climate model for two transects at latitudes of 8 and 15°S. Longitudes are between 30°E to 11.5°W.


1590

1595



**Figure 8.** Vertical profiles of aerosol extinction coefficient (in  $\text{km}^{-1}$ ) at three different wavelengths from ALADIN-Climate and HSRL-2 instrument (red: 1064 nm / purple: 550 nm / blue: 440 nm), associated with the mean values (note that the wavelengths are not exactly similar, especially in ultraviolet). Also reported are the vertical profiles of RH simulated by ALADIN-Climate (dotted black) and MERRA2 (black), as well as the ALADIN-Climate cloud fraction (dotted yellow).

1635



1645

**Figure 9.** Daily-mean in-situ SSA estimated at the surface at Ascension Island (LASIC) and simulated with the ALADIN-Climat model at two different altitudes (0.2 and 3 km).


1650

1655

1660

1665





1670

1675

1680

1685


1695

1700

**Figure 10.** Monthly-mean (September 2016) aerosol SW heating rate vertical profiles simulated by the ALADIN-Climate model for two transects at latitudes of 8 (top) and 15°S (bottom). Longitudes are between 30°E to 11.5°W.

1705

1710



1715



**Figure 11.** Instantaneous SW heating rate (12:00 UTC) only due to smoke aerosols, obtained from ORACLES aircraft data and simulated by the ALADIN-Climate model in clear-sky (red dashed) and all-sky (blue dashed) conditions.

1720

1725



1730

1735





1740

1745



1750

1755




1760

1765 **Figure 12.** Vertical profiles of aerosol extinction coefficient (at 550 nm) obtained from ALADIN-Climate for the SMK and SMK\_SN simulations, and derived from the HSRL-2 instrument for the 24/09, associated to the vertical mean (left panels). Also reported the RH obtained from the same simulations and MERRA2 data (right panels).

1770


ALADIN-Climat Monthly-mean DRF (SW) at TOA - September 2016

1775



1780

1785




1790

**Figure 13.** Monthly-mean SW DRE ( $W.m^{-2}$ ) exerted by smoke particles at TOA for the September 2016 period in clear-sky (left up) and all-sky (right up) conditions for the SMK simulation and for the (bottom) SMK\_SSA run.


1795

1800

*Direct Radiative Forcing at the surface ( $W.m^{-2}$ )*




*Surface Temperature over land (K)*




1805

*Sensible Heat Fluxes ( $W.m^{-2}$ )*



*PBL altitude (m)*



1810


1815

**Figure 14.** Differences between the CTL and SMK ALADIN-Climate runs in the monthly-mean (September 2016) SW surface radiations (top left), 2 meter continental temperature (top right), sensible heat fluxes (bottom left) and PBL height (bottom right).


1820

1825

1830




1835



1840

1845



1850

1855

**Figure 15.** Daily-mean Surface Temperature, Sensible Heat Fluxes and PBL height obtained from the CTL (black lines) and SMK (dashed black) ALADIN-Climate simulations for september 2016. The AOD (red dashed lines) and the difference between the two simulations (CTL and SMK) for each variables are also reported (blue lines).

1860