

1 **Response to referee comments on “An evaluation of the OMI satellite instrument’s ability to**
2 **observe boundary layer ozone pollution across China: application to 2005-2017 ozone trends”**

3
4 We thank the referees for their careful reading of the manuscript and the valuable comments. This
5 document is organized as follows: the Referee’s comments are in *italic*, our responses are in plain
6 text, and all the revisions in the manuscript are shown **in blue**. The line numbers in this document
7 refer to the updated manuscript.

8

9 **Referee 1**

10
11 *The authors have conducted a thorough revision of their manuscript and I appreciate their efforts*
12 *to evaluate OMI against IAGOS profiles, additional ozonesondes and long-term surface*
13 *observations. The results are much more defensible, especially since the authors now focus on*
14 *southern China and have clarified that their analysis is representative of the boundary layer rather*
15 *than the surface. The new Figure 4 is also very helpful as it shows that ozone variability in the*
16 *boundary layer is greater than in the mid-troposphere (in southern China), which can help to*
17 *distinguish boundary layer pollution episodes. I still have a few concerns, as described below, and I*
18 *could recommend the paper for publication if the authors can address these concerns.*

19 **Response.** We thank the reviewer for this detailed feedback, which has helped improved our work.

20
21 *The paper has been revised to address the concerns of the referees that OMI does not detect*
22 *surface ozone but rather has some sensitivity to ozone in the deeper boundary layer, provided that*
23 *the enhancements are relatively strong compared to the mid-troposphere. However, this revised*
24 *focus is incomplete as the paper still states or implies, in many places throughout the manuscript,*
25 *that OMI is detecting surface ozone. The most important statement in the revision is this sentence at*
26 *the beginning of section 4:*

27 *“The correlation of OMI with the MEE surface ozone data likely does not reflect a direct sensitivity*
28 *of OMI to surface ozone, which is very weak, but rather a sensitivity to boundary layer ozone*
29 *extending up to a certain depth and correlated with surface ozone.”*

30 *The reader should not have to wait until the middle of the paper to read this very important*
31 *statement. It needs to appear in the Introduction and then the authors need to thoroughly revise*
32 *their paper to make sure that the discussion and the conclusions are consistent with this statement.*

33 **Response.** Thanks for pointing this out. We have revised a lot of sentences to make sure the
34 statements are consistent. We also have this sentence in the introduction.

35 **P2L22.** Even if sensitivity to the boundary layer is low, the enhancements can be sufficiently large
36 to enable detection.

37
38 *For example, the authors changed the title of the paper so that it reflects the analysis in the revised*
39 *manuscript, but it’s still not quite right. In their response to the comments of all three referees the*
40 *authors clarified that the OMI retrievals are reflective of the boundary layer rather than the*
41 *surface. But the title gives the impression that OMI has the ability to observe surface ozone*
42 *pollution. At a minimum the title needs to be revised, with “surface” replaced with “boundary*
43 *layer”. But even then the reader has the impression that the product can be used for all of China,*
44 *whereas the product is not effective in northern China. A better title would be something like:*
45 *“An evaluation of the OMI satellite instrument’s ability to observe boundary layer ozone pollution*
46 *across China: application to 2005-2017 ozone trends”*

47 **Response.** We have revised the title.

48 **New title.** An evaluation of the OMI satellite instrument’s ability to observe boundary layer ozone
49 pollution across China: application to 2005-2017 ozone trends

50
51 *Other statements in the paper that are not consistent with the first sentence at the beginning of*
52 *Section 4 are:*

53 *1) First sentence of the abstract: “Nadir-viewing satellite observations of tropospheric ozone in the*
54 *UV have been shown to detect surface ozone pollution episodes but no quantitative evaluation of*

55 *this ability has been done so far.” This isn’t entirely accurate. A better summary statement would*
56 *be “...have been shown to have some sensitivity to surface or near-surface ozone pollution*
57 *episodes”*

58 **Response.** We have revised accordingly.

59
60 *2) Second sentence of the abstract: “Here we use 2013-2017 surface ozone data from the new*
61 *China Ministry of Ecology and Environment (MEE) network of ~1000 sites, together with vertical*
62 *profiles from ozonesondes and aircraft, to quantify the ability of OMI tropospheric ozone retrievals*
63 *to characterize surface ozone pollution in China.”*

64 *This also gives the impression that OMI can detect surface ozone, when you should be talking*
65 *about the correlation between OMI and surface ozone. A better statement would be:*

66 *“...to quantify the correlation between OMI tropospheric ozone retrievals and surface ozone*
67 *pollution events in China.”*

68 **Response.** We have revised the sentence.

69
70 *3) Third sentence of the abstract:*

71 *“After subtracting the Pacific background, the 2013-2017 mean OMI ozone enhancements over*
72 *eastern China can quantify the spatial distribution of mean summer afternoon surface ozone with a*
73 *precision of 8.4 ppb and a spatial correlation coefficient $R=0.73$.” This sentence would be more*
74 *accurate as:*

75 *“After subtracting the Pacific background, the correlation between observed surface ozone and the*
76 *2013-2017 mean OMI ozone enhancements over eastern China can be used to estimate the spatial*
77 *distribution of mean summer afternoon surface ozone with a precision of 8.4 ppb and a spatial*
78 *correlation coefficient $R=0.73$.”*

79 **Response.** We have revised the sentence.

80
81 *4) Fifth sentence of the abstract:*

82 *“OMI is much more successful at capturing the day-to-day variability of surface ozone at sites in*
83 *southern China $<34^{\circ}N$ ($R = 0.3-0.6$) than in northern China ($R = 0.1-0.3$) because of weaker*
84 *retrieval sensitivity and larger upper tropospheric variability in the north.”*

85 *Would be better as:*

86 *“In terms of day-to-day variability, OMI has a higher correlation with observed surface ozone at*
87 *sites in southern China $<34^{\circ}N$ ($R = 0.3-0.6$) than in northern China ($R = 0.1-0.3$) because of*
88 *weaker retrieval sensitivity and larger upper tropospheric variability in the north.”*

89 **Response.** We have revised the sentence.

90
91 *There are many more statements like this throughout the paper that need to be revised, following*
92 *the examples above.*

93
94 *In Section 2 the following important statement appears:*

95 *“We focus on summer when ozone pollution in China is most severe and when OMI has the*
96 *strongest sensitivity”*

97 *A similar statement needs to appear in the abstract and introduction so that the reader is aware*
98 *that the evaluation has so far only focused on a region and time of year when boundary layer ozone*
99 *is very high.*

100 **Response.** Now we say this in the abstract.

101 P1L16. We focus on summer when ozone pollution in China is most severe and when OMI has the
 102 strongest sensitivity.

103
 104 *In Section 3 you provide the following equation and state that “With such a precision, OMI can
 105 provide useful information on mean summer afternoon levels of surface ozone in polluted regions.”:*

106
 107 $[O_3] = 6.9 \Delta\Omega + 24.6 \pm 8.4$

108
 109 *This presentation (and in the abstract as well) gives the reader the impression that they can take
 110 this equation and an OMI retrieval above a given location on a given day, and estimate the
 111 afternoon ozone on that day. For example, an OMI enhancement of 5 DU would indicate a surface
 112 ozone mixing ratio of 59.1 ± 8.4 ppbv, or a range of 50.7 to 67.5 ppbv. This sounds quite good, but
 113 when I look at Figure 1e, I can see that the full range of surface ozone values for an enhancement
 114 of 5 DU is 35-80 ppbv. And of course, this equation is not derived for estimating ozone on a given
 115 day, it is derived from data averaged over 5 years. Therefore the text needs to clearly state that the
 116 equation can only provide an estimate of multi-year average ozone. Accordingly, the text in the
 117 abstract also needs to be modified to more accurately represent the limitations of the equation. The
 118 abstract should provide a statement similar to:*

119
 120 *“After subtracting the Pacific background, the correlation between observed surface ozone and the
 121 2013-2017 mean OMI ozone enhancements over eastern China can be used to estimate the broad
 122 multi-year spatial distribution of mean summer afternoon surface ozone with a precision of 8 ppb
 123 and a spatial correlation coefficient $R=0.73$.” Note that I emphasized “multi-year”, and changed
 124 8.4 to 8 (8.4 is far too precise for such a rough estimate).*

125 **Response.** Thanks. We have revised it in the abstract and also in these places.

126 P4L27. From there one can estimate **multi-year average** surface ozone (ppb) on the basis of the
 127 observed OMI enhancement (DU) as...

128 P9L17. After subtracting the contribution from the North Pacific background, we find that the OMI
 129 enhancement over eastern China can reproduce the observed spatial distribution of **multi-year**
 130 mean summer afternoon ozone concentrations at the MEE sites, with a correlation coefficient $R =$
 131 **0.73** and a precision of 8 ppb.

132
 133 *Section 4*

134 *Here you state: “The Hong Kong ozonesonde data thus indicate that OMI can quantify the
 135 frequency of high-ozone episodes in the boundary layer even if it may not be reliable for individual
 136 events.”*

137 *This is a good summary of OMI’s capability, and it’s this type of statement that your paper should
 138 focus on, rather than emphasizing the equation in Section 3.*

139 **Response.** We have it in the conclusion now.

140 P9L27. The Hong Kong ozonesonde data also indicates that OMI can quantify the frequency of
 141 high-ozone episodes in the boundary layer even if it may not be reliable for individual events.

142
 143 *In the last paragraph of the Introduction you state:
 144 “However, no quantitative comparison of the satellite data to surface observations has so far been
 145 done.” The implication of this statement is that no one has ever bothered to compare satellite
 146 retrievals to surface ozone, when it could be done very easily. But the reason this has not been done*

147 *before is because the available tropospheric column ozone products don't have a true sensitivity to*
148 *surface ozone. A better statement would be:*

149 *"While the available tropospheric column ozone products don't have a true sensitivity to surface*
150 *ozone, here we demonstrate that the strong ozone enhancements in the summertime boundary layer*
151 *of southern China can be used to derive a relationship between OMI tropospheric ozone and*
152 *observed surface ozone "*

153 *Also, the above statement discounts the important advances made by the scientists who are*
154 *combining IASI and GOME-2 to produce retrievals of tropospheric ozone between the surface and*
155 *3 km. Figure 8 and Figure 9 of Gaudel et al. [2018] show tropospheric column ozone between the*
156 *surface and 3 km above China and Europe, based on retrievals from the IASI and GOME-2*
157 *instruments. These retrievals compared well to IAGOS ozone profiles above China (see Figure S8*
158 *in the supplement to Gaudel et al. 2018). (Please also see Dufour et al [2018] who look at ozone*
159 *across China from the surface to 6 km using IASI). These papers should be referenced in the*
160 *Introduction.*

161 **Response.** We have added these two references. About the ability of OMI to detect surface ozone
162 pollution, we think it is better to briefly mention it in the introduction and then give more details in
163 later discussion. Now we say this in the introduction.

164 P2L23. Even if sensitivity to the boundary layer is low, the enhancements can be sufficiently large
165 to enable detection.

166
167

168 *The following sentence appears in the Introduction:*

169 *"Retrieval of tropospheric ozone (only ~10% of the column) from these instruments has mostly*
170 *been done in the past by subtracting independent satellite measurements of stratospheric ozone*
171 *(Fishman et al., 1987; Ziemke et al., 2011)."*

172 *Another common method that should be mentioned is the cloud-slicing technique. This technique*
173 *was recently used in an excellent paper by Ziemke et al. [2019] who show 40 years of ozone*
174 *increases above southern Asia from the TOMS and OMI instruments.*

175 **Response.** Now we say.

176 P2L14. Retrieval of tropospheric ozone (only ~10% of the column) from these instruments has
177 mostly been done in the past by subtracting independent satellite measurements of stratospheric
178 ozone (Fishman et al., 1987; Ziemke et al., 2011) or using the convective cloud differential method
179 (Ziemke et al., 1998, 2019).

180
181 *The IASI+GOME2 results for China [Gaudel et al., 2018] show that high ozone in spring and*
182 *summer can extend over the North Pacific Ocean. Why do you not show OMI data above the ocean*
183 *in Figure 6 and Figure S5? However, you do show ozone above Taiwan in Figure 6 and compare*
184 *OMI to the TOAR observations from Taiwan, even though you don't use Taiwanese surface*
185 *observations in your equation 8.*

186 **Response.** Thanks. Now we say this in text.

187 P8L22. Here we have extended the trend analysis to Taiwan because of the opportunity to compare
188 to surface records.

189

190 *I don't understand Figure 5b. Figure 5a shows some clear differences between the observed and*
191 *predicted high ozone episodes for a threshold of 82 ppbv. Yet Figure 5b with higher thresholds*
192 *shows nearly perfect agreement between the observed and predicted. Is this for all sites averaged*

193 *together? What is the scatter for the individual sites?*

194 **Response.** Now we make it clear the Figure 5b uses all data in eastern China south of 34°N. We
195 also reported the spatial correlation of ozone episode days in surface sites and from OMI.

196 P8L15. However, we find that the model can also accurately estimate the probability of exceedance
197 above higher thresholds (Figure 5b) **for the ensemble of eastern China sites south of 34°N**, which
198 confirms the property of threshold invariance of an extreme value model (Cole, 2001).

199 P8L12. The results show that the predicted fraction of ozone episodes resembles that observed, with
200 a spatial correlation of 0.62 (Figure 5a).

201
202 Dufour, G., Eremenko, M., Beekmann, M., Cuesta, J., Foret, G., Fortems-Cheiney, A., Lachâtre, M.,
203 Lin, W., Liu, Y., Xu, X., and Zhang, Y.: Lower tropospheric ozone over the North China Plain:
204 variability and trends revealed by IASI satellite observations for 2008–2016, *Atmos. Chem. Phys.*,
205 18, 16439-16459, <https://doi.org/10.5194/acp-18-16439-2018>, 2018.

206
207 Gaudel, A., et al. (2018), Tropospheric Ozone Assessment Report: Present-day distribution and
208 trends of tropospheric ozone relevant to climate and global atmospheric chemistry model evaluation,
209 *Elem. Sci. Anth.*, 6(1):39, DOI: <https://doi.org/10.1525/elementa.291>

210
211 Ziemke, J. R., Oman, L. D., Strode, S. A., Douglass, A. R., Olsen, M. A., McPeters, R. D., Bhartia,
212 P. K., Froidevaux, L., Labow, G. J., Witte, J. C., Thompson, A. M., Haffner, D. P., Kramarova, N.
213 A., Frith, S. M., Huang, L.-K., Jaross, G. R., Seftor, C. J., Deland, M. T., and Taylor, S. L.: Trends
214 in global tropospheric ozone inferred from a composite record of TOMS/OMI/MLS/OMPS satellite
215 measurements and the MERRA-2 GMI simulation, *Atmos. Chem. Phys.*, 19, 3257-3269,
216 <https://doi.org/10.5194/acp-19-3257-2019>, 2019.

217

218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239

Referee 2

The authors well addressed the reviewers' comments and concerns. It's a nice addition to clarify the limitation of satellite data for detecting ozone pollution. The additional evaluation with ground-based observations makes the results more robust. However, I have some concerns with the evaluation with TOAR sites (Page 9, Line 10). There is large discrepancy between OMI-derived and observed trends in ozone, especially at the Hok Tsui site, where OMI underestimates the change by more than a factor of two. I'd suggest the authors provide possible explanations for such discrepancy, instead of making the questionable statement that 'they are fairly consistent'. Given the discrepancy, I think it's necessary to provide an uncertainty estimate for the derived trends.

Response. Thanks. Now we have made these changes in text.

P9L8. For Lin'an, Hong Kong, and the 5 sites in Taiwan (we report the mean value here), the changes of mean ozone concentrations from 2005-2009 to 2013-2017 are 1.1 ± 3.6 , 2.3 ± 3.3 , and -0.18 ± 2.9 ppbv as estimated from OMI, compared to 0.7 ± 3.6 , 5.6 ± 3.9 (or 5.8 ± 1.3 in Hok Tsui station), and -0.75 ± 2.5 ppbv for MDA8 ozone at the TOAR sites. The changes in the number of ozone episodes per summer are 1.2 ± 0.7 , 1.9 ± 0.24 , and -0.17 ± 0.14 days in OMI, compared to 2.1 ± 4.4 , 1.8 ± 1.7 (or 2.1 ± 1.1 in Hok Tsui station), and -3.5 ± 1.8 days at the TOAR sites. The standard errors are obtained by applying a parametric bootstrap method. The OMI inferred trends are generally consistent with the long-term records available from surface sites.

An evaluation of the OMI satellite instrument's ability to observe boundary layer ozone pollution across China: application to 2005-2017 ozone trends

Lu Shen¹, Daniel J. Jacob¹, Xiong Liu², Guanyu Huang³, Ke Li¹, Hong Liao⁴, Tao Wang⁵

¹John A. Paulson School of Engineering and Applied Sciences, Harvard University, Cambridge, MA 02138, USA

²Harvard-Smithsonian Center for Astrophysics, Cambridge, Massachusetts 02138, USA

³Environmental & Health Sciences, Spelman College, Atlanta, Georgia 30314, USA

⁴School of Environmental Science and Engineering, Nanjing University of Information Science & Technology, Nanjing 210044, China

⁵Department of Civil and Environmental Engineering, The Hong Kong Polytechnic University, Hong Kong

10 Correspondence to: Lu Shen (lshen@fas.harvard.edu)

Abstract. Nadir-viewing satellite observations of tropospheric ozone in the UV have been shown to have some sensitivity to boundary layer ozone pollution episodes but no quantitative evaluation of this ability has been done so far. Here we use 2013-2017 surface ozone data from the new China Ministry of Ecology and Environment (MEE) network of ~1000 sites, together with vertical profiles from ozonesondes and aircraft, to quantify the ability of tropospheric ozone retrievals from the OMI satellite instrument and to detect boundary layer ozone pollution in China. We focus on summer when ozone pollution in China is most severe and when OMI has the strongest sensitivity. After subtracting the Pacific background, we find that the 2013-2017 mean OMI ozone enhancements over eastern China have strong spatial correlation with the corresponding multiyear means in the surface afternoon observations ($R = 0.73$), and that OMI can estimate these multiyear means in summer afternoon surface ozone with a precision of 8 ppb. The OMI data show significantly higher values on observed surface ozone episode days (>82 ppb) than on non-episode days. Day-to-day correlations with surface ozone are much weaker due to OMI instrument noise, and are stronger for sites in southern China (<34°N; $R = 0.3-0.6$) than in northern China ($R = 0.1-0.3$) because of weaker retrieval sensitivity and larger upper tropospheric variability in the north. Ozonesonde data show that much of the variability of OMI ozone over southern China in summer is driven by the boundary layer. Comparison of 2005-2009 and 2013-2017 OMI data indicates that mean summer afternoon surface ozone in southern China (including urban and rural regions) has increased by 3.5 ± 3.0 ppb over the 8-year period and that the number of episode days per summer has increased by 2.2 ± 0.4 (as diagnosed by an extreme value model), generally consistent with the few long-term surface records. Ozone increases have been particularly large in the Yangtze River Delta and in the Hubei, Guangxi, and Hainan provinces.

- Lu Shen 4/18/19 8:00 AM
Deleted: detect surface
- Jacob, Daniel J. 4/20/19 3:08 PM
Deleted: surface or near-surface
- Jacob, Daniel J. 4/20/19 3:10 PM
Deleted: ability correlation of between OMI
- Lu Shen 4/18/19 8:07 AM
Deleted: to characterize
- Jacob, Daniel J. 4/20/19 3:11 PM
Deleted: surface
- Jacob, Daniel J. 4/20/19 3:14 PM
Deleted: correlation between observed surface ozone and the
- Jacob, Daniel J. 4/20/19 3:16 PM
Deleted: can quantify estimate the broad multi-year spatial distribution of mean
- Lu Shen 4/18/19 8:16 AM
Deleted: .4
- Jacob, Daniel J. 4/20/19 3:17 PM
Deleted: and a spatial correlation coefficient $R=0.73$
- Jacob, Daniel J. 4/20/19 3:19 PM
Deleted: In terms of day-to-day variability, OMI is much more successful at capturing the day-to-day variability of has a higher correlation with surface ozone at
- Jacob, Daniel J. 4/20/19 3:19 PM
Deleted: (
- Jacob, Daniel J. 4/20/19 3:49 PM
Deleted: M
- Jacob, Daniel J. 4/20/19 3:49 PM
Deleted: appears to be
- Jacob, Daniel J. 4/20/19 8:20 PM
Deleted: fairly

1 Introduction

Ozone in surface air is harmful to public health (Bell et al., 2004). It is produced by photochemical oxidation of volatile organic compounds (VOCs) in the presence of nitrogen oxides ($\text{NO}_x \equiv \text{NO} + \text{NO}_2$). Both VOCs and NO_x are emitted in large amounts in polluted regions by fuel combustion and industry. Ozone pollution is a particularly severe problem in China, where the air quality standard of 82 ppb (maximum 8-h daily average) is frequently exceeded (Wang et al., 2017). Observations in eastern China have reported increasing ozone trends of 1-3 ppb a^{-1} over the past decade (Sun et al., 2016; Gao et al., 2017; Ma et al., 2017; Li et al., 2019). The surface observations were very sparse until 2013, when data from a national network of ~1000 sites operated by the China Ministry of Ecology and Environment (MEE) started to become available. Here we use the MEE network data to evaluate the ability of the space-based Ozone Monitoring Instrument (OMI) to observe ozone pollution in China, and we use the OMI data going back to 2005 to infer long-term ozone pollution trends.

OMI measures atmospheric ozone absorption by solar backscatter in the UV (270-365 nm) (Levelt et al., 2006). It follows a long lineage of UV satellite instruments (TOMS series starting in 1979, GOME series starting in 1995) directed primarily at monitoring the total ozone column. Retrieval of tropospheric ozone (only ~10% of the column) from these instruments has mostly been done in the past by subtracting independent satellite measurements of stratospheric ozone (Fishman et al., 1987; Ziemke et al., 2011) or using the convective cloud differential method (Ziemke et al., 1998, 2019). OMI has sufficiently fine spectral resolution to allow direct retrieval of tropospheric ozone, although the sensitivity decreases strongly toward the surface because of Rayleigh scattering (Liu et al., 2010). The direct retrieval typically provides one piece of information for the tropospheric ozone column weighted towards the middle troposphere (Zhang et al., 2010).

A number of previous studies have shown that satellite observations of ozone can detect boundary layer ozone pollution events (Fishman et al., 1987; Shim et al., 2009; Eremenko et al., 2008; Hayashida et al., 2008;), including for Chinese urban plumes (Kar et al., 2010; Hayashida et al., 2015; Gaudel et al., 2018; Dufour et al., 2018). Even if sensitivity to the boundary layer is low, the enhancements can be sufficiently large to enable detection. However, no quantitative comparison of the satellite data to surface observations has so far been done. Surface ozone network data are available in the US and Europe but levels are generally too low to enable effective comparison. Ozone levels in China are much higher (Lu et al., 2018). The high density of the MEE network, combined with vertical profile information from ozonesondes and aircraft, provides a unique opportunity for evaluating quantitatively the ability of OMI to observe ozone pollution.

2 Data and Methods

We use the OMI Ozone Profile retrieval (PROFOZ v0.9.3, level 2) product (Liu et al., 2010; Kim et al., 2013; Huang et al.,

Jacob, Daniel J. 4/20/19 3:55 PM

Deleted: surface

Jacob, Daniel J. 4/20/19 3:55 PM

Deleted: lower troposphere

Jacob, Daniel J. 4/20/19 3:59 PM

Deleted: statistically meaningful validation

2017, 2018) from the Smithsonian Astrophysical Observatory (SAO). OMI is in polar sun-synchronous orbit with a 1330 local observation time, and provides daily global mapping with $13 \times 24 \text{ km}^2$ nadir pixel resolution (Levelt et al., 2006). Partial ozone columns are retrieved by PROFOZ for 24 vertical layers, of which 3-7 are in the troposphere with pressure levels dependent on tropopause and surface pressure (Liu et al., 2010). The retrieval uses a Bayesian optimization algorithm with prior information from the McPeters et al. (2007) climatology varying only by latitude and month. Averaging kernel matrices quantifying retrieval sensitivity are provided for individual retrievals. The trace of the averaging kernel matrix below a given retrieval pressure (degrees of freedom for signal or DOFS) estimates the number of independent pieces of information on the ozone profile below that pressure. The DOFS for the tropospheric ozone column in summer as retrieved by PROFOZ is about 1 (Zhang et al., 2010). The PROFOZ tropospheric retrievals have been successfully validated with ozonesonde data (Huang et al., 2017).

We focus on summer when ozone pollution in China is most severe and when OMI has the strongest sensitivity (Zhang et al., 2010). Since 2009, certain cross-track OMI observations have degraded because of the so-called row anomaly (Kroon et al., 2011; Huang et al., 2017, 2018). We only use pixels that (1) pass the reported quality checks, (2) have a cloud fraction less than 0.3, and (3) have a solar zenith angle less than 60° .

The DOFS below 400 hPa over eastern China are in the range 0.3-0.6 (Figure 1a). The DOFS are higher in the south than in the north due to higher solar elevation in the south, and higher over China than in background air at the same latitude due to higher ozone abundances. We use $\text{DOFS} > 0.3$ in Figure 1a as criterion for further analysis; this excludes northern and western China. Even though a DOFS of 0.3 is still low, it is based on the prior estimate of low boundary layer ozone in the McPeters et al. (2007) zonal mean climatology. As we will see, the retrieval is sensitive to ozone enhancements in the boundary layer when these are sufficiently high.

The prior estimate from McPeters et al. (2007) includes a latitudinal gradient of ozone concentrations that may be retained in the retrieval. To remove this background gradient and also any long-term uniform drift in the data, we subtract the monthly mean Pacific background (150°E - 150°W) from the OMI data over China for the corresponding latitude and month. The residual defines an OMI enhancement over China that we use for further analysis. This subtraction requires that we use a common pressure range for the OMI observations over China and the Pacific, but the OMI retrievals have variable pressure ranges depending on the local tropopause and surface pressure (Liu et al., 2010). The three lowest layers in the retrieval (L24-L22) have pressure ranges of approximately 1000-700 hPa, 700-500 hPa, and 500-350 hPa for a column based at sea level, and all contain some information on boundary layer ozone (Figure S1). Here we choose the pressure range 850-400 hPa to define the OMI enhancement relative to the Pacific background, and compute OMI columns for that pressure range by weighting the local L24-22 retrievals. Using 850 hPa as a bottom pressure avoids complications from variable topography in eastern China. The 850-400 hPa retrievals capture all of the OMI sensitivity below 850 hPa in any case. We examined

different spatial and temporal averaging domains for the North Pacific background and found little effect on the residual.

We compare the OMI ozone enhancements to ozone measurements from surface sites, ozonesondes, and aircraft. We use surface ozone measurements from the MEE network available for 2013-2017 (<http://datacenter.mep.gov.cn/index>). We select the summer (JJA) data at 12-15 local solar time (LT), corresponding to the OMI overpass. The network had 450 sites in 2013 and 1500 sites as of 2017, most located in large cities. We also use 2005-2016 summertime ozonesonde data at 12-15 LT for Hanoi (105.8°E, 21.0°N), Hong Kong (114.2°E, 22.3°N), Naha (127.7°E, 26.2°N), Tsukuba (140.1°E, 36.1°N), and Sapporo (141.3°E, 43.1°N), available from the World Ozone and Ultraviolet Radiation Data Center (WOUDC) (<http://woudc.org/>). We further use take-off/landing vertical profiles at 12-15 LT over East Asia from the In-Service Aircraft for the Global Observing System (IAGOS, <http://www.iagos-data.fr/>). For evaluating the long-term surface ozone trends inferred from OMI, we use 2005-2014 trend statistics for maximum daily 8-hour average (MDA8) ozone from the Tropospheric Ozone Assessment Report (TOAR) (Schultz et al., 2018). We also have 2005-2017 JJA 12-15 LT mean ozone at the Hok Tsui station in Hong Kong (Wang et al., 2009).

3 Inference of surface ozone from OMI observations

Figure 1b shows the mean midday (12-15 LT) surface ozone for the summers of 2013-2017 as measured by the MEE network. Concentrations exceed 70 ppb over most of the North China Plain with particularly high values in the Beijing-Tianjin-Hebei (BTH) megacity cluster. Values are also high in the Yangtze River Delta (YRD), Pearl River Delta (PRD), Sichuan Basin (SCB), and the city of Wuhan in central China. High values extend to the region west of the North China Plain, which is less densely populated but has elevated terrain.

OMI mean ozone abundances at 850-400 hPa for the summers of 2013-2017 are shown in Figure 1c. Values are partial column concentrations in Dobson units ($1 \text{ DU} = 2.69 \times 10^{16} \text{ molecules cm}^{-2}$). After subtracting the North Pacific background for the corresponding latitude in month, we obtain the OMI ozone enhancements shown in Figure 1d. The spatial correlation coefficient between the OMI ozone enhancements and the MEE surface network is $R = 0.73$ over eastern China. The correlation is driven in part by the latitudinal gradient but also by the enhancements in the large megacity clusters identified as rectangles in Figure 1b. Thus the correlation coefficient is $R = 0.55$ for the 26-34°N latitude band including YRD, SCB, and Wuhan. Figure 1e shows the corresponding scatterplot and the reduced major axis (RMA) regression relating the OMI enhancement $\Delta\Omega$ to the 12-15 LT surface concentration $[\text{O}_3]$ (the slope is 0.14 DU/ppb). From there one can estimate [multi-year average](#) surface ozone (ppb) on the basis of the observed OMI enhancement (DU) as

$$[\text{O}_3] = 6.9 \Delta\Omega + 24.6 \pm 8.4 \quad (1)$$

where the error standard deviation (precision) of 8 ppb is inferred from the scatterplot. With such a precision, OMI can

Lu Shen 4/19/19 2:04 PM

Deleted: 4

provide useful information on mean summer afternoon levels of surface ozone in polluted regions.

Capturing the day-to-day variability of surface ozone leading to high-ozone pollution episodes is far more challenging because of noise in individual retrievals. Figure 1f shows the OMI vs. MEE temporal correlation for the daily data. Correlation coefficients are consistently positive and statistically significant, but relatively weak. They are higher in southern China ($R = 0.3-0.6$) than in northern China ($R = 0.1-0.3$), consistent with the pattern of OMI information content (DOFS) in Figure 1a. This implies that OMI can only provide statistical rather than deterministic temporal information on ozone pollution episodes, and may be more useful in South than in North China. We return to this point in Section 4.

Figure 2 (top panel) shows the relationship of OMI enhancements with daily MEE surface ozone concentrations averaged spatially in each of the five megacity clusters identified in Figure 1. Consistent with the distribution of DOFS (Figure 1a), the correlations are higher in PRD, SCB, and Wuhan ($0.42-0.53$, $p < 0.05$) than in YRD (0.35 , $p < 0.05$), and lowest in BTH (0.27 , $p < 0.05$). The correlations indicate some capability for OMI to predict ozone daily variability on a statistical basis. The reduced major axis (RMA) regression slopes are consistent across the five regions and average 0.15 DU/ppb. We define an ozone episode day by afternoon concentrations exceeding 82 ppb, corresponding to the Chinese air quality standard. The bottom panel of Figure 2 compares the OMI ozone enhancements between episode and non-episode days as measured by the surface network. OMI is significantly higher ($p < 0.05$) on episode days for all five regions.

4 OMI boundary layer sensitivity inferred from ozonesondes

The correlation of OMI with the MEE surface ozone data likely does not reflect a direct sensitivity of OMI to surface ozone, which is very weak, but rather a sensitivity to boundary layer ozone extending up to a certain depth and correlated with surface ozone. We examined in more detail the sensitivity of OMI to boundary layer ozone and its day-to-day variability by comparing to summertime 2005-2016 Hong Kong ozonesonde data. Figure 3a shows the measured ozonesonde profiles (in ppb) mapped on a 100 hPa grid and selecting only the days when concurrent OMI retrievals are available ($n = 57$). The boundary layer ozone ($950-850$ hPa) in the ozonesonde data has large day-to-day variability, ranging from 20 to 120 ppb with a mean of 47 ppb. The variability in the free troposphere is much less.

Figure 3b shows the ozonesonde data smoothed by the OMI averaging kernel sensitivities for the corresponding retrievals. The retrievals over Hong Kong have a mean DOFS of 0.46 below 400 hPa. We see from Figure 3c that the OMI information is weighted toward the free troposphere but there is sensitivity in the boundary layer, and since boundary layer variability is much larger it can make a major contribution to OMI variability. The L23 ozone smoothed from the ozonesonde data in Figure 3b has a correlation coefficient of 0.75 with the $950-850$ hPa ozone in the original data. The temporal correlation coefficients of the OMI retrievals at different levels with the $950-850$ hPa ozonesonde data are given in Figure 3d. The correlation coefficient with L23 OMI ozone is 0.51 ($p < 0.05$), and the correlation coefficient with the $850-400$ hPa OMI

ozone constructed by weighting the L24-L23-L22 retrievals is 0.50. Figure 3e shows a scatterplot of the latter. We see that high-ozone episodes in the 950-850 hPa sonde data are systematically associated with high OMI values, though the converse does not always hold because free tropospheric enhancements affecting OMI can also occur. For the 8 boundary layer episode days (> 82 ppb), the average OMI 850-400 hPa ozone is 23.7 ± 3.1 DU, significantly higher than for the non-episode days (18.2 ± 4.1 DU). The Hong Kong ozonesonde data thus indicate that OMI can quantify the frequency of high-ozone episodes in the boundary layer even if it may not be reliable for individual events.

We applied the same daily correlation analysis to the other ozonesonde datasets and IAGOS aircraft measurements during 2005-2017 summers. For the 54 IAGOS vertical profiles coincident with OMI observations, the correlation coefficient of the 950 hPa in situ ozone and 850-400 hPa OMI ozone is $R = 0.59$ ($p < 0.05$) (Figure S2). For the five ozonesonde sites with long-term observations, the correlation coefficients are 0.4-0.6 for Hanoi, Hong Kong, and Naha (south of 30°N), and 0-0.3 for Sapporo and Tsukuba (north of 35°N) (Figure S3), consistent with the patterns of daily correlations for the MEE data (Figure 1f).

The correlation between boundary layer ozone pollution and the OMI ozone retrievals could be due in part to correlation between boundary layer and mid-tropospheric ozone, considering that both tend to be driven by the same weather systems. We used the ozonesonde data to examine what correlation with boundary layer (950-850 hPa) ozone would be observed if OMI were sensitive only to the free troposphere at ~ 500 hPa (where its sensitivity is maximum, Figure 3c) and not to the boundary layer. In that case the correlation coefficient $R_{1,3}$ of boundary layer ozone and the OMI 850-400 hPa retrievals would be given by (Vos, 2009):

$$R_{1,3} = R_{1,2}R_{2,3} \pm \sqrt{(1 - R_{1,2}^2)(1 - R_{2,3}^2)} \quad (2)$$

where $R_{1,2}$ is the correlation coefficient between boundary layer and 500 hPa ozone in the ozonesonde data, and $R_{2,3}$ is that between 500 hPa ozone and the OMI 850-400 hPa retrievals. As seen from Figure S4, $R_{1,3}$ at the five sonde sites is only ~ 0.2 , implying that direct sensitivity to the boundary layer dominates the correlation of OMI with surface ozone at least in southern China. Further evidence for this is the ability of OMI to detect the ozone enhancements in megacity clusters (Figure 1).

We find that the low correlation of OMI with boundary layer ozone in the northern ozonesonde data is due not only to the low DOFS but also to a large variability of ozone in the upper troposphere. Figure 4 (left panel) shows the standard deviation of daily OMI 400-200 hPa ozone during 2005-2017 summers, indicating that upper tropospheric ozone has much higher variability in the north ($> 34^\circ\text{N}$) than in the south. This is related to the location of the jet stream and more active stratospheric influence (Hayashida et al., 2015). Figure 4 (right panel) displays the vertical profiles of ozone standard deviations for the five ozonesonde sites. For the two sites north of 34°N , the ozone variability becomes very large above 8

km. Since the OMI 850-400 hPa retrieval also contains information from above 400 hPa, this upper tropospheric variability causes a large amount of noise that masks the signal from boundary layer variability. For the three sites south of 34°N, the ozone variability in the boundary layer is much higher than in the free troposphere and the upper tropospheric ozone variability still remains low even above 8 km. In the rest of this paper we focus our attention on ozone episodes and the long-term trends in southern China (south of 34°N).

5. Using extreme value theory to predict the occurrence of high-ozone episodes from OMI data

We construct a point process (PP) model from extreme value theory (Cole, 2001) to estimate the likelihood of surface ozone exceeding a high-ozone threshold u (here $u = 82$ ppb at 12-15 LT) at a given site i and day t given the observed OMI ozone enhancement $x_{i,t}$ for that day. The model describes the high tail of the ozone probability density function (pdf) as a Poisson process limit, conditioned on the local OMI observation. Such a model has been used previously to relate the probability of extreme air pollution conditions to meteorological predictor variables (Rieder et al., 2013; Shen et al., 2016, 2017; Pendergrass et al., 2019) but here we use the OMI enhancement as predictor variable. We fit the model to all daily concurrent observations of surface ozone and OMI ozone enhancements for the ensemble of eastern China sites south of 34°N in Figure 5 (90,601 observations for summers 2013-2017). The probability of exceeding the threshold at a site i should depend not only on $x_{i,t}$ but also on its time-averaged value \bar{x}_i , because a high value of \bar{x}_i means that a higher $x_{i,t}$ is less anomalous and more likely to represent an actual ozone exceedance than for a site with low \bar{x}_i . Thus the model has two predictor variables, $x_{i,t}$ and \bar{x}_i .

Details of the PP model can be found in Cole (2001). The model fits three parameters that control the shift, spread and shape of the high-tail pdf. The fit minimizes a cost function L given by

$$L = \prod_{i=1}^m L_i(\mu_i, \sigma_i, \xi) \quad (3)$$

with

$$L_i(\mu_i, \sigma_i, \xi) = \exp\left\{-\frac{1}{n_o} \sum_{t=1}^n \left[1 + \frac{\xi(u - \mu_i)}{\sigma_{i,t}}\right]^{-1/\xi}\right\} \prod_{t=1}^n \left\{\frac{1}{\sigma_{i,t}} \left[1 + \frac{\xi(y_{i,t} - \mu_i)}{\sigma_{i,t}}\right]^{-1/\xi - 1}\right\}^{I(y_{i,t} > u)} \quad (4)$$

$$\mu_i = \alpha_0 + \alpha_1 \bar{x}_i \quad (5)$$

$$\bar{x}_i = \frac{1}{n} \sum_{t=1}^n x_{i,t} \quad (6)$$

$$\sigma_{i,t} = \exp(\beta_0 + \beta_1 x_{i,t}) \quad (7)$$

Here $L_i(\mu_i, \sigma_i, \xi)$ is the cost function for site i and L is for the total cost function for all m sites, $y_{i,t}$ is the daily 12-15 LT MEE surface ozone from each individual site i on day t , $n_a = 92$ is the number of days in summer, μ_i is the location parameter for site i conditioned on the 2013-2017 summertime mean OMI enhancements \bar{x}_i , $\sigma_{i,t}$ is the scale parameter conditioned on the local OMI ozone enhancements $x_{i,t}$, ξ is the shape factor, and $I_{[y_{i,t} > u]}$ is one if observed ozone is above the threshold and zero otherwise. Minimization of the cost function optimizes the values of the parameters α_0 , α_1 , β_0 , β_1 , and ξ given the 90,601 $(x_{i,t}, y_{i,t})$ data pairs. The resulting values are $\alpha_0 = 103$ ppb, $\alpha_1 = 6.0$, $\beta_0 = 2.8$ ppb, $\beta_1 = -0.033$, $\xi = -0.12$. The probability of daily ozone exceeding the threshold u is then calculated as

$$P(y_{i,t} \geq u | x_{i,t}) = \frac{1}{n_a} \left[1 + \xi \left(\frac{u - \mu_i}{\sigma_{i,t}} \right)^{-1/\xi} \right]^{-1/\xi} \quad (8)$$

The model is optimized using the *extRemes* package in R (Gilleland and Katz, 2011). We performed a 10-fold cross validation of the model, in which we partitioned the sites into 10 equal subsets and repeatedly used one subset as testing data and the rest as training data. The results show that the predicted fraction of ozone episodes resembles that observed, with a spatial correlation of 0.62 (Figure 5a). The model tends to underestimate the probability of episodes in polluted regions due to the noise of daily OMI ozone. 82 ppb corresponds to the 84th percentile of the data, which is a relatively low threshold for application of extreme value theory. However, we find that the model can also accurately estimate the probability of exceedance above higher thresholds (Figure 5b) [for the ensemble of eastern China sites south of 34°N](#), which confirms the property of threshold invariance of an extreme value model (Cole, 2001). We also tested the model with uniform location or scale factors, but neither could reproduce the observed spatial distribution of ozone episodes.

6. 2005-2017 trends in surface ozone inferred from OMI data

We used the long-term OMI ozone record for 2005-2017 to infer trends in surface ozone over southern China, not including any tropospheric background trends (removed by our subtraction of the North Pacific). Figure 6 shows the changes between 2005-2009 and 2013-2017 [\(an 8-year period\)](#) in mean summer afternoon ozone concentrations and in the number of high-ozone episode days per summer. [Here we have extended the trend analysis to Taiwan because of the opportunity to compare to surface records.](#) The changes in mean summer afternoon ozone concentrations are obtained from the difference in the mean OMI ozone enhancements [between the two time periods](#) (Figure S5) and applying equation (1). The changes in the number of high-ozone episode days [per summer](#) are obtained by applying the probability of exceeding 82 ppb (equation 8) to each pair of 5 years of OMI data. When averaged across southern China (including urban and rural regions), the mean summer afternoon ozone concentrations have increased by 3.5 ± 3.0 ppb between the two periods (Figure 6a) and the number of ozone episodes (> 82 ppb) has increased by 2.2 ± 0.4 days per summer (Figure 6b). Conditions have become particularly

worse in YRD and in Hubei, Guangxi, and Hainan provinces where the number of high-ozone days per summer has increased by more than 5.

We compared the OMI trends in Figure 6 to the trends of MDA8 ozone and number of high-ozone days reported by the long-term TOAR sites (Schultz et al., 2018) and our own analysis for the Hok Tsui station in Hong Kong (Wang et al., 2009). For Lin'an, Hong Kong, and the 5 sites in Taiwan (we report the mean value here), the changes of mean ozone concentrations from 2005-2009 to 2013-2017 are 1.1 ± 3.6 , 2.3 ± 3.3 , and -0.18 ± 2.9 ppbv, as estimated from OMI, compared to 0.7 ± 3.6 , 5.6 ± 3.9 (or 5.8 ± 1.3 in Hok Tsui station), and -0.75 ± 2.5 ppbv for MDA8 ozone at the TOAR sites. The changes in the number of ozone episodes per summer are 1.2 ± 0.7 , 1.9 ± 0.24 , and -0.17 ± 0.14 days in OMI, compared to 2.1 ± 4.4 , 1.8 ± 1.7 (or 2.1 ± 1.1 in Hok Tsui station), and -3.5 ± 1.8 days at the TOAR sites. The standard errors are obtained by applying a parametric bootstrap method. The OMI inferred trends are generally consistent with the long-term records available from surface sites.

5 Discussion and conclusions

Satellite observations of tropospheric ozone in the UV could provide an indicator of surface ozone pollution if the associated boundary layer enhancement is large enough. We presented a quantitative evaluation of this capability for OMI ozone retrievals in China by comparison to the extensive 2013-2017 ozone network data from the China Ministry of Ecology and Environment (MEE), together with vertical profiles from ozonesondes and aircraft. We went on to use the long-term OMI record (2005-2017) to infer surface ozone pollution trends over that period.

After subtracting the contribution from the North Pacific background, we find that the OMI enhancement over eastern China can reproduce the observed spatial distribution of multi-year mean summer afternoon ozone concentrations at the MEE sites, with a correlation coefficient $R = 0.73$ and a precision of 8 ppb. Even though OMI has little sensitivity to surface ozone, the high ozone levels seen at surface sites propagate deep enough in the boundary layer to be observed by OMI. Day-to-day correlation at individual sites is weaker ($R = 0.1-0.3$ north of 34°N , $0.3-0.6$ south of 34°N) because of noise in individual OMI retrievals. But we find that OMI is statistically enhanced in urban areas when surface ozone exceeds an 8-h maximum daily average (MDA8) value of 82 ppb (the Chinese air quality standard).

To better understand the correlation of OMI with surface ozone we examined vertical ozone profiles from Hong Kong and other ozonesondes, and from the IAGOS commercial aircraft program. Some of the correlation is driven by similar meteorology influencing ozone in the mid-troposphere (where OMI sensitivity is maximum) and the boundary layer, but most of the correlation is driven by direct sensitivity to the boundary layer. The Hong Kong ozonesonde data also indicates that OMI can quantify the frequency of high-ozone episodes in the boundary layer even if it may not be reliable for individual events. In southern China ($< 34^\circ\text{N}$), we find that ozone variability in the tropospheric column is dominated by the

Lu Shen 4/18/19 8:31 PM

Deleted: 2

Lu Shen 4/18/19 8:33 PM

Deleted: (standard deviation among the 5 sites)

Lu Shen 4/19/19 12:54 PM

Deleted: in

Lu Shen 4/18/19 8:33 PM

Deleted: 3.4

Lu Shen 4/18/19 8:33 PM

Deleted: 74

Jacob, Daniel J. 4/20/19 7:54 PM

Deleted: se

Jacob, Daniel J. 4/20/19 8:23 PM

Deleted: fairly

Jacob, Daniel J. 4/20/19 7:58 PM

Deleted: near-

Lu Shen 4/19/19 2:04 PM

Deleted: .4

boundary layer, explaining the stronger correlations there. The lower correlation of OMI with surface ozone further north is due to large upper tropospheric variability in addition to lower sensitivity.

We went on to use the 2005-2017 OMI record to diagnose long-term trends of surface ozone in southern China (<34 °N). This involved the development of a point process model from extreme value theory to infer the probability of surface ozone exceeding 82 ppb and higher thresholds on the basis of the daily observed OMI ozone enhancements. The OMI record shows a general increase across southern China (including urban and rural regions) from 2005-2009 to 2013-2017 (8-year period) in mean summertime afternoon ozone (+3.5±3.0 ppb) and in the frequency of high-ozone episodes (+2.2±0.4 days per summer). Increases are particularly large in the Yangtze River Delta and in Hubei, Guangxi, and Hainan provinces. The trends are generally consistent with the few long-term records available from surface sites.

Our method for inferring ozone pollution and the frequency of high-ozone episodes from the OMI satellite data may be applied to other regions of the world where surface ozone is expected to be high but where in situ observations are lacking. The next generation of UV satellite instruments may improve this capability. The TROPOMI instrument launched in October 2017 is now providing daily observations with 3.5×7 km² pixel resolution, much finer than OMI (Theys et al., 2017). The GEMS (Geostationary Environment Monitoring Spectrometer) instrument is expected to launch in late 2019 and will observe East Asia with hourly frequency and sensitivity similar to OMI (Bak et al., 2013). The TEMPO (Tropospheric Emissions: Monitoring of Pollution) geostationary satellite instrument to be launched around 2020 over North America will have a spectral range extending to the visible Chappuis bands where ozone detection sensitivity remains high down to the surface (Zoogman et al., 2011, 2017). This should allow for improved observations of surface ozone, particularly where concentrations are not as high as they are presently in China.

Author contribution

L. Shen and D. Jacob designed the experiments and L. Shen carried them out. X. Liu and G. Huang provided the satellite data. K. Li and H. Liao provided the surface observations. L. Shen and D. Jacob prepared the manuscript with contributions from all co-authors.

Acknowledgments

This work was funded by the Harvard Global Institute (HGI), by the NASA Earth Science Division, and by the Joint Laboratory for Air Quality and Climate (JLAQC) between Harvard and the Nanjing University for Information Sciences and Technology (NUIST).

Jacob, Daniel J. 4/20/19 8:20 PM

Deleted: fairly

References

- Bell, M.L., McDermott, A., Zeger, S.L., Samet, J.M. and Dominici, F.: Ozone and short-term mortality in 95 US urban communities, 1987-2000, *JAMA*, 292(19), 2372-2378, 2004.
- Bak, J., Kim, J. H., Liu, X., Chance, K., and Kim, J.: Evaluation of ozone profile and tropospheric ozone retrievals from GEMS and OMI spectra, *Atmos. Meas. Tech.*, 6, 239-249, <http://sci-hub.tw/10.5194/amt-6-239-2013>, 2013.
- Coles, S. G.: *An Introduction to Statistical Modeling of Extreme Values*, Springer, New York, 2001.
- [Dufour, G., Eremenko, M., Beekmann, M., Cuesta, J., Foret, G., Fortems-Cheiney, A., Lachâtre, M., Lin, W., Liu, Y., Xu, X., and Zhang, Y.: Lower tropospheric ozone over the North China Plain: variability and trends revealed by IASI satellite observations for 2008–2016, *Atmos. Chem. Phys.*, 18, 16439-16459, <https://doi.org/10.5194/acp-18-16439-2018>, 2018.](#)
- Eremenko, M., Dufour, G., Foret, G., Keim, C., Orphal, J., Beekmann, M., Bergametti, G., and Flaud, J.-M.: Tropospheric ozone distributions over Europe during the heat wave in July 2007 observed from infrared nadir spectra recorded by IASI, *Geophys. Res. Lett.*, 35, L18805, doi:10.1029/2008GL034803, 2008.
- Fishman, J., Vukovich, F. M., Cahoon, D. R., and Shipham, M. C.: The characterization of an air pollution episode using satellite total ozone measurements, *J. Clim. Appl. Meteorol.*, 26, 1638–1654, 1987.
- Gao W, et al.: Long-term trend of O₃ in a mega City (Shanghai), China: Characteristics, causes, and interactions with precursors, *Sci. Total. Environ.*, 603, 425-433, 2017.
- [Gaudel, A., Cooper, O. R., Ancellet, G., Barret, B., Boynard, A., Burrows, J. P., Clerbaux, C., Coheur, P. F., Cuesta, J., Cuevas Agulló, E., and Doniki, S.: Tropospheric Ozone Assessment Report: Present-day distribution and trends of tropospheric ozone relevant to climate and global atmospheric chemistry model evaluation, *Elem. Sci. Anth.*, 6, 39, <https://doi.org/10.1525/elementa.291>, 2018.](#)
- Gilleland, E., Katz, R. W.: New software to analyze how extremes change over time, *Eos Trans. AGU*, 92(2), 13–14, doi:10.1029/2011EO020001, 2011.
- Hayashida, S., Urita, N., Noguchi, K., Liu, X., and Chance, K.: Spatiotemporal Variation in Tropospheric Column Ozone over East Asia Observed by GOME and Ozonesondes, *Sola*, 4, 117–120, 2008.
- Hu, L., Jacob, D. J., Liu, X., Zhang, Y., Zhang, L., and Kim, P. S.: Global budget of tropospheric ozone: evaluating recent model advances with satellite (OMI), aircraft (IAGOS), and ozonesonde observations, *Atmos. Environ.*, 1–36, 2017.
- Huang, G., Liu, X., Chance, K., Yang, K., Bhartia, P. K., Cai, Z., Allaart, M., Ancellet, G., Calpini, B., Coetzee, G. J. R., Cuevas-Agulló, E., Cupeiro, M., De Backer, H., Dubey, M. K., Fuelberg, H. E., Fujiwara, M., Godin-Beekmann, S., Hall, T. J., Johnson, B., Joseph, E., Kivi, R., Kois, B., Komala, N., König-Langlo, G., Laneve, G., Leblanc, T., Marchand, M., Minschwaner, K. R., Morris, G., Newchurch, M. J., Ogino, S.-Y., Ohkawara, N., Piters, A. J. M., Posny, F., Querel, R., Scheele, R., Schmidlin, F. J., Schnell, R. C., Schrems, O., Selkirk, H., Shiotani, M., Skrivánková, P., Stübi, R., Taha, G., Tarasick, D. W., Thompson, A. M., Thouret, V., Tully, M. B., Van Malderen, R., Vömel, H., von der Gathen, P., Witte, J. C., and Yela, M.: Validation of 10-year SAO OMI Ozone Profile (PROFOZ)

- product using ozonesonde observations, *Atmos. Meas. Tech.*, 10, 2455-2475, <http://sci-hub.tw/10.5194/amt-10-2455-2017>, 2017.
- Huang, G., Liu, X., Chance, K., Yang, K., and Cai, Z.: Validation of 10-year SAO OMI ozone profile (PROFOZ) product using Aura MLS measurements, *Atmos. Meas. Tech.*, 11, 17-32, <http://sci-hub.tw/10.5194/amt-11-17-2018>, 2018.
- 5 Kar, J., Fishman, J., Creilson, J. K., Richter, A., Ziemke, J., and Chandra, S.: Are there urban signatures in the tropospheric ozone column products derived from satellite measurements?, *Atmos. Chem. Phys.*, 10, 5213-5222, <http://sci-hub.tw/10.5194/acp-10-5213-2010>, 2010.
- Kim, P. S., Jacob, D. J., Liu, X., Warner, J. X., Yang, K., Chance, K., Thouret, V., and Nedelec, P.: Global ozone-CO correlations from OMI and AIRS: constraints on tropospheric ozone sources, *Atmos. Chem. Phys.*, 13, 9321-9335, <http://sci-hub.tw/10.5194/acp-13-9321-2013>, 2013.
- 10 Kroon, M., de Haan, J. F., Veeffkind, J. P., Froidevaux, L., Wang, R., Kivi, R., and Hakkarainen, J. J.: Validation of operational ozone profiles from the Ozone Monitoring Instrument, *J. Geophys. Res.*, 116, D18305, doi:10.1029/2010JD015100, 2011.
- Levelt, P. F., Van den Oord, G. H. J., Dobber, M. R., Malkki, A., Visser, H., de Vries, J., Stammes, P., Lundell, J. O. V., and Saari, H.: The Ozone Monitoring Instrument, *IEEE T. Geosci. Remote*, 44, 1093-1101, 2006.
- 15 Liu, X., Bhartia, P. K., Chance, K., Spurr, R. J. D., and Kurosu, T. P.: Ozone profile retrievals from the Ozone Monitoring Instrument, *Atmos. Chem. Phys.*, 10, 2521-2537, <http://sci-hub.tw/10.5194/acp-10-2521-2010>, 2010.
- Li, K., Jacob, D.J., Liao, H., Shen, L., Zhang, Q. and Bates, K.H.: Anthropogenic drivers of 2013–2017 trends in summer surface ozone in China, *Proc. Natl. Acad. Sci. U.S.A.*, 116(2), 422-427, 2019.
- 20 Lu X., Hong J., Zhang L., Cooper OR.,Schultz MG., Xu X., Wang T., Gao M., Zhao Y., and Zhang Y., Severe surface ozone pollution in China: a global perspective, *Environ. Sci. Technol. Lett.*, 5, 487–494, 2018.
- Ma, Z., Xu, J., Quan, W., Zhang, Z., Lin, W., and Xu, X.: Significant increase of surface ozone at a rural site, north of eastern China, *Atmos. Chem. Phys.*, 16, 3969-3977, <http://sci-hub.tw/10.5194/acp-16-3969-2016>, 2016.
- McPeters, R. D., Labow, G. J., and Logan, J. A.: Ozone climatological profiles for satellite retrieval algorithms, *J. Geophys. Res.*, 112, D05308, doi:10.1029/2005JD006823, 2007.
- 25 Rieder, H. E., Fiore, A. M., Polvani, L. M., Lamarque, J. F., and Fang, Y.: Changes in the frequency and return level of high ozone pollution events over the eastern United States following emission controls, *Environ. Res. Lett.*, 8, 014012, doi:10.1088/1748-9326/8/1/014012, 2013.
- Pendergrass, D. C., Shen, L., Jacob, D.J., and Mickley, L. J.: Predicting the impact of climate change on severe wintertime particulate pollution events in Beijing using extreme value theory, *Geophys. Res. Lett.*, 46, 1824–1830, <https://doi.org/10.1029/2018GL080102>, 2019.
- 30 Shen, L., Mickley, L. J., Gilleland E.: Impact of increasing heat waves on U.S. ozone episodes in the 2050s: Results from a multimodel analysis using extreme value theory, *Geophys. Res. Lett.*, 43, 4017–4025, doi:10.1002/2016GL068432, 2016

- Shen, L., Mickley, L. J., Leibensperger, E.M., Li, M.: Strong dependence of U.S. summertime air quality on the decadal variability of Atlantic sea surface temperatures, *Geophys. Res. Lett.*, 44, 12,527–12,535, <https://doi.org/10.1002/2017GL075905>, 2017
- Shim, C., Li, Q., Luo, M., Kulawik, S., Worden, H., Worden, J., Eldering, A., Diskin, G., Sachse, G., Weinheimer, A., Knapp, D., Montzka, D., Campos, T.: Satellite observations of Mexico City pollution outflow from the Tropospheric Emissions Spectrometer (TES), *Atmos. Environ.*, 43(8), 1540–1547, 2009.
- Sun, L., Xue, L., Wang, T., Gao, J., Ding, A., Cooper, O. R., Lin, M., Xu, P., Wang, Z., Wang, X., Wen, L., Zhu, Y., Chen, T., Yang, L., Wang, Y., Chen, J., and Wang, W.: Significant increase of summertime ozone at Mount Tai in Central Eastern China, *Atmos. Chem. Phys.*, 16, 10637–10650, <http://sci-hub.tw/10.5194/acp-16-10637-2016>, 2016.
- 10 Theys, N., De Smedt, I., Yu, H., Danckaert, T., van Gent, J., Hörmann, C., Wagner, T., Hedelt, P., Bauer, H., Romahn, F., Pedernana, M., Loyola, D., and Van Roozendael, M.: Sulfur dioxide retrievals from TROPOMI onboard Sentinel-5 Precursor: algorithm theoretical basis, *Atmos. Meas. Tech.*, 10, 119–153, <https://doi.org/10.5194/amt-10-119-2017>, 2017.
- Vos P.: Pearson's correlation between three variables; using students' basic knowledge of geometry for an exercise in mathematical statistics, *Int. J. Sci. Math. Educ.*, 40(4), 533–54, 2009.
- 15 Wang, T., Wei, X.L., Ding, A.J., Poon, C.N., Lam, K.S., Li, Y.S., et al., 2009a. Increasing surface ozone concentrations in the background atmosphere of Southern China, 1994–2007. *Atmos. Chem. Phys.* 9, 6217–6227.
- Wang, T., Xue, L. K., Brimblecombe, P., Lam, Y. F., Li, L., and Zhang, L.: Ozone pollution in China: A review of concentrations, meteorological influences, chemical precursors, and effects, *Sci. Total Environ.*, 575, 1582–1596, 2017.
- 20 Zhang, L., Jacob, D. J., Liu, X., Logan, J. A., Chance, K., Eldering, A., and Bojkov, B. R.: Intercomparison methods for satellite measurements of atmospheric composition: application to tropospheric ozone from TES and OMI, *Atmos. Chem. Phys.*, 10, 4725–4739, <http://sci-hub.tw/10.5194/acp-10-4725-2010>, 2010.
- Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L., Zhang, Y., Zhao, H., Zheng, Y., He, K., and Zhang, Q.: Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18, 14095–14111, <http://sci-hub.tw/10.5194/acp-18-14095-2018>, 2018.
- 25 [Ziemke, J. R., Chandra, S., and Bhartia, P. K.: Two new methods for deriving tropospheric column ozone from TOMS measurements: The assimilated UARS MLS/HALOE and convective-cloud differential techniques, *J. Geophys. Res.*, 103, 22115–22127, 1998.](#)
- Ziemke, J. R., Chandra, S., Labow, G. J., Bhartia, P. K., Froidevaux, L., and Witte, J. C.: A global climatology of tropospheric and stratospheric ozone derived from Aura OMI and MLS measurements, *Atmos. Chem. Phys.*, 11, 9237–9251, <http://sci-hub.tw/10.5194/acp-11-9237-2011>, 2011.
- 30 [Ziemke, J. R., Oman, L. D., Strode, S. A., Douglass, A. R., Olsen, M. A., McPeters, R. D., Bhartia, P. K., Froidevaux, L., Labow, G. J., Witte, J. C., Thompson, A. M., Haffner, D. P., Kramarova, N. A., Frith, S. M., Huang, L.-K., Jaross, G. R., Seftor, C. J., Deland, M. T., and Taylor, S. L.: Trends in global tropospheric ozone inferred from a composite record](#)

[of TOMS/OMI/MLS/OMPS satellite measurements and the MERRA-2 GMI simulation](#), *Atmos. Chem. Phys.*, **19**, 3257–3269, <https://doi.org/10.5194/acp-19-3257-2019>, 2019.

- Zoogman, P., Liu, X., Suleiman, R. M., Pennington, W. F., Flittner, D. E., Al, Saadi, J. A., et al.: Tropospheric emissions: Monitoring of pollution (TEMPO), *J. Quant. Spectrosc. Radiat. Transf.*, **186**, 17–39. <https://doi.org/10.1016/j.jqsrt.2016.05.008>, 2017.
- 5 Zoogman, P., Jacob, D. J., Chance, K., Zhang, L., Sager, P. L., Fiore, A. M., Eldering, A., Liu, X., Natraj, V., and Kulawik, S. S.: Ozone air quality measurement requirements for a geostationary satellite mission, *Atmos. Environ.*, **45**, 7143–7150, 2011.

Summer (JJA) 2013-2017 ozone over China from the MEE network and the OMI instrument





Figure 1. Summertime observations of ozone over China (JJA 2013-2017) from the MEE surface network and the OMI satellite instrument. (a) Mean degrees of freedom for signal (DOFS) of OMI ozone retrievals below 400 hPa. We limit our attention to the China domain with DOFS > 0.3 (south of dashed line) and to sites with at least 100 concurrent surface and OMI observations for the 2013-2017 period. (b) Mean midday (12-15 local time) ozone concentrations from the MEE surface network. Rectangles identify high-ozone regions discussed in the text including Beijing-Tianjing-Hebei (BTH, 114°-121°E, 34-41°E), Yangtze River Delta (YRD, 119.5°-121.5°E, 30-32.5°E), Pearl River Delta (PRD, 112.5°-114.5°E, 22-24°E), Sichuan Basin (SCB, 103.5°-105.5°E, 28-31.5°E), and Wuhan (113.5°-115.5°E, 29.5-31.5°E). (c) Mean OMI partial columns at 850-400 hPa. (d) Mean OMI ozone enhancements at 850-400 hPa after subtraction of the latitude-dependent mean background over the Pacific (150°E-150°W). (e) Spatial correlation of mean JJA 2013-2017 MEE ozone with the OMI ozone enhancement at 850-400 hPa. The correlation coefficient and the fitted reduced major axis (RMA) regression equation are shown inset. (f) Temporal correlation coefficients (R) of daily MEE surface ozone with OMI at individual sites, measuring the ability of OMI to capture the day-to-day variability of surface ozone.

15



5 | **Figure 2.** Ability of daily OMI observations to detect high-ozone episodes in the five megacity clusters of Figure 1. Daily surface afternoon (12-15 local time) observations from the MEE network in summer (JJA) 2013-2017 averaged over the megacity clusters are compared to the corresponding OMI enhancements relative to the Pacific background. The top panels show the correlations in the daily data, with correlation coefficients inset. Reduced-major-axis (RMA) linear regression lines are also shown. The bottom panels show the distributions of OMI enhancements for episode (≥ 82 ppbv) and non-episode (< 82 ppbv) days. The top and bottom of each box are the 25th and 75th percentiles, the centerline is the median, the vertical bars are the 2nd and 98th percentiles, and the dots are outliers.




Figure 3. Ozone vertical profiles over Hong Kong in summer (JJA) 2015-2016. (a) Ozonesonde data coincident with OMI observations ($n=57$), averaged over a 100 hPa grid and arranged in chronological order. (b) The same ozonesonde data but smoothed by the OMI averaging kernels. Mean pressures for each OMI retrieval level are indicated. (c) Mean averaging kernel sensitivities for each OMI retrieval level, as described by the rows of the averaging kernel matrix; values are shown for August 2015 but are similar in other summer months and years. The dashed lines are boundaries between retrieval levels. (d) OMI ozone observations coincident with the ozonesondes. The correlations of unsmoothed 950-850 hPa ozonesonde data with the OMI retrievals for different levels are shown inset. (e) Relationship of unsmoothed 950-850 hPa ozonesonde data and OMI 850-400 hPa ozone. The correlation is shown inset. The dashed line corresponds to the Chinese ozone air quality standard (82 ppb).

Standard deviation of ozone concentrations, summers 2005-2017





Figure 4. (a) Standard deviation of daily OMI 400-200 hPa ozone in East Asia during 2005-2017 summers. The triangles are the locations of ozonesonde sites with observations during this period. (b) Vertical profiles of daily ozone standard deviation in 1-km bins (DU/km) in the ozonesonde data for the
5 2005-2017 summers.

Evaluation of extreme value point process model for high-ozone episodes

(a) Probability of exceeding 82 ppb in the daily 12-15 LT data



(b) Probabilities of exceeding 82-130 ppb ozone thresholds




Figure 5. Evaluation of the extreme value point process (PP) model for predicting the probability of occurrence of summertime high-ozone episodes from the OMI daily data. The episodes are defined by exceedance of a given ozone threshold in the 3-hour average data at 12-15 local time. (a) Observed and predicted probability of ozone episode days exceeding a 82 ppb threshold. The predicted probability is calculated from equation (8). (b) Observed and predicted probabilities of exceeding higher thresholds from 82 to 130 ppb.

Changes in summertime surface ozone pollution inferred from OMI
(2005-2009 to 2013-2017)




Figure 6. Changes in surface ozone pollution between the 2005-2009 and 2013-2017 periods (separated by 8 years) as inferred from OMI afternoon observations at around 13:30 local time. (a) Change in mean summer afternoon concentrations, obtained from the difference in the mean OMI ozone enhancements and applying equation (1). Also shown with symbols are observed changes in mean MDA8 ozone from in situ observations in Lin'an, Hong Kong, and Taiwan reported by TOAR (Schultz et al., 2018). Because the TOAR observations are only reported for 2005-2014, we estimate the changes from 2005-2009 to 2013-2017 on the basis of the reported linear trends during 2005-2014 (ppb a^{-1}). (b) Change in the number of high-ozone days ($> 82 \text{ ppb}$) per summer, calculated by applying the probability of exceeding 82 ppb (equation 8) to the daily OMI enhancements. Also shown with symbols are observed changes of the number of days with MDA8 ozone exceeding 80 ppb at the TOAR sites, similarly adjusted as the change from 2005-2009 to 2013-2017.