Characteristics of wintertime VOCs in suburban and urban Beijing: concentrations, emission ratios, and festival effects

Kun Li^{1,2,3}, Junling Li^{1,2}, Shengrui Tong^{1,2}, Weigang Wang^{1,2}, Ru-Jin Huang⁴, and Maofa Ge^{1,2,5}

5

¹Beijing National Laboratory for Molecular Sciences (BNLMS), State Key Laboratory for Structural Chemistry of Unstable and Stable Species, CAS Research/Education Center for Excellence in Molecular Sciences, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100190, China

²University of Chinese Academy of Sciences, Beijing 100049, China.

10 ³Air Quality Research Division, Environment and Climate Change Canada, Toronto, Ontario M3H 5T4, Canada.

⁴Key Laboratory of Aerosol Chemistry & Physics, State Key Laboratory of Loess and Quaternary Geology, Institute of Earth Environment, Chinese Academy of Sciences, Xi'an 710061, China

⁵Center for Excellence in Regional Atmospheric Environment, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen 361021, China

15 Correspondence: Maofa Ge (gemaofa@iccas.ac.cn)



Figure S1. Time series of wind speed, wind direction, temperature, and relative humidity (RH) at the suburban site (a) and the urban site (b).



Figure S2. Scatterplots of representative inert primary VOCs (acetonitrile and benzene) over CO at the suburban site.