

MAX-DOAS measurements of tropospheric NO₂ and HCHO in Nanjing and the comparison to OMI observations

Ka Lok Chan¹, Zhuoru Wang¹, Aijun Ding², Klaus-Peter Heue¹, Yicheng Shen², Jing Wang³, Feng Zhang⁴, Yining Shi⁴, Nan Hao⁵, and Mark Wenig⁶

¹Remote Sensing Technology Institute (IMF), German Aerospace Center (DLR), Oberpfaffenhofen, Germany

²School of Atmospheric Sciences, Nanjing University, Nanjing, China

³Key Laboratory for Aerosol-Cloud-Precipitation of China Meteorological Administration, Nanjing University of Information Science and Technology, Nanjing, China

⁴Key Laboratory of Meteorological Disaster, Ministry of Education, Nanjing University of Information Science and Technology, Nanjing, China

⁵European Organisation for the Exploitation of Meteorological Satellites (EUMETSAT), Darmstadt, Germany

⁶Meteorological Institute, Ludwig-Maximilians-Universität München, München, Germany

Correspondence: Ka Lok Chan (ka.chan@dlr.de)

Abstract.

In this paper, we present long term observations of atmospheric nitrogen dioxide (NO₂) and formaldehyde (HCHO) in Nanjing using a Multi-AXis Differential Optical Absorption Spectroscopy (MAX-DOAS) instrument. Ground based MAX-DOAS measurements were performed from April 2013 to February 2017. The MAX-DOAS measurements of NO₂ and HCHO vertical column densities (VCDs) are used to validate OMI satellite observations over Nanjing. The comparison shows that the OMI observations of NO₂ correlate well with the MAX-DOAS data with Pearson correlation coefficient (R) of 0.91. However, OMI observations are on average a factor of 3 lower than the MAX-DOAS measurements. Replacing the a priori NO₂ profiles by the MAX-DOAS profiles in the OMI NO₂ VCD retrieval would increase the OMI NO₂ VCDs by $\sim 30\%$ with correlation nearly unchanged. The comparison result of MAX-DOAS and OMI observations of HCHO VCD shows a good agreement with R of 0.75 and the slope of the regression line is 0.99. An age weighted backward propagation approach is applied to the MAX-DOAS measurements of NO₂ and HCHO to reconstruct the spatial distribution of NO₂ and HCHO over Yangtze River Delta during summer and winter time. The reconstructed NO₂ fields show a distinct agreement with OMI satellite observations. However, due to the short atmospheric lifetime of HCHO, the backward propagated HCHO data does not show a strong spatial correlation with the OMI HCHO observations. The result shows the MAX-DOAS measurements are sensitive to the air pollution transportation in the Yangtze River Delta, indicating the air quality in Nanjing is significantly influenced by regional transportation of air pollutants. The MAX-DOAS data are also used to evaluate the effectiveness of air pollution control measures implemented during the Youth Olympic Games 2014. The MAX-DOAS data show a significant reduction of ambient aerosol, NO₂ and HCHO (30% - 50%) during the Youth Olympic Games. Our results provide a better understanding of the transportation and sources of pollutants in over the Yangtze River Delta as well as the effect of emission control measures during large international event, which are important for the future design of air pollution control policies.

1 Introduction

Nitrogen dioxide (NO_2) and formaldehyde (HCHO) are major atmospheric pollutants playing crucial roles in atmospheric chemistry. NO_2 is a catalyst for ozone (O_3) formation in the troposphere, while also participating in the catalytic destruction of stratospheric O_3 (Crutzen, 1970). Major NO_2 sources include fossil fuel combustion, biomass burning, lightning and oxidation of ammonia (Bond et al., 2001; Zhang et al., 2003). The emissions of NO_2 show a significant increasing trend in China due to the rapid industrialization and economy development in the last two decades (Zhang et al., 2007; van der A et al., 2008; Zhao et al., 2013), making it one of the most severe air pollution problems. HCHO is an intermediate product of the oxidation of almost all volatile organic compounds (VOCs). Therefore, it is widely used as an indicator of non methane volatile organic compounds (NMVOCs) (Fried et al., 2011). VOCs also have significant impacts on the abundance of hydroxyl (OH) radicals in the atmosphere, which is the major oxidant in the tropospheric. Major HCHO sources over the continents include the oxidation of VOCs emitted from plants, biomass burning, traffic and industrial emissions. Oxidation of methane (CH_4) emitted from the ocean is the main source of HCHO over water. Both NO_2 and HCHO contribute to the formation of secondary aerosols (Jang and Kamens, 2001) and they are toxic to human in high concentration. The spatial distribution of NO_2 and HCHO is strongly related to their emissions due to their short atmospheric lifetime. Consequently, it is important to understand the spatial and temporal variations of atmospheric NO_2 and HCHO for better air pollution management and control.

Multi-AXis Differential Optical Absorption Spectroscopy (MAX-DOAS) is a powerful remote sensing measurement technique which provides valuable vertical distribution information of atmospheric aerosols and trace gases (Platt and Stutz, 2008). Information of tropospheric aerosols and trace gases are obtained from the molecular absorption in the ultraviolet and visible spectral bands by applying the differential optical absorption spectroscopy (DOAS) technique to the observations of scattered sun light spectrum in several different viewing directions. As the experimental setup of MAX-DOAS is rather simple and inexpensive, it has been widely used for the observation of atmospheric aerosols and trace gases in the past decade (Hönninger and Platt, 2002; Hönninger et al., 2004; Wittrock et al., 2004; Frieß et al., 2006; Irie et al., 2008; Li et al., 2010; Clémer et al., 2010; Halla et al., 2011; Li et al., 2013; Ma et al., 2013; Chan et al., 2015; Jin et al., 2016; Wang et al., 2016; Chan et al., 2018). MAX-DOAS measurements are highly sensitive to aerosols and trace gases in the lower troposphere and provide valuable information of the vertical distribution of aerosol extinction and trace gases. The information of the aerosol and trace gas vertical profiles are particularly important for the study of the physical and chemical processes in the atmosphere.

Satellite based remote sensing measurements provide indispensable spatial information of air pollutants (Burrows et al., 1999; Bovensmann et al., 1999; Callies et al., 2000; Levelt et al., 2006). Trace gas columns are derived from the satellite observations of Earth's reflected solar spectrum for the investigation of atmospheric dynamics and emissions from both anthropogenic and natural sources (Beirle et al., 2003; Wenig et al., 2003; Beirle et al., 2004; Richter et al., 2005; Zhang et al., 2007; van der A et al., 2008). Satellite measurements can also be used to determine the effectiveness of emission control measures (Mijling et al., 2009; Witte et al., 2009; Wu et al., 2013; Chan et al., 2015). However, the uncertainties of satellite trace gas column retrieval are strongly dependent on the accuracy of the assumptions of trace gas vertical distributions. In addition, the temporal resolution of satellite measurements is often limited to single observation per day. Therefore, it is useful to compare

and integrate ground based and satellite observations for the interpretation of the spatial and temporal variation of NO₂ and HCHO.

Nanjing is the second largest city in the Yangtze River Delta and Eastern China. It is also the provincial capital of Jiangsu province. The population of Nanjing is about 8 million. Yangtze River is running through the city of Nanjing making it the largest inland port in China. Industrial, water and road transportation are the major anthropogenic sources of air pollutions in Nanjing. Due to its rapid development as well as its surrounding cities in the Yangtze River Delta, Nanjing is facing a series of air pollution problems in recent years. In addition, Nanjing has also hosted several important international events including the summer Youth Olympic Games in 2014. Therefore, it is important to have a better understanding of the air pollution sources in order to support the design of air quality related environmental policies in the future.

In this paper, we present long term MAX-DOAS observations of NO₂ and HCHO in Nanjing. Ground based MAX-DOAS measurements were performed from April 2013 to February 2017. Details of the MAX-DOAS experimental setup, the spectral analysis as well as the retrieval of the aerosol extinction, NO₂ and HCHO profiles are presented in section 2. Section 3.1 shows the validation of MAX-DOAS aerosol observations by comparing the reported aerosol optical depths (AODs) to sun photometer measurements. The comparison of NO₂ and HCHO VCDs measured by the MAX-DOAS and OMI satellite is presented in Section 3.2. An analysis of regional transport of pollutants over Yangtze River Delta is shown in section 3.4. In section 3.5, we presented an evaluation of the pollution reduction observed during the Youth Olympic Games in 2014.

2 Methodology

2.1 MAX-DOAS measurements

2.1.1 Experimental setup

A MAX-DOAS instrument was set up at a meteorological station of Nanjing University (32.12°N, 118.95°E) which is located on a small hill in the University campus at about 45 m above sea level. The meteorological station is located about 17 km northeast of the Nanjing city center and about 5 km south of the Yangtze River. The MAX-DOAS instrument for scattered sun-light measurements consists of a scanning telescope, a stepping motor controlling the viewing zenith angle of the telescope and a spectrometer. Scattered sun-light collected by the telescope is redirected by a quartz fiber to the spectrometer for spectral analysis. The field of view of the telescope is about 0.6°. An Ocean Optic USB2000 spectrometer equipped with a Sony ILX511 charge-coupled device (CCD) detector is used to cover the wavelength range from 288 nm to 434 nm. The full width half maximum (FWHM) spectral resolution of the spectrometer is 0.6 nm (at 360 nm).

A complete measurement cycle consists of scattered sun-light observations at elevation angle (α) of 1°, 2°, 3°, 6°, 10°, 18°, 30° and the zenith (90°). The viewing azimuth angle is adjusted to 320° (northwest). The exposure time and the number of scan of each measurement is adjusted automatically depending on the received intensity of the scattered sun-light spectrum in order to achieve a similar intensity level for all the measurements. A full measurement sequence takes about 5 - 10 minutes.

2.1.2 Spectral retrieval




Figure 1. An example of the DOAS retrieval of NO₂ and HCHO DSCDs from a MAX-DOAS spectrum taken on 12th May 2013 at 08:48 (local time) with viewing elevation angle of 1°. The left panels show the DOAS fit in the wavelength range of 338 - 370 nm, while the right panels show DOAS fit in the wavelength range of 324.5 - 359 nm.

All the measurement spectra were first corrected for offset and dark current and then analyzed using the spectral analyzing software QDOAS version 3.2 (<http://uv-vis.aeronomie.be/software/QDOAS/>). The MAX-DOAS spectral fit was performed at 2

different wavelength ranges of 338 - 370 nm for NO₂ retrieval and 324.5 - 359 nm for HCHO retrieval. In this study, the zenith spectrum ($\alpha=90^\circ$) taken in the same measurement cycle is used as reference spectrum in the spectral analysis to retrieve the differential slant column densities (DSCDs). The differential slant column density (DSCD) is defined as the difference between the slant column density (SCD) of the measured spectrum and the corresponding zenith reference spectrum. The broad band spectral structures caused by Rayleigh and Mie scattering are removed by including a low order polynomial in the DOAS fit. Absorption cross section of several trace gases were included in the DOAS fit at different fitting ranges, details of the DOAS retrieval settings for each wavelength range are listed in Table 1. The DOAS fit settings follow the one used in the Quality Assurance for Essential Climate Variables (QA4ECV) project (<http://www.qa4ecv.eu/>). These settings have also been adopted for the Second Cabauw Intercomparison campaign for Nitrogen Dioxide measuring Instruments (CINDI-2) campaign (<http://www.tropomi.eu/data-products/cindi-2>). Small shift and squeeze of the wavelengths are allowed in the wavelength mapping process in order to compensate small uncertainties caused by the instability of the spectrograph. Figure 1 shows an example of the spectral fit retrieval of NO₂ and HCHO DSCDs from a MAX-DOAS spectrum taken on 12th May 2013 at 08:48 (local time) with viewing elevation angle of 1°.

Table 1. The DOAS retrieval settings for different wavelength bands.

Species	Temperature	Wavelength Range		Reference
		324.5 - 359 nm	338 - 370 nm	
BrO	223 K	✓	✓	Fleischmann et al. (2003)
HCHO	298 K	✓	✓	Meller and Moortgat (2000)
NO ₂ ^(a)	298 K	✓	✓	Vandaele et al. (1998)
NO ₂ ^(a)	220 K		✓	Vandaele et al. (1998)
O ₃ ^(b)	223 K	✓	✓	Serdyuchenko et al. (2014)
O ₃ ^(b)	243 K	✓	✓	Serdyuchenko et al. (2014)
O ₄	293 K	✓	✓	Thalman and Volkamer (2013)
Ring		✓	✓	
Polynomial		5 th order	5 th order	

^(a) I₀ correction is applied with SCD of 10¹⁷ molec/cm² (Aliwell et al., 2002).

^(b) I₀ correction is applied with SCD of 10²⁰ molec/cm² (Aliwell et al., 2002).

As clouds are not included in the radiative transfer calculation of the aerosol and trace gas profile retrieval, the retrieval result can be significantly influenced by the present of cloud in the atmosphere. Therefore, the retrieved DSCDs were first filtered by removing cloudy scenes before proceeding to the aerosol and trace gas profile retrieval. The vertical profile of the oxygen collision complex O₄ only varies in a small range with atmospheric pressure and temperature, the retrieved O₄ DSCDs and (relative) intensities are assumed to vary smoothly with time with the solar and viewing geometry. Rapid change of O₄ DSCDs and intensities indicates a sudden change in the radiative transport condition which is likely due to the present of clouds. Therefore, we applied a locally weighted regression smoothing filter (LOWESS) (Cleveland, 1981) with a regression window

of 3 hours to the O₄ DSCDs and intensities time series at each elevation angle to filter data influenced by inhomogeneous and/or rapid changes of radiation transport conditions. Data with fast varying O₄ DSCDs and intensities were filter out. Only data with slow varying O₄ DSCDs and intensities were used in the aerosol and trace gas retrieval. The limitation of this cloud screening algorithm is that the algorithm is not able to distinguish continuous and homogeneous cloud condition. However, it is rare that the cloud does not change for a long time (within an hour) and the cloud layer is homogeneous for all viewing directions.

2.1.3 Aerosol and trace gas profile retrieval

Previous studies show that there is a systematic discrepancy between observation and model simulation of O₄ DSCDs (Wagner et al., 2009; Cl mer et al., 2010; Wagner et al., 2011; Chan et al., 2015; Wang et al., 2016; Chan et al., 2018; Zhang et al., 2018). A scaling correction factor has to be applied to the measured O₄ DSCDs in order to bring measurement and model results into agreement. However, the physical meaning of this observation is still not well understood (Ortega et al., 2016; Wagner et al., 2019). Theoretically, the optical path should be the longest under aerosol free condition for off zenith measurement. Thus, the MAX-DOAS measurement of O₄ DSCDs should be smaller than the one simulated with pure Rayleigh atmosphere. In this study, we compared the forward simulation of O₄ DSCDs under aerosol free condition to the O₄ DSCDs retrieved from the MAX-DOAS observations to examine the necessity of O₄ correction. Our result shows that the measured O₄ DSCDs occasionally exceeded the forward simulation results which implies that correction of O₄ DSCDs is necessary. The ratio between simulated and measured O₄ DSCD varies from 0.75 to 1.0 for cases where the measured O₄ DSCDs exceeded the forward simulation results. Our finding agrees with previous reported scaling correction factors of O₄ DSCDs which is ranging from 0.7 up to 1.0. In order to avoid over-correction due to the outliers, we take the 10th percentile instead of the minimum value of the simulated and measured O₄ DSCD ratio as the correction factor which is ~0.8. All MAX-DOAS observations of O₄ DSCD are corrected by multiplying the correction factor of 0.8. From hereafter, all O₄ DSCDs are referring to the corrected O₄ DSCDs.

The conversion of the MAX-DOAS observations to aerosol extinction and trace gas profile requires inversion of the underlying radiative transfer equations. These equations cannot be linearized, therefore, it is suggested to fit the measurement quantities to the forward calculation of radiation transfer (Wagner et al., 2004; H nninger et al., 2004; Sinreich et al., 2005; Frie  et al., 2006; Hartl and Wenig, 2013; Chan et al., 2018). As the vertical distribution profile of O₄ is very stable and it has several absorption bands in the ultraviolet and visible spectral range, it is commonly used as fitting quantity for the aerosols retrieval.

In this study, aerosol vertical profiles are retrieved at the 360 nm O₄ absorption band using the Munich Multiple wavelength MAX-DOAS retrieval algorithm (M³) (Chan et al., 2018). The algorithm is developed based on the optimal estimation method (Rodgers, 2000) and utilizes the Library for Radiative Transfer (LibRadTran) model (Emde et al., 2016) as the forward model. In this study, the U.S. standard mid-latitude atmosphere profiles for winter (January) and summer (July) are temporally interpolated to each month of the year for the radiative transfer simulations. A brief description of the aerosol and trace gases

vertical profile retrieval algorithm and the parameterization used in this study are presented in the following. A more detailed description of the 3M retrieval algorithm can be found in Chan et al. (2018).

In this study, all valid MAX-DOAS observations within a single measurement cycle are grouped together for the aerosol vertical profile retrieval. Assuming the set of measurement can be reproduced by the forward model and the forward model results are dependent on the aerosol extinction profile, we can then retrieve the aerosol extinction profile by fitting the forward model results to the MAX-DOAS O_4 observations using the iterative Newton-Gauss method. As the information contained in the MAX-DOAS observations are most likely not enough to retrieve a unique aerosol extinction profile, we have to supply the necessary information to the aerosol inversion in a form of a priori aerosol profile. As the aerosol load in Nanjing varies in a wide range, using a fix a priori could result in over regularizing the retrieval under higher aerosol load conditions. Therefore, we have implemented an iterative approach to avoid over regularizing the retrieval. We first use a fixed initial a priori to retrieve the aerosol extinction profile. The fixed a priori profile is then scaled to have the same aerosol optical depth retrieved from the previous run. Then the new a priori is used in the next run to retrieve the aerosol extinction profile. This procedure is repeated until the difference between the retrieved and a priori aerosol optical depth is less than 10 % or the number of iteration reaches the limit.

As aerosols are typically emitted and formed close to the surface in urban areas, we assume the a priori aerosol extinction profile follows an exponentially decreasing function with a scale height of 1.0 km. The aerosol optical depth (AOD) of the a priori aerosol profile is set to 0.5 for the retrieval at 360 nm. The uncertainties of the a priori aerosol profile are set to 100 % and the correlation length of the aerosol inversion is assumed to be 0.5 km. As MAX-DOAS measurements are more sensitive to the aerosol and trace gases close to the instrument, therefore, we divided the lowest 3.0 km of the troposphere into 15 layers with thickness of each layer of 200 m. A fix set of single scattering albedo of 0.92, asymmetry parameter of 0.68 and ground albedo of 0.04 is assumed in the radiative transfer calculations. An example of aerosol extinction profile retrieval is indicated in Figure 2a-c.

The aerosol information obtained from the procedure described above were used for the differential box air mass factor (Δ DAMF) calculation for the trace gas profile inversion. The Δ DAMFs were calculated at a single wavelength for the retrieval of trace gas profile using the radiative transfer model LibRadTran with the Monte Carlo simulation module MYSTIC (Emde et al., 2016). The Δ DAMF was assumed to be constant within the DOAS spectral fitting windows. In this study, vertical distribution profiles of NO_2 are retrieved at 360 nm. An example of NO_2 profile retrieval is indicated in Figure 2d-f. For HCHO vertical distribution profile retrieval, aerosol extinction profiles obtained at the 360 nm O_4 band are converted to 340 nm assuming a fixed Ångström coefficient (Ångström, 1929) of 1. The Ångström coefficient can varies significantly with time. Sun photometer measurements in Nanjing (see Section 2.2) show that the Ångström coefficient mostly varies between 0.7 (10th percentile) and 1.4 (90th percentile). Based on the sun photometer observations, we estimated the error caused by fixed Ångström coefficient on aerosol extinction is less than 2 %. The Δ DAMFs are subsequently calculated at 340 nm for the HCHO profile retrieval. An example of HCHO profile retrieval is indicated in Figure 2g-i.

The atmosphere layer settings of the trace gas profile retrieval are the same as the one used in the aerosol profile retrieval. NO_2 and HCHO concentrations above the retrieval height (3 km) are assumed to follow the U.S. standard atmosphere. As the




Figure 2. An example of vertical profile retrieval of aerosol extinction, NO₂ and HCHO from MAX-DOAS measurements taken on 4th July 2013 at around 8:47 (local time). The left panels show the (a) aerosol extinction, (d) NO₂ and (g) HCHO profiles. Panels in the middle column indicate the average kernel of (b) aerosol extinction, (e) NO₂ and (h) HCHO profile retrieval. The measured (blue curves) and modeled (green curves) DSCDs of (c) O₄, (f) NO₂ and (i) HCHO are shown in the right column.

trace gas profile cannot be fully reconstructed by the small number of measurements, therefore, we use the optimal estimation method (Rodgers, 2000) with iterative approach to regularize the inversion (Chan et al., 2015, 2018). The first step uses a

fixed initial a priori to retrieve the trace gas distribution. The fixed a priori profile is then scaled to have the vertical column retrieved from the first run. The scaled a priori is used in the next run to retrieve the trace gas profile. The process repeats until the difference between the retrieved and a priori trace gas vertical column is less than 10 % or the number of iteration reaches the limit. In this study, the a priori is assumed to follow an exponential decrease function with a scale height of 1.0 km. The uncertainty of the a priori profile is set to 100 % of the a priori and correlation length is set to 0.5 km in the trace gas profile inversion. The NO₂ vertical column density (VCD) of the a priori is set to 2×10^{16} molec/cm² while the a priori HCHO VCD is set to 1×10^{16} molec/cm².

2.2 Sun photometer measurements

A sun photometer was installed on the roof of a building of Nanjing University of Information Science and Technology (32.20°N, 118.70°E), which is located at the north bank of the Yangtze River. The sun photometer site is ~25 km northwest of the MAX-DOAS measurement site. The two instruments are separated by the Yangtze River. An industrial park is about 3 km east of the sun photometer site. Heavy industry factories, i.e., steel and cement plants are located in the industrial park. In this study, AODs measured by the sun photometer were used for the inter-comparison study to validate the MAX-DOAS aerosol retrieval. Cloud screened data were used. The data consist of AOD measurements at 7 different wavelength channels from 340 nm to 1020 nm. AOD measurements at 340 nm and 380 nm are interpolated to 360 nm for comparison.

2.3 Meteorological data

Meteorological parameters such as temperature, wind speed and wind direction are taken from a weather station in Nanjing which is located about 20 km south of the MAX-DOAS measurement site. The weather station is operated by the National Meteorological Center of China Meteorological Administration. These data are available on the webpage of China Meteorological Administration (<https://data.cma.cn/site/index.html>). The meteorological data are mainly used for the analysis of meteorological effects on air quality in Nanjing during the Youth Olympic Games in 2014.

2.4 OMI satellite observations

The Ozone Monitoring Instrument (OMI) is a passive nadir-viewing satellite borne imaging spectrometer (Levelt et al., 2006) on board the Earth Observing System's (EOS) Aura satellite. The EOS Aura satellite orbit at an altitude of ~710 km with a local equator overpass time of 13:45 LT (local time). The instrument consists of two charge-coupled devices (CCDs) covering a wavelength range from 264 nm to 504 nm. A scan provides measurements at 60 positions across the orbital track covering a swath of approximately 2600 km. The spatial resolution of OMI varies from ~320 km² (at nadir) to ~6400 km² (at both edges of the swath). The instrument scans along 14.5 sun-synchronous polar orbits per day providing daily global coverage.

The NASA's OMI NO₂ and HCHO standard product version 3 (Krotkov et al., 2017; González Abad et al., 2015) are used in this study. In the NO₂ product, the slant column densities (SCDs) of NO₂ are derived from Earth's reflected spectra in the visible range (402 - 465 nm) using an iterative sequential algorithm (Marchenko et al., 2015). Previous studies show the updated

SCDs are on average 10-40% lower compared to the previous version of retrieval (Marchenko et al., 2015). The OMI NO₂ SCDs are converted to VCDs by using the concept of air mass factor (AMF) (Solomon et al., 1987). The AMFs are calculated using NO₂ profile simulated by the Global Modeling Initiative (GMI) chemistry transport model. The horizontal resolution of GMI is 1° (latitude) × 1.25° (longitude) (Rotman et al., 2001). Separation of stratospheric and tropospheric columns is achieved by the local analysis of the stratospheric field over unpolluted areas (Bucsela et al., 2013).

The OMI HCHO retrieval algorithm uses the direct fit of radiances in the spectral range from 328.5 nm to 356.5 nm for the SCD retrieval. In the current version of HCHO product, OMI radiance measurement over the remote Pacific is used as reference in the fitting process. This approach is reported to reduce the interferences from unresolved spectral structures in the retrieval of weak absorbers like HCHO (De Smedt et al., 2018). The retrieved SCDs are then converted to VCDs using the AMF approach. The AMFs are calculated based on climatological HCHO profiles.

2.5 Back trajectory modeling

The National Oceanic and Atmospheric Administration Air Resources Laboratory (NOAA ARL) developed HYbrid Single Particle Lagrangian Integrated Trajectory (HYSPLIT) model (Stein et al., 2015) (<http://www.arl.noaa.gov/HYSPLIT.php>) was used to investigate the transportation pollutants over Yangtze River Delta. Meteorological data from the Global Data Assimilation System (GDAS) with a spatial resolution of 0.5° × 0.5° and 24 vertical levels was used in the model for trajectory simulations. Backward trajectories are computed for air masses arriving at the mid point of each MAX-DOAS retrieval layer.

3 Results and discussions

3.1 Comparison of MAX-DOAS and sun photometer AODs

Aerosol optical depths (AODs) retrieved from the MAX-DOAS observations are compared to the sun photometer measurements. As the sampling resolution of the MAX-DOAS and the sun photometer are different, individual measurements are averaged to hourly, daily and monthly values for comparison. Figure 3 shows the scatter plot of AOD measured by the sun photometer and MAX-DOAS. Both datasets are cloud filtered. AODs measured by both MAX-DOAS and sun photometer are in general in good agreement. The Pearson correlation coefficient (R) of daily averaged data is 0.73. The slope of the total least squares regression between the two datasets is 0.56 with an offset of 0.13. The discrepancies between the two datasets can be explained by the difference of measurement technique and measurement location. The sun photometer derives AOD from direct sun measurements, while the MAX-DOAS retrieves AOD using the O₄ absorption information from scattered sun light. The sun photometer is sensitive to the entire column while the MAX-DOAS is mostly sensitive to aerosol at the lowest few kilometers of the troposphere. Therefore, the MAX-DOAS is likely to underestimate the AOD when there is an elevated aerosol layer in the upper troposphere. This explanation matches with the observations that the MAX-DOAS agrees better with the sun photometer under low aerosol conditions (AOD < 1) and underestimate the AOD during high AOD conditions. Another source of error is the assumption of aerosol optical properties in the MAX-DOAS retrieval. In order to quantify the error caused by




Figure 3. Comparison of AOD measured by the MAX-DOAS and sunphotometer. The total least squares regression line is calculated based on the daily data.

the fixed asymmetry parameter and single scatter albedo used in the aerosol retrieval, we have performed a sensitivity analysis by retrieving the aerosol profile with different asymmetry parameter and single scattering albedo. As the sun photometer in Nanjing does not provide full inversion of aerosol optical properties, aerosol optical properties are taken from the Taihu AERONET station which is about 150 km southeast of Nanjing. The 10th and 90th percentile of asymmetry parameter and single scatter albedo are used for the sensitivity study. The result shows that the error caused by the fixed asymmetry parameter on AOD retrieval can be up to 4 %, which the single scattering albedo shows a smaller impact of up to 1.5 %. In addition, the MAX-DOAS and sun photometer are separated by a distance of ~ 25 km and the sun photometer site is very close to an industrial park, where the heavy polluted industries are located, i.e., steel and cement plants. Therefore, the sun photometer measurements are expect to be higher than the MAX-DOAS observations. Accounting for these effects, the MAX-DOAS and sun photometer observation agrees well with each other and the MAX-DOAS derived aerosol profiles are reliable for the retrieval of NO₂ and HCHO profiles.

3.2 Comparison of MAX-DOAS and OMI observations

Vertical column densities of NO₂ and HCHO retrieved from the ground based MAX-DOAS measurements are compared to OMI satellite observations. MAX-DOAS data are temporally averaged around the OMI overpass time from 12:00 to 14:00




Figure 4. (a) Time series of the MAX-DOAS (red markers) and NASA’s OMI (blue markers) tropospheric NO_2 VCDs over Nanjing from 2013 to 2017. MAX-DOAS data are temporally averaged around the OMI overpass time. OMI measurements are spatially averaged for pixels within 20 km of the MAX-DOAS measurement site. OMI NO_2 VCDs retrieved using MAX-DOAS profile as a priori information are indicated as green markers. Tropospheric NO_2 VCDs taken from KNMI product are also shown as gray marker for reference. (b) Comparison of tropospheric NO_2 VCDs between the MAX-DOAS and OMI satellite observations. The total least squares regression line is calculated based on the monthly average data.




Figure 5. (a) Time series of the MAX-DOAS (red markers) and OMI (blue markers) HCHO VCDs over Nanjing from 2013 to 2017. MAX-DOAS data are temporally averaged around the OMI overpass time. OMI measurements are spatially averaged for pixels within 20 km of the MAX-DOAS measurement site. OMI NO_2 VCDs retrieved using MAX-DOAS profile as a priori information are indicated as green markers. (b) Comparison of HCHO VCDs between the MAX-DOAS and OMI satellite observations. The total least squares regression line is calculated based on the monthly average data.

(local time), while OMI VCDs are spatially averaged for pixels within 20 km of the MAX-DOAS measurement site. Time series of MAX-DOAS and OMI observations of NO₂ and HCHO VCDs are shown in Figure 4 and Figure 5, respectively. Both OMI NO₂ and HCHO datasets are filtered for cloud radiance fraction smaller than 0.4. Daily and monthly averaged data are shown. Missing data for some months are due to cloud filtering or instrumental issues of the MAX-DOAS.

5 NO₂ VCDs measured by the MAX-DOAS and the OMI satellite shows a similar seasonal pattern with higher NO₂ columns during winter and lower NO₂ values over summer. Higher NO₂ levels are due longer atmospheric lifetime of NO₂ during winter and higher emissions in winter, e.g., higher power consumption and emissions from individual domestic heating. MAX-DOAS and OMI observations show good temporal consistency with each other with Pearson correlation coefficient (*R*) of 0.91. Despite the strong correlation between the two datasets, the OMI observations is systematically underestimating the NO₂ columns. The slope and offset of the total least squares regression between the two datasets is 0.50 and -1.89×10^{15} molec/cm², respectively. Averaged OMI NO₂ column over Nanjing is 5.67×10^{15} molec/cm² which is about 60 % lower than the MAX-DOAS averaged VCD of 14.96×10^{15} molec/cm² (see Figure 4). The underestimation of NO₂ VCDs from OMI measurements agree with previous studies in the Yangtze Delta (Chan et al., 2015). The discrepancies can be related to the differences in spatial coverage between the ground based and satellite observations and the uncertainties related to the NO₂ vertical distribution profile shape used in the air mass factor calculation. We quantified the influence of NO₂ vertical distribution profile on the air mass factor calculation by recomputed the OMI tropospheric NO₂ air mass factor using the MAX-DOAS NO₂ profile as a priori information. OMI NO₂ VCDs retrieved with MAX-DOAS NO₂ profiles are also indicated in Figure 4. Using the MAX-DOAS profile as a priori information in the OMI NO₂ VCDs retrieval on average increased the OMI NO₂ VCDs by ~30 % with correlation nearly unchanged. The averaged OMI NO₂ column over Nanjing increased to 7.29×10^{15} molec/cm². In addition, the slope of the regression line increases from 0.50 to 0.57 while the offset reduces from -1.89 to -1.35×10^{15} molec/cm². The result indicates that a better estimation of NO₂ vertical distribution in the satellite VCD retrieval reduces the discrepancies. Our findings agree with previous studies that using better estimated NO₂ vertical profile in the satellite air mass factor calculation enhanced the OMI NO₂ columns and reduced discrepancies between OMI and ground based observations (Chan et al., 2012; Lin et al., 2014; Chan et al., 2015; Wang et al., 2017). However, the improved OMI NO₂ VCDs are still about 50 % lower than the MAX-DOAS observations indicating that there are still remaining issues. Previous study shows that the OMI NO₂ tropospheric vertical columns processed by the Royal Netherlands Meteorological Institute (KNMI) (DOMINO version 2.0) (Boersma et al., 2011) only underestimated the tropospheric NO₂ VCDs by ~20 % compared to ground based MAX-DOAS observations in Wuxi (another city in Yangtze River Delta) (Wang et al., 2017). In order to investigate the differences between different OMI NO₂ products, we have also plotted the KNMI OMI NO₂ product in Figure 4. The average NO₂ column reported from the KMI product is 10.10×10^{15} molec/cm², which is about a factor of 2 higher than the NASA product. The result shows the KNMI product only underestimate the NO₂ columns in Nanjing by ~30 % which is consistent with previous study in Yangtze River Delta (Wang et al., 2017). The difference between the two OMI products can be related to different spectral analysis techniques, stratospheric tropospheric separation algorithms as well as a priori profiles used in the retrievals. Investigation of the differences between the two algorithms is however beyond the scope of this study. Another source of discrepancy is related to the difference in spatial coverage of OMI and MAX-DOAS observations. The spatial coverage of

the OMI measurement footprint is ranging from 330 km² up to 4600 km² with an average of 920 km². Measurements with such large spatial coverage are probably difficult to capture the spatial gradient of NO₂ and resulted in an underestimation over pollution hot spots due to the averaging of large OMI footprint. This effect is especially significant over Nanjing, as it is a local pollution hot spot surrounded by rather clean areas. The underestimation of NO₂ columns is coherent with previous measurements over urban pollution hot spots, i.e., Washington DC and Shanghai (Wenig et al., 2008; Chan et al., 2015). In addition, previous measurements over suburban area in Shanghai show better agreement with OMI observations compared to the measurements in the city (Chan et al., 2015) indicating the effect of spatial inhomogeneity of NO₂ on the satellite data comparison. However, the impact of this effect is difficult to quantify due to lack of high spatial resolution data. Cloud contamination could be an important source of error in the satellite retrieval. Previous study show that the cloud effect on OMI NO₂ VCD retrieval is only significant for cloud radiance fraction > 0.4 (Wang et al., 2017). The OMI NO₂ data used in this study are filtered for cloud radiance fraction smaller than 0.4. Therefore, cloud contamination should only show a negligible effect in the comparison.

We have also compared the MAX-DOAS and OMI observations of HCHO and both datasets show a similar seasonal variation pattern. The seasonal variation pattern of HCHO is opposite to NO₂ with higher HCHO columns during summer and lower in winter. Higher HCHO levels in summer are related to the increases of biogenic emission of precursor VOCs from plants and higher oxidation rate of VOCs. The MAX-DOAS measurements of HCHO VCDs agree well with the OMI observations with Pearson correlation coefficient (*R*) of 0.75. In addition, the absolute value of the columns show a perfect agreement. The average HCHO VCD measured by the MAX-DOAS is 8.04×10^{15} molec/cm² while the averaged OMI HCHO VCD is 7.89×10^{15} molec/cm². The slope of the total least squares regression between the two datasets is 0.99 with an offset of 0.31×10^{15} molec/cm² (see Figure 4b). In order to quantify the influence of HCHO vertical distribution profile on the air mass factor calculation, we have recomputed the OMI HCHO VCDs using the MAX-DOAS HCHO profiles as a priori information. OMI HCHO VCDs retrieved with MAX-DOAS HCHO profiles are also indicated in Figure 4. Using the MAX-DOAS profile as a priori information in the OMI HCHO VCDs retrieval on average increased the OMI HCHO VCDs by ~25 % with correlation nearly unchanged. The averaged OMI HCHO column over Nanjing increased to 9.95×10^{15} molec/cm². The slope of the regression line also increases from 0.99 to 1.39 while the offset reduces from 0.31 to -0.72×10^{15} molec/cm². Higher HCHO VCDs retrieved from OMI using MAX-DOAS profiles as a priori is mainly due to the MAX-DOAS measurements are not sensitive to HCHO at higher altitudes. Sources of HCHO in the troposphere include the oxidation of various VOCs, including methane. Some of these VOCs have relatively long atmospheric lifetime, e.g., methane, therefore, they are rather well mixed in the atmospheric and resulting in a larger portion of HCHO in the upper troposphere. The MAX-DOAS could not capture HCHO at higher altitudes. Therefore, using the MAX-DOAS profiles for OMI HCHO retrieval is likely underestimated the air mass factors and overestimated the HCHO total columns.

3.3 Seasonal variation of aerosol, NO₂ and HCHO vertical profiles

The seasonal variations of pollutant are closely related to meteorological conditions as well as the characteristics of different emission sources. Analyzing the seasonal variation patterns of different atmospheric pollutants can provide further information




Figure 6. Vertical distribution of (a) aerosol, (b) NO₂ and (c) HCHO for different seasons. Shaded areas represent 1 σ standard deviation of variation.

on the emission source characteristic as well as the atmospheric processes. Figure 6 shows the vertical profiles of NO₂ and HCHO for all four seasons. Significant seasonal patterns are observed from the NO₂ and HCHO data. The NO₂ vertical profiles show higher NO₂ mixing ratios in autumn and winter and lower in spring and summer. NO₂ profiles from all seasons show that the NO₂ mixing ratios decrease with increasing altitude. The result indicates most of the measured NO₂ is produced close to the surface which agrees with the fact that traffic emission is one of the largest source of atmospheric NO₂ in Nanjing. HCHO measurements show a different seasonal variation pattern with lower values during winter and higher mixing ratios in summer. The seasonal pattern of HCHO indicates the significant contribution from biogenic emissions from vegetation. Although HCHO profiles for all seasons also show decreasing mixing ratio with height, the decreasing rate is in general much lower than that of NO₂. Larger fraction of HCHO is located at higher altitudes. This is probably related to the source characteristic of HCHO. Majority of the tropospheric HCHO is secondary produced from the oxidation of VOCs and resulting in a larger portion of HCHO in the upper altitudes.

3.4 Regional pollution transport

Air pollutants released to the atmosphere do not only cause local impacts, but also can influence regions far from the source through regional transport. In order to investigate the influences from regional transportation of air pollutants, we use the backward trajectories calculated by the HYSPLIT model to back correlate the possible source regions of the MAX-DOAS measured NO₂. Similar approaches have also been applied for other source appointment studies (Stohl et al., 2009, 2011;




Figure 7. The upper panels show spatial distribution of NO₂ in (a) summer and (b) winter reconstructed from the MAX-DOAS measurements using backward trajectories calculated by the HYSPLIT model. Backward trajectories are calculated over 6 h for summer measurements and 12 h for winter measurements. The lower panels indicates the average OMI NO₂ vertical column densities over the Yangtze River Delta during (c) summer and (d) winter.

Brunner et al., 2012, 2017; Wang et al., 2019). Backward trajectories are calculated at the center point of each MAX-DOAS retrieval layer below 2 km. The MAX-DOAS measurement of NO₂ partial columns are then assigned to the grid points along the backward trajectories. We assume the partial columns represent a 200 m thick layer of the assigned grid point and the vertical distribution of NO₂ in the assigned grid point is assumed to be homogeneous from the surface to 2 km above ground.

- 5 In this study, the backward propagated NO₂ data are binned in a resolution of $0.2^\circ \times 0.2^\circ$ grid. As NO₂ is a relatively short life pollutants in the atmosphere, it is less likely to stay in the atmospheric for a long time and being transported far from the sources. Data are more reliable with shorter atmospheric lifetime. In order to account for this effect, an age weighting factor (w_τ) is used in the backward propagation. The age weighting factor is defined by Eq. 1 where τ is the assumed lifetime of NO₂




Figure 8. The upper panels show spatial distribution of HCHO in (a) summer and (b) winter reconstructed from the MAX-DOAS measurements using backward trajectories calculated by the HYSPLIT model. Backward trajectories are calculated over 6 h for summer measurements and 12 h for winter measurements. The lower panels indicate the average OMI HCHO vertical column densities over the Yangtze River Delta during (c) summer and (d) winter.

and t represents the time for the air mass to travel to the measurement site. This weighting scheme is useful when multiple trajectories overlapping with each other within a single grid point.

$$w_{\tau} = e^{-\frac{t}{\tau}} \quad (1)$$

As the lifetime of NO_2 has a strong seasonal variability, we separate the measurements into summer (June, July and August) and winter (December, January and February) in our analysis. We assume the NO_2 lifetime in summer and winter is 3 h and 6 h, respectively. Backward trajectories are calculated over 6 h for summer measurements and 12 h for winter measurements. As the assumed lifetime is only used for the calculation of the age factor, the accuracy of the lifetime does not show a strong impact on the reconstructed spatial pattern. In order to get a balance between having better spatial coverage and the reliabil-

ity of the reconstructed pollution maps, these numbers are used in this study. NO₂ fields reconstructed from the MAX-DOAS measurements in summer and winter using backward trajectories calculated by the HYSPLIT model are shown in Figure 7a&b. Figure 7c&d shows the averaged OMI satellite observations of NO₂ during summer (June, July and August) and winter (December, January and February) from June 2013 to February 2017. The OMI measurements show that Nanjing has serious NO₂ pollution problem in winter.

The reconstructed NO₂ fields show a good spatial agreement with OMI satellite observations. However, the reconstructed NO₂ fields are on average 3 times higher than the OMI observations. As the MAX-DOAS NO₂ VCDs are on average 3 times higher than the OMI data, the large discrepancies of the absolute values between the reconstructed data and OMI observations are expected. Elevated NO₂ levels are observed during winter when air mass coming from the east, e.g., Shanghai, Suzhou and Nantong, where OMI observations also show high NO₂ levels. The summer time NO₂ levels are in general lower. However, elevated values can still be observed along the Yangtze River and over some cities, e.g., Nantong and Wuhu.

We have also applied the same approach to the MAX-DOAS HCHO measurements with HCHO lifetime of 1 h and 2 h for summer and winter measurements, respectively. HCHO fields reconstructed from the MAX-DOAS measurements are shown in Figure 8a&b. OMI observations of HCHO are also shown in Figure 8c&d for reference. Summer time HCHO data are in general higher than the observations during winter. However, the spatial distribution of the back propagated HCHO data does not show a strong correlation with the OMI satellite observations as the NO₂ data. This is probably due to HCHO is mostly secondary produced in the atmosphere. In addition, the atmospheric lifetime of HCHO is much shorter compared to NO₂ and therefore more dependent on the local productions. The result suggests that the MAX-DOAS measurements are sensitive to the regional transport of air pollutants. Despite the strong local contributions, the air quality of Nanjing can also be significantly influenced by the air pollution transportation, especially during winter.

3.5 Assessments of emission reduction during Youth Olympic

The summer Youth Olympic Games was held from 16 to 28 August 2014 in Nanjing, China. During the event, the government implemented a series of air pollution control measures. Heavy emission vehicles were strictly prohibited. Local heavy industries, e.g., petrochemical and steel industries were required to limit their production during the Youth Olympic Games. These air pollution control measures were often implemented when such an international events held in China (Wu et al., 2013; Wang et al., 2014; Chan et al., 2015; Liu et al., 2016; Sun et al., 2016). In order to study the influences of the emission control measures implemented during the Youth Olympic Games on the local air quality, we compare the MAX-DOAS observations of aerosol, NO₂ and HCHO taken before, during and after the Youth Olympic Games. We define the pre-Olympic period from a month before the Youth Olympic Games (16 July 2014) to 1 day before the Youth Olympic Games (15 August 2014). During the pre-Olympic period, the government gradually started to implement some of the emission control measures. During the Youth Olympic Games all those air pollution control measures were strictly implemented. The post-Olympic period is defined from 29 August 2014 to 28 September 2014 where the emission control was gradually getting back to normal.

Time series of aerosol optical depths, HCHO and NO₂ VCDs and surface mixing ratios measured by the MAX-DOAS around the Youth Olympic Games are shown in Figure 9. The gray shadowed area indicates the Youth Olympic Games period




Figure 9. Time series of (a) aerosol optical depths, (b) HCHO and NO₂ VCDs and (c) HCHO and NO₂ surface mixing ratios measured by the MAX-DOAS around the Youth Olympic Games. Hourly and daily averaged values are shown. The gray shadowed area indicates the Youth Olympic Games period (16 to 28 August 2014).

(16 to 28 August 2014). Meteorological parameters such as temperature, wind speed and wind direction measured by a weather station in Nanjing during the same period are shown in Figure 10. The AODs, HCHO and NO₂ measurements show a strong




Figure 10. (a) Time series of wind speed (blue curve) and wind direction (red curve). (b) Time series of daily minimum (blue curve), daily average (green curve) and daily maximum temperature. The gray shadowed area indicates the Youth Olympic Games period (16 to 28 August 2014).

temporal variability. In order to investigate the effect of the Youth Olympic Games, we analyzed the statistic of AODs, HCHO and NO₂ VCDs for the pre-Olympic, Olympic and post-Olympic periods. Boxplots of the AODs, HCHO and NO₂ VCDs for the three periods are shown in Figure 11. The results show that AOD, NO₂ and HCHO VCD are the lowest during the Olympic period among those 3 periods. The MAX-DOAS measurements of AOD reduced from 0.9 during the pre-Olympic period to 0.6 for the Olympic period and bounced back to about 0.8 after the Youth Olympic Games. A reduction of NO₂ columns can also be observed during the Youth Olympic Games. Averaged NO₂ VCDs are 14, 11 and 19 × 10¹⁵ molec/cm² for periods before, during and after the Youth Olympic Games, respectively. Similar reduction pattern is also observed from the NO₂ surface mixing ratios. Averaged NO₂ surface mixing ratios are 7.5, 4.7 and 9.0 ppbv for periods before, during and after the Youth Olympic Games, respectively. The HCHO columns measured during the Youth Olympic Games also decreased




Figure 11. Boxplots of (a) aerosol optical depths, (b) NO₂ and (c) HCHO VCDs before, during and after the Youth Olympic Games.

by more than $\sim 50\%$ from 17×10^{15} molec/cm² before the Olympic down to 8×10^{15} molec/cm² during the Youth Olympic Games. Surface HCHO mixing ratios also show similar reduction. The surface HCHO mixing ratios measured also reduced from 7.0 ppbv before the Youth Olympic to 3.3 ppbv during the Youth Olympic. As the meteorological conditions are very similar during the three periods, the decrease of pollutant concentrations are mainly attributed to the reduction of emissions during the Youth Olympic Games.

4 Summary and conclusions

In this paper, we present long term observations of atmospheric NO₂ and HCHO in Nanjing using a MAX-DOAS instrument. Ground based MAX-DOAS measurements were performed from April 2013 to February 2017. Differential slant columns of O₄, NO₂ and HCHO were retrieved by applying the differential optical absorption technique to the scattered sun light spectra at the UV band. The results are served as inputs for the retrieval of aerosol extinction, NO₂ and HCHO profiles in the lower troposphere.

Aerosol results derived from the MAX-DOAS observations are validated by comparing the AODs to sun photometer observations. The MAX-DOAS and sun photometer measurements show a good agreement with each other with Pearson correlation coefficient (R) of 0.73. Considering the differences in measurement technique and measurement location, we concluded that the MAX-DOAS aerosol measurements are reliable for the NO₂ and HCHO profile retrieval.

Tropospheric vertical column densities (VCDs) of NO₂ derived from MAX-DOAS measurements are used to validate OMI observations. The comparison shows that the OMI observations correlate well with the MAX-DOAS data with Pearson correlation coefficient (R) of 0.91. However, OMI observations are on average 60% lower than the MAX-DOAS measurements. Using the MAX-DOAS NO₂ profile as a priori information in the OMI retrieval on average increased the OMI NO₂ VCDs by $\sim 30\%$ with correlation nearly unchanged. However, the improved OMI NO₂ VCDs are still about 50% lower than the

MAX-DOAS observations. The KNMI OMI NO₂ product shows a better agreement with the MAX-DOAS observations with an underestimation of ~30%. Large difference between the two OMI product is related to the different spectral analysis techniques, stratospheric tropospheric separation algorithms as well as a priori profiles used in the retrievals. Another source of discrepancy between MAX-DOAS and OMI observations is related to the difference in spatial coverage between the two
5 measurements. We also compared the OMI observation of HCHO VCDs to our MAX-DOAS data. The result shows a good agreement between the two datasets with R of 0.75 and the slope of the regression line is 0.99.

The MAX-DOAS measurement of NO₂ and HCHO profiles are analyzed together with the backward trajectory simulations to assess the regional transportation and possible source regions of the MAX-DOAS measured NO₂ and HCHO. The age weighted backward propagation approach is used to reconstruct the spatial distribution of NO₂ and HCHO over the Yangtze
10 River Delta during summer and winter time. The reconstructed NO₂ fields show a distinct agreement with OMI satellite observations. The result shows the MAX-DOAS measurements are sensitive to the air pollution transportation in the Yangtze River Delta. However, due to the short atmospheric lifetime of HCHO, the backward propagated HCHO data does not show a strong spatial correlation with the OMI HCHO observations. Our result suggested the air quality of Nanjing are significantly influenced by air pollution transportation, especially during winter.

15 We have also used the MAX-DOAS observations of aerosol, NO₂ and HCHO for the investigation of the effectiveness of air pollution control measures implemented during the Youth Olympic Games 2014. Our results show a significant reduction (30% - 50%) of ambient aerosol, NO₂ and HCHO compared to measurements before and after the Youth Olympic games. The results indicate the effects of the reduction of emissions during the Youth Olympic Games. Our findings provide a better understanding of the transportation and sources of pollutants in the Yangtze River Delta as well as the effects of emission
20 control measures during large international event, which are important for the future design of air pollution control policies.

Author contributions. KLC, AD, KPH and NH designed the experiment. KLC, ZW, AD, YS and NH carried out the experiment. JW, FZ and YS provided auxiliary data. MW provided useful comments for the discussion. KLC analyzed the measurement data and prepared the manuscript with contributions from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

25 *Acknowledgements.* The authors would also like to thank the National Oceanic and Atmospheric Administration (NOAA) Air Resources Laboratory (ARL) for the provision of the HYSPLIT transport and dispersion model used in this publication. The work described in this paper was partly supported by the ESA-MOST Dragon 3 Cooperation Programme under the framework of the East Asian monsoon and air quality project (project ID: 10455).

References

- Aliwell, S. R., Roozendael, M. V., Johnston, P. V., Richter, A., Wagner, T., Arlander, D. W., Burrows, J. P., Fish, D. J., Jones, R. L., Tørnkvist, K. K., Lambert, J.-C., Pfeilsticker, K., and Pundt, I.: Analysis for BrO in zenith-sky spectra: An intercomparison exercise for analysis improvement, *Journal of Geophysical Research: Atmospheres*, 107, ACH 10–1–ACH 10–20, <https://doi.org/10.1029/2001JD000329>, 2002.
- 5 Ångström, A.: On the Atmospheric Transmission of Sun Radiation and on Dust in the Air, *Geografiska Annaler*, 11, 156–166, <https://doi.org/10.1080/20014422.1929.11880498>, 1929.
- Beirle, S., Platt, U., Wenig, M., and Wagner, T.: Weekly cycle of NO₂ by GOME measurements: a signature of anthropogenic sources, *Atmospheric Chemistry and Physics*, 3, 2225–2232, <https://doi.org/10.5194/acp-3-2225-2003>, 2003.
- Beirle, S., Platt, U., Wenig, M., and Wagner, T.: NO_x production by lightning estimated with GOME, *Advances in Space Research*, 34, 793–797, <https://doi.org/10.1016/j.asr.2003.07.069>, 2004.
- 10 Boersma, K. F., Eskes, H. J., Dirksen, R. J., van der A, R. J., Veefkind, J. P., Stammes, P., Huijnen, V., Kleipool, Q. L., Sneep, M., Claas, J., Leitão, J., Richter, A., Zhou, Y., and Brunner, D.: An improved tropospheric NO₂ column retrieval algorithm for the Ozone Monitoring Instrument, *Atmospheric Measurement Techniques*, 4, 1905–1928, <https://doi.org/10.5194/amt-4-1905-2011>, 2011.
- Bond, D. W., Zhang, R., Tie, X., Brasseur, G., Huffman, G., Orville, R. E., and Boccippio, D. J.: NO_x production by lightning over the continental United States, *Journal of Geophysical Research: Atmospheres*, 106, 27 701–27 710, <https://doi.org/10.1029/2000JD000191>, 2001.
- 15 Bovensmann, H., Burrows, J., Buchwitz, M., Frerick, J., Noël, S., Rozanov, V., Chance, K., and Goede, A.: SCIAMACHY: Mission objectives and measurement modes, *Journal of the Atmospheric Sciences*, 56, 127–150, 1999.
- Brunner, D., Henne, S., Keller, C. A., Reimann, S., Vollmer, M. K., O’Doherty, S., and Maione, M.: An extended Kalman-filter for regional scale inverse emission estimation, *Atmospheric Chemistry and Physics*, 12, 3455–3478, <https://doi.org/10.5194/acp-12-3455-2012>, 2012.
- 20 Brunner, D., Arnold, T., Henne, S., Manning, A., Thompson, R. L., Maione, M., O’Doherty, S., and Reimann, S.: Comparison of four inverse modelling systems applied to the estimation of HFC-125, HFC-134a, and SF₆ emissions over Europe, *Atmospheric Chemistry and Physics*, 17, 10 651–10 674, <https://doi.org/10.5194/acp-17-10651-2017>, 2017.
- Bucsela, E. J., Krotkov, N. A., Celarier, E. A., Lamsal, L. N., Swartz, W. H., Bhartia, P. K., Boersma, K. F., Veefkind, J. P., Gleason, J. F., and Pickering, K. E.: A new stratospheric and tropospheric NO₂ retrieval algorithm for nadir-viewing satellite instruments: applications to OMI, *Atmospheric Measurement Techniques*, 6, 2607–2626, <https://doi.org/10.5194/amt-6-2607-2013>, 2013.
- 25 Burrows, J. P., Weber, M., Buchwitz, M., Rozanov, V., Ladstätter-Weissenmayer, A., Richter, A., DeBeek, R., Hoogen, R., Bramstedt, K., Eichmann, K.-U., et al.: The global ozone monitoring experiment (GOME): Mission concept and first scientific results, *Journal of the Atmospheric Sciences*, 56, 151–175, 1999.
- 30 Callies, J., Corpaccioli, E., Eisinger, M., Hahne, A., and Lefebvre, A.: GOME-2-Metop’s second-generation sensor for operational ozone monitoring, *ESA bulletin*, 102, 28–36, 2000.
- Chan, K. L., Pöhler, D., Kuhlmann, G., Hartl, A., Platt, U., and Wenig, M. O.: NO₂ measurements in Hong Kong using LED based long path differential optical absorption spectroscopy, *Atmospheric Measurement Techniques*, 5, 901–912, <https://doi.org/10.5194/amt-5-901-2012>, 2012.
- 35 Chan, K. L., Hartl, A., Lam, Y. F., Xie, P. H., Liu, W. Q., Cheung, H. M., Lampel, J., Pöhler, D., Li, A., Xu, J., Zhou, H. J., Ning, Z., and Wenig, M.: Observations of tropospheric NO₂ using ground based MAX-DOAS and OMI measurements during the Shanghai World Expo 2010, *Atmospheric Environment*, 119, 45–58, <https://doi.org/10.1016/j.atmosenv.2015.08.041>, 2015.

- Chan, K. L., Wiegner, M., Wenig, M., and Pöhler, D.: Observations of tropospheric aerosols and NO₂ in Hong Kong over 5 years using ground based MAX-DOAS, *Science of The Total Environment*, 619-620, 1545 – 1556, <https://doi.org/10.1016/j.scitotenv.2017.10.153>, 2018.
- Clémer, K., Van Roozendael, M., Fayt, C., Hendrick, F., Hermans, C., Pinardi, G., Spurr, R., Wang, P., and De Mazière, M.: Multiple wavelength retrieval of tropospheric aerosol optical properties from MAXDOAS measurements in Beijing, *Atmospheric Measurement Techniques*, 3, 863–878, <https://doi.org/10.5194/amt-3-863-2010>, 2010.
- Cleveland, W. S.: LOWESS: A Program for Smoothing Scatterplots by Robust Locally Weighted Regression, *The American Statistician*, 35, <https://doi.org/10.2307/2683591>, 1981.
- Crutzen, P. J.: The influence of nitrogen oxides on the atmospheric ozone content, *Quarterly Journal of the Royal Meteorological Society*, 96, 320–325, <https://doi.org/10.1002/qj.49709640815>, 1970.
- De Smedt, I., Theys, N., Yu, H., Danckaert, T., Lerot, C., Compernelle, S., Van Roozendael, M., Richter, A., Hilboll, A., Peters, E., Pedergnana, M., Loyola, D., Beirle, S., Wagner, T., Eskes, H., van Geffen, J., Boersma, K. F., and Veefkind, P.: Algorithm theoretical baseline for formaldehyde retrievals from S5P TROPOMI and from the QA4ECV project, *Atmospheric Measurement Techniques*, 11, 2395–2426, <https://doi.org/10.5194/amt-11-2395-2018>, 2018.
- Emde, C., Buras-Schnell, R., Kylling, A., Mayer, B., Gasteiger, J., Hamann, U., Kylling, J., Richter, B., Pause, C., Dowling, T., and Bugliaro, L.: The libRadtran software package for radiative transfer calculations (version 2.0.1), *Geoscientific Model Development*, 9, 1647–1672, <https://doi.org/10.5194/gmd-9-1647-2016>, 2016.
- Fleischmann, O. C., Burrows, J. P., and Orphal, J.: Time-windowing Fourier transform absorption spectroscopy for flash photolysis investigations, *Journal of Photochemistry and Photobiology A: Chemistry*, 157, 127–136, [https://doi.org/10.1016/S1010-6030\(03\)00069-8](https://doi.org/10.1016/S1010-6030(03)00069-8), atmospheric Photochemistry, 2003.
- Fried, A., Cantrell, C., Olson, J., Crawford, J. H., Weibring, P., Walega, J., Richter, D., Junkermann, W., Volkamer, R., Sinreich, R., Heikes, B. G., O’Sullivan, D., Blake, D. R., Blake, N., Meinardi, S., Apel, E., Weinheimer, A., Knapp, D., Perring, A., Cohen, R. C., Fuelberg, H., Shetter, R. E., Hall, S. R., Ullmann, K., Brune, W. H., Mao, J., Ren, X., Huey, L. G., Singh, H. B., Hair, J. W., Riemer, D., Diskin, G., and Sachse, G.: Detailed comparisons of airborne formaldehyde measurements with box models during the 2006 INTEX-B and MILAGRO campaigns: potential evidence for significant impacts of unmeasured and multi-generation volatile organic carbon compounds, *Atmospheric Chemistry and Physics*, 11, 11 867–11 894, <https://doi.org/10.5194/acp-11-11867-2011>, 2011.
- Frieß, U., Monks, P. S., Remedios, J. J., Rozanov, A., Sinreich, R., Wagner, T., and Platt, U.: MAX-DOAS O₄ measurements: A new technique to derive information on atmospheric aerosols: 2. Modeling studies, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006618>, 2006.
- González Abad, G., Liu, X., Chance, K., Wang, H., Kurosu, T. P., and Suleiman, R.: Updated Smithsonian Astrophysical Observatory Ozone Monitoring Instrument (SAO OMI) formaldehyde retrieval, *Atmospheric Measurement Techniques*, 8, 19–32, <https://doi.org/10.5194/amt-8-19-2015>, 2015.
- Halla, J. D., Wagner, T., Beirle, S., Brook, J. R., Hayden, K. L., O’Brien, J. M., Ng, A., Majonis, D., Wenig, M. O., and McLaren, R.: Determination of tropospheric vertical columns of NO₂ and aerosol optical properties in a rural setting using MAX-DOAS, *Atmospheric Chemistry and Physics*, 11, 12 475–12 498, <https://doi.org/10.5194/acp-11-12475-2011>, 2011.
- Hartl, A. and Wenig, M. O.: Regularisation model study for the least-squares retrieval of aerosol extinction time series from UV/VIS MAX-DOAS observations for a ground layer profile parameterisation, *Atmospheric Measurement Techniques*, 6, 1959–1980, <https://doi.org/10.5194/amt-6-1959-2013>, 2013.

- Hönninger, G. and Platt, U.: Observations of BrO and its vertical distribution during surface ozone depletion at Alert, *Atmospheric Environment*, 36, 2481–2489, [https://doi.org/10.1016/S1352-2310\(02\)00104-8](https://doi.org/10.1016/S1352-2310(02)00104-8), 2002.
- Hönninger, G., von Friedeburg, C., and Platt, U.: Multi axis differential optical absorption spectroscopy (MAX-DOAS), *Atmospheric Chemistry and Physics*, 4, 231–254, <https://doi.org/10.5194/acp-4-231-2004>, 2004.
- 5 Irie, H., Kanaya, Y., Akimoto, H., Iwabuchi, H., Shimizu, A., and Aoki, K.: First retrieval of tropospheric aerosol profiles using MAX-DOAS and comparison with lidar and sky radiometer measurements, *Atmospheric Chemistry and Physics*, 8, 341–350, <https://doi.org/10.5194/acp-8-341-2008>, 2008.
- Jang, M. and Kamens, R. M.: Characterization of Secondary Aerosol from the Photooxidation of Toluene in the Presence of NO_x and 1-Propene, *Environmental Science & Technology*, 35, 3626–3639, <https://doi.org/10.1021/es010676+>, 2001.
- 10 Jin, J., Ma, J., Lin, W., Zhao, H., Shaiganfar, R., Beirle, S., and Wagner, T.: MAX-DOAS measurements and satellite validation of tropospheric NO₂ and SO₂ vertical column densities at a rural site of North China, *Atmospheric Environment*, 133, 12–25, <https://doi.org/10.1016/j.atmosenv.2016.03.031>, 2016.
- Krotkov, N. A., Lamsal, L. N., Celarier, E. A., Swartz, W. H., Marchenko, S. V., Bucsela, E. J., Chan, K. L., Wenig, M., and Zara, M.: The version 3 OMI NO₂ standard product, *Atmospheric Measurement Techniques*, 10, 3133–3149, <https://doi.org/10.5194/amt-10-3133-2017>,
15 2017.
- Levelt, P., Van den Oord, G. H. J., Dobber, M., Malkki, A., Visser, H., de Vries, J., Stammes, P., Lundell, J., and Saari, H.: The ozone monitoring instrument, *Geoscience and Remote Sensing, IEEE Transactions on*, 44, 1093–1101, <https://doi.org/10.1109/TGRS.2006.872333>, 2006.
- Li, X., Brauers, T., Shao, M., Garland, R. M., Wagner, T., Deutschmann, T., and Wahner, A.: MAX-DOAS measurements in southern China:
20 retrieval of aerosol extinctions and validation using ground-based in-situ data, *Atmospheric Chemistry and Physics*, 10, 2079–2089, <https://doi.org/10.5194/acp-10-2079-2010>, 2010.
- Li, X., Brauers, T., Hofzumahaus, A., Lu, K., Li, Y. P., Shao, M., Wagner, T., and Wahner, A.: MAX-DOAS measurements of NO₂, HCHO and CHOCHO at a rural site in Southern China, *Atmospheric Chemistry and Physics*, 13, 2133–2151, <https://doi.org/10.5194/acp-13-2133-2013>, 2013.
- 25 Lin, J.-T., Martin, R. V., Boersma, K. F., Sneep, M., Stammes, P., Spurr, R., Wang, P., Van Roozendaal, M., Clémer, K., and Irie, H.: Retrieving tropospheric nitrogen dioxide from the Ozone Monitoring Instrument: effects of aerosols, surface reflectance anisotropy, and vertical profile of nitrogen dioxide, *Atmospheric Chemistry and Physics*, 14, 1441–1461, <https://doi.org/10.5194/acp-14-1441-2014>, 2014.
- Liu, H., Liu, C., Xie, Z., Li, Y., Huang, X., Wang, S., Xu, J., and Xie, P.: A paradox for air pollution controlling in China revealed by "APEC Blue" and "Parade Blue", *Scientific Reports*, 6, 34408, 2016.
- 30 Ma, J. Z., Beirle, S., Jin, J. L., Shaiganfar, R., Yan, P., and Wagner, T.: Tropospheric NO₂ vertical column densities over Beijing: results of the first three years of ground-based MAX-DOAS measurements (2008-2011) and satellite validation, *Atmospheric Chemistry and Physics*, 13, 1547–1567, <https://doi.org/10.5194/acp-13-1547-2013>, 2013.
- Marchenko, S., Krotkov, N. A., Lamsal, L. N., Celarier, E. A., Swartz, W. H., and Bucsela, E. J.: Revising the slant column density retrieval of nitrogen dioxide observed by the Ozone Monitoring Instrument, *Journal of Geophysical Research: Atmospheres*, 120, 5670–5692, <https://doi.org/10.1002/2014JD022913>, 2014JD022913, 2015.
- 35 Meller, R. and Moortgat, G. K.: Temperature dependence of the absorption cross sections of formaldehyde between 223 and 323 K in the wavelength range 225–375 nm, *Journal of Geophysical Research: Atmospheres*, 105, 7089–7101, <https://doi.org/10.1029/1999JD901074>, 2000.

- Mijling, B., van der A, R. J., Boersma, K. F., Van Roozendael, M., De Smedt, I., and Kelder, H. M.: Reductions of NO₂ detected from space during the 2008 Beijing Olympic Games, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2009GL038943>, 2009.
- Ortega, I., Berg, L. K., Ferrare, R. A., Hair, J. W., Hostetler, C. A., and Volkamer, R.: Elevated aerosol layers modify the O₂-O₂ absorption measured by ground-based MAX-DOAS, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 176, 34–49, <https://doi.org/10.1016/j.jqsrt.2016.02.021>, 2016.
- Platt, U. and Stutz, J.: *Differential optical absorption spectroscopy - principles and applications*, Springer, 2008.
- Richter, A., Burrows, J. P., Nüß, H., Granier, C., and Niemeier, U.: Increase in tropospheric nitrogen dioxide over China observed from space, *Nature*, 437, 129–132, 2005.
- Rodgers, C. D.: *Inverse methods for atmospheric sounding: Theory and practice*, vol. 2, World scientific, 2000.
- 10 Rotman, D. A., Tannahill, J. R., Kinnison, D. E., Connell, P. S., Bergmann, D., Proctor, D., Rodriguez, J. M., Lin, S. J., Rood, R. B., Prather, M. J., Rasch, P. J., Considine, D. B., Ramaroson, R., and Kawa, S. R.: Global Modeling Initiative assessment model: Model description, integration, and testing of the transport shell, *Journal of Geophysical Research: Atmospheres*, 106, 1669–1691, <https://doi.org/10.1029/2000JD900463>, 2001.
- Serdyuchenko, A., Gorshelev, V., Weber, M., Chehade, W., and Burrows, J. P.: High spectral resolution ozone absorption cross-sections - Part 2: Temperature dependence, *Atmospheric Measurement Techniques*, 7, 625–636, <https://doi.org/10.5194/amt-7-625-2014>, 2014.
- 15 Sinreich, R., Frieß, U., Wagner, T., and Platt, U.: Multi axis differential optical absorption spectroscopy (MAX-DOAS) of gas and aerosol distributions, *Faraday Discussions*, 130, 153, <https://doi.org/10.1039/b419274p>, 2005.
- Solomon, S., Schmeltekopf, A. L., and Sanders, R. W.: On the interpretation of zenith sky absorption measurements, *Journal of Geophysical Research: Atmospheres*, 92, 8311–8319, <https://doi.org/10.1029/JD092iD07p08311>, 1987.
- 20 Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Bulletin of the American Meteorological Society*, 96, 2059–2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
- Stohl, A., Seibert, P., Arduini, J., Eckhardt, S., Fraser, P., Grealley, B. R., Lunder, C., Maione, M., Mühle, J., O'Doherty, S., Prinn, R. G., Reimann, S., Saito, T., Schmidbauer, N., Simmonds, P. G., Vollmer, M. K., Weiss, R. F., and Yokouchi, Y.: An analytical inversion method for determining regional and global emissions of greenhouse gases: Sensitivity studies and application to halocarbons, *Atmospheric Chemistry and Physics*, 9, 1597–1620, <https://doi.org/10.5194/acp-9-1597-2009>, 2009.
- 25 Stohl, A., Prata, A. J., Eckhardt, S., Clarisse, L., Durant, A., Henne, S., Kristiansen, N. I., Minikin, A., Schumann, U., Seibert, P., Stebel, K., Thomas, H. E., Thorsteinsson, T., Tørseth, K., and Weinzierl, B.: Determination of time- and height-resolved volcanic ash emissions and their use for quantitative ash dispersion modeling: the 2010 Eyjafjallajökull eruption, *Atmospheric Chemistry and Physics*, 11, 4333–4351, <https://doi.org/10.5194/acp-11-4333-2011>, 2011.
- 30 Sun, Y., Wang, Z., Wild, O., Xu, W., Chen, C., Fu, P., Du, W., Zhou, L., Zhang, Q., Han, T., Wang, Q., Pan, X., Zheng, H., Li, J., Guo, X., Liu, J., and Worsnop, D. R.: "APEC Blue": Secondary Aerosol Reductions from Emission Controls in Beijing, *Scientific Reports*, 6, 20668, 2016.
- Thalman, R. and Volkamer, R.: Temperature dependent absorption cross-sections of O₂-O₂ collision pairs between 340 and 630 nm and at atmospherically relevant pressure, *Phys. Chem. Chem. Phys.*, 15, 15371–15381, <https://doi.org/10.1039/C3CP50968K>, 2013.
- van der A, R. J., Eskes, H. J., Boersma, K. F., van Noije, T. P. C., Van Roozendael, M., De Smedt, I., Peters, D. H. M. U., and Meijer, E. W.: Trends, seasonal variability and dominant NO_x source derived from a ten year record of NO₂ measured from space, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD009021>, 2008.

- Vandaele, A., Hermans, C., Simon, P., Carleer, M., Colin, R., Fally, S., Merienne, M., Jenouvrier, A., and Coquart, B.: Measurements of the NO₂ absorption cross-section from 42000 cm⁻¹ to 10000 cm⁻¹ (238-1000 nm) at 220 K and 294 K, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 59, 171 – 184, [https://doi.org/10.1016/S0022-4073\(97\)00168-4](https://doi.org/10.1016/S0022-4073(97)00168-4), *atmospheric Spectroscopy Applications* 96, 1998.
- Wagner, T., Dix, B., Friedeburg, C. v., Fries, U., Sanghavi, S., Sinreich, R., and Platt, U.: MAX-DOAS O₄ measurements: A new technique to derive information on atmospheric aerosols - Principles and information content, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2004JD004904>, 2004.
- Wagner, T., Deutschmann, T., and Platt, U.: Determination of aerosol properties from MAX-DOAS observations of the Ring effect, *Atmospheric Measurement Techniques*, 2, 495–512, <https://doi.org/10.5194/amt-2-495-2009>, 2009.
- Wagner, T., Beirle, S., Brauers, T., Deutschmann, T., Frieß, U., Hak, C., Halla, J. D., Heue, K. P., Junkermann, W., Li, X., Platt, U., and Pundt-Gruber, I.: Inversion of tropospheric profiles of aerosol extinction and HCHO and NO₂ mixing ratios from MAX-DOAS observations in Milano during the summer of 2003 and comparison with independent data sets, *Atmospheric Measurement Techniques*, 4, 2685–2715, <https://doi.org/10.5194/amt-4-2685-2011>, 2011.
- Wagner, T., Beirle, S., Benavent, N., Bösch, T., Chan, K. L., Donner, S., Dörner, S., Fayt, C., Frieß, U., García-Nieto, D., Gielen, C., González-Bartolome, D., Gomez, L., Hendrick, F., Henzing, B., Jin, J. L., Lampel, J., Ma, J., Mies, K., Navarro, M., Peters, E., Pinardi, G., Puentedura, O., Puķīte, J., Remmers, J., Richter, A., Saiz-Lopez, A., Shaiganfar, R., Sihler, H., Van Roozendael, M., Wang, Y., and Yela, M.: Is a scaling factor required to obtain closure between measured and modelled atmospheric O₄ absorptions? An assessment of uncertainties of measurements and radiative transfer simulations for 2 selected days during the MAD-CAT campaign, *Atmospheric Measurement Techniques*, 12, 2745–2817, <https://doi.org/10.5194/amt-12-2745-2019>, 2019.
- Wang, S., Gao, J., Zhang, Y., Zhang, J., Cha, F., Wang, T., Ren, C., and Wang, W.: Impact of emission control on regional air quality: An observational study of air pollutants before, during and after the Beijing Olympic Games, *Journal of Environmental Sciences*, 26, 175 – 180, [https://doi.org/10.1016/S1001-0742\(13\)60395-2](https://doi.org/10.1016/S1001-0742(13)60395-2), 2014.
- Wang, S., Cuevas, C. A., Frieß, U., and Saiz-Lopez, A.: MAX-DOAS retrieval of aerosol extinction properties in Madrid, Spain, *Atmospheric Measurement Techniques*, 9, 5089–5101, <https://doi.org/10.5194/amt-9-5089-2016>, 2016.
- Wang, Y., Beirle, S., Lampel, J., Koukouli, M., De Smedt, I., Theys, N., Li, A., Wu, D., Xie, P., Liu, C., Van Roozendael, M., Stavrou, T., Müller, J.-F., and Wagner, T.: Validation of OMI, GOME-2A and GOME-2B tropospheric NO₂, SO₂ and HCHO products using MAX-DOAS observations from 2011 to 2014 in Wuxi, China: investigation of the effects of priori profiles and aerosols on the satellite products, *Atmospheric Chemistry and Physics*, 17, 5007–5033, <https://doi.org/10.5194/acp-17-5007-2017>, 2017.
- Wang, Y., Dörner, S., Donner, S., Böhnke, S., De Smedt, I., Dickerson, R. R., Dong, Z., He, H., Li, Z., Li, Z., Li, D., Liu, D., Ren, X., Theys, N., Wang, Y., Wang, Y., Wang, Z., Xu, H., Xu, J., and Wagner, T.: Vertical profiles of NO₂, SO₂, HONO, HCHO, CHOCHO and aerosols derived from MAX-DOAS measurements at a rural site in the central western North China Plain and their relation to emission sources and effects of regional transport, *Atmospheric Chemistry and Physics*, 19, 5417–5449, <https://doi.org/10.5194/acp-19-5417-2019>, 2019.
- Wenig, M., Spichtinger, N., Stohl, A., Held, G., Beirle, S., Wagner, T., Jähne, B., and Platt, U.: Intercontinental transport of nitrogen oxide pollution plumes, *Atmospheric Chemistry and Physics*, 3, 387–393, <https://doi.org/10.5194/acp-3-387-2003>, 2003.
- Wenig, M. O., Cede, A. M., Bucsele, E. J., Celarier, E. A., Boersma, K. F., Veefkind, J. P., Brinksma, E. J., Gleason, J. F., and Herman, J. R.: Validation of OMI tropospheric NO₂ column densities using direct-Sun mode Brewer measurements at NASA Goddard Space Flight Center, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD008988>, 2008.

- Witte, J. C., Schoeberl, M. R., Douglass, A. R., Gleason, J. F., Krotkov, N. A., Gille, J. C., Pickering, K. E., and Livesey, N.: Satellite observations of changes in air quality during the 2008 Beijing Olympics and Paralympics, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2009GL039236>, 2009.
- Wittrock, F., Oetjen, H., Richter, A., Fietkau, S., Medeke, T., Rozanov, A., and Burrows, J. P.: MAX-DOAS measurements of atmospheric trace gases in Ny-Alesund - Radiative transfer studies and their application, *Atmospheric Chemistry and Physics*, 4, 955–966, <https://doi.org/10.5194/acp-4-955-2004>, 2004.
- Wu, F. C., Xie, P. H., Li, A., Chan, K. L., Hartl, A., Wang, Y., Si, F. Q., Zeng, Y., Qin, M., Xu, J., Liu, J. G., Liu, W. Q., and Wenig, M.: Observations of SO₂ and NO₂ by mobile DOAS in the Guangzhou eastern area during the Asian Games 2010, *Atmospheric Measurement Techniques*, 6, 2277–2292, <https://doi.org/10.5194/amt-6-2277-2013>, 2013.
- 10 Zhang, J., Wang, S., Guo, Y., Zhang, R., Qin, X., Huang, K., Wang, D., Fu, Q., Wang, J., and Zhou, B.: Aerosol vertical profile retrieved from ground-based MAX-DOAS observation and characteristic distribution during wintertime in Shanghai, China, *Atmospheric Environment*, 192, 193 – 205, <https://doi.org/10.1016/j.atmosenv.2018.08.051>, 2018.
- Zhang, Q., Streets, D. G., He, K., Wang, Y., Richter, A., Burrows, J. P., Uno, I., Jang, C. J., Chen, D., Yao, Z., and Lei, Y.: NO_x emission trends for China, 1995-2004: The view from the ground and the view from space, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2007JD008684>, 2007.
- 15 Zhang, R., Tie, X., and Bond, D. W.: Impacts of anthropogenic and natural NO_x sources over the U.S. on tropospheric chemistry, *Proceedings of the National Academy of Sciences*, 100, 1505–1509, <https://doi.org/10.1073/pnas.252763799>, 2003.
- Zhao, B., Wang, S. X., Liu, H., Xu, J. Y., Fu, K., Klimont, Z., Hao, J. M., He, K. B., Cofala, J., and Amann, M.: NO_x emissions in China: historical trends and future perspectives, *Atmospheric Chemistry and Physics*, 13, 9869–9897, <https://doi.org/10.5194/acp-13-9869-2013>,
20 2013.