Originally 'Contributions from intrinsic low-frequency climate variability to the accelerated decline in Arctic sea ice in recent decades'

Title changed to 'Changes in sea surface temperature and atmospheric circulation patterns associated with reductions in Arctic sea-ice cover in recent decades

- 5 Referee report:
 - General comments:

The authors have provided more information and revised the discussions in response to the referees' comments, and have improved their manuscript.

Some aspects of the SOM patterns are still not clear to me, and the discussion of mechanisms needs more detail to be clearer.

I am not convinced that asymmetries in North Pacific SST anomalies are as important as the authors claim, so that aspect could be more speculative.

Thus I recommend further revision as follows:

15 Specific comments:

30

Are composite results sensitive to the selection of the SOM matrix of patterns? As in the example in the supplement, patterns similar to node1 and node 9 also occur and dominate other choices of matrix, and the selection of years for composites is likewise similar for other array choices. The composite results are not sensitive to the selection of the SOM matrix of patterns.

- 20 On reading again the section on potential mechanisms, I think it could be made clearer and improved by providing more precise information about geographic regions mentioned, and linking more clearly to the sea ice concentration features in the nodes. e.g. line 162 (and 187): specify latitude/longitude for the region of interest in the western tropical Pacific; likewise line 168 for the 'region between two centers', line 171 for region of reduced sea ice concentration,
- 25 line 174 for the region of anomalous high pressure, line 176 for the region of anomalous low pressure, etc.

Specific geographic locations have been added following the suggestion.

Line 207/208: apart from opposite sign, the differences in SST patterns are rather small, and it is difficult to attribute the differences in high-latitude circulation to the SST patterns without more direct evidence, so this should be stated as a hypothesis. (The SST anomaly sign difference alone

- and nonlinear effects might be the main cause of any SST-related circulation differences.) It would be informative if one or two example scatterplots could be added to illustrate how particular regional values in ice cover and in atmospheric variables are related in in years making up composites.
- 35 This sentence has been revised, as suggested.

A scatter plot is produced (Fig. S1). During the years of Node 1 when sea ice concentration anomalies over Barents and Kara Seas (30°E-100°E, 70°N-85°N) are negative, the 500-hPa height anomalies in the region (0°-160°E, 60°N-90°N) are mostly positive. The opposite occurs during the years of Node 9.





45 Further minor comments:

It would be helpful to mention the use of composites in the abstract.

Added

line 25: '...remain a subject of ..'

Changed

50 line 31: '...On larger scales, ...'

Changed

line 50: '... (ENSO) events to the ...'

```
Added
```

55

line 78: what is the resolution of the OLR data grid?

The resolution of the OLR grid has been added.

Line 81: it would be helpful to add a statement about what is optimised in SOM Added

Line 85: I am not sure what the sentence 'All patterns in ...' means, as the patterns represent a subset of the input data.

60 Changed 'all patterns' to 'all vectors'

Line 94: it would be clearer to say 'The best matching SOM pattern for each autumn is determined on the basis of minimum Euclidean distance'. It would be useful to provide an equation for this. It would also be better to place this later in the paragraph (e.g. after line 105), after defining the patterns.

65 The sentence is modified, but we feel it is better to leave the sentence at where it is. Instead of adding an equation, a reference is added.

Line 107: It would be clearer to define frequency of occurrence as the number of autumns for which a pattern is selected as 'best matching' divided by the total number of autumns. Clarified 70 Line 110: I am not sure what is meant by 'linear regression of the number of the projections' : an equation would be useful.

Replace the word 'projections' by 'occurrences' and added an equation

Fig. 1: are the patterns normalised?

(It might be worth noting that the sign of the pattern is important, unlike EOFs for which the sign is arbitrary. An EOF analysis would give one pattern like nodes 1 and 9, with different signs of projection in early/late years in the data.)

The patterns were not normalized, which represent the realistic anomalous sea ice concentration.

Line 122: it would help to say that 24% corresponds to 9/38.

80 Added

75

90

Line 127: omit 'much' (the occurrence difference is between 5/38 and 3/38) Changed Line 130: ... Figs 2 and 3 Added

Fig. 4: are the labels correct? The SOM-associated trend seems larger than the observed trend. Also, the 'dots' indicating significance are very hard to see in the figure.

The label is right. The SOM-related trend is smaller than the observed trend. The dots are made less dense to make it easier to see.

Line 155: Suggest better to put 'To explore the processes associated with the spatial patterns ...' Changed

Line 213: better to also state the years used in the before and after 2000 composites. Added

Line 217: to me, the dominant feature of the SST patterns associated with node1 and node9 is their similarity, with small differences. Further evidence is required if circulation and sea-ice

95 concentration differences are to be attributed to the influence of the small SST pattern differences – perhaps you could state that further investigation is required to support your statements, particularly as this effect forms a large part of your conclusions in section 8. We added some discussions about the impact of SST.

Fig. 2 caption: Better to say that the figure indicates which mode best matched the observed sea ice concentration in each autumn.

Changed

100

Fig. 3 caption: Rather than 'explained', this figure just shows the linear trend associated with each pattern when projected on the observations?

Changed

105 Fig. 4: check the content, labelling and dots.

We revised fig. 4a.

Fig. 5 and Fig. 8: It would be helpful to repeat in the caption the actual years used for each composite.

Added

Changes in sea-surface temperature and atmospheric circulation patterns associated with reductions in Arctic sea-ice cover in recent decades

115 Lejiang Yu¹, Shiyuan Zhong²

¹SOA Key Laboratory for Polar Science, Polar Research Institute of China, Shanghai, China
²Department of Geography, Environment and Spatial Sciences, Michigan State University, East Lansing, MI, USA

4

Correspondence to: Dr. Shiyuan Zhong (zhongs@msu.edu)

Abstract. In recent decades, the Arctic sea ice has been declining at a rapid pace as the Arctic is warmed at a rate of twice the global average. The underlying physical mechanisms for the Arctic warming and accelerated sea ice retreat are not fully understood. In this study, we apply a relatively novel statistical method called Self-Organizing Maps (SOM) <u>along with composite analysis</u> to examine the trend and

- variability of autumn Arctic sea ice in the past three decades and their relationships to large-scale atmospheric circulation changes. Our statistical results show that the anomalous autumn Arctic Dipole (AD) (Node 1) and the Arctic Oscillation (AO) (Node 9) could explain in a statistical sense as much as 50% of autumn sea ice decline between 1979 and 2016. The Arctic atmospheric circulation anomalies associated with anomalous sea surface temperature patterns over the North Pacific and North Atlantic
 influence Arctic sea ice primarily through anomalous temperature and water vapour advection and
 - associated radiative feedback.

1 Introduction

135

In recent decades, the Arctic sea ice has been decreasing at an unprecedented rate (Rothrock et al., 1999; Parkinson, 2014). The accelerated retreat in Arctic sea ice exerts a significant impact not only on the marine and terrestrial ecological systems of the Arctic (Post et al., 2013), but also on the environment of

the mid latitude (Mori et al., 2014; Kug et al., 2015).

The underline mechanisms for the Arctic sea ice decline remain to be a subject of active research. Studies have suggested that both anthropogenic forcing due to greenhouse gas and aerosol emissions (Min et al., 2008; Notz and Marotzke, 2012; Gagné et al., 2015) and natural mechanisms at a wide range of scales

- contribute to the observed Arctic sea ice decline. Local processes, including surface thermal inversion (Bintanja et al., 2011), atmospheric lapse-rate (Pithan and Mauritsen, 2014), ice-albedo feedback (Flanner et al., 2011) and water vapour and cloud radiative feedback (Sedlar et al., 2011), have been found to affect Arctic sea ice. On the other endlarger scales, global sea-surface temperature (SST) and pressure oscillations, such as the Arctic Dipole (AD) (Wang et al., 2009), the Atlantic Multidecadal
- 145 Oscillation (AMO) (Park and Latif, 2009), the Arctic Oscillation (AO) (Rigor et al., 2002; Deser and Teng, 2008), the North Atlantic Oscillation (NAO) (Koenigk et al., 2009), and the Pacific Decadal Oscillation (PDO) (Ding et al., 2014), have also been linked to the Arctic sea ice variations and the recent declining trend. Understanding the relative contributions from these multi-scale processes to Arctic sea ice decline is vital not only for forecasting sea ice conditions but also for projecting climate change and

its impact on the Arctic environment and beyond (Stroeve et al., 2007; Day et al., 2012; Swart et al., 2015; Ding et al., 2017).

In this study, we examine the contributions of changes in large-scale atmosphere and ocean circulations to the trends in Arctic sea ice by applying the Self Organizing Maps (SOM) method (Kohonen, 2001). A relatively new neural network-based method, SOM is superior to some other feature-extracting or

- 155 clustering methods in that it describes the continuum of atmospheric and oceanic states with a manageable number of representative patterns as compared to only providing useful information on aggregate patterns by Empirical Orthogonal Function (EOF) and other similar methods. Over the past decade, SOM has been widely used in atmosphere and ocean sciences (Hewitson and Crane, 2002; Leloup et al., 2007; Johnson et al., 2008; Lee et al., 2011; Chu et al., 2012). For example, SOM was used
- to explain the eastward shift of NAO since the late 1970s (Johnson et al., 2008). The SOM approach was also used to examine the contributions of different El Niño Southern Oscillation (ENSO) events to the SST trend in the tropical Pacific Ocean (Johnson, 2013).

We apply SOM to a monthly sea ice concentration dataset for the period 1979 - 2016. We will show how much <u>of</u> the recent declining trend in the Arctic sea ice concentrations may be associated with the low

165 frequency atmospheric circulation modes related to SST anomalies over the Pacific and Atlantic Oceans. Although the analyses have been carried out for all four seasons, we will show the results for autumn only since Arctic sea ice reduction in autumn has strongest influence on the wintertime atmospheric circulations over Eurasia and North America (Francis et al., 2009; Petoukhov and Semenov, 2010; Peings and Magnusdottir, 2014).

170 2 Methods

The period of this study is from 1979-2016. The Arctic sea ice anomaly analysis utilizes the monthly sea ice concentration dataset produced by the United States National Snow and Ice Data Centre (NSIDC) (http://nsidc.org/data/NSIDC-0051). The NSIDC data are on a horizontal grid of 25 km \times 25 km with a polar stereographic projection. The corresponding SST patterns are analyzed using the 1° latitude \times 1°

175 longitude SST data from the Hadley Centre (Rayner et al., 2003) (<u>http://www.metoffice.gov.uk/hadobs/hadisst/</u>). The SST anomaly patterns are linked to well-known modes of climate variability characterized by several climate indices including the Pacific Decadal Oscillation (PDO) (Mantua et al., 1997) defined as the first EOF mode of SST anomalies in the North Pacific Ocean, the Atlantic Multidecadal Oscillation (AMO) (Enfield et al., 2001) defined as the
10-year running mean of detrended SST anomalies in the Atlantic Ocean north of the equator, and the
Arctic Oscillation (AO) and Arctic Dipole (AD) defined as the first and second EOF modes of the
1000-hPa geopotential height anomalies north of 70°N (Wang et al., 2009).

The atmospheric circulations are determined using data from the 1.5° latitude × 1.5° longitude European Centre for Medium-Range Weather Forecasts (ECMWF) Reanalysis (ERA-Interim) (Dee et al., 2011). Serreze et al. (2012) assessed humidity data from ERA-Interim reanalyses and found only small bias of

185 Serreze et al. (2012) assessed humidity data from ERA-Interim reanalyses and found only small bias of less than 8% at 1000 hPa for boreal autumn and smaller bias above 1000 hPa. Ding et al. (2017) showed that the ERA-Interim reanalyses can represent reliably the observed circulation, radiation flux, temperature, and water vapour. The surface and upper-air atmospheric variables examined in this study are all extracted from ERA-Interim, except for outgoing long-wave radiation (OLR) that is
190 obtained from the gridded monthly OLR data with a 2.5° latitude ×2.5° longitude grid produced by the US National Oceanic and Atmospheric Administration (NOAA)

(http://www.esrl.noaa.gov/psd/data/gridded/data.interp_OLR.html) (Liebmann and Smith, 1996).

The SOM methods, which combines some features of clustering analysis and neural network, is more advantageous over traditional clustering and pattern extracting methods (Grotjahn et al., 2016).

- 195 The SOM technique is utilized to extract patterns of Arctic sea ice concentrations. A neural network-based method, SOM uses unsupervised learning to determine generalized patterns in complex data. The technique can reduce multidimensional data into two-dimensional array consisting of a matrix of nodes. Each node in the array has a reference vector that displays a spatial pattern of the input data. All patterns-vectors in the two-dimensional array represent the full continuum of states in the input data. The
- 200 SOM algorithm also is a clustering technique, but unlike other clustering technique, it does not need a priori decisions on data distribution. Unlike the EOF analysis, the SOM technique does not require the orthogonality of two spatial patterns. A detailed description of the SOM algorithm can be found in Kohonen (2001).

In this study, the SOM technique is used to categorize anomalous seasonal sea ice concentration patterns north of 50°N for autumn (September-October-November). Seasonal anomalies are calculated by subtracting the climatology for the season relative to the study period 1979-2016 from the seasonal means. <u>The best matching SOM pattern to the sea ice anomaly pattern</u><u>The_anomalous sea ice pattern</u> for each autumn is <u>determined assigned to the best matching SOM pattern</u> on the basis of minimum 210

calculated by the product of root-mean-square error and the square root of the number of grid points (Hewitson and Crane, 2002). Spatial correlations between sea ice field for each autumn and its corresponding best-matching SOM pattern are calculated and the overall mean spatial correlation coefficients obtained by averaging over the 38 autumns are used to determine the number of SOM nodes or grids to be used in the current analysis (Lee and Feldstein, 2013). Table 1 shows the overall mean

Euclidean distance, a measure of straight-line distance between two points in Euclidean space and

- 215 spatial correlation coefficients for several SOM grid configurations with nodes ranging from 2×2 to 4×5. As expected, the overall correlation increases as the number of nodes increases. The largest jump between any two grid configurations occurs from a 2×4 grid to a 3×3 grid. Thus, a 3×3 grid is selected for the SOM analysis. While a higher spatial correlation can be achieved with larger number of grids, the 3x3 grid is able to capture the main variability pattern in the autumn Arctic sea ice at the sacrifice of some
- 220 details that can be depicted by larger number of grids. The frequency of occurrence for each SOM node is defined as the number of autumns that node represents for which a pattern is selected as 'best matching' divided by the total number of autumns (38) in the study period. The contribution of each SOM pattern to the trends in the Arctic sea ice concentrations is calculated by the product of each SOM pattern and a rate determined by temporal linear regression of the number of the projectionsoccurrences of each pattern for each autumn (Johnson, 2013) represented by $Y(y_1, y_2, \Box, y_i, \Box, y_n)$ onto the
- 225

time series of autumns between 1979 and 2016 represented by $X(x_1, x_2, \Box, x_i, \Box, x_n)$ with n

equal to 38. The linear regression between Y and X is calculated using

 $y_i = \alpha + \beta x_i + \varepsilon_i$

<u>Where α is the intercept</u>, β is the slope that we focus on, and ε_i is a residual term.

230 -The sum of these contributions presents the trends in Arctic sea ice explained in the statistical sense by the SOM patterns, which in this case indicates trends resulting from low-frequency variability. The significance of the trends in the time series for each SOM pattern is tested using the Student's t-test. Residual trends are calculated by subtracting SOM-explained trends from the total trends.

3 SOM results

235 Sea ice concentration anomalies occur mainly in the marginal seas from the Barents Sea to the Beaufort Sea, with maximum anomalies in the Barents Sea and the Kara Sea, and over the Beaufort Sea and the 8

Χ	域代码已更改
	域代码已更改
	域代码已更改
_	域代码已更改
	域代码已更改
	域代码已更改
	域代码已更改

域代码已更改

East Siberian Sea (Fig. 1). Nodes 1, 2 and 4 depict all negative anomalies, whereas Nodes 5, 6, 8 and 9 exhibit all positive anomalies. Nodes 3 and 7, on the other hand, show a mixed pattern with opposite changes in the Barents Sea in the Atlantic sector of the Arctic Ocean and the Beaufort Sea and East

- 240 Siberian Sea over the Pacific sector. The largest change with similar strength over the Pacific and the Atlantic sectors are depicted by Nodes 1 and 9, and both nodes also have the highest (24% corresponding to 9/38) frequency of occurrences. The other nodes that are either all positive or all negative exhibit larger spatial variability in the strengths of the signals with much stronger signal in either the Atlantic sector (Nodes 2, 5 and 8) or the Pacific sector (Nodes 4 and 6). The two mixed patterns show similar 245 strength over the two sectors, but the pattern with positive anomalies over the Beaufort Sea and the East
- Siberian Sea and negative anomalies over the Barents Sea, as depicted by Node 3, occurs much-more frequently (13%) than the opposite pattern represented by Node 7 (8%).

The trends in the frequency of occurrence for each SOM pattern and their contribution to the trends in the Arctic autumn sea ice concentrations are examined and the results are shown in Figs. 2 and 3. The nodes

250 with spatially uniform changes appear to be separated into two clusters with those showing all positive anomalies (Nodes 5, 6, 8 and 9) appearing in the 80s and 90s and those having all negative anomalies (Nodes 1 and 4) appearing after 2000. The transition from all positive anomalies in the earlier part, to all negative anomalies in the later part, of the time series is consistent with the trends in the observed Arctic sea ice concentration during the same time period. Not surprisingly, the transition appears to be 255 dominated by the two strongest and most frequent patterns denoted by Node 9 (all positive and occurring from the 1980s through mid 1990s) and Node 1 (all negative and occurring exclusively after 2005). Only

these two nodes have linear trends that are statistically significant at above 95% confidence level. The slopes of the trend lines for these two nodes are opposite but the values are similar (0.027 yr⁻¹ for Node 1 and -0.021 yr⁻¹ for Node 9). The other nodes have statistically insignificant trends with magnitudes less 260 than 0.01 yr^{-1} .

The spatial patterns of the autumn sea ice concentration trends explained in a statistical sense by each node and by all nodes are shown in Figs. 3 and 4. Together, the nine SOM nodes explain statistically about 60% of total trends in the autumn sea ice loss (Fig. 4b). Among them Node 1 explains statistically the largest portion (33%) of the total trend, followed by Node 9 (21%), with the other 7 nodes together accounting for only 6% of the loss.

For comparison, the SOM patterns and their occurrence time series for a larger grid (3x5) are provided in

Supplement (Figs. S1 and S2). As expected, with more nodes, the 3x5 grid depict more details and each node has smaller frequency compared to the 3x3 grid. However, the dominant nodes (also nodes 1 and 9) show nearly identical patterns as those in the 3x3 grid. Like the 3×3 grid, nodes 1 and 9 in the 3×5 grid make greater contributions to the trend in autumn Arctic sea ice than other nodes (Fig. S3). Also as

expected, the trend explained by nodes 1 and 9 in the 3x5 grid is smaller (46%) compared to 54% by the same two nodes in the 3×3 grid.

4 Potential Mechanisms

270

To explain statistically explore the processes associated with the spatial patterns depicted by the two dominant nodes, we made composite maps over the years of occurrences for Nodes 1 and 9, respectively, and Figs. 5-7 show composite patterns of the anomalous SST and OLR (as a proxy for tropical convection) and anomalous atmospheric circulations represented by the 500-hPa geopotential height, 850-hPa wind, surface-750-hPa specific humidity, surface downward longwave radiation, and surface air temperature. The composite results are insensitive to the selection of the size of the SOM matrix. For

- Node 1, the SST composite pattern resembles negative phase of the PDO in the North Pacific and positive phase of the AMO in the North Atlantic (Fig. 5). The positive SST anomalies over the tropical western Pacific (140°E, 10°N) produce anomalous local convection, exciting a wave train (Hoskins and Karoly, 1981; Jin and Hoskins, 1995; Higgins and Mo, 1997), which propagates northeastwards to North Pacific, North America, North Atlantic, and Eurasia (Fig. 5). In mid-latitudes, the anomalous SST
- and the nodes of the wave train interacts to produce atmospheric circulation anomalies. Over the Arctic, the SST and height patterns are similar to those associated with a negative phase AD, with a negative 500-hPa height anomaly center over Greenland and the Baffin Bay, and a positive anomaly center over the Kara Sea. The zonal pressure gradient between the two centers (60°W, 75°N; 100°E, 75°N) induces anomalous low-troposphere southwesterly and southerly winds over the North Atlantic Ocean (Fig. 6)
- 290 that transport warm and moist air from North Atlantic into the Arctic Ocean north of Eurasia, thus increasing surface air temperature and humidity and reducing sea ice concentration in the the whole Arctic (Fig. 7). The higher moisture content in the Arctic surface air also facilitates the occurrence of water vapour and cloud radiative feedback process (Sedlar et al., 2011) during which increased downward longwave radiation (Fig. 7) enhances surface warming and sea ice melting. The anomalous

high pressure (100°E, 75°N) produces subsidence and the adiabatic warming associated with the sinking

air contributes to the sea ice loss (Ding et al., 2017) (Not shown). The opposite may occur in the region under the anomalous low pressure center ($60^{\circ}W$, $75^{\circ}N$). But the warm southwesterly winds over the northeastern Canada also favor the sea ice loss there. Gong et al. (2017) and Lee et al. (2017) showed that the increasing downward longwave radiation related to the changes of mid-latitude atmospheric

300 circulation leads to the wintertime Arctic warming. Our focus here is on autumn sea ice trend and we use the low-level water vapour field that is found to resemble more closely to the surface downward radiation fields compared to the total column water (not shown) used in the two studies. Like their studies, the North Atlantic is the main passage of water vapour into the Arctic. But over the Bering Sea and Europe no water vapour is transported into the Arctic in our studies. Their study period ranges from 1980 to 2010, which is different from our study period (1979-2016).

For Node 9, the SST composite is characterized by negative anomalies in both the Pacific and the Atlantic except for areas of the tropical Pacific and the west coast of North America (Fig.5). The negative SST anomalies over the tropical western Pacific (140°E, 10°N), corresponding to suppressed convective activities, generate one wave train that is in opposite phase to the wave train associated with Node 1.

- 310 Similarly, the nodes of the wave train interact with the local SST anomalies over the mid-latitude regions, producing a nearly opposite (but not symmetrical) anomalous atmospheric circulation pattern to that of Node 1. Over the Arctic, the patterns resemble a positive phase AO (Fig.5). Anomalous northeasterly winds in the North Atlantic Ocean induced by anomalous Icelandic low are unfavourable for warm air intrusion into the Arctic Ocean (Fig. 6), a result also indicated in a previous study (Kim et al.,
- 315 2017). Despite the occurrences of anomalous southwesterly winds over the Barents Sea and southeasterly winds over the Sea of Okhotsk, the cold advection from anomalous cold sea water prevents the Arctic sea ice from melting. Meanwhile the cold advection also reduces water vapor content in the lower troposphere and the resulting smaller downward longwave radiation facilitates the occurrence of negative surface air temperature and positive Arctic sea ice anomalies north of Eurasia (Fig. 7). The
- 320 northerly winds over the Beaufort Sea, northeastern Canada and Greenland Sea also contribute to the decrease in surface air temperature and increase in the sea ice concentration. The opposite patterns in Nodes 1 and 9 can be largely associated with the differences in the water vapor-radiation feedback process resulting from anomalous temperature and especially water vapour transport by anomalous atmospheric circulations associated with different patterns of SST anomalies
- 325 over the North Pacific and Atlantic. The patterns of SST anomalies are nearly symmetric for the two 11

nodes over the North Atlantic, but they are somewhat asymmetric over the North Pacific. For Node 1, there is one center of high SST over the central North Pacific, whereas Node 9 is associated with two centers of low SST: one over the Coast of Japan and another in central North Pacific. <u>The opposite in the sign of the SST anomalies and the non-linear effects might be the main cause of SST-related circulation differences</u>These differences in the SST anomaly patterns lead to different wave trains and

330

335

high latitude atmospheric circulations.

For the composite analysis often used in previous literature, lower (higher) Arctic sea concentration years refer to after (before) 2000.

The SOM-based composites in the above discussion, which are made using years indicated (Fig. 2) by the occurrences for Node 1 (2007-2013, 2015-2016), and, separately, for node 9 (1980-1982,

- 1986-1989, 1992, 1996), are compared to composites of years of high (before 20001979-2000) and low (after 20002001-2016) Arctic sea ice coverage according to the Autumn Arctic sea ice time series (Figure 8). The composited SST patterns over the North Pacific are almost a mirror image to each other, indicating a positive (negative) phase PDO before (after) 2000 (Fig. 8). On the other hand, the
- 340 composited SST patterns corresponding to Node 1 and Node 9 are not symmetrical (Fig. 5). Similar situations occur over North Atlantic. The magnitude and significant level of SOM-based SST composites in Fig. 5 are also higher than the time-series-based composites in Fig. 8. Similarly, in mid-latitudes, the composites of the anomalous 500hPa geopotential heights before and after 2000 are nearly symmetrical (Fig. 8) and the significant level and amplitude of the wave train are also lower
- 345 than those in Fig. 5. At high latitudes, the anomalous atmospheric circulations in Fig. 8 show a mixed pattern of AO and AD and have no extreme centers. However, Fig. 5 exhibits clear AO and AD patterns and extreme centers, revealing more clearly the relationship between the anomalous sea ice concentration and anomalous atmospheric circulations. Based on these examples, the SOM-based composites allows for better depiction of atmospheric and SST conditions corresponding to sea ice anomaly patterns compared to typical time-series-based composite approach.
- One drawback of the SOM-based composite is the relatively small number of composite members (Node 1 and Node 9 each has 9 members), which makes the composite more vulnerable to the influence of extreme cases. To examine this issue, the standard deviations of the SOM-based composites are compared to those of time-series-based composites that have much larger members (17 355 members for after-2000- composites and 21 members for before-2000-composites) and the results for 12

the anomalous SST and 500-hPa height are shown in Figs. S4 and S5. The spatial patterns and magnitudes of the standard deviations are similar between the Node-1 and after-2000 composites and between Node-9 and before-2000 composites. The F-test indicates that except for small patches there are no statistically significant differences in the standard deviations between the small member SOM-based composites and the large-member typical composites (Fig. S6), suggesting it is unlikely the SOM-based composites are strongly influenced by extreme cases as a result of small composite members.

5 Conclusions

360

We investigate the potential mechanisms for the autumn arctic sea ice decline for the period 1979-2016
using the SOM method. Our results show that more than half of the autumn Arctic sea ice loss may be associated with the changes in the temperature and water vapour transport and the associated water vapour radiation feedback resulting from anomalous atmospheric circulations, which might be linked to SST anomalies over the North Pacific and North Atlantic. This result provides further evidence that the mid-latitude SST anomalies play a vital role in the accelerated Arctic sea ice decline in recent decades

- 370 (Ding et al., 2017; Yu et al., 2017). The main result is that the opposite pattern of the Arctic sea ice anomalies during the early (positive) and later (negative) parts of the 1979-2016 period is not associated with opposite phase of an atmospheric circulation mode, but instead the change can be largely associated with two different atmospheric circulation patterns (AO and AD) associated with an asymmetry in the anomalous SST distributions over the North Pacific. The teleconnections between the Arctic sea ice
- 375 variability and mid-latitude SST anomalies suggest that on a decadal or longer time scale it may be necessary to include the Arctic sea ice and mid-latitude SST interactions or feedbacks in any investigations of Arctic warming and sea ice decline and their potential influence on mid-latitude weather and climate, an area of active research in recent years (Barnes and Screen, 2015; Overland and Wang, 2015; Francis and Skific, 2015). Our results are based solely on statistical analyses, which do not

380

suggest causal relationship. Studies using coupled ocean-atmosphere models are necessary to confirm the impact of the SST anomalies over North Pacific on northern high-latitude atmospheric circulations. Finally, the results here help advance the knowledge about the relatively large contributions from the decadal-scale SST variability to Arctic climate change, though further studies using coupled global atmosphere-ocean-sea ice models are warranted to fully understand the physical mechanisms.

385 Data Availability

All data used in the current study are publicly available. The monthly sea ice concentration data are available from the National Snow and Ice Data Center (NSIDC) (http://nsidc.org/data/NSIDC-0051), the ERA-Interim reanalysis data are available from the European Center for Mid-Range Weather Forecasting (https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era-interim) and the SST

390 data are available from the Hadley Centre for Climate Prediction and Research (http://www.metoffice.gov.uk/hadobs/hadisst/).

Competing interests

The authors declare that they have no conflict of interest.

Author Contributions

395 L. Yu designed the study, with input from S. Zhong, and carried out the analyses. L. Yu and S. Zhong prepared the manuscript.

Acknowledgements

This study is supported by National Key R&D Program of China (No.2017YFE0111700).

References

- Barnes, E. A., and Screen, J.: The impact of Arctic warming on the midlatitude jet stream: Can it? Has it?
 Will it? WIREs Clim. Change, 6, 277-286, <u>https://doi.org/10.1002/wcc.337</u>, 2015.
 - Bintanja, R., Graversen, R. G., and Hazeleger, W.: Arctic winter warming amplified by the thermal inversion and consequent low infrared cooling to space, Nat. Geosci., 4, 758-761, <u>https://doi.org/10.1038/ngeo1285</u>, 2011.
- 405 Chu, J.-E., Hameed, S. N., and Ha, K.-J.: Non-linear intraseasonal phases of the East Asian summer monsoon: Extraction and analysis using self-organizing maps, J. Climate, 25, 6975-6988, <u>https://doi.org/10.1175/JCLI-D-11-00512.1</u>, 2012.
 - Day, J. J., Hargreaves, J. C., Annan, J. D., and Abe-Ouchi, A.: Sources of multidecadal variability in Arctic sea ice extent, Environ. Res. Lett., 7, 034011, <u>https://doi.org/10.1088/1748-9326/7/7/034011</u>,
- 410 2012.
 - Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U.,

Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L.,
Bidlot, J., Bormaan, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy,
S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally,
A. P., Mong-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C.,
Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the

- data assimilation system, Q. J. R. Meteorol. Soc., 137, 553-597, <u>https://doi.org/10.1002/qj.828</u>, 2011.
- Deser, C., and Teng, H.: Evolution of Arctic sea ice concentration trends and the role of atmospheric circulation forcing, Geophys. Res. Lett., 35, L02504, <u>https://doi.org/10.1029/2007GL032023</u>, 2008.
 - Ding, Q., Wallace, J. M., Battisti, D. S., Steig, E. J., Gallant, A. J. E., Kim, H.-J., and Geng, L.: Tropical forcing of the recent rapid Arctic warming in northeastern Canada and Greenland, Nature, 509, 209–212, <u>https://doi.org/10.1038/nature13260</u>, 2014.
- 425 Ding, Q., Schweiger, A., L'Heureux, M., Battisti, D. S., Po-Chedley, S., Johnson, N. C., Blanchard-Wrigglesworth, E., Harnos, K., Zhang, Q., Eastman, R., and Steig, E. J.: Influence of high-latitude atmospheric circulation changes on summertime Arctic sea ice, Nat. Clim. Change, 7, 289-295, <u>https://doi.org/10.1038/NCLIMATE3241</u>, 2017.

Enfield, D. B., Mestas-Nunez, A. M., and Trimble, P. J.: The Atlantic Multidecadal Oscillation and its

- relationship to rainfall and river flows in the continental US, Geophys. Res. Lett., 28, 2077–2080,
 https://doi.org/10.1029/2000GL012745, 2001.
 - Flanner, M. G., Shell, K. M., Barlage, M., Perovich, D. K., and Tschudi, M. A.: Radiative forcing and albedo feedback from the Northern Hemisphere cryosphere between 1979 and 2008, Nat. Geosci., 4, 151–155, <u>https://doi.org/10.1038/NGEO1062</u>, 2011.
- 435 Francis, J. A., Chan, W., Leathers, D. J., Miller, J. R., and Veron, D. E.: Winter Northern Hemisphere weather patterns remember summer Arctic sea-ice extent, Geophys. Res. Lett., 36, L07503, <u>https://doi.org/10.1029/2009GL037274</u>, 2009.
 - Francis, J. A., and Skific, N.: Evidence linking rapid Arctic warming to mid-latitude weather patterns, Phil. Trans. R. Soc. A., 373, 20140170, <u>https://doi.org/10.1098/rsta.2014.0170</u>, 2015.
- Gagné, M. –È., Gillett, N. P., and Fyfe, J. C.: Impact of aerosol emission controls on future arctic se ice cover, Geophys. Res. Lett., 42, 8481-8488, <u>https://doi.org/10.1002/2015GL065504</u>, 2015.

415

- Gong, T., Feldstein, S., and Lee, S.: The Role of Downward Infrared Radiation in the Recent Arctic Winter Warming Trend. J. Climate, 30, 4937–4949, <u>https://doi.org/10.1175/JCLI-D-16-0180.1</u>, 2017.
- Grotjahn, R., Black, R., Leung, R., Wehner, M. F., Barlow, M., Bosilovich, M., Gershunov, A.,
 <u>Gutowski Jr, W. J., Gyakum, J. R., Katz, R. W., Lee. Y.-Y., Lim, Y.-K., and Prabhat: North</u>
 <u>American extreme temperature events and related large scale meteorological patterns: a review</u>
 <u>of statistical methods, dynamics, modelling, and trends, Clim. Dyn., 46, 1151-1184, 2016,</u>
 <u>doi:10.1007/s00382-015-2638-6.</u>
- Hewitson, B. C., and Crane, R. G.: Self-organizing maps: Applications to synoptic Climatology, Clim.
 Res., 22, 13-26, 2002.
 - Higgins, R. W., and Mo, K. C.: Persistent North Pacific circulation anomalies and the tropical intraseasonal circulation, J. Climate, 10, 223-244, 1997.

Hoskins, B. J., and Karoly, D. J.: The steady linear response of a spherical atmosphere to thermal and

455 orographic forcing, J. Atmos. Sci., 38, 1179-1196.

Jin, F., and Hoskins, B. J.: The direct response to tropical heating in a baroclinic atmosphere, J. Atmos. Sci., 50, 1661-1671, <u>https://doi.org/10.1175/1520-0469(1995)052<0307:TDRTTH>2.0.CO;2.</u> 1995.

Johnson, N. C., Feldstein, S. B., and Tremblay, B.: The continuum of Northern Hemisphere

teleconnection patterns and a description of the NAO shift with the use of self-organizing maps, J.
 Climate, 21, 6354-6371, <u>https://doi.org/10.1175/2008JCLI2380.1</u>, 2008.

Johnson, N. C.: How many ENSO flavors can we distinguish?, J. Climate, 26, 4816-4827, https://doi.org/10.1175/JCLI-D-12-00649.1, 2013.

Kim, B.-M., Hong, J.-Y., Jun, S.-Y., Zhang, X., Kwon, H., Kim, S.-J., Kim, J.-H., Kim, S.-W., and

- Kim, H.-K.: Major cause of unprecedented arctic warming in January 2016: Critical role of an Atlantic windstorm, Sci. Rep., 7, 40051, <u>https://doi.org/10.1038/srep40051</u>, 2017.
 - Koenigk, T., Mikolajewicz, U., Jungclaus, J., and Kroll, A.: Sea ice in the Barents Sea: Seasonal to interannual variability and climate feedbacks in a global coupled model, Clim. Dyn., 32, 1119–1138, <u>https://doi.org/10.1007/s00382-008-0450-2</u>, 2009.
- Kohonen, T. (Eds): Self-organizing maps, 3rd ed. Springer, Berlin, Germany, 501pp., 2001.
 Kug, J.-S., Jeong, J.-H., Jang, Y.-S., Kim, B.-M., Folland, C. K., Min, S.-K., and Son, S.-W.: Two 16

distinct influences of Arctic warming on cold winters over North America and East Asia, Nat. Geosci., 8, 759-762, <u>https://10.1038/NGEO2517</u>, 2015.

Lee, S., and Feldstein, S. B.: Detecting ozone-and greenhouse gas-driven wind trends with observational

475 data, Science, 339, 563-567, <u>https://doi.org/10.1126/science.1225154</u>, 2013.

Lee, S. T., Gong, T., Johnson, N. C., Feldstein, S. B., and Polland, D.: On the possible link between tropical convection and the Northern Hemisphere Arctic surface air temperature change between 1958 and 2001, J. Climate, 24, 4350-4367, <u>https://doi.org/10.1175/2011JCLI4003.1</u>, 2011.

Lee, S., Gong , T., Feldstein, S. B., Screen, J. A., and Simmonds, I.: Revisiting the cause of the 1989-

- 2009Arctic surface warming using the surface energy budget: Downward infrared radiation dominates the surface fluxes. Geophysical Research Letters, 44, 10654–10661. <u>https://doi.org/10.1002/2017GL075375</u>, 2017.
 - Leloup, J., Lachkar, Z., Boulanger, J.-P., and Thiria, S.: Detecting decadal changes in ENSO using neural networks, Clim. Dyn., 28, 147-162, <u>https://10.1007/s00382-006-0173-1</u>, 2007.
- 485 Liebmann, B., and Smith, C.A.: Description of a Complete (Interpolated) Outgoing Longwave Radiation Dataset, *Bulletin of the American Meteorological Society*, **77**, 1275-1277, 1996.
 - Mantua, N. J., Hare, S. R., Zhang, Y., Wallace, J. M., and Francis, R. C.: A Pacific interdecadal climate oscillation with impacts on salmon production, Bull. Am. Meteorol. Soc., 78, 1069–1079, https://10.1175/1520-0477(1997)078<1069:APICOW>2.0.CO;2, 1997.
- 490 Min, S.-K., Zhang, X., Zwiers, F. W., and Agnew, T.: Huamn influence on arctic sea ice detectable from early 1990s onwards, Geophys. Res. Lett., 35, 213-36, <u>https://10.1029/2008GL035725</u>, 2008.
 - Mori, M., Watanabe, M., Shiogama, H., Inoue, J., and Kimoto, M.: Robust Arctic sea-ice influence on the frequent Eurasian cold winters in past decades, Nat. Geosci., 7, 869-873, <u>https://doi.org/10.1038/NGEO2277</u>, 2014.
- 495 Notz, D., and Marotzke, J.: Observations reveal external driver for Arctic sea-ice retreat, Geophys. Res. Lett., 39, 89-106, <u>https://doi.org/10.1029/2012GL051094</u>, 2012.

Overland, J. E., and Wang, M.: Increased variability in the early winter subarctic North American atmospheric circulation, J. Climate, 28, 7297-7305, <u>https://doi.org/10.1175/JCLI-D-15-0395.1</u>,2015.

Park, W., and Latif, M.: Multidecadal and multicentennial variability of the meridional overturning circulation, Geophys. Res. Lett., 35, L22703, <u>https://doi.org/10.1029/2008GL035779</u>, 2008.

- Parkinson, C. L.: Global sea ice coverage from satellite data: Annual cycle and 35-Yr trends, J. Climate, 27, 9377–9382, <u>https://doi.org/10.1175/JCLI-D-14-00605.1</u>, 2014.
- Peings, Y., and Magnusdottir, G.: Response of the wintertime Northern Hemisphere atmospheric
- circulation to current and projected Arctic sea ice delcine: a numerical study with CAM5, J.
 Climate, 27, 244-264, <u>https://doi.org/10.1175/JCLI-D-13-00272.1</u>, 2014.
 - Petoukhov, V., and Semenov, V. A.: A link between reduced Barents-Kara sea ice and cold winter extremes over northern continents, J. Geophys. Res., 115, D211111, <u>https://doi.org/10.1029JD013568</u>, 2010.
- 510 Pithan, F., and Mauritsen, T.: Arctic amplification dominated by temperature feedbacks in contemporary climate models, Nat. Geosci., 7, 181-184, <u>https://doi.org/10.1038/NGEO2071</u>, 2014.
 - Post, E., Bhatt, U. S., Bitz, C. M., Brodie, J. F., Fulton, T. L., Hebblewhite, M., Kerby, J., Kutz, S. J., Stirling, I., and Walker, D. A.: Ecological consequences of sea-ice decline, Science, 341, 519–524, <u>https://doi.org/10.1126/science.1235225</u>, 2013.
- Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P., Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, J. Geophys. Res., 108, 4407, https://doi.org/10.1029/2002JD002670, 2003.

Rigor, I., Wallace, J. M., and Colony, R. L.: Response of sea ice to the Arctic Oscillation, J. Climate, 15,

- 520 2648–2663, <u>https://doi.org/10.1175/1520-0442(2002)015<2648:ROSITT>2.0.CO:2</u>, 2002.
 - Rothrock, D. A., Yu, Y., and Maykut, G. A.: Thinning of the Arctic sea-ice cover, Geophys. Res. Lett., 26, 3469–3472, 1999.
 - Sedlar, J., Tjernström, M., Mauritsen, T., Shupe, M. D., Brooks, I. M., Persson, P. Ola G., Birch, C. E., Leck, C., Sirevaag, A., and Nicolaus, M.: A transitioning Arctic surface energy budget: the
- impacts of solar zenith angle, surface albedo and cloud radiative forcing, Clim. Dyn., 37, 1643-1660, <u>https://doi.org/10.1007/s00382-010-0937-5</u>, 2011.
 - Serreze, M. C., Barrett, A. P., and Stroeve, J.: Recent changes in tropospheric water vapor over the Arctic as assessed from radiosondes and atmospheric reanalyses, J. Geophys. Res., 117, D10104, doi:10.1029/2011JD017421, 2012.
- Stroeve, J., Holland, M., Meier, W., Scambos, T., and Serreze, M.: Arctic sea ice decline: Fast than forecast, Geophys. Res. Lett., 34, L09501, <u>https://doi.org/10.1029/2007GL029703</u>, 2007.

Swart, N. C., Fyfe, J. C., Hawkins, E., Kay, J. E., and Jahn, A.: Influence of internal variability on Arctic sea-ice trends, Nat. Clim. Change, 5, 86-89, 2015.

Wang, J., Zhang, J., Watanabe, E., Ikeda, M., Mizobata, K., Walsh, J. E., Bai, X., and Wu, B.: Is the

535 Dipole anomaly a major driver to record lows in Arctic summer sea ice extent?, Geophys. Res.
 Lett., 36, L05706, <u>https://doi.org/10.1029/2008GL036706</u>, 2009.

Yu, L., Zhong, S., Winkler, J. A., Zhou, M., Lenschow, D. H., Li, B., Wang, X., and Yang, Q.: Possible connections of the opposite trends in Arctic and Antarctic sea-ice cover, Sci. Rep., 7, 45804, https://doi.org/10.1038/srep45804, 2017.

Table 1.	The 38	8-year (1	1979-201	6) averag	e of spatia	l correlations	between	the seasonal	Arctic sea
					, ,				

ice concentration and the corresponding SOM pattern for each year.

SOM grids	2×2	2×3	2×4	3×3	3×4	3×5	4×4	4×5	
Correlation coefficients	0.59	0.56	0.59	0.64	0.66	0.69	0.68	0.70	

Figure captions

Fig.1. The SOM patterns of autumn (September-November) Arctic sea ice concentration for a 3×3 grid for the 1979-2016 period. The percentages at the top left of each panel indicate the frequency of occurrence of the pattern.

Fig. 2. <u>Which mode best matches the observed sea ice concentration in each autumn</u>Occurrence time series for each SOM pattern in Fig. 1.

Fig. 3. In the statistical sense, <u>the linear</u> trends in the autumn Arctic sea ice concentration explained byassociated with each SOM pattern <u>when projected on the observations</u> (Units: yr⁻¹).

Fig. 4. Total (a), SOM-explained (b) and residual (c) autumn Arctic sea ice concentration trends (Units: yr⁻¹). Dots in (a) indicate above 90% confidence level.

Fig. 5. Composites of anomalous sea surface temperature (°C) and 500-hPa geopotential height (gpm) for nodes 1 (2007-2013, 2015-2016) and 9(1980-1982, 1986-1989, 1992, 1996). Dotted regions denote above 90% confidence level.

Fig. 6. The same as Fig. 5, but for 850-hPa wind field. Shaded regions denote above 90% confidence level.

Fig. 7. The same as Fig. 5, but for integrated atmospheric water vapor from surface to 750 hPa (g kg⁻¹), accumulated surface downward longwave radiation (W $m^{-2} \times 10^5$) and surface air temperature (°C). Dotted regions denote above 90% confidence level.

Fig. 8. Composites of anomalous sea surface temperature (°C) and 500-hPa geopotential height (gpm) for after 2001-2016 and before 20001979-2000. Dotted regions denote above 90% confidence level.



Figure 1 The SOM patterns of the autumn (September-November) Arctic sea ice concentration for a 3×3 grid for the 1979-2016 period. The percentages at the top left of each panel indicate the frequency of occurrence of the pattern.







Figure 3. In the statistical sense, <u>the linear</u> trends in the autumn Arctic sea ice concentration <u>explained by associated with</u> each SOM pattern <u>when projected on the observations</u> (Units: yr^{-1}).



Figure 4. Total (a), SOM-explained (b) and residual (c) autumn Arctic sea ice concentration trends. (Units: yr^{-1}). Dots in (a) indicate above 95% confidence level.



Figure 5. Composites of anomalous sea surface temperature (°C), outgoing long wave radiation (OLR) (W m^{-2}) and 500-hPa geopotential height (gpm) for nodes 1 (2007-2013, 2015-2016) and 9 (1980-1982, 1986-1989, 1992, 1996). Dotted regions denote above 90% confidence level.



Figure 6. The same as Figure 5, but for 850 hPa wind field. Shaded regions denote above 90% confidence level.



Figure 7. The same as Figure 5, but for integrated atmospheric water vapor from surface to 750 hPa (g kg⁻¹), accumulated surface downward longwave radiation $(10^5 Wm^{-2})$, and surface air temperature (°C). Dotted regions denote above 90% confidence level.



Figure 8. Composites of anomalous sea surface temperature (°C) and 500-hPa geopotential height (gpm) for <u>after-2001-2016</u> and <u>before 20001979-2000</u>. Dotted regions denote above 90% confidence level.