Supplement of

Modeling the impact of heterogeneous reactions of chlorine on summertime nitrate formation in Beijing, China

Xionghui Qiu^{1,2}, Qi Ying^{3*}, Shuxiao Wang^{1,2*}, Lei Duan^{1,2}, Jian Zhao⁴, Jia

Xing^{1,2}, Dian Ding^{1,2}, Yele Sun⁴, Baoxian Liu⁵, Aijun Shi⁶, Xiao Yan⁶,

Qingcheng Xu^{1,2}, Jiming Hao^{1,2}

¹ State Key Joint Laboratory of Environmental Simulation and Pollution Control, School of Environment, Tsinghua University. Beijing 100084, China.

² State Environmental Protection Key Laboratory of Sources and Control of Air Pollution Complex, Beijing 100084, China

³ Zachry Department of Civil Engineering, Texas A&M University, College Station Texas, United States.

⁴ State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

⁵ Beijing Environmental Monitoring Center, Beijing 100048, China

⁶ Beijing Municipal Research Institute of Environmental Protection, Beijing 100037, China

*Corresponding author: shxwang@tsinghua.edu.cn & qying@civil.tamu.edu

Table S1 Emissions of conventional chemical species in beijing in 2017 (Unit: t)					
Sector	SO_2	NO _X	PM _{2.5}	PM ₁₀	VOCs
Power Plant	842	6267	675	823	70
Industry	7421	19867	4545	7059	222046
Residential	12121	10472	14614	19424	53938
Transportation	2365	92024	5080	5255	69908

 Table S1 Emissions of conventional chemical species in Beijing in 2017 (Unit: t)

Figure S1 The spatial distribution of pH in June

