



Fakultät für Physik und Geowissenschaften Institut für Meteorologie Johannes Mülmenstädt

Universität Leipzig, Institut für Meteorologie, PF 232101, 04081 Leipzig

July 8, 2019

Dear Xiahong,

attached, please find our responses to the reviewers as well as a marked-up version of the manuscript showing the changes we have made. In addition to the recommendations of the reviewers, a number of small improvements to the language have found their way into the new version.

Best regards,

Johannes (for the authors)

We thank Steve Ghan for his thorough reading of the manuscript and helpful comments. Please find our responses inline below.

Page 1, line 4. Insert "by anthropogenic cloud droplet number change" after "radiative forcing".

Thank you for suggesting this clarification. We have adopted the change in the revised manuscript.

Page 3, lines 15–16. Consider the decomposition expressed by equations 6–8 in Ghan et al. PNAS 2016.

Thank you for pointing out the imprecise wording in this sentence. Our intended meaning was to differentiate between, on the one hand, "exact" methods that use the time-varying, three-dimensional state of the model (i.e., the model's direct knowledge of the anthropogenic perturbations); and, on the other hand, methods that require idealizing the cloud as globally homogeneous or performing statistical analysis such as linear regression on the model output. We have noted this in the revised manuscript. We have also changed "exact" to "direct", since we show later on that our method still carries uncertainties on the order of 0.1 W m⁻².

Page 4. I'm concerned about substituting a cloud property from one run into diagnostic radiation calculations from another run, since cloud properties vary in time. What is done when clouds at a particular time are simulated in one run but not in the other. How is the cloud property determined then? Using time mean property will work if cloud forms at least once at that point, but what if it never forms at that point in one simulation but does in the other? This issue is mentioned later: large artifacts that occur due to the decorrelated cloud property fields, and tested in the Appendix, but it does not address the question of how to specify properties of clouds not present in one simulation.

We were concerned, too, and we suspect this problem has dissuaded others from trying PRP earlier. When clouds are absent in one run and present in the other, we let the radiative transfer resolve the conflicting cloud properties in the same way as it does when the cloud microphysics and cloud cover schemes produce conflicting cloud properties, i.e., effective radii can only vary within the limits of the cloud optics lookup table. This is guaranteed to produce incorrect results for the model column in which the mismatch of cloud properties occurs; in fact, the correct result is probably undefined. However, we would consider this to be the heart (or perhaps the logical extreme) of the decorrelation problem. This is the reason we designed the tests in Sec. A2, where the correct forcing components are known, and found that the forward–backward PRP results agree with the correct values to within 0.1 W m^{-2} accuracy, as you point out in your comment.

We agree that the issue of cloud presence in one run and absence in the other should be discussed in the text, and that the discussion should include a prescription for what to do when this case occurs. We have expanded the revised manuscript accordingly.

Page 6, line 7. Insert "global mean" before "forcing".

Thank you for suggesting this clarification. We have adopted the change in the revised manuscript.

Page 9, lines 29–32. Is PRP the most direct method? Is it more direct than the method described

by Ghan et al. PNAS 2016? Why not compare the two methods? The Ghan method is simple to implement.

Thank you for pointing out the imprecise wording in this sentence. "Direct" was meant in the same way that "exact" was meant on p. 3, l. 15–16; we have clarified this in the revised manuscript. The suggestion of an intercomparison of methods is a good one, in particular as several additional decomposition methods are close to publication (Gryspeerdt et al., submitted, and at least one other study, private comm.). In the revised manuscript, we mention that the Ghan et al. (2016) estimate of $F_{\mathcal{L}}/F_{N_d} \approx 5$ is much greater than our result of ≈ 1 . However, we feel that tracking down the sources of differences between methods is best left for a dedicated intercomparison study.

Figure 2 Caption has a question.

Thank you for pointing out this leftover editing detritus. We have removed it from the manuscript.

We thank the reviewer for his or her thorough reading of the manuscript and helpful comments. Please find our responses inline below.

Page 2, line 17: In RFari you are using the ARI abbreviation for the first time. You should more clearly spell out what this abbreviation means (even though you mention the phrase "aerosol–radiation" in the line above).

Thank you for pointing out this omission. We have defined the abbreviation in the revised manuscript.

Page 3, Line 1: On a similar note, please formally define f_c as cloud fraction. I don't see it defined anywhere.

Thank you for pointing out this omission, as well. We have defined the variable in the revised manuscript.

Page 3, line 17: "... but this decomposition does not correspond to the forcing-and- adjustment decomposition." More or clearer explanation about why APRP does not fit the forcingadjustment framework would be helpful. This was a bit vague.

We thank the reviewer for pointing out the vague language. In the revised manuscript, we now explain that APRP decomposes the cloud property changes into changes in area fraction, cloud albedo, and cloud absorption. The change in area fraction maps well onto the cloud fraction adjustment in the forcing-adjustment framework, but the APRP cloud albedo change includes both the effect of the anthropogenic N_d change and the \mathcal{L} adjustment.

Page 6, line 20. A specific example reference of the observational studies you talk about would be helpful here.

We have expanded the discussion in this paragraph, also in light of the reviewer's comment on Fig. 1 below. We now separately discuss changes over ocean and over land and compare the patterns we have derived for each to a number of observational or observationally constrained modeling studies. We have also factorized the geographic distributions of the ERFaci components into an anthropogenic N_d perturbation and a model sensitivity to the perturbation, to facilitate comparison to observational estimates of aerosol susceptibilities.

Table 1: Does RFari account perfectly for the difference between the sum of the ERFaci and the total ERF? Or is there some error associated with the PRP method in that difference? It would be good to quantify RFari. Perhaps with double-call calculations or the Ghan method.

Very good point; the RFari and ice-cloud ACI are not part of our decomposition, but they are estimated for a very similar model run in Gryspeerdt et al. (submitted; should be in ACPD by the time this manuscript is published). The total ERF also includes the ice-phase ACI effects (-0.59 W m^{-2} in the SW, 0.88 W m⁻² in the LW), RFari (-0.17 W m^{-2} in the SW), and a negligible surface-albedo contribution (-0.01 W m^{-2}). The sum of the components thus balances at approximately the 0.2 W m⁻² level, a relative error similar to the 0.1 W m⁻² estimated uncertainty on the ERFaci components.

Figure 1: Any explanation for the local maximum in forcing/adjustment terms along the eastern

boundary currents? Right along the west coast of N. America, S. America and Europe? It seems these are also regions where the backwards and forwards PRP calculations differ notably (Figure A1)

We have expanded the discussion in Sec. 3.1 to better describe and explain these local maxima: these are regions where low clouds are abundant (\mathcal{L} is large) and anthropogenic aerosols mix from the continents into the cleaner maritime air masses.

Regarding the last sentence of the comment, the forcing estimate, by construction, is highest where the differences between forward and backward PRP are greatest.

Figure 2: The caption seems to include an editing note by accident.

Thank you for pointing out this leftover editing detritus. We have removed it from the manuscript.

A3: I'd prefer the appendix discussion and figure about temporal averaging to be included in the main section of the note, especially since it is given a prominent spot in the abstract. Given the recent push for large model comparison projects to include forcing diagnosis (where temporally averaged data is the norm), this result seems important.

We agree; it is a bit strange to have to consult the appendix for one of the main points of the paper. We have moved Appendix 3 into the results section (Sec. 3).

Separating radiative forcing by aerosol–cloud interactions and fast rapid cloud adjustments in the ECHAM–HAMMOZ aerosol–climate model using the method of partial radiative perturbations

Johannes Mülmenstädt¹, Edward Gryspeerdt², Marc Salzmann¹, Po–Lun Ma³, Sudhakar Dipu¹, and Johannes Quaas¹

¹Universität Leipzig, Leipzig, Germany

²Space and Atmospheric Physics Group, Imperial College London, London, United Kingdom ³Pacific Northwest National Laboratory, Richland, Washington, USA

Correspondence: Johannes Mülmenstädt (johannes.muelmenstaedt@uni-leipzig.de)

Abstract. Using the method of offline radiative transfer modelling within the partial radiative perturbations (PRP) approach, the effective radiative forcing (ERF) by aerosol-cloud interactions (ACIERF_{aci}) in the ECHAM–HAMMOZ aerosol climate model is decomposed into a radiative forcing by anthropogenic cloud droplet number change and adjustments of the liquid water path and cloud fraction. The simulated radiative forcing by anthropogenic cloud droplet number change and liquid water

- 5 path adjustment are of approximately equal magnitude at -0.52 W m^{-2} and -0.53 W m^{-2} , respectively, while the cloud fraction adjustment is somewhat weaker at -0.31 W m^{-2} (constituting 38%, 39%, and 23% of the total ERFaciERF_{aci}, respectively); geographically, all three ERF-ERF_{aci} components in the simulation peak over China, the subtropical eastern ocean boundaries, the northern Atlantic and Pacific Oceanoceans, Europe, and eastern North America (in order of prominence). Spatial correlations indicate that the temporal-mean liquid water path adjustment is proportional to the temporal-mean radiative forcing,
- 10 while the relationship between cloud fraction adjustment and radiative forcing is less direct. While the estimate of warm-cloud ACI-ERF_{aci} is relatively insensitive to the treatment of ice and mixed-phase cloud overlying warm cloud, there are indications that more restrictive treatments of ice in the column result in a low bias in the estimated magnitude of the liquid water path adjustment and a high bias in the estimated magnitude of the droplet number forcing. Since the present work is the first PRP decomposition of the aerosol ERF into and fast effective radiative forcing into radiative forcing and rapid cloud adjustments,
- 15 idealized experiments are conducted to provide evidence that the PRP results are accurate. The experiments show that using low-frequency (daily or monthly) time-averaged model output of the cloud property fields underestimates the ERF, but 3-hourly three-hourly mean output is sufficiently frequent.

Copyright statement. TEXT

1 Introduction

Following Boucher et al. (2013), it has become common to distinguish between radiative forcing (RF) by aerosol-cloud interactions (ACI) and fast-rapid adjustments to this radiative forcing, with the sum of the forcing (RF_{aci} and the fast-) and the rapid adjustments denoted as the effective radiative forcing (ERF_{aci}). In liquid-water clouds, RF_{aci} arises from the increased

5 availability of cloud condensation nuclei (CCN) in a polluted atmosphere leading to higher droplet number N_d and smaller effective radius r_e at constant cloud liquid water path \mathcal{L} (Twomey, 1977). The atmosphere responds to the higher- N_d , lower- r_e clouds by various processes occurring on short timescales, leading to adjustments in other cloud properties, including \mathcal{L} and cloud vertical and horizontal geometric extent.

Physically, the most important adjustment mechanisms are suppression of precipitation formation (Albrecht, 1989) and en-

- 10 hanced cloud-edge dry-air entrainment and droplet evaporation in the smaller-droplet clouds (Ackerman et al., 2004). The former mechanism, in isolation, would lead to an increase in cloud condensate and cloud fraction, and thus a negative adjustment to the radiative forcing; the latter, in isolation, would lead to a decrease in cloud condensate and thus a positive adjustment to the radiative forcing. Since no mechanism occurs in isolation in a coupled system (Stevens and Feingold, 2009), the question of whether the net adjustment is positive or negative is both difficult and unresolved (e.g., Mülmenstädt and Feingold, 2018; ?)(e.g., Mülmenst
- 15 One method by which aerosol effective radiative forcing is estimated is to calculate the difference in top-of-atmosphere (TOA) radiative fluxes in general-circulation models (GCMs) between runs with fixed sea-surface temperature and present-day or preindustrial aerosol concentrations or emissions (e.g., Lohmann et al., 2010; Forster et al., 2016). This requires a GCM that includes the relevant aerosol-radiation and aerosol-cloud interaction mechanisms. By representing the optical properties of aerosol in the radiative transfer, the radiative forcing by aerosol-radiation interactions (RF_{ari}-) can be estimated; by representing
- 20 the N_d dependence on aerosol activation during cloud formation, RF_{aci} can be estimated; and by representing precipitation suppression and enhanced evaporation in smaller- r_e clouds, the adjustments to RF_{aci} can be estimated. (We exclude ice and mixed-phase cloud processes, which introduce further complications, from this discussion.) These processes occur on scales far below the resolved scale, so their representation in the GCM requires parameterization. Thus, the model is only imperfectly (if at all) aware of subgrid-scale variability in the process rates and feedbacks between the processes; relies on imperfect base-
- 25 state cloud properties (e.g., Penner et al., 2006); and only considers effects that are amenable to parameterization, meaning that precipitation suppression is included in many models but enhanced evaporation is not (e.g., Salzmann et al., 2010; Michibata et al., 2016; Zhou and Penner, 2017). Based on these considerations, a prevalent view is that, from the standpoint of achieving GCM fidelity, ACI are more difficult than aerosol–radiation interactions, and ACI adjustments are more difficult than the ACI forcing. On top of this, the usual concerns about parametric uncertainty apply, so that the overall uncertainty on GCM estimates
- 30 of ERF_{aci} is large (Boucher et al., 2013). Nevertheless, the ability of GCMs to produce a global estimate will assure their star will continue to shine brightly in the firmament of ERF estimation methods until competing methods overcome their own significant drawbacks.

In general, one might argue that knowing the uncertainty on each term in a sum is a good first step towards attacking the uncertainty on the total; certainly, this is consistent with the GCM paradigm of building up the total forcing from parameteri-

zations for each of the contributing processes, even if it is less applicable to "top-down" estimates from the historical evolution of the climate system. Thus, we write the effective radiative forcing by aerosol as

$$F = F_{\rm ari} + F_{N_d} + F_{\mathcal{L}} + F_{f_c},\tag{1}$$

where F_{ari} is the RF_{ari}, F_{N_d} is the RF_{aci} due to the increase in N_d , and $F_{\mathcal{L}}$ and F_{f_c} are the \mathcal{L} and <u>cloud fraction</u> (f_c) ad-

- 5 justments to the RF_{ari} and RF_{aci}. Other adjustments, e.g., due to rapid changes in land surface temperatures or atmospheric temperature and humidity profiles, have been estimated as small in previous studies a previous study (Heyn et al., 2017). Each of these terms maps fairly well onto a parameterization in the GCM: RF_{ari} is parameterized in the radiative transfer, F_{N_d} is parameterized in a droplet activation scheme, the ACI part of $F_{\mathcal{L}}$ is parameterized in the precipitation microphysics (and, if enhanced evaporation becomes tractable in the future, that component will presumably be parameterized in the turbulence
- 10 scheme; e.g., Guo et al., 2011; Neubauer et al., 2014), and F_{f_c} is parameterized in the cloud cover scheme (although in our model the response to the perturbation is indirect, via relative humidity changes subsequent to precipitation rate changes); the only component that emerges from the model dynamics rather than from an explicit parameterization are is the adjustments of temperature-temperature and moisture profiles that entail further adjustments to aerosol-cloud interactions, and to aerosol-radiation aerosol-cloud interactions and to aerosol-radiation interactions (formerly known as "semi-direct effect").
- 15 Both; both of these terms are small (Heyn et al., 2017; Stjern et al., 2017). A natural first step would then be to ask how large each of these terms is, and a natural second step would be to ask how much uncertainty each contributes to the total. One of the benefits of such a decomposition would be that it would provide a more solid footing for—or falsify—the notion that models agree fairly well on the "simpler" problem of RF_{aci} and not well at all on the "harder" problem of the adjustments. However, performing the decomposition is quite difficult in practice. Ghan (2013) addresses the issue of separating
- 20 F_{ari} precisely from $F_{N_d} + F_{\mathcal{L}} + F_{f_c}$, but methods using the model's intrinsic knowledge of the anthropogenic perturbations. Methods to separate the latter three are far less exact three ERF_{aci} components using the intrinsic model knowledge of the time-varying, three-dimensional aerosol perturbation and resulting perturbation of cloud properties need to contend with the problems that the ERF_{aci} is diagnosed from two separate runs that cannot easily share fields online, so that double radiation calls as in Ghan (2013) are not feasible; and that the adjustments, by definition, do not respond to the aerosol perturbations
- 25 instantaneously. One possibility is APRP decomposition into cloud amount, scattering, and absorption (Zelinka et al., 2014), but this decomposition does not correspond to the foreing-and-adjustment decompositionthe approximate partial radiative perturbation (APRP) decomposition (Taylor et al., 2007; Zelinka et al., 2014). APRP decomposes the cloud property changes into changes in area fraction, cloud albedo, and cloud absorption; the change in area fraction maps well onto the f_c adjustment in the forcing–adjustment framework, but the APRP cloud albedo change includes both the effect of the anthropogenic N_d
- 30 change and the L adjustment. Another possibility is to deactivate the parameterized precipitation suppression (and, if models include a parameterization of the enhanced evaporation, deactivate that as well); however, the model without with parameterized adjustments will produce the a different climate than the model with (Penner et al., 2006). without (Penner et al., 2006). Due to the complications arising in methods that directly use the model state, less direct methods have been developed that

idealize the cloud field as a globally homogeneous single layer (Ghan et al., 2016) or use similar regression-based statistical techniques as satellite studies (Gryspeerdt et al., submitted).

In this technical notework, we apply the method of partial radiative perturbations (PRP, Wetherald and Manabe, 1988; Colman and McAv the ERF decomposition problem. PRP falls in the category of methods that directly use the intrinsic model knowledge of the

- 5 time-varying, three-dimensional aerosol perturbation and resulting perturbation of cloud properties. The starting point for PRP is a perturbed and an unperturbed model run. One then introduces the fields of the perturbed run into the unperturbed run, one at a time, and reruns the radiative transfer scheme on the "partially perturbed" state to derive the resulting change in radiative fluxes. In our application, the two runs are fixed-SST runs (simulations with an atmospheric GCM with prescribed climatological, seasonally varying sea surface temperature (SST) and sea ice cover distributions) (SIC) distributions with present-day and
- 10 preindustrial aerosol emissions, nudged to reanalysis-present-day large-scale upper-level winds to reduce the internal variability without overconstraining the behavior of lower-tropospheric warm cloud and allow significant changes in cloud property to emerge after a shorter integration time than would otherwise be required (e.g., Kooperman et al., 2012; Zhang et al., 2014). The perturbed fields are N_d , \mathcal{L} , and f_c ; the corresponding changes in radiative fluxes are F_{N_d} , $F_{\mathcal{L}}$, and F_{f_c} .

In Section 2, we describe the PRP method and ECHAM-HAMMOZ model in detail; in Section 3, we use PRP to estimate

15 the ERF components in the ECHAM–HAMMOZ model , and determine whether the adjustments are a simple proportional response to the forcing.

2 Methods

25

We first give a brief formal description of the PRP method; then we we then describe the model configurations to which we will apply the method.

20 2.1 Partial radiative perturbations

We denote the shortwave TOA flux as Q and the longwave flux as R (all-sky, positive downward in both cases). For the purposes of this analysis, the radiative flux in each spectral range is considered a function of the cloud properties N_d , \mathcal{L} (or the vertically resolved analogue q_l), and f_c . The dependence of the fluxes on other climate state variables water variables – water variables –

 $Q(\lambda,\phi,t) = Q(N_d(\lambda,\phi,p,t),q_l(\lambda,\phi,p,t),f_c(\lambda,\phi,p,t))$ ⁽²⁾

$$R(\lambda,\phi,t) = R(N_d(\lambda,\phi,p,t),q_l(\lambda,\phi,p,t),f_c(\lambda,\phi,p,t))$$
(3)

Let $\mathbf{x}^A = \{N_d^A, \mathcal{L}^A, f_c^A\}$ and $\mathbf{x}^B = \{N_d^B, \mathcal{L}^B, f_c^B\}$ denote the cloud properties in runs *A* and *B*. We then define *forward* and *backward* PRP as inserting one cloud property at a time from one run into the cloud field of the other and recalculating the

radiative fluxes:

$$\delta_{A \to B} Q_{i\xi} = Q(\{x_{i\xi}^{A}, x_{j \neq i\zeta \neq \xi}^{B}\}) - Q(\mathbf{x}^{B})$$

$$\tag{4}$$

$$\delta_{B \to A} Q_{\underline{i}\underline{\xi}} = Q(\{x_{\underline{i}\underline{\xi}}^{B}, x_{\underline{j} \neq \underline{i}\underline{\zeta} \neq \underline{\xi}}^{A}\}) - Q(\mathbf{x}^{A}), \tag{5}$$

where cloud property $i \notin \xi$ is substituted from run *A* into run *B* or run *B* into run *A*, respectively, and *Q* (or *R*, analogously) is 5 recalculated using the offline version of the model's radiative transfer scheme. *Forward–backward* PRP is simply the average of the two, taking into consideration that reversing direction reverses the sign of the radiative-flux perturbation (e.g., Klocke et al., 2013):

2

• •

10

25

When A denotes the PI-emissions preindustrial (PI)-emissions run and B denotes the PD-emissions present-day (PD)-emissions run, the components of ERF_{aci} correspond to

$$F_{N_d} = \overline{\delta_{\text{PI}\leftrightarrow\text{PD}}Q_{N_d} + \delta_{\text{PI}\leftrightarrow\text{PD}}R_{N_d}} \tag{7}$$

15
$$F_{\mathcal{L}} = \overline{\delta_{\text{PI} \leftrightarrow \text{PD}} Q_{\mathcal{L}} + \delta_{\text{PI} \leftrightarrow \text{PD}} R_{\mathcal{L}}}$$
(8)

$$F_{f_c} = \delta_{\text{PI}\leftrightarrow\text{PD}}Q_{f_c} + \delta_{\text{PI}\leftrightarrow\text{PD}}R_{f_c}.$$
(9)

For other meanings of A and B, as in the additional experiments performed in Sections 3.1–3.3, the equivalent expressions to Equations (7)–(9) describe pseudo-forcing components rather than forcing components; we denote them as \tilde{F}_{N_d} , $\tilde{F}_{\mathcal{L}}$, and \tilde{F}_{f_c} . In Equations (7)–(9),

20
$$\overline{\mathcal{F}} = \frac{1}{N} \sum_{i=1}^{N} \mathcal{F}(t_i)$$
 (10)

indicates averaging over the time dimension of a field \mathcal{F} evaluated at the *N* time steps $\{t_1, \ldots, t_N\}$. "Evaluated" can, itself, refer to a temporal average over the interval between evaluation time steps, as in a 3-hourly or daily mean, or it can refer to the instantaneous value of the field at that time step; when the distinction matters (because *Q* and *R* are not linear functions of their input variables), we will indicate the averaging interval as $\overline{\mathcal{F}}^{(\Delta t)}$. Thus, $\overline{\mathcal{F}}^{(\text{inst})}$ denotes the temporal mean of instantaneous model output, while $\overline{\mathcal{F}}^{(3 \text{ h})}$ denotes the temporal mean of 3-hourly-averaged model output.

5

2.2 Model description

We use several model runs performed with the ECHAM–HAMMOZ model, version echam6.1–ham2.2–moz0.9 (Neubauer et al., 2014). The model is based on the ECHAM atmospheric general circulation model (Stevens et al., 2013), the HAM interactive aerosol module (Stier et al., 2005; Zhang et al., 2012), and the trace-gas chemistry module MOZ (Kinnison et al., 2007) (the latter

- 5 is disabled in these runs). Of most direct relevance to our study, the parameterized processes contributing to warm-cloudaerosol interactions are aerosol activation into cloud droplets according to Lin and Leaitch (1997); diagnostic warm rain processes (autoconversion and accretion) according to Khairoutdinov and Kogan (2000); and aerosol scavenging according to Croft et al. (2009, 2010). The stratiform cloud scheme is that of Lohmann and Roeckner (1996), extended to double-moment microphysics by Lohmann et al. (2007); Lohmann and Hoose (2009)Lohmann et al. (2007) and Lohmann and Hoose (2009),
- 10 with the Sundqvist et al. (1989) critical-relative-humidity cloud cover parameterization.

To reduce internal variability and achieve low statistical uncertainty on the forcing components within a reasonable integration time, we use monthly varying but yearly repeating sea surface temperatures (SST) and sea ice concentrations (SIC) SST and SIC from the observed climatology and nudge the large-scale wind fields to the present-day ERA-Interim reanalysis (?) wind fields of the years 2000–2010 (in some sensitivity studies, only the year 2000 is used).

15 Estimates of radiative forcing are computed by performing a pair of model runs with present-day SST, SIC, and wind fields, and aerosol (precursor) emissions estimates for either the year 2000 or the year 1850. Emissions are from the AEROCOM-II ACCMIP dataset; in particular, anthropogenic emissions follow Lamarque et al. (2010).

To perform PRP on the model output, we have updated the offline version of the RRTM-based ECHAM6 radiative transfer code (Pincus and Stevens, 2013) that was originally used in Klocke et al. (2013). We neglect time-varying aerosol-radiation

20 interactions to reduce technical complexity. To the extent that aerosol overlying cloud is a small effect, this choice mainly affects our estimate of the f_c adjustment (Ghan, 2013; Zelinka et al., 2014), which, unlike the N_d forcing and \mathcal{L} adjustment, is straightforward to compute without the PRP machinery; comparison to ? Gryspeerdt et al. (submitted) shows that the f_c adjustment estimate is not strongly affected by this simplification.

When clouds are absent (or the cloud fraction is very low) in one run and present in the other, perturbing cloud properties

- 25 can yield unrealistically large or small q_l or N_d (and thus r_e); this "decorrelation" problem is well known from the application to climate feedbacks (Colman and McAvaney, 1997) in the context of the correlation between water vapor and cloudiness. We allow the radiative transfer code to resolve the conflicting cloud properties in the same way as it does when the cloud microphysics and cloud cover schemes produce conflicting cloud properties; in particular, r_e can only vary within the limits of the cloud optics lookup table used by the model (2×10^{-6} to 32×10^{-6} m). Appendices A1 and A2 describe tests we performed
- 30 to verify that forward-backward PRP ERF_{aci} estimates are not impacted by the decorrelation problem.

3 Results

Since the components of the ERF_{aci} have not been diagnosed before in ECHAM–HAM by any method, we begin by presenting their global-mean values and geographic distributions in Section 3.1. In Section 3.2, we investigate whether fast responses rapid

adjustments to the Twomey forcing are proportional to F_{Nd} in terms of their spatial patternsare proportional to F_{Nd} . Section 3.3 investigates the sensitivity of the PRP results to the treatment of model columns containing ice and mixed-phase clouds. In Section 3.4, we determine how much temporal averaging is permissible before the PRP estimate becomes inaccurate. Sections A1 and A2 discuss whether PRP diagnoses the ERF_{aci} components correctly in the presence of decorrelation effect effects

5 (i.e., effects of introducing one cloud property from one run into an uncorrelated cloud field in another run). In Section 3.4, we determine what temporal averaging is permissible before the PRP estimate becomes inaccurate. Section 3.3 investigates the sensitivity of the PRP results to the treatment of model columns containing ice and mixed-phase clouds.

3.1 What are the ERF_{aci} components in ECHAM–HAM?

- Using the PRP decomposition, Equations (7)–(9), we can diagnose the contributions to the ERF_{aci} from PD and PI-emission fixed-SST model runs. This is shown in Figure 1 and Table 1. The longwave effective forcing due to warm-cloud-aerosol interactions is small, as could be expected. No single forcing or adjustment dominates the shortwave ERF_{aci}; the global-mean forcing F_{N_d} and \mathcal{L} adjustment $F_{\mathcal{L}}$ are of comparable magnitude at -0.52 W m^{-2} and -0.53 W m^{-2} . The cloud-fraction adjustment F_{f_c} at -0.31 W m^{-2} is the smallest of the components, consistent with other models (Zelinka et al., 2014; ?)CMIP5 models but not with other AeroCom models (Zelinka et al., 2014; Ghan et al., 2016; Gryspeerdt et al., submitted). Our decomposition
- 15 agrees with the results of Gryspeerdt et al. (submitted) to within 0.1 W m⁻² but disagrees with Ghan et al. (2016) (who estimate $F_L/F_{N_d} \approx 5$, in contrast to our estimate of ≈ 1), which both use a very similar configuration of the same version of ECHAM–HAM. Research to understand the differences between our decomposition and others is underway. All results in this section are based on 3-hourly mean output, $\overline{F}^{(3 \text{ h})}$. This is a common commonly used model output configuration, albeit at the expensive end of the spectrum from the standpoint of storage space requirements. We will justify this choice in Section 3.4.
- 20 The geographic patterns of all components exhibit similar features that result from a convolution of the distributions of susceptible clouds and CCN perturbations. There are large interhemispheric differences. In the northern hemisphere, fairly strong forcing prevails over both oceans and over most of the continents, with the exception of desert regions, northern Asia, and the Arctic; over the continents, a plume of high forcing components over China, extending eastwards into the Pacific Ocean, and of a magnitude far greater than over Europe and North America, is especially pronounced. In the southern hemisphere, on
- 25 the other hand, sizable forcing components are largely limited to the subtropical southern Pacific and southern Atlantic in the vicinity of the persistent stratocumulus decks; smaller local maxima in the forcing components also exist in the outflow regions of the midlatitude westerlies downwind of South America, Africa, and Australia.

While observational studies often find that the strongest forcing components occur in highly susceptible warm oceanic clouds where continental pollution intrudes on relatively clean conditions (but see Gryspeerdt et al., 2017, as a counterexample), the

30 These geographic patterns result from a convolution of the distributions of susceptible clouds and N_d perturbations. Figure 2 shows the ERF_{aci} sensitivity, defined as the forcing or adjustment strength per *e*-folding of the N_d burden (with the bar denoting temporal averaging over the length of the run),

$$S_{\xi} = \frac{F_{\xi}}{\Delta \ln \overline{N}_d}$$

(11)

where the N_d burden is defined as

$$\mathcal{N}_d = \int N_d \, dz;$$

Equation (11) is similar to the factorization of Bellouin et al. (submitted). The sensitivity for all ERF_{aci} components is greater over ocean than over land. Over land, the most susceptible clouds can be found over tropical Africa, southeastern Asia, and

(12)

- 5 Central America; the anthropogenic N_d perturbation over South America is too small to determine reliable sensitivities. Over ocean, the largest sensitivities occur in the midlatitudes, near the equator, and over the eastern ocean boundaries; again, N_d perturbations in the southern hemisphere are too small to estimate sensitivities reliably. The regions of maximum sensitivity are the regions where the base-state \mathcal{L} is high (not shown), which is intuitive in light of the strong influence of \mathcal{L} on cloud albedo.
- Figure 3 shows the anthropogenic perturbation in N_d . The perturbation is generally stronger over the continents than over ocean; over ocean, it is generally strongest in the subtropics. Thus, the highest perturbations tend to coincide with the least susceptible clouds, which explains why the ERF_{aci} components exhibit far less variability over a wide latitudinal range over the northern hemisphere oceans than either the sensitivity or the N_d perturbation, and why the remote northern hemisphere oceans are comparable to the polluted continental regions in ERF_{aci} strength. The exceptions to this pattern are the near-shore
- 15 eastern ocean boundaries, where a high sensitivity and reasonably strong N_d perturbation coincide, and eastern China, where a reasonably high sensitivity and very strong N_d perturbation coincide. The location where ECHAM–HAMMOZ simulates both the strongest N_d forcing and the strongest \mathcal{L} adjustment is over land in China, extending downwind into the northwestern Pacific Ocean.

In observational studies or observationally constrained modeling studies, it is common to define susceptibilities analogously

- 20 to Eq. (11) based on PD variability in cloud and aerosol variables and then multiply those susceptibilities by wholly or partially (Bellouin et al., 2013; Kinne, 2019) model-derived estimates of anthropogenic aerosol perturbations. (In the terminology we adopt here, "sensitivity" is a change in cloud property or cloud radiative effect in response to a climatological change in an aerosol variable, whereas "susceptibility" is a change in response to an instantaneous change in an aerosol variable.) There is disagreement among these studies on whether (Quaas et al., 2008; Alterskjær et al., 2012; Engström et al., 2015; Gryspeerdt et al., 2016; A
- 25 not (Lebsock et al., 2008; Chen et al., 2014; Gryspeerdt et al., 2017) the most susceptible oceanic clouds (or strongest ERF_{aci} components over ocean, in studies that do not report susceptibilities) occur in warm oceanic clouds where continental pollution intrudes on relatively clean conditions over the eastern ocean boundaries, as in our results. Of those studies that are not restricted to oceanic clouds, some agree with our finding of strong forcing due to relatively susceptible clouds over China (Gryspeerdt et al., 2017; McCoy et al., 2017) and some do not (Quaas et al., 2008; Alterskjær et al., 2012; Gryspeerdt et al., 2016).
- 30

3.2 Are the adjustments proportional to the forcing?

An intriguing aspect of the ACI problem is whether the adjustments may be described approximately as a proportional response to the forcing (?, where the question is also addressed from a multimodel perspective)(Gryspeerdt et al., 2019). On the

one hand, we do not necessarily expect proportionality in the physical atmosphere, since the processes responsible for the adjustments carry memory of the cloud evolution over various time scales; the parameterized cloud processes in GCMs share this feature, at least in principle, since the anthropogenic N_d perturbation seen by the precipitation parameterization at one time step could be the result of a CCN perturbation at some point in the past, carried to another point in space by advection,

- 5 and influenced by any of the other parameterized cloud processes. On the other hand, complex systems oftentimes exhibit simple emergent behaviors (e.g., Mülmenstädt and Feingold, 2018, and references therein). If the adjustments were to follow proportionally from the forcing, one consequence for the ACI problem would be that the total ERF_{aci} uncertainty should not be estimated as the uncertainty on the sum of uncorrelated RF_{aci} and adjustments but rather take the correlation between the forcing and adjustments into account, which would result in a smaller ERF_{aci} uncertainty estimate.
- In this study, we can test for proportionality in terms of the geographic distribution using the spatial variability in the temporal-mean ERF_{aci} components. Figure 4 shows that the zonal mean of the ratio between \mathcal{L} adjustment and N_d forcing is relatively stable around unity between the southern and northern midlatitudes with fairly small interhemispheric differences except in the Southern Ocean. The picture is somewhat different for the ratio between f_c adjustment and N_d forcing, which is more latitudinally variable and more different between the northern and southern hemisphere. Figure 5 reinforces these
- 15 conclusions, showing that $F_{\mathcal{L}}$ and F_{N_d} are fairly tightly correlated with a regression relationship remarkably close to one-toone, while the relationship between F_{f_c} and F_{N_d} is much looser.

One interpretation of these results is that the ERF_{aci} components share a geographic pattern due to the fact that large effects result from the coincidence of large anthropogenic aerosol sources and susceptible clouds; the shared geographic pattern then leads to an approximately proportional relationship that breaks down farther from the source regions or where a different

- 20 mixture of cloud processes dominates the cloud response (e.g., the Southern Ocean). The cloud cover scheme, which diagnoses f_c from the grid mean relative humidity, to some extent decouples f_c from the other cloud properties, which attenuates the influence of N_d on f_c . Nevertheless, the vagueness of this argument is unsatisfactorily mismatched against the precision of the $F_{\mathcal{L}}$ - F_{N_d} relationship, which suggests a deeper mechanism at play, e.g., that precipitation acts as a common sink process for both N_d and \mathcal{L} .
- 25 Further evidence for proportionality comes from (Gryspeerdt et al., submitted), who find an intermodel proportional relationship between global-mean forcing and rapid adjustments.

3.3 How should we treat columns containing ice?

30

In attempting to diagnose warm-cloud ACI forcing components, the question arises how ice-containing clouds should be handled. We can conduct the following set of experiments to determine the range of forcing strengths associated with different thermodynamic-phase treatments:

- 1. Perturb cloud properties in all cloudy model levels.
- 2. Perturb cloud properties in any liquid-containing cloudy model levels.
- 3. Perturb cloud properties in liquid-only cloudy model levels (default).

- 4. Perturb cloud properties in liquid-only cloudy columns.
- 5. Perturb cloud properties in liquid-only cloudy columns, correcting the result by their temporal occurrence fraction.

Table 2 summarizes the results. (For reasons of efficiency, we performed these sensitivity experiments on daily-mean output, SW-shortwave flux only. We did not perform experiment 2 because we expect the result to lie between experiments 1 and 3, whose separation is already in the noise.) We conclude that how we choose to treat mixed-phase and ice clouds makes little difference in ECHAM–HAM, so long as we do not restrict ourselves to columns containing only warm clouds. In the latter case, correcting the forcing by the temporal occurrence fraction of liquid-only columns in each model latitude–longitude box approximately recovers the results in which when ice-containing columns are retained; however, there is some indication of diverging trends in F_{N_d} (which decreases in magnitude as the ice filtering becomes more restrictive) and $F_{\mathcal{L}}$ (which increases in magnitude as the ice filtering becomes more restrictive). The ice-free column requirement is often made in passive remote

sensing studies to prevent contamination from ice clouds overlying warm clouds and uncertainties in multilayer cloud retrievals.

3.4 Does temporal averaging bias the results?

As Table 3 shows, longer averaging periods underestimate the forcing, but the differences between instantaneous output (the model time step is 7.5 minutes, but we sample every 3 h to reduce the data volume) and 3 h averages is minimal. Multimodel

15 ensembles which archive 3 h average output or 3 h subsampled instantaneous of column cloud properties, e.g., AeroCom and CFMIP2, are therefore amenable to treatment by the PRP method.

4 Conclusions

We have presented the first decomposition of the ACI effective forcing in ECHAM–HAM into a Twomey forcing component and rapid adjustments of \mathcal{L} and f_c . In ECHAM–HAM, no single component dominates: $F_{N_d} = -0.52 \text{ W m}^{-2}$, $F_{\mathcal{L}} = -0.53 \text{ W m}^{-2}$, and $F_{f_c} = -0.31 \text{ W m}^{-2}$; the Twomey forcing and \mathcal{L} adjustment are approximately equally strong, and the f_c adjustment is somewhat weaker, as in many other models. The global ERF is dominated by the northern-hemisphere forcing. Within the northern hemisphere, the strongest forcing components occur over land in China in F_{N_d} and $F_{\mathcal{L}}$. As expected, the stratocumulus sheets over the eastern ocean basins also show strong responses in both hemispheres, as do the midlatitude North Atlantic and North Pacific.

- The temporal-mean spatial patterns of F_{N_d} and $F_{\mathcal{L}}$ are highly correlated, suggesting an effective proportionality in the \mathcal{L} adjustment to the Twomey forcing even though the precipitation-suppression mechanism by which the \mathcal{L} adjustment is parameterized in the model has inherent memory that could decouple it from the Twomey effect. The spatial patterns of the temporal-mean F_{N_d} and $F_{\mathcal{L}}$, while sharing some of the same gross features, have a much less tight relationship than F_{N_d} and $F_{\mathcal{L}}$.
- 30 In our study of ECHAM–HAMMOZ, the forcing components are fairly insensitive to how we treat columns containing both ice and liquid cloud condensate. Requiring that columns be free of ice and then correcting for the temporal fractional

occurrence of ice cloud, a technique that is often necessary in observational studies, largely reproduces the results we obtain when we do not filter out such columns, albeit possibly causing an overestimate of the \mathcal{L} adjustment and an underestimate of the N_d forcing. (In interpreting the bearing of these results on analyses of satellite cloud retrievals, note that these studies do not necessarily apply the ice-free requirement at the coarse GCM scales of the present work, depending on whether they use gridded "level 3" data or the "level 2" native resolution of the ratioval algorithms.)

5 gridded "level 3" data or the "level 2" native resolution of the retrieval algorithms.)

Through idealized sensitivity studies presented in the Appendix, we have showed that PRP is a viable method for accurately decomposing ERF_{aci} into a N_d forcing and \mathcal{L} and f_c adjustments. This is the case despite large artifacts that occur due to the decorrelated cloud property fields; the forward–backward technique advocated by Colman and McAvaney (1997) is capable of removing these artifacts.

- PRP is the most direct method of diagnosing directly uses the intrinsic model knowledge of the time-varying, three-dimensional aerosol perturbation and resulting perturbation of cloud properties to diagnose the ERF_{aci} components and their spatial patterns, making it a valuable tool. This makes it useful a useful tool for providing context to other decomposition methods (e.g., Zelinka et al., 2014; ?) or less resource-intensive decomposition methods (e.g., Ghan et al., 2016; Gryspeerdt et al., submitted) or to intercomparison studies (e.g., Pincus et al., 2016; Smith et al., 2018) despite its demand for high-frequency vertically re-
- 15 solved model output.

20

Appendix A: Validation of the PRP method for ACI decomposition

A1 What is the effect of decorrelating the cloud properties?

Consider the results of forward and backward PRP plotted separately for the PD–PI experiment in Figure A1. Not only are the magnitudes grotesque, but taken at face value, they would imply a positive forcing in one direction and a negative forcing in the other. Furthermore, the spatial patterns bear no resemblance to that expected for ERF_{aci} . In this section, we investigate the consequences of these features for the ERF_{aci} decomposition.

Any given atmospheric property is often correlated with many others. Substituting cloud properties one at a time breaks these correlations. For example, since ECHAM–HAM parameterizes precipitation suppression by aerosol, we expect a positive correlation between N_d and \mathcal{L} within a model run. If we substitute \mathcal{L} from another run, the mechanistic link between N_d and

25 \mathcal{L} through precipitation suppression, by which higher N_d at a given point in time leads to higher \mathcal{L} at later times, is broken, and, therefore, the correlation between N_d and \mathcal{L} is altered.

We estimate the strength of such decorrelation effects by performing two model runs with (almost exactly) the same model physics, both nudged to the same large-scale dynamics and with the same fixed SST; the only difference between the runs is that a parameter in the Khairoutdinov and Kogan (2000) formulation for the autoconversion rate, tuned for ECHAM–HAM,

30
$$Q_{\text{aut}} = \gamma q_l^{\alpha} \left(\frac{N_d}{1 \text{ cm}^{-3}} \right)^{-\beta}, \tag{A1}$$

has been slightly perturbed from $\beta = 1.79$ to $\beta' = 1.79 + 10^{-5}$ ($\alpha = 2.47$ and $\gamma = 4 \times 1350$ s⁻¹ are unchanged). Even over short integration times (a year), these model runs will converge on the same climate, with nearly identical forcing components. (The

small perturbation in β does not result in a significant change in model sensitivity.) However, at any given elapsed integration time and geographic location, the cloud properties in the two runs are essentially uncorrelated. We refer to this pair of runs as the same-climate-different-weather experiment. Knowing that the true climatological TOA flux difference between this pair of runs is zero, we can use these runs to estimate decorrelation effects between any other decorrelated pair of runs, including the

PD and PI emissions runs. 5

> We find that decorrelation effects cause the PRP method to estimate enormous TOA flux perturbations when we perform forward or backward substitution of any single cloud property; this is shown in panels (a) and (b) of Figure A2.

> Unlike forward PRP or backward PRP individually, forward-backward PRP is unaffected by decorrelation, both in the global mean and locally in the temporal mean: panel (c) of Figure A2 shows that the fluctuations in ΔQ rapidly (i.e., within a year)

average to zero. This confirms that the Colman and McAvaney (1997) prescription is successful at minimizing the spurious 10 effects of decorrelation.

A2 Does PRP give the right answer?

The preceding section provides evidence that strong decorrelation effects do not lead to a spurious offset in forward-backward PRP results. Next, we show that decorrelation effects also do not lead to spurious scale factors. To do so, we scale N_d and \mathcal{L} by a globally constant factor of 1.1 at all timesteps and scale f_c by 0.99. We use PRP to diagnose the forcing associated with 15 each of these perturbations; the results are shown in the first three rows of Table A1. We can then estimate the strength of the decorrelation effects by performing PRP on the $\beta' = 1.79 + 10^{-5}$ run and the scaled-{ N_d , \mathcal{L} , f_c } $\beta = 1.79$ run. This is shown in the middle three rows of Table A1; the correct results are recovered to good approximation, with generally small attribution to incorrect ERF_{aci} components (the largest is -0.05 W m⁻² incorrectly diagnosed as f_c adjustment in the $\mathcal{L} \times 1.1$ experiment) and

generally small differences between the actual and diagnosed (the largest is a diagnosed $F_{\mathcal{L}} = -0.48 \text{ W m}^{-2}$ when the correct 20 value is -0.53 W m⁻²). The final test is an experiment in which all cloud properties are perturbed simultaneously and the clouds are decorrelated by using a $\beta' = 1.79 + 10^{-5}$ baseline run. The results are shown on the last line of Table A1; the correct ERFaci components are recovered in the presence of the confounding effects of decorrelation and of perturbing multiple cloud properties simultaneously with 0.1 W m⁻² or better accuracy, the largest discrepancy being the diagnosed $F_{f_c} = 0.14$ W m⁻² when the correct value is 0.24 W m^{-2}). 25

Thus, we find that forward-backward PRP can diagnose the forcing components correctly in the presence of decorrelations, in addition to diagnosing the absence of forcing correctly in the same-climate-different-weather case.

A3 **Does temporal averaging bias the results?**

See Table 3. Longer averaging periods underestimate the forcing, but the differences between instantaneous output (the model 30 time step is 7.5, but we sample every 3 to reduce the data volume) and 3 averages is minimal. As the latter is a standard model configuration, while subsampled instantaneous output requires modifications to the model code, we use the latter. Note that 3 average output of column cloud properties is also available in some multimodel ensembles, e.g., CFMIP2.

A3 Are the results sensitive to choosing grid-mean or in-cloud perturbations?

5

10

Perturbing in-cloud or grid-mean N_d and \mathcal{L} would be equivalent in the limit in which TOA flux perturbations are linear in the cloud properties. While individual model columns do not satisfy this linearity requirement, the temporal mean apparently exhibits sufficient effective linearity that the choice of in-cloud or grid-mean perturbations has little effect on the ERF component estimate; compare Tables 1 and A2.

Code and data availability. The PRP code is available in a GitHub repository that will be made public and receive a DOI concurrently with publication of the final paper. Similarly for the analysis code. ECHAM–HAMMOZ is available from hammoz.ethz.ch subject to acknowl-edgement of a licensing agreement. The PRP output on which the manuscript is based will be made available and receive a DOI on Zenodo concurrently with publication of the final paper. Due to the large data volume of 3-hourly vertically resolved fields, the model output itself was not archived, but model configuration files that can be used to replicate the output are available as part of the PRP code.

Author contributions. j_{μ} designed the study, performed model runs, computed the PRP diagnostics, and drafted the manuscript. EG and j_{μ} validated and debugged the results. EG, MS, PM, JQ, and j_{μ} proposed various sensitivity studies. EG, MS, PM, SD, JQ, and j_{μ} worked on the interpretation of the results. All authors contributed text to or comments on the manuscript.

Competing interests. The authors declare that they have no competing interests.

- 15 Acknowledgements. We thank Christina Sackmann, Steve Ghan, and an anonymous reviewer for their insightful comments that have improved the manuscript. The ECHAM-HAMMOZ model is developed by a consortium composed of ETH Zurich, Max-Planck-Institut für Meteorologie, Forschungszentrum Jülich, University of Oxford, and the Finnish Meteorological Institute and managed by the Center for Climate Systems Modeling (C2SM) at ETH Zurich. This work was funded by the FLASH project (project number QU 311/14-1) in the HALO Priority Program (SPP 1294) of the German Research Foundation (Deutsche Forschungsgemeinschaft, DFG); by the European Union through European Research Council (ERC) Starting Grant QUAERERE (grant agreement 306284); by the German Federal Ministry of Education and Research within the Research for Sustainable Development (FONA) framework program under the HD(CP)² project (project number 01LK1504C). EG was supported by an Imperial College London Junior Research Fellowship. PM's contributions were supported
 - by a travel grant from the Leibniz Invitations program at Universität Leipzig. Computing resources were provided by the German Climate Computing Center (Deutsches Klimarechenzentrum, DKRZ).

References

Ackerman, A., Kirkpatrick, M., Stevens, D., and Toon, O.: The impact of humidity above stratiform clouds on indirect aerosol climate forcing, NATURE, 432, 1014–1017, https://doi.org/10.1038/nature03174, 2004.

Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, SCIENCE, 245, 1227–1230, 1989.

- 5 Alterskjær, K., Kristjánsson, J. E., and Seland, Ø.: Sensitivity to deliberate sea salt seeding of marine clouds observations and model simulations, Atmospheric Chemistry and Physics, 12, 2795–2807, https://doi.org/10.5194/acp-12-2795-2012, 2012.
 - Andersen, H., Cermak, J., Fuchs, J., Knutti, R., and Lohmann, U.: Understanding the drivers of marine liquid-water cloud occurrence and properties with global observations using neural networks, Atmos. Chem. Phys., 17, 9535–9546, https://doi.org/10.5194/acp-17-9535-2017, 2017.
- 10 Bellouin, N., Quaas, J., Morcrette, J.-J., and Boucher, O.: Estimates of aerosol radiative forcing from the MACC re-analysis, Atmos. Chem. Phys., 13, 2045–2062, https://doi.org/10.5194/acp-13-2045-2013, 2013.
 - Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K., Christensen, M., Daniau, A.-L.,
 Dufresne, J.-L., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J. M., Malavelle, F., Lohmann, U., Mauritsen, T., McCoy,
 D., Myhre, G., Mülmenstädt, J., Neubauer, D., Possner, A., Rugenstein, M., Sato, Y., Schulz, M., Schwartz, S. E., Sourdeval, O., Storelvmo,
- 15 T., Toll, V., Winker, D., and Stevens, B.: Bounding aerosol radiative forcing of climate change, Rev. Geophys., submitted. Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S., Sherwood, S., Stevens, B., and Zhang, X.: Clouds and Aerosols, book section Chapter 7, pp. 571–658, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, https://doi.org/10.1017/CBO9781107415324.016, 2014.
- Chen, Y.-C., Christensen, M. W., Stephens, G. L., and Seinfeld, J. H.: Satellite-based estimate of global aerosol-cloud radiative forcing by
 marine warm clouds, NATURE GEOSCIENCE, 7, 643–646, https://doi.org/10.1038/NGEO2214, 2014.
- Christensen, M. W., Neubauer, D., Poulsen, C., Thomas, G., McGarragh, G., Povey, A. C., Proud, S., and Grainger, R. G.: Unveiling aerosolcloud interactions Part 1: Cloud contamination in satellite products enhances the aerosol indirect forcing estimate, Atmos. Chem. Phys. Discuss., https://doi.org/10.5194/acp-2017-450, 2017.

Colman, R.: A comparison of climate feedbacks in general circulation models, CLIMATE DYNAMICS, 20, 865–873, https://doi.org/10.1007/s00382-003-0310-z, 2003.

- Colman, R. A. and McAvaney, B. J.: A study of general circulation model climate feedbacks determined from perturbed sea surface temperature experiments, JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES, 102, 19383–19402, https://doi.org/10.1029/97JD00206, 1997.
 - Croft, B., Lohmann, U., Martin, R. V., Stier, P., Wurzler, S., Feichter, J., Posselt, R., and Ferrachat, S.: Aerosol size-dependent
- 30 below-cloud scavenging by rain and snow in the ECHAM5-HAM, ATMOSPHERIC CHEMISTRY AND PHYSICS, 9, 4653–4675, https://doi.org/10.5194/acp-9-4653-2009, 2009.
 - Croft, B., Lohmann, U., Martin, R. V., Stier, P., Wurzler, S., Feichter, J., Hoose, C., Heikkila, U., van Donkelaar, A., and Ferrachat, S.: Influences of in-cloud aerosol scavenging parameterizations on aerosol concentrations and wet deposition in ECHAM5-HAM, ATMO-SPHERIC CHEMISTRY AND PHYSICS, 10, 1511–1543, https://doi.org/10.5194/acp-10-1511-2010, 2010.
- 35 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Holm, E. V., Isaksen, L., Kallberg, P., Koehler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M.,

Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thepaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, QUARTERLY JOURNAL OF THE ROYAL METEOROLOGICAL SOCIETY, 137, 553–597, https://doi.org/10.1002/qj.828, 2011.

- 5 Engström, A., Bender, F. A.-M., Charlson, R. J., and Wood, R.: Geographically coherent patterns of albedo enhancement and suppression associated with aerosol sources and sinks, Tellus B, 67, 26442, https://doi.org/10.3402/tellusb.v67.26442, 2015.
 - Forster, P. M., Richardson, T., Maycock, A. C., Smith, C. J., Samset, B. H., Myhre, G., Andrews, T., Pincus, R., and Schulz, M.: Recommendations for diagnosing effective radiative forcing from climate models for CMIP6, JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES, 121, 12460–12475, https://doi.org/10.1002/2016JD025320, 2016.
- 10 Ghan, S., Wang, M., Zhang, S., Ferrachat, S., Gettelman, A., Griesfeller, J., Kipling, Z., Lohmann, U., Morrison, H., Neubauer, D., Partridge, D. G., Stier, P., Takemura, T., Wang, H., and Zhang, K.: Challenges in constraining anthropogenic aerosol effects on cloud radiative forcing using present-day spatiotemporal variability, PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, 113, 5804–5811, https://doi.org/10.1073/pnas.1514036113, 2016.

Ghan, S. J.: Technical Note: Estimating aerosol effects on cloud radiative forcing, ATMOSPHERIC CHEMISTRY AND PHYSICS, 13,

15 9971–9974, https://doi.org/10.5194/acp-13-9971-2013, 2013.

20

- Gryspeerdt, E., Quaas, J., and Bellouin, N.: Constraining the aerosol influence on cloud fraction, JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES, 121, 3566–3583, https://doi.org/10.1002/2015JD023744, 2016.
- Gryspeerdt, E., Quaas, J., Ferrachat, S., Gettelman, A., Ghan, S., Lohmann, U., Morrison, H., Neubauer, D., Partridge, D. G., Stier, P., Takemura, T., Wang, H., Wang, M., and Zhang, K.: Constraining the instantaneous aerosol influence on cloud albedo, Proc. Nat. Acad. Sci. USA, 119, 4899–4904, https://doi.org/10.1073/pnas.1617765114, 2017.
- Gryspeerdt, E., Goren, T., Sourdeval, O., Quaas, J., Mülmenstädt, J., S., D., Unglaub, C., Gettelman, A., and Christensen, M.: Constraining the aerosol influence on cloud liquid water path, Atmos. Chem. Phys., 19, 5331–5347, https://doi.org/10.5194/acp-19-5331-2019, 2019a.
 - Gryspeerdt, E., Mülmenstädt, J., Gettelman, A., Malavelle, F. F., Morrison, H., Neubauer, D., Partridge, D. G., Stier, P., Takemura, T., Wang, H., Wang, M., and Zhang, K.: Surprising similarities in model and observational aerosol radiative forcing estimates, Atmos. Chem. Phys.
- 25 Discuss., pp. 1–18, https://doi.org/10.5194/acp-2019-533, 2019b.
 - Guo, H., Golaz, J.-C., and Donner, L. J.: Aerosol effects on stratocumulus water paths in a PDF-based parameterization, GEOPHYSICAL RESEARCH LETTERS, 38, L17 808, https://doi.org/10.1029/2011GL048611, 2011.
 - Heyn, I., Block, K., Mülmenstädt, J., Gryspeerdt, E., Kühne, P., Salzmann, M., and Quaas, J.: Assessment of simulated aerosol effective radiative forcings in the terrestrial spectrum, Geophys. Res. Lett., 44, 1001–1007, https://doi.org/10.1002/2016GL071975, 2017.
- 30 Khairoutdinov, M. and Kogan, Y.: A new cloud physics parameterization in a large-eddy simulation model of marine stratocumulus, MONTHLY WEATHER REVIEW, 128, 229–243, https://doi.org/10.1175/1520-0493(2000)128<0229:ANCPPI>2.0.CO;2, 2000.
 - Kinne, S.: Aerosol radiative effects with MACv2, Atmos. Chem. Phys. Discuss., pp. 1–50, https://doi.org/10.5194/acp-2018-949, 2019.
 - Kinnison, D. E., Brasseur, G. P., Walters, S., Garcia, R. R., Marsh, D. R., Sassi, F., Harvey, V. L., Randall, C. E., Emmons, L., Lamarque, J. F., Hess, P., Orlando, J. J., Tie, X. X., Randel, W., Pan, L. L., Gettelman, A., Granier, C., Diehl, T., Niemeier, U., and Simmons, A. J.: Sen-
- 35 sitivity of chemical tracers to meteorological parameters in the MOZART-3 chemical transport model, JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES, 112, D20 302, https://doi.org/10.1029/2006JD007879, 2007.
 - Klocke, D., Quaas, J., and Stevens, B.: Assessment of different metrics for physical climate feedbacks, Clim. Dyn., 41, 1173–1185, https://doi.org/10.1007/s00382-013-1757-1, 2013.

Kooperman, G. J., Pritchard, M. S., Ghan, S. J., Wang, M., Somerville, R. C. J., and Russell, L. M.: Constraining the influence of natural variability to improve estimates of global aerosol indirect effects in a nudged version of the Community Atmosphere Model 5, JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES, 117, D23 204, https://doi.org/10.1029/2012JD018588, 2012.

Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G.,

5 Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850-2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, ATMOSPHERIC CHEMISTRY AND PHYSICS, 10, 7017–7039, https://doi.org/10.5194/acp-10-7017-2010, 2010.

Lin, H. and Leaitch, R.: Development of an in-cloud aerosol activation parameterization for climate modeling, in: Proc. WMO Workshop on Measurements of Cloud Properties for Forecasts of Weather and Climate, edited by Baumgardner, D. and Raga, G., pp. 328–335, World Meteorological Organization, 1997.

Lohmann, U. and Hoose, C.: Sensitivity studies of different aerosol indirect effects in mixed-phase clouds, ATMOSPHERIC CHEMISTRY

15 AND PHYSICS, 9, 8917–8934, https://doi.org/10.5194/acp-9-8917-2009, 2009.

20

- Lohmann, U. and Roeckner, E.: Design and performance of a new cloud microphysics scheme developed for the ECHAM general circulation model, CLIMATE DYNAMICS, 12, 557–572, https://doi.org/10.1007/s003820050128, 1996.
 - Lohmann, U., Stier, P., Hoose, C., Ferrachat, S., Kloster, S., Roeckner, E., and Zhang, J.: Cloud microphysics and aerosol indirect effects in the global climate model ECHAM5-HAM, ATMOSPHERIC CHEMISTRY AND PHYSICS, 7, 3425–3446, https://doi.org/10.5194/acp-7-3425-2007, 2007.
- Lohmann, U., Rotstayn, L., Storelvmo, T., Jones, A., Menon, S., Quaas, J., Ekman, A. M. L., Koch, D., and Ruedy, R.: Total aerosol effect: radiative forcing or radiative flux perturbation?, ATMOSPHERIC CHEMISTRY AND PHYSICS, 10, 3235–3246, https://doi.org/10.5194/acp-10-3235-2010, 2010.

McCoy, D. T., Bender, F. A.-M., Mohrmann, J. K. C., Hartmann, D. L., Wood, R., and Grosvenor, D. P.: The global aerosol-cloud first

- 25 indirect effect estimated using MODIS, MERRA, and AeroCom, JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES, 122, 1779–1796, https://doi.org/10.1002/2016JD026141, 2017.
 - Michibata, T., Suzuki, K., Sato, Y., and Takemura, T.: The source of discrepancies in aerosol-cloud-precipitation interactions between GCM and A-Train retrievals, ATMOSPHERIC CHEMISTRY AND PHYSICS, 16, 15413–15424, https://doi.org/10.5194/acp-16-15413-2016, 2016.
- 30 Mülmenstädt, J. and Feingold, G.: The radiative forcing of aerosol-cloud interactions in liquid clouds: Wrestling and embracing uncertainty, Curr. Clim. Change Rep., 4, 23–40, https://doi.org/10.1007/s40641-018-0089-y, 2018.
 - Neubauer, D., Lohmann, U., Hoose, C., and Frontoso, M. G.: Impact of the representation of marine stratocumulus clouds on the anthropogenic aerosol effect, ATMOSPHERIC CHEMISTRY AND PHYSICS, 14, 11 997–12 022, https://doi.org/10.5194/acp-14-11997-2014, 2014.
- 35 Penner, J. E., Quaas, J., Storelvmo, T., Takemura, T., Boucher, O., Guo, H., Kirkevag, A., Kristjansson, J. E., and Seland, O.: Model intercomparison of indirect aerosol effects, ATMOSPHERIC CHEMISTRY AND PHYSICS, 6, 3391–3405, https://doi.org/10.5194/acp-6-3391-2006, 2006.

Lebsock, M. D., Stephens, G. L., and Kummerow, C.: Multisensor satellite observations of aerosol effects on warm clouds, JOURNAL OF
 GEOPHYSICAL RESEARCH-ATMOSPHERES, 113, D15 205, https://doi.org/10.1029/2008JD009876, 2008.

- Pincus, R. and Stevens, B.: Paths to accuracy for radiation parameterizations in atmospheric models, JOURNAL OF ADVANCES IN MOD-ELING EARTH SYSTEMS, 5, 225–233, https://doi.org/10.1002/jame.20027, 2013.
- Pincus, R., Forster, P. M., and Stevens, B.: The Radiative Forcing Model Intercomparison Project (RFMIP): experimental protocol for CMIP6, GEOSCIENTIFIC MODEL DEVELOPMENT, 9, 3447–3460, https://doi.org/10.5194/gmd-9-3447-2016, 2016.
- 5 Quaas, J., Boucher, O., Bellouin, N., and Kinne, S.: Satellite-based estimate of the direct and indirect aerosol climate forcing, J. Geophys. Res., 113, 05 204, https://doi.org/10.1029/2007JD008962, 2008.
 - Salzmann, M., Ming, Y., Golaz, J.-C., Ginoux, P. A., Morrison, H., Gettelman, A., Kraemer, M., and Donner, L. J.: Two-moment bulk stratiform cloud microphysics in the GFDL AM3 GCM: description, evaluation, and sensitivity tests, ATMOSPHERIC CHEMISTRY AND PHYSICS, 10, 8037–8064, https://doi.org/10.5194/acp-10-8037-2010, 2010.
- 10 Smith, C. J., Kramer, R. J., Myhre, G., Forster, P. M., Soden, B. J., Andrews, T., Boucher, O., Faluvegi, G., Fläschner, D., Hodnebrog, O., Kasoar, M., Kharin, V., Kirkevåg, A., Lamarque, J.-F., Mülmenstädt, J., Olivié, D., Richardson, T., Samset, B. H., Shindell, D., Stier, P., Takemura, T., Voulgarakis, A., and Watson-Parris, D.: Understanding rapid adjustments to diverse forcing agents, Geophys. Res. Lett., 45, 12 023–12 031, https://doi.org/10.1029/2018GL079826, 2018.

Stevens, B. and Feingold, G.: Untangling aerosol effects on clouds and precipitation in a buffered system, NATURE, 461, 607-613,

- 15 https://doi.org/10.1038/nature08281, 2009.
- Stevens, B., Giorgetta, M., Esch, M., Mauritsen, T., Crueger, T., Rast, S., Salzmann, M., Schmidt, H., Bader, J., Block, K., Brokopf, R., Fast, I., Kinne, S., Kornblueh, L., Lohmann, U., Pincus, R., Reichler, T., and Roeckner, E.: Atmospheric component of the MPI-M Earth System Model: ECHAM6, JOURNAL OF ADVANCES IN MODELING EARTH SYSTEMS, 5, 146–172, https://doi.org/10.1002/jame.20015, 2013.
- 20 Stier, P., Feichter, J., Kinne, S., Kloster, S., Vignati, E., Wilson, J., Ganzeveld, L., Tegen, I., Werner, M., Balkanski, Y., Schulz, M., Boucher, O., Minikin, A., and Petzold, A.: The aerosol-climate model ECHAM5-HAM, ATMOSPHERIC CHEMISTRY AND PHYSICS, 5, 1125– 1156, https://doi.org/10.5194/acp-5-1125-2005, 2005.
 - Stjern, C. W., Samset, B. H., Myhre, G., Forster, P. M., Hodnebrog, O., Andrews, T., Boucher, O., Faluvegi, G., Iversen, T., Kasoar, M., Kharin, V., Kirkevag, A., Lamarque, J.-F., Olivie, D., Richardson, T., Shawki, D., Shindell, D., Smith, C. J., Takemura, T., and Voul-
- 25 garakis, A.: Rapid Adjustments Cause Weak Surface Temperature Response to Increased Black Carbon Concentrations, JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES, 122, 11 462–11 481, https://doi.org/10.1002/2017JD027326, 2017.
 - Sundqvist, H., Berge, E., and Kristjansson, J.: Condensation and cloud parameterization studies with a mesoscale numerical weather prediction model, MONTHLY WEATHER REVIEW, 117, 1641–1657, https://doi.org/10.1175/1520-0493(1989)117<1641:CACPSW>2.0.CO;2, 1989.
- 30 Taylor, K. E., Crucifix, M., Braconnot, P., Hewitt, C. D., Doutriaux, C., Broccoli, A. J., Mitchell, J. F. B., and Webb, M. J.: Estimating Shortwave Radiative Forcing and Response in Climate Models, Journal of Climate, 20, 2530–2543, https://doi.org/10.1175/JCLI4143.1, 2007.

Twomey, S.: Influence of pollution on shortwave albedo of clouds, JOURNAL OF THE ATMOSPHERIC SCIENCES, 34, 1149–1152, https://doi.org/10.1175/1520-0469(1977)034<1149:TIOPOT>2.0.CO;2, 1977.

- 35 Wetherald, R. and Manabe, S.: Cloud feedback processes in a general-circulation model, JOURNAL OF THE ATMOSPHERIC SCIENCES, 45, 1397–1415, https://doi.org/10.1175/1520-0469(1988)045<1397:CFPIAG>2.0.CO;2, 1988.
 - Wood, S. N.: Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models, Journal of the Royal Statistical Society (B), 73, 3–36, 2011.

- Zelinka, M. D., Andrews, T., Forster, P. M., and Taylor, K. E.: Quantifying components of aerosol-cloud-radiation interactions in climate models, JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES, 119, 7599–7615, https://doi.org/10.1002/2014JD021710, 2014.
 Zhang, K., O'Donnell, D., Kazil, J., Stier, P., Kinne, S., Lohmann, U., Ferrachat, S., Croft, B., Quaas, J., Wan, H., Rast, S., and Feichter,
- 5 J.: The global aerosol-climate model ECHAM-HAM, version 2: sensitivity to improvements in process representations, ATMOSPHERIC CHEMISTRY AND PHYSICS, 12, 8911–8949, https://doi.org/10.5194/acp-12-8911-2012, 2012.
- 545 Zhang, K., Wan, H., Liu, X., Ghan, S. J., Kooperman, G. J., Ma, P.-L., Rasch, P. J., Neubauer, D., and Lohmann, U.: Technical Note: On the use of nudging for aerosol-climate model intercomparison studies, ATMOSPHERIC CHEMISTRY AND PHYSICS, 14, 8631–8645, https://doi.org/10.5194/acp-14-8631-2014, 2014.
 - Zhou, C. and Penner, J. E.: Why do general circulation models overestimate the aerosol cloud lifetime effect?: A case study comparing CAM5 and a CRM, ATMOSPHERIC CHEMISTRY AND PHYSICS, 17, 21–29, https://doi.org/10.5194/acp-17-21-2017, 2017.



Figure 1. ERFaci components in ECHAM-HAM estimated by forward-backward PRP



Figure 2. Sensitivity of ERF_{aci} components to anthropogenic N_d change (shown only where $\Delta \ln \overline{N}_d > 0.05$)



Figure 3. Anthropogenic $\ln N_d$ change



Figure 4. ERFaci adjustments relative to the Twomey forcingshould the F's in the figure have overbars to indicate they are temporal averages?



Figure 5. Correlation plots between the temporal-mean Twomey forcing and the adjustments; color indicates the number of grid boxes within each 0.05 W m⁻² × 0.05 W m⁻² bin; the red line is a linear least-squares regression; the blue line is a generalized additive model regression (Wood, 2011), with 95% confidence interval shaded in light blue; and the dashed gray line is the one-to-one line

Table 1. ERF_{aci} components in ECHAM–HAM estimated by forward–backward PRP. The total ERF also includes the ice-phase ACI effects (-0.59 W m⁻² in the SW, 0.88 W m⁻² in the LW), RF_{ari} (-0.17 W m⁻² in the SW), and a negligible surface-albedo contribution (-0.01 W m⁻²), estimated for a very similar model run in Gryspeerdt et al. (submitted). The sum of the components thus balances at approximately the 0.2 W m⁻² level, a relative error similar to the 0.1 W m⁻² estimated uncertainty on the ERF_{aci} components.

	ERF components (W m ⁻²)		Sum (W m ⁻²)	Total ERF (W m ⁻²)	
Spectrum	F_{N_d}	F_{f_c}	$F_{\mathcal{L}}$	$F_{N_d} + F_{f_c} + F_{\mathcal{L}}$	
LW	0.00	0.04	0.03	0.07	0.72
SW	-0.52	-0.35	-0.57	-1.44	-2.03



Figure A1. Forward (a) and backward (b) PRP estimates of the ERF_{aci} components. Note the significantly wider color scale than in Figure 1.



Figure A2. Forward (a), backward (b), and (c) forward–backward PRP performed on the same-climate–different-weather case. Note the significantly wider color scale than in Figure 1.

Table 2. Dependence of diagnosed ERF_{aci} components on treatment of thermodynamic phase

	ERF components (W m ⁻²)		
Phase treatment	δQ_{N_d}	δQ_{f_c}	$\delta Q_{\mathcal{L}}$
All phases	-0.29	-0.29	-0.34
Liquid-only cloudy model levels	-0.27	-0.27	-0.35
Liquid-only cloudy model columns	-0.15	-0.17	-0.21
Liquid-only cloudy model columns (corrected for occurrence fraction)	-0.26	-0.29	-0.38

	ERF components (W m ⁻²)		
Averaging period	FNd~	Flow	FL
1_month	-0.09	-0.09	-0.11
l_d	-0.35	-0.33	-0.30
<u>3</u> _h	-0.52	-0.31	-0.53
instantaneous	-0.55	-0.30	-0.51

Table 3. Dependence of diagnosed ERFaci components on temporal averaging

Table A1. ERF_{aci} components resulting from idealized perturbations to N_d , \mathcal{L} , and f_c ; estimate of the same ERF_{aci} components by forward–backward PRP in the presence of decorrelation effects.

	ERF components (W m ⁻²)		
Forcing Perturbation	F_{N_d}	F_{f_c}	$F_{\mathcal{L}}$
$N_d \times 1.1$	-0.38	-0.00	-0.00
$f_c imes 0.99$	-0.00	0.24	-0.00
$\mathcal{L} \times 1.1$	-0.00	-0.00	-0.53
$N_d \times 1.1$ with $\beta' = \beta + 10^{-5}$	-0.37	-0.01	-0.01
$f_c \times 0.99$ with $\beta' = \beta + 10^{-5}$	0.01	0.21	0.02
$\mathcal{L} \times 1.1$ with $\beta' = \beta + 10^{-5}$	0.01	-0.05	-0.48
$N_d \times 1.1, \mathcal{L} \times 1.1, f_c \times 0.99$ with $\beta' = \beta + 10^{-5}$	-0.31	0.14	-0.49

Dependence of diagnosed components on temporal averaging Averaging period F_{N_d} F_{f_c} $F_{\mathcal{L}}$ 1 - 0.09 - 0.09 - 0.11 1 - 0.35 - 0.33 - 0.30 3 - 0.52 - 0.31 - 0.53 instantaneous - 0.55 - 0.30 - 0.51-

Table A2. ERF_{aci} components calculated by PRP on f_c and in-cloud N_d and q_l

ERF components (W m ⁻²)			
F_{N_d}	F_{f_c}	$F_{\mathcal{L}}$	
-0.48	-0.30	-0.48	