Atmos. Chem. Phys. Discuss., https://doi.org/10.5194/acp-2018-1309-RC1, 2019 © Author(s) 2019. This work is distributed under the Creative Commons Attribution 4.0 License.





Interactive comment

Interactive comment on "Contributions of biomass-burning, urban, and biogenic emissions to the concentrations and light-absorbing properties of particulate matter in central Amazonia during the dry season" by Suzane S. de Sá et al.

Anonymous Referee #1

Received and published: 5 February 2019

1. Line 155. As reported by (Zotter et al 2017) at 370 nm there maybe a non-negligible light-absorption contribution from SOA compounds, I would use the wavelength at 430 nm. 2. Line 166-170. I would use some references to support the assumptions made. 3. Line 528-537. When you describe Figure 11 you don't comment on the Angström exponent 4. Figure 4 and Figure 11: I would write the values of the interquartile ranges: 25, 75 or 10, 90? 5. Figure 11: I would write the wavelengths also on the graph to be clear, in particular on the Angstrom exponent 6. Figure 13: I would draw the correlation

Printer-friendly version

Discussion paper



Interactive comment on Atmos. Chem. Phys. Discuss., https://doi.org/10.5194/acp-2018-1309, 2019.

ACPD

Interactive comment

Printer-friendly version

Discussion paper

