



The importance of crystalline phases in ice nucleation by volcanic ash

Elena C. Maters¹, Donald B. Dingwell², Corrado Cimarelli², Dirk Müller², Thomas F. Whale^{1,3}, Benjamin J. Murray¹

¹School of Earth and Environment, University of Leeds, Leeds LS2 9JT, United Kingdom

5 ²Department of Earth and Environmental Sciences, Ludwig-Maximilians University, 80333 Munich, Germany

³School of Chemistry, University of Leeds, Leeds LS2 9JT, United Kingdom

Correspondence to: Elena C. Maters (e.c.maters@leeds.ac.uk)

Abstract. Volcanic ash is known to nucleate ice when immersed in supercooled water droplets. This process may impact the properties and dynamics of the eruption plume and cloud, as well as those of meteorological clouds once the ash is dispersed in the atmosphere. However, knowledge of what controls the ice-nucleating effectiveness (INE) of ash remains limited, although it has been suggested that crystalline components in ash may play an important role. Here we adopted a novel approach using nine pairs of tephra and their remelted and quenched glass equivalents to investigate the influence of chemical composition, crystallinity and mineralogy on ash INE in the immersion mode. For all nine pairs studied, the crystal-bearing tephra nucleated ice at higher temperatures than the corresponding crystal-free glass, demonstrating that crystalline phases are key to ash INE. Similar to findings for desert dust from arid and semi-arid regions, the presence of feldspar minerals characterises the four most ice-active tephra samples, although a high INE is observed even in the absence of alkali feldspar in samples bearing plagioclase feldspar and orthopyroxene. There is evidence of a potential indirect relationship between chemical composition and ash INE, whereby a magma of felsic to intermediate composition may generate ash containing ice-active feldspar minerals. This complex interplay between chemical composition, crystallinity, and mineralogy could help partly to explain the variability in volcanic ash INE reported in the literature. Overall, by categorically demonstrating the importance of crystalline phases in the INE of volcanic ash, our study contributes insights essential for better appraising the role of airborne ash in ice formation. Among these is the inference that glass-dominated ash emitted by the largest explosive eruptions may be less effective at impacting ice-nucleating particle populations than crystalline ash generated by smaller, more frequent eruptions.

25 1 Introduction

Volcanic ash produced by explosive eruptions can act as ice-nucleating particles (INPs), promoting heterogeneous freezing of supercooled water in the vertical eruption plume, the laterally dispersed eruption cloud, and the wider atmosphere (Isono et al., 1959a; 1959b; Hobbs et al., 1971; Rose et al., 2003). Ice formation in these contexts is poorly understood yet may exert a profound influence on eruption plume/cloud dynamics and electrification (e.g., Herzog et al., 1998; Cimarelli et al., 2016), sequestration of gaseous species (e.g., Textor et al., 2003; Guo et al., 2004a) and ash aggregation and sedimentation (e.g., Guo et al., 2004b; Van Eaton et al., 2015), as well as atmospheric cloud properties and lifetime (e.g., Komabayasi, 1957; Seifert et al., 2011) and thereby the hydrological cycle and climate (e.g., Isono and Komabayasi, 1954). Ongoing volcanic activity generates a recurrent flux of ash particles into the atmosphere (176–256 Tg a⁻¹; Durant et al., 2010), whereas sporadic large eruptions can result in ash loadings greatly exceeding annual averages over very short (hour to day) time scales and transiently dominating INP populations (e.g., Isono et al., 1959a; 1959b; Hobbs et al., 1971).

By definition, volcanic ash consists of pyroclastic particles <2 mm in diameter usually dominated by aluminosilicate glass derived from the melt and/or aluminosilicate and Fe(-Ti) oxide minerals in the form of crystals suspended within the original melt (Heiken and Wohletz, 1992). Field and laboratory measurements present conflicting evidence as to the ice-nucleating effectiveness (INE) of ash, even for samples from the same volcano, and it is far from clear what drives this variation (Durant



et al., 2008, and references therein; Mangan et al., 2017). Studies on desert dust from arid and semi-arid regions (1000-3000 Tg a⁻¹ emitted; Penner et al., 2001) - considered one of the most important INP types globally (Hoose et al., 2010; Vergara-Temprado et al., 2017) - suggest that chemical composition, crystallinity and mineralogy can influence the abundance of ice-active surface sites on the solid particles (Murray et al., 2012, and references therein). Specifically, the presence of K-rich feldspar is thought to dominate the INE of dust (Atkinson et al., 2013; Yakobi-Hancock et al., 2013; Kaufmann et al., 2016). Similar factors might influence ice nucleation by volcanic ash (Kulkarni et al., 2015; Schill et al., 2015; Genareau et al., 2018). To date however, the roles and potential interplay of differing physicochemical attributes in determining a solid particle's INE remain poorly understood, having rarely been systematically investigated for any ice-nucleating material.

Here we examine the influence of three properties dictated primarily by the state of the erupted source magma - chemical composition, crystallinity, and mineralogy - on the INE of volcanic ash in the immersion mode which is likely relevant to ash particles in the water-rich eruption plume/cloud and in mixed-phase atmospheric clouds (Textor et al., 2006; McNutt and Williams, 2010; Pruppacher and Klett, 2010; Murray et al., 2012). To assist in disentangling the individual effects of these properties on ice nucleation, we have adopted a novel approach of sample selection by using a wide range of natural tephra and their remelted and quenched glass equivalents. In this manner, we aim to contribute to improving understanding of the potential for airborne ash from different eruptions to impact ice formation above the volcanic vent and/or once dispersed in the ambient atmosphere.

2 Materials and methods

2.1 Volcanic tephra and glass samples

Eighteen tephra and (remelted and quenched) glass powders spanning a range of compositions associated with volcanic activity were studied (Table 1). The powders are represented by an abbreviated sample code with the subscripts 'teph' or 'glass' used to designate tephra or glass material, respectively. The tephra samples correspond to ash or pumice originating from different eruptions. The glass samples were synthesised by melting a portion of the tephra at 1400 to 1600 °C, homogenising the melts by stirring for 12 to 72 h, and quenching the melts at room temperature to form glasses. This technique for generating volcanic glass has been used previously in studies of volcanic ash reactivity (e.g., Ayris et al., 2013; 2014; Maters et al., 2016). All samples were crushed to fine powders in a ball mill using a zirconia ceramic ball and vial, to ensure consistent treatment of the tephra and glass materials prior to ice nucleation experiments. The specific surface area (SSA_{BET}; Table 1) of the samples after overnight degassing was obtained from a ten-point N₂ adsorption isotherm at -196 °C based on the Brunauer, Emmet and Teller model (Brunauer et al., 1938) using a Micromeritics TriStar 3000 instrument.

The chemical composition of volcanic ash predominantly reflects the state of the source magma at the point of eruption but can also be influenced by lithic material (pre-existing country rock) entrained during the explosive eruption (Heiken and Wohletz, 1992). The composition of the tephra and glass samples studied here (Table S1) was measured by Nora Groschopf (Institute of Geosciences, Johannes Gutenberg University Mainz, Germany) by X-ray fluorescence at 3.2 kW ionisation energy using a Philips Analytical MagiX PRO spectrometer. The conventional classification of these materials according to total alkali (Na₂O + K₂O) versus silica (SiO₂) content is shown on a Total Alkali versus Silica diagram in Fig. 1a (Le Maitre et al., 2002).

As magma ascends to the surface prior to a volcanic eruption, the aluminosilicate melt typically carries a cargo of mineral species in the form of crystals predominantly originating from the melt. Upon magma fragmentation, the ash generated comprises a corresponding mixture of glass and crystal components (Heiken and Wohletz, 1992). The crystallinity refers to the relative abundance of crystals in ash (i.e., crystal mass/total mass) and typically ranges from 0 to 65 wt.%, depending on factors such as the prior state of the magma (e.g., chemical composition, temperature) and even the dynamics of the conduit such as the ascent rate of the magma (Heiken and Wohletz, 1992; Wright et al., 2012). The incorporation of lithic material



from the conduit or vent and/or the interaction with ground or surface water can also influence the crystallinity of the bulk erupted ash.

The mineralogy refers to the identities and abundances of crystalline phases in ash. Among the factors that influence crystallisation from the melt, the chemical composition of the magma is a key determinant of the mineral phases that can form (Heiken and Wohletz, 1992). Common minerals in basaltic ash include pyroxene, olivine, amphibole, and (Ca-rich) plagioclase feldspar, whereas those in rhyolitic ash include quartz, mica, amphibole, (Na-rich) plagioclase feldspar, and (K-rich) alkali feldspar (Fig. 1b; Rogers, 2015). Note that in this study, we use the terms ‘plagioclase’ and ‘alkali’ to refer specifically to Na-/Ca-rich and K-rich feldspars, respectively.

The proportions of various crystalline phases in the tephra studied here (Table 2) were determined by X-ray diffraction (XRD) with a $\text{Cu}_{\text{K}\alpha 1}$ X-ray beam using a GE diffractometer (XRD 3003 TT) and the Profex software program for Rietveld refinement (Döbelin and Kleeberg, 2015). Briefly, this involved crushing the tephra and spiking it with a known mass (~17 wt.%) of pure Si powder, as crystalline internal standard, to quantify the crystallinity and mineralogy of the samples. While the LIP_{teph} and CID_{teph} crystallinities cannot be quantified below the ~2 wt.% detection limit of the technique, this does not rule out the possibility of smaller amounts of crystals and/or nanoscale crystallites being present in these samples. The glasses were also analysed by XRD to confirm their amorphous nature (i.e., the absence of crystalline minerals within the ~2 wt.% detection limit).

2.2 Immersion mode ice nucleation experiments

The INE of the tephra and glass samples was assessed using a microlitre Nucleation by Immersed Particles Instrument (μL -NIPI). This instrument has been described in detail elsewhere (Whale et al., 2015) and has been used previously to study heterogeneous ice nucleation by mineral and ash material (e.g., Atkinson et al., 2013; Harrison et al., 2016; Mangan et al., 2017; Whale et al., 2017). Briefly, a 1 wt.% sample suspension in Milli-Q water ($18.2 \text{ M}\Omega \cdot \text{cm}$) was shaken for a few minutes by a vortex mixer, and then pipetted in an array of 30 to 40 $1 \mu\text{L}$ droplets onto a hydrophobic silanised glass cover slip placed on a temperature-controlled stage (Grant-Asymptote EF600 Stirling Cryocooler). The stage was cooled from room temperature at a rate of $-5 \text{ }^\circ\text{C min}^{-1}$ down to $0 \text{ }^\circ\text{C}$, and subsequently at a rate of $-1 \text{ }^\circ\text{C min}^{-1}$ until all droplets were frozen. A dry nitrogen flow ($\sim 0.2 \text{ L min}^{-1}$) over the droplets prevented condensation and frost accumulation on the cover slip and hence served to avoid frozen droplets affecting neighbouring liquid droplets. A digital camera was used to observe the droplets throughout the experiment and determine the fraction of droplets frozen as a function of temperature $f_{\text{ice}}(T)$ according to:

$$f_{\text{ice}}(T) = n_{\text{ice}}(T)/n \quad (1)$$

where $n_{\text{ice}}(T)$ is the cumulative number of droplets frozen at temperature (T) and n is the total number of droplets in the experiment. At least three replicate experiments were conducted for each sample. In addition, ice nucleation of the background water at higher temperatures than those predicted by classical nucleation theory (Murray et al., 2010a; Koop and Murray, 2016), due to impurities in the water and/or effects of the cover slip (Polen et al., 2018), was assessed by acquiring baseline droplet freezing measurements of water containing no added particles.

To facilitate comparison of different materials including across literature studies, their ability to nucleate ice is often expressed in terms of the ice nucleation active site density $n_s(T)$, which represents the number of active sites per unit surface area of a solid sample on cooling from $0 \text{ }^\circ\text{C}$ down to temperature (T) (Connolly et al., 2009):

$$n_{\text{ice}}(T)/n = 1 - \exp(-n_s(T)A) \quad (2)$$

where A is the total surface area of the solid sample per droplet. Although the fundamental nature of ice-active sites remains unclear, and may vary across different materials, $n_s(T)$ allows us to empirically define the INE of a range of solid substrates (Vali, 2014). The uncertainty in $n_s(T)$ was calculated using simulations of possible active site distributions propagated with the



uncertainty in surface area of nucleant per droplet, as outlined in Harrison et al., (2016). The uncertainty in temperature for the $\mu\text{L-NIPI}$ is estimated to be ± 0.4 °C (Whale et al., 2015).

3 Results

The $f_{\text{ice}}(T)$ and $n_s(T)$ values for the eighteen samples studied are shown in Fig. 2. For the sake of clarity, droplet freezing events across replicate experiments have been combined here to generate a single dataset for each sample. The samples display wide variation in freezing temperatures with generally higher values associated with the tephra (-3 to -25 °C; Fig. 2a, c) than with the glass (-12 to -30 °C; Fig. 2b, d).

For the tephra samples, the $f_{\text{ice}}(T)$ curves are separate from those of the background water (Fig. 2a). This gives confidence to attribution of the observed freezing to ice nucleation by tephra particles. In terms of ice nucleation active site densities (Fig. 2c), taking the temperature at which $n_s \approx 1 \text{ cm}^{-2}$ as a simple single-number proxy for INE ($T_{n_s \approx 1 \text{ cm}^{-2}}$), the most active tephra are the trachyphonolite samples NUO_{teph} and AST_{teph} with $T_{n_s \approx 1 \text{ cm}^{-2}}$ values of -5.0 °C and -6.4 °C, respectively, followed by the andesite samples COL_{teph} and TUN_{teph} with $T_{n_s \approx 1 \text{ cm}^{-2}}$ values of -8.1 °C and -10.5 °C, respectively. The least active tephra are the basalt and phonolite samples KIL_{teph} and LAC_{teph} with $T_{n_s \approx 1 \text{ cm}^{-2}}$ values of -17.5 °C and -17.9 °C, respectively.

For the glass samples in contrast, there is significant overlap of the $f_{\text{ice}}(T)$ curves and those of the background water, spanning temperatures from -18 °C to -35 °C (Fig. 2b). This prevents attribution of the observed freezing to ice nucleation by glass particles and comparison of individual glass activities. It also likely explains the poorer reproducibility seen in replicate experiments of the glass material relative to the tephra material (Fig. 3). Hence, in most cases the reported $n_s(T)$ values should be regarded as upper limits. However, the trachyte and andesite samples $\text{CID}_{\text{glass}}$ and $\text{COL}_{\text{glass}}$ stand out with only partial overlap of $f_{\text{ice}}(T)$ curves and the freezing temperature range of the background water (Fig. 2b), and thus are identified as being the most active glasses in terms of ice nucleation active site densities (Fig. 2d), with $T_{n_s \approx 1 \text{ cm}^{-2}}$ values of -16.8 °C and -17.0 °C, respectively.

Consideration of the compositionally analogous tephra-glass pairs clearly illustrates the observation of tephra nucleating ice more effectively than the equivalent remelted and quenched glass (Fig. 3). The $n_s(T)$ curves of each tephra sample fall at higher temperatures compared to those of its counterpart glass sample, albeit displaying varying temperature differences between them.

4 Discussion

The eighteen samples were chosen to encompass a variety of chemical compositions, crystallinities and mineralogies encountered in volcanic ash, with the aim of investigating the influence of these physicochemical properties on ash INE. The results of our ice nucleation experiments are examined in relation to each of these properties below.

4.1 Crystallinity

As noted above, our sample pairs of crystal-bearing tephra and crystal-free glass of nearly identical chemical composition (Table S1) constitute a novel approach to studying controls on volcanic ash ice nucleation. These pairs were chosen to disentangle variation in crystallinity from that in composition, which together might complicate interpretation of INE trends in natural ash collections. For all nine pairs studied, the tephra nucleates ice at higher temperatures than the corresponding glass (Fig. 3), overall implying a positive effect of crystals on INE. However, the difference between tephra and glass $T_{n_s \approx 1}$



cm^{-2} values across the sample pairs ($\Delta T_{n_s \approx 1 \text{ cm}^{-2}}$, ranging from ~ 2 to 19 °C) does not vary simply with respect to tephra crystallinity (<2 to 66 wt.%; Fig. S1a). Additionally, a plot of $T_{n_s \approx 1 \text{ cm}^{-2}}$ values versus crystallinity of the tephra samples shows no correlation between ice nucleation and crystalline content (Fig. S1b). The compositionally quite similar NUO_{teph} and AST_{teph} display the highest INE ($T_{n_s \approx 1 \text{ cm}^{-2}}$ values of -5.0 and -6.4 °C, respectively) and are characterised by markedly contrasting crystallinities (60 and 28 wt.%, respectively). A difference in crystallinity was proposed as one of a number of potential explanations for the variable INE of two ash samples from Soufrière Hills volcano, Montserrat, with the 100% crystalline material produced by a dome collapse showing a higher INE than the 89% crystalline material produced by a magmatic eruption (Schill et al., 2015; Mangan et al., 2017). However, in light of our findings, it seems unlikely that this slight difference in crystallinity of two Soufrière Hills ash samples can adequately explain the large disparity in their INE.

10 A study comparing ice nucleation by crystalline and glassy anorthite ($\text{CaAl}_2\text{Si}_2\text{O}_8$), with the former displaying a $n_s(T)$ curve reaching higher freezing temperatures than the latter, suggested that crystals may introduce rarer, more ice-active surface sites but are not required for ice nucleation (Harrison et al., 2016). In addition, aqueous organic solutions in a glassy state have been shown to nucleate ice (Murray et al., 2010b; Wagner et al., 2012; Ignatius et al., 2016). In contrast, our observations strongly suggest that the presence of crystals is crucial in making volcanic ash an effective ice nucleant, although the abundance of crystalline phases in ash may be less important than the mere presence and specific properties of those phases in determining the INE.

4.2 Mineralogy

Consideration of the nine tephra samples might provide insight into the influence of mineralogy on the INE of volcanic ash. As noted above, comparison of the compositionally analogous tephra-glass pairs points to a role of crystalline phases in promoting freezing, and we infer that the properties of those phases have a strong effect on a sample's INE. A study of ash from Fuego, Soufrière Hills, and Taupo volcanoes attributed differences in their INE to their contrasting mineralogies (Schill et al., 2015). Overall, when the $T_{n_s \approx 1 \text{ cm}^{-2}}$ values are plotted against the content of various minerals in the tephra samples studied here, no clear correlations are evident (Fig. 4). However, certain features do stand out; the two most ice-active NUO_{teph} and AST_{teph} have the highest contents of alkali feldspar (60 and 19 wt.%, respectively; Fig. 4a), while the next most ice-active COL_{teph} and TUN_{teph} are characterised by an abundance of plagioclase feldspar (55 and 43 wt.%, respectively; Fig. 4b).

The $n_s(T)$ curves of our tephra samples are compared with the ice-nucleating activity of the two feldspar mineral groups in Fig. 5. The $n_s(T)$ curves of NUO_{teph} and AST_{teph} span temperatures consistent with the alkali feldspar parameterisation compiled from literature data (Harrison et al., in prep.), supporting the notion that the INE of these two samples relates to the presence of this mineral phase. The next two most ice-active materials COL_{teph} and TUN_{teph} contain no appreciable quantity of alkali feldspar, instead being rich in plagioclase feldspar. However, the $n_s(T)$ curves of these two samples are inconsistent with the relatively low INE of plagioclase feldspar reported in the literature (Fig. 5). This may point to the presence of very ice-active plagioclase feldspar or to the influence of some other component in COL_{teph} and TUN_{teph} (e.g., orthopyroxene, discussed below). Harrison et al. (2016) have found that some alkali and plagioclase feldspar samples are hyper-active relative to the majority of materials tested, hence our observations may simply reflect the natural variability in INE of feldspar minerals.

35 Other tephra samples containing feldspar are comparatively less effective at nucleating ice, in particular; the intermediately ice-active ETN_{teph} (44 wt.% plagioclase feldspar; Fig. 4b) and the least ice-active KIL_{teph} (3 wt.% plagioclase feldspar; Fig. 4b) and LAC_{teph} (9 wt.% alkali feldspar; Fig. 4a). Such differences may relate to the specific chemistry of the mineral phases present in the tephra. Harrison et al. (2016) showed that the INE of feldspar generally decreases from the K end-member (KAlSi_3O_8) to the Na end-member ($\text{NaAlSi}_3\text{O}_8$) to the Ca end-member ($\text{CaAl}_2\text{Si}_2\text{O}_8$), with the exception of a hyper-active NaAlSi₃O₈ specimen (Amelia albite). Based on electron microprobe analysis (Text S1, Table S2), the Na₂O/CaO ratio in



plagioclase feldspar in COL_{teph} and TUN_{teph} (both ~0.5) is higher than in ETN_{teph} and KIL_{teph} (both ~0.2), reflecting a greater proportion of the more ice-active NaAlSi₃O₈ relative to CaAl₂Si₂O₈ in the former samples. On the other hand, the K₂O/Na₂O ratio in alkali feldspar in NUO_{teph}, AST_{teph} and LAC_{teph} (1.2, 5.0, 1.9, respectively) does not support a link between these samples' INE and the proportion of KAlSi₃O₈ relative to NaAlSi₃O₈. This is consistent with the results of Whale et al. (2017),

5 who found that the INE of alkali feldspar does not relate to K content, but rather to the presence of perthitic intergrowth microtexture arising from phase separation (exsolution) into Na- and K-rich regions. Strain at the boundary of these regions gives rise to nanoscale topographic features that are suggested to be important in generating sites for ice nucleation (Whale et al., 2017).

Perthite in alkali feldspar develops in metamorphic and plutonic contexts during slow cooling at temperatures <700 °C (Parsons, 2010), and is generally not expected in volcanic ash which cools rapidly from magmatic down to ambient temperatures during eruption (Parsons et al., 2015). The low INE of LAC_{teph} could hence reflect a lack of perthitic microtexture, as noted for the alkali feldspar mineral sanidine (KAlSi₃O₈) sourced from the same geological setting (Eifel volcanic field), and similarly showing a poor ability to nucleate ice (Whale et al., 2017). In contrast, the high INE of NUO_{teph} and AST_{teph} is surprising in the absence of perthitic microtexture, and perhaps some other textural feature underlies their ability to nucleate

15 ice as effectively as alkali feldspar of non-pyroclastic origin (Fig. 5). Pyroclastic material from both the 1538 Monte Nuovo eruption (i.e., the origin of NUO_{teph}) and the ~4 ka Astroni eruption (i.e., the origin of AST_{teph}) has been found to contain anti-rapakivi overgrowth texture characterised by plagioclase feldspar cores rimmed by alkali feldspar (D'Oriano et al., 2005; Astbury et al., 2016; 2018). Such textures are challenging to resolve optically in powdered samples. Further optical and microanalytical (Scanning- and Transmission Electron Microscopy) observations (e.g., Whale et al., 2017; Holden et al., in

20 press) will be needed to explore whether the boundary between Na- and K-rich regions in anti-rapakivi texture may give rise to nanoscale topography that induces effects analogous to perthitic microtexture in promoting ice nucleation.

In addition to feldspar, several of the tephra samples contain pyroxene (Table 2). Specifically, the presence of orthopyroxene distinguishes COL_{teph} and TUN_{teph} from the other tephra (Fig. 4d), raising the question of whether it may underlie the high INE of these two samples (i.e., rather than plagioclase feldspar). We are aware of only a few early studies on ice nucleation by

25 ortho- and clinopyroxene minerals (hypersthene, augite; Hama and Itoo, 1956; Isono and Ikebe, 1960), but these studies have only reported semi-quantitative onset freezing temperatures (between -8 and -15 °C). Therefore, at present we cannot rule out a potential influence of pyroxene minerals on ice nucleation by volcanic ash. Additional research is needed to quantify the INE of a range of pyroxene minerals in order to better inform this assessment.

4.3 Chemical composition

To explore a potential link between chemical composition and INE, the $T_{n_g \approx 1 \text{ cm}^{-2}}$ values of the tephra and glass samples are plotted as a function of SiO₂, Al₂O₃, Fe₂O₃, MgO, CaO, Na₂O, K₂O, TiO₂, MnO and P₂O₅ contents in Fig. S2. No clear correlations are observed in any of these scatter plots to indicate a compositional dependency of the tephra or glass INE. This stands in apparent contrast to the recent work of Genereau et al. (2018) (based on a sample set of two rhyolites and three basalts), who reported that the $n_s(T)$ of volcanic ash correlates positively with K₂O content at -25 °C and negatively with TiO₂

35 and MnO contents from -30 to -35 °C.

The glass samples, in lacking crystalline minerals, are well-suited to assess for any direct relationships between INE and specific element oxide abundances. However, due to the overlap of droplet freezing temperatures of the glass suspensions and the background water (Fig. 2b), our ability to distinguish differences in INE across the nine samples is impeded. While CID_{glass} and COL_{glass} are the most ice-active glass samples, with signals clearly above the background (Fig. 2b, d), they represent

40 intermediate chemical compositions and thus their behaviour does not support any simple link between ash INE and chemical composition.



In contrast, the tephra samples are characterised by variations in crystallinity and mineralogy as well as composition, which convolutes the assessment of relationships between INE and specific element oxide abundances. However, if the crystallinity and mineralogy of the tephra samples are taken into consideration, a broad pattern emerges in the plots of $T_{n_s \approx 1 \text{ cm}^{-2}}$ versus Fe_2O_3 , MgO , and CaO contents (Fig. S2c-e). Excluding a cluster of three samples with comparatively low crystallinities (LIP_{teph}, CID_{teph}, LAC_{teph}), the $T_{n_s \approx 1 \text{ cm}^{-2}}$ decreases with increasing Fe_2O_3 , MgO , and CaO contents for NUO_{teph}, AST_{teph}, COL_{teph}, TUN_{teph}, ETN_{teph}, and KIL_{teph} (Fig. 6), in an order consistent with interpretations relating to their feldspar contents and chemistries (see discussion Sect. 4.2). This conforms to the notion of an indirect relationship between chemical composition and volcanic ash INE, whereby $\text{FeO}/\text{Fe}_2\text{O}_3$, MgO , and CaO contents increase from felsic to mafic magma, influencing the mineral phases that can crystallise from the magma and hence exist in the resultant ash (Fig. 1b).

10 5 Conclusions and implications

Here we used nine compositionally analogous pairs of natural tephra and remelted and quenched glass to investigate the influence of chemical composition, crystallinity and mineralogy on the INE of volcanic ash. The higher INE of the tephra relative to the glass strongly suggests that the presence of crystalline phases promotes ice nucleation. The large variability in INE of the tephra is inferred to reflect an influence of mineralogy - and hence an indirect influence of magma composition - on ice nucleation. As in desert dust, alkali feldspar is probably the most ice-active component in volcanic ash, conferring the highest INE to NUO_{teph} and AST_{teph} in this study. However, the ability of alkali feldspar in ash to nucleate ice likely cannot unequivocally be attributed to perthitic microtexture, as has been done for alkali feldspar of non-pyroclastic origin. Additional research is needed to explore whether other textural features in ash may elicit a similar effect in promoting ice nucleation. Further, the presence of alkali feldspar is neither always sufficient nor necessary for effective ice nucleation by ash. The high INE of COL_{teph} and TUN_{teph} may alternatively reflect very ice-active plagioclase feldspar, or possibly orthopyroxene which is present exclusively in these studied tephra. Previous studies on Soufrière Hills ash, also lacking alkali feldspar and containing plagioclase feldspar and orthopyroxene, have reported both a low and high INE of this ash (Schill et al., 2015; Mangan et al., 2017). Future studies quantifying the INE of individual crystalline phases found in ash will be necessary to unravel the precise role of mineralogy in volcanic ash ice nucleation.

An improved knowledge of the link between particular ash properties and ash INE may ultimately enhance predictive capability regarding volcanic eruptions likely to generate ice-active material. For example, as crystalline phases are primarily controlled by magma composition and storage/ascent conditions (Rogers, 2015), we speculate that highly ice-active ash particles might be erupted by volcanoes with intermediate to felsic alkaline magmas *and* possessing a history of magma mixing leading to feldspar crystals featuring overgrowth textures (e.g., Astbury et al., 2016; 2018). In addition, an eruption producing an abundance of crystal-bearing particles is expected to elicit a greater impact on heterogeneous ice nucleation than an eruption producing an abundance of crystal-free glass particles, all else being equal. Accordingly, massive outputs from the largest and most explosive eruptions, corresponding to violent caldera-forming ignimbrite events that generate ash clouds dominated by the glassy component (Sparks et al., 1997; Cather et al., 2009), may be less efficient in affecting INP populations than ash emissions from smaller eruptions. Further, since airborne ash typically becomes enriched in glassy fragments during long-range transport due to earlier gravitational settling of crystalline fragments (Hinkley et al., 1982), the INE of a suspended ash population is expected to decrease over time and distance from the volcano.

Lastly, it must be noted that once ash particles are generated, their surface properties can be altered by interactions with gases and condensates (e.g., H_2O , SO_2 , H_2SO_4 , HCl , HF) at variable temperatures in the eruption plume/cloud and ambient atmosphere (Delmelle et al., 2007; Ayris et al., 2013; 2014; Maters et al., 2016; 2017). The effects of such interactions on ash INE are not known, although it has been suggested from field measurements that volcanic gases may deactivate INPs (Schnell



and Delany, 1976; Schnell et al., 1982). Laboratory studies on desert dust show that ‘aging’ of dust particle surfaces by exposure to H₂SO₄ vapours at elevated temperatures reduces dust INE (Sullivan et al., 2010; Niedermeier et al., 2011). Therefore, exploring potential eruptive and atmospheric controls on ash INE is an important next step towards developing a better understanding of the capacity of volcanic ash emissions to affect heterogeneous ice nucleation during their airborne
5 lifetime.

Data availability: e.c.maters@leeds.ac.uk

Author contribution: E.M. designed the study and carried out the experiments. D.D. and C.C. provided the tephra samples and produced the glass samples. D.M. performed chemical and mineralogical analyses of these samples. T.W. performed the Poisson Monte Carlo error calculations. B.M supervised the project and provided insight on data interpretation. E.M. wrote
10 the manuscript with contributions from all co-authors.

Competing interests: The authors declare that they have no conflict of interest.

Acknowledgements: E.M. is funded by the European Union’s Horizon 2020 Research and Innovation Programme under the Marie Skłodowska-Curie Actions grant agreement No. 746695 (INoVA project). B.M. also acknowledges the European Research Council (MarineIce: No. 648661; CryoProtect: No. 713664) for funding. The authors wish to thank Ulrich Küppers
15 for collecting many of the tephra samples used in this study, and Bruce Houghton who made possible the collection of Kilauea achneliths. We are also grateful to William Orsi for enabling access to ultrapure water facilities at LMU, Alex Harrison for providing the plagioclase and alkali feldspar parameterisations, and to Sebastien Sikora, Lesley Neve, Fiona Keay and Andy Connelly for laboratory assistance.

References

- 20 Astbury, R. L., Petrelli, M., Arienzo, I., D’Antonio, M., Morgavi, D., and Perugini, D.: Using trace element mapping to identify discrete magma mixing events from the Astroni 6 eruption, in: American Geophysical Union Fall Meeting 2016, San Francisco, United States of America, 12-16 December 2016, 2016AGUFM.V33E3176A, 2016.
- Astbury, R. L., Petrelli, M., Ubide, T., Stock, M. J., Arienzo, I., D’Antonio, M., and Perugini, D.: Tracking plumbing system dynamics at the Campi Flegrei caldera, Italy: High-resolution trace element mapping of the Astroni crystal cargo, *Lithos*,
25 318-319, 464-477, <https://doi.org/10.1016/j.lithos.2018.08.033>, 2018.
- Atkinson, J. D., Murray, B. J., Woodhouse, M. T., Whale, T. F., Baustian, K. J., Carslaw, K. S., Dobbie, S., O’Sullivan, D., and Malkin, T. L.: The importance of feldspar for ice nucleation by mineral dust in mixed-phase clouds, *Nature*, 498, 355-358, doi:10.1038/nature12278, 2013.
- Ayris, P. M., Lee, A. F., Wilson, K., Kueppers, U., Dingwell, D. B., and Delmelle, P.: SO₂ sequestration in large volcanic
30 eruptions: high-temperature scavenging by tephra, *Geochim. Cosmochim. Ac.*, 110, 58-69, <https://doi.org/10.1016/j.gca.2013.02.018>, 2013.
- Ayris, P. M., Delmelle, P., Cimarelli, C., Maters, E. C., Suzuki, Y. J., and Dingwell, D. B.: HCl uptake by volcanic ash in the high temperature eruption plume: Mechanistic insights, *Geochim. Cosmochim. Ac.*, 144, 188-201, <https://doi.org/10.1016/j.gca.2014.08.028>, 2014.
- 35 Brunauer, S., Emmett, P. H., and Teller, E.: Adsorption of gases in multimolecular layers, *J. Am. Chem. Soc.*, 60, 309-319, doi:10.1021/ja01269a023, 1938.



- Cather, S. M., Dunbar, N. W., McDowell, F. W., McIntosh, W. C., and Scholle, P. A.: Climate forcing by iron fertilization from repeated ignimbrite eruptions: The icehouse silicic large igneous province (SLIP) hypothesis, *Geosphere*, 5, 315–324, <https://doi.org/10.1130/GES00188.1>, 2009.
- Cimarelli, C., Alatorre-Ibargüenogitia, M. A., Aizawa, K., Yokoo, A., Díaz-Marina, A., Iguchi, M., and Dingwell D. B.:
5 Multiparametric observation of volcanic lightning: Sakurajima Volcano, Japan, *Geophys. Res. Lett.*, 43, 4221–4228, <https://doi.org/10.1002/2015GL067445>, 2016.
- Connolly, P. J., Möhler, O., Field, P. R., Saathoff, H., Burgess, R., Choulaton, T., and Gallagher, M.: Studies of heterogeneous freezing by three different desert dust samples, *Atmos. Chem. Phys.*, 9, 2805–2824, <https://doi.org/10.5194/acp-9-2805-2009>, 2009.
- 10 Delmelle, P., Lambert, M., Dufrière, Y., Gerin, P., and Óskarsson, N.: Gas/aerosol-ash interaction in volcanic plumes: New insights from surface analysis of fine ash particles, *Earth Planet Sc. Lett.*, 259, 159–170, <https://doi.org/10.1016/j.epsl.2007.04.052>, 2007.
- Döbelin, N., and Kleeberg, R.: Profex: a graphical user interface for the Rietveld refinement program BGMN, *J. Appl. Crystallogr.*, 48, 1573–1580, <https://doi.org/10.1107/S1600576715014685>, 2015.
- 15 D’Oriano, C., Poggianti, E., Bertagnini, A., Cioni, R., Landi, P., Polacci, M., and Rosi, M.: Changes in eruptive style during the A.D. 1538 Monte Nuovo eruption (Phlegrean Fields, Italy): the role of syn-eruptive crystallization, *B. Volcanol.*, 67, 601–621, doi:10.1007/s00445-004-0397-z, 2005.
- Durant, A. J., Shaw, R. A., Rose, W. I., Mi, Y., and Ernst, G. G.: Ice nucleation and overseeding of ice in volcanic clouds, *J. Geophys. Res.*, 113, D09206, <https://doi.org/10.1029/2007JD009064>, 2008.
- 20 Durant, A. J., Bonadonna, C., Horwell, C. J.: Atmospheric and environmental impacts of volcanic particulates, *Elements*, 6, 235–240, <https://doi.org/10.2113/gselements.6.4.235>, 2010.
- Genareau, K., Cloer, S. M., Primm, K., Tolbert, M. A., and Woods, T. W.: Compositional and mineralogical effects on ice nucleation activity of volcanic ash, *Atmosphere*, 9, 238, <https://doi.org/10.3390/atmos9070238>, 2018.
- Guo, S., Bluth, G. J. S., Rose, W. I., Watson, I. M., and Prata, A. J.: Re-evaluation of SO₂ release of the 15 June 1991 Pinatubo
25 eruption using ultraviolet and infrared satellite sensors, *Geochem. Geophys. Geosy.*, 5, Q04001, <https://doi.org/10.1029/2003GC000654>, 2004a.
- Guo, S., Rose, W. I., Bluth, G. J. S., and Watson, I. M.: Particles in the great Pinatubo volcanic cloud of June 1991: The role of ice, *Geochem. Geophys. Geosy.*, 5, Q05003, <https://doi.org/10.1029/2003GC000655>, 2004b.
- Hama, K., and Ito, K.: Freezing of supercooled water-droplets (II), *Pap. Meteorol. Geophys.*, 7, 99–106,
30 https://doi.org/10.2467/mripapers1950.7.2_99, 1956.
- Harrison, A. D., Lever, K., Sanchez-Marroquin, A., Holden, M. A., Whale, T. F., McQuaid, J. B., and Murray, B. J.: The ice nucleating efficiency of a variety of quartz samples and their atmospheric importance compared to K-feldspar. In preparation for *Atmos. Chem. Phys.*
- Harrison, A. D., Whale, T. F., Carpenter, M. A., Holden, M. A., Neve, L., O’Sullivan, D., Vergara Temprado, J., and Murray
35 B. J.: Not all feldspars are equal: a survey of ice nucleating properties across the feldspar group of minerals, *Atmos. Chem. Phys.*, 16, 10927–10940, <https://doi.org/10.5194/acp-16-10927-2016>, 2016.
- Heiken, G., and Wohletz, K. (Eds.): *Volcanic Ash*, University of California Press, London, United Kingdom, 1992.
- Hinkley, T. K., Smith, K. S., Taggart, J. E., and Brown, J. T.: Chemical and mineralogic aspects of observed fractionation of ash from May 18, 1980 eruption of Mount St. Helens, U.S. Geological Survey Professional Paper 1397-A, 10–22, 1982.
- 40 Hobbs, P. V., Fullerton, C. M., and Bluhm, G. C.: Ice nucleus storms in Hawaii, *Nat. Phys.*, 230, 90–91, [10.1038/physci230090a0](https://doi.org/10.1038/physci230090a0), 1971.
- Holden, M. A., Whale, T. F., Tarn, M. D., O’Sullivan, D., Walshaw, R. D., Murray, B. J., Meldrum, F. C., and Christenson, H. K.: High-speed imaging of ice nucleation in water proves the existence of active site, *Science Advances*, in press.



- Hoose, C., Kristjánsson, J. E., Chen, J.-P., Hazra, A.: A classical-theory-based parameterization of heterogeneous ice nucleation by mineral dust, soot, and biological particles in a global climate model, *J. Atmos. Sci.*, 67, 2483-2503, <https://doi.org/10.1175/2010JAS3425.1>, 2010.
- Herzog, M., Graf, H. F., Textor, C., and Oberhuber, J. M.: The effect of phase changes of water on the development of volcanic plumes. *J. Volcanol. Geoth. Res.*, 87, 55-74, [https://doi.org/10.1016/S0377-0273\(98\)00100-0](https://doi.org/10.1016/S0377-0273(98)00100-0), 1998.
- 5 Ignatius, K., Kristensen, T. B., Järvinen, E., Nichman, L., Fuchs, C., Gordon, H., Herenz, P., Hoyle, C. R., Duplissy, J., Garimella, S., Dias, A., Frege, C., Höppl, N., Tröstl, J., Wagner, R., Yan, C., Amorim, A., Baltensperger, U., Curtius, J., Donahue, N. M., Gallagher, M. W., Kirkby, J., Kulmala, M., Möhler, O., Saathoff, H., Schnaiter, M., Tomé, A., Virtanen, A., Worsnop, D., and Stratmann, F.: Heterogeneous ice nucleation of viscous secondary organic aerosol produced from ozonolysis of α -pinene. *Atmos. Chem. Phys.*, 16, 6495-6509, <https://doi.org/10.5194/acp-16-6495-2016>, 2016.
- 10 Isono, K., and Ikebe, Y.: On the ice-nucleating ability of rock-forming minerals and soil particles, *J. Meteorol. Soc. Jpn.*, 38, 213-230, https://doi.org/10.2151/jmsj1923.38.5_213, 1960.
- Isono, K., and Komabayasi, M.: The influence of volcanic dust on precipitation, *J. Meteorol. Soc. Jpn.*, 32, 345-353, https://doi.org/10.2151/jmsj1923.32.11-12_345, 1954.
- 15 Isono, K., Komabayasi, M., and Ono, A.: Volcanoes as a source of atmospheric ice nuclei. *Nature*, 183, 317-318, <https://doi.org/10.1038/183317a0>, 1959a.
- Isono, K., Komabayasi, M., and Ono, A.: The nature and origin of ice nuclei in the atmosphere, *J. Meteorol. Soc. Jpn.*, 37, 211-233, https://doi.org/10.2151/jmsj1923.37.6_211, 1959b.
- Kaufmann, L., Marcolli, C., Hofer, J., Pinti, V., Hoyle, C. R., and Peter, T.: Ice nucleation efficiency of natural dust samples in the immersion mode, *Atmos. Chem. Phys.*, 16, 11177-11206, <https://doi.org/10.5194/acp-16-11177-2016>, 2016.
- 20 Komabayasi, M.: The suppression of thunder cloud occurrence by frequent volcanic eruptions, *J. Meteorol. Soc. Jpn.*, 35A, 25-30, https://doi.org/10.2151/jmsj1923.35A.0_25, 1957.
- Koop, T., and Murray, B. J.: A physically constrained classical description of the homogeneous nucleation of ice in water, *Journal of Chemical Physics*, 145, 211915, 2016.
- 25 Kulkarni, G., Nandasiri, M., Zelenyuk, A., Beranek, J., Madaan, N., Devaraj, A., Shutthanandan, V., Thevuthasan, S., and Varga, T.: Effects of crystallographic properties on the ice nucleation properties of volcanic ash particles, *Geophys. Res. Lett.*, 42, 3048-3055, <https://doi.org/10.1002/2015GL063270>, 2015.
- Le Maitre, R. W., Streckeisen, A., Zanettin, B., Le Bas, M. J., Bonin, B., and Bateman, P. (Eds.) *Igneous Rocks: A Classification and Glossary of Terms, Recommendations of the International Union of Geological Sciences Subcommission on the Systematics of Igneous Rocks*, Cambridge University Press, Cambridge, United Kingdom, 2002.
- 30 Mangan, T. P., Atkinson, J. D., Neuberger, J. W., O'Sullivan, D., Wilson, T. W., Whale, T. F., Neve, L., Umo, N. S., Malkin, T. L., and Murray, B. J.: Heterogeneous ice nucleation by Soufriere Hills volcanic ash immersed in water droplets, *PLoS ONE*, 12, e0169720, <https://doi.org/10.1371/journal.pone.0169720>, 2017.
- Maters, E.C., Delmelle, P., Rossi, M.J., Ayriss, P.M., and Bernard, A.: Controls on the surface chemical reactivity of volcanic ash investigated with probe gases, *Earth Planet Sc. Lett.*, 450, 254-262, <https://doi.org/10.1016/j.epsl.2016.06.044>, 2016.
- 35 Maters, E.C., Delmelle, P., Rossi, M.J., and Ayriss, P.M.: Reactive uptake of sulfur dioxide and ozone on volcanic glass and ash at ambient temperature, *J. Geophys. Res. - Atmos.*, 122, 10077-10088, <https://doi.org/10.1002/2017JD026993>, 2017.
- McNutt, S. R., and Williams, E. R.: Volcanic lightning: global observations and constraints on source mechanisms, *B. Volcanol.*, 72, 1153-1167, doi: 10.1007/s00445-010-0393-4, 2010.
- 40 Murray, B. J., Broadley, S. L., Wilson, T. W., Bull, S. J., Wills, R. H., Christenson, H. K., and Murray, E. J.: Kinetics of homogeneous freezing of water, *Phys. Chem. Chem. Phys.*, 12, 10380-10387, doi:10.1039/c003297b, 2010a.



- Murray, B. J., Wilson, T. W., Dobbie, S., Cui, Z., Al-Jumur, S. M. R. K., Möhler, O., Schnaiter, M., Wagner, R., Benz, S., Niemand, M., Saathoff, H., Ebert, V., Wagner, S., and Kärcher, B.: Heterogeneous nucleation of ice particles on glassy aerosols under cirrus conditions, *Nat. Geosci.*, 3, 233-237, <https://doi.org/10.1038/ngeo817>, 2010b.
- Murray, B. J., O'Sullivan, D., Atkinson, J. D., and Webb, M. E.: Ice nucleation by particles immersed in supercooled cloud droplets, *Chem. Soc. Rev.*, 41, 6519-6554, doi:10.1039/c2cs35200a, 2012.
- Niedermeier, D., Hartmann, S., Clauss, T., Wex, H., Kiselev, A., Sullivan, R. C., DeMott, P.J., Petters, M. D., Reitz, P., Schneider, J., Mikhailov, E., Sierau, B., Stetzer, O., Reimann, B., Bundke, U., Shaw, R. A., Buchholz, A., Mentel, T. F., and Stratmann, F.: Experimental study of the role of physicochemical surface processing on the IN ability of mineral dust particles, *Atmos. Chem. Phys.*, 11, 11131-11144, <https://doi.org/10.5194/acp-11-11131-2011>, 2011.
- 10 Parsons, I.: Feldspars defined and described: a pair of posters published by the Mineralogical Society. Sources and supporting information, *Mineral. Mag.*, 74, 529-551, <https://doi.org/10.1180/minmag.2010.074.3.529>, 2010.
- Parsons, I., Fitz Gerald, J. D., Lee, M. R.: Routine characterization and interpretation of complex alkali feldspar intergrowths, *Am. Mineral.*, 100, 1277-1303, <https://doi.org/10.2138/am-2015-5094>, 2015.
- Penner, J. E., Andreae, M., Annegarn, H., Barrie, L., Feichter, J., Hegg, D., Jayaraman, A., Leitch, R., Murphy, D., Nganga, J., and Pitari, G.: Aerosols, their direct and indirect effects, in: *Climate Change 2001: The Scientific Basis, Contribution of Working Group I to IPCC*, Cambridge University Press, Cambridge, United Kingdom, 290-347, 2001.
- 15 Polen, M., Brubaker, T., Somers, J., and Sullivan, R. C.: Cleaning up our water: reducing interferences from non-homogeneous freezing of "pure" water in droplet freezing assays of ice nucleating particles, *Atmos. Meas. Tech.*, 11, 5315-5334, <https://doi.org/10.5194/amt-11-5315-2018>, 2018.
- 20 Pruppacher, H. R., and Klett, J. D. (Eds.) *Microphysics of Clouds and Precipitation*, Second Edition, Springer, Netherlands, 2010.
- Rogers, N.: The Composition and Origin of Magmas, in: *The Encyclopedia of Volcanoes*, Second Edition, edited by: Sigurdsson, H., Houghton, B., McNutt, S. R., Rymer, H., and Stix, J., Academic Press, London, United Kingdom, 93-112, <https://doi.org/10.1016/B978-0-12-385938-9.00004-3>, 2015.
- 25 Rose, W. I., Gu, Y., Watson, M., Yu, T., Bluth, G. J. S., Prata, A. J., Krueger, A. J., Krotkov, N., Carn, S., Fromm, M. D., Hunton, D. E., Ernst, G. G. J., Viggiano, A. A., Miller, T. M., Ballenthin, J. O., Reeves, J. M., Wilson, J. C., Anderson, B. E., and Flittner, D. E.: The February-March 2000 Eruption of Hekla, Iceland from a Satellite Perspective, in: *Volcanism and the Earth's Atmosphere*, Volume 139, edited by: Robock A., and Oppenheimer, C., American Geophysical Union, United States of America, 107-132, <https://doi.org/10.1029/139GM07>, 2003.
- 30 Schill, G. P., Genareau, K., and Tolbert, M. A.: Deposition and immersion-mode nucleation of ice by three distinct samples of volcanic ash, *Atmos. Chem. Phys.*, 15, 7523-7536, <https://doi.org/10.5194/acp-15-7523-2015>, 2015.
- Schnell, R. C., and Delany, A. C.: Airborne ice nuclei near an active volcano, *Nature*, 264, 535-536, <https://doi.org/10.1038/264535a0>, 1976.
- Schnell, R. C., Pueschel, R. F., and Wellman, D. L.: Ice nucleus characteristics of Mount St. Helens effluents, *J. Geophys. Res.*, 87, 11109-11112, <https://doi.org/10.1029/JC087iC13p11109>, 1982.
- 35 Seifert, P., Ansmann, A., Groß, S., Freudenthaler, V., Heinold, B., Hiesch, A., Mattis, I., Schmidt, J., Schnell, F., Tesche, M., Wandinger, U., and Wiegner, M.: Ice formation in ash-influenced clouds after the eruption of the Eyjafjallajökull volcano in April 2010, *J. Geophys. Res.*, 116, D00U04, <https://doi.org/10.1029/2011JD015702>, 2011.
- Sparks, R. S. J., Bursik, M. I., Carey, S. N., Gilbert, J. S., Glaze, L. S., Sigurdsson, H., and Woods, A. W.: *Volcanic Plumes*, John Wiley & Sons, University of California, United States of America, 1997.
- 40 Sullivan, R. C., Petters, M. D., DeMott, P. J., Kreidenweis, S. M., Wex, H., Niedermeier, D., Hartmann, S., Clauss, T., Stratmann, F., Reitz, P., Schneider, J., and Sierau, B.: Irreversible loss of ice nucleation active sites in mineral dust particles



- caused by sulphuric acid condensation, *Atmos. Chem. Phys.*, 10, 11471-11487, <https://doi.org/10.5194/acp-10-11471-2010>, 2010.
- Textor, C., Graf, H. -F., Herzog, M., and Oberhuber, J. M.: Injection of gases into the stratosphere by explosive volcanic eruptions, *J. Geophys. Res.*, 108, 4606, <https://doi.org/10.1029/2002JD002987>, 2003.
- 5 Textor, C., Graf, H. -F., Herzog, M., Oberhuber, J. M., Rose, W. I., and Ernst, G. G. J.: Volcanic particle aggregation in explosive eruption columns. Part I: Parameterization of the microphysics of hydrometeors and ash, *J. Volcanol. Geoth. Res.*, 150, 359-377, <https://doi.org/10.1016/j.jvolgeores.2005.09.007>, 2006.
- Vali, G.: Interpretation of freezing nucleation experiments: singular and stochastic; sites and surfaces, *Atmos. Chem. Phys.*, 14, 5271-5294, <https://doi.org/10.5194/acp-14-5271-2014>, 2014.
- 10 Van Eaton, A. R., Mastin, L. G., Herzog, M., Schwaiger, H. F., Schneider, D. J., Wallace, K. L., and Clarke, A. B.: Hail formation triggers rapid ash aggregation in volcanic plumes, *Nat. Commun.*, 6, 7860, <https://doi.org/10.1038/ncomms8860>, 2015.
- Vergara-Temprado, J., Murray, B. J., Wilson, T. W., O'Sullivan, D., Browse, J., Pringle, K. J., Ardon-Dryer, K., Bertram, A. K., Burrows, S. M., Ceburnis, D., DeMott, P. J., Mason, R. H., O'Dowd, C. D., Rinaldi, M., and Carslaw, K. S.: Contribution of feldspar and marine organic aerosols to global ice nucleating particle concentrations, *Atmos. Chem. Phys.*, 17, 3637-3658, <https://doi.org/10.5194/acp-17-3637-2017>, 2017.
- 15 Whale, T. F., Murray, B. J., O'Sullivan, D., Wilson, T. W., Umo, N. S., Baustian, K. J., J. D. Atkinson, J. D., Workneh, D. A., and Morris, G. J.: A technique for quantifying heterogeneous ice nucleation in microlitre supercooled water droplets, *Atmos. Meas. Tech.*, 8, 2437-2447, <https://doi.org/10.5194/amt-8-2437-2015>, 2015.
- 20 Whale, T. F., Holden, M. A., Kulak, A. N., Kim, Y.-Y., Meldrum, F. C., Christenson, H. K., and Murray, B. J.: The role of phase separation and related topography in the exceptional ice-nucleating ability of alkali feldspars, *Phys. Chem. Chem. Phys.*, 19, 31186-31193, doi:10.1039/c7cp04898j2, 017.
- Wagner, R., Möhler, O., Saathoff, H., Schnaiter, M., Skrotzki, J., Leisner, T., Wilson, T. W., Malkin, T. L., Murray, B. J.: Ice cloud processing of ultra-viscous/glassy aerosol particles leads to enhanced ice nucleation ability, *Atmos. Chem. Phys.*, 12, 8589-8610, <https://doi.org/10.5194/acp-12-8589-2012>, 2012.
- 25 Wright, H. M. N., Cashman, K. V., Mothes, P. A., Hall, M. L., Ruiz, A. G., and Le Pennec, J.-L.: Estimating rates of decompression from textures of erupted ash particles produced by 1999–2006 eruptions of Tungurahua volcano, Ecuador, *Geology*, 40, 619-622, <https://doi.org/10.1130/G32948.1>, 2012.
- 30 Yakobi-Hancock, J. D., Ladino, L. A., and Abbat, J. P. D.: Feldspar minerals as efficient deposition ice nuclei, *Atmos. Chem. Phys.*, 13, 11175-11185, <https://doi.org/10.5194/acp-13-11175-2013>, 2013.

**Table 1.** Details of the volcanic tephra and glass samples used in this study.

Sample code	Source volcano	Eruption date ^a	Classification ^b	SSA _{BET} ^c (tephra/glass) (m ² g ⁻¹)
LIP _{teph/glass}	Lipari (Italy)	1230	rhyolite	1.8/1.1
COL _{teph/glass}	Colima (Mexico)	Jan-Feb 2017	andesite	1.9/0.9
TUN _{teph/glass}	Tungurahua (Ecuador)	Feb 2014	andesite	1.4/1.1
CID _{teph/glass}	Sete Cidades (Portugal)	16 ka	trachyte	1.6/1.4
AST _{teph/glass}	Astroni (Italy)	3.8-4.4 ka	trachyphonolite	3.7/1.4
NUO _{teph/glass}	Monte Nuovo (Italy)	Sept-Oct 1538	trachyphonolite	4.6/1.3
LAC _{teph/glass}	Laacher See (Germany)	12.9 ka	phonolite	3.3/0.9
ETN _{teph/glass}	Mount Etna (Italy)	July 2014	trachybasalt	1.7/1.1
KIL _{teph/glass}	Kilauea (Hawaii)	July 2018	basalt	2.1/1.1

^aRefers to the eruption of origin of the tephra material. This does not apply to the glass material as it has been synthesised (from tephra) in the laboratory by a melting, homogenising and quenching protocol. ^bAccording to the Total Alkali versus Silica igneous rock classification diagram (Fig. 1) based on chemical composition (Table S1). ^cUncertainty is in the range of 0.5-1.2 %.

5

Table 2. Crystallinity and mineralogy of the tephra samples used in this study, in wt. %.

Sample ^a	Crystallinity	Alkali (K-rich) feldspar	Plagioclase (Na-/Ca-rich) feldspar	Clino- pyroxene	Ortho- pyroxene	Quartz	Fe(-Ti) oxide	Olivine
LIP _{teph}	<2	-	-	-	-	-	m.c.	-
COL _{teph}	62	m.c.	55	-	7	m.c.	m.c.	-
TUN _{teph}	54	-	43	6	5	-	m.c.	-
CID _{teph}	<2	m.c.	-	m.c.	-	-	m.c.	-
AST _{teph}	28	19	7	2	-	-	m.c.	-
NUO _{teph}	60	60	-	-	-	-	m.c.	-
LAC _{teph}	11	9	-	-	-	2	-	-
ETN _{teph}	66	-	44	22	-	-	m.c.	-
KIL _{teph}	3	-	3	-	-	-	-	m.c.

^aSample codes are listed in Table 1. m.c. = minor component; below ~2 wt. % detection limit.




Figure 1. (a) Total Alkali versus Silica diagram showing the classification of the tephra (red symbols) and glass (blue symbols) used in this study. Sample codes are listed in Table 1. Modified after Le Maitre et al. (2002). (b) Schematic summarising the mineralogy of common igneous rock types. Modified after Rogers (2015).




Figure 2. Droplet fraction frozen (f_{ice}) as a function of temperature for 1 wt.% suspensions of (a) tephra or (b) glass in water. The grey bands represent the spread of $f_{ice}(T)$ measurements (mean values \pm standard deviation) of the background water (i.e., containing no added sample). Ice nucleation active site density (n_s) as a function of temperature for 1 wt.% suspensions of (c)

- 5 tephra or (d) glass in water. For the sake of comparison, the grey curves represent theoretical upper limit $n_s(T)$ values of the background water, calculated using the upper limit $f_{ice}(T)$ measurements (mean values + standard deviation) of the background water, and assuming it contains particles with SSA_{BET} values equal to the lowest from the tephra and glass sets (1.4 and 0.9 m² g⁻¹, respectively). The tephra $n_s(T)$ values are well above this background but most of the glass $n_s(T)$ values should be regarded as upper limits. The uncertainty in $n_s(T)$ is shown as error bars for a subset of data points (of NUO_{teph} and CID_{glass}) and omitted from remaining data points for clarity, but is typical of all tephra and glass samples. Sample codes are listed in Table 1.
- 10




Figure 3. Ice nucleation active site density (n_s) as a function of temperature for 1 wt.% suspensions of tephra or glass in water. Each plot shows replicates of a compositionally analogous pair of tephra (red triangles) and glass (blue circles). The grey curves with triangle and circle symbols represent the detection limit for $n_s(T)$ based on the background water runs accompanying the tephra and glass experiments, respectively (see Fig. 2 caption for details). Sample codes are listed in Table 1.




Figure 4. The INE ($T_{n_s \approx 1 \text{ cm}^{-2}}$) of the tephra versus their content of (a) alkali feldspar, (b) plagioclase feldspar, (c) clinopyroxene, and (d) orthopyroxene. Note that minor components below the detection limit are plotted at 1 wt.%. Ice nucleation experiments were conducted with 1 wt.% suspensions of tephra in water. Sample codes are listed in Table 1.




Figure 5. Ice nucleation active site density (n_s) as a function of temperature for 1 wt.% suspensions of tephra in water. The blue and red lines represent parameterisations for, respectively, alkali and plagioclase feldspar of non-pyroclastic origin reported in Harrison et al. (in prep.) from a compilation of literature data. The solid lines indicate mean values and the dashed lines indicate lower and upper limits corresponding to the standard deviation of the mean. Sample codes are listed in Table 1.




Figure 6. The INE ($T_{n_s \approx 1 \text{ cm}^{-2}}$) of NUO_{teph}, AST_{teph}, COL_{teph}, TUN_{teph}, ETN_{teph}, and KIL_{teph} versus their (a) Fe_2O_3 , (b) MgO, and (c) CaO contents. The grey triangles correspond to tephra samples with comparatively low crystallinities (LIP_{teph}, CID_{teph}, LAC_{teph}) which are excluded from the trendline. Ice nucleation experiments were conducted with 1 wt.% suspensions of tephra in water. Sample codes are listed in Table 1.

5