

# Computation and analysis of atmospheric carbon dioxide annual mean growth rates from satellite observations during 2003-2016

Michael Buchwitz<sup>1</sup>, Maximilian Reuter<sup>1</sup>, Oliver Schneising<sup>1</sup>, Stefan Noël<sup>1</sup>, Bettina Gier<sup>1,2</sup>, Heinrich Bovensmann<sup>1</sup>, John P. Burrows<sup>1</sup>, Hartmut Boesch<sup>3,4</sup>, Jasdeep Anand<sup>3,4</sup>, Robert J. Parker<sup>3,4</sup>, Peter Somkuti<sup>3,4</sup>, Rob G. Detmers<sup>5</sup>, Otto P. Hasekamp<sup>5</sup>, Ilse Aben<sup>5</sup>, André Butz<sup>2,6,#</sup>, Akihiko Kuze<sup>7</sup>, Hiroshi Suto<sup>7</sup>, Yukio Yoshida<sup>8</sup>, David Crisp<sup>9</sup>, Christopher O'Dell<sup>10</sup>

<sup>1</sup>Institute of Environmental Physics (IUP), University of Bremen, Bremen, Germany

10 <sup>2</sup>Deutsches Zentrum für Luft- und Raumfahrt e.V. (DLR), Institut für Physik der Atmosphäre, Oberpfaffenhofen, Germany

<sup>3</sup>Earth Observation Science, University of Leicester, Leicester, UK

<sup>4</sup>NERC National Centre for Earth Observation, Leicester, UK

<sup>5</sup>SRON Netherlands Institute for Space Research, Utrecht, The Netherlands

15 <sup>6</sup>Meteorologisches Institut, Ludwig-Maximilians-Universität (LMU), Munich, Germany

<sup>7</sup>Japan Aerospace Exploration Agency (JAXA), Tsukuba, Japan

<sup>8</sup>National Institute for Environmental Studies (NIES), Tsukuba, Japan

<sup>9</sup>Jet Propulsion Laboratory (JPL), Pasadena, CA, USA

<sup>10</sup>Colorado State University (CSU), Fort Collins, CO, USA

20 <sup>#</sup>Now at: Institut für Umweltphysik, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

*Correspondence to:* Michael Buchwitz (Michael.Buchwitz@iup.physik.uni-bremen.de)

**Abstract.** The growth rate of atmospheric carbon dioxide (CO<sub>2</sub>) reflects the net effect of emissions and uptake resulting from anthropogenic and natural carbon sources and sinks. Annual mean CO<sub>2</sub> growth rates have been determined from satellite retrievals of column-average dry-air mole fractions of CO<sub>2</sub>, i.e., XCO<sub>2</sub>, for the years 2003 to 2016. The XCO<sub>2</sub> growth rates agree with National Oceanic and Atmospheric Administration (NOAA) growth rates from CO<sub>2</sub> surface observations within the uncertainty of the satellite-derived growth rates (mean difference ± standard deviation: 0.0±0.3 ppm/year; R: 0.82). This new and independent data set confirms record large growth rates around 3 ppm/year in 2015 and 2016, which are attributed to the 2015/2016 El Niño. Based on a comparison of the satellite-derived growth rates with human CO<sub>2</sub> emissions from fossil fuel combustion and with El Niño Southern Oscillation (ENSO) indices, we estimate by how much the impact of ENSO dominates the impact of fossil fuel burning related emissions in explaining the variance of the atmospheric CO<sub>2</sub> growth rate. Our analysis shows that the ENSO impact on CO<sub>2</sub> growth rate variations dominates over that of human emissions throughout the period 2003-2016 but in particular during the period 2010-2016 due to strong La Niña and El

25  
30

Niño events. Using the derived growth rates and their uncertainty, we estimate the probability that the impact of ENSO on the variability is larger than the impact of human emissions to be 63% for the time period 2003-2016. If the time period is restricted to 2010-2016 this probability increases to 94%.

## 5 1 Introduction

Atmospheric carbon dioxide (CO<sub>2</sub>) is an important greenhouse gas that causes global warming (IPCC 2013). Sources that emit CO<sub>2</sub> into the atmosphere include anthropogenic and natural sources at the surface, and the oxidation of carbon monoxide and hydrocarbons in the atmosphere. The sinks that remove CO<sub>2</sub> primarily at the surface include biological (photosynthesis) and physical (solubility) processes. Anthropogenic emissions of CO<sub>2</sub>, primarily from fossil fuel combustion, have increased the atmospheric CO<sub>2</sub> mixing ratios at the surface by more than 40% since pre-industrial times, from less than 280 parts per million (ppm) to 402.8±0.1 ppm in 2016 (Dlugokencky and Tans, 2017a). A global increase of atmospheric CO<sub>2</sub> by 1 ppm in a one-year time period corresponds to an annual increase of 2.12 GtC/year (Ballantyne et al., 2012). However, this increase in mass does not directly correspond to the emissions. The reason is that only a fraction of the emitted CO<sub>2</sub> remains in the atmosphere as CO<sub>2</sub> is partitioned between the atmosphere and ocean and land carbon sinks. On average, somewhat less than half of the emitted CO<sub>2</sub> remains in the atmosphere but this “airborne fraction” varies substantially from year to year (Le Quéré et al., 2016, 2018). Variations of the airborne fraction are not well understood primarily because of an inadequate understanding of the terrestrial carbon sink, which introduces large uncertainties for climate prediction (e.g., IPCC 2013; Peylin et al., 2013; Wieder et al., 2015; Huntzinger et al., 2017). Identification of the origin of changes of the growth rate requires additional information for the attribution to particular sources or sinks (Peters et al., 2017). Atmospheric CO<sub>2</sub> growth rates inferred from in-situ CO<sub>2</sub> surface measurements are regularly determined and published, for example, by the National Oceanic and Atmospheric Administration (NOAA) (see <https://www.esrl.noaa.gov/gmd/ccgg/trends/gr.html>). In this study, we present and interpret atmospheric growth rates determined from the remote sensing of CO<sub>2</sub> vertical columns from space, which are described in the following section.

## 2 Global satellite observations of atmospheric CO<sub>2</sub> columns

Satellites provide retrievals of CO<sub>2</sub> vertical columns in terms of the CO<sub>2</sub> column-average dry-air mole fraction, denoted XCO<sub>2</sub>. Although a relatively new field, satellite-based XCO<sub>2</sub> data products have already been used to improve our knowledge of natural (e.g., Basu et al., 2013; Maksyutov et al., 2013; Chevallier et al., 2014; Reuter et al., 2014a; Schneising et al., 2014; Houweling et al., 2014; Parker et al., 2016; Heymann et al., 2017; Liu et al., 2017; Kaminski et al., 2017) and anthropogenic (e.g., Schneising et al., 2013; Reuter et al., 2014b; Kort et al., 2012; Hakkarainen et al., 2016; Nassar et al., 2017) CO<sub>2</sub> sources and sinks but only a few studies explicitly present and discuss CO<sub>2</sub> growth rates. Buchwitz et al., 2007, analyzed the first three years (2003-2005) of XCO<sub>2</sub> retrievals from SCIAMACHY/ENVISAT (Burrows et al., 1995; Bovensmann et al., 1999) generated using the WFM-DOAS retrieval algorithm (Buchwitz et al., 2006). They computed year-to-year CO<sub>2</sub> variations and compared the XCO<sub>2</sub> increase with the XCO<sub>2</sub> increase computed from the output of NOAA’s CO<sub>2</sub> assimilating

system CarbonTracker (Peters et al., 2007) and found agreement within 1 ppm/year. Schneising et al., 2014, computed growth rates from the 2003-2011 SCIAMACHY XCO<sub>2</sub> record. They compared the derived annual growth rates with surface temperature and found that years having higher temperatures during the vegetation growing season are associated with larger growth rates in atmospheric CO<sub>2</sub> at northern mid-latitudes. Growth rates from GOSAT (Kuze et al., 2016) are published by the National Institute for Environmental Studies (NIES), Tsukuba, Japan (NIES 2017).

In this study, we analyze a new satellite XCO<sub>2</sub> data set covering 14 years (2003-2016) generated from SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT. We use the XCO<sub>2</sub> data product Obs4MIPs (Observations for Model Intercomparisons Project) version 3 (O4Mv3), which is a gridded (Level 3) monthly data product at 5° latitude by 5° longitude spatial resolution in Obs4MIPs format (Buchwitz et al., 2017a). Obs4MIPs (<https://www.earthsystemcog.org/projects/obs4mips/>) is an activity to make observational products more accessible for climate model intercomparisons (e.g., Lauer et al., 2017). The O4Mv3 XCO<sub>2</sub> data product was generated by gridding (averaging) the XCO<sub>2</sub> Level 2 (i.e., individual soundings) product generated with the Ensemble Median Algorithm (EMMA, Reuter et al., 2013). EMMA uses as input an ensemble of XCO<sub>2</sub> Level 2 data products (Buchwitz et al., 2015, 2017a, 2017b; Reuter et al., 2013) from SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT. To generate the O4Mv3 product, the EMMA version 3.0 (EMMAv3, Reuter et al., 2017e) product was used. The list of satellite products used for the generation of the EMMAv3 Level 2 product - and therefore also for the O4Mv3 Level 3 data product used in this study - is provided in Tab. 1. The quality of this product relative to Total Carbon Column Observing Network (TCCON) ground-based observations (Wunch et al., 2011, 2015) can be summarized as follows (Buchwitz et al., 2017c): +0.23 ppm overall (global) bias, relative accuracy 0.3 ppm (1-sigma), and very good stability in terms of linear bias trend (-0.02±0.04 ppm/year).

Figure 1 presents an overview of the O4Mv3 product in terms of time series and global XCO<sub>2</sub> maps. The maps show the typical coverage of XCO<sub>2</sub> from SCIAMACHY (until April 2012) and GOSAT (since mid 2009). As can be seen, the time series for the three latitude bands shown in Fig. 1 have very similar slopes. They mainly differ in the amplitude of the seasonal cycle, which reflects the latitudinal dependence of uptake and release of atmospheric CO<sub>2</sub> by the terrestrial biosphere (Schneising et al., 2014). These time series have been used to compute annual mean CO<sub>2</sub> growth rates as will be explained in the following section.

### 3 Atmospheric CO<sub>2</sub> growth rates from satellite observations

National Oceanic and Atmospheric Administration (NOAA) defines the annual mean CO<sub>2</sub> growth rate for a given year as the CO<sub>2</sub> concentration difference at the end of that year minus the CO<sub>2</sub> concentration at the beginning of that year (Thoning et al., 1989; see also additional explanations as given on the NOAA/ESRL website ([https://www.esrl.noaa.gov/gmd/ccgg/about/global\\_means.html](https://www.esrl.noaa.gov/gmd/ccgg/about/global_means.html))). As described below, our method involves the following three steps: (i) Computation of an XCO<sub>2</sub> time series (at monthly resolution and sampling) by averaging the XCO<sub>2</sub> in the region of interest (e.g., a latitude band, see Annex A, Fig. A1). (ii) Computation of monthly sampled XCO<sub>2</sub> annual growth rates by computing the difference of the XCO<sub>2</sub> value of month *i* minus the XCO<sub>2</sub> value of month *i*-12 and computation of the corresponding uncertainty estimate. (iii) Computation of annual mean growth rates and their corresponding uncertainties from the monthly sampled annual growth rates.

In the following, this method is described in detail using Fig. 2 for illustration. In Figure 2a monthly satellite XCO<sub>2</sub> (O4Mv3), as obtained by globally averaging all the individual (5°x5°) XCO<sub>2</sub> values, is plotted. To compute the spatially averaged XCO<sub>2</sub> time series (shown in Fig. 2a), we first longitudinally average the XCO<sub>2</sub> followed by the computation of the area-weighted latitudinal average of XCO<sub>2</sub> by using the cosine of latitude as weight. We consider surface area because surface fluxes are linked to mass of CO<sub>2</sub> (or number of CO<sub>2</sub> molecules) rather than molecular mixing ratios or mole fractions. As can be seen, the computed time series does not start at the beginning of 2003 but in April 2003. As explained in Buchwitz et al., 2017d (see discussion of their Fig. 6.1.1.1) the underlying SCIAMACHY BESD v02.01.02 XCO<sub>2</sub> data product (see Tab. 1) apparently suffers from an approximately 1 ppm high bias in the first few months of 2003. The exact magnitude of this bias has not been quantified due to lack of TCCON validation data in this early time period. As this bias in early 2003 is critical for the year 2003 growth rate, we have omitted the first three months of 2003 for the computation of the growth rates shown in this publication.

Figure 2b shows monthly sampled annual growth rates as computed from the monthly XCO<sub>2</sub> values shown in Fig. 2a. Each value is the difference of two monthly XCO<sub>2</sub> values corresponding to the same month (e.g., January) but different years (e.g., 2004 and 2005). For example, the first data point (first diamond symbol) shown in Fig. 2b is the difference of the April 2004 XCO<sub>2</sub> value minus the April 2003 XCO<sub>2</sub> value. The second data point corresponds to May 2004 minus May 2003, etc. The time difference between the monthly XCO<sub>2</sub> pairs is always one year and the time assigned to each XCO<sub>2</sub> difference is the time in the middle of that year. Therefore, the time series shown in Fig. 2b starts six months later and ends six months earlier as compared to the time series shown in Fig. 2a. Each XCO<sub>2</sub> difference shown in Fig. 2b therefore corresponds to an estimate of the XCO<sub>2</sub> annual growth rate and the position on the time-axis corresponds to the middle of the corresponding one-year time period.

A 1-sigma uncertainty estimate has been computed for each of the monthly sampled annual growth rates shown in Fig. 2b (see grey vertical bars). They have been computed such that they reflect the following aspects: (i) the standard error of the O4Mv3 XCO<sub>2</sub> values as given in the O4Mv3 data product file for each of the 5°x5° grid cells, (ii) the spatial variability of the XCO<sub>2</sub> within the selected region, (iii) the temporal variability of the annual growth rates in the one year time interval, which corresponds to the annual growth rate, and (iv) the number of months (N) with data located in that one year time interval. The uncertainties have been computed as the mean value of three terms divided by the square root of N. The first term is the mean value of the standard error, the second term is the standard deviation of the XCO<sub>2</sub> values in the selected region and the third term is the standard deviation of the monthly sampled annual growth rates in the corresponding one-year time interval. We aimed at providing realistic error estimates but we acknowledge that our uncertainty estimates are not based on full error propagation, which would be difficult especially due to unknown or not well enough known systematic errors and error correlations. The reported uncertainty estimates should therefore be interpreted as error indications rather than fully rigorous error estimates.

Figure 2c shows the final result, i.e., the annual mean XCO<sub>2</sub> growth rates and their estimated (1-sigma) uncertainties. The annual mean growth rates have been computed by averaging all the monthly sampled annual growth rates (shown in Fig. 2b), which are located in the year of interest (e.g., 2003). For most years, 12 annual

growth rate values are available for averaging but there are some exceptions. For example, for the year 2003 only 3 values are present as can be seen from Fig. 2b and for the years 2014 and 2015 there are only 11 values as no data are available for January 2015 due to issues with the GOSAT satellite. The uncertainty of the annual mean growth rate has been computed by averaging the uncertainties assigned to each of the monthly sampled annual growth rates (shown as grey vertical bars in Fig. 2b) scaled with a factor, which depends on the number of months (N) available for averaging. This factor is the square root of 12/N. It ensures that the uncertainty is larger, the less data points are available for averaging. Overall, our uncertainty estimate is quite conservative, as we do not assume that errors improve upon averaging. As a result of this procedure, the error bar of the year 2003 growth rate is quite large (0.76 ppm/year, see Tab. A1 in Annex A, where all numerical values are listed). This is because the monthly sampled annual growth rate varies significantly in 2003 (see Fig. 2b) and because only N=3 data points are available for averaging in 2003. In contrast, the year 2005 growth rate uncertainty is much smaller (0.28 ppm/year) because the growth rates vary less during 2005 and because N=12 data points are available for averaging.

In Fig. 2c also the NOAA global growth rates (Dlugokencky and Tans, 2017b) are shown. As can be seen, the satellite-derived growth rates agree well with the NOAA growth rates obtained from CO<sub>2</sub> surface observations. For the time period 2003-2016 the linear correlation coefficient R is 0.82 and the difference is  $-0.02 \pm 0.28$  ppm/year (mean difference  $\pm$  standard deviation). Perfect agreement is not to be expected as these two growth rate time series have been obtained from CO<sub>2</sub> observations, which represent very different vertical sampling of the atmosphere (surface (NOAA) versus entire vertical column (satellite)) (see Fig. A3b in Annex A for a comparison of XCO<sub>2</sub> and surface CO<sub>2</sub> growth rates obtained using a global re-analysis CO<sub>2</sub> data product). Perfect agreement is also not to be expected because we use different time periods for the computation of the annual growth rates compared to NOAA (see Fig. A3c in Annex A for a comparison of two different methods to compute annual XCO<sub>2</sub> growth rates).

As can also be seen from Fig. 2c, the largest growth rates are approximately 3 ppm/year during 2015 and 2016. These record large growth rates (Peters et al., 2017) are attributed to the consequences of the strong 2015/2016 El Niño event, which produced large CO<sub>2</sub> emissions from fires and enhanced net biospheric respiration in the tropics relative to normal conditions (Heymann et al., 2017; Liu et al., 2017). Many of these fires are initiated by humans, for example, to clear tropical forests. In this study, human emissions of CO<sub>2</sub> are defined as emissions from fossil fuel combustion and industry (Le Quéré et al., 2016, 2018) but do not include, for example, CO<sub>2</sub> emissions originating from slash and burn agriculture.

#### **4 Correlation of CO<sub>2</sub> growth rates with fossil CO<sub>2</sub> emissions and ENSO indices**

It is well known that changes of the growth rate of atmospheric CO<sub>2</sub> have anthropogenic and natural causes (e.g., Jones et al., 2001; Betts et al., 2016; Kim et al., 2016; Liu et al., 2017; Chylek et al., 2018). In this section we are aiming at answering the following question: “Assuming that the variability of the CO<sub>2</sub> growth rate is dominated by ENSO and by human emissions, which of the two considered causes dominates the growth rate variability given the satellite-derived growth rates and their uncertainty?”. To answer this question we are using a simple

linear statistical model and time series of human emissions and two ENSO indices assuming that these indices are appropriate proxies for ENSO related effects in the context of providing a reliable answer.

Figure 3 shows a comparison of the CO<sub>2</sub> annual mean growth rates (Fig. 3a) with annual global CO<sub>2</sub> emissions from fossil fuel combustion and industry (Fig. 3b) (Le Quéré et al., 2018; GCP 2017) (correlation of growth rate and human emissions:  $R^2 = 31\%$ ). As can be seen, the growth rates vary significantly in recent years despite nearly constant human emissions. Figure 3d shows two ENSO indices: the Southern Oscillation Index (SOI, blue lines) (NOAA 2017a; Ropelewski and Jones, 1987) and the Oceanic Niño Index (ONI, green lines) (NOAA 2017b). Whereas SOI is defined as the normalized pressure difference between Tahiti and Darwin (values less than -1 indicate the presence of a strong El Niño), ONI is based on Sea Surface Temperature (SST) differences (positive values correspond to El Niño). The dotted lines correspond to the original (i.e., unshifted) annual mean indices and the solid lines correspond to time shifted ENSO indices. Time shifts have been investigated to consider the delay in atmospheric response to ENSO-induced changes. As shown in Fig. 3c, the growth rate response as quantified by  $R^2$  is largest after 4 months for ONI ( $R^2 = 35\%$ ) and after 7 months for SOI ( $R^2 = 30\%$ ). These maxima have been adopted for the solid (shifted) lines in Fig 3d. This finding is consistent with results from other studies, where lags in the range 3-9 months have been reported (Jones et al., 2001; Kim et al., 2016; Chylek et al., 2018).

Figure 3 shows that the anthropogenic emission variability is mostly linked to a trend whereas the El Niño signal is variable on much shorter time scales. Thus, the relative impact of the anthropogenic and natural contributions depends on the length of the time series. The shorter the time series, the smaller the anthropogenic variability is. It is therefore expected that the natural contribution to the variability of the growth rate gets larger for a shorter time series. In order to separate and quantify the contributions of the human CO<sub>2</sub> emissions and of ENSO, as described by the two indices SOI and ONI, to the growth rate variations, we employ the method of “variation partitioning” (Peres-Neto et al., 2006). To achieve this, we have fitted three basis functions to the 2003-2016 growth rate time series via linear least-squares minimization (we explain the method in this paragraph using SOI but the method does not depend on which ENSO index is used): (i) a constant offset (variance zero), (ii) the human CO<sub>2</sub> emissions (Fig. 3b) and (iii) SOI shifted by 7 months (blue solid line in Fig. 3d). The variance of the scaled emission, i.e., of the human emission scaled with the corresponding fit parameter, is 0.0758 ppm<sup>2</sup>/year<sup>2</sup> (note that in this section we report numerical values with four digital places but this shall not imply that all decimal places are significant). The variance of the scaled SOI is 0.1070 ppm<sup>2</sup>/year<sup>2</sup> and the variance of the fit residual is 0.0728 ppm<sup>2</sup>/year<sup>2</sup>. The sum of the three individual variances is 0.2557 ppm<sup>2</sup>/year<sup>2</sup> whereas the variance of the annual mean growth rate is 0.2307 ppm<sup>2</sup>/year<sup>2</sup>. This shows that the sum of the variances is 10.8% larger than the variance of the growth rate, i.e., the sum of the variances is not exactly equal to the variance of the sum. The reason for this is that the CO<sub>2</sub> emission and the SOI time series are not uncorrelated ( $R = 0.14$ ). To account for correlations, we subtract the variance of the residual from the variance of the growth rate. The result is the part of the variance to be explained by the emissions and by the SOI. The ratio of this to be explained variance (0.1579 ppm<sup>2</sup>/year<sup>2</sup>) and the sum of the variances of the emissions and SOI ((0.0758 + 0.1070) ppm<sup>2</sup>/year<sup>2</sup> = 0.1828 ppm<sup>2</sup>/year<sup>2</sup>) is 0.8638. The latter is then used as a scaling factor applied to the variances of the emissions and of the SOI. The scaled variances are 0.0655 ppm<sup>2</sup>/year<sup>2</sup> for the emissions and 0.0924 ppm<sup>2</sup>/year<sup>2</sup> for SOI (note that the sum of these scaled variances and the variance of the residual is equal to the

variance of the growth rate). From this we conclude that the human emissions explain 28% ( $= 0.0655/0.2307$ ) of the variance of the growth rate and that ENSO as quantified by the SOI explains 40% ( $= 0.0924/0.2307$ ). We computed (1-sigma) uncertainties of these estimates by numerically perturbing the satellite-derived annual mean growth rates taking into account their uncertainty (see Figs. 2c and 3) and by subsequently repeating the  
5 computations as explained above 10,000 times. The perturbations correspond to random perturbations of the annual mean growth rates assuming normal distributions for each year and no correlation between the different years. This analysis yields that  $40\pm 13\%$  of the growth rate variation results from the impact of ENSO and that  $28\pm 14\%$  is due to the human emissions of CO<sub>2</sub>. Using these simulations, we also computed the fraction of cases where the ENSO impact dominates over the human emissions. This fraction is 63% in this case, i.e., when using  
10 SOI and when the analysis is applied to the entire time period 2003-2016. This fraction is interpreted as the probability that ENSO-induced impacts on the variation of the growth rate dominates that of human emissions.

When using ONI instead of SOI, ENSO explains  $37\pm 14\%$  of the growth rate variance during 2003-2016, human emissions explain  $24\pm 14\%$  and the fraction where ENSO dominates is again 63%. When restricting the time period to 2010-2016, which is dominated by strong 2010/2012 La Niña events (Boening et al., 2012; Rodrigues et al., 2014) and by the strong 2015/2016 El Niño, the results are the following: Using the SOI analysis, we find that  
15 ENSO explains  $58\pm 19\%$  of the variance, human emissions explain  $2\pm 9\%$  and the probability that ENSO dominates is 94%. For the ONI analysis, we find that ENSO explains  $59\pm 20\%$  of the variance, human emissions explain  $3\pm 9\%$  and the probability that ENSO dominates is 94%. This analysis shows that the ENSO impact on CO<sub>2</sub> growth rate variations dominates over that of human emissions throughout the period 2003-2016 but in  
20 particular in the second half of this period, i.e., during 2010-2016.

## 5 Conclusions

We presented a method for the computation of atmospheric CO<sub>2</sub> column annual mean growth rates from satellite  
25 XCO<sub>2</sub> retrievals. The satellite XCO<sub>2</sub> data product used is the Obs4MIPs version 3 (O4Mv3) XCO<sub>2</sub> data product based on SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT satellite data. This product covers the time period 2003-2016 and has monthly time and  $5^\circ \times 5^\circ$  spatial resolution.

The estimated uncertainty of the satellite-derived annual mean growth rates is typically 0.3 ppm/year (1-sigma) with the exception of the first year 2003, where the uncertainty is 0.76 ppm/year, and of the last year 2016, where  
30 the uncertainty is 0.50 ppm/year. The growth rates agree with NOAA within the uncertainty of the satellite-derived growth rates (mean difference  $\pm$  standard deviation:  $0.0\pm 0.3$  ppm/year; R: 0.82). In agreement with NOAA, we find that the growth rates are largest in the years 2015 and 2016. These growth rates are around 3 ppm/year and are attributed to the 2015/2016 El Niño resulting in large CO<sub>2</sub> emissions from fires and enhanced net biospheric respiration in the tropics relative to normal conditions (Heymann et al., 2017; Liu et al., 2017). Our  
35 analysis also shows that the ENSO impact on CO<sub>2</sub> growth rate variations dominates over that of human emissions throughout the period 2003-2016 (14 years) but in particular during the period 2010-2016 (second half of the investigated time period) due to strong La Niña and El Niño events. We estimate the probability that the impact

of ENSO on the variability is larger than the impact of human emissions to be 63% for the time period 2003-2016. If the time period is restricted to 2010-2016 this probability increases to 94%.

In the future, we plan to regularly update the satellite-derived XCO<sub>2</sub> growth rates to monitor this important quantity. This will also include satellite XCO<sub>2</sub> retrievals from other satellite instruments such as XCO<sub>2</sub> from  
5 NASA's OCO-2 mission (e.g., Eldering et al., 2017; Reuter et al., 2017c, 2017d).

## References

- Ballantyne, A. P., Alden, C. B., Miller, J. B., Tans, P. P., and White, J. W. C.: Increase in observed net carbon dioxide uptake by land and oceans during the last 50 years, *Nature* 488, 70–72, 2012.
- 10 Basu, S., Guerlet, S., Butz, A., Houweling, S., Hasekamp, O., Aben, I., Krummel, P., Steele, P., Langenfelds, R., Torn, M., Biraud, S., Stephens, B., Andrews, A., and Worthy, D.: Global CO<sub>2</sub> fluxes estimated from GOSAT retrievals of total column CO<sub>2</sub>, *Atmos. Chem. Phys.*, 13, 8695–8717, doi:10.5194/acp-13-8695-2013, 2013.
- Betts, R. A., Jones, C. D., Knight, J. R., Keeling, R. F., and Kennedy, J. J., El Niño and a record CO<sub>2</sub> rise, *Nature Climate Change*, vol. 6, p. 806–810, <https://www.nature.com/articles/nclimate3063.pdf>, 2016.
- 15 Boening, C., Willis, J. K., Landerer, F. W., Nerem, R. S., and J. Fasullo, J.: The 2011 La Niña: So strong, the oceans fell, *Geophys. Res. Lett.*, 38, pp. 5, doi:10.1029/2012GL053055, 2012.
- Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noël, S., Rozanov, V. V., Chance, K. V., and Goede, A. H. P.: SCIAMACHY - Mission objectives and measurement modes, *J. Atmos. Sci.*, 56 (2), 127-150, 1999.
- Buchwitz, M., de Beek, R., Noël, S., Burrows, J. P., Bovensmann, H., Schneising, O., Khlystova, I., Bruns, M.,  
20 Bremer, H., Bergamaschi, P., Körner, S., and Heimann, M.: Atmospheric carbon gases retrieved from SCIAMACHY by WFM-DOAS: version 0.5 CO and CH<sub>4</sub> and impact of calibration improvements on CO<sub>2</sub> retrieval, *Atmos. Chem. Phys.*, 6, 2727–2751, 2006.
- Buchwitz, M., Schneising, O., Burrows, J. P., Bovensmann, H., Reuter, M., and Notholt, J.: First direct observation of the atmospheric CO<sub>2</sub> year-to-year increase from space, *Atmos. Chem. Phys.*, 7, 4249-4256, 2007.
- 25 Buchwitz, M., Reuter, M., Schneising, O., Boesch, H., Guerlet, S., Dils, B., Aben, I., Armante, R., Bergamaschi, P., Blumenstock, T., Bovensmann, H., Brunner, D., Buchmann, B., Burrows, J. P., Butz, A., Chédin, A., Chevallier, F., Crevoisier, C. D., Deutscher, N. M., Frankenberg, C., Hase, F., Hasekamp, O. P., Heymann, J., Kaminski, T., Laeng, A., Lichtenberg, G., De Mazière, M., Noël, S., Notholt, J., Orphal, J., Popp, C., Parker, R., Scholze, M., Sussmann, R., Stiller, G. P., Warneke, T., Zehner, C., Bril, A., Crisp, D., Griffith, D. W. T., Kuze,  
30 A., O'Dell, C., Oshchepkov, S., Sherlock, V., Suto, H., Wennberg, P., Wunch, D., Yokota, T., and Yoshida, Y.: The Greenhouse Gas Climate Change Initiative (GHG-CCI): comparison and quality assessment of near-surface-sensitive satellite-derived CO<sub>2</sub> and CH<sub>4</sub> global data sets, *Remote Sensing of Environment*, 162, 344-362, doi:10.1016/j.rse.2013.04.024, 2015.
- Buchwitz, M., Reuter, M., Schneising-Weigel, O., Aben, I., Detmers, R. G., Hasekamp, O. P., Boesch, H.,  
35 Anand, J., Crevoisier, C., and Armante, R.: Product User Guide and Specification (PUGS) – Main document, Technical Report Copernicus Climate Change Service (C3S), available from C3S website



- (<https://climate.copernicus.eu/>) and from [http://www.iup.uni-bremen.de/sciamachy/NIR\\_NADIR\\_WFM\\_DOAS/C3S\\_docs/C3S\\_D312a\\_Lot6.3.1.5-v1\\_PUGS\\_MAIN\\_v1.3.pdf](http://www.iup.uni-bremen.de/sciamachy/NIR_NADIR_WFM_DOAS/C3S_docs/C3S_D312a_Lot6.3.1.5-v1_PUGS_MAIN_v1.3.pdf), 20/10/2017, pp. 91, 2017a.
- Buchwitz, M., Reuter, M., Schneising, O., Hewson, W., Detmers, R. G., Boesch, H., Hasekamp, O. P., Aben, I., Bovensmann, H., Burrows, J. P., Butz, A., Chevallier, F., Dils, B., Frankenberg, C., Heymann, J., Lichtenberg, G., De Mazière, M., Notholt, J., Parker, R., Warneke, T., Zehner, C., Griffith, D. W. T., Deutscher, N. M., Kuze, A., Suto, H., and Wunch, D.: Global satellite observations of column-averaged carbon dioxide and methane: The GHG-CCI XCO<sub>2</sub> and XCH<sub>4</sub> CRDP3 data set, *Remote Sensing of Environment*, 203, 276-295, <http://dx.doi.org/10.1016/j.rse.2016.12.027>, 2017b.
- 10 Buchwitz, M., Reuter, M., Schneising-Weigel, O., Aben, I., Detmers, R. G., Hasekamp, O. P., Boesch, H., Anand, J., Crevoisier, C., and Armante, R.: Product Quality Assessment Report (PQAR) – Main document, Technical Report Copernicus Climate Change Service (C3S), available from C3S website (<https://climate.copernicus.eu/>) and from [http://www.iup.uni-bremen.de/sciamachy/NIR\\_NADIR\\_WFM\\_DOAS/C3S\\_docs/C3S\\_D312a\\_Lot6.3.1.7-v1\\_PQAR\\_MAIN\\_v1.1.pdf](http://www.iup.uni-bremen.de/sciamachy/NIR_NADIR_WFM_DOAS/C3S_docs/C3S_D312a_Lot6.3.1.7-v1_PQAR_MAIN_v1.1.pdf), 20/10/2017, pp. 103, 2017c.
- 15 Buchwitz, M., Dils, B., Boesch, H., Brunner, D., Butz, A., Crevoisier, C., Detmers, R., Frankenberg, C., Hasekamp, O., Hewson, W., Laeng, A., Noël, S., Notholt, J., Parker, R., Reuter, M., Schneising, O., Somkuti, P., Sundström, A.-M., De Wachter, E.: ESA Climate Change Initiative (CCI) Product Validation and Intercomparison Report (PVIR) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG) for data set Climate Research Data Package No. 4 (CRDP#4), Technical Report, version 5, 9-Feb-2017, link: [http://www.esa-ghg-cci.org/?q=webfm\\_send/352](http://www.esa-ghg-cci.org/?q=webfm_send/352), pp. 253, 2017d.
- Burrows, J. P., Hölzle, E., Goede, A. P. H., Visser, H., and Fricke, W.: SCIAMACHY—Scanning Imaging Absorption Spectrometer for Atmospheric Chartography, *Acta Astronaut.*, 35(7), 445–451, doi:10.1016/0094-5765(94)00278-t, 1995.
- 25 Butz, A., Guerlet, S., Hasekamp, O., Schepers, D., Galli, A., Aben, I., Frankenberg, C., Hartmann, J.-M., Tran, H., Kuze, A., Keppel-Aleks, G., Toon, G., Wunch, D., Wennberg, P., Deutscher, N., Griffith, D., Macatangay, R., Messerschmidt, J., Notholt, J., and Warneke, T.: Toward accurate CO<sub>2</sub> and CH<sub>4</sub> observations from GOSAT, *Geophys. Res. Lett.*, doi:10.1029/2011GL047888, 2011.
- Chevallier, F., Palmer, P. I., Feng, L., Boesch, H., O'Dell, C. W., and Bousquet, P.: Towards robust and consistent regional CO<sub>2</sub> flux estimates from in situ and space-borne measurements of atmospheric CO<sub>2</sub>, *Geophys. Res. Lett.*, 41, 1065–1070, doi:10.1002/2013GL058772, 2014.
- 30 Chevallier, F., Validation report for the inverted CO<sub>2</sub> fluxes, v17r1, Technical Report Copernicus Atmosphere Monitoring Service (CAMS), version 1.0 (06/07/2018), available from CAMS website ([https://atmosphere.copernicus.eu/sites/default/files/2018-10/CAMS73\\_2015SC3\\_D73.1.4.2-1979-2017-v1\\_201807\\_v1-1.pdf](https://atmosphere.copernicus.eu/sites/default/files/2018-10/CAMS73_2015SC3_D73.1.4.2-1979-2017-v1_201807_v1-1.pdf)), 2018.
- 35 Chylek, P., Tans, P., Christy, P., and Dubey, M. K.: The carbon cycle response to two El Niño types: an observational study, *Environ. Res. Lett.*, 13, pp. 8, <https://doi.org/10.1088/1748-9326/aa9c5b>, 2018.

- Cogan, A. J., Boesch, H., Parker, R. J., Feng, L., Palmer, P. I., Blavier, J.-F. L., Deutscher, N. M., Macatangay, R., Notholt, J., Roehl, C., Warneke, T., and Wunsch, D.: Atmospheric carbon dioxide retrieved from the Greenhouse gases Observing SATellite (GOSAT): Comparison with ground-based TCCON observations and GEOS-Chem model calculations, *J. Geophys. Res.*, 117, D21301, doi:10.1029/2012JD018087, 2012.
- 5 Dlugokencky, E., and Tans, P.: Trends in atmospheric carbon dioxide. National Oceanic & Atmospheric Administration, Earth System Research Laboratory (NOAA/ESRL), 2017a.
- Dlugokencky, E., and Tans, P.: Trends in atmospheric carbon dioxide, National Oceanic & Atmospheric Administration, Earth System Research Laboratory (NOAA/ESRL), file: [ftp://aftp.cmdl.noaa.gov/products/trends/co2/co2\\_gr\\_gl.txt](ftp://aftp.cmdl.noaa.gov/products/trends/co2/co2_gr_gl.txt) (access: 24 November 2017), 2017b.
- 10 Eldering, A. et al. The Orbiting Carbon Observatory-2: first 18 months of science data products, *Atmos. Meas. Tech.*, 10, 549-563, <https://doi.org/10.5194/amt-10-549-2017>, 2017.
- GCP: Global CO<sub>2</sub> emissions from Global Carbon Project (GCP) website (<http://www.globalcarbonproject.org/carbonbudget/17/data.htm>), main data source: Thomas Boden, Gregg Marland, Robert Andres 2017 (CDIAC), GCP-CICERO, UNFCCC, BP Statistical Review of World Energy (BP, 15 2017), US Geological Survey (USGS, 2017), 2017.
- Hakkarainen, J., Ialongo, I., and Tamminen, J.: Direct space-based observations of anthropogenic CO<sub>2</sub> emission areas from OCO-2, *Geophys. Res. Lett.*, 43, 11,400–11,406, doi:10.1002/2016GL070885, 2016.
- Heymann, J., Reuter, M., Buchwitz, M., Schneising, O., Bovensmann, H., Burrows, J. P., Massart, S., Kaiser, J. W., and Crisp, D.: CO<sub>2</sub> emission of Indonesian fires in 2015 estimated from satellite-derived atmospheric CO<sub>2</sub> 20 concentrations, *Geophys. Res. Lett.*, doi:10.1002/2016GL072042, pp. 18, 2017.
- Houweling, S., Baker, D., Basu, S., Boesch, H., Butz, A., Chevallier, F., Deng, F., Dlugokencky, E. J., Feng, L., Ganshin, A., Hasekamp, O., Jones, D., Maksyutov, S., Marshall, J., Oda, T., O'Dell, C. W., Oshchepkov, S., Palmer, P. I., Peylin, P., Poussi, Z., Reum, F., Takagi, H., Yoshida, Y., and Zhuralev, R.: An intercomparison of inverse models for estimating sources and sinks of CO<sub>2</sub> using GOSAT measurements, *J. Geophys. Res. Atmos.*, 25 120, 5253–5266, doi:10.1002/2014JD022962, 2015.
- Huntzinger, D. N., Michalak, A. M., Schwalm, C., Ciais, P., King, A. W., Fang, Y., Schaefer, K., Wei, Y., Cook, R. B., Fisher, J. B., Hayes, D., Huang, M., Ito, A., Jain, A. K., Lei, H., Lu, C., Maignan, F., Mao, J., Parazoo, N., Peng, S., Poulter, B., Ricciuto, D., Shi, X., Tian, H., Wang, W., Zeng, N., and Zhao, F.: Uncertainty in the response of terrestrial carbon sink to environmental drivers undermines carbon-climate feedback predictions, 30 *Nature, Scientific Reports*, 7: 4765, doi:10.1038/s41598-017-03818-2, 2017.
- IPCC: Climate Change 2013: The Physical Science Basis, Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Report on Climate Change, <http://www.ipcc.ch/report/ar5/wg1/>, Cambridge University Press, 2013.
- Jones, C. D., Collins, M., Cox, P. M., and Spall, S. A.: The Carbon Cycle Response to ENSO: A Coupled 35 Climate-Carbon Cycle Model Study, *Journal of Climate*, Vol. 14, 4113-4129, 2001.

- Kaminski, T., Scholze, M., Voßbeck, M., Knorr, W., Buchwitz, M., and Reuter, M.: Constraining a terrestrial biosphere model with remotely sensed atmospheric carbon dioxide, *Remote Sensing of Environment* 203, 109-124, 2017.
- Kim, J.-S., Kug, J.-S., Yoon, J.-H., Jeong, S.-J., Increased Atmospheric CO<sub>2</sub> Growth Rate during El Niño Driven by Reduced Terrestrial Productivity in the CMIP5 ESMs, *Journal of Climate*, 8783-8805, 29. 10.1175/JCLI-D-14-00672.1, 2016.
- Kort, E., Frankenberg, C., Miller, C. E., and Oda, T.: Space-based observations of megacity carbon dioxide. *Geophys. Res. Lett.*, 39, doi:10.1029/2012GL052738, 2012.
- Kuze, A., Suto, H., Shiomi, K., Kawakami, S., Tanaka, M., Ueda, Y., Deguchi, A., Yoshida, J., Yamamoto, Y., Kataoka, F., Taylor, T. E., and Buijs, H. L.: Update on GOSAT TANSO-FTS performance, operations, and data products after more than 6 years in space, *Atmos. Meas. Tech.*, 9, 2445-2461, doi:10.5194/amt-9-2445-2016, 2016.
- Lauer, A., Eyring, V., Righi, M., Buchwitz, M., Defourny, P., Evaldsson, M., Friedlingstein, P., de Jeu, R., de Leeuw, G., Loew, A., Merchant, C. J., Müller, B., Popp, T., Reuter, M., Sandven, S., Senfleben, D., Stengel, M., Van Roozendaal, M., Wenzel, S., and Willén, U.: Benchmarking CMIP5 models with a subset of ESA CCI Phase 2 data using the ESMValTool, *Remote Sensing of Environment*, 203, 9-39, <http://dx.doi.org/10.1016/j.rse.2017.01.007>, 2017.
- Le Quéré, C., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Peters, G. P., Manning, A. C., Boden, T. A., Tans, P. P., Houghton, R. A., Keeling, R. F., Alin, S., Andrews, O. D., Anthoni, P., Barbero, L., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Currie, K., Delire, C., Doney, S. C., Friedlingstein, P., Gkritzalis, T., Harris, I., Hauck, J., Haverd, V., Hoppema, M., Klein Goldewijk, K., Jain, A. K., Kato, E., Körtzinger, A., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Lombardozzi, D., Melton, J. R., Metzl, N., Millero, F., Monteiro, P. M. S., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S. I., O'Brien, K., Olsen, A., Omar, A. M., Ono, T., Pierrot, D., Poulter, B., Rödenbeck, C., Salisbury, J., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I., Stocker, B. D., Sutton, A. J., Takahashi, T., Tian, H., Tilbrook, B., van der Laan-Luijkx, I. T., van der Werf, G. R., Viovy, N., Walker, A. P., Wiltshire, A. J., and Zaehle, S.: Global Carbon Budget 2016, *Earth Syst. Sci. Data*, 8, 605-649, 2016.
- Le Quéré, C., Andrew, R. M., Friedlingstein, P., Sitch, S., Pongratz, J., Manning, A. C., Korsbakken, J. I., Peters, G. P., Canadell, J. G., Jackson, R. B., Boden, T. A., Tans, P. P., Andrews, O. D., Arora, V. K., Bakker, D. D. E., Barbero, L., Becker, M., Betts, R. A., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Cosca, C. E., Cross, J., Currie, K., Gasser, T., Harris, I., Hauck, J., Haverd, V., Houghton, R. A., Hunt, C. W., Hurtt, G., Ilyina, T., Jain, A. K., Kato, E., Kautz, M., Keeling, R. F., Goldewijk, K. K., Körtzinger, A., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Lima, I., Lombardozzi, D., Metzl, N., Millero, F., Monteiro, P. M. S., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S., Nojiri, Y., Padín, X. A., Pregon, A., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G., Reimer, J., Rödenbeck, C., Schwinger, J., Séférian, R., Skjelvan, I., Stocker, B. D., Tian, H., Tilbrook, B., van der Laan-Luijkx, I. T., van der Werf, G. R., van Heuven, S., Viovy, N., Vuichard, N., Walker, A. P., Watson, A.

- J., Wiltshire, A. J., Zaehle, S., and Zhu, D.: Global Carbon Budget 2017, *Earth System Science Data*, 10, 405-448, DOI: 10.5194/essd-10-405-2018, 2018.
- Liu, J., Bowman, K. W., Schimel, D. S., Parazoo, N. C., Jiang, Z., Lee, M., Bloom, A. A., Wunch, D., Frankenberg, C., Sun, Y., O'Dell, C. W., Gurney, K. R., Menemenlis, D., Gierach, M., Crisp, D., and Eldering, A.: Contrasting carbon cycle responses of the tropical continents to the 2015–2016 El Niño, *Science*, 358, eaam5690, pp. 7, 2017.
- Maksyutov, S., Takagi, H., Valsala, V. K., Saito, M., Oda, T., Saeki, T., Belikov, D. A., Saito, R., Ito, A., Yoshida, Y., Morino, I., Uchino, O., Andres, R. J., and Yokota, T.: Regional CO<sub>2</sub> flux estimates for 2009–2010 based on GOSAT and ground-based CO<sub>2</sub> observations, *Atmos. Chem. Phys.*, 13, 9351-9373, 2013.
- 10 Nassar, R., Hill, T. G., McLinden, C. A., Wunch, D., Jones, D. B. A., and Crisp, D.: Quantifying CO<sub>2</sub> emissions from individual power plants from space, *Geophys. Res. Lett.*, doi:10.1002/2017GL074702, 2017.
- NIES: Recent global CO<sub>2</sub> (<http://www.gosat.nies.go.jp/en/recent-global-co2.html>), 2017.
- NOAA: Southern Oscillation Index (SOI), file [https://www.esrl.noaa.gov/psd/gcos\\_wgsp/Timeseries/Data/soi.long.data](https://www.esrl.noaa.gov/psd/gcos_wgsp/Timeseries/Data/soi.long.data), data based on the method of
- 15 Ropelewski and Jones, 1987, using data from the Climate Research Unit (CRU); <http://www.cru.uea.ac.uk/cru/data/soi/>), 2017a.
- NOAA: Oceanic Niño Index (ONI), file <https://www.esrl.noaa.gov/psd/data/correlation/oni.data> obtained from <https://www.esrl.noaa.gov/psd/data/climateindices/list>, 2017b.
- O'Dell, C. W., Connor, B., Bösch, H., O'Brien, D., Frankenberg, C., Castano, R., Christi, M., Eldering, D.,
- 20 Fisher, B., Gunson, M., McDuffie, J., Miller, C. E., Natraj, V., Oyafuso, F., Polonsky, I., Smyth, M., Taylor, T., Toon, G. C., Wennberg, P. O., and Wunch, D.: The ACOS CO<sub>2</sub> retrieval algorithm – Part 1: The ACOS CO<sub>2</sub> retrieval algorithm – Part 1: Description and validation against synthetic observations, *Atmos. Meas. Tech.*, 5, 99–121, doi:10.5194/amt-5-99-2012, 2012.
- Parker, R. J., Boesch, H., Wooster, M. J., Moore, D. P., Webb, A. J., Gaveau, D., and Murdiyarso, D.:
- 25 Atmospheric CH<sub>4</sub> and CO<sub>2</sub> enhancements and biomass burning emission ratios derived from satellite observations of the 2015 Indonesian fire plumes, *Atmos. Chem. Phys.*, 16, 10111-10131, doi:10.5194/acp-16-10111-2016, 2016.
- Peres-Neto, P. R., Legendre, P., Dray, S., and Borcard, D.: Variation partitioning of species data matrices: Estimation and comparison of fractions, *Ecology*, 87(19), pp. 2614-2625, 2006.
- 30 Peters, W., Jacobson, A. R., Sweeney, C., Andrews, A. E., Conway, T. J., Masarie, K., Miller, J. B., Bruhwiler, L. M. P., Pétron, G., Hirsch, A. I., Worthy, D. E. J., van der Werf, G. R., Randerson, J. T., Wennberg, P. O., Krol, M. C., and Tans, P. P.: An atmospheric perspective on North American carbon dioxide exchange: CarbonTracker, *Proc. Natl. Acad. Sci. USA*, 104, 18925–18930, doi:10.1073/pnas.0708986104, 2007.
- Peters, G. P., Le Quéré, C., Andrew, R. M., Canadell, J. G., Friedlingstein, P., Ilyina, T., Jackson, R., Joos, F.,
- 35 Korsbakken, J. I., McKinley, G. A., Sitch, S., and Tans, P.: Towards real-time verification of CO<sub>2</sub> emissions, *Nature Climate Change*, 13 November 2017, doi:10.1038/s41558-017-0013-9, 2017.

- Peylin, P., Law, R. M., Gurney, K. R., Chevallier, F., Jacobson, A. R., Maki, T., Niwa, Y., Patra, P. K., Peters, W., Rayner, P. J., Rödenbeck, C., van der Laan-Luijkx, I. T., and Zhang, X.: Global atmospheric carbon budget: results from an ensemble of atmospheric CO<sub>2</sub> inversions, *Biogeosciences*, 10, 6699–6720, doi:10.5194/bg-10-6699-2013, 2013.
- 5 Reuter, M., Bovensmann, H., Buchwitz, M., Burrows, J. P., Connor, B. J., Deutscher, N. M., Griffith, D. W. T., Heymann, J., Keppel-Aleks, G., Messerschmidt, J., Notholt, J., Petri, C., Robinson, J., Schneising, O., Sherlock, V., Velasco, V., Warneke, W., Wennberg, P. O., and Wunch, D.: Retrieval of atmospheric CO<sub>2</sub> with enhanced accuracy and precision from SCIAMACHY: Validation with FTS measurements and comparison with model results, *J. Geophys. Res.*, 116, D04301, doi:10.1029/2010JD015047, 2011.
- 10 Reuter, M., H. Boesch, H. Bovensmann, A. Bril, M. Buchwitz, A. Butz, J. P. Burrows, C. W. O'Dell, S. Guerlet, O. Hasekamp, J. Heymann, N. Kikuchi, S. Oshchepkov, R. Parker, S. Pfeifer, O. Schneising, T. Yokota, and Y. Yoshida: A joint effort to deliver satellite retrieved atmospheric CO<sub>2</sub> concentrations for surface flux inversions: the ensemble median algorithm EMMA, *Atmos. Chem. Phys.*, 13, 1771-1780, 2013.
- Reuter, M., Buchwitz, M., Hilker, M., Heymann, J., Schneising, O., Pillai, D., Bovensmann, H., Burrows, J. P.,  
 15 Bösch, H., Parker, R., Butz, A., Hasekamp, O., O'Dell, C. W., Yoshida, Y., Gerbig, C., Nehrkorn, T., Deutscher, N. M., Warneke, T., Notholt, J., Hase, F., Kivi, R., Sussmann, R., Machida, T., Matsueda, H., and Sawa, Y.: Satellite-inferred European carbon sink larger than expected, *Atmos. Chem. Phys.*, 14, 13739-13753, 2014a.
- Reuter, M., Buchwitz, M., Hilboll, A., Richter, A., Schneising, O., Hilker, M., Heymann, J., Bovensmann, H., and Burrows, J. P.: Decreasing emissions of NO<sub>x</sub> relative to CO<sub>2</sub> in East Asia inferred from satellite  
 20 observations, *Nature Geoscience*, 28 Sept. 2014, doi:10.1038/ngeo2257, pp. 4, 2014b.
- Reuter, M., Buchwitz, M., Schneising, O., Noël, S., Rozanov, V., Bovensmann, H., and Burrows, J. P.: A Fast Atmospheric Trace Gas Retrieval for Hyperspectral Instruments Approximating Multiple Scattering - Part 1: Radiative Transfer and a Potential OCO-2 XCO<sub>2</sub> Retrieval Setup, *Remote Sens.*, 9, 1159, doi:10.3390/rs9111159, 2017c.
- 25 Reuter, M., Buchwitz, M., Schneising, O., Noël, S., Bovensmann, H., and Burrows, J. P.: A Fast Atmospheric Trace Gas Retrieval for Hyperspectral Instruments Approximating Multiple Scattering - Part 2: Application to XCO<sub>2</sub> Retrievals from OCO-2, *Remote Sens.*, 9, 1102, doi:10.3390/rs9111102, 2017d.
- Reuter, M., Buchwitz, M., and Schneising-Weigel, O.: Algorithm Theoretical Basis Document (ATBD) – ANNEX D for products XCO<sub>2</sub>\_EMMA and XCH<sub>4</sub>\_EMMA. Technical Report Copernicus Climate Change  
 30 Service (C3S), available from C3S website (<https://climate.copernicus.eu/>) and from [http://www.iup.uni-bremen.de/sciamachy/NIR\\_NADIR\\_WFM\\_DOAS/C3S\\_docs/C3S\\_D312a\\_Lot6.2.1.2-v1\\_ATBD\\_ANNEX-D\\_v1.1.pdf](http://www.iup.uni-bremen.de/sciamachy/NIR_NADIR_WFM_DOAS/C3S_docs/C3S_D312a_Lot6.2.1.2-v1_ATBD_ANNEX-D_v1.1.pdf), 20/10/2017, pp. 32, 2017e.
- Rodrigues, R. R., and McPhaden, M. J.: Why did the 2011-2012 La Niña cause a severe drought in the Brazilian Northeast? *Geophys. Res. Lett.*, 41, 1012–1018, doi:10.1002/2013GL058703, 2014.
- 35 Ropelewski, C. F., and Jones, P. D.: An extension of the Tahiti-Darwin Southern Oscillation Index. *Monthly Weather Review*, 115, 2161-2165, 1987.

- Schneising, O., Heymann, J., Buchwitz, M., Reuter, M., Bovensmann, H., and Burrows, J. P.: Anthropogenic carbon dioxide source areas observed from space: assessment of regional enhancements and trends, *Atmos. Chem. Phys.*, 13, 2445-2454, doi:10.5194/acp-13-2445-2013, 2013.
- Schneising, O., Reuter, M., Buchwitz, M., Heymann, J., Bovensmann, H., and Burrows, J. P.: Terrestrial carbon sink observed from space: variation of growth rates and seasonal cycle amplitudes in response to interannual surface temperature variability, *Atmos. Chem. Phys.*, 14, 133-141, 2014.
- Thoning, K. W., Tans, P. P., and Komhyr, W. D.: Atmospheric carbon dioxide at Mauna Loa Observatory, 2. Analysis of the NOAA/GMCC data. 1974-1985, *J. Geophys. Res.*, 94, 8549-8565, 1989.
- Wieder, W. R., Cleveland, C. C., Smith, W. K., and Todd-Brown, K.: Future productivity and carbon storage limited by terrestrial nutrient availability, *Nature Geoscience*, doi:10.1038/NGEO2313, 2015.
- Wunch, D., Toon, G. C., Blavier, J.-F. L., Washenfelder, R. A., Notholt, J., Connor, B. J., Griffith, D. W. T., Sherlock, V., and Wennberg, P. O.: The Total Carbon Column Observing Network. *Phil. Trans. R. Soc. A*, 369, 2087–2112, doi:10.1098/rsta.2010.0240, 2011.
- Wunch, D., Toon, G. C., Sherlock, V., Deutscher, N. M, Liu, X., Feist, D. G., and Wennberg, P. O.: The Total Carbon Column Observing Network's GGG2014 Data Version, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA, available at: doi:10.14291/tccon.ggg2014.documentation.R0/1221662, 2015.
- Yoshida, Y., Kikuchi, N., Morino, I., Uchino, O., Oshchepkov, S., Bril, A., Saeki, T., Schutgens, N., Toon, G. C., Wunch, D., Roehl, C. M., Wennberg, P. O., Griffith, D. W. T., Deutscher, N. M., Warneke, T., Notholt, J., Robinson, J., Sherlock, V., Connor, B., Rettinger, M., Sussmann, R., Ahonen, P., Heikkinen, P., Kyrö, E., Mendonca, J., Strong, K., Hase, F., Dohe, S., and Yokota, T.: Improvement of the retrieval algorithm for GOSAT SWIR XCO<sub>2</sub> and XCH<sub>4</sub> and their validation using TCCON data, *Atmos. Meas. Tech.*, 6, 1533–1547, doi:10.5194/amt-6-1533-2013, 2013.

## Acknowledgements

This study has been funded in parts by the European Space Agency (ESA) (via the GHG-CCI project of ESA's Climate Change Initiative (CCI, <http://www.esa-ghg-cci.org/>)), by the European Union (EU) (via the Copernicus Climate Change Service (C3S, <https://climate.copernicus.eu/>) managed by the European Centre for Medium-range Weather Forecasts (ECMWF)) and by the State and the University of Bremen. The University of Leicester GOSAT retrievals used the ALICE High Performance Computing Facility at the University of Leicester. We thank ESA/DLR for providing us with SCIAMACHY Level 1 data products and JAXA for GOSAT Level 1B data. We also thank ESA for making these GOSAT products available via the ESA Third Party Mission archive.

10 We thank NIES for the operational GOSAT XCO<sub>2</sub> Level 2 product and the NASA/ACOS team for the GOSAT ACOS Level 2 XCO<sub>2</sub> product. We also thank NOAA for the global CO<sub>2</sub> growth rates (file: [ftp://aftp.cmdl.noaa.gov/products/trends/co2/co2\\_gr\\_gl.txt](ftp://aftp.cmdl.noaa.gov/products/trends/co2/co2_gr_gl.txt); access: 24-Nov-2017) and for the Mauna Loa (MLO) CO<sub>2</sub> growth rates (file: [ftp://aftp.cmdl.noaa.gov/products/trends/co2/co2\\_gr\\_mlo.txt](ftp://aftp.cmdl.noaa.gov/products/trends/co2/co2_gr_mlo.txt); access: 9-Aug-2018). The fossil fuel and industry CO<sub>2</sub> emissions have been obtained from the Global Carbon Project website

15 (<http://www.globalcarbonproject.org/carbonbudget/17/data.htm>; access: 20-Nov-2017). The Southern Oscillation Index (SOI) data have been obtained from NOAA (file [https://www.esrl.noaa.gov/psd/gcos\\_wgsp/Timeseries/Data/soi.long.data](https://www.esrl.noaa.gov/psd/gcos_wgsp/Timeseries/Data/soi.long.data); access: 20-Nov-2017). The Oceanic Niño Index (ONI) data have also been obtained from NOAA (file <https://www.esrl.noaa.gov/psd/data/correlation/oni.data>). The CAMS CO<sub>2</sub> re-analysis data set has been obtained

20 from <http://apps.ecmwf.int/datasets/data/cams-ghg-inversions/> (access: 13-Nov-2018). Finally, we would like to thank four anonymous referees for helpful comments.

## Author contributions

M.B. supported by O.S., M.R., H.Boe., S.N., B.G., J.P.B.: design, data analysis, interpretation and writing of the

25 paper. The paper has been significantly improved by: H.Boe., J.A., R.J.P., P.S., R.G.D., O.P.H., I.A., A.B., A.K., H.S., Y.Y., D.C., C.O'D. Satellite input data have been provided by: M.R., M.B., O.S. (SCIAMACHY products) and H.Boe., J.A., R.J.P., P.S., R.G.D., O.P.H., I.A., A.B., A.K., H.S., Y.Y., D. C., C.O'D. (GOSAT products).

**Data availability.** The O4Mv3 XCO<sub>2</sub> data product (but also the underlying EMMAv3 product and those


30 individual sensor Level 2 input products which have been generated with European retrieval algorithms) are available via the Copernicus Climate Change Service (C3S, <https://climate.copernicus.eu/>) Climate Data Store (CDS, <https://cds.climate.copernicus.eu/>). Earlier versions are available from the GHG-CCI website (<http://www.esa-ghg-cci.org/>) of the European Space Agency (ESA) Climate Change Initiative (CCI, e.g., Obs4MIPs version 2 (O4Mv2) covering the years 2003-2015).

### **Competing financial interests**


The authors declare no competing financial interests.




## Carbon dioxide SCIAMACHY/ENVISAT & TANSO-FTS/GOSAT



**Figure 1.** Time series and global maps of satellite-derived column-average dry-air mole fractions of carbon dioxide, i.e.,  $\text{XCO}_2$ . Shown is data product Obs4MIPs version 3 (O4Mv3) based on an ensemble of 5 SCIAMACHY/ENVISAT (until April 2012) and TANSO-FTS/GOSAT (since mid 2009) individual sensor / individual soundings (Level 2) data products. The three time series correspond to three latitude bands: 30°N-60°N (red), 30°S-30°N (green) and 60°S-30°S (blue). The maps in the top left show monthly  $\text{XCO}_2$  for April and September 2003 (SCIAMACHY, land only) and the maps on the bottom right show monthly  $\text{XCO}_2$  for April and September 2016 (TANSO-FTS, land and ocean glint).



**Figure 2.** Atmospheric CO<sub>2</sub> and corresponding growth rates. (a) Monthly mean XCO<sub>2</sub> (red line) as obtained from averaging XCO<sub>2</sub> data product O4Mv3 globally for each month. (b) Monthly sampled annual CO<sub>2</sub> growth rates as computed from the red curve shown in (a) including 1-sigma uncertainty (grey vertical bars). (c) Annual mean growth rates computed from averaging the values shown in (b) including 1-sigma error estimates (vertical bars) (the numerical values are listed in Tab. A1 of Annex A). The NOAA annual mean global growth rate is also shown in (c) for comparison (in blue). Also listed in (c) is the linear correlation coefficient (R), the mean difference and the standard deviation of the difference of the satellite and the NOAA growth rates for 2003-2016 and for 2004-2016.



**Figure 3.** Carbon dioxide global annual mean growth rates compared with human emissions and ENSO indices. (a) Satellite-derived global annual mean growth rates (with 1-sigma uncertainty range shown as vertical lines). (b) CO<sub>2</sub> emissions from fossil fuel and industry (the correlation with the growth rate is  $R^2 = 31\%$ ). (c) Correlation in terms of  $R^2$  of growth rate and annual SOI (blue curve) and ONI (green curve) as a function of time shift in months. (d) Annual SOI for no shift (blue dotted line,  $R^2 = 10\%$ ) and for a shift of 7 months (blue solid line,  $R^2 = 30\%$ ) and annual ONI for no shift (green dotted line,  $R^2 = 13\%$ ) and for a shift of 4 months (green solid line,  $R^2 = 35\%$ ).

**Table 1.** Satellite XCO<sub>2</sub> data products. Individual satellite sensor XCO<sub>2</sub> algorithms and corresponding Level 2 data products used for generating the EMMAv3 Level 2 (i.e., individual soundings) data product, which has been gridded to obtain the O4Mv3 Level 3 data product used in this study. GHG-CCI refers to the GHG-CCI project of ESA’s Climate Change Initiative (<http://www.esa-ghg-cci.org/>) and C3S is the Copernicus Climate Change Service (<https://climate.copernicus.eu/>).

<b>Algorithm (Version)</b>	<b>Sensor</b>	<b>Comment</b>	<b>Reference</b>
BESD (v02.01.02)	SCIAMACHY / ENVISAT	GHG-CCI / C3S product ID: CO2_SCI_BESD	Reuter et al., 2011
RemoTeC (v2.3.8)	TANSO-FTS / GOSAT	GHG-CCI / C3S product ID: CO2_GOS_SRF	Butz et al., 2011
UoL-FP (v7.1)	TANSO-FTS / GOSAT	GHG-CCI / C3S product ID: CO2_GOS_OCFP	Cogan et al., 2012
ACOS (v7.3.10a)	TANSO-FTS / GOSAT	NASA’s GOSAT XCO <sub>2</sub> product	O’Dell et al., 2012
NIES (v02)	TANSO-FTS / GOSAT	Operational GOSAT product	Yoshida et al., 2013

## Annex A


Growth rate time series have also been computed for several latitude bands as shown in Fig. A1. As can be seen, the growth rates agree within their 1-sigma uncertainty range in all latitude bands including the global results (for numerical values see Tab. A1).

The reason for this is that atmospheric CO<sub>2</sub> is long-lived and therefore well-mixed. Because of this we expect similar annual mean CO<sub>2</sub> growth rates, i.e., agreement within measurement error, for the different latitude bands and globally. Identical growth rates are not expected due to differences in the sources and sinks and the time needed for transport and mixing. The expectation of similar growth rates is corroborated by Fig. A2, which shows a comparison of the uncertainty of the satellite-derived growth rates (red bars) with the difference of two annual mean CO<sub>2</sub> growth rate time series from NOAA, namely the time series from Mauna Loa, Hawaii, and the global time series obtained from globally averaged marine surface data (both obtained from <https://www.esrl.noaa.gov/gmd/ccgg/trends/gr.html>). As shown in Fig. A2, the uncertainty of the satellite data is similar (mean value: 0.34 ppm/year) as the difference between the two NOAA time series (standard deviation: 0.21 ppm/year). We acknowledge that the maximum difference between any two latitude bands may be somewhat larger than the difference between the two NOAA time series shown in Fig. A2, but it is assumed that the difference shown in Fig. A2 is at least a reasonable approximation.


The agreement shown in Fig. A1 is interpreted as an indication of the good quality of the satellite XCO<sub>2</sub> data product and of the adequacy of the method used to compute the annual mean CO<sub>2</sub> growth rates because we do not find “strange values” in certain latitude bands or certain years, which would be an indication for a potential problem.

Figure A3 shows a comparison of XCO<sub>2</sub> and surface CO<sub>2</sub> annual growth rates as computed from a Copernicus Atmosphere Monitoring Service (CAMS) global re-analysis CO<sub>2</sub> data set (Chevallier, 2018). This CAMS atmospheric CO<sub>2</sub> data set does not (in contrast to satellite data) suffer from data gaps and measurement noise. Therefore, the annual growth rate can simply be computed from the difference of the (XCO<sub>2</sub> or surface CO<sub>2</sub>) values at the end of a year and the beginning of that year (“method M1”). Figure A3b confirms that growth rates computed (using method M1) from XCO<sub>2</sub> and from surface CO<sub>2</sub> are very similar but not exactly identical. Figure A3c shows that the satellite method (“M2”) described in this publications provides annual XCO<sub>2</sub> growth rates, which are very similar to those obtained with the M1 method.


30



**Figure A1.** Satellite-derived annual mean XCO<sub>2</sub> growth rates: Global (black), Northern Hemisphere (NH) mid latitudes (“NHmidlat” (30°N - 60°N), red), Tropics (30°S - 30°N, green), and Southern Hemisphere mid latitudes (5 “SHmidlat” (60°S - 30°S), blue). The corresponding numerical values are listed in Tab. A1.



5 **Figure A2.** Comparison of the 1-sigma uncertainty range of the satellite-derived growth rates (red bars) with the difference of two annual mean growth rate time series obtained from NOAA, namely the time series from Mauna Loa (MLO), Hawaii, and the global time series obtained from globally averaged marine surface data (black line and symbols).



**Figure A3.** (a) Monthly mean global time series of XCO<sub>2</sub> (red) and surface CO<sub>2</sub> (blue) as computed from a Copernicus Atmosphere Monitoring Service (CAMS) CO<sub>2</sub> re-analysis data product (version v17r1 obtained from <http://apps.ecmwf.int/datasets/data/cams-ghg-inversions/>; Chevallier, 2018). The symbols correspond to the values at the beginning/end of each year. (b) Annual XCO<sub>2</sub> (red) and surface CO<sub>2</sub> (blue) growth rates as computed from the time series shown in (a) by computing for each year the difference of the values at the end and the beginning of that year (method M1). Also listed is the linear correlation coefficient R, the mean difference and the standard deviation of the difference. (c) The red symbols (and the red curve) show the same values as the red symbols shown in (b), i.e., they show annual XCO<sub>2</sub> growth rates computed using method M1. The green symbols also show XCO<sub>2</sub> annual growth rates but computed using the “satellite method” (M2), which is described in this publication.



**Table A1.** Satellite-derived annual mean XCO<sub>2</sub> growth rates in ppm/year including 1-sigma uncertainty (in brackets). Abbreviations: NH is Northern Hemisphere and SH is Southern Hemisphere.

Year	Latitude band / region			
	Global	NH mid-latitudes (30°N-60°N)	Tropics (30°S-30°N)	SH mid-latitudes (60°S-30°S)
<b>2003</b>	1.66 (0.76)	1.99 (0.72)	1.54 (0.74)	1.77 (0.62)
<b>2004</b>	1.59 (0.30)	1.52 (0.29)	1.71 (0.29)	1.30 (0.23)
<b>2005</b>	2.16 (0.28)	2.51 (0.26)	1.99 (0.28)	2.17 (0.22)
<b>2006</b>	2.21 (0.27)	2.13 (0.25)	2.22 (0.27)	2.33 (0.21)
<b>2007</b>	2.26 (0.27)	2.33 (0.25)	2.20 (0.26)	2.34 (0.21)
<b>2008</b>	1.67 (0.29)	1.60 (0.27)	1.81 (0.28)	1.41 (0.20)
<b>2009</b>	1.77 (0.30)	1.75 (0.30)	1.86 (0.28)	1.70 (0.21)
<b>2010</b>	2.22 (0.29)	2.67 (0.29)	2.08 (0.27)	2.14 (0.20)
<b>2011</b>	1.86 (0.28)	1.69 (0.27)	1.86 (0.27)	2.19 (0.19)
<b>2012</b>	2.46 (0.29)	2.64 (0.28)	2.44 (0.27)	2.38 (0.21)
<b>2013</b>	2.27 (0.30)	2.38 (0.28)	2.27 (0.28)	2.10 (0.22)
<b>2014</b>	1.74 (0.31)	1.53 (0.30)	1.80 (0.29)	1.84 (0.23)
<b>2015</b>	2.89 (0.34)	2.89 (0.31)	2.97 (0.32)	2.54 (0.25)
<b>2016</b>	3.23 (0.50)	3.28 (0.46)	3.23 (0.48)	3.41 (0.36)