

**Transport of trace gases via eddy shedding from the Asian summer monsoon anticyclone
and associated impacts on ozone heating rates**

Suvarna Fadnavis¹, Chaitri Roy¹, Rajib Chattopadhyay¹, Christopher E. Sioris², Alexandru
Rap³, Rolf Müller⁴, K. Ravi Kumar⁵ and Raghavan Krishnan¹

¹Indian Institute of Tropical Meteorology, Pune, India

²Environment and Climate Change, Toronto, Canada

³School of Earth and Environment, University of Leeds, Leeds, United Kingdom

⁴Forschungszentrum Jülich GmbH, IEK-7, Jülich, Germany

⁵King Abdullah University of Science and Technology, Thuwal, Saudi Arabia

*Email of corresponding author: suvarna@tropmet.res.in

Abstract:

The highly vibrant Asian Summer Monsoon (ASM) anticyclone plays an important role in efficient transport of Asian tropospheric air masses to the extratropical upper troposphere and lower stratosphere (UTLS). In this paper, we demonstrate long-range transport of Asian trace gases via eddy shedding events using MIPAS (Michelson Interferometer for Passive Atmospheric Sounding) satellite observations, ERA-Interim re-analysis data and the ECHAM5–HAMMOZ global chemistry–climate model. Model simulations and observations consistently show that Asian boundary layer trace gases are lifted to UTLS altitudes in the monsoon anticyclone and are further transported horizontally eastward and westward by eddies detached from the anticyclone. We present an event of eddy shedding during 1-8 July 2003 and discuss a 1995-2016 climatology of eddy shedding events. Our analysis indicates

that eddies detached from the anticyclone contribute to the transport of Asian trace gases away from the Asian region to the West-Pacific (20°-30° N; 120°-150° E) and West-Africa (20°-30° N, 0°-30° E). Over the last two decades, the estimated frequency of occurrence of eddy shedding events is ~68 % towards West-Africa and ~25 % towards the West-Pacific.

Model sensitivity experiments considering a 10 % reduction in Asian emissions of non-methane volatile organic compounds (NMVOCs) and nitrogen oxides (NO_x) were performed with ECHAM5-HAMMOZ to understand the impact of Asian emissions on the UTLS. The model simulations show that transport of Asian emissions due to eddy shedding significantly affects the chemical composition of the upper troposphere (~100-400 hPa) and lower stratosphere (~100-80 hPa) over West-Africa and the West-Pacific. The 10 % reduction of NMVOCs and NO_x Asian emissions leads to decreases in peroxyacetyl nitrate (PAN) (2-10 % near 200-80 hPa), ozone (1-4.5 % near ~150 hPa) and ozone heating rates (0.001-0.004 K·day⁻¹ near 300-150 hPa) in the upper troposphere over West-Africa and the West-Pacific.

Key Words: Asian summer monsoon anticyclone; Eddy shedding from the monsoon anticyclone, Transport of Asian trace gases, Ozone heating rates; ECHAM5-HAMMOZ model.

1. Introduction

Rapid industrialization, traffic growth, and urbanization resulted in significant increases in the concentrations of tropospheric trace gases, such as carbon dioxide (CO_2), carbon monoxide (CO) and methane (CH_4) over Asia. There is global concern about rising levels of these trace gases (due to their global warming potential) as they are projected to increase further over the coming years despite efforts to implement several mitigation strategies (Ohara et al., 2007). In situ observations, satellite measurements, trajectory analysis and model simulations show long range transport of Asian trace gases to remote locations (e.g. North America, Europe) (Liang et al., 2004). The transported trace gases change the radiative balance, dynamics and chemical composition at the respective locations (Vogel et al., 2016). Satellite observations show increasing trends in several tropospheric Asian trace gases over the last decade, e.g. ozone at $\sim 1\text{-}3\%$ year^{-1} (Verstraeten et al., 2015), CO at 3% year^{-1} (Strode and Pawson, 2013), NO_x at $\sim 3.8\text{-}7.3\%$ year^{-1} (Schneider and van der A, 2012; Ghude et al., 2013). Biomass burning is another major contributor to the observed growth in these trace gases (van der Werf et al., 2006). Peroxyacetyl nitrate (PAN), a powerful pollutant formed in biomass burning plumes (Wayne, 2000), is a secondary pollutant produced through the oxidation of hydrocarbons released from anthropogenic and biogenic sources. It is a reservoir of reactive nitrogen and plays a fundamental role in the global ozone budget (Tereszchuk et al., 2013; Payne et al., 2017). PAN can also be formed in the upper troposphere through the production of NO_x from lightning (Zhao et al., 2009). Simulations of the Model of Ozone and Related Tracers (MOZART) show an increase of 20-30 % of PAN concentrations in the upper troposphere and lower stratosphere (UTLS) over the Asian summer monsoon (ASM) region produced from

64 lightning (Tie et al., 2002). While in the lower troposphere, PAN has a short lifetime (a few
65 hours), in the UTLS it has a longer lifetime (3-5 months), and can therefore act as a reservoir
66 and carrier of NO_x (Tereszchuk et al., 2013). Recent satellite observations show an increasing
67 trend in PAN ($\sim 0.1 \pm 0.05$ to 2.7 ± 0.8 ppt year⁻¹) in the UTLS over Asia (Fadnavis et al.,
68 2014).

69 Monsoon convection plays an important role in lofting of boundary layer Asian air masses to
70 the UTLS (e.g., Randel et al., 2010; Fadnavis et al., 2015; Santee et al., 2017). The uplifted air
71 masses become confined into the anticyclone enclosed by jets (westerly and easterly jets to the
72 north and south, respectively), which act as a strong transport-barrier and restrict isentropic
73 mixing into the extra-tropical lower stratosphere or the equatorial tropics (Ploeger et al., 2015;
74 Ploeger et al., 2017). Confinements of high concentrations of trace gases, including ozone
75 precursors (e.g., hydrogen cyanide (HCN), CO, hydrochloric acid (HCl), NO_x and PAN), and
76 low ozone in the anticyclone are evident in satellite and aircraft observations, (Randel et al.,
77 2010; Vogel et al., 2014; Fadnavis et al., 2015; Ungermann et al., 2016; Santee et al., 2017).
78 The observed ozone minimum in spite of high amounts of its precursors in the anticyclone is
79 still an open question. A fraction of these trace gases enters the lower stratosphere and affects
80 the UTLS chemical composition (Randel et al., 2010; Fadnavis et al., 2015, 2016; Garny and
81 Randel, 2016), with associated radiative forcing impacts (Riese et al., 2012). Cross-tropopause
82 transport associated with the Asian monsoon is evident in a number of species, including
83 aerosols, hydrogen cyanide (HCN) and PAN (Randel et al. 2010; Fadnavis et al. 2014, 2015;
84 Bourassa et al., 2012).

The ASM anticyclone is highly dynamic in nature (e.g., Hsu and Plumb, 2000; Popovic and Plumb, 2001; Vogel et al., 2016). On the sub-seasonal scale, it shows variation in strength and location (Garny and Randel, 2016). It frequently sheds eddies and on occasions, it splits into two anticyclones, namely the Tibetan and Iranian anticyclones (Zhang et al., 2002; Nützel et al., 2016). An eddy detached from the anticyclone carries Asian air masses (trace gases) away from the ASM region. There are scattered studies indicating eddy shedding to the west (Popovic and Plumb, 2001) and east (Ungermann et al., 2016; Vogel et al., 2014) of the anticyclone. An eddy shedding event causes irreversible mixing in the surrounding air changing the chemical composition and radiative balance of that region (Garny and Randel, 2016). Here, we analyze in detail transport of Asian trace gases via eddies, subsequent mixing into the extra-tropics and radiative impact of eddy shedding events on decadal scales. In this study, we answer the following questions: (1) how frequent were eddy shedding events during the last two decades? (2) Which regions are the most affected? (3) Does the transport of Asian trace gases arising from eddy shedding affect UTLS ozone concentrations and heating rates at remote locations?

To address these questions, we first consider an eddy shedding event demonstrating eastward and westward shedding from the ASM anticyclone during 1-8 July 2003. This year was chosen since the monsoon season was quite normal (i.e., no evidence of El Niño or Indian Ocean dipole phenomenon influencing the monsoon circulation). We then present a climatology of eddy shedding events and lead-lag relations of eddies with the anticyclone. We also evaluate the impact of increasing Asian emissions of NO_x and NMVOCs on ozone and PAN during the eddy shedding event, using model sensitivity simulations. Finally, we estimate the associated

changes in ozone heating rates in the UTLS due to Asian trace gases transported via eddy shedding events.

2. Model set-up and satellite observations

2.1 Satellite observations

The [Michelson Interferometer for Passive Atmospheric Sounding \(MIPAS\)](#) on-board the [European ENVironmental SATellite \(ENVISAT\)](#) (MIPAS-E) was launched in March 2002 into a polar orbit of 800 km altitude. Its orbital period is about 100 min. MIPAS-E provided continual limb emission measurements in the mid-infrared over the range 685– 2410 cm^{-1} (14.6–4.15 μm) (Fischer et al., 2008). MIPAS monitored many atmospheric trace constituents including CO, PAN, and O₃. The details of the general retrieval method and setup, error estimates and use of averaging kernel and visibility flag are documented by von Clarmann et al. (2009). Here, [we analyze the MIPAS observations of CO, PAN, and O₃ during 1-8 July 2003.](#)

To account for the comparatively low, and altitude-dependent vertical resolution of MIPAS, the model data were convolved with the MIPAS averaging kernel to be directly comparable to MIPAS measurements of CO, PAN, and ozone. MIPAS vertical resolution for CO, O₃ and PAN in the UTLS is 5, 3.5 and 5 km, respectively. [The data are contoured and gridded. For each grid point, the surrounding MIPAS data points are averaged while applying a distance weighting. The maximum distance for which MIPAS data points are considered is \$\pm 7\$ deg in latitude and \$\pm 15\$ deg in longitude \(covering a box of 14 deg in latitude and 30 deg in longitude\), and a minimum number of 2 data points per interpolation grid point is required.](#)

The data quality specifications as documented at <http://share.lsd.fkit.edu/imk/asf/sat/mipas-export/Documentation/> were employed, namely: only data with a visibility flag equal to 1 and a diagonal value of averaging kernel greater than 0.03 were used for ozone and PAN, while 0.008 was used for CO (Glatthor et al., 2007; Funke et al., 2009)..

2.2 Model set-up

We employ the ECHAM5-HAMMOZ (Roeckner et al., 2003) aerosol-chemistry-climate model to understand re-distribution of Asian trace gases via eddy shedding from the anticyclone. ECHAM5-HAMMOZ comprises of the general circulation model ECHAM5 (Roeckner et al., 2003), the tropospheric chemistry module MOZ (Horowitz et al., 2003), and the aerosol module, Hamburg Aerosol Model (HAM) (Stier et al., 2005). The chemistry of ozone, VOCs, NO_x, and other gas-phase species is based on the MOZART-2 chemical scheme (Horowitz et al., 2003). It includes O_x-NO_x-hydrocarbons with 63 tracers and 168 reactions. The details of the parameterizations and emissions used in the model as well as a validation of the results are described by Fadnavis et al. (2013, 2014, 2015) and Pozzoli et al. (2011).

The model simulations were performed with a T42 spectral resolution corresponding to about 2.8° × 2.8° in the horizontal dimension and 31 vertical hybrid σ -p levels from the surface up to 10 hPa. Here, we note that our base year for aerosol and trace gas emissions is 2000. We performed two simulations: (i) a control experiment (CTRL), and (ii) a sensitivity experiment (Asia10), where emissions of both NO_x and NMVOCs were simultaneously reduced by 10 % over Asia (10° S–50° N, 60–130° E) similar as in earlier publication (Naik et al., 2005; Fadnavis et al., 2015). This fixed 10% reduction was chosen due to the spatial-temporal

variability of NMVOCs over Asia and the inherent difficulty in obtaining a common trend value (Li et al., 2014). The impacts of this NMVOCs and NO_x emission perturbation are investigated by analyzing the associated anomalies (Asia10 – CTRL) in ozone, PAN and ozone heating rates.

Both simulations were performed for the year 2003 driven by European Centre for Medium-Range Weather Forecasts operational analyses (Integrated Forecast System (IFS) cycle-32r2) meteorological fields (available every six hours) (Uppala et al., 2005). All simulations include lightning NO_x and the subsequent PAN production. Since the lightning parameterization is the same in the CTRL and sensitivity simulations, its impact may be negligible. However, there may be an indirect impact of changed emissions on lightning and thus on NO_x or PAN production. The model simulations used here are the same as those used by Fadnavis et al. (2015). The climatology of ozone mass mixing ratio, winds and potential vorticity (PV) are obtained from ERA-Interim reanalysis data for the period 1995-2016. The anomalies are obtained from difference between daily mean values of July 2003 and daily climatology. Power spectral analysis and lag/lead correlations have been carried out on PV data for the period 1995-2016 to show climatological features.

Instantaneous ozone heating rates are calculated using the Edwards and Slingo (1996) radiative transfer model. We used the off-line version of the model, with six shortwave and nine longwave bands, and a delta-Eddington 2-stream scattering solver at all wavelengths, in a set-up similar to other recent studies (Rap et al., 2015, Roy et al., 2017).

3. Results

3.1 A typical case study of eddy shedding from the monsoon anticyclone

The dynamics of the monsoon anticyclone is best portrayed at the 370 K potential temperature surface and the monsoon anticyclone is obvious as an area of low PV values ($PV < 2$ PVU, $1 \text{ PVU} = 10^{-6} \text{ K m}^2 \text{ kg}^{-1} \text{ s}^{-1}$) (indicating tropospheric air-mass) at this surface (Garny and Randel, 2016). Eddies are identified as air with low PV emanating from the monsoon anticyclone (Popovic and Plumb, 2001; Vogel et al., 2014). Past studies have shown that during the monsoon season (June to September), the bulk of the low PV air at the isentropic level of 370 K, is confined between about $20\text{--}35^\circ \text{ N}$ and $20\text{--}120^\circ \text{ E}$ indicating the spatial extent of the anticyclone (Popovic and Plumb, 2001; Vogel et al., 2014; Garny and Randel, 2016). A pocket of low PV air-mass detached from the boundary of the anticyclone (outside the anticyclone, $20\text{--}35^\circ \text{ N}$ and $20\text{--}120^\circ \text{ E}$) is considered as an eddy. **Figures 1a-h** show the distribution of PV at 370 K during 1-8 July 2003. It can be seen that during this period the anticyclone was wobbling and shed eddies eastward and westward over West-Africa ($20\text{--}30^\circ \text{ N}$, $0\text{--}30^\circ \text{ E}$) and the West-Pacific ($20\text{--}30^\circ \text{ N}$; $120\text{--}150^\circ \text{ E}$). Initially, during 2-5 July 2003, the ASM anticyclone shed an eddy westward over West-Africa. The eddy moved further west with the progression of time. Later during 4-8 July 2003, eddy shedding occurred to the east of the anticyclone, over the West-Pacific and the air detached from the anticyclone moved further eastward with time. The longitude-pressure section of PV shows that the eddy protrudes down to 400 hPa (not shown).

Previous studies have shown that eddy shedding events are associated with Rossby wave breaking (RWB) (Hsu and Plumb, 2000; Popovic and Plumb, 2001; Fadnavis and Chattopadhyay, 2017). The RWB is manifested as a rapid and large-scale irreversible overturning of PV contours on the 350 K isentropic surface. It is accompanied with a cyclonic circulation at 200 hPa (Strong and Magnusdottir, 2008; Fadnavis and Chattopadhyay, 2017). **Figures 2a-h** show the distribution of PV at the 350 K surface and the circulation at 200 hPa during 1-8 July 2003. It can be seen that, during 1-8 July 2003, three RWB events occurred: one near 30° E (referred as RWB-1), one near 70° E (referred as RWB-2) and another one near 120° E (referred to as RWB-3). Since RWB-3 was outside the region of the ASM anticyclone (over the West-Pacific ~150-170° E) it did not play a role in the eddy shedding event of 1-8 July. If we track the location of these RWB events (indicated by the black and red arrows), one can see that, with the progression of time, the RWB feature moved eastward. The eastward migration of RWB is linked to its movement along the subtropical westerly jet (Fadnavis and Chattopadhyay, 2017). Initially during 1-5 July RWB-1 was strong ($PV > 2$ PVU) while RWB-2 ($PV < 2$ PVU) was weak. During this period the southward and westward moving RWB-1 leads to eddy shedding over West Africa. Later, during 4-8 July, RWB-2 strengthened while RWB-1 weakened and disappeared. The southward and eastward moving RWB-2 was responsible for the eddy shedding event near the Western Pacific (see **Figs. 2d-h**).

3.2. Climatology of eddy shedding from the monsoon anticyclone

A power spectrum analysis (PSA) has been performed on the PV data (averaged for 300-100 hPa) during 1995-2016 for West-Africa (20-30° N, 0-30° E) and the West-Pacific (20-30° N,

140-150° E). The PSA uses the temporal-to-frequency fast Fourier transform in order to identify dominant signal frequencies. It provides information of signal power (square of variance) associated with the frequency components of the signal, with the dominant signal periodicity being the inverse of the dominant signal frequency. **Figures 3a-b** show the distribution of power spectral variance over the West-Africa and West-Pacific regions. The variances corresponding to the periodicities of 3-5 days, 12-15, and 18-21 days are significant at 95 % confidence level for both the regions indicating that the eddy shedding activity is dominated in the range of synoptic frequency (~10 days). Popovic and Plumb (2001) also indicated a typical duration of an eddy shedding event of ~4-8 days. We compute the frequency of eddy shedding days ($PV < 1$ PVU) occurring over West-Africa and the Western Pacific. The ERA-Interim data for the last two decades show that eddy shedding is quite frequent over west-Africa (~68 %) and the West-Pacific (~25 %). The lag-lead correlation of PV (averaged for 200-100 hPa) for the centre region of the anticyclone (85-90° E, 28-30° N) with PV averaged over the West-Pacific shows a maximum positive lead correlation at 3-4 days (**Fig. 3c**). Similarly, PV over West-Africa shows a maximum positive lead correlation for 5-6 days with the PV averaged over the monsoon anticyclone (**Fig. 3d**). This indicates that the transport of the eddies from the anticyclone (source region) has a typical duration of three to four days over the West Pacific and five to six days over West Africa. This transport time is the timescale over which the trace gases are moved to remote locations from the ASM anticyclone.

3.3. Long range transport of trace gases

3.3.1 Horizontal transport of ozone, CO and PAN via eddies

Biomass burning over south-east Asia and East Asia produces large amounts of CO, NO_x, VOCs, PAN, ozone and aerosols (e.g., Streets et al., 2003, Fadnavis et al., 2014). The monsoon convection over the Bay of Bengal, southern slopes of Himalaya and South China Sea (see **Fig. S1**) lifts up these species into the anticyclone where they may get dispersed in the UTLS by the vibrant anticyclone and its associated eddies. **Figures 4a-h** show the distribution of ozone during 1-8 July 2003 (MIPAS O₃ is binned for 2 days and simulated O₃ is plotted for alternate days) in the anticyclone at 16 km (~100 hPa). Ozone concentrations from MIPAS satellite measurements and model simulations (CTRL) are plotted at 16 km and from ERA-Interim reanalysis at 100 hPa. For comparison, we have interpolated the model data to the MIPAS altitude grid and smoothed with the averaging kernel. The ASM anticyclone is marked by minimum ozone although its precursors (e.g. CO, NO_x and CH₄) show maxima (Randel et al., 2010; Roy et al., 2017). The spatial pattern of low ozone amounts in the anticyclone and the associated eddies is evident in all of the data sets during 1-8 July 2003. The locations of ozone local minima in the model are slightly shifted relative to the locations of eddies and relative to the locations of ozone local minima in MIPAS and ERA. During 1-5 July, ozone concentrations in the eddy over West-Africa are ~40-200 ppb in MIPAS, ~60-180 ppb in ERA-Interim and 100-200 ppb in the model simulations. During 4-8 July, the eddy over the west Pacific shows ozone amounts of ~60-180 ppb in MIPAS, ~60-180 ppb in ERA-

252 Interim and ~120-200 ppb in the model simulations. In general, model overestimates ozone
253 amounts by ~60 ppb than ERA-Interim and MIPAS measurements.

254 **Figures 5a-h** show the distribution of CO from MIPAS observations and model simulations
255 during 1-8 July 2003 (MIPAS CO is binned for 2 days and simulated CO is plotted for
256 alternate days). The confinement of high concentrations of CO in the anticyclone and in eddies
257 is seen in both MIPAS observations and model simulations. During 1-5 July, eddies over west-
258 Africa and west-Pacific show CO volume mixing ratios of ~85-95 ppb in MIPAS, and ~70-95
259 ppb in the model simulations. Similar to ozone, a location of maximum in CO is not collocated
260 with eddies and slightly different in MIPAS and Model.

261 **Figures 6a-h** show the distribution of PAN from MIPAS measurements and the model
262 simulation (CTRL) at 16 km during 1-8 July 2003 (MIPAS PAN mixing ratios are binned for
263 2 days and simulated PAN is plotted for alternate days). A confinement of high amounts of
264 PAN in the anticyclone and the associated eddies is seen both in the MIPAS measurements
265 and the model simulations. During 1-5 July, MIPAS observed PAN amounts are ~120-240 ppt
266 in eddies over west-Africa, while the model simulation shows ~180-240 ppt of PAN at the
267 same location. The eddy over the west-Pacific shows PAN amounts of ~160-240 ppt both in
268 MIPAS measurements and model simulations.

269 There are differences in ozone, CO and PAN amounts from model simulation, satellite
270 observations and ozone from ERA-Interim. These differences may be due to a number of
271 reasons e.g. different grid sizes of MIPAS, ERA-Interim and model data, binning of MIPAS
272 data for two days to accommodate better spatial coverage, uncertainties in the model emission

inventory, and retrieval errors in the satellite data. A maximum in PAN near the location of eddy differ in MIPAS and model. Comparison of Fig. 1 and Figs 4-6 shows that minimum in ozone and maximum in CO and PAN is not collocated at eddies. The location varies slight in species and data sets (in MIPAS, ERA and model). This may be due to differences in data sets and production and loss processes of each species.

3.3.2 Vertical distribution of CO, PAN and ozone

Further, we analyze the vertical distribution of CO and PAN as an indication of Asian biomass burning emissions. **Figure 7** shows longitude-pressure cross-sections (averaged for 20°-40° N) of CO and PAN from the CTRL simulation, with wind vectors depicting circulation patterns. It illustrates that during 1-5 July 2003 a plume of CO/PAN has been uplifted from the Asian region (80°-120° E), moving further upward into the UTLS. The location of the plume (**Fig. 7**) coincides with a strong convection region - see **Fig. S1**, showing combined cloud droplet (CDNC) and ice crystal (ICNC) number concentrations from the CTRL simulation. **Figure 7** and **Fig. S1**, together, indicate that surface emissions are lifted up by the monsoon convection. In the upper troposphere (~120 hPa), westward horizontal transport of CO/PAN towards West-Africa is obvious as a result of eddy shedding during the respective days. In particular, during 2-4 July high amounts of CO/PAN are observed near 0°-30° E at 100 hPa (**Figs. 7a-b** and **7e-f**). On 2 July, there is some PAN transport over the west-Pacific. During 4-8 July 2003, eddy shedding occurs to the east of the anticyclone over the West-Pacific (120°-150° E) (see **Figures 1e-f**). East-ward horizontal transport of CO/PAN in the regions of eddy shedding is evident in **Figs. 7c-d** and **7g-h**. The Asian trace gases then disperse downward deep into the

294 troposphere (to ~500 hPa over the West Pacific and to ~200 hPa over West-Africa) and are
295 partially lifted into the lower stratosphere.

296 The vertical distribution of ozone shows low ozone amounts extending from the convective
297 regions of Bay of Bengal and South China Sea (~15-25° N) upward in the upper troposphere
298 (Figs. S2a-d), with ozone amounts of ~100-200 ppb near the tropopause (see also Figs. 4-i-l).
299 The lower ozone amounts over the Asian troposphere may be due to clean marine air masses
300 during the monsoon season (Zhao et al., 2009). The feature of low ozone air-mass ascent is
301 less evident than the CO and PAN vertical ascent, due to a number of factors which are
302 influencing ozone production and loss processes at different altitudes in the troposphere and
303 lower stratosphere, such as stratospheric intrusions, lightning etc. (see discussions in section
304 3.4).

305 3.4 Influence of Asian emissions on extra-tropical UTLS

306 In this section, we investigate the influence of Asian anthropogenic emissions of
307 NMVOCs and NO_x on the distribution of PAN and ozone in the tropical/extra-tropical UTLS
308 from sensitivity experiments. Figures 8a-d show anomalies of PAN (Asia10-CTRL) at 16km
309 during 1-8 July 2003 (plotted on alternate days). The negative anomalies in PAN are seen
310 confined to the region of the anticyclone and the associated eddies (1-5 July over West-Africa
311 and 4-8 July over West-Pacific). These anomalies portray the effect of Asian boundary layer
312 emissions (NMVOCs and NO_x) on the upper level anticyclone and the associated eddies. A
313 number of studies (Randel et al., 2010; Fadnavis et al., 2013; 2015; Vogel et al., 2014) have
314 shown lifting of Asian emissions to the UTLS by the monsoon convection and its confinement

in the anticyclone. A 10% decrease in Asian NMVOCs and NO_x emissions decreases PAN amounts by ~5-23 % in the ASM anticyclone and the associated eddies over West-Africa and the West-Pacific.

Further, we analyze the vertical distribution of anomalies of PAN and ozone. **Figures 8e-h** show longitude-pressure sections of anomalies of PAN. It shows negative anomalies (in response to reduced Asian emissions) along the transport pathways (**Fig. S1**), i.e. from the boundary layer of the Asian region (80°-120° E) into the upper troposphere and westward/eastward transport from the anticyclone owing to eddy shedding. These anomalies extend above the tropopause, indicating cross-tropopause transport. Upward transport across the tropopause in monsoon season has been demonstrated to occur in recent tracer studies (Ploeger et al., 2017; Vogel et al., 2018). PAN is rather long lived in the cold tropopause region and should therefore behave similar as inert trace gases in the model studies (Fadnavis et al., 2014; 2015). Our simulations show that a 10 % reduction in Asian emissions of both NMVOCs and NO_x, results in a decrease in the amount of PAN by ~2-10 % over North-West Africa during 1-5 July and over the Western Pacific during 4-8 July 2003.

The vertical distribution of ozone anomalies (**Figs. 8i-l**) show negative values (–1 to –4.5 %) in the troposphere extending from the surface up to ~180 hPa along the transport pathways (~90° E) and in the region from where cross tropopause transport occurs. Near the tropopause (except in the region of cross-tropopause transport; indicated by boxes) ozone anomalies are positive, varying between 1 to 8 % (**Figs. 8i-l**). In contrast to PAN, ozone will be chemically active during the slow ascent over the monsoon area for several months (Vogel

et al., 2018). Ozone loss rates are likely to be affected in the Asia-10 simulations. For example reduced NO_x will lead to a lower efficiency of ozone loss providing a reason for higher ozone in the Asia-10 runs. Further, less NMVOCs in Asia-10 simulations might lead to lower OH concentrations in the lowermost stratosphere above the monsoon region. The major ozone loss cycle in the lowermost stratosphere in the tropics is driven by HO_x radicals with the rate limiting step being the reaction of OH with ozone. The anomalies of OH concentrations are negative near the tropopause indicating lower ozone loss rates (**Fig. S3**). The changes in dynamics (e.g stratospheric intrusions and lightning) due to emission sensitivity may also partially contribute to positive anomalies of ozone near the tropopause. Ozone distribution of from CTRL simulations show stratospheric intrusion in the northern part of the anticyclone $\sim 30^\circ\text{N}$ (**Fig. S2**) which is enhanced (positive anomalies) in Asia-10 simulations (**Figs. S4 a-d**). The spatial distribution of ozone anomalies (**Fig. S4 e-h**) indicate that the response to emission reductions generates negative anomalies of ozone in the southern part of anticyclone ($15\text{--}25^\circ\text{N}$; $60\text{--}120^\circ\text{E}$) (may be due to cross tropopause of monsoon air), while ozone anomalies are positive in the northern part of the anticyclone (which may be associated with stratospheric intrusions). The ozone variability near the tropopause is generally driven by the strong mixing of tropospheric and stratospheric air-masses.

In **Fig. 8i-l**, negative values of ozone anomalies extending from surface to ~ 180 hPa may be likely to be related to the vertical extent of transport and associated outflow. A plume of high values of CO (~ 95 ppb) and PAN (~ 260 ppt) (**Fig. 7**), together with relatively low ozone amounts ($70\text{--}80$ ppb) (**Fig. S2**) reaching to ~ 180 hPa and leads to a strong gradient near the tropopause. It also indicates that the outflow of uplifted trace gases in the upper

troposphere reaches to ~250-180 hPa. The moderate concentrations of CO and PAN between 180-70 hPa may also be due to the slow ascent into the lower stratosphere of these Asian pollutants (Park et al., 2008).

During the monsoon season, marine air masses containing low amounts of ozone prevail over the Asian land mass. The monsoon air mass gathers Asian boundary layer ozone precursors (and other trace gases) and are uplifted to the UTLs by the monsoon circulation. It should be noted that a decrease in emissions of NO_x and NMVOCs in the Asia10 simulations produces lower ozone amounts in the troposphere than CTRL. Therefore, in the regions of eddy shedding, negative anomalies near 200-300 hPa indicate transport of monsoon air (via eddies) towards West-Africa during 1-5 July and to the West-Pacific during 4-8 July.

.

3.5 Influence of Asian emission of trace gases on ozone heating rates

Ozone is a dominant contributor to radiative heating in the tropical lower stratosphere, impacting the local heating budget and non-local forcing of the troposphere below (Gilford and Solomon, 2017). We estimate changes in ozone heating rates caused by a 10 % decrease in Asian NMVOCs and NO_x emissions. **Figures 9a-d**, show anomalies of ozone heating rates on 1-8 July (plotted on alternate days), indicating a reduction in ozone heating rates in response to a decrease in Asian NMVOCs and NO_x emissions, coincident with the region of convective transport (also see Fig. S5). In the upper troposphere (300-180 hPa), the negative anomalies in ozone heating rates vary between -0.001 and -0.0045 K·day⁻¹. Interestingly, reduced Asian emissions (NMVOCs and NO_x), lead to a reduction in ozone, which leads to a

reduction in ozone heating rates (-0.001 to $-0.003 \text{ K}\cdot\text{day}^{-1}$) in the region of eddy shedding over West-Africa (1-5 July) and the West-Pacific (4-8 July). The ozone poor Asian air mass trapped within eddies has reduced the heating over West-Africa and the West-Pacific. Influence of Asian NO_x emissions on ozone heating rates (mean for June-September $\sim 0.0001 - 0.0012 \text{ K}\cdot\text{day}^{-1}$ for 38 % increase over India) in the upper troposphere (300-200 hPa) have been reported in the past (Roy et al., 2017). Near the tropopause ozone heating rates are positive ($0.001 - 0.005 \text{ K}\cdot\text{day}^{-1}$) except in the region of cross-tropopause transport (marked in **Figs. 8i-l**). The positive anomalies of ozone heating rates are associated with positive anomalies of ozone near the tropopause. The ECMWF dataset for 44 years (1958-2001) shows an inter-annual amplitude of the ozone heating rate $\pm 0.00025 \text{ K}\cdot\text{day}^{-1}$ near the tropopause over 30° S - 30° N (Wang et al., 2008).

4. Summary and Discussion

In this study, we show evidence of eddy shedding from the ASM anticyclone to both its eastern and western edge, during 1-8 July 2003 based on MIPAS satellite observations and ERA Interim re-analysis data as well as the associated transport patterns of trace gases from the ASM region to remote regions. The transport diagnostic based on ERA-Interim data shows that eddy shedding events are associated with RWB in the subtropical westerly jet. The RWB feature moves eastward in the subtropical westerly jet. Initially, during 1-5 July 2003, RWB occurs in the western part of the anticyclone and then sheds over West-Africa (20° - 30° N , 0° - 30° E). Later, during 5-8 July 2003, RWB moves to the eastern part of the anticyclone and

400 sheds an eddy over the West-Pacific (20°-30° N; 120°-150° E). Analysis of ERA-Interim PV
401 data for the last two decades (1995-2016) shows that the occurrence frequency of eddy
402 shedding from the ASM anticyclone over West-Africa is ~68 % and ~25 % over the West-
403 Pacific. In the UTLS (300-100 hPa), eddies (PV<2 PUV) over West-Africa/West-Pacific
404 shows highest correlation with the PV in the anticyclone after accounting for 3-4 days/5-6days
405 of lag. This indicates that the anticyclone sheds eddies with transport duration of typically
406 three to four days to West Africa and five-six days to the Western Pacific.

407 We employ the chemistry climate model ECHAM5-HAMMOZ to investigate transport of
408 Asian boundary layer trace gases (CO, ozone and PAN) into the monsoon anticyclone and the
409 associated eddies. The model simulations show that Asian trace gases transported into the
410 monsoon anticyclone are further carried away horizontally towards West-Africa and the West-
411 Pacific by eddies which detach from the anticyclone. These eddies protrude down to ~200 hPa
412 over West-Africa and ~500 hPa over the West Pacific. They re-distribute Asian trace gases
413 downward into the troposphere over these regions. Moreover, part of this air-mass is also
414 transported upward into the lower stratosphere. A higher frequency of eddy shedding over
415 West-Africa (68 %) during the last two decades (1995-2016) indicates a greater influence of
416 Asian trace gases on the UTLS over West-Africa than the West-Pacific over this period.

417 We evaluate the impact of Asian NO_x and NMVOCs emissions on ozone and PAN in
418 the regions of the ASM anticyclone and the associated eddies. The model sensitivity
419 simulations for a 10 % reduction in Asian emissions of NMVOCs and NO_x indicate significant
420 reduction (~2-10 %) in the concentration of PAN in the UTLS (300-80 hPa) over West-Africa

and the West-Pacific. The vertical distribution of anomalies of PAN shows negative values along the transport pathways, i.e., rising from the Asian region (80°-120° E) into the upper troposphere and both westward and eastward transport towards the region of eddy shedding. Tropospheric ozone (1000-180 hPa) shows a decrease of up to -4.5 % in response to a 10 % decrease in Asian emissions of NMVOCs and NO_x, while positive ozone anomalies (up to 8 %) are seen near the tropopause. In general, negative ozone anomalies in response to 10% reduction of NO_x and NMVOCs in the region of convective transport are seen in Figs. 8i-l. However, positive anomalies of ozone are observed near the tropopause (except in the region of cross tropopause transport) which may be due to reduction in the efficiency of ozone loss induced by reduced of NO_x and OH in the Asia-10 simulations and changes in dynamics due to emission changes, e.g. stratospheric intrusions and lightning. Mixing of tropospheric and stratospheric air-masses near the tropopause generates ozone variability. However, such an analysis is beyond the scope of the paper.

Our analysis indicates that transport of Asian trace gases from the anticyclone to West-Africa and the West-Pacific via eddies causes a change in the chemical composition of the UTLS and may therefore impact the radiative balance of the UTLS. We also estimate that a 10 % reduction in Asian NMVOCs and NO_x emissions leads to a decrease of ozone heating rates of 0.001 to 0.004 K·day⁻¹ in the region of transport into the troposphere and an increase of 0.001 to 0.005 K·day⁻¹ near the tropopause and lower stratosphere (180-50 hPa) over Asia (20°-150° E; 20°-40° N). Previous studies show that ozone changes in the lower stratosphere have the largest impact on the ozone radiative forcing (Riese et al., 2012). Interestingly, in the upper troposphere (200-300 hPa) negative anomalies of ozone heating rates (~0.001-0.003 K·day⁻¹)

are seen in the region of eddy shedding over West-Africa and the West-Pacific. Thus transport of Asian air masses via eddies eventually alters the heating rates in the UTLS in the regions of eddy shedding and may thus affect radiative forcing and local temperature. However, such questions are beyond the scope of this study. It should be noted that the distributions of MIPAS concentration fields look different from those of ERA-Interim and ECHAM5-HAMMOZ.. The ozone heating rates estimated from the model simulations will vary accordingly. Notwithstanding, we suggest further scrutiny of long range transport of Asian trace gases via eddies shedding from the anticyclone and its impact on ozone heating rates in the respective regions.

Acknowledgements: Dr. S. Fadnavis and C. Roy acknowledge with gratitude of Prof. Ravi Nanjundiah, Director of IITM, for his encouragement during the course of this study. We are grateful to B. Vogel for helpful discussions. This work was partly funded by the European Community's Seventh Framework Programme (FP7/2007–2013) as part of the StratoClim project (grant agreement no. 603557). We thank the European Centre for Medium-Range Weather Forecasts (ECMWF) for providing meteorological data sets. The authors are also thankful to Dr. Bernd Funke, Dr. Michael Kiefer and Dr. Gabriele Stiller, Karlsruhe Institute of Technology, Germany, for providing MIPAS data and for helpful discussions. We thank the anonymous reviewers for their valuable suggestions.

462 **References:**

- 463 Bourassa, A. E., Robock, A., Randel, W. J., Deshler, T., Rieger, L. A., Lloyd, N. D.,
 464 Llewellyn, E. J. and Degenstein, D. A.: Large volcanic aerosol load in the stratosphere
 465 linked to Asian monsoon transport, *Science*, 336(6090), 78–81,
 466 doi:10.1126/science.1219371, 2012.
- 467 Edwards, J. M. and Slingo, A.: Studies with a flexible new radiation code. I: Choosing a
 468 configuration for a large-scale model, *Q. J. R. Meteorol. Soc.*, 122(531), 689–719,
 469 doi:10.1256/smsqj.53106, 1996.
- 470 Fadnavis, S., Semeniuk, K., Pozzoli, L., Schultz, M. G., Ghude, S. D., Das, S. and Kakatkar,
 471 R.: Transport of aerosols into the UTLS and their impact on the asian monsoon region
 472 as seen in a global model simulation, *Atmos. Chem. Phys.*, 13(17), 8771–8786,
 473 doi:10.5194/acp-13-8771-2013, 2013.
- 474 Fadnavis, S., Schultz, M. G., Semeniuk, K., Mahajan, A. S., Pozzoli, L., Sonbawne, S., Ghude,
 475 S. D., Kiefer, M. and Eckert, E.: Trends in peroxyacetyl nitrate (PAN) in the upper
 476 troposphere and lower stratosphere over southern Asia during the summer monsoon
 477 season: Regional impacts, *Atmos. Chem. Phys.*, 14(23), 12725–12743,
 478 doi:10.5194/acp-14-12725-2014, 2014.
- 479 Fadnavis, S., Semeniuk, K., Schultz, M. G., Kiefer, M., Mahajan, A., Pozzoli, L. and
 480 Sonbawane, S.: Transport pathways of peroxyacetyl nitrate in the upper troposphere
 481 and lower stratosphere from different monsoon systems during the summer monsoon
 482 season, *Atmos. Chem. Phys.*, 15(20), 11477–11499, doi:10.5194/acp-15-11477-2015,
 483 2015.
- 484 Fadnavis, S. and Chattopadhyay, R.: Linkages of subtropical stratospheric intraseasonal
 485 intrusions with Indian summer monsoon deficit rainfall, *J. Clim.*, 30(13), 5083–5095,
 486 doi:10.1175/JCLI-D-16-0463.1, 2017.
- 487 Fischer, H., Birk, M., Blom, C., Carli, B., Carlotti, M., von Clarmann, T., Delbouille, L.,
 488 Dudhia, A., Ehhalt, D., Endemann, M., Flaud, J. M., Gessner, R., Kleinert, A.,
 489 Koopman, R., Langen, J., López-Puertas, M., Mosner, P., Nett, H., Oelhaf, H., Perron,
 490 G., Remedios, J., Ridolfi, M., Stiller, G. and Zander, R.: MIPAS: An instrument for
 491 atmospheric and climate research, *Atmos. Chem. Phys.*, 8(8), 2151–2188,
 492 doi:10.5194/acp-8-2151-2008, 2008.
- 493 Funke, B., López-Puertas, M., García-Comas, M., Stiller, G. P., von Clarmann, T., Höpfner,
 494 M., Glatthor, N., Grabowski, U., Kellmann, S., and Linden, A.: Carbon monoxide
 495 distributions from the upper troposphere to the mesosphere inferred from 4.7 μm non-
 496 local thermal equilibrium emissions measured by MIPAS on Envisat, *Atmos. Chem.*
 497 *Phys.*, 9, 2387–2411, doi: 10.5194/acp-9-2387-2009, 2009.

498 Garny, H. and Randel, W. J.: Transport pathways from the Asian monsoon anticyclone to the
 499 stratosphere, *Atmos. Chem. Phys.*, 16(4), 2703–2718, doi: 10.5194/acp-16-2703-2016,
 500 2016.

501 Ghude, S. D., Kulkarni, S. H., Jena, C., Pfister, G. G., Beig, G., Fadnavis, S. and van Der, R.
 502 J.: Application of satellite observations for identifying regions of dominant sources of
 503 nitrogen oxides over the indian subcontinent, *J. Geophys. Res. Atmos.*, 118(2), 1075–
 504 1089, doi: 10.1029/2012JD017811, 2013.

505 Gilford, D. M. and Solomon, S.: Radiative effects of stratospheric seasonal cycles in the
 506 tropical upper troposphere and lower stratosphere, *J. Clim.*, 30(8), 2769–2783,
 507 doi:10.1175/JCLI-D-16-0633.1, 2017.

508 Glatthor, N., von Clarmann, T., Fischer, H., Funke, B., Grabowski, U., Höpfner, M.,
 509 Kellmann, S., Kiefer, M., Linden, A., Milz, M., Steck, T., and Stiller, G. P.: Global
 510 peroxyacetyl nitrate (PAN) retrieval in the upper troposphere from limb emission
 511 spectra of the Michelson Interferometer for Passive Atmospheric Sounding (MIPAS),
 512 *Atmos. Chem. Phys.*, 7, 2775– 2787, doi: 10.5194/acp-7-2775-2007, 2007.

513 Horowitz, L. W., Walters, S., Mauzerall, D. L., Emmons, L. K., Rasch, P. J., Granier, C., Tie,
 514 X., Lamarque, J.-F., Schultz, M. G., Tyndall, G. S., Orlando, J. J. and Brasseur, G. P.:
 515 A global simulation of tropospheric ozone and related tracers: Description and
 516 evaluation of MOZART, version 2, *J. Geophys. Res. Atmos.*, 108(D24), doi:
 517 10.1029/2002JD002853, 2003.

518 Hsu, C. J. and Plumb, R. A.: Nonaxisymmetric thermally driven circulations and upper-
 519 tropospheric monsoon dynamics, *J. Atmos. Sci.*, 57(1977), 1255–1276,
 520 doi:https://doi.org/10.1175/1520-0469(2000)057<1255:NTDCAU>2.0.CO;2, 2000.

521 Li, M., Zhang, Q., Streets, D. G., He, K. B., Cheng, Y. F., Emmons, L. K., Huo, H., Kang, S.
 522 C., Lu, Z., Shao, M., Su, H., Yu, X., and Zhang, Y., Mapping Asian anthropogenic
 523 emissions of non-methane volatile organic compounds to multiple chemical
 524 mechanisms, *Atmos. Chem. Phys.*, 14, 5617–5638, 2014, doi:10.5194/acp-14-5617-
 525 2014.

526 Liang, Q., Jaeglé, L., Jaffe, D. A., Weiss-Penzias, P., Heckman, A. and Snow, J. A.: Long-
 527 range transport of Asian pollution to the northeast Pacific: Seasonal variations and
 528 transport pathways of carbon monoxide, *J. Geophys. Res. D Atmos.*, 109(23), 1–16,
 529 doi:10.1029/2003JD004402, 2004.

530 Nützel, M., Dameris, M., and Garny, H.: Movement, drivers and bimodality of the South
 531 Asian High, *Atmos. Chem. Phys.*, 16, 14755–14774, https://doi.org/10.5194/acp-16-
 532 14755-2016, 2016.

533 Ohara, T., Akimoto, H., Kurokawa, J., Horii, N., Yamaji, K., Yan, X. and Hayasaka, T.: An
 534 Asian emission inventory of anthropogenic emission sources for the period 1980–2020,

535 Atmos. Chem. Phys., 7, 4419–4444, doi:10.5194/acp-7-4419-2007, 2007.

552 Park, M., W. J. Randel, L. K. Emmons, P. F. Bernath, K. A. Walker, and C. D. Boone,
 553 Chemical isolation in the Asian monsoon anticyclone observed in Atmospheric
 554 Chemistry Experiment (ACE-FTS) data, Atmos. Chem. Phys., 8, 757 – 764, 2008.

555 Payne, V. H., Fischer, E. V., Worden, J. R., Jiang, Z., Zhu, L., Kurosu, T. P. and Kulawik, S.
 556 S.: Spatial variability in tropospheric peroxyacetyl nitrate in the tropics from infrared
 557 satellite observations in 2005 and 2006, Atmos. Chem. Phys., 17(10), 6341–6351,
 558 doi:10.5194/acp-17-6341-2017, 2017.

559 Ploeger, F., Gottschling, C., Griessbach, S., Grooß, J.-U., Guenther, G., Konopka, P., Müller,
 560 R., Riese, M., Stroh, F., Tao, M., Ungermann, J., Vogel, B. and von Hobe, M.: A
 561 potential vorticity-based determination of the transport barrier in the Asian summer
 562 monsoon anticyclone, Atmos. Chem. Phys., 15(22), 13145–13159, doi:10.5194/acp-
 563 15-13145-2015, 2015.

564 Ploeger, F., Konopka, P., Walker, K. and Riese, M.: Quantifying trace gases transport from the
 565 Asian monsoon anticyclone into the lower stratosphere, Atmos. Chem. Phys., 17(11),
 566 7055–7066, doi: 10.5194/acp-17-7055-2017, 2017.

567 Popovic, J. M. and Plumb, R. A.: Eddy Shedding from the Upper-Tropospheric Asian
 568 Monsoon Anticyclone, J. Atmos. Sci., 58(1), 93–104, doi:https://doi.org/10.1175/1520-
 569 0469(2001)058<0093:ESFTUT>2.0.CO;2, 2001.

570 Pozzoli, L., Janssens-Maenhout, G., Diehl, T., Bey, I., Schultz, M. G., Feichter, J., Vignati, E.
 571 and Dentener, F.: Re-analysis of tropospheric sulfate aerosol and ozone for the period
 572 1980-2005 using the aerosol-chemistry-climate model ECHAM5-HAMMOZ, Atmos.
 573 Chem. Phys., 11(18), 9563–9594, doi:10.5194/acp-11-9563-2011, 2011.

574 Randel, W. J., Park, M., Emmons, L., Kinnison, D., Bernath, P., Walker, K. A., Boone, C., and
 575 Pumphrey, H.: Asian monsoon transport of trace gases to the stratosphere, Science,
 576 328, 611-613, 10.1126/science.1182274, 2010.

577 Rap A., Richards N.A.D.; Forster P.M.; Monks S.; Arnold S.R.; Chipperfield, M., Satellite
 578 constraint on the tropospheric ozone radiative effect, Geophys. Res. Lett., 42, 5074-
 579 5081, doi: 10.1002/2015GL064037, 2015.

580 Riese, M., F. Ploeger, A. Rap, B. Vogel, P. Konopka, M. Dameris, and P. Forster, Impact of
 581 uncertainties in atmospheric mixing on simulated UTLS composition and related
 582 radiative effects, J. Geophys. Res., 117, D16305, doi: 10.1029/2012JD017751, 2012.

583 Roeckner, E., Bauml, G., Bonaventura, L., Brokopf, R., Esch, M., Giorgetta, M., Hagemann,
 584 S., Kirchner, I., Kornblueh, L., Manzini, E., Rhodin, A., Schlese, U., Schulzweida, U.,
 585 and Tompkins, A.: The atmospheric general circulation model ECHAM5: Part 1, Tech.
 586 Rep. 349, Max Planck Institute for Meteorology, Hamburg, 2003.

587 Roxy, M. K., Ghosh, S., Pathak, A., Athulya, R., Mujumdar, M., Murtugudde, R., Terray, P.
588 and Rajeevan, M.: A threefold rise in widespread extreme rain events over central
589 India, *Nat. Commun.*, 8(1), 1–11, doi:10.1038/s41467-017-00744-9, 2017.

590 Roy, C., Fadnavis, S., Müller, R., Chaudhary, A. D. and Ploeger, F.: Influence of enhanced
591 Asian NO_x emissions on ozone in the upper troposphere and lower stratosphere
592 (UTLS) in chemistry climate model simulations, *Atmos. Chem. Phys.*, 17, 1297–1311,
593 doi: <https://doi.org/10.5194/acp-17-1297-2017>.

594 Santee, M. L., G. L. Manney, N. J. Livesey, M. J. Schwartz, J. L. Neu, and W. G. Read, A
595 comprehensive overview of the climatological composition of the Asian summer
596 monsoon anticyclone based on 10 years of Aura Microwave Limb Sounder
597 measurements, *J. Geophys. Res. Atmos.*, 122, 5491–5514, doi:[10.1002/2016JD026408](https://doi.org/10.1002/2016JD026408),
598 2017.

599 Schneider, P. and van Der A, R. J.: A global single-sensor analysis of 2002–2011 tropospheric
600 nitrogen dioxide trends observed from space, *J. Geophys. Res. Atmos.*, 117(16), 1–17,
601 doi: [10.1029/2012JD017571](https://doi.org/10.1029/2012JD017571), 2012.

602 Stier, P., Feichter, J., Kinne, S., Kloster, S., Vignati, E., Wilson, J., Ganzeveld, L., Tegen, I.,
603 Werner, M., Balkanski, Y., Schulz, M., Boucher, O., Minikin, A. and Petzold, A.: The
604 aerosol-climate model ECHAM5-HAM, *Atmos. Chem. Phys.*, 5, 1125–1156, doi:
605 [10.1029/2005-5-1125](https://doi.org/10.1029/2005-5-1125), 2005.

606 Streets, D. G., Yarber, K. F., Woo, J.-H. and Carmichael, G. R.: Biomass burning in Asia:
607 Annual and seasonal estimates and atmospheric emissions, *Global Biogeochem.*
608 *Cycles*, 17(4), doi:10.1029/2003GB002040, 2003.

609 Strode, S. A. and Pawson, S.: Detection of carbon monoxide trends in the presence of
610 interannual variability, *J. Geophys. Res. Atmos.*, 118(21), 12257–12273, doi:
611 [10.1002/2013JD020258](https://doi.org/10.1002/2013JD020258), 2013.

612 Strong, C. and Magnusdottir, G.: Tropospheric Rossby wave breaking and the NAO/NAM, *J.*
613 *Atmos. Sci.*, 65(9), 2861–2876, doi:10.1175/2008JAS2632.1, 2008.

614 Tereszchuk, K. A., Moore, D. P., Harrison, J. J., Boone, C. D., Park, M., Remedios, J. J.,
615 Randel, W. J. and Bernath, P. F.: Observations of peroxyacetyl nitrate (PAN) in the
616 upper troposphere by the Atmospheric Chemistry Experiment-Fourier Transform
617 Spectrometer (ACE-FTS), *Atmos. Chem. Phys.*, 13(11), 5601–5613, doi:10.5194/acp-
618 13-5601-2013, 2013.

619 Tie, X., Zhang, R., Brasseur, G. and Lei, W.: Global NO_x Production by Lightning, *J. Atmos.*
620 *Chem.*, 43(1), 61–74, doi:10.1023/A:1016145719608, 2002.

621 Ungermann, J., Ern, M., Kaufmann, M., Müller, R., Spang, R., Ploeger, F., Vogel, B. and
622 Riese, M.: Observations of PAN and its confinement in the Asian summer monsoon

623 anticyclone in high spatial resolution, *Atmos. Chem. Phys.*, 16(13), 8389–8403, doi:
624 10.5194/acp-16-8389-2016, 2016.

625 Uppala, S. M., Kållberg, P. W., Simmons, A. J., Andrae, U., da Costa Bechtold, V., Fiorino,
626 M., Gibson, J. K., Haseler, J., Hernandez, A., Kelly, G. A., Li, X., Onogi, K., Saarinen,
627 S., Sokka, N., Allan, R. P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C.
628 M., van de Berg, L., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A.,
629 Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Hólm, E., Hoskins, B. J.,
630 Isaksen, I., Janssen, P. A. E. M., Jenne, R., McNally, A. P., Mahfouf, J. F., Morcrette,
631 J. J., Rayner, N. A., Saunders, R. W., Simon, P., Sterl, A., Trenberth, K. E., Untch, A.,
632 Vasiljevic, D., Viterbo, P. and Woollen, J.: The ERA-40 re-analysis, *Q. J. R. Meteorol.*
633 *Soc.*, 131(612), 2961–3012, doi:10.1256/qj.04.176, 2005.

634 Verstraeten, W. W., Neu, J. L., Williams, J. E., Bowman, K. W., Worden, J. R. and Boersma,
635 K. F.: Rapid increases in tropospheric ozone production and export from China, *Nat.*
636 *Geosci.*, 8(9), 690–695, doi:10.1038/ngeo2493, 2015.

637 Vogel, B., Günther, G., Müller, R., Grooß, J.-U., Hoor, P., Krämer, M., Müller, S., Zahn, A.
638 and Riese, M.: Fast transport from Southeast Asia boundary layer sources to northern
639 Europe: Rapid uplift in typhoons and eastward eddy shedding of the Asian monsoon
640 anticyclone, *Atmos. Chem. Phys.*, 14(23), 12745–12762, doi:10.5194/acp-14-12745-
641 2014, 2014.

642 Vogel, B., Günther, G., Müller, R., Grooß, J.-U., Afchine, A., Bozem, H., Hoor, P., Krämer,
643 M., Müller, S., Riese, M., Rolf, C., Spelten, N., Stiller, G. P., Ungermann, J., and Zahn,
644 A.: Long-range transport pathways of tropospheric source gases originating in Asia
645 into the northern lower stratosphere during the Asian monsoon season 2012, *Atmos.*
646 *Chem. Phys.*, 16, 15301-15325, <https://doi.org/10.5194/acp-16-15301-2016>, 2016.

647 Vogel B., Müller R., Günther G. , Spang R., Hanumanthu S., Li D. , Riese M , and Stiller G.P.,
648 [Lagrangian simulations of the transport of young air masses to the top of the Asian](#)
649 [monsoon anticyclone and into the tropical pipe, submitted to ACPD, 2018.](#)

650 von Clarmann, T., De Clercq, C., Ridolfi, M., Höpfner, M. and Lambert, J. C.: The horizontal
651 resolution of MIPAS, *Atmos. Meas. Tech.*, 2(1), 47–54, doi:10.5194/amt-2-47-2009,
652 2009.

653 van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S. and Arellano,
654 Avelino F., J.: Interannual variability in global biomass burning emissions from 1997
655 to 2004, *Atmos. Chem. Phys.*, 6(11), 3423–3441, doi:10.5194/acpd-6-3175-2006,
656 2006.

657 Wang, W.-G., Yuan, M., Wang, H.-Y., Sun, J.-H., Xie, Y.-Q., Fan, W.-X. and Chen, X.-M., A
658 Study of Ozone Amount in the Transition Layer Between Troposphere and
659 Stratosphere and Its Heating Rate. *Chinese J. Geophys.*, 51: 916–930.
660 doi:10.1002/cjg2.1287, 2008.

661 Wayne, R. P.: Chemistry of atmospheres, 3rd Edn., Oxford science publications, Clarendon
662 Press, Oxford, 337, 2000.

663 Zhang, Q., Wu, G. and Qian, Y.: The Bimodality of the 100 hPa South Asia High and its
664 Relationship to the Climate Anomaly over East Asia in summer. J. Meteorol. Soc.
665 Japan, 80(4), 733–744, doi:10.2151/jmsj.80.733, 2002.

666 Zhao, C., Wang, Y., Choi, Y. and Zeng, T.: Summertime impact of convective transport and
667 lightning NO_x production over North America: Modeling dependence on
668 meteorological simulations, Atmos. Chem. Phys., 9(13), 4315–4327, doi: 10.5194/acp-
669 9-4315-2009, 2009.

670




Figure 1: Spatial distribution of potential vorticity (PVU) ($1 \text{ PVU} = 10^{-6} \text{ K m}^2 \text{ kg}^{-1} \text{ s}^{-1}$) (color shades) at 370 K potential temperature surface and wind anomalies at 200 hPa from ERA-Interim reanalysis for (a) 01 July, (b) 02 July, (c) 03 July, (d) 04 July, (e) 05 July, (f) 06 July, (g) 07 July, (h) 08 July, 2003. Wind vectors are represented by black arrows ($\text{m}\cdot\text{s}^{-1}$). Eddies are shown with white circles.




Figure 2: Spatial distribution of potential vorticity (PVU) (color shades) at 350 K potential temperature surface and wind anomalies in m s^{-1} (thin black vectors) at 200 hPa from ERA-Interim reanalysis for (a) 01 July, (b) 02 July, (c) 03 July, (d) 04 July, (e) 05 July, (f) 06 July, (g) 07 July, (h) 08 July, 2003. The events of RBW-1, RBW-2 and RBW-3 are indicated by solid black, red and blue arrows, respectively.




Figure 3: Power spectral analysis of ERA-Interim PV averaged for 100-300 hPa and in June-September during 1995-2015 (a) West-Africa (20-30° N, 0-30° E) and (b) West-Pacific (20-30° N, 140-150° E) and lag-lead Pearson correlation coefficient of PV in the monsoon anticyclone (85-90° E, 28-30° N) with (c) West-Pacific (20-30° N, 140-150° E), (d) West Africa (20-30° N, 0-30° E). In Fig. a-b dotted green line indicates spectrum and blue and red line indicates 5% and 95% confidence levels respectively for lag-1 autocorrelation. Any spectral peak above red line is statistically significant at 95% confidence level.




Figure 4: Spatial distribution of ozone mixing ratios (ppb) (color shades) corresponding to MIPAS satellite observations at 16 km for (a) 1-2 July, (b) 3-4 July, (c) 5-6 July, (d) 7-8 July, 2003; ERA-Interim reanalysis at 100 hPa for (e) 2 July, (f) 4 July, (g) 6 July, (h) 8 July, 2003, and ECHAM5-HAMMOZ CTRL simulations at 16 km for (i) 2 July, (j) 4 July, (k) 6 July, (l) 8 July, 2003. Black arrows in panels (e)-(h) show wind anomalies (m s^{-1}) at 200 hPa. Minimum ozone amounts near the location of eddies are shown with black circles.




Figure 5: Spatial distribution of CO mixing ratios (ppb) at 16 km: MIPAS satellite observations for (a) 1-2 July, (b) 3-4 July, (c) 5-6 July, (d) 7-8 July, 2003 and ECHAM5-HAMMOZ CTRL simulations for (e) 02 July, (f) 04 July, (g) 06 July, (h) 08 July, 2003. Maximum CO amounts near the location of eddies are shown with black circles.




Figure 6: Spatial distribution of PAN mixing ratios (ppt) at 16 km: MIPAS satellite observations for (a) 1-2 July, (b) 3-4 July, (c) 5-6 July, (d) 7-8 July, 2003, and ECHAM5-HAMMOZ CTRL simulations for (e) 02 July, (f) 04 July, (g) 06 July, (h) 08 July, 2003. Maximum PAN amounts near the location of eddies are shown with black circles.




Figure 7: Longitude-pressure section (averaged for 20°-40° N) of CO (ppb) from ECHAM5-HAMMOZ CTRL simulation for (a) 02 July, (b) 04 July, (c) 06 July, (d) 08 July, 2003. (e)-(h) same as (a)-(d) but for PAN (ppt). Black thick line indicates the tropopause and black dotted circles indicate maximum in CO and PAN amounts near eddies. Pressure (hPa) is indicated on left y-axis and altitudes (km) on the right y-axis. Wind vectors ($\text{m}\cdot\text{s}^{-1}$) are shown by black arrows. Vertical velocity field is scaled by a factor of 300.




Figure 8: Spatial distribution of anomalies (Asia10-CTRL) of PAN mixing ratios (ppt) (color shades) at 16 km from ECHAM5-HAMMOZ model simulations for (a) 02 July, (b) 04 July, (c) 06 July, (d) 08 July, 2003. Longitude-pressure distribution (averaged for 20°-40° N) of anomalies of PAN (%) for (e) 02 July, (f) 04 July, (g) 06 July, (h) 08 July, 2003. (i)-(l) same as (e)-(h) but for ozone anomalies (%) (averaged for 18°-20° N). Black thick line indicates the tropopause. Pressure (hPa) is indicated on left y-axis and altitudes (km) on right y-axis. Black boxes in the bottom panels indicate regions of cross tropopause transport.




Figure 9: Longitude-pressure distribution (averaged for 18°-20° N) of anomalies of ozone heating rates $((K \cdot day^{-1}) \times 10^{-2})$ for (a) 02 July, (b) 04 July, (c) 06 July, (d) 08 July, 2003. Pressure (hPa) is indicated on left y-axis and altitudes (km) on right y-axis. The black thick line indicates the tropopause.