

Modeling the effect of non-ideality, dynamic mass transfer and viscosity on SOA formation in a 3-D air quality model

Youngseob Kim¹, Karine Sartelet¹, and Florian Couvidat²

¹CEREA, Joint Laboratory École des Ponts ParisTech / EDF R&D, Université Paris-Est, 77455 Champs-sur-Marne, France

²Institut National de l'Environnement Industriel et des Risques, 60550 Verneuil-en-Halatte, France

Correspondence to: Y. Kim (youngseob.kim@enpc.fr)

Abstract.

In this study, assumptions (ideality and thermodynamic equilibrium) commonly made in 3-dimensional (3-D) air quality models were reconsidered to evaluate their impacts on secondary organic aerosol (SOA) formation over Europe.

To investigate the effects of non-ideality, dynamic mass transfer and aerosol viscosity on the SOA formation, the Secondary Organic Aerosol Processor (SOAP) model was implemented in the 3-D air quality model Polyphemus. This study presents the first 3-D modeling simulation which describes the impact of aerosol viscosity on the SOA formation. The model uses either the equilibrium approach or the dynamic approach with a method specially designed for 3-D air quality models to solve efficiently particle-phase diffusion when particles are viscous.

Sensitivity simulations using two organic aerosol models implemented in Polyphemus to represent mass transfer between gas and particle phases show that the computation of the absorbing aerosol mass strongly influences the SOA formation. In particular, taking into account the concentrations of inorganic aerosols and hydrophilic organic aerosols in the absorbing mass of the aqueous phase increases the average SOA concentration by 5% and 6%, respectively. However, inorganic aerosols influence the SOA formation not only because they constitute an absorbing mass for hydrophilic SOA, but also because they interact with organic compounds. Non-ideality (short, medium and long-range interactions) was found to influence SOA concentrations by about 30%.

Concerning the dynamic mass transfer for the SOA formation, if the viscosity of SOA is not taken into account and if ideality of aerosols is assumed, the dynamic approach is found to give generally similar results than the equilibrium approach (indicating that equilibrium is an efficient hypothesis for inviscid and ideal aerosols). However, when a non-ideal aerosol is assumed, taking into account the dynamic mass transfer leads to a decrease of concentrations of the hydrophilic compounds (compared to equilibrium). This decrease is due to differences in the values of activity coefficients, which are different between values computed for bulk aerosols and those for each size section. This result indicates the importance of non-ideality on the dynamic evolution of SOA.

For viscous aerosols, assuming a highly viscous organic phase leads to an increase in SOA concentrations during daytime (by preventing the evaporation of the most volatile organic compounds). The partitioning of non-volatile compounds is not affected by viscosity, but the aging of more volatile compounds (that leads to the formation of the less volatile compounds) slows down as the evaporation of those compounds is stopped due to the viscosity of the particle. These results imply that

aerosol concentrations may deviate significantly from equilibrium as the gas-particle partitioning could be higher than predicted by equilibrium. Furthermore, although a compound evaporates in the simulation using the equilibrium approach, the same compound can condense in the simulation using the dynamic approach if the particles are viscous.

The results of this study emphasize the need for 3-D air quality models to take into account the effect of non-ideality on SOA formation and the effect of aerosol viscosity for the more volatile fraction of semi-volatile organic compounds.

1 Introduction

Inorganic and organic aerosols constitute an important fraction of aerosols (Putaud et al., 2004; Jimenez et al., 2009), which influence climate and health (Rattanavaraha et al., 2011; Boucher, 2015; Shrivastava et al., 2017). A large fraction of inorganic and organic aerosols are not directly emitted, but they are formed in the atmosphere by the condensation of condensable compounds, which are often semi volatile, i.e. they exist both in the gas and in the particle phases. The modeling of the mass transfer of condensable compounds (inorganic and organic) from the gas phase to the particle phase is important, because it determines the fraction of the condensable compounds in the particle phase, and therefore the particle concentration. It is usually modeled by three approaches: the dynamic approach, the equilibrium approach, and the hybrid approach. In the dynamic approach, the mass transfer between the gas and the particle phases is explicitly calculated by solving the mass flux equation (e.g., Wexler and Seinfeld, 1990; Bowman et al., 1997). In the equilibrium approach, instantaneous equilibrium is assumed between the gas and particle phases (Pankow, 1994). The dynamic approach provides a more accurate representation of the gas/particle mass transfer but is computationally more expensive than the equilibrium approach. The hybrid approach combines these two approaches. As gas-phase molecules condense more rapidly on fine than on coarse particles (and therefore reach more rapidly equilibrium with fine particles), the mass transfer is explicitly computed for coarse particles using the dynamic approach and instantaneous equilibrium is assumed for fine particles in the hybrid approach (Capaldo et al., 2000; Zhang et al., 2004; Debry et al., 2007).

Several previous studies showed that organic aerosols can be highly viscous (Virtanen et al., 2010; Cappa and Wilson, 2011; Pfrang et al., 2011; Shiraiwa et al., 2011; Vaden et al., 2011; Shiraiwa and Seinfeld, 2012; Abramson et al., 2013; Renbaum-Wolff et al., 2013; Shiraiwa et al., 2013; Reid et al., 2018). The diffusion of organic compounds from the particle surface inside the particle is influenced by the viscosity of organic aerosols, which depends on relative humidity and aerosol composition (Song et al., 2016a; O'Meara et al., 2017). The diffusion is very slow when the particle phase state is semi-solid, solid or glassy solid (Shiraiwa et al., 2017). By influencing the diffusion inside the particle, viscosity influences the mass transfer between gas and particle phases, which is much slower than for non-viscous particles (Shrivastava et al., 2017). Several models explicitly treat the particle-phase diffusion of organic compounds (Shiraiwa et al., 2012; Roldin et al., 2014). However, the use of these models is limited in 3-dimensional (3-D) air quality models because particles need to be discretized with a high number of particle layers, which leads to an expensive computational cost.

Although the equilibrium approach is widely used in 3-D air quality models because of its computational efficiency, the dynamic approach is also sometimes used in 3-D air quality models for inorganic aerosols (Jacobson, 1997; Meng et al., 1998;

Sun and Wexler, 1998; Sartelet et al., 2007a). However, to our knowledge, the impact of viscosity of particles on gas/particle phase partitioning and organic aerosol concentrations has yet not been taken into account in 3-D air quality models.

The mass transfer of condensable organic compounds between the gas and particle phases is influenced by interactions with other compounds. The activity coefficients reflect the non-ideality of aerosols and the influence of the interactions between
5 compounds on the mass transfer between the gas and particle phases.

Organic aerosol models often assume ideality, and they do not take into account the influence of activity coefficients on the formation of secondary organic aerosols (SOA). Activity coefficients may be determined by thermodynamic models. For example, the UNIQUAC Functional-group Activity Coefficients (UNIFAC) thermodynamic model (Fredenslund et al., 1975) is based on a functional group method, which estimates short-range activity coefficients (interactions between uncharged
10 molecules) by using the structure of the molecules present in the particles. However, in the aqueous phase, for hydrophilic organic compounds, due to the presence of ions, such as inorganic ions, medium and long-range activity coefficients (resulting from electrostatic interactions) may also influence activity coefficients. These medium and long-range activity coefficients are described by the Aerosol Inorganic–Organic Mixtures Functional groups Activity Coefficient AIOMFAC model (Zuend et al., 2008). The effect of activity coefficients was already investigated by a previous study (Couvidat et al., 2012) by using
15 the UNIFAC model. Compared to assuming ideality, computing activity coefficients was found to decrease the concentrations of hydrophobic SOA (condensing onto the organic phase of particles) but also to increase the concentrations of hydrophilic SOA (condensing onto the aqueous phase of particles). AIOMFAC and UNIFAC are used in this study to compute the activity coefficients for organic-inorganic mixture. These models have been developed using a group contribution method. 18 main functional groups and 45 subgroups in AIOMFAC are used in this study. The SOA surrogates are splitted into these functional
20 groups. The computation of activity coefficients depends on the functional groups that are present in the SOA surrogates. It is assumed here that the SOA surrogates represent the major SOA compound types in terms of functional groups. However, considering more compounds in the model may affect the computation of activity coefficients, and enhance their effect as a stronger variability of composition would be simulate. Pye et al. (2017) obtained a reduction of the bias in SOA for routine monitoring stations taking into account the non-ideality via activity coefficients. However, this study did not take into account
25 the effect of interactions between inorganic and organic compounds.

The Secondary Organic Aerosol Processor (SOAP) model (Couvidat and Sartelet, 2015) has been developed to represent the condensation/evaporation of organic aerosols using both the equilibrium and dynamic approaches. The SOAP model was designed to be implemented in 3-D air quality models and can be used to implicitly represent the diffusion of organic com-
30 pounds inside the particle phase, using a low number of particle layers. Compared to an explicit representation of the diffusion of organic compounds with a high number of particle layers, the SOAP model showed good agreements of modeled organic concentrations of viscous particles, using a lower number of aerosol layers.

In this study, we present the implementation of the SOAP model in the 3-D air quality model Polyphemus, and present differences between SOAP and the hydrophilic/hydrophobic organic (H²O) model (Couvidat et al., 2012) and how the new processes implemented in the SOAP model influence SOA formation (absorbing mass, non-ideality, viscosity). Jathar et al.
35 (2016) showed that organic-phase water uptake leads to an increase in total organic aerosol concentration. Water uptake is

taken into account in the SOAP model to estimate the absorbing mass. We estimate for the first time in a 3-D air quality model the maximum influence of aerosol viscosity on particle organic concentrations over Europe. To do so, we compare simulations assuming inviscid or extremely viscous aerosols. We also estimate the influence of non-ideality, in particular the influence of inorganic concentrations via medium and long-range activity coefficients on SOA concentrations. The SOAP model and differences with H²O, the previously used SOA model in Polyphemus, are described in section 2. Section 3 details the modeling of the newly added processes studied here: medium and long-range activity coefficients, aerosol dynamic, viscosity. Finally, section 4 presents the simulations, and the sensitivity to these processes.

2 Description and implementation of SOAP in Polyphemus

The SOAP model was implemented in the chemistry transport model Polair3D (Sartelet et al., 2007b) of the air quality platform Polyphemus version 1.8 (Mallet et al., 2007). The aerosol dynamics (coagulation, condensation/evaporation) is modeled with the Size REsolved Aerosol Model (SIREAM, Debry et al. (2007)). The particle size distribution is divided into sections, each section corresponding to a range of diameters. Similarly to H²O, the SOAP model is based on the molecular surrogate approach. It distinguishes hydrophobic compounds from hydrophilic compounds.

In the molecular surrogate approach, organic compounds are represented by surrogates, which are model compounds chosen depending on their sources (anthropogenic vs biogenic) and their properties, such as their affinity with water (hydrophilic vs hydrophobic) and their volatility. Oxidation of the SOA precursors differs depending on the regime of nitrogen oxides (NO_x) (low NO_x regime vs high NO_x regime). Different reactions (Kim et al., 2011) were added to the gas-phase chemistry model of Polyphemus (CB05 is used here, Yarwood et al. (2005)) to model the formation of organic compounds from five classes of SOA precursors (intermediate and semi-volatile organic compounds of anthropogenic emissions, aromatic compounds, isoprene, monoterpenes and sesquiterpenes). As detailed in Couvidat et al. (2012), surrogates from anthropogenic precursors are mostly hydrophobic, while those from biogenic precursors are mostly hydrophilic. Table 1 summarizes the surrogates and their properties (volatility and hydrophilicity).

In previous studies using Polyphemus (Couvidat et al., 2012; Sartelet et al., 2012; Couvidat et al., 2013; Zhu et al., 2016), the H²O model was used to partition organics between the gas and particle phases: instantaneous equilibrium was assumed between the gas and particle phases, and only short-range activity coefficients were taken into account. They were computed with UNIFAC (Fredenslund et al., 1975), i.e. without taking into account the impact of inorganic compounds, as if the aqueous phase is only constituted of water and organics. In other words, H²O only takes into account solvents into the computation of short range interactions and H²O implicitly assumed that organics have no interaction with inorganics.

However, AIOMFAC (Zuend et al., 2008) is a thermodynamic model designed for the calculation of activity coefficients of different chemical species in inorganic-organic mixtures. It takes into account the short-range, middle-range and long-range interactions between molecules and ions. For short-range interactions, AIOMFAC differs from UNIFAC because it takes inorganics into account in short-range interactions by taking relative van der Waals subgroup volume and surface area

Table 1. Description of the SOA surrogate compounds (Couvidat and Sartelet, 2015).

Surrogate	Type*	Precursors	Conditions of formation [†]	Volatility [‡]
BiMT	A	isoprene	Oxidation by OH (low NO _x)	high
BiPER	A	isoprene	Oxidation by OH (low NO _x)	high
BiDER	A	isoprene	Oxidation by OH (low NO _x)	medium
BiMGA	A	isoprene	Oxidation by OH (high NO _x)	medium
BiNGA	B	isoprene	Oxidation by OH (high NO _x)	high
BiNIT3	B	isoprene	Oxidation by NO ₃	high
BiA0D	A	monoterpenes	Oxidation by OH and O ₃	very low if the aqueous aerosol is acidic
BiA1D	A	monoterpenes	Oxidation by OH and O ₃	medium
BiA2D	A	monoterpenes	Oxidation by OH and O ₃	medium
BiNIT	B	monoterpenes	Oxidation by NO ₃	high
BiBIP	B	sesquiterpenes	Oxidation by OH	very low
BiBmP	B	sesquiterpenes	Oxidation by OH	medium
AnBIP	B	aromatics	Oxidation by OH (low NO _x)	low
AnBmP	B	aromatics	Oxidation by OH (low NO _x)	high
AnCIP	B	aromatics	Oxidation by OH (high NO _x)	non-volatile
POAIP	B	-	Primary SVOC	low
POAmP	B	-	Primary SVOC	high
POAhP	B	-	Primary SVOC	very high
SOAIP	B	POAIP	Oxidation by OH	very low
SOAmP	B	POAmP	Oxidation by OH	low
SOAhP	B	POAhP	Oxidation by OH	high

*: Type A and B correspond to hydrophilic and hydrophobic compounds, respectively.

[†]: hydroxyl radical (OH), nitrogen oxides (NO_x), ozone (O₃) and semi-volatile organic compounds (SVOC).

[‡]: very low for compounds with $K_p > 100 \text{ m}^3 \mu\text{g}^{-1}$, low for compounds with $100 \text{ m}^3 \mu\text{g}^{-1} \geq K_p > 1 \text{ m}^3 \mu\text{g}^{-1}$, medium for compounds with $1 \text{ m}^3 \mu\text{g}^{-1} \geq K_p > 0.1 \text{ m}^3 \mu\text{g}^{-1}$, high for compounds with $0.1 \text{ m}^3 \mu\text{g}^{-1} \geq K_p > 0.01 \text{ m}^3 \mu\text{g}^{-1}$ and very high for compounds with $K_p \leq 0.01 \text{ m}^3 \mu\text{g}^{-1}$.

UNIFAC parameters. It assumes that interaction parameters of inorganics with organics for short-range interactions are zero, i.e., the short range organic-inorganic interactions are ideal.

The SOAP model inherits all the characteristics of the H₂O model and new processes (such as modeling of inorganic-organic interactions via activity coefficients, and dynamic evolution of gas/particle partitioning) are added (Couvidat and Sartelet, 2015). However, SOAP differs from H₂O not only because of the possibility to model inorganic-organic interactions via activity coefficients, and to model dynamically the gas/particle partitioning of viscous aerosols, but also differences occur in the gas/particle partitioning due to the computation of the absorbing mass.

In SOAP and H²O, ideality is defined by reference to the pure state for hydrophobic compounds (activity coefficients are equal to one when the compound is pure) and to the infinite dilution state for hydrophilic compounds (activity coefficients are equal to one when the compounds is diluted into an infinite amount of water). The partitioning is computed according to Raoult's law for hydrophobic compounds and to Henry's law for hydrophilic compounds.

- 5 The differences between SOAP and H²O are now detailed and their impact on previously published simulations of Polyphemus/H²O is quantitatively assessed.

2.1 Gas-particle partitioning for the aqueous and organic phases

The equilibrium approach is used in the H²O model, and it can be used in the SOAP model. In this approach, the partitioning between the gas and particle organic phases is done following Pankow (1994):

$$10 \quad \frac{c_{p,i}}{c_{g,i}} = K_{p,i} c_p \quad (1)$$




Figure 1. Computation steps for the gas-particle partitioning and the water absorption.

where $K_{p,i}$ is the organic-phase gas/particle partitioning coefficient ($\text{m}^3/\mu\text{g}$), $c_{p,i}$ is concentration of the compounds i in the organic phase ($\mu\text{g}/\text{m}^3$), $c_{g,i}$ is the gas-phase concentration ($\mu\text{g}/\text{m}^3$), c_p is the total concentration of the particles in the organic phase ($\mu\text{g}/\text{m}^3$). Whereas in the H^2O model, c_p is only the concentration of the organic compounds in the particles, in the SOAP model the absorption of water by the organic phase, $c_{water,p}$ ($\mu\text{g}/\text{m}^3$), is also included in c_p .

- 5 The absorption of water by the organic phase is computed using Equation 2 following Couvidat and Sartelet (2015).

$$c_{water,p} = \frac{c_p M_{water} RH}{\gamma_{water,p} M_p} \quad (2)$$

where M_{water} is the molar mass of water (g/mol), RH is the relative humidity, $\gamma_{water,p}$ is the activity coefficient of water in the organic phase and M_p is the averaged molar mass of the organic phase (g/mol). Figure 1 shows the computation steps for the gas-particle partitioning and the water absorption.

- 10 The partitioning between the gas and the aqueous phases is done similarly as in the organic phase:

$$\frac{c_{aq,i}}{c_{g,i}} = K_{aq,i} c_{aq} \quad (3)$$

- where $c_{aq,i}$ is the aqueous-phase concentration of the compound i ($\mu\text{g}/\text{m}^3$), $K_{aq,i}$ is the aqueous-phase gas/particle partitioning coefficient ($\text{m}^3/\mu\text{g}$), and c_{aq} is the total concentration of the particles in the aqueous phase ($\mu\text{g}/\text{m}^3$). $K_{aq,i}$ is computed as detailed in Couvidat and Sartelet (2015) and depends on the activity coefficient. In the H^2O model, c_{aq} corresponds only to the liquid water content (LWC) calculated using a thermodynamic model, e.g., ISORROPIA (Nenes et al., 1999), for inorganic aerosols. However, c_{aq} includes inorganic aerosols, hydrophilic organic aerosols, and absorbed water by hydrophilic organic aerosols in addition to LWC in the SOAP model. The larger concentrations of c_{aq} in the SOAP model than in the H^2O model lead in return to larger compounds concentrations in the aqueous phase ($c_{aq,i}$).
- 15

2.2 Impact on SOA concentrations

- 20 Sensitivity simulations are performed to quantify the impact on organic concentrations of the differences between the H^2O and SOAP models in the formulation of the absorbing mass used in the modeling (c_p and c_{aq} : the total particle concentrations of the organic and aqueous phases respectively). Within the Polyphemus platform, the two SOA models are implemented with the Size Resolved Aerosol Model (SIREAM) aerosol module (Debry et al., 2007). The simulations of Couvidat et al. (2012) are rerun using the SOAP model instead of H^2O . The model configuration is detailed in Couvidat et al. (2012). The simulation domain covers Europe (see Figure 2) with a horizontal resolution of $0.5^\circ \times 0.5^\circ$ and 9 vertical levels (20 m, 80 m, 210 m, 550 m, 1150 m, 1950 m, 2950 m, 4750 m, 9000 m). The initial and boundary conditions are calculated using data from global models MOZART (gas) and ECHAM5-HAMMOZ (particles). Anthropogenic emissions are taken from the EMEP (European Monitoring and Evaluation Programme) inventory (<http://www.ceip.at/>) and biogenic emissions are estimated with MEGAN (Model of Emissions of Gases and Aerosols from Nature) (Guenther et al., 2006).
- 25

- 30 Sensitivity simulations are conducted for June 2002 and detailed in Table 2. Domain-averaged concentrations of SOA are used to compare the sensitivity simulations in Figure 3. In the SOAP-Reference simulation, c_{aq} is computed by considering




Figure 2. Simulation domain and location of observation stations.

the water absorbed by inorganic aerosols and by hydrophilic aerosols (inorganic aerosols and hydrophilic organic aerosols) while c_p is computed by considering hydrophobic organic aerosols and the water absorbed by hydrophobic organic aerosols. Overall, the average difference in SOA concentrations between the SOAP-Reference and H²O-Reference simulations is 15%. The differences between these two simulations are mostly due to the influence of the different compounds included in the absorbing mass used for the partitioning of the gas and particle phases, i.e. in the computation of c_p and c_{aq} (the total particle concentrations of the organic and aqueous phases). Simulations using the same absorbing mass in SOAP and H²O (water absorbed by inorganic aerosols for c_{aq} and hydrophobic organic aerosol for c_p) lead to similar concentrations (see the comparison of the simulations H²O-ideal and SOAP-ideal in Figure 3). Adding water absorbed by organic aerosols in the absorbing mass leads to a slight increase in SOA concentration (comparisons of the simulations SOAP-no_inorg and SOAP-no_water).

10 Adding inorganic aerosols in the absorbing mass of hydrophilic aerosols (c_{aq}) has a larger impact than the addition of water absorbed by organic aerosols (5%, see the comparison of the simulations SOAP-Reference and SOA-no_inorg). Adding organic aerosols in the absorbing mass of c_{aq} has an impact as large as inorganic aerosols (6%, see the comparison of the simulations SOAP-no_inorg and SOAP-basic).

Not only the absorbing mass strongly influences the SOA concentrations, but also the interactions between compounds, as modeled by activity coefficients. The influence of taking into account organic-organic interactions by short-range activity coefficients is as high as 18% (see the comparison between the simulations H²O-Reference and H²O-Ideal). The difference between the SOAP-Reference and SOAP-Ideal simulations is much larger (35%), because of differences in the computation of the absorbing mass between SOAP and H²O.

15

An additional sensitivity simulation SOAP-sr is used to estimate UNIFAC sensitivities when inorganic aerosols are added in the computation of short-range activity coefficient as in AIOMFAC. The averaged SOA concentrations in SOAP-sr increase by 15% compared to those of SOAP-Reference. This difference is further discussed in section 4.3. As inorganic-organic interaction parameters are set to zero in UNIFAC, taking into account inorganics in the computation of short-range activity coefficients

Table 2. List of the sensitivity simulations to compare the H²O and SOAP models.

Simulation name	SOA model	Aqueous-phase particle included in c_{aq}^{\dagger}	Organic-phase particle included in c_p^{\dagger}	Activity coefficient
SOAP-sr	SOAP	- Water absorbed by inorganic aerosol - Inorganic aerosol - Hydrophilic organic aerosol - Water absorbed by hydrophilic organic aerosol	- Hydrophobic organic aerosol - Water absorbed by hydrophobic organic aerosol	UNIFAC-sr**
SOAP-Reference	SOAP	- Water absorbed by inorganic aerosol - Inorganic aerosol - Hydrophilic organic aerosol - Water absorbed by hydrophilic organic aerosol	- Hydrophobic organic aerosol - Water absorbed by hydrophobic organic aerosol	UNIFAC
SOAP-no_inorg	SOAP	- Water absorbed by inorganic aerosol - Hydrophilic organic aerosol - Water absorbed by hydrophilic organic aerosol	- Hydrophobic organic aerosol - Water absorbed by hydrophobic organic aerosol	UNIFAC
SOAP-no_water	SOAP	- Water absorbed by inorganic aerosol - Hydrophilic organic aerosol	- Hydrophobic organic aerosol	UNIFAC
SOAP-basic	SOAP	- Water absorbed by inorganic aerosol	- Hydrophobic organic aerosol	UNIFAC
SOAP-ideal	SOAP	- Water absorbed by inorganic aerosol	- Hydrophobic organic aerosol	ideal*
H ² O-Reference	H ² O	- Water absorbed by inorganic aerosol	- Hydrophobic organic aerosol	UNIFAC
H ² O-Ideal	H ² O	- Water absorbed by inorganic aerosol	- Hydrophobic organic aerosol	ideal*

[†]: total particle concentration of the organic phase (c_p) and aqueous phase (c_{aq}).

*: ideal mixture, activity coefficient is set to 1.0.

** : short range activity coefficients are calculated taking into account inorganic aerosols.

(simulation SOAP-sr) leads to activity coefficients closer to the pure compound state and therefore to a decrease of activity coefficients (as organics are generally more stable at pure state than in water). As activity coefficients are lower in SOAP-sr than in SOAP-Reference, organic concentrations are higher.

- The differences between the simulations SOAP-basic and SOAP-ideal are larger during nighttime than those during daytime.
- 5 This shows that the effect of ideality is larger during nighttime than daytime. This is due to the lower temperature, leading to the condensation of a larger number of organic compounds (some compounds are too volatile to condense during daytime but condense during nighttime).

Figure 4 shows the horizontal distribution of (a) SOA concentrations obtained by the SOAP-Reference simulation, and (b) the differences between the SOAP-Reference and H²O-Reference simulations. As expected, the SOA concentrations are higher

10 in the SOAP-Reference simulation than in the H²O-Reference simulation. Depending on the location, the differences in the SOA concentrations between the simulations are due to different compounds used to compute the partitioning between the gas and particle phases. Over North-Eastern Europe, the differences are due to the large hydrophilic biogenic organic aerosol concentrations. Taking them into account in the computation of c_{aq} strongly increases the concentrations of monoterpenes SOA over South-Western Europe (especially in northern Italy where simulated concentrations of nitrate are high). Over north-

15 ern Italy, large differences are also observed for anthropogenic aromatic organic aerosol concentrations. Even though these compounds are hydrophobic, taking into account the water they absorbed when computing c_p leads to an increase in their concentrations. Similarly, near North Africa and the Iberian Peninsula, the concentrations of hydrophilic surrogates from isoprene oxidation are higher with SOAP than with H²O, because of the large concentrations of sulfate from shipping emissions. Taking sulfate into account (but without taking into account its influence on activity coefficients) when computing the partitioning




Figure 3. Temporal variation of the average SOA concentrations over the domain (see Table 2 for the description of simulations).

between the gas and particle phases leads to an increase in the concentrations of hydrophilic organic compounds in the particle phase.

The coupling of inorganic and organic aerosol formation influences the water absorption by particles. This coupling consists of two effects: the influence of organic species on the inorganic aerosol formation and the influence of inorganic species on the organic aerosol formation. The latter is implemented in the SOAP model, but the influence of organic species on the inorganic




Figure 4. (a) SOA concentrations in the SOAP-Reference simulation ($\mu\text{g m}^{-3}$), (b) the differences of the SOA concentrations between the SOAP-Reference and H²O-Reference simulations ($\mu\text{g m}^{-3}$) and (c) concentration of total condensed water in the SOAP-Reference simulation ($\mu\text{g m}^{-3}$).

aerosol formation is not included. According to Choi and Chan (2002), the organic species can either reduce or enhance the water absorption of inorganic species, which in turns can lead to a change in the condensation of organic species.

Figure 4c shows the concentration of total condensed water in the SOAP-Reference simulation. The coupling of inorganic and organic aerosol formation may lead to changes in the aerosol concentrations in the regions where both the concentrations of total condensed water and hydrophilic organic species are large, e.g., Barcelona, Milano and eastern Spain.

3 Description of the newly added processes

3.1 Interaction of inorganic/organic aerosols using the AIOMFAC model

Although activity coefficients are computed with the UNIFAC model for H²O, depending on the user's choice, in the SOAP model, activity coefficients can be calculated using the UNIFAC or the AIOMFAC model. UNIFAC was developed to reproduce the short-range interactions between water and organic compounds, which are dominant for a non-electrolyte liquid mixture. In UNIFAC, organic compounds are represented by different functional groups including alkane, aromatic carbon, alcohol, carbonyl. Interaction coefficients between water and these functional groups are calculated. However, for an electrolyte liquid mixture, the mixed organic and inorganic system may influence activity coefficients, by middle-range and long-range interactions in addition to the short-range interaction. This influence of inorganic aerosols on the calculation of activity coefficients in the SOAP model can be estimated by the AIOMFAC model that considers this mixed organic/inorganic system.

The activity coefficient in the AIOMFAC model is calculated by the following equation:

$$\gamma = \gamma_{LR} \gamma_{MR} \gamma_{SR} \quad (4)$$

where γ_{LR} , γ_{MR} , γ_{SR} are the contributions of long-range interactions (electrostatic force between ions), middle-range interactions (interactions between ions and molecular dipoles), and short-range interactions (group-contribution method as in UNIFAC).

3.2 Equilibrium and dynamic approaches

Typically, 3-D air quality models mostly use an equilibrium approach to represent condensation/evaporation of aerosols. However, using a dynamic approach may be necessary if the kinetic effects are large (for example if the diffusion in the organic phase is low due to the high particle viscosity or if condensation over coarse particles occurs). In the SOAP model, depending on the user's choice, either the equilibrium approach or the dynamic approach can be used to model condensation/evaporation. An explicit representation of diffusion inside particles, which would involve discretizing the particle along the radius of the particle, can not be used in 3-D air quality models, due to the heavy computation time of such a method. To solve this issue, a method was developed by Couvidat and Sartelet (2015) to implicitly represent the condensation/evaporation/diffusion of organic compounds for a specified organic phase diffusion coefficient. This method separates the particle into a low number of layers that represent different areas of the particle (the gas/particle interface, the core of the particle and intermediate layers).

To use the dynamic approach in this study, several simplifications are carried out for hydrophilic compounds. As a dynamic approach is not used to simulate the formation of inorganic aerosols, the thermodynamic model ISORROPIA (Nenes et al.,

1998) with the equilibrium approach is used to estimate the partitioning of inorganics, the aerosol liquid water content and the pH. The pH given by ISORROPIA is used for each size section and the liquid water content is redistributed over sections proportionally to the amount of inorganics.

In the dynamic approach, the mass transfer rate, J ($\mu\text{g}/\text{m}^3/\text{s}$) by condensation/evaporation at the gas/particle interface is calculated as follows:

$$J_{\text{cond/evap}} = k_{\text{absorption}} (c_g - c_{eq}) \quad (5)$$

where $k_{\text{absorption}}$ is the kinetic rate of absorption (s^{-1}), c_g is the gas-phase concentration ($\mu\text{g}/\text{m}^3$) and c_{eq} is the gas-phase concentration at the interface of particles ($\mu\text{g}/\text{m}^3$).

c_{eq} is calculated taking into account a deviation from an equilibrium concentration. This deviation is presented in Couvidat and Sartelet (2015) (Equations 52 to 55).

The kinetic rate of absorption $k_{\text{absorption}}$ is defined as follows (Seinfeld and Pandis, 1998):

$$k_{\text{absorption}} = 2\pi d_p D_{\text{air}} N f(Kn, \alpha) \quad (6)$$

where d_p is the particle mean diameter (m), D_{air} is the diffusivity of the condensing compounds in air (m^2/s), and N is the number concentration of particles ($\#/ \text{m}^3$). The function $f(Kn, \alpha)$ depends on the Knudsen number ($Kn = \frac{2\lambda}{d_p}$), which is calculated using the mean free path in air λ (m), and the accommodation coefficient α , which accounts for imperfect surface accommodation. It is taken equal to 0.5 following Saleh et al. (2013) and Krechmer et al. (2017).

For viscous particles, the condensation/evaporation is limited by the diffusion flux in the internal layers of the particles.

We assume that in each particle layer the evolution of concentration $c_{p,i}^{\text{layer}}$ of species i can be described as a deviation of an equilibrium concentration ($c_{g,i} K_{p,i}^{\text{layer}} c_p^{\text{layer}}$) when the condensation/evaporation of the species is limited by the diffusion of organic compounds in the organic phase.

This deviation can be described by taking into account the flux of diffusion with the mass transfer rate by condensation/evaporation for each particle layer (Equation 36 of Couvidat and Sartelet (2015)).

$$J_{\text{diff}}^{\text{layer}} = k_{\text{diff}}^{\text{layer}} (c_{g,i} K_{p,i}^{\text{layer}} c_p^{\text{layer}} - c_{p,i}^{\text{layer}}) \quad (7)$$

The kinetic rate of diffusion $k_{\text{diff}}^{\text{layer}}$ (s^{-1}) is computed as follows (Couvidat and Sartelet, 2015):

$$k_{\text{diff}}^{\text{layer}} \propto \frac{1}{\tau_{\text{diff}}} \quad (8)$$

τ_{diff} is the characteristic time (s) for diffusion in the particle:

$$\tau_{\text{diff}} = \frac{R_p^2}{\pi^2 D_{\text{org}}} \quad (9)$$

where R_p is the radius of the particle (m) and D_{org} is the organic-phase diffusivity (m^2/s).

The sum of the diffusion fluxes over all aerosol layers is obtained as follows:

$$J_{\text{diff}} = \sum_{\text{layer}} J_{\text{diff}}^{\text{layer}} \quad (10)$$

The final mass flux by the mixed phenomenon condensation/evaporation/diffusion for the particle is computed by assuming that the characteristic time of the combined effect of condensation/evaporation and diffusion is equal to the sum of the characteristic time of condensation/evaporation ($J_{\text{cond/evap}}$) and the sum of the diffusion fluxes over all aerosol layers (J_{diff}) as follows:

$$\frac{1}{J_{\text{tot}}} = \frac{1}{J_{\text{cond/evap}}} + \frac{1}{J_{\text{diff}}} \quad (11)$$

More details on the model are obtained in Couvidat and Sartelet (2015).

3.3 Particle-phase diffusion cases and impact of viscosity on SOA formation

10 To assess the maximum impact of viscosity on SOA formation: two theoretical studies are studied. The first case, referred hereafter as the “Dynamic inviscid” simulation, assumes that particles are inviscid, i.e. SOA formation is not limited by the particle-phase diffusion: the particle-phase diffusion is so fast that there is no difference in concentrations inside the particle. In this case, compounds condense or evaporate until reaching equilibrium over the whole particle.

The other case, referred as the “Dynamic viscous” simulation, assumes that the particle is “Infinitely viscous” (i.e. too
15 viscous for diffusion to occur inside the particle even at high relative humidity). A very low diffusivity of $10^{-30} \text{ m}^2 \text{ s}^{-1}$ is assumed in order to investigate the maximum deviation of SOA concentrations from the inviscid condition. The diffusivity of organic species is modeled using a bulk viscosity of the mixture estimated by the Refutas method (Maples, 2000). According to measurement studies, the diffusivity of organic species in SOA can be lower than $10^{-21} \text{ m}^2 \text{ s}^{-1}$ (e.g., Pfrang et al., 2011; Abramson et al., 2013). Song et al. (2016b) and DeRieux et al. (2018) showed that scaled values from measured viscosities
20 and predicted values can pass through a viscosity of 10^{12} Pa s , which is the order of a diffusivity of $10^{-30} \text{ m}^2 \text{ s}^{-1}$, at low relative humidity. In addition, Couvidat and Sartelet (2015) reported that at a diffusivity of $10^{-24} \text{ m}^2 \text{ s}^{-1}$, diffusivity does not influence the mass of the condensed organic species as the diffusion is too low to significantly affect the formation of organic aerosol that still occur by condensation/evaporation of organic compounds at the interface. Therefore a diffusivity lower than
25 $10^{-24} \text{ m}^2 \text{ s}^{-1}$ may not affect the concentrations of organic aerosols compared to simulation results with a diffusivity of $10^{-24} \text{ m}^2 \text{ s}^{-1}$.

Practically, for simplification purposes, two aerosol layers (the interface and an internal layer) are used in the “Dynamic viscous” simulation. The internal layer and the interface represent 99% and 1% of the aerosol mass, respectively following the method of Couvidat and Sartelet (2015) in which condensation at interface is not limited by particle-phase diffusion.

The SOA formation for a highly viscous particle is complex. The evolution of the concentration of an organic compound
30 depends on the volatility of the compound with respect to the other compounds. The SOA formation for a highly viscous particle is schematized in Figure 5 in the case of the growth and the shrinking of an extremely viscous particle.

Figure 5 theoretically presents different behaviors of volatile and low volatile organic species in a highly viscous aerosol. For these theoretical cases, mass transfer between the interface and the core is neglected because of an extremely low diffusion flux. The condensation of low-volatility compounds influences the behavior of higher volatility compounds.

In the case of an organic particle growth (mass increase), the condensation of a low-volatility organic compound **B** (its behavior is described by the red curved arrows in Figure 5) onto a particle can favor the condensation of a compound **A** of higher volatility (its behavior is described by blue curved arrows in Figure 5) at the interface of the particle (even if the total concentration of **A** inside the particle exceeds equilibrium). The compound **A** condenses onto the new layer created by the compound **B** to respect Raoult's law at the interface. Even though the compound **A** would evaporate if the particle was inviscid




Figure 5. Schematic representation of SOA formation for a growing (top) and shrinking (down) highly viscous aerosol. The blue curved arrows describe the behavior of an organic compound **A** and the red curved arrows describe the behavior of a low-volatility organic compound **B**.

and the concentration of **A** exceeds equilibrium, for the extremely viscous case, the condensation of the compound **B** at the interface can prevent the evaporation of the compound **A** stuck in the core of the particle (because of the absence of diffusion) and can lead to its “entrapment”.

In the case of a shrinking particle (mass decrease), a volatile compound **A** would evaporate from the inner layers to meet the equilibrium condition if the particle is inviscid and concentration of **A** in the particle exceeds equilibrium. However, if the particle is viscous, this evaporation can strongly be slowed down, because there is no diffusion of the compound **A** from the core to the interface.

Even though the total particle mass reduces, a low-volatility compound **B** may condense at the interface and may therefore slow down the shrinking of the particle. This condensation at the interface prevents the evaporation of the compound **A** from the core of the particle.

In SOAP, a redistribution is done every time step to keep the interface thin and the mass fraction of layers constant (to prevent numerical issues: only the mass of the interface would change for a very viscous particle). This redistribution represents the fact that if the particle grows the compounds that have previously condensed are not anymore at the interface (because other compounds have condensed onto the particle) or that if the particle shrinks the compounds that were previously at the core of the particle will be eventually at the interface. Using two layers, compounds are immediately transferred between the core and the interface. A more accurate representation of the particle dynamics would be obtained using more inner layers to better represent the position of the compounds inside the particle. Nonetheless, the simulation using two layers should give a good estimation on the effect of viscosity on SOA formation.

These SOA formations for a highly viscous particle are more discussed in section 4.5.

20 **4 Impact on SOA formation**

4.1 Simulation setup

The Polair3D model coupled to the SOAP model is evaluated during summer 2012. The modeling domain covers Europe with a horizontal resolution of $0.5^\circ \times 0.5^\circ$ (see Figure 2). Anthropogenic emissions are generated with the EMEP inventory for 2012. Intermediate and semi-volatile organic compounds (I/S-VOC) emissions are estimated as detailed in Couvidat et al. (2012), by multiplying the primary organic aerosol emissions by a factor 4 and by assigning them to compounds of different volatilities (POAIP, POAmP and POAhP). Biogenic emissions are generated with the MEGAN model (Guenther et al., 2006). ECMWF (European Centre for Medium-Range Weather Forecasts) meteorological reanalysis data (<http://www.ecmwf.int/>, ERA Interim) are used to calculate meteorological fields. Initial and boundary conditions are obtained from the simulation data of MOZART-4/GEOS-5 (<http://www.acom.ucar.edu/wrf-chem/mozart.shtml>). The aerosols are assumed internally mixed in this model. The number of aerosol bins is 5 covering from 0.01 to 10 μm . An adaptative time-step is used to solve the dynamics of organics. The minimum time step is 1 s and the maximum time step is set to 10 min in the simulations of this study. 10 min correspond to the time step used to split the different processes in the 3-D model (advection, diffusion and chemistry). When concentrations are computed by the dynamic approach, the second-order Rosenbrock scheme is used for

time integration (Couvidat and Sartelet, 2015). Further details about the model configuration may be found in Couvidat et al. (2012).

Six sensitivity simulations are conducted over Europe to study the effect of non-ideality and non-equilibrium phenomena on SOA formation. The list of the simulations is presented in Table 3. The reference simulation (named “Equilibrium UNIFAC”) uses the default model options: thermodynamic equilibrium between the gas and particle phases is assumed and activity coefficients are computed with UNIFAC. To evaluate the impact of activity coefficients on concentrations, a simulation (named “Equilibrium Ideal”) is run. The impact of inorganic aerosols on the short-range activity coefficients using UNIFAC is estimated with a simulation (named “Equilibrium UNIFAC-sr”). To evaluate the impact of inorganic concentrations on activity coefficients, a simulation (named “Equilibrium AIOMFAC”) using AIOMFAC to compute activity coefficients instead of UNIFAC is run.

To evaluate the impact of the particle viscosity on SOA concentrations, two other simulations are run: one with a dynamic approach to model condensation/evaporation of inviscid particles (simulation named “Dynamic inviscid”), and one with a dynamic approach but extremely viscous particles (simulation named “Dynamic viscous”). The simulations that use the equilibrium approach for condensation/evaporation are run from 1 June to 31 August 2012. However, the sensitivity simulations using the dynamic approach are run for only 3 weeks starting 1 June 2012, because of expensive computational time.

The used absorption approaches for the simulations are presented in Table 3. The absorbing mass includes inorganic aerosol, hydrophilic organic aerosol and water absorbed by inorganic aerosol/hydrophilic organic aerosol for c_{aq} . c_p includes hydrophobic organic aerosol and water absorbed by hydrophobic organic aerosol as listed in Table 2.

The algorithm of SOAP was developed in order to consider both the organic and the aqueous phases inside a particle. It assumes that the organic and the aqueous phases co-exist in a particle but evolve separately in different regions of the particle. For example, for the dynamic representation, if a compound tends to go from the aqueous to the organic phases, it has first to evaporate to the gas phase and then condense to the organic phases instead of a direct mass transfer between the phases. It is

Table 3. List of the sensitivity simulations using different options in SOAP.

Simulation name	Absorption approach	Activity coefficient	Viscous aerosol	Number of aerosol layers
Equilibrium UNIFAC	equilibrium	UNIFAC	No	1
Equilibrium Ideal	equilibrium	ideal mixture	No	1
Equilibrium UNIFAC-sr	equilibrium	UNIFAC-sr**	No	1
Equilibrium AIOMFAC	equilibrium	AIOMFAC	No	1
Dynamic inviscid	dynamic	UNIFAC	No	1
Dynamic viscous	dynamic	UNIFAC	Yes	2

** : activity coefficients are calculated taking into account inorganic aerosols.

due to the complexity of representing properly these transfers. This assumption is discussed in more details in section 2.3 of Couvidat and Sartelet (2015).

4.2 Model evaluation

To evaluate the general performances of the model, the concentrations of organic aerosols given by the “Equilibrium UNIFAC” simulation are compared to the concentrations of organic matter (OM), organic carbon (OC) and inorganic aerosols measured at stations of the ACTRIS observation network (<http://actris.nilu.no>) in Europe. OC concentrations are measured by high/low volume samplers and those of OM and inorganic aerosols are measured by aerosol mass spectrometers. The locations of stations are presented in Figure 2. To compare the simulated OM concentrations with the measured OC, the simulated OM concentrations are converted into OC concentrations using the OM/OC ratio for each surrogate of the organic aerosols, as described in Couvidat et al. (2012).

To evaluate the model ability to reproduce SOA concentrations, the standard metrics of the model performance for particulate matter of Boylan and Russell (2006) are used: the mean fractional bias (MFB), the mean fractional error (MFE), the root mean square error (RMSE), and the correlation. Boylan and Russell (2006) proposed a performance evaluation criteria (IMFBI < 60% and MFE < 75%) and a goal evaluation criteria (IMFBI < 30% and MFE < 50%). Model performance statistics are presented in Table 4. For organic compounds, the model performance and goal criteria are satisfied for the stations Kosetice, Melpitz, Ispra, and Cabauw (IMFBI < 30% and MFE < 50%, see Figure 6). For the Birkenes station, the performance criteria are satisfied, but the goal criteria are almost satisfied, although the concentrations of OM₁ are quite underestimated (MFB: -33%). For the Aspveten station, the model performance criteria are satisfied but the OC₁₀ concentrations are significantly underestimated (MFB: -56%). For the Montseny stations, the goal criteria are satisfied for OC and the performance criteria are satisfied for OM₁ with a significant underestimation.

For inorganic PM₁ aerosols, the performance biases and errors are satisfied at the Cabauw station. However, inorganic PM₁ concentrations are underestimated at the Montseny and Melpitz stations, even though the MFE satisfies the model performance criteria.

4.3 Impact of inorganic-organic interactions

Figure 7 shows the modeled hydrophilic SOA concentrations. The choice of thermodynamic model affects the spatial distribution of hydrophilic SOA with an decrease of concentrations when using AIOMFAC over Netherlands, Belgium, part of Italy, Spain over the Mediterranean coast and southeastern Europe and an increase of concentrations over Northern Europe, part of the Alps, southern France, part of Italy and part of Spain. The area with the strongest decrease of concentrations corresponds to areas with strong inorganic concentrations. For example, the decrease of concentrations over Netherlands corresponds to high ammonium nitrate concentrations while the decrease in southeastern Europe corresponds to high ammonium sulfate concentrations.

The changes in concentration of specific SOA compounds using AIOMFAC and UNIFAC are illustrated by Figure 8. The local increases of concentrations can be due to non-linear effects. Indeed, while the concentrations of the less oxidized hy-

Table 4. Comparison of the simulated concentrations to the measurements. Performance statistics are calculated with daily mean concentrations.

Station	Particle type*	Measurement [†] ($\mu\text{g m}^{-3}$)	Simulation [†] ($\mu\text{g m}^{-3}$)	RMSE ($\mu\text{g m}^{-3}$)	MFB	MFE	Correlation
Kosetice	OC _{2.5}	2.36	1.94	0.86	-26%	37%	0.73
Melpitz	OC _{2.5}	1.41	1.67	0.56	16%	27%	0.85
	OC ₁₀	2.05	1.67	0.63	-23%	28%	0.86
	OM ₁	3.83	2.75	2.21	-20%	38%	0.81
	NH _{4,1}	0.66	0.46	0.32	-30%	36%	0.72
	NO _{3,1}	0.84	0.53	0.59	-47%	60%	0.56
	SO _{4,1}	1.60	0.84	0.91	-62%	63%	0.74
Montseny	OC ₁	1.89	2.06	0.57	6%	20%	0.76
	OC _{2.5}	2.41	2.20	0.70	-10%	23%	0.66
	OC ₁₀	2.72	2.06	0.99	-30%	35%	0.64
	OM ₁	7.56	3.90	4.80	-60%	64%	0.26
	NH _{4,1}	1.14	0.59	0.78	-60%	60%	0.49
	NO _{3,1}	0.58	0.36	0.77	-51%	66%	0.04
	SO _{4,1}	2.54	1.23	1.84	-65%	66%	0.44
Ispra	OC _{2.5}	2.79	3.16	0.88	16%	27%	0.75
Aspvreten	OC ₁₀	2.26	1.29	1.14	-56%	56%	0.63
Birkenes	OM ₁	1.55	1.07	0.54	-39%	39%	0.80
Cabauw	OM ₁	2.86	2.59	1.11	0%	20%	0.94
	NH _{4,1}	1.04	1.09	0.52	11%	30%	0.78
	NO _{3,1}	2.82	2.65	2.00	12%	49%	0.75
	SO _{4,1}	0.88	0.95	0.31	8%	21%	0.79

*: subscripts are used for the particle size. For example, OC_{2.5} is organic carbon of aerodynamic diameter lower than 2.5 μm . For ammonium (NH₄), sulfate (SO₄) and nitrate (NO₃), e.g., SO_{4,1} is sulfate of aerodynamic diameter lower than 1 μm .

[†]: mean concentration from June 1 to August 31, 2012.

drophilic compounds (BiA0D with a ratio of oxygen to carbon atoms (O/C) of 0.2 and BiA1D with a O/C of 0.5) mainly decrease over Europe (-2% for BiA0D and -27% for BiA1D), the concentrations of the more oxidized compounds (BiPER with a O/C of 1.2 and BiDER with a O/C of 0.8) mainly increase (6% for BiPER and 16% for BiDER). This finding is in agreement with Pye et al. (2018) who found that in the eastern US, particle-phase interactions of organic and inorganic compounds increase partitioning toward the particle phase (vs. gas phase) for highly oxidized compounds (O/C \geq 0.6) but decrease particle-phase partitioning for low O/C.




Figure 6. Comparison of modeled SOA concentrations (blue) with observations (black) for (a) $OC_{2.5}$ concentrations at Ispra, (b) OM_1 concentrations at Cabauw.

Figure 7d shows the temporal evolution of the domain-averaged concentration of hydrophilic SOA for different simulations. Modeling organic interactions by activity coefficients strongly influences hydrophilic SOA. It leads on average to a concentration increase of 33% (simulation “Equilibrium-UNIFAC” compared to simulation “Equilibrium-Ideal”). When the computation of short-range interactions between inorganic and organic aerosols is taken into account in UNIFAC, the SOA concentrations increase because of a decrease of activity coefficient (see “Equilibrium UNIFAC-sr” simulation in Figure 7d). Long-range and medium-range interactions in the “Equilibrium AIOMFAC” simulation lead to an increase of activity coefficients as concentrations decrease compared to the “Equilibrium UNIFAC-sr” simulation by 28%. This evaporation of hydrophilic organic concentrations by the medium and long-range inorganic-organic interactions agrees with the results of Zuend et al. (2008), who showed that the activity coefficients of hydrophilic organic aerosols increase because of the interactions with inorganic aerosols.

The SOA concentrations simulated with the AIOMFAC model are close to the concentrations simulated with the UNIFAC model (without taking into account inorganics in the computation of short-range interactions). It suggests that computing activity coefficients for hydrophilic organic compounds by only taking water and organic compounds (and therefore by ignoring inorganics) could give a good first approximation of activity coefficients. Medium and long-range interactions compensate the decrease of activity coefficients due to the inclusion of inorganic ions in short-range interactions. We estimated the contributions of long-range and medium-range interactions in this decrease by an additional simulation. In this additional simulation, only the medium-range interactions are taken into account in the AIOMFAC model. According to the results of this simulation, the differences in the concentrations of hydrophilic SOA are due to the medium-range interactions by 65% and the long-range interactions by 35%.




Figure 7. Modeled hydrophilic SOA concentrations in (a) the Equilibrium AIOMFAC simulation, (b) the Equilibrium UNIFAC simulation, (c) the differences between the simulations ($\mu\text{g m}^{-3}$) and (d) the temporal evolutions of domain averaged concentrations.

4.4 Impact of the thermodynamic equilibrium assumption

The aqueous phase of the particles is assumed inviscid and organics are strongly influenced by inorganic concentrations, because they constitute an absorbing mass for hydrophilic organics. However, in the organic phase, the particles may be viscous, and the dynamic evolution of the SOA concentrations by condensation/evaporation may be limited by diffusion due to the particle viscosity (Couvidat and Sartelet, 2015).

To evaluate the impact of particle viscosity on SOA concentrations, condensation/evaporation need to be solved using the dynamic approach. Because condensation/evaporation is solved using the equilibrium approach in the previous simulations,




Figure 8. Differences in concentrations of several SOA compounds between AIOMFAC and UNIFAC over Europe (AIOMFAC - UNIFAC in $\mu\text{g m}^{-3}$). The definition of the compounds is given in Table 1.

the impact of using the dynamic approach, while still assuming particles to be inviscid, is assessed by running the “Dynamic inviscid” simulation.

Differences between the “Equilibrium-UNIFAC” and the “Dynamic inviscid” simulations are very low for hydrophobic compounds (less than 1%) whereas a decrease of concentrations by about 6% is found for hydrophilic compounds in the

5 “Dynamic inviscid” simulation. The differences are due to the non-ideality of the aerosols as low differences are found when assuming ideality (3%). In the “Equilibrium-UNIFAC” simulation, activity coefficients are computed by taking the composition of the aerosols averaged over size sections. However, for the “Dynamic inviscid” simulation, activity coefficients are computed








Figure 9. Modeled hydrophobic SOA concentrations in (a) the Dynamic inviscid simulation, (b) the Dynamic viscous simulation, (c) the differences between the simulations ($\mu\text{g m}^{-3}$) and (d) the temporal evolutions.

for each size section. The section activity coefficients of the “Dynamic inviscid” simulation are therefore different from the activity coefficients of the “Equilibrium-UNIFAC” simulation.



For hydrophobic compounds, the differences are mainly due to the variations of the mass transfer rate computed by Equation 5. In the dynamic approach, the condensation/evaporation is slower than in the equilibrium approach. Therefore, using the dynamic approach reduces the magnitudes of the peaks in the temporal variations of the SOA concentrations, although the average concentrations do not change much with the temporal and spatial resolutions used here. In the dynamic approach, in opposition to the equilibrium approach, low-volatility secondary compounds formed by gas-phase chemistry are found to not be totally into the particle phase due to the kinetic of condensation. For example, 97% of SOAIP is absorbed inside the particle and 3% of SOAIP is still present in the gas-phase.



(a) SOAhP



(b) SOAIP



(c) POAIP

Figure 10. Temporal evolution (left) and differences between the Dynamic viscous and Dynamic inviscid simulations (right) of modeled SOA concentrations for (a) SOAhP, (b) SOAIP, and (c) POAIP ($\mu\text{g m}^{-3}$).

4.5 Impact of viscosity of the organic phase

In the simulation Dynamic-viscous, as expected, the dynamic evolution of hydrophilic SOA concentration does not change from those of the “Dynamic inviscid” simulation, but the organic hydrophobic phase is strongly influenced by the viscosity.

Assuming that the organic phase is very viscous leads to an increase in concentrations of hydrophobic SOA: 6% on average of the total concentrations (see Figure 9). The increase can exceed 20% over areas with low concentrations in the “Dynamic inviscid” simulation (Spain and Northern Europe). This increase of concentrations of hydrophobic SOA is due to the absence of evaporation (because of the absence of diffusion) when concentrations exceed equilibrium. The hydrophobic SOA concentrations in the “Dynamic viscous” simulation decrease where they are very high in the “Dynamic inviscid” simulation (the Straits of Gibraltar and Istanbul). As shown in Figure 9d, the increase of concentrations in the “Dynamic viscous” simulation happens mainly during daytime.

The influence of viscosity differs depending on the volatility of the surrogate. For example, in the model, the emitted anthropogenic I/S-VOC are represented by surrogates of different volatility classes: high volatility (POAhP, $K_p = 0.00031 \text{ m}^3 \mu\text{g}^{-1}$), average volatility (POAmP, $K_p = 0.0116 \text{ m}^3 \mu\text{g}^{-1}$) and low volatility (POAIP, $K_p = 1.1 \text{ m}^3 \mu\text{g}^{-1}$). The chemical kinetic mechanism used for the SOAP model includes the oxidation of these surrogates to other surrogates with lower volatilities: SOAhP ($K_p = 0.031 \text{ m}^3 \mu\text{g}^{-1}$), SOAmP ($K_p = 1.16 \text{ m}^3 \mu\text{g}^{-1}$) and SOAIP ($K_p = 110 \text{ m}^3 \mu\text{g}^{-1}$) with the following equations (Couvidat et al., 2012).



In Figure 10a, the concentrations of SOAhP (one of the most volatile compounds of the mechanism) strongly increase in the “Dynamic viscous” simulation (by 44% in average). This increase is especially strong in Southeastern Europe where concentrations double and increase by $0.1 \mu\text{g m}^{-3}$. In the “Dynamic viscous” simulation, concentrations increase strongly at the beginning of the day and reach a maximum during daytime. On the contrary, in the “Dynamic inviscid” simulation, concentrations decrease at the beginning of the day and reach a minimum during daytime (as the volatility of the compound increases during daytime). In the “Dynamic viscous” simulation, the diurnal variations of SOAhP follow those of the low-volatility compound SOAIP (Figure 10b). Figure 11 show the deviation of the particle/gas partitioning from equilibrium for SOAhP (defined as the particle/gas concentration ratio in the “Dynamic viscous” simulation over that in the “Dynamic inviscid” simulation, which is close to equilibrium). This deviation often exceeds 50% and particle-phase concentrations exceeds equilibrium over most of Europe. As presented in section 3.3, condensation of a semi-volatile compound can happen without respecting equilibrium as long as the particle is growing (growth that can be due to the condensation of a low-volatile compound such as SOAIP). The condensation during the day of low-volatile compounds formed during daytime stops the evaporation of SOAhP captured inside the particle (evaporation that would occur for an inviscid organic phase) and is even able to bring further condensation of the compound.

Concentrations for the low-volatile compound SOAIP slightly decrease (see Figure 10b). This decrease is mainly due to the increase of the POAIP particle-phase concentration during daytime (see Figure 10c). In the chemical kinetic mechanism used in this study, SOAIP is formed from the gas-phase oxidation of POAIP by OH radical (mainly present daytime). The increase of POAIP in the particle-phase during daytime slows down the formation of the compound SOAIP. On the contrary, at the end of the day, concentrations of POAIP become higher in the “Dynamic inviscid” simulation due to the decrease of volatility (because of the decrease of temperature). However, in the “Dynamic viscous” simulation, the decrease of the volatility has a low effect on concentrations (because the inner layer cannot absorb more compounds to reach equilibrium due to the absence of diffusion).

The large deviations from equilibrium suggested by this study agree with the measurements of Yatavelli et al. (2014) and Lopez-Hilfiker et al. (2015) who observed that the concentrations of pinonic acid in SOA are much higher than predicted with the equilibrium assumption using saturation vapor pressures. It could also be possible that this phenomenon is due to non-ideal effect and the possibility for pinonic acid to be absorbed onto an aqueous phase with an acidic dissociation.

The viscosity effect is very low for low-volatile compounds (Couvidat and Sartelet, 2015). Here, extremely low volatile compounds from the oxidation of monoterpenes are not modeled (Chrit et al., 2017). Taking them into account may decrease the viscosity effect estimated in this study.

4.6 Comparison of computation times

Table 5 presents the time elapsed for each simulation. The elapsed time for the “Equilibrium UNIFAC” simulation is set to a reference time. The time elapsed for the “Equilibrium AIOMFAC” simulation increases by 45% compared to the reference time. Using the dynamic approach leads to an increase of the computation time by a factor ten, making it difficult to represent viscous aerosols in long-term 3-D simulations. However, these computation times are acceptable for short-term 3-D simulations.

5 Conclusions

The SOAP model, which uses either the equilibrium approach or the dynamic approach for the mass transfer of organic compounds from the gas phase to the particle phases, was implemented in the 3-D air quality model of Polyphemus. Compared to its predecessor, SOAP provides a more complete description of the partitioning of semi-volatile compounds, in particular, by taking into account the effect of inorganic aerosols on SOA formation based on the computation of activity coefficients given by AIOMFAC. Sensitivity simulations indicate that including inorganic aerosols and hydrophilic organic aerosols in

Table 5. Comparison of time elapsed for the simulations. The elapsed time for the Equilibrium UNIFAC simulation is set to a reference time and ratios between the reference time and the time elapsed for other simulations are presented.

	Equilibrium Ideal	Equilibrium UNIFAC	Equilibrium AIOMFAC	Dynamic Inviscid	Dynamic Viscous
Ratio of computation time	0.97	1	1.45	9.74	10.02




Figure 11. Equilibrium deviation for compound SOAhP. A deviation close to 1 means that the compound reaches equilibrium, above 1 means that particle-phase concentrations are above equilibrium and under 1 that concentrations have not reached equilibrium.

the absorbing mass of the aqueous-phase can lead to an increase of concentrations around 5% and 6%, respectively. Overall, hydrophilic SOA concentrations using AIOMFAC are higher than those with the ideality assumption by about 33%. The results of this study suggest that non-ideality via organic-organic and inorganic-organic interactions influence strongly the condensation of hydrophilic organic compounds.

5 For an inviscid aerosol, the results of this study show that the equilibrium assumption is an efficient approximation, when assuming ideality for organic aerosols. However, assuming equilibrium can lead to significant differences in the concentrations of hydrophilic compounds when non-ideality is taken into account. Indeed, with a dynamic approach, different values of activity coefficients can be simulated for the different size sections. These results indicate that differences in the composition of particles with particle size can impact the formation of SOA. Note that in this study, an equilibrium approach is used for the
 10 condensation of inorganics. Using a dynamic approach to model the condensation/evaporation of both inorganic and organic compounds may be necessary to properly estimate the formation of hydrophilic SOA.

The dynamic approach in the SOAP model is used to account for the viscosity of aerosol to study SOA formation via two theoretical cases: the “inviscid case” where diffusion is extremely fast and concentrations inside the particle are homogeneous and the “infinite viscosity case” where viscosity is too high for diffusion to occur inside the particle but where condensation
 15 or evaporation of compounds at the gas/particle interface can still occur. Even if the two cases presented in this study are

theoretical, the results provide a first insight on how viscosity may affect SOA formation. For the “inviscid case”, concentrations of hydrophobic compounds are close to those in the equilibrium simulation. However, assuming a highly viscous organic-phase leads to an increase of hydrophobic SOA concentration during daytime (by stopping the evaporation of the most volatile compounds without stopping their condensation). SOA formation for a highly viscous particle can therefore significantly deviate from thermodynamic equilibrium, e.g., condensation can happen when evaporation occurs if equilibrium is assumed. This deviation may explain why some observed concentrations in the literature are significantly different to concentrations calculated with the equilibrium assumption and saturation vapor pressures.

Those results emphasize the need to study the effect of the dynamics of SOA formation. Next modeling studies should focus on the sensitivity of results to the organic-phase diffusion coefficient and try to take into account the effect of temperature, the aerosol water content and also aerosol composition on this parameter.

The estimation of the computation time shows that the dynamic approach used in the SOAP model can be applicable to the 3-D air quality modeling for a short period or with high computation time capability. Although, the results emphasize the need to study the effect of a dynamic approach compared to an equilibrium approach, the computation-time issue is probably a limiting factor in the possibility for the implementation of dynamic approaches in 3-D air quality models.

Finally, the effect of morphology for a highly viscous aerosol may be critical for a highly viscous aerosol. The coagulation of two highly viscous spherical particles may form a non-spherical particle composed of two spheres stuck together. Non-spherical particles may lead to higher surface-to-volume ratio and faster condensation/evaporation/diffusion.

Acknowledgements. This work was funded by the French Minister of Environment and Sustainable Development under the project NA-TORGA of the COPERNICUS-MDD program. The authors would like to thank the principal investigators of the ACTRIS measurements used for the model evaluation (Rupert Holzinger, Patrick Schlag and Astrid Kiendler-Scharr for the station Cabauw; Wenche Aas and Chris Lunder for the station Birkenes; Maria Cruz Minguillon, Anna Ripoll and Andres Alastuey for the station Montseny; Laurent Poulain for the station Melpitz; F. Cavalli and J.P. Putaud for the station Ispra; Milan Vana for the station Kosetice; Peter Tunved and Hans Areskoug for the station Aspöreten).

References

- Abramson, E., Imre, D., Beranek, J., Wilson, J., and Zelenyuk, A.: Experimental determination of chemical diffusion within secondary organic aerosol particles, *Phys. Chem. Chem. Phys.*, 15, 2983–2991, doi:10.1039/C2CP44013J, 2013.
- Boucher, O.: *Atmospheric Aerosols, Properties and Climate Impact*, Springer-Netherlands, doi:10.1007/978-94-017-9649-1, 2015.
- 5 Bowman, F. M., Odum, J. R., Seinfeld, J. H., and Pandis, S. N.: Mathematical model for gas-particle partitioning of secondary organic aerosols, *Atmos. Environ.*, 31, 3921 – 3931, doi:10.1016/S1352-2310(97)00245-8, 1997.
- Boylan, J. W. and Russell, A. G.: PM and light extinction model performance metrics, goals, and criteria for three-dimensional air quality models, *Atmos. Environ.*, 40, 4946–4959, doi:10.1016/j.atmosenv.2005.09.087, 2006.
- Capaldo, K. P., Pilinis, C., and Pandis, S. N.: A computationally efficient hybrid approach for dynamic gas/aerosol transfer in air quality
10 models, *Atmos. Environ.*, 34, 3617 – 3627, doi:10.1016/S1352-2310(00)00092-3, 2000.
- Cappa, C. D. and Wilson, K. R.: Evolution of organic aerosol mass spectra upon heating: implications for OA phase and partitioning behavior, *Atmos. Chem. Phys.*, 11, 1895–1911, doi:10.5194/acp-11-1895-2011, 2011.
- Choi, M. Y. and Chan, C. K.: The effects of organic species on the hygroscopic behaviors of inorganic aerosols, *Environ. Sci. Technol.*, 36, 2422–2428, doi:10.1021/es0113293, 2002.
- 15 Chrit, M., Sartelet, K., Sciare, J., Pey, J., Marchand, N., Couvidat, F., Sellegri, K., and Beekmann, M.: Modelling organic aerosol concentrations and properties during ChArMEx summer campaigns of 2012 and 2013 in the western Mediterranean region, *Atmos. Chem. Phys.*, 17, 12 509–12 531, doi:10.5194/acp-17-12509-2017, 2017.
- Couvidat, F. and Sartelet, K.: The Secondary Organic Aerosol Processor (SOAP v1.0) model: a unified model with different ranges of complexity based on the molecular surrogate approach, *Geosci. Model Dev.*, 8, 1111–1138, doi:10.5194/gmd-8-1111-2015, 2015.
- 20 Couvidat, F., Debry, É., Sartelet, K., and Seigneur, C.: A hydrophilic/hydrophobic organic (H²O) model: Model development, evaluation and sensitivity analysis, *J. Geophys. Res.*, 117, D10 304, doi:10.1029/2011JD017214, 2012.
- Couvidat, F., Kim, Y., Sartelet, K., Seigneur, C., Marchand, N., and Sciare, J.: Modeling secondary organic aerosol in an urban area: application to Paris, France, *Atmos. Chem. Phys.*, 13, 983–996, doi:10.5194/acp-13-983-2013, 2013.
- Debry, É., Fahey, K., Sartelet, K., Sportisse, B., and Tombette, M.: Technical Note: A new Size REsolved Aerosol Model (SIREAM), *Atmos. Chem. Phys.*, 7, 1537–1547, doi:10.5194/acp-7-1537-2007, 2007.
- 25 DeRieux, W.-S. W., Li, Y., Lin, P., Laskin, J., Laskin, A., Bertram, A. K., Nizkorodov, S. A., and Shiraiwa, M.: Predicting the glass transition temperature and viscosity of secondary organic material using molecular composition, *Atmos. Chem. Phys.*, 18, 6331–6351, doi:10.5194/acp-18-6331-2018, 2018.
- Fredenslund, A., Jones, R. L., and Prausnitz, J. M.: Group-contribution estimation of activity coefficients in nonideal liquid mixtures, *AIChE Journal*, 21, 1086–1099, doi:10.1002/aic.690210607, 1975.
- 30 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, doi:10.5194/acp-6-3181-2006, 2006.
- Jacobson, M. Z.: Development and application of a new air pollution modeling system—II. Aerosol module structure and design, *Atmos. Environ.*, 31, 131 – 144, doi:10.1016/1352-2310(96)00202-6, 1997.
- 35

- Jathar, S. H., Mahmud, A., Barsanti, K. C., Asher, W. E., Pankow, J. F., and Kleeman, M. J.: Water uptake by organic aerosol and its influence on gas/particle partitioning of secondary organic aerosol in the United States, *Atmos. Environ.*, 129, 142 – 154, doi:10.1016/j.atmosenv.2016.01.001, 2016.
- 5 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. D., Ulbrich, I., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, E. J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, 10 L. R., Wood, E. C., Kolb, C. E., Middlebrook, A. M., Baltensperger, U., and Worsnop, D. R.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525–1529, doi:10.1126/science.1180353, 2009.
- Kim, Y., Couvidat, F., Sartelet, K., and Seigneur, C.: Comparison of different gas-phase mechanisms and aerosol modules for simulating particulate matter formation, *J. Air Waste Manage. Assoc.*, 61, 1–9, doi:10.1080/10473289.2011.603999, 2011.
- Krechmer, J. E., Day, D. A., Ziemann, P. J., and Jimenez, J. L.: Direct measurements of gas/particle partitioning and mass accommodation 15 coefficients in environmental chambers, *Environ. Sci. Technol.*, 51, 11 867–11 875, doi:10.1021/acs.est.7b02144, 2017.
- Lopez-Hilfiker, F. D., Mohr, C., Ehn, M., Rubach, F., Kleist, E., Wildt, J., Mentel, T. F., Carrasquillo, A. J., Daumit, K. E., Hunter, J. F., Kroll, J. H., Worsnop, D. R., and Thornton, J. A.: Phase partitioning and volatility of secondary organic aerosol components formed from α -pinene ozonolysis and OH oxidation: the importance of accretion products and other low volatility compounds, *Atmos. Chem. Phys.*, 15, 7765–7776, 10.5194/acp-15-7765-2015, 2015.
- 20 Mallet, V., Qu  lo, D., Sportisse, B., Ahmed de Biasi, M., Debry,  . E., Korsakissok, I., Wu, L., Roustan, Y., Sartelet, K., Tombette, M., and Foudhil, H.: Technical Note: The air quality modeling system Polyphemus, *Atmos. Chem. Phys.*, 7, 5479–5487, 2007.
- Maples, R. E.: *Petroleum Refinery Process Economics*, Pennwell Books, 2nd ed., 2000.
- Meng, Z., Dabdub, D., and Seinfeld, J. H.: Size-resolved and chemically resolved model of atmospheric aerosol dynamics, *J. Geophys. Res.*, 103, doi:10.1029/97JD02796, 1998.
- 25 Nenes, A., Pandis, S. N., and Pilinis, C.: ISORROPIA: A new thermodynamic equilibrium model for multiphase multicomponent inorganic aerosols, *Aquat. Geochem.*, 4, 123–152, doi:10.1023/A:1009604003981, 1998.
- Nenes, A., Pandis, S. N., and Pilinis, C.: Continued development and testing of a new thermodynamic aerosol module for urban and regional air quality models, *Atmos. Environ.*, 33, 1553–1560, doi:10.1016/S1352-2310(98)00352-5, 1999.
- O’Meara, S., Topping, D., Zaveri, R., and McFiggans, G.: An efficient approach for treating composition-dependent diffusion within organic 30 particles., *Atmos. Chem. Phys.*, 17, 10 477–10 494, doi:10.5194/acp-17-10477-2017, 2017.
- Pankow, J. F.: An absorption model of gas/particle partitioning of organic compounds in the atmosphere, *Atmos. Environ.*, 28, 185–188, doi:10.1016/1352-2310(94)90093-0, 1994.
- Pfrang, C., Shiraiwa, M., and P  schl, U.: Chemical ageing and transformation of diffusivity in semi-solid multi-component organic aerosol particles, *Atmos. Chem. Phys.*, 11, 7343–7354, doi:10.5194/acp-11-7343-2011, 2011.
- 35 Putaud, J.-P., Raes, F., Van Dingenen, R., Br  ggemann, E., Facchini, M.-C., Decesari, S., Fuzzi, S., Gehrig, R., H  glin, C., Laj, P., Lorbeer, G., Maenhaut, W., Mihalopoulos, N., M  ller, K., Querol, X., Rodriguez, S., Schneider, J., Spindler, G., ten Brink, H., T  rseth, K., and Wiedensohler, A.: A european aerosol phenomenology–2: chemical characteristics of particulate matter at kerbside, urban, rural and background sites in Europe, *Atmos. Environ.*, 38, 2579–2595, doi:10.1016/j.atmosenv.2004.01.041, 2004.

- Pye, H., Zuend, A., Fry, J., Isaacman-VanWertz, G., Capps, S., Appel, K., Foroutan, H., Xu, L., Ng, N., and Goldstein, A.: Coupling of organic and inorganic aerosol systems and the effect on gas–particle partitioning in the southeastern US., *Atmos. Chem. Phys.*, 18, 357–370, doi:10.5194/acp-18-357-2018, 2018.
- Pye, H. O. T., Murphy, B. N., Xu, L., Ng, N. L., Carlton, A. G., Guo, H., Weber, R., Vasilakos, P., Appel, K. W., Budisulistiorini, S. H., Surratt, J. D., Nenes, A., Hu, W., Jimenez, J. L., Isaacman-VanWertz, G., Misztal, P. K., and Goldstein, A. H.: On the implications of aerosol liquid water and phase separation for organic aerosol mass, *Atmos. Chem. Phys.*, 17, 343–369, doi:10.5194/acp-17-343-2017, 2017.
- Rattanavaraha, W., Rosen, E., Zhang, H., Li, Q., Pantong, K., and Kamens, R. M.: The reactive oxidant potential of different types of aged atmospheric particles: An outdoor chamber study, *Atmos. Environ.*, 45, 3848–3855, doi:10.1016/j.atmosenv.2011.004.02, 2011.
- 10 Reid, J. P., Bertram, A. K., Topping, D. O., Laskin, A., Martin, S. T., Petters, M. D., Pope, F. D., and Rovelli, G.: The viscosity of atmospherically relevant organic particles, *Nature communications*, 9, doi:10.1038/s41467-018-03027-z, 2018.
- Renbaum-Wolff, L., Grayson, J. W., Bateman, A. P., Kuwata, M., Sellier, M., Murray, B. J., Shilling, J. E., Martin, S. T., and Bertram, A. K.: Viscosity of α -pinene secondary organic material and implications for particle growth and reactivity, *Proc. Natl. Acad. Sci.*, 110, 8014–8019, doi:10.1073/pnas.1219548110, 2013.
- 15 Roldin, P., Eriksson, A. C., Nordin, E. Z., Hermansson, E., Mogensen, D., Rusanen, A., Boy, M., Swietlicki, E., Svenningsson, B., Zelenyuk, A., and Pagels, J.: Modelling non-equilibrium secondary organic aerosol formation and evaporation with the aerosol dynamics, gas- and particle-phase chemistry kinetic multilayer model ADCHAM, *Atmos. Chem. Phys.*, 14, 7953–7993, doi:10.5194/acp-14-7953-2014, 2014.
- Saleh, R., Donahue, N. M., and Robinson, A. L.: Time scales for gas-particle partitioning equilibration of secondary organic aerosol formed from alpha-pinene ozonolysis, *Environ. Sci. Technol.*, 47, 5588–5594, doi:10.1021/es400078d, 2013.
- 20 Sartelet, K., Hayami, H., and Sportisse, B.: Dominant aerosol processes during high-pollution episodes over Greater Tokyo., *J. Geophys. Res.*, 112, doi:10.1029/2006JD007885, 2007a.
- Sartelet, K. N., Debry, E., Fahey, K., Roustan, Y., Tombette, M., and Sportisse, B.: Simulation of aerosols and gas-phase species over Europe with the Polyphemus system: Part I–Model-to-data comparison for 2001, *Atmos. Environ.*, 41, 6116–6131, doi:10.1016/j.atmosenv.2007.04.024, 2007b.
- 25 Sartelet, K. N., Couvidat, F., Seigneur, C., and Roustan, Y.: Impact of biogenic emissions on air quality over Europe and North America, *Atmos. Environ.*, 53, 131 – 141, doi:10.1016/j.atmosenv.2011.10.046, 2012.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, Wiley-Interscience, New York, 1998.
- 30 Shiraiwa, M. and Seinfeld, J. H.: Equilibration timescale of atmospheric secondary organic aerosol partitioning, *Geophys. Res. Lett.*, 39, doi:10.1029/2012GL054008, 2012.
- Shiraiwa, M., Ammann, M., Koop, T., and Pöschl, U.: Gas uptake and chemical aging of semisolid organic aerosol particles, *Proc. Natl. Acad. Sci.*, 108, 11 003–11 008, doi:10.1073/pnas.1103045108, 2011.
- Shiraiwa, M., Pfrang, C., Koop, T., and Pöschl, U.: Kinetic multi-layer model of gas-particle interactions in aerosols and clouds (KM-GAP): linking condensation, evaporation and chemical reactions of organics, oxidants and water, *Atmos. Chem. Phys.*, 12, 2777–2794, doi:10.5194/acp-12-2777-2012, 2012.
- 35 Shiraiwa, M., Zuend, A., Bertram, A. K., and Seinfeld, J. H.: Gas-particle partitioning of atmospheric aerosols: interplay of physical state, non-ideal mixing and morphology, *Phys. Chem. Chem. Phys.*, 15, 11 441–11 453, doi:10.1039/C3CP51595H, 2013.

- Shiraiwa, M., Li, Y., Tsimpidi, A. P., Karydis, V. A., Berkemeier, T., Pandis, S. N., Lelieveld, J., Koop, T., and Pöschl, U.: Global distribution of particle phase state in atmospheric secondary organic aerosols, *Nature Communications*, 8, 15 002, doi:10.1038/ncomms15002, 2017.
- Shrivastava, M., Cappa, C., Fan, J., Goldstein, A., Guenther, A., Jimenez, J., Kuang, C., Laskin, A., Martin, S., Ng, N., Petaja, T., Pierce, J., Rasch, P., Roldin, P., Seinfeld, J., Shilling, J., Smith, J., Thornton, J., Volkamer, R., Wang, J., Worsnop, D., Zaveri, R., Zelenyuk, A., and Zhang, Q.: Recent advances in understanding secondary organic aerosol: Implications for global climate forcing, *Rev. Geophys.*, 55, 509–559, doi:10.1002/2016RG000540, 2017.
- Song, M., Liu, P., Hanna, S., Zaveri, R., Potter, K., You, Y., Martin, S., and Bertram, A.: Relative humidity-dependent viscosity of secondary organic material from toluene photo-oxidation and possible implications for organic particulate matter over megacities., *Atmos. Chem. Phys.*, 16, 8817–8830, doi:10.5194/acp-16-8817-2016, 2016a.
- 10 Song, Y. C., Haddrell, A. E., Bzdek, B. R., Reid, J. P., Bannan, T., Topping, D. O., Percival, C., and Cai, C.: Measurements and predictions of binary component aerosol particle viscosity, *J. Phys. Chem. A*, 120, 8123–8137, doi:10.1021/acs.jpca.6b07835, 2016b.
- Sun, Q. and Wexler, A. S.: Modeling urban and regional aerosols - condensation and evaporation near acid neutrality, *Atmos. Environ.*, 32, 3527 – 3531, doi:10.1016/S1352-2310(98)00059-4, 1998.
- Vaden, T. D., Imre, D., Beránek, J., Shrivastava, M., and Zelenyuk, A.: Evaporation kinetics and phase of laboratory and ambient secondary organic aerosol, *Proc. Natl. Acad. Sci.*, 108, 2190–2195, doi:10.1073/pnas.1013391108, 2011.
- 15 Virtanen, A., Joutsensaari, J., Koop, T., Kannosto, J., Yli-Pirilä, P., Leskinen, J., Mäkelä, J. M., Holopainen, J. K., Pöschl, U., Kulmala, M., Worsnop, D. R., and Laaksonen, A.: An amorphous solid state of biogenic secondary organic aerosol particles, *Nature*, 467, 824–827, doi:10.1038/nature09455, 2010.
- Wexler, A. S. and Seinfeld, J. H.: The distribution of ammonium salts among a size and composition dispersed aerosol, *Atmos. Environ.*, 24, 1231 – 1246, doi:10.1016/0960-1686(90)90088-5, 1990.
- 20 Yarwood, G., Rao, S., Yocke, M., and Whitten, G.: Updates to the carbon bond chemical mechanism: CB05. Rep. RT-0400675, available at http://www.camx.com/files/cb05_final_report_120805.aspx, last access 21 September 2018, 2005.
- Yatavelli, R. L. N., Stark, H., Thompson, S. L., Kimmel, J. R., Cubison, M. J., Day, D. A., Campuzano-Jost, P., Palm, B. B., Hodzic, A., Thornton, J. A., Jayne, J. T., Worsnop, D. R., and Jimenez, J. L.: Semicontinuous measurements of gas–particle partitioning of organic acids in a ponderosa pine forest using a MOVI-HRToF-CIMS, *Atmos. Chem. Phys.*, 14, 1527–1546, doi:10.5194/acp-14-1527-2014, 2014.
- 25 Zhang, Y., Pun, B., Vijayaraghavan, K., Wu, S.-Y., Seigneur, C., Pandis, S. N., Jacobson, M. Z., Nenes, A., and Seinfeld, J. H.: Development and application of the Model of Aerosol Dynamics, Reaction, Ionization, and Dissolution (MADRID), *J. Geophys. Res.*, 109, D01 202, doi:10.1029/2003JD003501, 2004.
- 30 Zhu, S., Sartelet, K., Healy, R., and Wenger, J.: Simulation of particle diversity and mixing state over Greater Paris: A model-measurement inter-comparison, *Faraday Discuss.*, 189, 547–566, doi:10.1039/C5FD00175G, 2016.
- Zuend, A., Marcolli, C., Luo, B. P., and Peter, T.: A thermodynamic model of mixed organic-inorganic aerosols to predict activity coefficients, *Atmos. Chem. Phys.*, 8, 4559–4593, doi:10.5194/acp-8-4559-2008, 2008.