

Responses to the Co-Editor Prof. Jayanarayanan Kuttippurath for his important suggestions of our final stage manuscript (acp – 2018-201) titled *“Climate and the September 2014 flood event over Mountainous Jammu and Kashmir, India: Physical explanations based on observations and modelling”* by Sumira et al. submitted for publication in the journal *Atmospheric Chemistry and Physics*, Copernicus publications.

General response:

We are indebted to the Co-Editor Prof. Jayanarayanan Kuttippurath for his important suggestions about improving the quality of the figures, adding information about trends in the rainfall and temperature in the abstract and correcting typos in the final stage of this manuscript. We hope that the Co-Editor will be convinced of this final stage manuscript. Manuscript with changes tracked version also enclosed for quick reference. If any further improvement is needed for figures, we will surely make our sincere attempts so that ACP journal quality figures will be produced.

One to one responses:

The responses are in bold faceted.

1. The figures are of very poor quality. Please improve the quality of the figures for the final upload. Figures 1-6, the font size of the axes are too small to read. Please consider to re-draw them. This is a very important study for that region. Therefore, please take care all aspects of the manuscript. Thank you.

The quality of all the figures is improved now and if any further improvement is needed, we will surely do it during production stages for ACP publication quality figures.

2. Abstract: You have mentioned the increase and decrease in precipitation, please provide the trend values with uncertainty there. Also, add uncertainty values (error range or stddev) with temperature trends.

The following sentences are now added

In the present study, the observed long-term trends in temperature (°C/year) and precipitation (mm/year) along with their respective standard errors during 1980-2016 are as follows: (1) 0.05 (0.01) and -16.7 (6.3) for Gulmarg, (2) 0.04 (0.01) and -6.6 (2.9) for Srinagar, (3) 0.04 (0.01) and -0.69 (4.79) for Kokernag, (4) 0.04 (0.01) and -0.13 (3.95) for Pahalgam, (5) 0.034 (0.01) and -5.5 (3.6) for Kupwara and (6) 0.01 (0.01) and -7.96 (4.5) for Quazigund.

3. There are still some technical errors, but this would be taken care of during the proof reading. However, if you can correct some, correct those to speed-up the publication process. e.g. Line 288, line 331, line 454,

All typos are now corrected.

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Analyses of temperature and precipitation in the Indian Jammu-Kashmir for the 1980–2016 period: Implications for remote influence and extreme events

Sumira Nazir Zaz¹, Romshoo Shakil Ahmad¹, Ramkumar Thokuluwa Krishnamoorthy^{2*}, and
Yesubabu Viswanadhapalli²

1. Department of Earth Sciences, University of Kashmir, Hazratbal, Srinagar,
Jammu and Kashmir-190006, India
2. National Atmospheric Research Laboratory, Dept. of Space, Govt. of India, Gadanki, Andhra Pradesh
517112, India

Email: zaz.sumira@gmail.com, shakilrom@kashmiruniversity.ac.in, tkram@narl.gov.in,
yesubabu@narl.gov.in;

***Corresponding author** (tkram@narl.gov.in)

Abstract

Local weather and climate of the Himalayas are sensitive and interlinked with global scale changes in climate, as the hydrology of this region is mainly governed by snow and glaciers. There are clear and strong indicators of climate change reported for the Himalayas, particularly the Jammu and Kashmir region situated in the western Himalayas. In this study, using observational data, detailed characteristics of long- and short-term as well as localized variations of temperature and precipitation are analysed for these six meteorological stations, namely, Gulmarg, Pahalgam, Kokernag, Qazigund, Kupwara and Srinagar of Jammu and Kashmir, India during 1980-2016. In addition to analysis of stations observations, we also utilized the dynamical downscaled simulations of WRF model and ERA-Interim (ERA-I) data for the study period. The annual and seasonal temperature and precipitation changes were analysed by carrying out Student's t-test, Mann-Kendall, Linear regression and Cumulative deviation statistical tests. The results show an increase of 0.8°C in average annual temperature over thirty seven years (from 1980 to 2016) with higher increase in maximum temperature (0.97°C) compared to minimum temperature (0.76°C). Analyses of annual mean temperature at all the stations reveal that the high-altitude stations of Pahalgam (1.13°C) and Gulmarg (1.04°C) exhibit a steep increase and statistical significant trends. ~~The overall p~~Precipitation patterns in the valley ~~such as at Gulmarg and Pahalgam~~ shows ~~significant~~a slight and definite decrease in the annual rainfall ~~at Gulmarg and Pahalgam stations~~. Seasonal analyses show significant increasing trends in the winter and spring temperatures at all stations with prominent decrease in spring precipitation. ~~In the present study, the observed long-term trends in temperature (°C/year) and precipitation (mm/year) along with their respective standard errors during 1980-2016 are as follows: (1) 0.05 (0.01) and -16.7 (6.3) for Gulmarg, (2) 0.04 (0.01) and -6.6 (2.9) for Srinagar, (3) 0.04 (0.01) and -0.69 (4.79) for Kokernag, (4) 0.04 (0.01) and -0.13 (3.95) for Pahalgam, (5) 0.034 (0.01) and -5.5~~

(3.6) for Kupwara and (6) 0.01 (0.01) and -7.96 (4.5) for Quazigund. The present study also reveals that variation in temperature and precipitation during winter (December - March) has a close association with the North Atlantic Oscillation (NAO). Further, the observed temperature data (monthly averaged data for 1980-2016) at all the stations show good correlation of 0.86 with the results of WRF and therefore the model downscaled simulations are considered as a valid scientific tool for the studies of climate change in this region. Though the correlation between WRF model and observed precipitation is significantly strong, the WRF model underestimates significantly the rainfall amount, which necessitates the need for the sensitivity study of the model using the various microphysical parameterization schemes. The potential vorticities in the upper troposphere troposphere are obtained from ERA-I over the Jammu and Kashmir region indicate that the extreme weather event of September 2014 occurred due to breaking of intense atmospheric Rossby wave activity over Kashmir. As the wave could transport a large amount of water vapour from both the Bay of Bengal and Arabian Sea and dump them over the Kashmir region through wave breaking, it is probably resulted in the historical devastating flooding of the whole Kashmir valley in the first week of September 2014. This was accompanied by extreme rainfall events measuring more than 620 mm in some parts of the Pir Panjal range in the South Kashmir.

1. Introduction

Climate change is a real Earth's atmospheric and surface phenomenon and the influences of which on all spheres of life are considered significant almost everywhere in the world in the past few decades. Extreme weather events like anomalously large floods and unusual drought conditions associated with changes in climate play havoc with livelihoods of even established civilizations particularly in the coastal and high-mountainous areas. Jammu and Kashmir, India, located in the Western Himalayan region, is one such cataclysmic mountainous region where significant influence of climate change on local weather has been observed for the last few decades; (1) shrinking and reducing glaciers, (2) devastating floods, (3) decreasing winter duration and rainfall, and (4) increasing summer duration and temperature (Solomon et al., 2007; Kohler and Maselli, 2009; Immerzeel et al., 2010; Romshoo et al., 2015; Romshoo et al., 2017). Western disturbances (WD) is considered as one of the main sources of winter precipitation for the Jammu and Kashmir region, which brings water vapour mainly from the tropical Atlantic Ocean, Mediterranean Sea, Caspian Sea and Black sea. Though WD is perennial, it is most intense during northern winter (December-February; Demri et al., 2015). Planetary-scale atmospheric Rossby-waves have potential to significantly alter the distribution and movement of WD according to their intensity and duration (few to tens of days). Since WD is controlled by planetary-scale Rossby waves in the whole troposphere of the subtropical region, diagnosing different kinds of precipitation characteristics is easier with the help of potential vorticity (PV) at 350K potential temperature (PT) and 200 hPa level pressure surface (PS) as they are considered as proxies for Rossby wave activities (Ertel, 1942; Bartels et al., 1998; Demri et al., 2015 and Hunt et al 2018a). Here onwards, it will be simply called PV at 350 K and 200 hPa surfaces. For example, (Postel and Hitchman, 1999; Hunt 2018b) studied the characteristics of Rossby wave breaking (RWB) events occurring at 350K surface transecting the subtropical

westerly jets. Similarly, Waugh and Polvani (2000) studied RWB characteristics at 350K surface in the Pacific region during northern fall–spring with emphasis on their influence on westerly ducts and their intrusion into the tropics. Since PV is a conserved quantity on isentropic and isobaric surfaces (ISOES & ISOBS) when there is no exchange of heat and pressure respectively, it is widely used for investigating large-scale dynamical processes associated with frictionless and adiabatic flows. Moreover, all other dynamical parameters, under a given suitable balanced-atmospheric-background condition, can be derived from PV and boundary conditions (Hoskins et al., 1985).

Divergence of the atmospheric air flows near the upper troposphere is larger during precipitation, leading to increase in the strength of PV. Because of which, generally there will be a good positive correlation between variations in the strength of PV in the upper troposphere and precipitation over the ground provided that the precipitation is mainly due to the passage of large-scale atmospheric weather systems like western disturbances, and monsoons. Wind flows over topography can significantly affect the vertical distribution of water vapour and precipitation characteristics. Because of this, positive correlation between variations in PV and precipitation can be modified significantly.. These facts need to be taken into account while finding long-term variations of precipitation near mountainous regions like the western Himalaya. The interplay between the flow of western disturbances and topography of the western Himalaya complicates further the identification of source mechanisms of extreme weather events (Das et al., 2002; Shekhar et al., 2010) like the ones that occurred in the western Himalayan region; Kashmir floods in 2014 and Leh floods in 2010 in the Jammu and Kashmir region and Uttarakhand floods in 2013. Kumar et al. (2015) also noted that major flood events in the Himalayas are related to changing precipitation intensity in the region. This necessitates making use of proper surrogate parameters like PV and distinguish between different source mechanisms of extreme weather events associated with both the long-term climatic impacts of remote origin and short-term localized ones like organized convection (Romatschke and Houze 2011; Rasmussen and Houze 2012; Houze and Rasmussen 2016; Martius et al., 2012).

The main aim of the present study is to investigate long-term (climate) variation of surface temperature and precipitation over the Jammu and Kashmir, India region of the western Himalayas in terms of its connections with NAO and atmospheric Rossby wave activity in the upper troposphere. Since PV is considered as a measure of Rossby wave activity, the present work analyses in detail, for a period of 37 years during 1980-2016, monthly variation of PV (ERA-interim reanalysis data, Dee et al., 2001) in the upper troposphere (at 350 K and 200 hPa surfaces) and compares it with observed surface temperature and rainfall (India Meteorological Department, IMD) at six widely separated mountainous locations with variable orographic features (Srinagar, Gulmarg, Pahalgam, Qazigund, Kokarnag and Kupwara). There exist several reports on climatological variation of meteorological parameters in various parts of the Himalayas. For example, Kumar and Jain (2009) and Bhutiyani et al. (2010) found an increase in the temperature in the north-western Himalayas with significant variations in precipitation patterns. Archer and Fowler (2004) examined temperature data of seven stations in the Karakoram and Hindu Kush Mountains of the Upper Indus River Basin (UIRB) in search of seasonal and annual trends using statistical test like

regression analysis. Their results revealed that mean winter maximum-temperature has increased significantly while mean summer minimum-temperature declined consistently. On the contrary, Lui et al. (2009) examined long-term trends in minimum and maximum temperatures over the Tibetan mountain range during 1961-2003 and found that minimum temperature increases faster than maximum temperature in all the months. Romshoo et al. (2015) observed changes in snow precipitation and snow-melt-runoff in the Kashmir valley and attributed the observed depletion of stream flow to the changing climate in the region. Bolch et al. (2012) reported that the glacier extent in the Karakoram range is increasing.

These contrasting findings of long-term variations in temperature and precipitation in the Himalayas need to be verified by analyzing long-term climatological data available in the region. However, the sparse and scanty availability of regional climate data pose challenges in understanding the complex microclimate in this region. Therefore, studying the relationship of recorded regional (Jammu and Kashmir) climatic variations in temperature and precipitation with remote and large-scale weather phenomena such as the North Atlantic Oscillation (NAO), and El Niño Southern Oscillation (ENSO) is necessary for understanding the physical processes that control the locally observed variations (Ghashmi, 2015). Archer and Fowler (2004) and Iqbal and Kashif (2013) found that large-scale atmospheric circulation like NAO influences significantly the climate of the Himalayas. However, detailed information about variation in temperature and precipitation and its teleconnection with observed variations of NAO is inadequately available for this part of the Himalayan region (Kashmir Valley).

2. Geographical setting of Kashmir

The inter mountainous valley of Kashmir has a unique geographical setting and it is located between the Greater Himalayas in the north and Pir Panjal ranges in the south, roughly within the latitude and longitude ranges of $33^{\circ} 55'$ to $34^{\circ} 50'$ and $74^{\circ} 30'$ to $75^{\circ} 35'$ respectively (Fig.1). The heights of these mountains range from about 3,000 to 5,000 m and the mountains strongly influence the weather and climate of the region. Generally the topographic setting of the six stations, though variable, could be broadly categorized into two; (1) stations located on plains (Srinagar, Kokarnag, Qazigund and even Kupwara) and (2) those located in the mountain setting (Gulmarg, Pahalgam). Physiographically, the valley of Kashmir is divided into three regions; Jhelum valley floor, Greater Himalayas and Pir Panjal. In order to represent all the regions of the valley, six meteorological stations located widely with different mean sea levels (msl), namely, Gulmarg (2740m), Pahalgam (2600m), Kokarnag (2000m), Srinagar (1600m), Kupwara (1670m) and Qazigund (1650m) were selected for analyses of observed weather parameters.

The Kashmir valley is one of the important watersheds of the upper Indus basin harbouring more than 105 glaciers and it experiences the Mediterranean type of climate with marked seasonality (Romshoo and Rashid, 2014). Broadly, four seasons (Khattak et al 2011; Rashid et al., 2015) are defined for the Kashmir valley; winter (December to February), spring (March to May), summer (June to August), and autumn (September to November). It is to be

clarified here that while defining the period of NAO (Fig. 4) it is considered December-March as winter months as defined by Archer and Fowler (2004) and Iqbal and Kashif (2013) and in all other parts of the manuscript it is December-February as per the IMD definition. The annual temperature in the valley varies from about -10°C to 35°C. The rainfall pattern in the valley is dominated by winter time precipitation associated with western disturbances (Dar et al., 2014) while the snow precipitation is received mainly in winter and early spring season (Kaul and Qadri, 1979).

2. Data and Methodology

India Meteorological Department (IMD) provided 37 years (1980-2016) of data of daily precipitation, maximum and minimum temperatures for all the six stations. Monthly averaged data were further analysed to find long-term variations of weather parameters. Statistical tests including Mann-Kendall, Spearman Rho, Cumulative deviation, Student's t-test were performed to determine long term-trends and turning point of weather parameters with statistical significances. Similar analyses and tests were performed also for the Weather and Research Forecasting (WRF) model simulated and ERA-Interim reanalyses data (0.75° by 0.75° spatial resolution in the horizontal plane, monthly averaged time resolution) of same weather parameters and for the NAO index. Brief information about these data sets is provided below.

3.1 Measurements and model simulations

The obtained observational data are analysed carefully for homogeneity and missing values. Analyses of ratios of temperature from the neighbouring stations with the Srinagar station were conducted using relative homogeneity test (WMO, 1970). It is found that there is no significant inhomogeneity and data gap for any station. Few missing data points were linearly interpolated and enough care was taken not to make any meaningful interpretation during such short periods of data gaps in the observations. Annual and seasonal means of temperature and precipitation were calculated for all the stations and years. To compute seasonal means, the data were divided into the following seasons: winter (December to February), spring (March to May), summer (June to August) and autumn (September to November). Trends in the annual and seasonal means of temperature and precipitation were determined using Mann-Kendall (non-parametric test) and linear regression tests (parametric test) at the confidence levels of $S = 99\%$ or (0.01), $S = 95\%$ or 0.05 and $S = 90\%$ or 0.1. These tests have been extensively used in hydro meteorological data analyses as they are less sensitive to heterogeneity of data distribution and least affected by extreme values or outliers in data series. Various methods have been applied to determine change points of a time series (Radziejewski et al., 2000; Chen and Gupta, 2012). In this study, change point in time series of temperature and precipitation was identified using cumulative deviation test and Student's t test (Pettitt, 1979). This method

detects the time of significant change in the mean of a time series when the exact time of the change is unknown (Gao et al., 2011).

Winter NAO index during 1980–2010 were obtained for further analyses from Climatic Research Unit through the web link <https://www.cru.uea.ac.uk/data>. The winter (December - March) NAO index is based on difference of normalized sea level pressure (SLP) between Lisbon, Portugal and Iceland, which is available from 1964 onwards. Positive NAO index is associated with stronger-than-average westerlies over the middle latitudes (Hurrell, 1997). Correlation between mean (December-March) temperature, precipitation and NAO index was determined using Pearson correlation coefficient method. To test whether the observed trends in winter temperature and precipitation are enforced by NAO, linear regression analysis (forecast) was performed (Fig. 4e and f). The following algorithm calculates or predicts a future value by using existing values. The predicted value is a y-value for a given w-value. The known values are existing w-values and y-values, and the new value is predicted by using linear regression.

The syntax is as follows

FORECAST(x, known_y's, known_w's)

W is the data point for which we want to predict a value.

Known_y's is the dependent array or range of data (rainfall or temperature).

Known_w's is the independent array or range of data (time).

The equation for FORECAST is $a + bw$, where:

$$a = \hat{y} - b\hat{w} \quad \text{and} \quad b = \frac{\sum(w - \hat{w})(y - \hat{y})}{\sum(w - \hat{w})^2}$$

and where \hat{w} and \hat{y} are the sample means AVERAGE (known_w's) and AVERAGE (known y's).

3.2. WRF Model configuration

The Advanced Research WRF version 3.9.1 model simulation was used in this study to downscale the ERA-Interim (European Centre for Medium Range Weather Forecasting Re Analysis) data over the Indian Monsoon region. The model is configured with 2 two-way nested domains (18 km and 9-km horizontal resolutions), 51 vertical levels and model top at 10 hPa level. The model first domain extends from longitude from 24.8516 E to 115.148E and latitude from 22.1127S to 46.7629 N while the second domain covers the longitudes from 56.3838E to 98.5722E and latitudes from 3.86047 S to 38.2874 N.

The initial and boundary conditions supplied to WRF model are obtained from ERA-Interim 6-hourly data. Model physics used in the study for boundary layer processes is Yonsei University's non-local diffusion scheme (Hong et al., 2006), the Kain-Fritsch scheme for cumulus convection (Kain and Fritsch, 1993), Thomson scheme for microphysical processes, the Noah land surface scheme (Chen and Dudhia, 2001) for surface processes, Rapid Radiation Transfer Model (RRTM) for long-wave radiation (Mlawer et al., 1997), and the Dudhia (1989) scheme for short-wave radiation. The physics options configured in this study are adopted based on the previous studies of heavy rainfall and Monsoon studies over the Indian region (Srinivas et al., 2013, Madala et al., 2016, Ghosh et al., 2016; Srinivas et al., 2018).

For the present study, the WRF model is initialized on daily basis at 12 UTC using ERA-Interim data and integrated for a 36-hour period using the continuous re-initialization method (Lo et al., 2008; Langodan, et al., 2016; and Viswanadhapalli et al., 2017). Keeping the first 12-hours as model spin-up time, the remaining 24-hour daily simulations of the model are merged to get the data during 1980-2016. To find out the skill of the model, the downscaled simulations of WRF model are validated for six IMD surface meteorological stations. The statistical skill scores such as bias, mean error (ME) and root mean square error (RMS) were computed for the simulated temperature against the observed temperature data of IMD.

4. Results and Discussion:

4.1. Trend in annual and seasonal temperature

Tables 1& 2 show the results of statistical tests (Mann-Kendall and linear regression, cumulative deviation and Student's t) carried out on the temperature and precipitation data respectively. All the parametric and nonparametric tests carried out for the trend analysis and abrupt changes in the trend showed almost similar results. Table 1, therefore shows results of representative tests where higher values of statistical significance between Mann-Kendall/linear regression test and Cumulative deviation/student's t test are considered. It is evident that there is an increasing trend at different confidence levels in annual and seasonal temperatures of all the six stations (Pahalgam, Gulmarg, Kokarnag, Srinagar, Kupwara and Qazigund), located in different topographical settings (Table 3). During 1980-2016, Pahalgam and Gulmarg, located at higher elevations of about 2500m amsl (above mean sea level), registered statistically significant increase in average annual temperature by 1.13°C and 1.04°C (Fig. 2a). It is to be noted that hereafter it will not be mentioned explicitly about the period 1980-2016 and statistically significant means the confidence level is about 90%. Kokarnag and Kupwara, located at the heights of about 1800-2000m amsl, showed an increase of 0.9°C and 1°C respectively (Fig. 2a). However, Srinagar and Qazigund, located at the heights of about 1700m-1600m amsl, exhibited an increase of 0.65°C and 0.44°C (Fig. 2a).

Analyses of maximum and minimum temperatures (Table 1 and Fig. 2b) for the six stations reveal higher rate increase in maximum temperature. Pahalgam and Kupwara recorded the highest rise of $\sim 1.3^{\circ}\text{C}$ followed by Kokarnag (1.2°C) and Srinagar (1.1°C). The exception is that Gulmarg and Qazigund (being a hilly station) shows less than 0.6°C in maximum temperature. The minimum temperature exhibits a lowest increase of 0.3°C at Srinagar and highest increase at Gulmarg station of 1.2°C (Fig. 2c). Analyses of composite seasonal mean of minimum and maximum temperatures in the valley reveal higher increase in maximum temperature in winter and spring seasons. Among four stations (Gulmarg, Pahalgam, Kokarnag and Kupwara), Gulmarg indicates an increase of less than 1°C while Pahalgam, Kokarnag and Kupwara shows an increase of 0.9°C , 0.9°C and less than 0.9°C respectively (Table 1 and Fig. 2d). On the contrary, Qazigund and Srinagar showed a slight increase of less than 0.4°C and 0.5°C respectively. Mean spring-temperature shows higher rise comparing to other seasons temperatures for all the stations. Gulmarg shows an increase of less than 1.4°C . Pahalgam, Kupwara, Kokarnag showed increase of 1.3°C at $S = 0.01$. Qazigund and Srinagar revealed 0.6°C and 1°C increase respectively as shown in the Table 1 and Fig. 2e. In summer, the temperature rise for Pahalgam is about less than 0.6°C and for Gulmarg and Qazigund, it is about 0.4°C and 0.2°C respectively (Table 1). Kupwara, Kokarnag and Srinagar reveal an increase of less than 0.3°C , 0.4°C and 0.1°C respectively (Fig. 2f). In Autumn, Gulmarg shows an increase of 0.9°C and Pahalgam exhibit less than 0.6°C . On the contrary Qazigund shows less than 0.4°C at while Srinagar shows no significant increase in observed temperatures (Fig. 2g and Table 1).

4.2 Trend in annual and seasonal precipitation

The annual precipitation pattern of the valley is comparable to that of temperature with higher decrease observed at the upper elevation stations of Gulmarg and Pahalgam (Fig. 3a and Table 2). Similar to temperature, Table 2 provides in detail the test results of Mann-Kendall, linear regression and Student's t . While Kokarnag and Kupwara show significant decrease, the lower elevated stations, Qazigund and Srinagar, exhibit insignificant decrease (Fig. 3a). Winter precipitation decrease is maximum at Gulmarg and Kokarnag followed by Kupwara and Pahalgam and it is insignificant decrease for Srinagar and Qazigund (Table 2 and Fig. 3b). The spring season precipitation exhibits decreasing trend for all the six stations with the lowest decrease of 42mm precipitation at Kupwara (Table 2).

During summer months also, precipitation shows decreasing trend for all stations except Qazigund that it is statistically insignificant (Fig. 3d, and Table 2). For Qazigund there is no apparent trend in summer precipitation. The autumn precipitation also shows insignificant decreasing trend for the stations (Fig. 3e and Table 2). Cumulative test was used to determine the "change point" of trend in the annual and seasonal variations of temperature and precipitation. Results reveal that the year 1995 is the year of abrupt increase (change point) in

temperature of the valley (Fig. 4a) and the same year is identified as the year of abrupt decrease for precipitation (Fig. 4b).

4.3 Influence of North Atlantic Oscillation (NAO) on the winter precipitation over the Kashmir valley

The present study also investigates the tele-connection between the activity of North Atlantic Oscillation (NAO) and the variations in temperature and precipitation over the Kashmir valley, particularly during winter season (December - March). It is found that there is a significant negative/positive correlation (-0.54/0.68) between NAO (NAO index) and precipitation/temperature (Fig. 4c). This suggests that winter precipitation and temperature over the Kashmir valley has a close association with the winter NAO. Higher precipitation over Kashmir is associated with positive phase of NAO. Further the “change point” year, 1995, in the trend of temperature and precipitation coincides with that of the NAO index. To test whether the trends in temperatures and precipitation over the Kashmir valley are forced by the NAO, regression analysis was performed on winter temperature and precipitation (Figs. 4e and f) and the results indicate that there is a significant connection between NAO and precipitation over Kashmir,

The observed annual and seasonal variation of temperature at all stations except Qazigund is strongly correlated with WRF downscaled simulations. Overall, the simulations show correlation of 0.66, 0.67, 0.72, 0.62, 0.79 and 0.47 for Srinagar, Gulmarg, Kokarnag, Kupwara, Pahalgam and Qazigund respectively. The annual mean simulated temperature shows very good correlation (0.85) with observations. Figure 5 shows annual and seasonal correlations between trends of observed and simulated temperatures (location of Kokarnag is considered for WRF data). However, root mean square error (RMSE) analysis indicates that model simulations underestimate slightly the observations by an average value of -0.43°C. Similar to Figure 5, Figure 6 shows the comparison between WRF model simulated and observed precipitation. Even though the trend is similar, WRF model severely underestimates the rainfall amount. A detailed study on this topic will be presented in a separate paper.

4.4. Discussion

The Himalayan mountain system is quite sensitive to global climate change as the hydrology of the region is mainly dominated by snow and glaciers, making it one of the ideal sites for early detection of global warming (Solomon et al., 2007; Kohler and Maselli, 2009). Various reports claim that in the Himalayas significant warming had occurred in the last century (Fowler and Archer, 2006; Bhutiyani et al., 2007). Shrestha et al. (1999) analysed

surface temperature at 49 stations located across the Nepal Himalayas and the results indicate warming trends in the range of 0.06 to 0.12°C per year. The observations of the present study are in agreement with the studies carried out by Shrestha et al. (1999), Archer and Fowler (2004) and Butiyani (2007). In the present study, it is observed that rise in temperature is larger at higher altitude stations of Pahalgam (1.13°C) and Gulmarg (1.04°C) and it is about 0.9°C, 0.99°C, 0.04°C, and 0.10°C for the other stations, Kokarnag, Kupwara, Srinagar and Qazigund respectively during 1980-2016. Liu et al. (2009) and Liu and Chen (2000) also report higher warming trends at higher altitudes in the Himalayan regions. In the future, the impacts of climate change will be intense at higher elevations and in regions with complex topography, which is consistent with the model results of Wiltshire (2013).

The noteworthy observation in the present study is that statistically significant steep increase in the temperature (change point) occurred in the year 1995 and it has been continuing thereafter. The mega El Niño in 1998 has been considered as one of the strongest El Niño's in history that led worldwide increase in temperature (Epstein et al., 1998). Contrastingly, the El Niño in 1992 led to a decrease in temperature throughout the northern hemisphere, which is ascribed to the Mt. Pinatubo volcanic eruption (Swanson et al., 2009; IPCC, 2013). Also this event interrupted the direct sunlight to reach on the surface of the earth for about two months (Barnes et al., 2016).

Studies of trends in seasonal-mean temperature in many regions across the Himalayas indicate higher warming trends in winter and spring months (Shrestha et al., 1999; Archer and Fowler, 2004; Butiyani, 2009). The seasonal difference found in the present study is consistent with other studies carried out for the Himalayas (Archer and Fowler, 2004; Sheikh et al., 2009 and Roe et al., 2003), Lancang Valley, China (Yunling and Yiping, 2005), Tibet (Liu and Chen, 2000) and the Swiss Alps (Beniston et al., 2010), where almost all stations recorded higher increase in the winter and spring temperatures comparing to autumn and summer temperatures. Recent studies found that reducing the extent depth of snow cover and shrinking glaciers may also be one of the contributing factors for the observed higher warming, as the reduction in the percentage of snow and glacier can alter the surface albedo over a region, which in turn can increase the surface air temperatures (Kulkarni et al., 2002; Groisman et al., 1994). Romshoo et al. (2015) and Murtaza and Romshoo (2016) have also reported that reduction of snow and glacier cover in the Kashmir regions of the Himalayas during the recent decades could be one of the reasons of occurrence of higher warming particularly on the higher elevated stations of Gulmarg and Pahalgam.

In the Himalayan mountain system, contrasting trends have been noted in precipitation over the recent decades (IPCC, 2001). Borgaonkar et al. (2001), Shrestha et al. (2000) and Archer and Fowler (2004) observed increasing precipitation patterns over the Himalayas while Mooley and Parthasarathy (1983), Kumar and Jain (2009) and Demri and Dash (2012) reported large-scale decadal variation with increasing and decreasing precipitation periods. The results of the present study indicate that decrease in annual precipitation is slightly insignificant at all the six stations except the spring season. Increasing trend in temperature can trigger large-scale energy exchanges that become more intricate as complex topography alters the precipitation type and intensity in many ways (Kulkarni et al., 2002; Groisman et al., 1994). Climate model simulations (Zarenistana et al., 2014; Rashid et al.,

2015) and empirical evidence (Vose et al., 2005; Romshoo et al., 2015) also confirm that increasing temperature results in increased water vapour leading to more intense precipitation events even when the total annual precipitation reduces slightly. The increase in temperature therefore enhances the risks of both floods and droughts. For example, the disaster flood event of September 2014 occurred in the Kashmir valley due to high frequency and high intense precipitation.

The North Atlantic Oscillation (NAO) is one of the strongest northern atmospheric weather phenomena occurring due to the difference of atmospheric pressure at sea level between the Iceland low and Azores high. It controls the strength and direction of westerly winds across the northern hemisphere. Surface temperatures have increased in the northern hemisphere in the past few decades (Mann et al., 1999; Jones et al., 2001; Hijioka et al., 2014), and the rate of warming has been especially high ($\sim 0.15^{\circ}\text{C decade}^{-1}$) in the past 40 years (Folland et al., 2001; Hansen et al., 2001; Peters et al., 2013; Knutti et al., 2016). NAO causes substantial fluctuations in the climate of the Himalayas (Hurrell, 1997; Syed et al., 2006; Archer and Fowler, 2004). Several workers found a strong connection between the NAO and temperature and precipitation in the north-western Himalayas (Archer and Fowler, 2004; Bhutiyani et al., 2007; Bookhagen, 2010; Sharif et al., 2012; Iqbal and Kashif, 2013). A substantial fraction of the most recent warming is linked to the behaviour of the NAO (Hurrell, 1997; Thompson et al., 2003; Madhura et al., 2015). The climate of the Kashmir Himalayas is influenced by western disturbances in winter and spring seasons. Figs. 4c and d show correlation between winter time NAO and temperature and precipitation over the Kashmir region. While temperature shows negative correlation of -0.54, precipitation shows positive correlation of 0.68. From linear regression analyses, it is found that considerable variation in winter precipitation and temperature over Kashmir is forced by winter NAO. The weakening link of NAO after 1995 has a close association with decreased winter precipitation and increased winter temperature in the valley. Similarly, Bhutiyani et al. (2009) and Dimri and Dash (2012) also found statistically significant decreasing trend in precipitation which they related to weakening of NAO index. However, for establishing a detailed mechanism incorporating these variations requires thorough investigation.

The WRF model simulations compare well with observations (significantly strong correlation of 0.85) and the correlation is more forelevated stations than valley stations of Srinagar and Kupwara. However, it is expected that the good correlation can result if more precise terrain information is incorporated in the WRF model simulations. Earlier studies (e.g. Kain and Fritsch, 1990, 1993; Kain, 2004) also found good correlation between observed and WRF simulated rainfall events. In conjunction with large-scale features such as NAO and ENSO, it can result in large-scale variability in the climate of this region (Ogura and Yoshizaku, 1988). Furthermore, incorporation of mesoscale teleconnections and their associations in the WRF model can further help in understanding large-scale weather forecasting over this region.

4.5. Physical mechanisms of climate and weather of Jammu & Kashmir

443 Large-scale spatial and temporal variations in the meridional winds could be due to the passage of
 444 planetary-scale Rossby waves (RW) in the atmospheric winds. When RWs break in the upper troposphere, it could
 445 lead to vertical transport of atmospheric air between the upper troposphere and lower stratosphere and an
 446 irreversible horizontal transport of air mass between the subtropics and extra tropics (McIntyre and Palmer, 1983).
 447 Rossby waves have the characteristic of remaining coherent over many days and propagate long distances of the
 448 order of synoptic to planetary scales leading to tele-connection of remote atmospheres of global extent. The study
 449 by Chang and Yu (1999) indicates that during northern winter months of December–January–February, Rossby
 450 wave packets can be most coherent over a large distance of from the northern Africa to the Pacific through the
 451 southern Asia. There are reports on extreme weather events connected to Rossby waves of synoptic to planetary
 452 scales in the upper troposphere (e.g. Screen and Simmonds, 2014). In northern India, there is an increasing trend in
 453 heavy rainfall events, particularly over the Himachal Pradesh, Uttarakhand and Jammu and Kashmir (Sinha Ray and
 454 Srivastava, 2000; Nibanupudi et al., 2015). Long-scale Rossby waves can lead to generation of alternating
 455 convergence and divergence in the upper troposphere that in turn can affect surface weather parameters like
 456 precipitation through generation of instabilities in the atmospheric air associated with convergence and divergence
 457 (Niranjankumar et al., 2016).
 458

459
 460 Using observations and MERRA (Modern-Era Retrospective Analysis for Research and Applications
 461 reanalysis data; <http://gmao.gsfc.nasa.gov/research/merra/>), Rienecker et al. (2011) showed strong correlation
 462 between 6-10 day periodic oscillations associated with Rossby waves in the upper tropospheric winds and surface
 463 weather parameters like atmospheric pressure, winds, temperature, relative humidity and rainfall during a severe
 464 weather event observed at the Indian extratropical station, Nainital (29.45° N, 79.5° E), in November–December
 465 2011. They also note that when the upper troposphere shows divergence, the lower troposphere shows convergence
 466 and as a result more moisture gets accumulated there leading to enhancement of relative humidity and hence
 467 precipitation. It was asserted that Rossby waves in the upper troposphere can lead to surface weather related events
 468 through the action of convergence or divergence in the atmospheric air. It is to be noted that a passing Rossby wave
 469 can cause fluctuations in divergence and convergence in the atmosphere at periodicities (typically 6-10 days, 12-20
 470 days) corresponding to the Rossby waves at a particular site.
 471

472 It was reported that Rossby waves account for more than 30% of monthly mean precipitation and more than
 473 60% of surface temperature over many extra tropical regions and influence short-term extreme weather phenomena
 474 (Schubert et al., 2011). Planetary waves affecting weather events severely for long duration of the order of months
 475 have been reported by many researchers (Petoukhov et al., 2013; Screen and Simmonds, 2014; Coumou et al.,
 476 2014). Screen and Simmonds (2014) found that in the mid latitudes, there was a strong association between
 477 enhanced Rossby wave activity, surface temperature and extreme precipitation events in 1979–2012. Since slowly
 478 propagating Rossby waves can influence weather at a particular site for long periods lasting more than few weeks, it
 479 is can be seen the imprint of climatic variations of Rossby waves in weather events from monthly mean atmospheric
 480 parameters.

481
482 To understand the present observation of different precipitation characteristics over different stations, it is
483 compared between monthly variation of PV in the upper troposphere and precipitation. Potential vorticity at 350K
484 surface is identified for investigating Rossby waves as their breakage (can be identified through reversal of
485 gradient in PV) at this level can lead to exchange of air at the boundary between the tropics and extra tropics
486 (Homeyer and Bowman, 2013). Similarly PV at 200 hPa pressure surface is more appropriate for identifying Rossby
487 wave breaking in the subtropical regions (Garfinkel and Waugh, 2014).

488
489 Since the Srinagar city is located on comparatively plain land than the other all six stations of the Kashmir
490 valley, precipitation associated with western disturbances here is under the direct influence of planetary-scale
491 Rossby waves. Accordingly, correlation between PV at the 350 K (located near the core of the subtropical jet,
492 Homeyer and Bowman, 2013) and 200 hPa pressure surfaces and precipitation is found significantly larger over
493 Srinagar than other stations. Orographic effects at other stations can have significant influence on planetary Rossby
494 waves. Therefore, PV (ERA-Interim data, Dee et al., 2011) in the upper troposphere varies in accordance with
495 precipitation, which is clearly depicted in Fig. 7, during the entire years of 1984, 1987, 1988, 1990, 1993, 1994,
496 1995, 1996, 1999, 2006 and 2009. In general, it is observed that sometimes PV at 350K surface and at other times at
497 200 hPa pressure surface follows precipitation. This would be due to the influence of Rossby waves generated due to
498 baroclinic or and barotropic instabilities. Particularly, the correlation between PV (sometimes either one or both) and
499 precipitation is significantly positive during the Indian summer monsoon months of June-September for all the years
500 from 1980 to 2009 except 1983, 1985, 1989, 2000-2005 and 2009. At present it is not known why this relation
501 became weak during 1999-2010.

502
503 For Kokarnag (Fig. 8), the topography of which is similar to Srinagar but it is located in the vicinity of high
504 mountains, the relation between PV and precipitation particularly during the Indian summer-monsoon is almost
505 similar to that of Srinagar during 1983, 1985, 1989, 1991, 1998, 1999, 2000-2005.. The deterioration of the link
506 between PV and rainfall over Kokarnag and Srinagar during 1999-2010 is intriguing and it may be associated with
507 climate change. In the northern Kashmir region of Kupwara (Fig. 9), msl higher by ~1 km than Srinagar, the relation
508 between PV and precipitation is good in the years 1982-1983, 1985-1988, 1990-1994, 1995-1996, 1999, and 2006.
509 Similar to Srinagar and Kokarnag, Kupwara also shows a poor link during 1999-2010. Particularly during the
510 summer monsoon period, the PV-precipitation relation is good in all the years except 1989, 1998, 2000-2005, and
511 2009. One interesting observation is that in 1983, 1985 and 1991 the correlation between PV and precipitation for
512 Kupwara is better than Srinagar and Kokarnag. Since Kupwara is located near elevated Greater Himalayan mountain
513 range, Rossby waves associated with topography would have contributed to the good correlation between PV and
514 precipitation here, which is not the case for Srinagar and Kokarnag. In the case of Pahalgam, (Fig.10), located near
515 the Greater Himalayas, generally the link between PV and precipitation is good in almost all the years 1980-2016
516 but with a difference that sometimes both the PVs and on other times only either of them follow precipitation.
517 Particularly during summer monsoon months, similar to Kupwara, these years 1989, 2000-2003, 2005 and 2009

show poor correlation. In general, precipitation near the Greater Himalayas is significantly influenced by Rossby waves associated with topography.

For the hilly station of Qazigund (Fig. 11), located in the south Kashmir region (above ~3 km msl) near the foot hills of Pir Panjal mountain range, the relation between PV and precipitation is better than that of the northern station Kupwara. For example, in 1988, the relation is much better over Qazigund than Kupwara. However the opposite is true in 1987. Interestingly, in 1985, both Kupwara and Qazigund show similar variation in PV and precipitation. This may be due to the effect of the nature of limited equatorward propagation of Rossby waves from mid-latitudes. In 1995, 1997 and 1998, PV and precipitation follow similar time variation at both Kupwara and Qazigund except for January-March during which precipitation over Qazigund but not Kupwara follows PV. Interestingly, in the whole year of 1999, precipitation at both the stations, follows exceedingly well with PV; however in 1998, only Qazigund but not Kupwara shows good relation. In 2009, precipitation does not follow PV for both the stations. Interestingly in all the months of 2006, PV follows well with precipitation for both Kupwara and Qazigund. However in September, Kupwara but not Qazigund shows good relation. In 2004, only PV at 350K surface follows well with precipitation for both the stations. For the summer monsoon period of June-September, these years, namely, 1983, 1985, 1989, 1990, 2000-2003, 2005, 2007-2009, do not show good correlation, which is almost similar to Srinagar and Kokarnag.

In the case of Gulmarg (Fig. 12), PV and precipitation follow each other well in the years of 1988, 1993, 1994 and 1995. In 1996, during the Indian summer monsoon period of June-September, only PV at 350K surface follows precipitation. Overall, during the summer monsoon period, the relationship between PV and precipitation is appreciable for all the years except for 1983, 1989, 1990, 1999 and 2000-2009, which is almost similar to Kupwara and Pahalgam. It may be noted that these stations are located near relatively elevated mountains and hence topographically induced Rossby waves could have contributed to this good relation. The observations suggest that high altitude mountains affect the precipitation characteristics through topography generated Rossby waves. The interesting finding here is that irrespective of the different heights of mountains, all the stations show that during 1999-2010 the correlation between upper tropospheric PV and surface precipitation found to be poor, indicating that some unknown new atmospheric dynamical concepts would have played significant role in disturbing the precipitation characteristics significantly over the western Himalayan region. This issue needs to be addressed in the near future by invoking suitable theoretical models so that predictability of extreme weather events can be improved in the mountainous Himalaya.

During 2011-2016 (Fig. 13), it may be observed that for Gulmarg the link between PV and precipitation holds good in general for all these years except around July 2012, July-December 2013 and 2015. It is interesting to note here that during the historical flood event of September 2014, the PV and precipitation follow each other but in the preceding and following years of 2013 and 2015 their linkage is poor as noted earlier. Similarly, all the other stations (Srinagar, Pahalgam, Kokarnag, Kupwara, and Qazigund) also show that the link between PV and

precipitation is good around September 2014. This would indicate clearly that the extreme weather event occurred during September 2014 is due to intense large-scale Rossby wave activity rather than any localized adverse atmospheric thermodynamical conditions such as local convection. In Srinagar, most of the times PV and precipitation follow each other very well as observed during January 2011-June 2012, January-July of 2013 & 2014, whole 2015 and 2016. In Qazigund, this relation is good only during January-July and September-October 2014, during the entire 2015 and 2016 (similar to Srinagar). For Kupwara, PV follows precipitation well during whole of 2011, January-July 2012, January-May 2013, January-November 2014, whole of 2015 and 2016. In the case of Kokarnag, good relation is observed during March-August 2012, January-June 2013 and 2014, around September 2014. In contrast, the relationship is very poor in the entire years of 2015 and 2016. Pahalgam interestingly shows good correlation between PV and precipitation during the whole years of 2011 and 2012. In 2013, 2014, 2015 and 2016, it is good only during January-June in addition to exceptionally good in September 2014.

Finally, it may be observed that the ERA-interim reanalysis data of meridional wind velocity (12UT) at ~3 km altitude above the mean sea level show alternating positive (southerly) and negative values, resembling the atmospheric Rossby waves in the subtropical region during 1-6 September 2014 (Fig. 14). The meridional winds associated with Rossby waves could be easily noted to have their extensions in both the Arabian Sea and Bay of Bengal, indicating that water vapour from both the regions was transported towards the Jammu and Kashmir, India region as the converging point of Rossby waves was located near this region. It may be easily noticed that the waves got strengthened on 4th and weakened on 5th and ultimately dissipated on 6th September. This dissipation of Rossby waves led to dumping of the transported water vapour over this region thus caused the historical-record heavy-flooding during this period. This is one clear example of how synoptic scale Rossby waves can reorganize water vapour over large scale and lead to extreme rainfall event. It is well known that subtropical westerly jet is one of many important sources of Rossby waves in the mid to tropical latitudes. If the subtropical jet drifts climatically northward then the surface weather events associated with them also will drift similarly, leading to unusual weather changes climatically.

Published reports (Barnes and Polvani, 2013; Lu et al., 2014) indicate that long-term variations in Rossby wave breaking activities and stratospheric dynamics have close association with global climate change. (Meridional shift of the center of subtropical jets, arising due to enhanced polar vortex and upper-tropospheric baroclinicity are possible due to the consequences of global warming, has been successfully linked to climatic changes in Rossby wave breaking events caused by baroclinic instabilities (Wittman et al., 2007; Kunz et al., 2009; Rivière, 2011; Wilcox et al., 2012). The long-term increase in the tropospheric warming arising due to baroclinic forcing of Rossby waves is more prominent in the mid-latitudes than in the tropical regions (Allen et al., 2012; Tandon et al., 2013). This mid-latitude warming plays a critical role in driving poleward shift of the subtropical jet responding to climate change (Ceppi et al., 2014). It is to be remembered that the combined effect of tropospheric baroclinic forcing (warming) and stratospheric polar vortex can gradually move the subtropical jet from about 27° to 54° (Garfinkel and Waugh, 2014). Using Global circulation models (GCM), linear wave theory predicts that in response to

increased greenhouse gas (GHG) forcing, mid-latitude eddy-driven jets, arising due to strong coupling between synoptic scale eddy activity and jet streams in both the hemispheres, will be shifted poleward (Fourth report of Intergovernmental Panel on Climate Change (IV-IPCC), Meehl et al., 2007). However, mid-latitude Rossby waves and the associated wave dissipation in the subtropical region are predicted to move climatologically towards equator due to the spherical geometry of the Earth (Hoskins et al., 1977; Edmon et al., 1980). This propagation of location of wave breaking towards the equator will have long-term (climatic) impact on relation between variations in upper tropospheric PV associated with Rossby waves and surface precipitation in the subtropical latitude regions. This may be one of the reasons that during 1999-2010, the relation between PV and precipitation became poor as observed in the present study.

Regarding surface temperature, except for its linear long-term trend, there is no clear evidence of strong link between variations in the upper tropospheric potential vorticities and surface temperature for all the six stations mentioned. It seems that long-term (climatic) variations in the upper tropospheric vorticities have significantly less influence on surface temperature variations.

5. Conclusions

In this study, trends and variations in surface temperature and precipitation over the Jammu and Kashmir, India region of the western Himalayas are carried out for a period of 37 years during 1980-2016. Analyses of the observations reveal that the annual temperature increased by 0.8°C during this period. Higher increase in annual temperature accompanied by insignificant decrease in annual precipitation is noted for stations located at higher altitudes. Long-term variation of winter temperature and precipitation has good correlation with winter NAO index. To provide more conclusive evidence on our observations, we employed WRF model simulations which show good correlation of 0.85 with the observed data. It is found that in the recent decades, precipitation associated with both the monsoons and western disturbances has been decreasing significantly. While the monsoon deficiency is associated with decreasing difference in surface temperature between the Indian landmass and nearby Indian Ocean, the deficiency associated with western disturbances during winter is due to the climatic northward displacement of the subtropical westerly jet. This subtropical jet wind helps to enhance the moisture transport associated with disturbances from the tropical Atlantic Ocean, Mediterranean and Caspian Seas to the Himalayan region. Regarding historical extreme weather event associated with September 2014 floods in Jammu and Kashmir, it is found that breaking of intense Rossby wave activity over Kashmir played an important role as the wave could transport lots of water vapor from both the Bay of Bengal and Arabian Sea and dump them here through its breaking during the first week of September, 2014, leading to the extreme rainfall event measuring more than 620 mm in southern parts of the Kashmir.

629

630

631 **Acknowledgements:**

632

633 Thanks are due to the India Meteorological Department, Pune, India, ERA-Interim reanalyses and WRF model
634 simulation teams for the data of meteorological parameters employed in the present work. Prof. Shakil Romshoo and
635 Dr. Sumaira Zaz gratefully acknowledge the support of the Department of Science and Technology (DST),
636 Government of India under the research project titled “Himalayan Cryosphere: Science and Society”. Dr. T. K.
637 Ramkumar and Dr. V. Yesubabu acknowledge the support of Dept. of Space, Govt. of India. The authors express
638 gratitude to the two anonymous reviewers and Editor for their valuable comments and suggestions on the earlier
639 version of the manuscript that has greatly improved its content and structure.

640

641

642 **References:**

643

644 Allen, R. J., Sherwood, S. C., Norris, J. R. and Zender, C. S.: Recent Northern Hemisphere tropical expansion
645 primarily driven by black carbon and tropospheric ozone, *Nature*, 485, 350–354, doi:10.1038/nature11097, 2012.

646

647 Archer, D. R. and Fowler, H. J.: Spatial and temporal variations in precipitation in the Upper Indus Basin, global
648 teleconnections and hydrological implications, *J. of Hyd. and Earth Sys. Sci.*, 8, 47–61, 2004.

649

650 Barnes, E. A., S. Solomon, and L. M. Polwani, Robust wind and precipitation responses to the Mount Pinatubo
651 eruption, as simulated in the CMIP5 Models, *J. of Clim.*, DOI: 10.1175/JCLI-D-15-0658.1, 2016.

652

653 Barnes, E. A. and Polvani, L.: Response of the mid latitude jets, and of their variability, to increased greenhouse
654 gases in the CMIP5 models, *J. Climate*, 26, 7117–7135, doi:10.1175/JCLI-D-12-00536.1, 2013.

655

656 Bartels, J., Peters, D. and Schmitz, G.: Climatological Ertel’s potential vorticity flux and mean meridional
657 circulation in the extratropical troposphere–lower stratosphere, *Ann. Geophys.*, 16, 250–265, 1998.

658

659 Beniston, M.: Impact of climatic change on water and associated economic activities in the Swiss Alps, *J. of*
660 *Hydrology*, 1-6, 2010.

661

662 Bhutiyani, M. R., Kale, V. S. and Pawar, N. J.: Long-term trends in maximum, minimum and mean annual air
 663 temperatures across the north western Himalaya during the 20th century, *Climatic Change*, 85, 159–177, 2007.
 664
 665 Bhutiyani, M. R., Kale, V. S. and Pawar, N. J.: Climate change and the precipitation variations in the north western
 666 Himalaya: 1866–2006, *Int. J. of Climatology*, 30(4), 535–548, 2009.
 667
 668 Bhutiyani, M. R., Kale, V. S. and Pawar, N. J.: Climate change and the precipitation variations in the north western
 669 Himalaya: 1866–2006, *Int. J. of Climatology*, 30, 535–548, 2010.
 670
 671 Bolch, T., Kulkarni, A., Kaabet. Al.: The state and fate of Himalayan glaciers, *Science*, 336, 310–314, 2012.
 672
 673 Bookhagen, B.: Appearance of extreme monsoonal rainfall events and their impact on erosion in the Himalaya.
 674 *Geomatics, Natural Hazards and Risk*, 1(1), 37–50. Doi: 10.1080/19475701003625737, 2010.
 675
 676 Borgaonkar, H. P. and Pant, G. B.: Long-term climate variability over monsoon Asia as revealed by some proxy
 677 sources, *Mausam*, 52, 9–22, 2001.
 678
 679 Ceppi, P., Zelinka, M. D. and Hartmann, D. L.: The response of the southern hemispheric eddy-driven jet to future
 680 changes in shortwave radiation in CMIP5, *Geophys. Res. Lett.*, 41, 3244–3250, doi:10.1002/2014GL060043, 2014.
 681
 682 Chang, E. K. M. and Yu, D. B.: Characteristics of Wave Packets in the Upper Troposphere. Part I: Northern
 683 Hemisphere Winter, *J. Atmos. Sci.*, 56, 1708–1728, 1999.
 684
 685 Chen, J. and Gupta, A. K.: *Parametric Statistical Change Point Analysis*, Birkhauser, Boston, MA, 240, 2012.
 686
 687 Collins, D.: Climatic warming, glacier recession and runoff from Alpine basins after the Little Ice Age maximum.
 688 *Ann. of Glac.*, 48(1), 119–124, 2008.
 689
 690 Coumou, D., Petoukhov, V., Rahmstorf, S., Petri, S. and Schellnhuber, H. J.: Quasi-resonant circulation regimes and
 691 hemispheric synchronization of extreme weather in boreal summer, *P. Natl. Acad. Sci. USA*, 111, 12331–12336,
 692 doi:10.1073/pnas.1412797111, 2014.
 693
 694 Dar, R. A., Romshoo, S. A., Chandra, R. and Ahmad, I.: Tectono-geomorphic study of the 4 Karewa Basin of
 695 Kashmir valley, *J. of Asian Earth Sciences*, 92, 143–156, 2014.
 696

697 Das, M. R., Mukhopadhyay, R. L., Dandekar, M. M. and Kshirsagar, S. R.: Pre-monsoon western disturbances in
 698 relation to monsoon rainfall, its advancement over NW India and their trends, *Current Science*, 82(11), 1320-1321,
 699 2002.
 700
 701 Dee, D. P. and Coauthors: The ERA-Interim reanalysis: Configuration and performance of the data assimilation
 702 system, *Quart. J. of Royal Met. Soc.*, 137, 553–597, doi:10.1002/qj.828, 2011.
 703
 704 Dimri, A. P., Niyogi, D., Barros, A. P., Ridley, J., Mohanty, U. C., Yasunari, T. and Sikka, D. R.: Western
 705 Disturbances: A Review, *Rev. of Geophys*, 2014RG000460, Doi: 10.1002/2014RG000460, 2015.
 706
 707 Dimri, A. P. and Dash, S. K.: "Winter time climatic trends in the western Himalayas," *Climatic Change*, Springer,
 708 111(3), pages 775-800, 2012.
 709
 710 Edmon, H. J., Hoskins, B. J. and McIntyre, M. E.: Eliassen-Palm cross sections for the troposphere, *J. Atmos. Sci.*,
 711 37, 2600–2615, 1980.
 712
 713 Epstein, P. R. et al.: Extreme Weather Events: The Health and Economic Consequences of the 1997/98 ElNiño and
 714 La Niña. Center for Health and the Global Environment, Harvard Medical School, Boston. Database available on
 715 website [<http://www.chge2.med.harvard.edu/enso/disease.html>], 1998.
 716
 717 Ertel, H.: Einneuer hydrodynamischer Wirbelsatz. *Meteor. Z.*, 59, 277–281, 1942.
 718
 719 Folland, C. K., Rayner, N. A., Brown, S. J., Smith, T. M., S. P. Shen, Parker, D. E., Macadam, I., Jones, P. D.,
 720 Nicholls, R. N. N. and Sexton, D. M. H.: "Global temperature change and its uncertainties since 1861," *Geophys.*
 721 *Res. Lett.*, 28, 2621–2624, DOI:10.1029/2001GL012877, 2001.
 722
 723 Gao, P., Mu, X. M., Wang, F., and Li, R.: Changes in stream flow and sediment discharge and the response to
 724 human activities in the middle reaches of the Yellow River, *Hydrology and Earth System Sciences*, 15, 1–10, 2011.
 725
 726 Garfinkel, C. I. and Waugh, D. W.: Tropospheric Rossby wave breaking and variability of the latitude of the eddy-
 727 driven jet, *J. of Cli.*, 27, 7069-7085, DOI: 10.1175/JCLI-D-14-00081.1, 2014.
 728
 729 Ghasemi, A. R.: Changes and trends in maximum, minimum and mean temperature series in Iran, *Atmos. Sci. Lett.*,
 730 16, 201-230, 2015.
 731
 732 Ghasemi, A. R.: Changes and trends in maximum, minimum and mean temperature series in Iran, DOI:
 733 10.1002/asl2.569, *Atmos. Sci. Lett.*, 16, 366–372, 2015.

734
735 Groisman, Pavel, Ya, Karl, T. R., Richard, Knight, W., Georgiy, L. and Stenchikov: Changes of snow cover,
736 temperature, and radiative heat balance over the Northern Hemisphere, *J. of Climate*, 7:1633–1656, 1994.
737
738 Hansen, A. R., Nastrom, G. D. and Eaton, F. D.: Seasonal variation of gravity wave activity at 5–20 km observed
739 with VHF radar at White Sands Missile Range, New Mexico. *J. of Geophys. Res.*, 106: doi:
740 10.1029/2001JD900137. issn: 0148-0227, 2001.
741
742 Hewitt, K.: ‘The Karakoram anomaly? Glacier expansion and the ‘elevation effect,’ Karakoram, Himalaya.’
743 *Mountain Research and Development*, 25(4), 332-340, 2005.
744
745 Hijikata, Y., Lin, E., Pereira, J. J., Corlett, R. T., Cui, X., Insarov, G. E., Lasco, R. D., Lindgren, E. and Surjan, A.:
746 Asia. In: Barros VR et al. (eds), *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional*
747 *Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel of*
748 *Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2014.
749
750 Homeyer, C. R. and Bowman, K. P.: Rossby Wave Breaking and Transport between the Tropics and Extratropics
751 above the Subtropical Jet, *J. of Atmos. Sci.*, 70, 607-626, DOI: 10.1175/JAS-D-12-0198.1, 2013.
752
753 Hoskins, B. J., Simmons, A. J. and Andrews, D. G.: Energy dispersion in a barotropic atmosphere, *Quart. J. Roy.*
754 *Meteor. Soc.*, 103, 553–567, 1977.
755
756 Hoskins, B. J., McIntyre, M. E. and Robertson, A. W.: On the use and significance of isentropic potential vorticity
757 maps, *Quart. J. Roy. Meteor. Soc.*, 111, 877–946, 1985.
758
759 Hunt, K. M. R., Turner, A. G. and Shaffrey, L. C.: Extreme daily rainfall in Pakistan and north India: scale-
760 interactions, mechanisms, and precursors, *Mon. Wea. Rev.*, 146 (4), 1005-1022. ISSN 0027-0644
761 DOI <https://doi.org/10.1175/MWR-D-17-0258.1>, 2018a.
762
763 Hunt, K. M. R., Turner, A. G. and Shaffrey, L. C.: The evolution, seasonality and impacts of western disturbances,
764 *Quart. J. Roy. Meteor. Soc.*, 144 (710), 278-29. ISSN 1477-870X, <https://doi.org/10.1002/qj.3200>, 2018b.
765
766 Hurrell, J. W. and van Loon, H.: Decadal variations in climate associated with the North Atlantic Oscillation,
767 *Climatic Change*, 36, 301–326, *Res. Lett.*, 23, 665–668, 1997.
768
769 Immerzeel, W., Van Beek, L. P. H. and Bierkens, M. F. P.: Climate change will affect the Asian water towers.
770 *Science*, 328, 1382-1385, 2010.

771
772 IPCC Climate Change 2013: . T.F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels,
773 Y. Xia, V. Bex, P.M. Midgley (Eds.), The Physical Science Basis. Contribution of Working Group I to the Fifth
774 Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge,
775 United Kingdom and New York, NY, USA, 1535, 2013.
776
777 IPCC Climate change (2001): Impacts, adaption and vulnerability. Contribution of Working Group II to the Third
778 Assessment Report of the Intergovernmental Panel of Climate Change, Intergovernmental Panel on Climate Change,
779 Cambridge, U.K, 2001.
780
781 Iqbal, M. J. and Kashif, I.: Influence of Icelandic Low pressure on winter precipitation variability over northern part
782 of Indo-Pak Region Arabian, J. of Geosc., 6, 543–548, DOI 10.1007/s12517-011-0355-y, 2013.
783
784 Jones, P. D., Osborn, T. J. and Briffa, K. R.: The evolution of climate over the last millennium, Science, 292, 662–
785 667, 2001.
786
787 Niranjana Kumar, K., Phanikumar, D. V., Ouada, T. M. B. J., Rajeevan, M., Naja, M. and Shukla, K. K.: Modulation
788 of surface meteorological parameters by extratropical planetary-scale Rossby waves, Ann. Geophys., 34, 123–132,
789 doi:10.5194/angeo-34-123-2016, 2016.
790
791 Kaul, V. and Qadri, B. A.: Seasons of Kashmir. Geographic Revision India. 41(2), pp123-130, 1979.
792
793 Khattak, M. S., Babel M. S., and Sharif, M.: Hydro-meteorological trends in the upper Indus River basin in
794 Pakistan,” Inter-Research, Climate Research, 46, 103–119, 2011 doi: 10.3354/cr00957, 2011.
795
796 Knutti, R., Rogelj, J., Sedláček, J., Fischer, E. M.: A scientific critique of the two-degree climate change target,
797 Nature Geoscience, 9(1), 13–18, 2016.
798
799 Kohler, T. and Maselli, D.: Mountains and climate change from understanding to action. Berne: Swiss Agency for
800 Development and Cooperation, 2009.
801
802 Kulkarni, A. V., Mathur, P., Rathore, B. P., Suja Alex., Thakur, N. and Manoj et al.: Effect of global warming on
803 snow ablation pattern in the Himalaya. Current Science, 83, 120– 123, 2002.
804
805 Kumar, N. Yadav, B. P., Gahlot, S and Singh, M.: Winter frequency of western disturbances and precipitation
806 indices over Himachal Pradesh, India: 1977-2007. Atmósfera 28(1), 63-70. Doi:
807 [http://dx.doi.org/10.1016/S0187\(6236\(15\)72160\)0](http://dx.doi.org/10.1016/S0187(6236(15)72160)0), 2015.

808

809 Kumar, V and Jain, S. K.: Trends in seasonal and annual rainfall and rainy days in Kashmir valley in the last
810 century, *Quaternary International*. doi:10.1016/j.quaint.2009.08.006, 2009.

811

812 Kunz, T., Fraedrich, K. and Lunkeit, F.: Response of idealized baroclinic wave life cycles to stratospheric flow
813 conditions, *J. Atmos. Sci.*, 66, 2288–2302, doi:10.1175/2009JAS2827.1, 2009.

814

815 Lau, W. K. M. and Kim, K-M.: The 2010 pakistan flood and Russian heat wave: Teleconnection of hydro
816 meteorological Extremes, *J. of Hydro Meteorological*, 13(1), 392-403, DOI:10.1175/jhm-D-11-016.1, 2012.

817

818 Langodan S., Yesubabu V., Hoteit I.: The impact of atmospheric data assimilation on wave simulations in the Red
819 Sea, *Ocean Engineering*, 116, 200-215, doi:10.1016/j.oceaneng.2016.02.020, 2016.

820

821 Liu, X, Cheng, Z., Yan, L., Yin, Z. Y.: Elevation dependency of recent and future minimum surface air temperature
822 trends in the Tibetan Plateau and Its surroundings. *Global Planet Change* 68: 164-174, 2009.

823

824 Liu, X. B. and Chen.: Climatic warming in the Tibetan Plateau during recent decades, *J. of Clim.*, 20, 1729–1742,
825 2000.

826

827 Lo, J. C. F., Yang, Z. L., Pielke, R. A. Sr.: Assessment of three dynamical climate downscaling methods using the
828 weather research and forecasting (WRF) model, *J. Geophys. Res.* 113. D09112, doi: 10.1029/2007JD009216, 2008.

829

830 Lu, J., Sun, L., Wu, Y. and Chen, G.: The role of subtropical irreversible PV mixing in the zonal mean circulation
831 response to global warming-like thermal forcing, *J. Climate*, 27, 2297–2316, doi:10.1175/JCLI-D-13-00372.1, 2014.

832

833 Madala, S., Satyanarayana, A. N. V., Narayana Rao, T.: Performance evaluation of PBL and cumulus
834 parameterization schemes of WRF ARW model in simulating severe thunderstorm events over Gadanki MST radar
835 facility — Case study, *Atmos. Res.*, 139, 1-17, doi:10.1016/j.atmosres.2013.12.017, 2014.

836

837 Madhura, R. K., Krishnan, R., Revadekar, J. V., Mujumdar, M. and Goswami, B. N.: Changes in western
838 disturbances over the Western Himalayas in a warming environment. *Climate Dynamics*, 44, 3-4, 1157-1168, Doi:
839 10.1007/s00382-014-2166-9, 2015.

840

841 Mann, M. E., Bradley, R. S. and Hughes, M. K.: Northern Hemisphere Temperature During Past Millennium:
842 Inferences, uncertainties and Limitations, *Geophys. Res. Lett.*, 26(6), 759-762, 1999.

843

844 Martius, O., Sodemann, H., Joos, H., Pfahl, S., Winschall, A., Croci-Maspoli, M., Graf, M., Madonna, E., Mueller,
 845 B., Schemm, S., Sedláček, J., Sprenger M., Wernli H.: The role of upper-level dynamics and surface processes for
 846 the Pakistan flood of July 2010, *Q. J. R. Meteorol. Soc.*, 139, 1780–1797, doi:10.1002/qj.2082, 2012.
 847
 848 McIntyre, M. E. and Palmer, T. N.: Breaking planetary waves in the stratosphere. *Nature*, 305, 593–600, 1983.
 849
 850 Meehl, G. A., Covey, C., Delworth, T., Latif, M., McAvaney, B., Mitchell, J. F. B., Stouffer, R. J. and Taylor, K. E.:
 851 The WRCM CMIP3 multi-model dataset: A new era in climate change research, *Bull. Amer. Meteor. Soc.*, 88, 1383–
 852 1394, 2007.
 853
 854 Mooley, D. A. and Parthasarthy, B.: Fluctuations of all India summer monsoon rainfall during 1871–1978. *Climate*
 855 *Change*, 6, 287–301, 1984.
 856
 857 Murtaza, K. O. and Romshoo, S. A.: Recent Glacier Changes in the Kashmir Alpine Himalayas, India, *Geocarto*
 858 *International*, 32 (2), 188-205, 2016.
 859
 860 Nibanupudi, H. K., Gupta, A. K., and Rawat, P. K.: Mountain Hazards and Disaster Risk, (2015): Mitigating
 861 Climatic and Human Induced Disaster Risks Through Ecosystem Resilience: Harmonizing Built and Natural
 862 Environments in the KHK Region, edited by: Nibanupudi, H. K. and Shaw, R., 139–158, doi:10.1007/978-4-431-
 863 55242-0, Springer, Tokyo, Japan, 2015.
 864
 865 Peters, G. P., Andrew, R. M., Boden, T., Canadell, J. G., Ciais, P., Le Quéré C., et al.: The challenge to keep global
 866 warming below 2°C, *Nature, Climate Change*, 3(1), 4–6, 2013.
 867
 868 Petoukhov, V., Rahmstorf, S., Petri, S., and Schellnhuber, H. J.: Quasi resonant amplification of planetary waves
 869 and recent Northern Hemisphere weather extremes, *P. Natl. Acad. Sci., USA*, 110, 5336–5341, 2013.
 870
 871 Pettitt, A. N.: A non-parametric approach to the change point problem, *App. Stats.*, 28, 126–135, 1979.
 872
 873 Postel, G. A., and Hitchman, M. H.: Climatology of Rossbywave breaking along the subtropical tropopause, *J.*
 874 *Atmos. Sci.*, 56, 359–373, 1999.
 875
 876 Priyanka Ghosh, Ramkumar, T. K., Yesubabu, V. and Naidu, C. V.: Convection-generated high-frequency gravity
 877 waves as observed by MST radar and simulated by WRF model over the Indian tropical station of Gadanki, *Q. J. R.*
 878 *Meteorol. Soc.*, DOI:10.1002/qj.2887, 2016.
 879

880 Radziejewski, M., Bardossy, A., Kundzewicz, Z. W.: Detection of change in river flow using phase randomization,
 881 Hydrological Sciences Journal, 45, 547–558, 2000.
 882
 883 Rashid, .I, Romshoo, A. S., Chaturvedi, R. K., Ravindranath, N. H., Raman Sukumar, Mathangi Jayaraman,
 884 Thatiparthi Vijaya Lakshmi and Jagmohan Sharma: Projected Climate Change Impacts on Vegetation Distribution
 885 over Kashmir Himalaya, Climatic Change, DOI: 10.1007/s10584-015-1456-5, 2015.
 886
 887 Rasmussen, K. L. R., and Houze, R.: A Flash-Flooding Storm At The Steep Edge Of High Terrain: Disaster in the
 888 Himalayas, Bull. Ame. Meteorol. Soc., 93, 1713-1724, doi:10.1175/BAMS-D-11-00236.1, 2012.
 889
 890 Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M. G., Schubert, S. D.,
 891 Takacs, L., Kim, G.-K., Bloom, S., Chen, J., Collins, D., Conaty, A., daSilva, A., Gu, W., Joiner, J., Koster, R. D.,
 892 Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick,
 893 A. G., Sienkiewicz, M., and Woollen, J.: MERRA: NASA's Modern-Era Retrospective Analysis for Research and
 894 Applications, J. Climate, 24, 3624–3648, doi:10.1175/JCLI-D-11-00015.1, 2011.
 895
 896 Rivière, G.: A dynamical interpretation of the poleward shift of the jet streams in global warming scenarios, J.
 897 Atmos. Sci., 68, 1253–1272, doi:10.1175/2011JAS3641.1, 2011.
 898
 899 Roe, G. H., Montgomery, D. R. and Hallet, B.: Orographic climate feedbacks and the relief of mountain ranges, J. of
 900 Geophys. Res., 108, doi: 10.1029/2001JB001521, 2003.
 901
 902 Romatschke, U., and Houze, R.: Characteristics of Precipitating Convective Systems in the Premonsoon Season of
 903 South Asia, J. Hydrometeorology, 12, 157-180, doi:10.1175/2010JHM1311.1, 2011.
 904
 905 Romshoo, S. A. and Rashid, I.: Assessing the impacts of changing land cover and climate on Hokersar wet land in
 906 Indian Himalayas, Arabian J. of Geoscience, DOI: 10.1007/s12517-012-0761-9, 7 (1): 143-160, 2014.
 907
 908 Romshoo, S. A., Altaf, S., Rashid, I, and Dar, R. A.: Climatic, geomorphic and anthropogenic drivers of the 2014
 909 extreme flooding in the Jhelum basin of Kashmir, India. Geomatics, Natural Hazards and Risk, 9 (1), 224-248, 2017.
 910
 911 Romshoo, S. A., Dar, R. A., Rashid, I., Marazi, A., Ali, N. and Zaz, S. N.: Implications of Shrinking Cryosphere
 912 under Changing Climate on the Stream flows of the Upper Indus Basin, Arctic, Antarctic and Alpine Research,
 913 47(4), 627-644, ISSN: 1938-4246, 2015.
 914
 915 Schubert, S., Wang, H., and Suarez, M.: Warm season subseasonal variability and climate extremes in the Northern
 916 Hemisphere: The Role of Stationary Rossby Waves, J. Clim., 24, 4773–4792, 2011.

917
918 Screen, J. A. and Simmonds, I.: Amplified mid-latitude planetary waves favour particular regional weather
919 extremes, *Nature Climate Change*, 4, 704–709, 2014.
920
921 Sharif, M., Archer, R. D., Fowler, J. H. and Forsythe, N.: Trends in timing and magnitude of flow in the Upper
922 Indus Basin. *Hydrology and Earth System, Science*. 9, 9931–9966, 2012.
923
924 Sheikh, M. M., Manzoor, N., Adnan, M., Ashraf, J. and Khan, A. M.: Climate Profile and pastclimate changes in
925 Pakistan GCISC-RR-01Global Change Impact studies Center Islamabad, Pakistan, ISBN: 978-969-9395-04, 2009.
926
927 Shekhar, M. S., Chand, H., Kumar, S., Ganju, Ashwagosh: Climate change studies in western Himalaya, *Annals of*
928 *Glaciology* 51(54):105-112, 2010.
929
930 Shrestha, A. B., Wake, C. P., Dibb, J. E. and Mayewski, P. A: Precipitation fluctuations in the Nepal Himalaya and
931 its vicinity and relationship with some large scale climatological parameters, *Int. J. of Climatology*, 20, 317–327,
932 1999.
933
934 Shrestha, M. L.: Interannual variation of summer monsoon rainfall over Nepal and its relation to Southern
935 Oscillation Index, *Meteor. and Atmos. Physics*, 75, 21–28, doi: 10.1007/s007030070012, 2000.
936
937 Singh, P., Kumar, V., Thomas, M., Arora et al.: Changes in rainfall and relative humidity in river basins in
938 northwest and central India *Hydrological Processes*, 22, 16, 2982-2992, 2008.
939
940 Sinha Ray, K. C. and Srivastava, A. K.: Is there any change in extreme events like drought and heavy rainfall? *Curr.*
941 *Sci. India*, 79, 155–158, 2000.
942
943 Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M. and Miller, H. L. (eds):
944 *Climate change 2007: the physical science basis*, 2007.
945
946 Srinivas, C. V., Hariprasad, D., Bhaskar Rao, D. V., Anjaneyulu, Y., Baskaran, R., Venkatraman, B.: Simulation of
947 the Indian summer monsoon regional climate using advanced research WRF model, *Int. J. Climatol.*, 33, 1195-1210.
948 doi:10.1002/joc.3505, 2013.
949
950 Srinivasa, C. V., Yesubabu, V., Hari Prasad, D., Hari Prasad, K. B. R. R., Greeshmaa, M. M., Baskarana, R.,
951 Venkatramana, B.: Simulation of an extreme heavy rainfall event over Chennai, India using WRF: Sensitivity to grid
952 resolution and boundary layer physics, 210: 66–82, 2018.
953

954 Swanson, D. K., Wooten and Orr, T.: Buckets of Ash Track Tephra Flux From Halema'uma'u Crater, Hawai'i, Eos
 955 Trans.. AGU, 90(46), 427, 2009.
 956
 957 Syed, F. S., Giorgi, F., Pal, J. S., King, M. P.: Effect of remote forcings on the winter precipitation of central
 958 southwest Asia part 1: observations, Theor. Appl. Climatol., doi:10.1007/200704-005-0217-1, 2006.
 959
 960 Tandon, N. F., Gerber, E. P. Sobel, A. H. and Polvani, L. M.: Understanding Hadley cell expansion versus
 961 contraction: Insights from simplified models and implications for recent observations, J. Climate, 26, 4304–4321,
 962 doi:10.1175/JCLI-D-12-00598.1, 2013.
 963
 964 Thompson, L. G., Mosley-Thompson, E., Davis, M. E., Lin, P. N., Henderson, K., et al.: "Tropical glacier and ice
 965 core evidence of climate change on annual to millennial time scales". Climatic Change 59: 137-155, 2003.
 966
 967 Viswanadhapalli, Y., Dasari, H. P., Langodan, S., Challa, V. S. and Hoteit, I.: Climatic features of the Red Sea from
 968 a regional assimilative model. Int. J. Climatol., 37: 2563-2581. doi:10.1002/joc.4865, 2017.
 969
 970 Vose, R. S., Easterling, D. R. and Gleason, B.: Maximum and minimum temperature trends for the globe: an update
 971 through 2004, Geophys. Res. Lett., 32, 1–5, 2005.
 972
 973 Waugh, D. W., and Polvani, L. M.: Climatology of intrusions in to the tropical upper troposphere, Geophys. Res.
 974 Lett., 27, 3857–3860, 2000.
 975
 976 Wilcox, L. J., Charlton-Perez, A. and Gray, L. J.: Trends in austral jet position in ensembles of high-and low-top
 977 CMIP5 models, J. Geophys. Res., 117, D13115, doi:10.1029/2012JD017597, 2012.
 978
 979 Wiltshire, A. J.: Climate change implications for the glaciers of the Hindu-Kush Karakoram and Himalayan region.
 980 Cryosphere, 7, 3717–3748, 2013.
 981
 982 Wittman, M. A., Charlton, A. J. and Polvani, L. M.: The effect of lower stratospheric shear on baroclinic instability,
 983 J. Atmos. Sci., 64, 479–496, doi:10.1175/JAS3828.1., 2007.
 984
 985 World Meteorological Organization: Guide to Meteorological practices, 2nd Ed. WMO No 168. Tech Paper, 82,
 986 Geneva, Switzerland, 1970.
 987
 988 Yunling, H. and Yiping, Z.: Climate change from 1960-2000 in the Lancang River Valley, China, Mountain
 989 Research and Development, 25, 341-348, 2005.
 990

991 Zarenistana, K. M., Dhorde, A. G., and Kripalani, R. H.: Temperature analysis over southwest Iran: trends and
 992 projections, Theoretical and Applied Climatology, 116, 103–117, 2014.

993

994 **Table:**

995

996 Table 1. Annual and Seasonal temperature trend in Kashmir Valley during 1980-2016

997 Table 2. Annual and Seasonal Precipitation trends in Kashmir valley during 1980-2016

998 Table 3: Mean temperature increase at each station from 1980 to 2016

999

Stations (Mann Kendall test)	Temperature Trends	Annual	Min	Max	Winter	Spring	Summer	Autumn	Abrupt Change (student' s T test
Gulmarg Critical Values a=0.10 (1.654) a=0.05(1.96) a=0.01(2.567)	Increasing trend	S=0.01	S=0.01	S=0.1	S=0.05	S=0.01	NS	S=0.05	1995
	Z statistics	3.976	3.059	1.564	2.43	2.806	0.486	2.159	

Pahalgam	Increasing trend	S=0.01	S=0.01	S=0.01	S=0.01	S=0.01	S=0.1	S=0.05	1995
	Z statistics	4.119	3.6	3.519	3.118	3.438	1.71	2.416	
Srinagar	Increasing trend	S=0.05	S=0.1	S=0.01	S=0.05	S=0.05	S=0.1	NS	1995
	Z statistics	2.108	1.392	2.804	1.992	2.413	0.374	0.198	
Kupwara	Increasing trend	S=0.01	S=0.1	S=0.01	S=0.05	S=0.01	S=0.1	S=0.1	1995
	Z statistics	3.433	1.819	3.246	1.988	2.719	1.78	1.865	
Kokarnag	Increasing trend	S=0.01	S=0.05	S=0.01	S=0.01	S=0.01	S=0.1	S=0.1	1995
	Z statistics	3.467	2.363	3.11	3.195	3.195	1.46	0.68	
Qazigund	Increasing trend	S=0.1	S=0.1	S=0.1	S=0.05	S=0.05	NS	S=0.1	1995
	Z statistics	1.717	1.77	1.68	2.026	2.236	-0.714	-1.501	
Stations (Mann Kendall test)	Precipitation Trends	Annual	Winter	Spring	Summer	Autumn	Abrupt Change (student' s T test		

Table 1 Annual and Seasonal temperature trend in Kashmir Valley during 1980-2016

Table 2. Annual and Seasonal Precipitation trends in Kashmir valley during 1980-2016

Gulmarg Critical Values a=0.10 (1.654) a=0.05(1.96) a=0.01(2.567)	decreasing trend	S=0.05	S=0.1	S=0.01	NS	NS	1995
	Z statistics	-1.988	-1.53	-2.515	-0.445	-0.394	
Pahalgam	decreasing trend	S=0.1	S=0.1	S=0.05	NS	NS	1995
	Z statistics	-1.442	-1.136	-2.151	-0.556	0.034	
Srinagar	decreasing trend	S=0.05	NS	S=0.01	NS	NS	1995
	Z statistics	-2.532	0.051	-2.060	-0.105	-1.003	
Kupwara	decreasing trend	S=0.1	S=0.1	S=0.01	NS	NS	1995
	Z statistics	-1.962	-0.817	-2.919	-0.986	-0.153	
Kokarnag	decreasing trend	S=0.1	S=0.1	S=0.05	NS	NS	1995
	Z statistics	-1.326	-1.53	-2.276	0.186	-0.119	
Qazigund	decreasing trend	S=0.05	NS	S=0.05	NS	NS	1995
	Z statistics	-1.275	-0.764	-2.413	0.359	-0.232	

Table 3: Mean temperature increase at each station from during 1980-2016.

Stations	Elevation in meters	Topography	Increase annual temperature in °C
Pahalgam	2600mts	Located on mountain top	1.13
Gulmarg	2740mts	Located on mountain top	1.04
Srinagar	1600mts	Located on plane	0.55

		surface in an urbanized area	1030
Kupwara	1670mts	Located on plane surface bounded on three sides by mountains	0.92 1031 1032 1033
Kokarnag	2000mts	Located on plane surface	0.99 1034
Qazigund	1650mts	Located on plane surface	0.78 1035

1036

1037

1038

1039

1040

1041

1042

1043 **Figure captions:**

1044

1045 Fig. 1 Geographical setting of the Kashmir valley (b) inside the Jammu and Kashmir state (a) of India (c) along with
1046 marked locations of six meteorological observation stations: Srinagar, Gulmarg, Pahalgam, Kokarnag, Qazigund and
1047 Kupwara

1048

1049 Fig. 2(a-g) Trends in surface temperature (°C) at the six interested locations of the Kashmir valley (a) for annual
1050 mean temperature, (b) maximum temperature, (c) minimum temperature, (d) winter mean temperature during
1051 December-February, (e) spring mean temperature (March-May), (f) summer mean temperature (June-August) and
1052 (g) autumn mean temperature (September-November).

1053

1054 Fig. 3(a-e) Same as Fig. 2 but for precipitation (mm) and only for means of (a) annual, (b) winter, (c) spring, (d)
1055 summer and (e) autumn.

1056

1057 Fig. 4(a) Cumulative testing for defining change point of temperature (averaged for all the six stations of the
1058 Kashmir valley), (b) same as (a) but for precipitation, (c) comparison of trends of Kashmir temperature with North

1059 Atlantic Ocean (NAO index (d) same as (c) but for precipitation, (e) regression analysis of winter temperature and
1060 (f) regression analysis of winter precipitation.
1061
1062 Fig. 5 (a) Comparison between observed and WRF model (location of Kokarnag is considered) simulated annually
1063 averaged temperature (averaged for all the stations) variations for the years 1980-2016, (b) same as (a) but for spring
1064 season, (c) for summer, (d) for autumn, (e) winter, (f) for minimum temperature and (g) maximum temperature
1065
1066 Fig. 6. Same as Fig. 5 but for precipitation. Here the minimum and maximum precipitation are not considered
1067 because it cannot be defined them properly in a day.
1068
1069 Fig. 7 (a-f) Observed monthly-averaged surface temperature and precipitation and ERA-interim potential vorticities
1070 at the 350 K potential temperature and 200 hPa level pressure surfaces for the station, Srinagar during the years
1071 1980-2016.
1072
1073 Fig. 8 (a-f) Same as the Fig. 6 but for Kokarnag.
1074
1075 Fig. 9 (a-f) Same as the Fig. 7 but for Kupwara.
1076
1077 Fig. 10 (a-f) Same as the Fig. 8 but for Pahalgam.
1078
1079 Fig. 11 (a-f) Same as the Fig. 9 but for Qazigund.
1080
1081 Fig. 12 (a-f) Same as the Fig. 10 but for Gulmarg.
1082
1083 Fig. 13 (a-f) Same as the Fig. 11 but for all the stations and during the years 2011-2016.
1084
1085 Fig. 14. (a-f) Synoptic scale ERA-interim meridional wind velocity covering the Jammu and Kashmir region for six
1086 days from 01 to 06 September 2014 (historical record flooding rainfall over this region).
1087
1088
1089
1090
1091
1092
1093
1094