

# No Robust Evidence of Future Changes in Major Stratospheric Sudden Warmings: A Multi-model Assessment from CCM1

Blanca Ayarzagüena<sup>1,2a</sup>, Lorenzo M. Polvani<sup>3</sup>, Ulrike Langematz<sup>4</sup>, Hideharu Akiyoshi<sup>5</sup>, Slimane Bekki<sup>6</sup>, Neal Butchart<sup>7</sup>, Martin Dameris<sup>8</sup>, Makoto Deushi<sup>9</sup>, Steven C. Hardiman<sup>7</sup>, Patrick Jöckel<sup>8</sup>, Andrew Klekociuk<sup>10, 11</sup>, Marion Marchand<sup>6</sup>, Martine Michou<sup>12</sup>, Olaf Morgenstern<sup>13</sup>, Fiona M. O'Connor<sup>7</sup>, Luke D. Oman<sup>14</sup>, David A. Plummer<sup>15</sup>, Laura Revell<sup>16,17</sup>, Eugene Rozanov<sup>18,16</sup>, David Saint-Martin<sup>12</sup>, John Scinocca<sup>15</sup>, Andrea Stenke<sup>16</sup>, Kane Stone<sup>19,20b</sup>, Yousuke Yamashita<sup>5c</sup>, Kohei Yoshida<sup>9</sup> and Guang Zeng<sup>13</sup>

<sup>1</sup> Dpto. Física de la Tierra y Astrofísica, Universidad Complutense de Madrid, Madrid, Spain.

<sup>2</sup> Instituto de Geociencias (IGEO), CSIC-UCM, Madrid, Spain.

10 <sup>3</sup> Columbia University, New York, USA.

<sup>4</sup> Institut für Meteorologie, Freie Universität Berlin, Berlin, Germany.

<sup>5</sup> National Institute for Environmental Studies (NIES), Tsukuba, Japan.

<sup>6</sup> LATMOS, Institut Pierre Simon Laplace (IPSL), Paris, France.

<sup>7</sup> Met Office Hadley Centre (MOHC), Exeter, UK.

15 <sup>8</sup> Institut für Physik der Atmosphäre, Deutsches Zentrum für Luft- und Raumfahrt (DLR), Oberpfaffenhofen, Germany.

<sup>9</sup> Meteorological Research Institute (MRI), Tsukuba, Japan.

<sup>10</sup> Australian Antarctic Division, Kingston Tasmania, Australia.

<sup>11</sup> Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Tasmania, Australia

<sup>12</sup> CNRM UMR 3589, Météo-France/CNRS, Toulouse, France.

20 <sup>13</sup> National Institute of Water and Atmospheric Research (NIWA), Wellington, New Zealand.

<sup>14</sup> National Aeronautics and Space Administration Goddard Space Flight Center (NASA GSFC), Greenbelt, Maryland, USA.

<sup>15</sup> Environment and Climate Change Canada, Montréal, Canada.

<sup>16</sup> Institute for Atmospheric and Climate Science, ETH Zürich (ETHZ), Switzerland.

<sup>17</sup> Bodeker Scientific, Christchurch, New Zealand

25 <sup>18</sup> Physikalisch-Meteorologisches Observatorium Davos/World Radiation Centre, Davos, Switzerland

<sup>19</sup> School of Earth Sciences, University of Melbourne, Melbourne, Australia

<sup>20</sup> ARC Centre of Excellence for Climate System Science, Sydney, Australia

30 <sup>a</sup> previously at: College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter, United Kingdom.

<sup>b</sup> now at: Massachusetts Institute of Technology (MIT), Boston, Massachusetts, USA

<sup>c</sup> now at: Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Yokohama, Japan

*Correspondence to:* Blanca Ayarzagüena (bayarzag@ucm.es)

## Abstract.

35 Major mid-winter stratospheric sudden warmings (SSWs) are the largest instance of wintertime variability in the Arctic stratosphere. Because SSWs are able to cause significant surface weather anomalies on intra-seasonal time scales, several previous studies have focused on their potential future change, as might be induced by anthropogenic forcings. However, a wide range of results have been reported, from a future increase in the frequency of SSWs to an actual decrease. Several factors might explain these contradictory results, notably the use of different metrics for the identification of SSWs,  
40 and the impact of large climatological biases in single-model studies. To bring some clarity, we here revisit the question of

future SSWs changes, using an identical set of metrics applied consistently across 12 different models participating in the Chemistry Climate Model Initiative. Our analysis reveals that no statistically significant change in the frequency of SSWs will occur over the 21<sup>st</sup> century, irrespective of the metric used for the identification of the event. Changes in other SSWs characteristics, such as their duration, deceleration of the polar night jet and the tropospheric forcing, are also assessed: again, we find no evidence of future changes over the 21<sup>st</sup> century.

## 1 Introduction

Stratospheric sudden warmings (SSWs) are the largest manifestation of the internal variability of the wintertime polar stratosphere in the Northern Hemisphere, consisting of a very rapid temperature increase accompanied by a reversal of the westerly wintertime circulation (the polar vortex). In observations, SSWs occur roughly with a frequency of 6 SSWs per decade (e.g., Charlton and Polvani, 2007). However, large variability on intra- and inter-decadal time scales has been reported (Labitzke and Naujokat, 2000; Schimanke et al., 2011).

SSWs also play an important role in the dynamical coupling between the stratosphere and troposphere. Although other mechanisms are possible, SSWs are usually related to precursors in the troposphere that lead to an anomalously high injection of tropospheric waves that propagate into the stratosphere where they deposit momentum and energy, decelerating the mean flow (Matsuno, 1971; Polvani and Waugh, 2004). More importantly, however, their effects are not restricted to the stratosphere: SSWs can also impact the tropospheric circulation and surface climate for up to two months (e.g., Baldwin and Dunkerton, 2001). Given their importance for seasonal forecasting, SSWs have been studied with great interest, as they are likely to provide a source of improved weather forecasts on intraseasonal scales (Sigmond et al., 2013).

One question of particular relevance is whether SSWs will change in the future, as a consequence of increasing greenhouse gases (GHG) concentrations and ozone recovery. The answer to this question has proven elusive since the first studies over two decades ago. While Mahfouf et al. (1994) found an increase in the frequency of SSWs under doubled CO<sub>2</sub> conditions, Rind et al. (1998) reported a decrease, and Butchart et al. (2000) did not find any change that might be attributed to increasing GHG concentrations. And, in spite of an improved stratospheric representation and more realistic model features in the last decade, a clear consensus as to future SSW changes is still missing (Charlton-Perez et al., 2008; Bell et al., 2010; SPARC CCMVal, 2010; Mitchell et al. 2012a and b; Hansen et al., 2014; Kim et al., 2017).

Several potential reasons that might explain the disparity in the projected SSW changes have been proposed in the literature. One is the combination of different aspects of future climate change with opposing effects on the Arctic stratosphere, such as the projected ozone recovery, increasing GHG concentrations and their induced changes in global sea surface temperatures. These result in a weak polar stratospheric response to climate change (Mitchell et al., 2012a, Ayarzagüena et al., 2013). Consequently, individual models yield different future projections of SSW changes, depending on the relative importance of these competing effects in each model. Hence, any result obtained with a single model needs to be taken with much caution.

Another potential explanation for the discrepancies stems from the criterion chosen for the identification of SSWs. As shown in Butler et al. (2015), the identification of SSWs can be sensitive to the method used. It was found to depend on the meteorological variable chosen for analysis, and also on whether the identification criterion entails total fields and a fixed threshold (absolute criterion), or anomalies relative to a changing climatology (relative criterion). For instance, the traditional criterion of the World Meteorological Criterion (hereafter WMO criterion, McInturff, 1978) requires the reversal of both zonal-mean zonal wind at 60°N and 10hPa and the meridional gradient of zonal mean temperature between 60°N and the pole at the same level. This criterion was empirically developed from the observations in the last several decades, and was applied in historical stratospheric analyses (e.g., Labitzke, 1981). Recent studies have continued using the WMO criterion although many of them have only imposed the reversal of the wind for the SSW identification (e.g., Charlton and Polvani, 2007). Because of its simplicity and its dynamical insight, the WMO criterion (and its recent simplified version) is the most commonly used criterion in modelling studies as well. However, such an absolute metric might not always be the best choice to measure the polar stratospheric variability in these studies, as it does not account for potential model biases in the polar vortex climatology, or possible changes in this climatology in the future projections (McLandress and Shepherd, 2009; Mitchell et al., 2012a; Butler et al., 2015; Kim et al., 2017). An analysis with the Canadian Middle Atmosphere Model by McLandress and Shepherd (2009) showed that the frequency of SSWs may or may not change depending on the detection index.

The purpose of this study, therefore, is to revisit the question of possible future SSW changes, taking these issues into consideration. Seeking a robust answer, we employ three different SSW identification criteria (both absolute and relative) and apply them consistently to the output from 12 state-of-the-art climate models (contributing to the Chemistry Climate Model Initiative, CCMI). Interactive stratospheric chemistry, which is present in all the CCMI models, makes them the most realistic in terms of stratospheric processes. In addition, the CCMI models are improved compared to their counterparts which participated in the previous Chemistry Climate Model Validation-2 programme (CCMVal-2). In particular, several CCMI models are coupled to interactive ocean modules, and the vertical resolution of many models has been increased (Morgenstern et al. 2017). Moreover, unlike other previous studies such as Kim et al. (2017), our analysis is not only restricted to the mean frequency of SSWs, but we also examine the possible future changes in other characteristics such as the duration of events, the related deceleration of the polar night jet or the wave activity preceding their occurrence. To our knowledge this is the first time that a multi-model assessment of these different SSWs features is performed. The structure of the paper is as follows: In Section 2 the data and methodology used in the analysis are described. The main results are shown in Section 3, and Section 4 includes the discussion and the most important conclusions derived from the analysis.

## **2 Data and methodology**

### **2.1. Data description**

Our study is based on the analysis of the transient REF-C2 simulation of 12 CCMI models (cf. Table 1; for more details see Morgenstern et al., 2017). The REF-C2 runs extend from 1960 to 2099 or 2100 for most models (except for the

105 IPSL-LMDZ-REPROBUS model that terminates the run in 2095), and include natural and anthropogenic forcings following the CCMI specifications (Eyring et al., 2013). In particular, GHG concentrations and surface mixing ratios of ozone depleting substances (ODS) are based on observations until 2000, and on the Representative Concentration Pathway 6.0 (RCP6.0, Meinshausen et al., 2011) and A1 (WMO, 2011) scenarios, respectively, from 2000 to 2100. Solar variability is included in most of the models. Depending on the characteristics and performance of the models, sea surface temperatures (SSTs) and the quasi-biennial oscillation (QBO) are prescribed or internally generated. Future changes in frequency and other features of SSWs are obtained by comparing the last 40 winters of each run (denoted as “the future”) to the first 40 winters (denoted as “the past”). Unless otherwise stated, anomalies are calculated from the climatology of the corresponding 40-year period. A Student’s t-test is applied to determine if the future changes are statistically significant in all cases except for the duration of SSWs where we applied a Wilcoxon ranked-sum test. The performance of the models in reproducing SSWs characteristics for the past period (1960-2000) is assessed by comparing the models to the ERA-40 and JRA-55 reanalyses (Uppala et al., 2005; Kobayashi et al., 2015). Both reanalyses extend back to before 1979; ERA-40 data starts in September 1957 and JRA-55 begins in January 1958. Thus, they cover the past period of our study. Among the few reanalyses that have available data in the pre-satellite era, ERA-40 and JRA-55 are the most suitable for middle atmosphere analyses because they have a higher top level and vertical resolution (Fujiwara et al., 2017).

## 120 **2.2 Criteria for the detection of SSWs**

As the detection of SSWs is somewhat sensitive to the chosen criterion, we use three different criteria to ensure that the conclusions regarding future changes are the same irrespective of the metric. The criteria we use are described in Butler et al. (2015) and as follows.

### *1) WMO (World Meteorological Organization) criterion*

125 SSWs are identified when the zonal-mean zonal wind at 10 hPa and 60°N and the zonal-mean temperature difference between 60°N and the pole at the same level are simultaneously reversed. Two events must be separated by at least 20 consecutive days of westerly winds. Only events from November to March are considered. Stratospheric final warmings are excluded by imposing at least 10 days with westerly winds after the occurrence of a SSW and before 30 April, to ensure the recovery of the polar vortex before its final breakup. The onset date of the event corresponds to the first day of the wind reversal.

### *2) Polar cap zonal wind reversal (u6090N)*

SSWs are identified when the area-weighted zonal wind at 10 hPa averaged over the polar cap (60°N-90°N) reverses. The separation of events and the exclusion of stratospheric final warmings are done in the same way as for the WMO criterion.

### *3) Polar cap 10hPa geopotential (ZPOL)*

135 SSWs are identified based on the polar cap standardized anomalies of 10 hPa geopotential height. The anomalies are detrended and computed following Gerber et al. (2010). A SSW is detected if the anomalies exceed three standard deviations of the climatological January to March geopotential height (Thompson et al. 2002).

Note that WMO and u6090N are absolute SSWs criteria, whereas ZPOL is a relative SSW one.

## 2.3 Other SSW characteristics

140 Beyond their frequency, we also study if the other key characteristics of SSWs, such as duration or tropospheric forcing, will change in the future. The considered events in all features are those identified by the WMO criterion, because it is a popular criterion and, as will be shown later, the conclusions relative to the frequency results are not different from those obtained for the other two criteria. These are the metrics/diagnostics applied:

### 1. *Duration:*

145 The duration of the events is computed by the number of consecutive days of easterly wind regime at 60°N and 10 hPa as in Charlton et al. (2007).

### 2. *Deceleration of the polar night jet*

The deceleration of the polar night jet associated with the occurrence of SSWs is defined as the difference in the zonal-mean zonal wind at 60°N and 10hPa, 15-5 days before the SSWs minus 0-5 days after the SSW as in Charlton and Polvani (2007).

### 150 3. *Tropospheric forcing*

The analysis of the tropospheric forcing is based on the evolution of the anomalous eddy heat flux at 100 hPa averaged between 45° and 75°N (aHF100) before and after the occurrence of SSWs. aHF100 is a measure of the injection of tropospheric wave activity into the stratosphere (Hu and Tung, 2003).

## 3. Future changes in the main characteristics of SSWs

### 155 3.1 Mean frequency

We start by considering the frequency of SSWs, and whether it is projected to change as a consequence of anthropogenic forcings. For this purpose, we have identified SSWs in the 12 models listed in Table 1, for the past and future periods, according to the three criteria presented in Section 2.2. Figure 1 shows the mean frequency of SSWs for each case.

160 In spite of some differences among the criteria, there appears to be a suggestion of a small increase in frequency in the multimodel mean (hereafter MM), but this tendency is not statistically significant at the 95% confidence level for any of the criteria, either absolute (WMO, U6090N) or relative (ZPOL). Also, while most models show a small increase in the frequency of SSWs in the future (10 of 12 models for the WMO criterion; 9 of 12 in the u6090N criterion; and 7 of 12 for the ZPOL), most of those changes are not statistically significant. Specifically, none of the models displays a statistically significant future change for the relative criterion (ZPOL) (Fig. 1c), only 3 out of 12 models show a significant increase for the WMO criterion (NIWA-UKCA, EMAC-L90 and CMAM) (Fig. 1a), and only 2 out of 12 models for the u6090N criterion (SOCOL3, EMAC-L90) (Fig. 1b). It is, however, important to note that the NIWA-UKCA and CMAM models do not simulate a realistic frequency of SSWs when compared to reanalyses for the current climate, so they may not be a reliable indicator of possible future changes. Additionally, none of the four models (NIWA-UKCA, SOCOL3, EMAC-L90 and CMAM) shows an

increase in SSWs for the three criteria simultaneously, indicating the lack of consistency for those models across the different methods. This confirms the absence of a robust future signal regarding changes in the frequency of SSWs.

A further comparison of the results for the different criteria for the past period confirms the findings of previous studies (e.g. McLandress and Shepherd, 2009) which showed that models' biases in mean state and variability affect the frequency values for the absolute criteria, since the different models show a wide range of SSW frequency values in the past period (see Fig. S1). For instance, CCSRNIES-MIROC3.2 and NIWA-UKCA show very low SSW frequencies in agreement with the fact that the polar vortex in these models is much stronger than in the reanalyses, and the opposite is seen for ACCESS CCM, CMAM and CNRM-CCM (Fig S2). Note the good agreement between the JRA-55 and ERA-40 reanalyses. Conversely, SSW frequencies computed with the relative ZPOL criterion are more similar across the models, as they are less affected by climatological model biases. Interestingly, note how the values for the relative criterion are somewhat lower in models than in the reanalyses. Since the threshold for selecting events is based on the latter, this suggests that models may be underestimating the variability of the Arctic polar stratosphere. Nevertheless, regardless of the biases of models and their different representations of the underlying processes, the null future change in the frequency of SSWs is a robust result across all examined models.

Finally, it is worth highlighting that nearly identical results to the ones obtained with the WMO criterion are found, for both past and future periods, when only the reversal of the wind at 60°N and 10hPa (Charlton and Polvani, 2007) is used as the identification criterion. It is reassuring to report that the additional temperature constraint imposed in the WMO criterion does not significantly alter the frequency of SSWs, even for the future climates. This means that most recent studies, which have used the simpler method and considered the reversal of the wind as the sole quantity for identifying SSWs, would have likely reached the same conclusions had they used the more precise WMO criterion, and can thus be considered valid.

### 3.2 Duration

Next, we turn to the duration of SSWs, for which the results are shown in Fig. 2a, for the past and future. In each period, we notice a considerable spread across the models; nonetheless, the MM value for the past period falls within the interval of reanalyses values  $\pm 1.5$  standard error. Note, however, the variability within each model is larger than that across the models. This is particularly true for the NIWA-UKCA and CCSRNIES-MIROC3.2 models, possibly as a consequence of the low number of SSWs simulated by these two models. MRI-ESM1r1 also shows a large variability in SSW duration, but only in the past period.

The key message from Fig. 2a is that the duration of SSWs does not change in the future, using the canonical 95% confidence level for each individual model. In fact, even at 90% confidence level SSWs show a statistically significant change in only one model (HaddGEM3-ES). Nevertheless, as in the case of the mean frequency, more than half of the models (7 out of 12) agree on the sign of the future change in the SSW duration (they indicate that it will be slightly shorter), but this change in the MM is not statistically significant at the 95% confidence level.

### 3.3 Deceleration of the polar night jet

The next step is the assessment of future changes in the deceleration of the polar night jet (PNJ) associated with SSWs (Fig. 2b). Similar to the duration and mean frequency, the MM value of the PNJ deceleration does not change in the future at the 95% confidence level, with only two models (EMAC-L90 and CMAM) showing a significant future reduction of the PNJ. 205 These are the same models that show a significant though small increase in SSWs in the future with the WMO criterion (an absolute criterion), but at least, in one of these models (CMAM) the climatological polar vortex is unrealistically weak.

It is also worth noting that the MM value for the past period falls out of the interval of reanalysis values  $\pm 1.5$  standard error. Thus, one could question the reliability of the future projections of the deceleration of the polar night jet during SSWs. Half of the models show values included in the reanalysis interval, and only two of these six models display a statistical change 210 in the future. Thus, we are quite confident in concluding that the PNJ deceleration during SSWs does not change in the future.

### 3.4 Tropospheric forcing

Since SSWs are usually triggered by anomalously high tropospheric wave activity entering the stratosphere in the weeks preceding the events (Matsuno, 1971; Polvani and Waugh 2004), we have analyzed the possible future changes in the injection of wave activity during the course of the occurrence of these events for the MM. Thus, as indicated in Section 2.3, 215 Fig. 3 displays the anomalous eddy heat flux at 100hPa averaged between 45°-75°N (aHF100). The results do not show a statistically significant change in any aspect of the anomalous wave activity preceding SSWs in the MM and in the individual models (not shown). In particular, neither the strong peak of aHF100 of the MM in the 10 days prior to the occurrence of events nor the general time evolution of the aHF100 are projected to change in the future (Fig.3a). Hence the common, but not statistically significant, trend of models towards shorter future SSWs mentioned above cannot be explained by changes in 220 tropospheric forcing. Additionally, when examining the two first zonal wavenumber components of the anomalous HF100, no significant future changes are found either (Fig.3b). This would also imply little change in the distribution of split and displacement SSWs.

Model projections of future aHF100 are reliable because models are able to simulate the tropospheric forcing of these events reasonably well (Fig.3). Only a few discrepancies can be seen between the MM and the mean of JRA-55 and ERA-40 225 reanalyses (Reanalyses Mean, RM, black curve). Note that we include the average of JRA-55 and ERA-40 because they show very similar results and we avoid confusion by including too many lines in the same plot. One of the discrepancies between MM and RM is that the strong peak in aHF100 in the 5 days prior to the occurrence of SSWs is weaker in the models than in observations. The reanalyses also show a secondary peak of aHF100 between -20 and -10 days that does not appear in the MM. Additionally, the contribution of the wavenumber 1 (WN1) component to the strongest wave pulse is similar or even 230 stronger than in the reanalyses (Fig.3b), but the wavenumber 2 (WN2) in the models is much weaker than in the RM. This explains the weaker total value of aHF100 in the MM than in the RM. Nevertheless, the RM is only one realization averaged over 40 years and the MM corresponds to the average over many more realizations. Thus, the multi-model/individual

realization spread possibly accounts for at least some of the mismatch between MM and RM. In any case, the models show no statistically significant changes between the past and the future.

#### 235 4. Discussion and conclusions

We have revisited the question of whether SSWs will change in the future, analysing 12 state-of-the-art stratosphere resolving models that participated in CCMI. To obtain robust results, we have used three different identification criteria (two absolute and one relative) and have applied them consistently across all 12 models. Moreover, unlike most previous multi-model comparison studies, we have not restricted our analysis to the mean frequency of SSWs, but we have also analysed  
240 other SSWs characteristics that are important for the stratosphere-troposphere coupling. In summary, our analysis reveals that:

- No statistically significant changes in the frequency of occurrence of SSWs are to be expected in the coming decades and until the end of the 21st century. This result is robust, as it is obtained with three different identification criteria.
- Other features of SSWs, such as their duration, deceleration of the polar night jet and the tropospheric precursor wave fluxes, do not change in the future either in the model simulations, in agreement with other studies such as  
245 McLandress and Shepherd (2009) or Bell et al. (2010).
- The absence of a future change in SSWs is a robust result across all models examined here, regardless of their biases or different representation of the QBO, coupling to the ocean, solar variability, etc..

Despite the lack of statistical significant changes in the frequency of SSWs, both the MM and the majority of the models analysed show a slight increase in frequency across all criteria (Fig. 1). A similar result was reported by Kim et al.  
250 (2017), who analysed the change in SSW frequency in some CMIP5 models by identifying the events based either on the reversal of the wind or the vortex deceleration. Looking at changes in the daily climatology of the zonal mean zonal wind at 10 hPa (Figs. 4a and S3), the MM and individual model simulations also provide a consistent picture, with a robust weakening of the polar night jet (PNJ) from mid-December until mid-March, the deceleration being particularly strong between mid-December and mid-February; this is in agreement with previous CMIP5 results (Manzini et al., 2014). This deceleration is,  
255 however, only statistically significant in less than half of the models (Fig. S3), explaining why we do not find a significant change in the tropospheric forcing of SSWs (Fig. 3). To determine whether these changes in the climatology of wintertime PNJ might be associated with changes in SSWs frequency, the future-minus-past difference plots of the climatological wind are shown separately for winters with and without SSWs (Fig. 4b and c, respectively). We find a weakening of the PNJ in midwinter in both cases: this allows us to conclude that the future deceleration of the PNJ is not a consequence of a higher  
260 frequency of SSWs. This deceleration might be related to a general increase in the total stratospheric variability that, in the case of winters without SSWs, would correspond to a higher frequency of minor warmings. However, this possibility is unlikely because we do not find a robust future increase in the standard deviation of zonal-mean zonal wind at 10hPa across the models or a change in the shape of the distribution of the zonal-mean zonal wind as shown in Fig. S4. Perhaps the future deceleration of the PNJ might explain the statistically significant increase in SSWs in a few models, using the absolute criteria



265 in agreement with McLandress and Shepherd (2009). In any case, these signals are small and it is nearly impossible to untangle the cause and the effect, as these changes occur simultaneously.

More importantly our findings dispel, to a large degree, the confusion in the literature regarding future SSW changes, and suggest that previous reports of significant changes are likely to be artefacts, caused by biases associated with individual models, or by flaws in the identification methods used (or both). In addition, the analysis of other features of SSWs besides  
270 the mean frequency supports the key finding of our study, i.e. that anthropogenic forcings will not affect SSWs over the 21<sup>st</sup> century. Our results confirm and expand the findings of Kim et al. (2017) who did not find a statistically significant future change in the frequency of SSWs in CMIP5 models. Note that although the key finding of our study is a null result, it is by no means uninteresting. Just to offer one example: Kang and Tziperman (2017) have recently proposed that future changes in the Madden Julian Oscillation (which are expected to occur with increased levels of CO<sub>2</sub> in the atmosphere) will cause an increased  
275 occurrence of SSWs. While their conclusion may be correct, our findings indicate that it can be misleading to project changes in the SSWs on the basis of a single mechanism: the complexity of the climate system is such that multiple mechanisms may be at play, with likely opposite effects which may result in net changes that are not statistically significant.

One may argue that the lack of a statistically significant future change in our study could be explained, at least partially, by the high interannual variability of the boreal polar stratosphere in 40-year periods (e.g., Langematz and Kunze,  
280 2006), or perhaps by the natural variability on longer time-scales coming from other subcomponents of the climate system (e.g.: Schimanke et al., 2011). As shown in a recent paper, 10 identically forced model simulations, over the 50-year period 1952-2003, exhibit great differences in the number of SSWs, and these differences are solely due to internal variability (Polvani et al, 2017). This means that the 40 years of observations at our disposal may not represent the mean of a distribution, but could happen to be an outlier. Needless to say, we have no means of determining whether this is the case, as we do not have  
285 long enough observations.

One might also object that the forcing in the scenario used of our runs (RCP6.0) is not extreme enough to produce a significant signal in the frequency and duration of SSWs, but that a significant change would occur with stronger forcing, such as the RCP8.5 scenario. Although we cannot rule out this possibility, it seems improbable based on a similar lack of significance in the results documented for that very extreme scenario by several previous studies (Mitchell et al. 2012a;  
290 Ayarzagüena et al. 2013, Hansen et al. 2014; Kim et al. 2017). Nevertheless, it would be hard to verify the hypothesis because of the low number of CCMI RCP8.5 simulations available.

Finally, in recent years much activity has been devoted to search for novel criteria for the identification of SSWs (Butler et al., 2015). One of the reasons given to justify the implementation of a new metric was that the traditional WMO criterion was not appropriate for modelling studies, as it was based on observationally chosen parameters, such as the location  
295 of the polar night jet. However, our results show that this criterion performs well under a changing climate, provided models are able to reproduce correctly the past stratospheric variability. Thus, considering the good agreement among the three criteria used here on the lack of change in future SSWs, and given the dynamical implications for the propagation of planetary waves into the stratosphere, we suggest that the WMO criterion is appropriate for the study of SSWs in the future if the model can

represent well the stratospheric variability. Furthermore, since the simplest (and most commonly used) criterion, involving  
300 only the zonal winds (Charlton and Polvani, 2007), yields identical results as the WMO criterion, one could argue that the  
simplest method may suffice in most cases for the study of SSWs, and that more complex criteria might not be worth the  
trouble. A similar conclusion was reached, independently, by Butler and Gerber (2018) who methodically assessed different  
metrics and concluded that the simplest algorithm is within the optimal range.

**Data availability.** Data of this manuscript has been mostly downloaded from the Centre for Environmental Data Analysis  
305 (CEDA, 2017; <http://catalogue.ceda.ac.uk/uuid/9cc6b94df0f4469d8066d69b5df879d5> ) or supplied directly by the co-authors.  
For instructions for access to this archive see <http://blogs.reading.ac.uk/ccmi/badc-data-access>. The data supplied by the co-  
authors will in due course be uploaded to the CEDA archive.

**Competing interests.** The authors declare that they have no conflict of interest.

### Acknowledgements

310 We acknowledge the modeling groups for making their simulations available for this analysis, the joint WCRP SPARC/IGAC  
Chemistry-Climate Model Initiative (CCMI) for organizing and coordinating the model data analysis activity, and the British  
Atmospheric Data Centre (BADC) for collecting and archiving the CCMI model output. BA was funded by the European  
Project 603557-STRATOCLIM under the FP7-ENV.2013.6.1-2 programme, and “Ayudas para la contratación de personal  
postdoctoral en formación en docencia e investigación en departamentos de la Universidad Complutense de Madrid”. BA and  
315 UL wish to acknowledge the “Deutsche Forschungsgemeinschaft” (DFG) within the research programme SHARP under the  
grant LA 1025/15-1. LMP is grateful for the continued support of the US National Science Foundation. The work of NB, SCH,  
and FOC was supported by the Joint BEIS/Defra Met Office Hadley Centre Climate Programme (GA01101). NB and SCH  
were supported by the European Community within the StratoClim project (Grant 603557). OM and GZ acknowledge the UK  
Met Office for use of the MetUM. This research was supported by the NZ Government’s Strategic Science Investment Fund  
320 (SSIF) through the NIWA programme CACV. OM acknowledges funding by the New Zealand Royal Society Marsden Fund  
(grant 12-NIW-006) and by the Deep South National Science Challenge (<http://www.deepsouthchallenge.co.nz>). The authors  
wish to acknowledge the contribution of NeSI high-performance computing facilities to the results of this research. New  
Zealand’s national facilities are provided by the New Zealand eScience Infrastructure (NeSI) and funded jointly by NeSI’s  
collaborator institutions and through the Ministry of Business, Innovation & Employment’s Research Infrastructure  
325 programme (<https://www.nesi.org.nz>). The EMAC simulations have been performed at the German Climate Computing Centre  
(DKRZ) through support from the Bundesministerium für Bildung und Forschung (BMBF). DKRZ and its scientific steering  
committee are gratefully acknowledged for providing the HPC and data archiving resources for the consortial project ESCiMo

(Earth System Chemistry integrated Modelling). CCSRNIES's research was supported by the Environment Research and Technology Development Funds of the Ministry of the Environment (2-1303) and Environment Restoration and Conservation Agency (2-1709), Japan, and computations were performed on NEC-SX9/A(ECO) and NEC SX-ACE computers at the CGER, NIES. The authors wish to thank two anonymous referees for their helpful comments.

## References

- Ayarzagüena, B., Langematz, U., Meul, S., Oberländer, S., Abalichin, J., and Kubin, A.: The role of climate change and ozone recovery for the future timing of major stratospheric warmings, *Geophys. Res. Lett.*, 40, 2460-2465, 2013.
- 335 Baldwin, M. P. and Dunkerton, T.J.: Stratospheric harbingers of anomalous weather regimes, *Science*, 294, 581-583, 2001.
- Bell, C. J., Gray, L. J., and Kettleborough, J.: Changes in Northern Hemisphere stratospheric variability under increased CO<sub>2</sub> concentrations, *Q. J. R. Meteorol. Soc.*, 136, 1181-1190, 2010.
- Butchart, N., Austin, J., Knight, J. R., Scaife, A. A., and Gallani, M. L.: The response of the stratospheric climate to projected changes in the concentrations of well-mixed greenhouse gases from 1992 to 2015, *J. Climate*, 13, 2142-2159, 2000.
- 340 Butler, A. H., Seidel, D. J., Hardiman, S. C., Butchart, N., Birner, T., and Match, A.: Defining sudden stratospheric warmings, *Bull. Amer. Meteor. Soc.*, 96, 1913-1928, 2015.
- Butler, A. H., Gerber, E. P.: Optimizing the definition of a sudden stratospheric warming. *J. Climate*, in press, 2018.
- Charlton, A. J., and Polvani, L. M.: A new look at stratospheric sudden warmings. Part I: Climatology and modelling benchmarks, *J. Climate*, 20, 449-469, 2007.
- 345 Charlton, A. J., Polvani, L. M., Perlwitz, J., Sassi, F., Manzini, E., Shibata, K., Pawson, S., Nielsen, J. E., and Rind, D.: A new look at stratospheric sudden warmings. Part II: Evaluation of numerical model simulations, *J. Climate*, 20, 470-488, 2007.
- Charlton-Perez, A. J., Polvani, L. M., Austin, J., and Li, F.: The frequency and dynamics of stratospheric sudden warmings in the 21<sup>st</sup> century, *J. Geophys. Res.*, 113, D16116, doi: 10.1029/2007JD009571, 2008.
- Eyring, V., Lamarque, J.-F., Hess, P., Arfeuille, F., Bowman, K., Chipperfield, M. P., Duncan, B., Fiore, A., Gettelman, A., Giorgetta, M.A., Granier, C., Hegglin, M., Kinnison, D., Kunze, M., Langematz, U., Luo, B., Martin, R., Matthes, K., Newman, P. A., Peter, T., Robock, A., Ryerson, T., Saiz-Lopez, A., Salawitch, R., Schultz, M., Shepherd, T. G., Shindell, D., Staehelin, J., Tegtmeier, S., Thomason, L., Tilmes, S., Vernier, J. -P., Waugh, D. W., and Young, P. J.: Overview of IGAC/SPARC Chemistry-Climate Model Initiative (CCMI) community simulations in support of upcoming ozone and climate assessments, *SPARC Newsletter*, 40, 48-66, 2013.
- 350 Fujiwara, M., et al.: Introduction to the SPARC Reanalysis Intercomparison Project (S-RIP) and overview of the reanalysis systems, *Atmos. Chem. Phys.*, 17, 1417-1452, 2017.
- Gerber, E. P., Baldwin, M. P., Akiyoshi, H., Austin, J., Bekki, S., Braesicke, P., Butchart, N., Chipperfield, M., Dameris, M., Dhomse, S., Frith, S. M., Garcia, R. R., Garny, H., Gettelman, A., Hardiman, S. C., Karpechko, A., Marchand, M., Morgenstern, O., Nielsen, J. E., Pawson, S., Peter, T., Plummer, D. A., Pyle, J. A., Rozanov, E., Scinocca, J. F., Shepherd,

- 360 T. G., and Smale, D.: Stratosphere-troposphere coupling and annular mode variability in chemistry-climate models, *J. Geophys. Res.*, 115, doi: 10.1029/2009JD013770, 2010.
- Hansen, F., Matthes, K., Petrick, C., and Wang, W.: The influence of natural and anthropogenic factors on major stratospheric sudden warmings, *J. Geophys. Res.*, 119, 8117-8136, 2014.
- Hu, Y., and Tung, K. K.: Possible ozone-induced long-term changes in planetary wave activity in late winter, *J. Climate*, 16,  
365 3027-3038, 2003.
- Kang, W. and Tziperman, E.: More frequent sudden stratospheric warming events due to enhanced MJO forcing expected in a warmer climate. *Journal of Climate*, 30, 8727-8743, 2017
- Kim, J., Son, S.-W., Gerber, E.P, and Park, H.-S.: Defining sudden stratospheric warming in climate models: Accounting for biases in model climatologies, *J. Climate*, 30, 5529-5546, 2017.
- 370 Kobayashi, S. et al.: The JRA-55 reanalysis: General specifications and basic characteristics, *J. Meteor. Soc. Japan*, 93, 5-48, 2015.
- Labitzke, K.: Stratospheric-mesospheric midwinter disturbances: A summary of observed characteristics, *J. Geophys. Res.* 86:9665-9678, 1981.
- Labitzke, K., and Naujokat, B.: The lower arctic stratosphere in winter since 1952, *SPARC Newsl.*, 15, 11-14, 2000.
- 375 Langematz, U., and Kunze, M.: An update on dynamical changes in the Arctic and Antarctic stratospheric polar vortices, *Clim. Dyn.*, 27, 647-660, 2006.
- Mahfouf, J. F., Cariolle, D., Geleyn, J.-F., Timbal, B.: Response of the Météo-France climate model to changes in CO<sub>2</sub> and sea surface temperature, *Clim. Dyn.*, 9, 345-362, 1994.
- Manzini, E., Karpechko, A. Y., Anstey, J., Baldwin, M. P., Black, R. X., Cagnazzo, C., Calvo, N., Charlton-Perez, A.,  
380 Christiansen, B., Davini, P., Gerber, E., Giorgetta, M., Gray, L., Hardiman, S. C., Lee, Y.-Y., Marsh, D. R., McDaniel, B. A., Purich, A., Scaife, A. A., Shindell, D., Son, S. -W., Watanabe, S., and Zappa, G.: Northern winter climate change: Assessment of uncertainty in CMIP5 projections related to stratosphere-troposphere coupling, *J. Geophys. Res.*, 119, 7979-7998, doi: 10.1002/2013JD021403, 2014.
- Matsuno, T.: A dynamical model of stratospheric sudden warming, *J. Atmos. Sci.*, 28, 1479-1494, 1971.
- 385 McLandress, C., and Shepherd, T. G.: Impact of climate change on stratospheric sudden warmings as simulated by the Canadian Middle Atmosphere Model, *J. Climate*, 22, 5449-5463, 2009.
- McInturff, R. M., Ed.: Stratospheric warmings: Synoptic, dynamic and general-circulation aspects. NASA Reference Publ. NASA-RP-1017, 174 pp, 1978.[Available online at <http://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/19780010687.pdf>.]
- Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M. L. T., Lamarque, J-F., Matsumoto K., Montzka, S. A.,  
390 Raper, S. C. B., Riahi, K., Thomson, A., Velders, G. J. M., and van Vuuren, D. P. P.: The RCP greenhouse gas concentrations and their extensions from 1765 to 2300, *Climatic Change*, 109, 213-241, 2011.

- Mitchell, D. M., Osprey, S. M., Gray, L. J., Butchart, N., Hardiman, S. C., Charlton-Perez, A. J., and Watson, P.: The effect of climate change on the variability of the Northern Hemisphere stratospheric polar vortex, *J. Atmos. Sci.*, 69, 2608-2618, 2012a.
- 395 Mitchell, D. M., Charlton-Perez, A. J., Gray, L. J., Akiyoshi, H., Butchart, N., Hardiman, S. C., Morgenstern, O., Nakamura, T., Rozanov, E., Shibata, K., Smale, D., and Yamashita, Y.: The nature of Arctic polar vortices in chemistry-climate models, *Q. J. R. Meteorol. Soc.*, 138, 1681-1691, 2012b.
- Morgenstern, O., Hegglin, M. I., Rozanov, E., O'Connor, F. M., Abraham, N.L., Akiyoshi, H., Archibald, A. T., Bekki, S., Butchart, N., Chipperfield, M. P., Deushi, M., Dhomse, S. S., Garcia, R. R., Hardiman, S. C., Horowitz, L. W., Jöckel, P.,  
400 Josse, B., Kinnison, D., Lin, M., Mancini, E., Manyin, M. E., Marchand, M., Marécal, V., Michou, M., Oman, L. D., Pitari, G., Plummer, D. A., Revell, L. E., Saint-Martin, D., Schofield, R., Stenke, A., Stone, K., Sudo, K., Tanaka, T. Y., Tilmes, S., Yamashita, Y., Yoshida, K., and Zeng, G.: Review of the global models used within the phase 1 of the Chemistry-Climate Model Initiative (CCMI), *Geosci. Model Dev.*, 10, 639-671, 2017.
- Polvani, L. M., Waugh, D. W.: Upward wave activity flux as a precursor to extreme stratospheric events and subsequent  
405 anomalous surface weather regimes, *J. Climate*, 17, 3548-3554, 2004.
- Polvani, L.M., Sun, L., Butler, A.H., Richter, J.H. and Deser, C.: Distinguishing stratospheric sudden warmings from ENSO as key drivers of wintertime climate variability over the North Atlantic and Eurasia, *J. Climate*, 30, 1959-1969, 2017.
- Rind, D., Shindell, D., Lonergan, P., Balachandran, N. K.: Climate change and the middle atmosphere. Part III: The doubled CO<sub>2</sub> climate revisited, *J. Climate*, 11, 876-894, 1998.
- 410 Schimanke, S., Körper, J., Spanghel, T., and Cubasch, U.: Multi-decadal variability of sudden stratospheric warmings in an AOGCM, *Geophys. Res. Lett.*, 38, L01801, doi: 10.1029/2010GL045756, 2011.
- Sigmond, M., Scinocca, J.F., Kharin V. V., and Shepherd, T. G.: Enhanced seasonal forecast skill following stratospheric sudden warmings, *Nat. Geos.*, 6, 98-102, 2013.
- SPARC CCMVal, SPARC Report on the Evaluation of Chemistry-Climate Models, edited by V. Eyring, T. G. Shepherd, and  
415 D. W. Waugh, SPARC Report No. 5, WCRP-132, WMO/TD-No. 1526, pp 109-148, 2010.
- Thompson, D. W. J., Baldwin, M. P., and Wallace, J. M.: Stratospheric connection to Northern Hemisphere wintertime weather: Implications for prediction, *J. Climate*, 15, 1421-1428, 2002.
- Uppala, S.M., et al.: The ERA-40 re-analysis, *Q. J. R. Meteor. Soc.*, 131, 2961-3012, 2005.
- WMO (World Meteorological Organization): Scientific assessment of ozone depletion: 2010, Geneva, Switzerland, 2011.

## Tables

**Table 1. Main characteristics relative to the models and their REF-C2 simulations used in this study.**

CCMI models	Model resolution	QBO	Solar variability	SSTs
GEOS-CCM	2.5° x 2°, L72 (top:0.01hPa)	Internally generated	Yes	Prescribed (CESM1)
CNRM-CCM	T42L60 (top: 0.07 hPa)	Internally generated	Yes	Prescribed (CNRM)
NIWA-UKCA	3.75° x 2.5°, L60 (top: 84 km)	Internally generated	No	Coupled to ocean model
CCSRNIES-MIROC 3.2	T42L34 (top: 0.012 hPa)	Nudged	Yes	Prescribed (MIROC 3.2)
IPSL-LMDZ-REPROBUS	3.75° x 2.5°, L39 (top: 70 km)	Nudged	Yes	Prescribed (SRES A1b IPSL)
ACCESS CCM	3.75° x 2.5°, L60 (top: 84 km)	Internally generated	No	Prescribed (HadGEM2-ES)
HadGEM3-ES	1.875°x1.25°, L85 (top: 85 km)	Internally generated	Yes	Coupled to ocean model
SOCOL3	T42L39 (top: 0.01hPa)	Nudged	Yes	Prescribed (CESM1(CAM5))
MRI-ESM1r1	TL159L80 (top: 0.01hPa)	Internally generated	Yes	Coupled to ocean model
EMAC-L47	T42L47 (top: 0.01hPa)	Nudged	Yes	Prescribed (HadGEM2-ES)
EMAC-L90	T42L90 (top: 0.01hPa)	Internally generated (slightly nudged)	Yes	Prescribed (HadGEM2-ES)
CMAM	T47L71 (top: 0.0575 hPa)	No	No	Prescribed (CanCM4)






Figure 1: (a) Mean frequency of stratospheric sudden warmings per decade for the past (blue bars) and the future (red bars) for all models, the multimodel mean (MM) and JRA-55 and ERA-40 reanalyses (black bars) according to the WMO criterion. (b) – (c) Same as (a) but for the u6090N and ZPOL, respectively. Green stars on top of the future bar denote a statistically significant change in the frequency of SSWs in the future at the 95% confidence level.



**Figure 2.** (a) Duration of SSWs (in days) and (b) deceleration of the PNJ associated with SSWs (in  $m s^{-1}$ ) in each model for both periods of study. Bars denote  $\pm 1.5$  standard error and green stars would indicate future values that are statistically significantly different from the past ones at the 95% confidence level.





435

**Figure 3. (a) Multimodel mean of anomalous heat flux ( $\text{K m s}^{-1}$ ) at 100hPa averaged over  $45^{\circ}\text{N}$ - $75^{\circ}\text{N}$  from 30 days before until 10 days after the occurrence of SSWs. (b) Same as (a) but for WN1 (solid lines) and WN2 (dashed lines) wave components. Thick lines denote statistically significant future values different from the past ones at the 95% confidence level. RM stands for the Reanalyses (JRA-55 and ERA-40) Mean.**

440




Figure 4. (a) Multimodel mean of future-minus-past differences in the daily climatology of 5-day running mean of zonal mean zonal wind at 10 hPa. (b) Same as (a) but only for winters with SSWs. (c) Same as (a) but for winters without SSWs. Shading interval: 1  $\text{m s}^{-1}$ . Dots indicate where at least 75% of the models coincide in sign with the multimodel mean.

445