



Direct measurements of NO₃-reactivity in and above the boundary layer of a mountain-top site: Identification of reactive trace gases and comparison with OH-reactivity.

Jonathan M. Liebmann¹, Jennifer B. A. Muller², Dagmar Kubistin², Anja Claude², Robert Holla², Christian Plaß-Dülmer², Jos Lelieveld¹ and John N. Crowley¹

¹Atmospheric Chemistry Department, Max Planck Institut für Chemie, 55128, Mainz, Germany

²Meteorologisches Observatorium Hohenpeissenberg, Deutscher Wetterdienst, 82383, Hohenpeissenberg, Germany

10

Correspondence to: John Crowley (john.crowley@mpic.de)

Abstract. We present direct measurements of the summertime, total reactivity of NO₃ towards organic trace gases, $k_{\text{OTG}}^{\text{NO}_3}$, at
a rural mountain site (988 m a.s.l.) in southern Germany in 2017. The diel cycle of $k_{\text{OTG}}^{\text{NO}_3}$ was strongly influenced by local
meteorology with reactivity high during the day (values of up to 0.3 s⁻¹) but usually close to the detection limit (0.005 s⁻¹) at
night when the measurement site was in the residual layer / free troposphere. Daytime values of $k_{\text{OTG}}^{\text{NO}_3}$ were sufficiently large
that the loss of NO₃ due to reaction with organic trace gases competed with its photolysis and reaction with NO. Within
experimental uncertainty, monoterpenes and isoprene accounted for all of the measured NO₃-reactivity. Averaged over the
daylight hours, more than 25% of NO₃ was removed via reaction with biogenic volatile organic compounds (BVOCs),
implying a significant daytime loss of NO_x and formation of organic nitrates due to NO₃ chemistry. Ambient NO₃
concentrations were measured on one night and were comparable to those derived from a stationary state calculation using
measured values of $k_{\text{OTG}}^{\text{NO}_3}$. We present and compare the first simultaneous, direct-reactivity measurements for the NO₃ and
OH radicals. The decoupling of the measurement site from ground level emissions resulted in lower reactivity at night for
both radicals, though the correlation between OH- and NO₃-reactivity was weak as would be anticipated given their
divergent trends in rate constants with many organic trace gases.





1 Introduction

Hydroxyl (OH) and nitrate radicals (NO₃) play a centrally important role in cleansing the atmosphere of trace gas emissions resulting from both anthropogenic and biogenic activity (Lelieveld et al., 2004; Lelieveld et al., 2016; Ng et al., 2017). Whereas OH is largely photochemically generated and present at its highest concentrations during the day, NO₃ is generated through the oxidation of NO₂ by O₃ and, due to its rapid photolysis and reaction with NO, is present mainly at night. A further important difference in the roles of OH and NO₃ in the atmosphere is related to the mechanism of their reactions. NO₃ reacts rapidly via electrophilic addition to unsaturated organic trace gases but reacts comparatively slowly (via H-abstraction) with saturated organics. In the presence of O₂, the initial addition step results in the formation of nitrooxyalkyl peroxy radicals, which can react with HO₂, NO, NO₂ or NO₃ to form multifunctional peroxides and organic nitrates (Fry et al., 2014; Ng et al., 2017).

OH can react both by addition and H-abstraction to organic and inorganic trace gases and may be considered to be more reactive and much less selective than the NO₃ radical. The distinct reaction modes leads to significant differences in the lifetimes of both radicals, which for OH are typically less than 1s and for NO₃ can exceed 1 hour (Wayne et al., 1991; Atkinson, 2000; Atkinson and Arey, 2003a; Brown and Stutz, 2012; Liebmam et al., 2018). Maximum daytime concentrations of OH are typically less than 1 pptv, whereas NO₃ has been observed at the 10s to 100s of pptv levels during nighttime (Noxon et al., 1978; Sobanski et al., 2016; Ng et al., 2017).

The large NO₃ mixing ratios at nighttime and the large rate constants for reaction of NO₃ with several unsaturated, biogenic VOCs result in NO₃ being the dominant sink of many BVOCs (Wayne et al., 1991; Atkinson, 2000; Atkinson and Arey, 2003a, b; Long et al., 2011; Brown and Stutz, 2012; Liebmam et al., 2017) especially those whose emission is mainly temperature dependent and continues at nighttime, e.g. monoterpenes (Hakola et al., 2012). The importance of NO₃ on a global scale is highlighted by the fact that forest ecosystems (covering around 9% of the world's surface) annually release ≈ 1000 Tg of biogenic volatile organic compounds (BVOC, e.g. isoprene (2-methyl-1,3-butadiene), monoterpenes (C₁₀H₁₆) and sesquiterpenes (C₁₅H₂₄)) into the Earth's atmosphere (Guenther et al., 2012; Bastin et al., 2017). BVOCs have a strong impact on the atmospheric radical budget, the NO_x cycle (Hakola et al., 2003; Holzke et al., 2006; Nölscher et al., 2013) as well as on the formation and growth of organic particles (Jaoui et al., 2013; Lee et al., 2016; Ng et al., 2017) hence understanding their lifetime and fate is essential for predicting atmospheric processes and climate change (Lelieveld et al., 2008; Lelieveld et al., 2016). In addition, NO₃ is an intermediate in the step-wise oxidation of NO to N₂O₅ (R1-R2, R4) and its lifetime with respect to reaction with biogenic trace gases (R6) impacts on NO_x levels and thus on photochemical O₃ formation from NO₂ photolysis.





The organic nitrates formed in the multi-step reaction (R6) can transfer to the particle phase or be lost through deposition;

5 N_2O_5 formed in (R4) can react with aqueous particles to form particulate nitrate and/or ClNO_2 (R7) (Osthoff et al., 2008; Phillips et al., 2012; Bannan et al., 2015; Phillips et al., 2016) thus reducing the rate of photochemical O_3 production (Dentener and Crutzen, 1993). The absolute and relative fluxes through (R6) and (R7) thus control to some extent the lifetime of NO_x .

10 Direct NO_3 -reactivity measurements have recently become possible (Liebmann et al., 2017) and the first deployment in a forested region revealed a large NO_3 -reactivity at canopy height, not all of which could be accounted for by simultaneous measurements of a large suite of organic trace gases (Long et al., 2011) pointing towards unmeasured monoterpenes as well as sesquiterpenes likely to be responsible. The difference between the observed (or derived) NO_3 -reactivity and that calculated from summing loss rates for a set of reactive trace gases is generally termed “missing reactivity” as frequently reported for OH (Nölscher et al., 2012). Previous work on NO_3 -reactivity has also revealed a strong meteorological influence

15 on the NO_3 lifetime, especially when air masses are decoupled from the surface layer in which reactive trace gases (NO and BVOC) are emitted at night (Brown et al., 2007b; Brown et al., 2011; Long et al., 2011; Sobanski et al., 2016).

In this paper we describe direct measurements of the NO_3 -reactivity in ambient air on a rural mountain site in southern Germany and interpret the data based on measured VOCs and in terms of the underlying meteorological situation. We also compare NO_3 -reactivity to simultaneous measurements of OH-reactivity over the same period.

20 2 Site description and methods

During the period 20.07.17 to 6.08.17 NO_3 -reactivity measurements were conducted in parallel with ongoing observations at the Meteorological Observatory Hohenpeissenberg (MOHp) in Bavaria, southern Germany. The observatory is a meteorological monitoring and Global Atmosphere Watch site, operated by the German Meteorological Service (DWD). It is located on the Hohenpeissenberg mountain (988 m a.s.l.), 300-400 m above the surrounding countryside about 40 km from

25 the northern rim of the Alps and has been the location of several intensive field campaigns (Plass-Dulmer et al., 2002; Birmili et al., 2003; Handisides et al., 2003; Mannschreck et al., 2004; Bartenbach et al., 2007; Hock et al., 2008; Novelli et al., 2017). The vegetation around the measurement site consists of coniferous trees and beeches growing on the slopes of the mountain while grassland and marshes are dominant in the valley. Tourism related vehicular emissions represent a potential source of local anthropogenic pollution especially at the weekends. The nearest city, Munich, is about 70 km to the

30 northeast.



Trace gases were sampled into the NO₃-reactivity and NO₂-CRD instruments through 1-inch outer-diameter PFA tubing (20 m long, operated at a flow of 40 dm³ min⁻¹) located 1.5 m above the roof, directly next to the VOC inlet. The inlet was circa 3 m distance from the inlet used for the other NO_x measurements and circa 2 m distance from the OH-reactivity inlet.

2.1 NO₃-reactivity measurements

- 5 The NO₃-reactivity instrument was operated in a laboratory located in the 3rd floor of the of the MOhp station building at the Hohenpeissenberg. Air samples were drawn at a flow rate of 2900 cm³ (STD) min⁻¹ through a 2 μm membrane filter (Pall Teflon) and 4 m of PFA tubing (6.35 mm OD) from the centre of the bypass flow (see above) resulting in 7.5 s residence time for the transport of air from the sampling point. During night time (≈19:00-03:50 UTC) ambient air samples were drawn through a heated glass flask (35 °C, residence time 20 s) to destroy ambient N₂O₅ and NO₃ which would potentially
- 10 interfere with the reactivity measurements. Operational details of the instrument were recently described by Liebmann et al. (2017). NO₃ radicals were generated by mixing NO₂ and O₃ at elevated pressure (1.5 bar, ≈ 5 minutes reaction time) and passing the mixture through an oven at ≈ 100 °C to convert all N₂O₅ to NO₃ (R2-R5). The effluent from the oven was mixed with either zero-air or ambient air in a flow-tube thermostatted to 21 °C to yield a typical (initial) NO₃ mixing ratios of 40-60 pptv.
- 15 After a fixed reaction time, the remaining NO₃ was detected by cavity-ring-down spectroscopy (CRDS) at 662 nm. The lower pressure at the top of the Hohenpeissenberg station (903 ± 8 hPa) meant that the reaction time was reduced from 10.5 s as previously reported (Liebmann et al., 2017), to 9.5 s. The measurement cycle was typically 400 s for synthetic air and 1200 s for ambient air, with intermittent signal zeroing (every ≈ 100 s) by addition of NO. The fractional loss of NO₃ in ambient air compared to zero air was converted to a reactivity via numerical simulation of a simple reaction scheme
- 20 (Liebmann et al., 2017) using measured amounts of NO, NO₂ and O₃. The parameter obtained, $k_{\text{OTG}}^{\text{NO}_3}$, is a NO₃ loss rate constant from which contributions by NO and NO₂ have been removed, and thus refers to reactive loss to organic trace gases (OTG) only. Throughout the manuscript, NO₃-reactivity and $k_{\text{OTG}}^{\text{NO}_3}$ are equivalent terms, with units of s⁻¹. The upper measurement limit to $k_{\text{OTG}}^{\text{NO}_3}$ was 45 s⁻¹, achieved by automated, dynamic dilution of the air sample. The lower limit was 0.005 s⁻¹, defined by the stability of the NO₃ source. Online calibration of the reactivity using an NO standard was performed every
- 25 ≈2 hours for 10 min. The uncertainty of the measurement was between 0.015 and 0.205 s⁻¹, depending mainly on dilution accuracy, NO levels and stability of the NO₃ source as described by Liebmann et al. (2017). Since its first description in Liebmann et al. (2017), the instrument has been extended with a further cavity to measure mixing ratios of NO₂ (see below).

2.2 NO₂, NO and O₃ measurements

- 30 Since its first deployment, the NO₃-reactivity instrument described by Liebmann et al. (2017, 2018) has been extended with a further cavity to measure NO₂. This is described here for the first time and thus in detail. The CRDS measurement of NO₂



uses a 2500 Hz, square-wave modulated, 40 mW laser-diode located in a Thor Labs LDM 21 housing and thermally stabilized at 36 °C using a Thor Labs ITC 510 Laser-Diode Combi Controller to produce light at 405 nm (0.5 nm full-width at half-maximum). The laser-diode emission is first directed through an optical isolator (Thorlabs IO-3D-405-PBS), focused by a lens (Thorlabs C340TMD-A) into the optical-fibre (0.22 NA, 50 µm core, 400-2400 nm) and then collimated (Thor Labs FiberPort Collimator PAF-X-7-A) to a beam diameter of about 6 mm before entering the cavity. Part of the laser emission was directed to an Ocean Optics spectrograph to continuously measure the laser emission spectrum.

The NO₂ cavity (Teflon-coated glass (DuPont, FEP, TE 9568), length 70 cm, volume 79 cm³) was operated at 30 °C at a flow rate of 3000 cm³ (STP) min⁻¹ resulting in a residence time of approximately 1 s. To remove particles, air was drawn through a 2 µm membrane filter (Pall Teflon) from the centre of the same high-flow bypass used for the NO₃-reactivity measurements. Light exiting the cavities through the rear mirror was detected by a photomultiplier (Hamamatsu E717-500) which was screened by a 405 nm interference filter. The pre-amplified PMT signal was digitized and averaged with a 10 MHz, 12 bit USB scope (Picoscope 3424) which was triggered at the laser modulation frequency of 2500 Hz.

The ring-down constant in the absence of NO₂ was obtained by adding zero air every 30 points of measurement for approximately 15 s. The L/d ratio (the ratio of the distance between the cavity mirrors, L , and the length of the cavity that is filled by absorber, d) was determined as described previously (Schuster et al., 2009; Crowley et al., 2010) and was 1.00 ± 0.03 . Inverse decay-constants in dry zero-air at 660 Torr were usually between 28 and 31 µs indicating optical path lengths of ≈ 8 -9 km. The measurement precision (6 s integration) was circa 150 pptv. The cavity was not pressure stabilized leading to a pressure difference of circa 2.5 Torr when switching from ambient air to zero measurements. The data was corrected for the change in Rayleigh scattering resulting from the pressure difference (typically 120 pptv) and also different relative humidities (typically 60 to 100 pptv) when switching from ambient to zero-air measurement as described by Thieser et al. (2016). The laser spectrum was measured every hour and used to calculate an effective cross-section ($\approx 6.00 \times 10^{-19}$ cm² molecule⁻¹) using a literature absorption spectrum (Voigt et al., 2002). The overall uncertainty of the measurement is mainly determined by the uncertainty in the cross section (6%). Other contributions are from NO₂ formation (from reaction of NO with O₃ in the inlet lines, $\approx 0.5\%$) and the correction for humidity and pressure differences (5%), and an error in the L/d ratio (2%), giving an estimated uncertainty of 9%. The detection limit of the instrument can be estimated from the variability in the zeros and was usually around 150 pptv.

NO₂ measurements were made from the 20th of July to the 4th of August with breaks from the 27th of July to the 2nd August and from the 4th to the 6th due to instrumental problems. NO₂ mixing ratios were corrected for its formation (R1) during transport from the roof top inlet to the cavity (≈ 7.5 s).

Two commercially available instruments operated permanently at the site also provided measurements of NO₂ and NO. These were a cavity-phase-shift (CAPS) instrument for NO₂ measurement and a chemiluminescence device (CLD) for NO₂ and NO. The CAPS (Aerodyne, Ambient Monitor Version 2012) had a detection limit of 270 pptv (3 σ in 1 min integration time) and an uncertainty of 10% (1 min integration time). The CLD (ECO PHYSICS, Model AL 770 pptv) uses chemiluminescence from the reaction of NO with ozone in combination with a blue-light converter to convert NO₂ to NO.



The instrument was routinely calibrated once a week ($10 \text{ ppmv} \pm 5\%$ NO in N_2 , Riessner, Germany. Deviations between two calibrations are typically well below 3%. Detection limits during the intensive were 11 pptv for NO, 16 pptv for NO_2 (3σ in 1 min integration time) and median uncertainties are 27 pptv (7%) for NO and 70 pptv (10%) for NO_2 (2σ at 1 ppbv in 1 min integration time). Corrections were applied to take into account NO loss and NO_2 formation due to further reactions involving ozone in the inlet tubing.

A comparison of the three NO_2 measurements instruments is given in Fig. S1 of the supplementary information which plots the NO_2 mixing ratios (averaged over 60 s) of the CLD and CAPS instruments versus the CRDS. A least-squares fit (considering errors in both parameters) to the plot of NO_2 (CLD) versus NO_2 (CRDS) has a slope of 0.94 ± 0.25 and an intercept of 0.00 ± 0.04 . For the plot of NO_2 (CAPS) versus NO_2 (CRDS) comparison we derive a slope of 0.95 ± 0.02 and an intercept of 0.00 ± 0.02 . Within combined uncertainty, the NO_2 measurements are thus in agreement. The NO_2 mixing ratios used as input to calculate the NO_3 -reactivity were taken from the CRDS instrument, with data gaps filled by CAPS measurements.

Ozone was monitored with a UV absorption instrument (Thermo Environmental Instruments Inc., Model TECO 49C) which is calibrated at regular intervals with a transfer standard (TECO 49 PS). The uncertainty in the ozone mixing ratio is 1.2 ppbv or 2% (2σ in 1 hour).

2.3 NO_3 measurements

For the measurement of ambient NO_3 , a 10 m length of PFA-tubing (3/8-inch outer diameter) was installed on the top of the building, circa 10 cm from the VOC inlet. A $2 \mu\text{m}$ pore PTFE filter (47 mm in diameter, replaced every hour) in a PFA filter holder was located at the end of the inlet. The tubing was connected to a bypass pump operated at $20 \text{ dm}^3 (\text{STD}) \text{ min}^{-1}$ to reduce the residence time. The sample flow through the cavity was increased to $8 \text{ dm}^3 (\text{STD}) \text{ min}^{-1}$ to reduce the NO_3 residence time within the cavity. NO_3 mixing ratios were recorded every 6 s (3600 ring-downs co-added) with zeroing by titration (no addition) every 15 data points. The NO_3 transmission through the inlet ($67 \pm 15\%$), filter and filter-holder ($84 \pm 10\%$) and cavity ($88 \pm 10\%$) were established post-campaign and used to correct the data. The overall uncertainty in the NO_3 measurements, including uncertainty in the absorption cross-section, was circa 35%.

2.4 OH-reactivity measurements

OH-reactivity measurements were conducted using a chemical ionization mass spectrometer (CIMS) in which OH radicals (generated by photolysis of H_2O at 184.95 nm) are converted to H_2SO_4 (Berresheim et al., 2000; Schlosser et al., 2009). For the derivation of OH-reactivity ($k_{\text{total}}^{\text{OH}}$), relative OH radical concentrations are measured at two fixed reaction times and a decay constant is derived assuming exponential behaviour. After correction for wall losses, as well as NO-induced HOx



recycling in the sample tube, ambient reactivities between 1 to 40 s⁻¹ are measureable. OH-reactivity measurements were made every 20 min throughout the measurement period. Measurements were discontinued during periods of precipitation and when the pinhole to the mass spectrometer vacuum system was blocked by insects. The instrument performs best if NO mixing ratios are below 4 ppbv and reactivities do not exceed 15 s⁻¹; for the measurements reported here, the mean
 5 uncertainty in the OH-reactivity was ± 1.2 s⁻¹ (or 46%). OH-reactivity calibration was carried out before and after the measurement period, and the calibration factor was applied to the whole dataset. The determination of the OH wall loss rate from zero reactivity measurement (null measurement) using synthetic air cylinders was not reliable and therefore the zero was estimated using nighttime measurements when sampling from above the boundary layer. Details are given in the supplementary information.

10 2.5 VOC measurements

A gas-chromatograph (GC-MS/FID model AGILENT 6890 with 5975 B inert XL MSD), was used for the detection of C₅–C₁₃ NMHCs and BVOC (Hoerger et al., 2015). In a custom-made pre-concentration unit, air was sampled at 30 °C on a 3-bed adsorption trap and, after a cryo-focussing step, injected onto the GC column (50m BPX-5). Subsequently, signals were detected with a mass spectrometer (MS) running in parallel with a flame ionization detector (FID). The instrument measured
 15 i.a. isoprene and a wide variety of monoterpenes with uncertainties (2σ) from 6 to 100% depending on the compound. For the detection of light NMHCs (C₂–C₈), a GC-FID system (GC-1, Varian 3600 CX, FID detector) described in detail by Plass-Dülmer et al. (2002) was used. In both systems, an ozone scrubber (impregnated filter with Na₂SO₃) was used and water was removed from the sample air either by hydrophobic adsorbents (C₅–C₁₃) or a cold trap (C₂–C₈) prior to the pre-concentration step.

20 VOCs were sampled every hour for 15 min (C₂–C₈) respectively 20 min (C₅–C₁₃) by both instruments. During the rainy period from the 24th of July 12:00 UTC to the 27th of July 12:00 UTC, VOCs were only measured twice daily.

3 Results and discussion

Figure 1 displays the time series of NO₃-reactivity ($k_{\text{OTG}}^{\text{NO}_3}$) along with related trace gases and meteorological data obtained during the intensive. Sunrise was around 3:50 UTC and sunset at ≈19:00 UTC. Two mild days ($T_{\text{max}} = 20\text{--}25$ °C) at the
 25 beginning of the campaign were followed by a 3 day period with heavy rains and maximum temperatures of around 10 °C followed by a warm period with temperatures up to 30 °C and occasional thunderstorms. The predominant wind direction was west-south-west with only minor contributions from other directions (Fig. S2). Wind speeds were generally around 2.5 to 7.5 ms⁻¹, increasing up to 15 ms⁻¹ during the rainy periods. The highest values of $k_{\text{OTG}}^{\text{NO}_3}$ were detected with north-easterly winds (Fig. S2) coincident with the warmest days of the campaign and the highest biogenic emissions (see section 4.2).
 30 Ozone mixing ratios were strongly correlated with temperature and ranged from 85 ppbv at the 1th of August during the warm, photochemically intense period to less than 20 ppbv during the cool, rainy period between 23rd and 28th of August.



NO_x levels ($\text{NO}_x = \text{NO} + \text{NO}_2$) during the intensive were generally between about 0.5 and 4 ppbv. The mixing ratios of NO, a trace gas which can potentially impact NO_3 lifetimes, were generally below the detection limit (≈ 12 pptv) during most of the nights, increasing to maximum values of < 1 ppbv during the day. Occasional maxima of more than 1 ppbv NO were observed due to local traffic.

5 3.1 NO_3 reactivity

NO_3 -reactivity was measured continuously during a three week intensive (20.07.2017 to 06.08.2017) with the exception of one night (2nd - 3rd August) when, using the same instrument, NO_3 mixing ratios were measured instead. The full data set of $k_{\text{OTG}}^{\text{NO}_3}$ is reproduced in Fig. S3 of the supplementary information together with the corresponding 95% uncertainty limits, which take into account drifts in the zero signal, the stability of the NO_3 source, uncertainty in the dilution factors, uncertainty of the NO and NO_2 mixing ratios as well as the corresponding rate constants.

As described above, during daytime the short NO_3 lifetime normally results in levels that are under the detection limit of most instruments, precluding estimation of the NO_3 -reactivity via stationary-state calculations based on its mixing ratio and production rate. In contrast, our direct measurement enables us to derive the NO_3 -reactivity over the full diel cycle. During the intensive, the 10-minute averaged values of $k_{\text{OTG}}^{\text{NO}_3}$ obtained ranged from below the detection limit ($< 0.005 \text{ s}^{-1}$) to values as high as 0.3 s^{-1} . Campaign averaged values were low ($\approx 0.01 \text{ s}^{-1}$) during nighttime but a factor of ten larger $\approx 0.1 \text{ s}^{-1}$ at 14:00 UTC (local 16:00) and more variable during daytime.

This observation is in stark contrast to the high relative night-time/daytime NO_3 -reactivities we observed in a Boreal forest (Liebmann et al., 2018) and is related to very different meteorological conditions at the two sites. In the boreal forest, the canopy-level NO_3 -reactivity was controlled by the rate of emission of biogenic VOCs into a nocturnal boundary layer of varying height and stability. The elevated location of the Hohenpeissenberg observatory, located at a mountain-top above the surrounding countryside favoured sampling from the residual layer/free troposphere at nighttime. In the absence of turbulent exchange, the residual layer/free troposphere may become disconnected from the planetary boundary layer (PBL) and thus from ground-level emissions of reactive trace gases and may thus contain low levels of biogenic trace gases as well as low(er) levels of NO_2 and higher levels of ozone (Aliwell and Jones, 1998; Allan et al., 2002; von Friedeburg et al., 2002; Stutz et al., 2004; Brown et al., 2007a; Brown et al., 2007b; Brown and Stutz, 2012). NO_3 lifetimes as long as 1 hour (using stationary-state analyses) have been reported for mountain sites when sampling air from above the nocturnal boundary layer (Brown et al., 2016; Sobanski et al., 2016).

During the Hohenpeissenberg intensive two distinct air-mass types were encountered at night, whereby values of $k_{\text{OTG}}^{\text{NO}_3}$ were either at (or below) the detection limit or well above it (named type 1 and type 2, respectively). Figure 2 displays a time series of $k_{\text{OTG}}^{\text{NO}_3}$ over a single night (29th -30th July) in which a switch from type 2 to type 1 was observed. From early evening until shortly before 12:00 UTC, NO_3 -reactivity was variable with values between circa 0.02 and 0.03 s^{-1} . A sharp reduction in $k_{\text{OTG}}^{\text{NO}_3}$ was then observed with values close to the detection limit until sunrise (04:00 UTC). The reduction in $k_{\text{OTG}}^{\text{NO}_3}$ was



accompanied by a drop in relative humidity (from 70% to 60%) and an increase in O_3 (46 to 52 ppbv), both clear indicators of sampling from the residual layer. At the same time, the wind speed increased and the temperature became more variable, indicating that the site was close to the inversion level. At first sunlight, turbulent mixing resulted in gradual connection of the boundary layer and overlying residual layer, leading to an increase in $k_{OTG}^{NO_3}$. Upslope winds caused by heating of the mountainside may also have enhanced transport of air masses with high reactivity to the measurement site. Median, diel profiles of NO_3 -reactivity on type 1 (altogether five) and type 2 nights (altogether 10) are shown in Fig. 3. With the exception of the very low reactivity during type1 nights, type 1 and type 2 have similar diel shapes and similar maximum reactivities.

3.2 NO_3 -reactivity calculated from VOC measurements

- In this section we assess the contribution of various VOCs to the observed NO_3 -reactivity. The most abundant BVOCs were isoprene, sabinene, α -pinene and β -pinene with maximum mixing ratios during the warm period around the 1st August. The time series of BVOC mixing ratios are displayed in Fig. S4 of the supplementary information. $k_{OTG}^{NO_3}$ is a total loss rate constant for chemical reactions of $[NO_3]$ with all organic trace gases present and can be compared to the summed loss rate constant ($k_{VOC}^{NO_3}$) (also in units of s^{-1}) obtained from the concentrations of individual VOCs in the same air mass, $[C_i]$, and the rate coefficient (k_i):

$$k_{VOC}^{NO_3} = \sum k_i^{NO_3} [C_i] \quad (1)$$

Where $[C_i]$ is the measured BVOC concentration and k_i the corresponding rate constant. Individual values of $k_{VOC}^{NO_3}$, calculated using rate constants from the IUPAC evaluation (IUPAC, 2017) or elsewhere in the literature (Shorees et al., 1991), are plotted with interpolated 20 min averages of $k_{OTG}^{NO_3}$ as a time series in the upper panel of Fig. 4.

- The data are also displayed as a pie chart in the lower panel of Fig. 4 in which the contribution of individual biogenic trace-gases to the NO_3 -reactivity are listed. Of the terpenoids, α -pinene contributed most to the overall NO_3 -reactivity ($\approx 16\%$) followed by sabinene ($\approx 12\%$) with other individual BVOCs contributing less than 10%. VOCs such as methanol, acetaldehyde, ethanol, acetone, methylethylketone, alkanes and aromatics were also measured but not included in calculations of $k_{VOC}^{NO_3}$ as their summed contribution reached max. $1.5 \times 10^{-4} s^{-1}$ and was on average $5 \times 10^{-5} s^{-1}$. As $k_{VOC}^{NO_3}$ and $k_{OTG}^{NO_3}$ show a similar dependence on wind direction (Fig. S2) and because only BVOCs were used for the derivation of $k_{VOC}^{NO_3}$, we conclude that the high NO_3 reactivities measured in air masses arriving from the east and northeast are mostly from trace-gases of biogenic origin.

Type1 nights were characterized by very low BVOC mixing ratios, sometimes below the detection limit, whereas isoprene was still present. This observation is consistent with a long lifetime for isoprene in the residual layer at night (Brown et al., 2007a) as the OH concentration is too low and the NO_3 reaction too slow to remove it efficiently. Under conditions of very low NO_3 -reactivity, the fractional contribution of isoprene to the overall reactivity could increase to $\approx 100\%$ (from typically



20% during the day). During type two nights (those with non-zero NO_3 -reactivity) isoprene and monoterpenes were always detected and monoterpenes were the dominant reaction partners for NO_3 .

The difference between $k_{\text{OTG}}^{\text{NO}_3}$ and $k_{\text{VOC}}^{\text{NO}_3}$ (i.e. the NO_3 -reactivity not accounted for by measured VOCs) may be defined as “missing” reactivity (s^{-1}):

$$5 \quad \text{missing } \text{NO}_3\text{-reactivity} = k_{\text{OTG}}^{\text{NO}_3} - k_{\text{VOC}}^{\text{NO}_3} \quad (2)$$

A plot of $k_{\text{VOC}}^{\text{NO}_3}$ versus $k_{\text{OTG}}^{\text{NO}_3}$ (see Fig. S5 of the supplementary information) has a slope of 1.55 ± 0.15 and an intercept of 0.005. This implies, on average, a missing reactivity of $\approx 34\%$ when $k_{\text{OTG}}^{\text{NO}_3} = 0.3 \text{ s}^{-1}$ and a missing reactivity of $\approx 50\%$ when $k_{\text{OTG}}^{\text{NO}_3} = 0.03 \text{ s}^{-1}$. However, both $k_{\text{OTG}}^{\text{NO}_3}$ and $k_{\text{VOC}}^{\text{NO}_3}$ are associated with some uncertainty, which needs to be rigorously assessed to test whether the missing reactivity is significant. To do this we propagated uncertainty in each of the terms $\sum k_i^{\text{NO}_3} [\text{C}_i]$ (mainly related to VOC measurements and assuming 15% uncertainty in the rate coefficients) and derived mean diel profiles of $k_{\text{OTG}}^{\text{NO}_3}$ and $k_{\text{VOC}}^{\text{NO}_3}$ for the whole campaign (hour averages). The results are shown in Fig. 5, where the red shaded area represents total uncertainty and variability in $k_{\text{VOC}}^{\text{NO}_3}$ and the black error bars are the total uncertainty in $k_{\text{OTG}}^{\text{NO}_3}$. Clearly, within combined uncertainty the data overlap, so that missing reactivity is not significantly distinct from zero.

15 3.3 NO_3 measurements and comparison with stationary state calculations

Although rough estimates of NO_3 concentrations at the Hohenpeissenberg have been made (Handisides et al., 2003; Bartenbach et al., 2007), no direct NO_3 measurement had been previously made. For this reason, on just one night during the intensive (2nd - 3rd of August), the instrument was modified to enable measurement of ambient NO_3 mixing ratios rather than NO_3 -reactivity. The NO_3 , O_3 and NO_2 mixing ratios and meteorological data are plotted in Fig. 6.

20 NO_3 mixing ratios slowly increased in the first half of the night, reaching a maximum of 13 pptv around 21:40 UTC. At this time the O_3 mixing ratios were also largest and highly variable. After $\approx 22:30$, O_3 was slowly removed, the NO_3 decreased by a factor of 10 or more indicating that we were sampling more reactive, boundary layer air. This is also evident in the increase in relative humidity and decrease in the temperature until about 01:30.

Given sufficient time, stationary state can be reached for NO_3 at night in which the production and loss terms are approximately balanced (Brown et al., 2003a; Crowley et al., 2010; Crowley et al., 2011). In this case NO_3 mixing ratios can be described by the ratio of their production rate and loss rate (Eq. 3).

$$[\text{NO}_3]_{\text{ss}} = \frac{\text{NO}_3 \text{ production rate}}{\text{NO}_3 \text{ loss rate}} \quad (3)$$

The production rate is governed by the $[\text{NO}_2]$ and $[\text{O}_3]$ mixing ratio and the corresponding rate constant k_2 . If the loss processes are due to reaction with VOCs only, this expression becomes:

$$30 \quad [\text{NO}_3]_{\text{ss}} = \frac{[\text{O}_3][\text{NO}_2]k_2}{k_{\text{VOC}}^{\text{NO}_3}} \quad (4)$$



During this night $k_{\text{OTG}}^{\text{NO}_3}$ was not measured so $k_{\text{VOC}}^{\text{NO}_3}$ was used to account for NO_3 losses.

Figure 7 shows the measured NO_3 mixing ratios (black) compared to those derived from Eq. (4) using the measured VOC concentrations (red curve). Clearly, the predicted, stationary-state NO_3 concentrations are too high (by a factor of up to 3-4), implying that other NO_3 loss processes must be considered. As the directly measured reactivity $k_{\text{OTG}}^{\text{NO}_3}$ agrees rather well with that derived from VOC measurements ($k_{\text{VOC}}^{\text{NO}_3}$) calculated on other campaign nights, it would seem unlikely that unmeasured VOCs contribute sufficiently to NO_3 losses to explain this large factor. Stationary-state concentrations of NO_3 are influenced not only by VOCs but also by NO (if present at night) and also indirectly via heterogeneous loss of N_2O_5 . Equation (5) can be extended to include these processes (Martinez et al., 2000; Geyer et al., 2001; Brown et al., 2003a; Brown et al., 2003b; Brown et al., 2009; Crowley et al., 2010; Sobanski et al., 2016).

$$10 \quad [\text{NO}_3]_{\text{ss}} = \frac{[\text{O}_3][\text{NO}_2]k_1}{k_{\text{VOC}}^{\text{NO}_3} + k_2[\text{NO}] + K_5[\text{NO}_2]f_{\text{het}}} \quad (5)$$

where K_5 is the equilibrium constant for the forward and reverse reactions (R4, R5). The loss frequency due to heterogeneous uptake of N_2O_5 to particles (f_{het}) can be calculated by equation 6:

$$f_{\text{het}} \approx \frac{\gamma \bar{c} A}{4} \quad (6)$$

which is approximately valid if the particles are less than $\approx 1 \mu\text{m}$ in diameter. In this expression, A is the aerosol surface area density ($\text{cm}^2 \text{cm}^{-3}$), \bar{c} is the mean, molecular velocity of N_2O_5 (26233 cm s^{-1} at 298 K) and γ is the dimensionless uptake coefficient. If we assume a large value for the uptake coefficient of 0.03 as characteristic for aerosol with low organic content (Bertram and Thornton, 2009; Bertram et al., 2009; Crowley et al., 2011; Phillips et al., 2016) and use the aerosol surface of $1.25\text{--}1.55 \times 10^{-6} \text{ cm}^2 \text{cm}^{-3}$ (measured by a scanning mobility particle sizer for 10 – 890 nm), we obtain values for f_{het} of $2.4\text{--}2.9 \times 10^{-4} \text{ s}^{-1}$. In this case an unrealistic value of $\gamma = 0.5$ would be required to lower the calculated, stationary-state NO_3 mixing ratio to between 3 and 10 pptv as observed.

Clearly, heterogeneous losses of N_2O_5 do not account for the missing NO_3 sinks during this night and we now consider the role of NO. As NO mixing ratios in this night did not exceed the detection limit (11 pptv) we used a constant value of 5 pptv to approximately align the calculated NO_3 mixing ratio with that measured for much of the night. Clearly, the calculation of NO_3 -reactivity from stationary-state calculations can be precarious and subject to large cumulative uncertainty from e.g. measurement uncertainty in NO_3 mixing ratios, uptake coefficients, aerosol surface area and NO mixing ratios close to instrumental detection limits.

To assess the NO_3 mixing ratios for the rest of the intensive, equation 6 can be augmented by adding the loss rate constant for NO_3 -photolysis J_{NO_3} and substituting $k_{\text{VOC}}^{\text{NO}_3}$ for $k_{\text{OTG}}^{\text{NO}_3}$.

$$[\text{NO}_3]_{\text{ss}} = \frac{[\text{O}_3][\text{NO}_2]k_1}{k_{\text{OTG}}^{\text{NO}_3} + k_2[\text{NO}] + J_{\text{NO}_3} + K_5[\text{NO}_2]f_{\text{het}}} \quad (7)$$

In the absence of a direct measurement of J_{NO_3} , the diel cycle of the relative NO_3 -to- NO_2 photolysis rate constant ($J_{\text{NO}_3}/J_{\text{NO}_2}$) was calculated using the TUV (tropospheric ultraviolet and visible radiation) model



(<https://www2.acom.ucar.edu/modeling/tropospheric-ultraviolet-and-visible-tuv-radiation-model>) and then put on an absolute basis using measured J_{NO_2} values.

Figure 8 shows the NO_3 production rate (lower panel, black curve) and total loss rate (lower panel, red curve) as well as the stationary-state NO_3 mixing ratios for the entire intensive period (Fig. 8, upper panel black curve). During nights in which the reactivity fell below the detection limit of the instrument $k_{\text{OTG}}^{\text{NO}_3}$ was set to 0.005 s^{-1} . Calculated NO_3 mixing ratios were in the sub-pptv range during daytime and around 1-15 pptv during nighttime. The NO_3 mixing ratios thus derived are comparable to those measured on a single night (Fig. 8, red curve) and are broadly consistent with previous estimates for this site (Handisides et al., 2003; Bartenbach et al., 2007).

10 3.4 Contribution to NO_x loss

At nighttime, in the absence of NO and sunlight, each NO_3 radical formed in the reaction of NO_2 with O_3 will either be removed indirectly via the uptake of N_2O_5 onto particles or will react with a biogenic hydrocarbon. The latter results in the formation of an organic nitrate at a yield of between 20 and 100%, depending on the specific VOC (Ng et al., 2017). The large daytime values for $k_{\text{OTG}}^{\text{NO}_3}$ obtained in this study suggest that even during sunlight hours (when NO_3 is generally considered to be of little significance owing to its rapid photolysis) significant amounts of NO_3 form organic nitrates rather than reforming NO_2 by reaction with NO, or photolysis.

The fraction, f , of NO_3 that will react with organic trace gases is given by:

$$f = \frac{k_{\text{OTG}}^{\text{NO}_3}}{([k_{\text{OTG}}^{\text{NO}_3}] + [J_{\text{NO}_3}] + [\text{NO}]k_3 + K_5 [\text{NO}_2]f_{\text{het}})} \quad (8)$$

where the denominator sums all loss processes for NO_3 . Figure 9 illustrates this via a diel cycle of the median for f . At nighttime, $\approx 99\%$ of the NO_3 will be lost to reaction with BVOCs, with indirect heterogeneous losses representing the remaining 1%. During daytime, at the peak of the actinic flux ($\max J_{\text{NO}_3} \approx 0.2 \text{ s}^{-1}$) and correspondingly high levels of NO ($k_{\text{NO}} = 0.1 - 0.2 \text{ s}^{-1}$), 20% of the formed NO_3 was lost due to reaction with organic trace gases, increasing up to 40% in the late afternoon. This result is comparable with reactivity measurements in a boreal forest in Finland during IBARN 2016 where a very similar diel profile for f was determined (Liebmann et al., 2018). The NO_3 -reactivity data from these measurements indicate that the role of NO_3 as a daytime oxidant of biogenic VOCs in forested regions may so far have been underestimated, which in turn has implications for understanding the diel cycle of organic nitrate and secondary organic aerosol formation in such environments.

4 Comparison of NO_3 and OH-reactivity

As mentioned above NO_3 radicals and OH radicals react with atmospheric trace gases via different mechanisms, resulting in profoundly different rate coefficients and thus reactivities. By combining the continuous, on-site measurements of the OH-



reactivity with the NO_3 -reactivity measurements during the intensive period, we were able to generate the first dataset of simultaneous, direct measurement of both OH-reactivity, $k_{\text{total}}^{\text{OH}}$, and NO_3 -reactivity at any location.

To aid comparison, we subtracted the contributions of several inorganic and organic trace gases (NO , NO_2 , SO_2 , CO , CH_4) that are not included in $k_{\text{OTG}}^{\text{NO}_3}$ or do not react to a significant extent with NO_3 from the total OH-reactivity and thus derived

5 $k_{\text{OTG}}^{\text{OH}}$.

$$k_{\text{OTG}}^{\text{OH}} = k_{\text{total}}^{\text{OH}} - k_{\text{NO}}^{\text{OH}}[\text{NO}] - k_{\text{NO}_2}^{\text{OH}}[\text{NO}_2] - k_{\text{SO}_2}^{\text{OH}}[\text{SO}_2] - k_{\text{CH}_4}^{\text{OH}}[\text{CH}_4] - k_{\text{CO}}^{\text{OH}}[\text{CO}] \quad (9)$$

Fig. 10 depicts the time series of $k_{\text{Total}}^{\text{OH}}$, $k_{\text{OTG}}^{\text{OH}}$ and $k_{\text{OTG}}^{\text{NO}_3}$. All display maximum values close to midday, though $k_{\text{OTG}}^{\text{OH}}$ averaged over the intensive are larger by a factor of 44 larger than $k_{\text{OTG}}^{\text{NO}_3}$, reflecting generally larger rate coefficients for OH.

The blue shaded areas for $k_{\text{Total}}^{\text{OH}}$ represents the 1σ uncertainty of the measurements. Total uncertainty in $k_{\text{OTG}}^{\text{OH}}$ and $k_{\text{OTG}}^{\text{NO}_3}$ is not shown to preserve clarity of presentation. The time series of the $k_{\text{OTG}}^{\text{OH}}$ can be found in the supplementary information (Fig. S6).

The measured reactivities of both radicals show a clear diel profile, with higher daytime and lower nighttime values. Figure 11 shows a correlation plot of OH and NO_3 -reactivity divided into day (red) and nighttime (black) data. During the day, the data are highly scattered, which can be understood when one considers the highly variable organic content of the air masses being sampled. To illustrate this we have drawn the expected correlation lines (based on the known, relative rate coefficients) for single component organic trace gases including isoprene and terpenes. The expected slopes for these individual VOCs are very different and encompass the full scatter in the observations, which is the result of changing atmospheric composition (i.e. the mix of reactive organic species) owing to changes in air mass age and source region (wind direction) during the campaign. The extremes are represented by α -terpinene (which favours NO_3) and CH_4 (which favours OH).

During nighttime (black points) the plot of $k_{\text{OTG}}^{\text{OH}}$ versus $k_{\text{OTG}}^{\text{NO}_3}$ is less scattered, indicating that the air masses sampled (often from the residual layer) are chemically less complex and variable. The data lay close to the line that marks the expected correlation if isoprene were the dominant sink of both NO_3 and OH at nighttime once molecules such as CO and CH_4 have been removed from the term describing OH-reactivity. This is in broad agreement with our observation that isoprene is the main sink of NO_3 during nights when the measurement site was decoupled from direct boundary layer emissions.

5. Summary and Conclusion

Direct measurements of the NO_3 -reactivity towards organic trace gases, $k_{\text{OTG}}^{\text{NO}_3}$ were conducted at the top of the Hohenpeissenberg mountain (988 m a.s.l.) during an intensive measurement campaign in the summer of 2017. NO_3 -reactivities had a distinct diel profile with values as large as 0.3 s^{-1} during daytime but close to or below the detection limit of the instrument during nighttime when the measurement site was frequently in the residual layer / free troposphere. Within experimental uncertainty, the high daytime NO_3 -reactivity was accounted for by BVOCs that were measured at the site, and



was dominated by monoterpenes especially α -pinene and sabinene. On average, the reaction with VOCs accounted for \approx 99% of the loss of NO_3 during nighttime and an average of 20% at noon, increasing to 30-50% during early morning and late evening. The reaction of NO_3 with BVOCS therefore represents a significant NO_x loss not only during the night but also during daytime and implies significant formation of organic nitrates via NO_3 reactions throughout the diel cycle. Stationary-
5 state, daytime and nighttime NO_3 mixing ratios were calculated using the production term and $k_{\text{OTG}}^{\text{NO}_3}$ and were broadly consistent with direct measurement made on one night. A comparison between directly measured OH- and NO_3 reactivities was performed, indicating a weak correlation during the day when chemically reactive, complex and variable air masses were encountered. A tighter correlation, consistent with isoprene dominating the (low) NO_3 -reactivities was observed at night.

10
Acknowledgements: We would like to thank the DWD for hosting and supporting this measurement campaign. We would like to thank the DWD personnel for the data evaluation, most notably Katja Michl for NMHCs, Jennifer Englert for OVOCs, Harald Flentje for the SMPS data as well as the great technical support, particularly we are grateful to Erasmus Tensing, Thomas Elste, Georg Stange. We thank Chemours for provision of the FEP sample used to coat the CRD-cavities.



References

- Aliwell, S. R., and Jones, R. L.: Measurements of tropospheric NO₃ at midlatitude, *J. Geophys. Res. -Atmos.*, 103, 5719-5727, 1998.
- 5 Allan, B. J., Plane, J. M. C., Coe, H., and Shillito, J.: Observations of NO₃ concentration profiles in the troposphere, *J. Geophys. Res. -Atmos.*, 107, 4588, doi: 10.1029/2002jd002112, 2002.
- Atkinson, R.: Atmospheric chemistry of VOCs and NO_x, *Atmos. Env.*, 34, 2063-2101, 2000.
- Atkinson, R., and Arey, J.: Atmospheric degradation of volatile organic compounds, *Chem. Rev.*, 103, 4605-4638, 10.1021/cr0206420, 2003a.
- 10 Atkinson, R., and Arey, J.: Gas-phase tropospheric chemistry of biogenic volatile organic compounds: a review, *Atmos. Env.*, 37, S197-S219, 2003b.
- Bannan, T. J., Booth, A. M., Bacak, A., Muller, J. B. A., Leather, K. E., Le Breton, M., Jones, B., Young, D., Coe, H., Allan, J., Visser, S., Slowik, J. G., Furger, M., Prevot, A. S. H., Lee, J., Dunmore, R. E., Hopkins, J. R., Hamilton, J. F., Lewis, A. C., Whalley, L. K., Sharp, T., Stone, D., Heard, D. E., Fleming, Z. L., Leigh, R., Shallcross, D. E., and Percival, C. J.: The
15 first UK measurements of nitryl chloride using a chemical ionization mass spectrometer in central London in the summer of 2012, and an investigation of the role of Cl atom oxidation, *J. Geophys. Res. -Atmos.*, 120, 5638-5657, 10.1002/2014jd022629, 2015.
- Bartenbach, S., Williams, J., Plass-Dülmer, C., Berresheim, H., and Lelieveld, J.: In-situ measurement of reactive hydrocarbons at Hohenpeissenberg with comprehensive two-dimensional gas chromatography (GC×GC-FID): use in
20 estimating HO and NO₃, *Atmos. Chem. Phys.*, 7, 1-14, 10.5194/acp-7-1-2007, 2007.
- Bastin, J.-F., Berrahmouni, N., Grainger, A., Maniatis, D., Mollicone, D., Moore, R., Patriarca, C., Picard, N., Sparrow, B., Abraham, E. M., Aloui, K., Atesoglu, A., Attore, F., Bassüllü, Ç., Bey, A., Garzuglia, M., García-Montero, L. G., Groot, N., Guerin, G., Laestadius, L., Lowe, A. J., Mamane, B., Marchi, G., Patterson, P., Rezende, M., Ricci, S., Salcedo, I., Diaz, A. S.-P., Stolle, F., Surappaeva, V., and Castro, R.: The extent of forest in dryland biomes, *Science*, 356, 635-638,
25 10.1126/science.aam6527, 2017.
- Berresheim, H., Elste, T., Plass-Dülmer, C., Eiseleb, F. L., and Tannerb, D. J.: Chemical ionization mass spectrometer for long-term measurements of atmospheric OH and H₂SO₄, *Int. J. Mass Spectrom.*, 202, 91-109, [https://doi.org/10.1016/S1387-3806\(00\)00233-5](https://doi.org/10.1016/S1387-3806(00)00233-5), 2000.
- Bertram, T. H., and Thornton, J. A.: Toward a general parameterization of N₂O₅ reactivity on aqueous particles: the
30 competing effects of particle liquid water, nitrate and chloride, *Atmos. Chem. Phys.*, 9, 8351-8363, 2009.
- Bertram, T. H., Thornton, J. A., Riedel, T. P., Middlebrook, A. M., Bahreini, R., Bates, T. S., Quinn, P. K., and Coffman, D. J.: Direct observations of N₂O₅ reactivity on ambient aerosol particles, *Geophys. Res. Lett.*, 36, L19803, doi:10.1029/2009GL040248, 2009.
- Birmili, W., Berresheim, H., Plass-Dülmer, C., Elste, T., Gilge, S., Wiedensohler, A., and Uhrner, U.: The Hohenpeissenberg aerosol formation experiment (HAFEX): a long-term study including size-resolved aerosol, H₂SO₄, OH, and monoterpenes measurements, *Atmos. Chem. Phys.*, 3, 361-376, 10.5194/acp-3-361-2003, 2003.
35
- Brown, S. S., Stark, H., and Ravishankara, A. R.: Applicability of the steady state approximation to the interpretation of atmospheric observations of NO₃ and N₂O₅, *J. Geophys. Res. -Atmos.*, 108, Art. 4539, 10.1029/2003JD003407, 2003a.
- Brown, S. S., Stark, H., Ryerson, T. B., Williams, E. J., Nicks, D. K., Trainer, M., Fehsenfeld, F. C., and Ravishankara, A. R.: Nitrogen oxides in the nocturnal boundary layer: Simultaneous in situ measurements of NO₃, N₂O₅, NO₂, NO, and O₃, *J. Geophys. Res. -Atmos.*, 108, art. 4299, 10.1029/2002JD002917, 2003b.
40



- Brown, S. S., Dube, W. P., Osthoff, H. D., Stutz, J., Ryerson, T. B., Wollny, A. G., Brock, C. A., Warneke, C., De Gouw, J. A., Atlas, E., Neuman, J. A., Holloway, J. S., Lerner, B. M., Williams, E. J., Kuster, W. C., Goldan, P. D., Angevine, W. M., Trainer, M., Fehsenfeld, F. C., and Ravishankara, A. R.: Vertical profiles in NO_3 and N_2O_5 measured from an aircraft: Results from the NOAA P-3 and surface platforms during the New England Air Quality Study 2004, *J. Geophys. Res. - Atmos.*, 112, D22304, doi: 10.1029/2007jd008893, 2007a.
- 5 Brown, S. S., Dube, W. P., Osthoff, H. D., Wolfe, D. E., Angevine, W. M., and Ravishankara, A. R.: High resolution vertical distributions of NO_3 and N_2O_5 through the nocturnal boundary layer, *Atmos. Chem. Phys.*, 7, 139-149, 2007b.
- Brown, S. S., Dube, W. P., Fuchs, H., Ryerson, T. B., Wollny, A. G., Brock, C. A., Bahreini, R., Middlebrook, A. M., Neuman, J. A., Atlas, E., Roberts, J. M., Osthoff, H. D., Trainer, M., Fehsenfeld, F. C., and Ravishankara, A. R.: Reactive uptake coefficients for N_2O_5 determined from aircraft measurements during the Second Texas Air Quality Study: Comparison to current model parameterizations, *J. Geophys. Res. -Atmos.*, 114, art. D00F10, 10.1029/2008JD011679, 2009.
- 10 Brown, S. S., Dube, W. P., Peischl, J., Ryerson, T. B., Atlas, E., Warneke, C., de Gouw, J. A., Hekkert, S. t. L., Brock, C. A., Flocke, F., Trainer, M., Parrish, D. D., Fehsenfeld, F. C., and Ravishankara, A. R.: Budgets for nocturnal VOC oxidation by nitrate radicals aloft during the 2006 Texas Air Quality Study, *J. Geophys. Res. -Atmos.*, 116, 10.1029/2011jd016544, 2011.
- 15 Brown, S. S., and Stutz, J.: Nighttime radical observations and chemistry, *Chem. Soc. Rev.*, 41, 6405–6447, 2012.
- Brown, S. S., Dube, W. P., Tham, Y. J., Zha, Q., Xue, L., Poon, S., Wang, Z., Blake, D. R., Tsui, W., Parrish, D. D., and Wang, T.: Nighttime chemistry at a high altitude site above Hong Kong, *J. Geophys. Res. -Atmos.*, 121, 2457-2475, 10.1002/2015jd024566, 2016.
- 20 Crowley, J. N., Schuster, G., Pouvesle, N., Parchatka, U., Fischer, H., Bonn, B., Bingemer, H., and Lelieveld, J.: Nocturnal nitrogen oxides at a rural mountain site in south-western Germany, *Atmos. Chem. Phys.*, 10, 2795-2812, 2010.
- Crowley, J. N., Thieser, J., Tang, M. J., Schuster, G., Bozem, H., Hasaynali Beygi, Z., Fischer, H., Diesch, J.-M., Drewnick, F., Borrmann, S., Song, W., Yassaa, N., Williams, J., Pöhler, D., Platt, U., and Lelieveld, J.: Variable lifetimes and loss mechanisms for NO_3 and N_2O_5 during the DOMINO campaign: Contrast between marine, urban and continental air, *Atmos. Chem. Phys.*, 11, 10863-10870, 2011.
- 25 Dentener, F. J., and Crutzen, P. J.: Reaction of N_2O_5 on tropospheric aerosols - Impact on the global distributions of NO_x , O_3 , and OH, *J. Geophys. Res. -Atmos.*, 98, 7149-7163, 1993.
- Fry, J. L., Draper, D. C., Barsanti, K. C., Smith, J. N., Ortega, J., Winkle, P. M., Lawler, M. J., Brown, S. S., Edwards, P. M., Cohen, R. C., and Lee, L.: Secondary Organic Aerosol Formation and Organic Nitrate Yield from NO_3 Oxidation of Biogenic Hydrocarbons, *Env. Sci. Tech.*, 48, 11944-11953, 10.1021/es502204x, 2014.
- 30 Geyer, A., Alicke, B., Konrad, S., Schmitz, T., Stutz, J., and Platt, U.: Chemistry and oxidation capacity of the nitrate radical in the continental boundary layer near Berlin, *J. Geophys. Res. -Atmos.*, 106, 8013-8025, 2001.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geoscientific Model Development*, 5, 1471-1492, 10.5194/gmd-5-1471-2012, 2012.
- 35 Hakola, H., Tarvainen, V., Laurila, T., Hiltunen, V., Hellén, H., and Keronen, P.: Seasonal variation of VOC concentrations above a boreal coniferous forest, *Atmos. Env.*, 37, 1623-1634, [http://dx.doi.org/10.1016/S1352-2310\(03\)00014-1](http://dx.doi.org/10.1016/S1352-2310(03)00014-1), 2003.
- Hakola, H., Hellén, H., Hemmilä, M., Rinne, J., and Kulmala, M.: In situ measurements of volatile organic compounds in a boreal forest, *Atmos. Chem. Phys.*, 12, 11665-11678, 10.5194/acp-12-11665-2012, 2012.
- 40 Handisides, G. M., Plass-Dulmer, C., Gilge, S., Bingemer, H., and Berresheim, H.: Hohenpeissenberg Photochemical Experiment (HOPE 2000): Measurements and photostationary state calculations of OH and peroxy radicals, *Atmos. Chem. Phys.*, 3, 1565-1588, 2003.



- Hock, N., Schneider, J., Borrmann, S., Römpf, A., Moortgat, G., Franze, T., Schauer, C., Pöschl, U., Plass-Dülmer, C., and Berresheim, H.: Rural continental aerosol properties and processes observed during the Hohenpeissenberg Aerosol Characterization Experiment (HAZE2002), *Atmos. Chem. Phys.*, 8, 603-623, 10.5194/acp-8-603-2008, 2008.
- 5 Hoerger, C. C., Claude, A., Plass-Duelmer, C., Reimann, S., Eckart, E., Steinbrecher, R., Aalto, J., Arduini, J., Bonnaire, N., Cape, J. N., Colomb, A., Connolly, R., Diskova, J., Dumitrescu, P., Ehlers, C., Gros, V., Hakola, H., Hill, M., Hopkins, J. R., Jäger, J., Junek, R., Kajos, M. K., Klemp, D., Leuchner, M., Lewis, A. C., Locoge, N., Maione, M., Martin, D., Michl, K., Nemitz, E., O'Doherty, S., Pérez Ballesta, P., Ruuskanen, T. M., Sauvage, S., Schmidbauer, N., Spain, T. G., Straube, E., Vana, M., Vollmer, M. K., Wegener, R., and Wenger, A.: ACTRIS non-methane hydrocarbon intercomparison experiment in Europe to support WMO GAW and EMEP observation networks, *Atmos. Meas. Tech.*, 8, 2715-2736, 10.5194/amt-8-2715-2015, 2015.
- 10 Holzke, C., Hoffmann, T., Jaeger, L., Koppmann, R., and Zimmer, W.: Diurnal and seasonal variation of monoterpene and sesquiterpene emissions from Scots pine (*Pinus sylvestris* L.), *Atmos. Env.*, 40, 3174-3185, <http://dx.doi.org/10.1016/j.atmosenv.2006.01.039>, 2006.
- IUPAC: Task Group on Atmospheric Chemical Kinetic Data Evaluation, (Ammann, M., Cox, R.A., Crowley, J.N., Herrmann, H., Jenkin, M.E., McNeill, V.F., Mellouki, A., Rossi, M. J., Troe, J. and Wallington, T. J.) <http://iupac.pole-ether.fr/index.html>, 2017.
- 15 Jaoui, M., Kleindienst, T. E., Docherty, K. S., Lewandowski, M., and Offenberg, J. H.: Secondary organic aerosol formation from the oxidation of a series of sesquiterpenes: alpha-cedrene, beta-caryophyllene, alpha-humulene and alpha-farnesene with O₃, OH and NO₃ radicals, *Environmental Chemistry*, 10, 178-193, 10.1071/en13025, 2013.
- 20 Lee, A. K. Y., Abbatt, J. P. D., Leaitch, W. R., Li, S.-M., Sjostedt, S. J., Wentzell, J. J. B., Liggio, J., and Macdonald, A. M.: Substantial secondary organic aerosol formation in a coniferous forest: observations of both day- and nighttime chemistry, *Atmos. Chem. Phys.*, 16, 6721-6733, 10.5194/acp-16-6721-2016, 2016.
- Lelieveld, J., Dentener, F. J., Peters, W., and Krol, M. C.: On the role of hydroxyl radicals in the self-cleansing capacity of the troposphere, *Atmos. Chem. Phys.*, 4, 2337-2344, 2004.
- 25 Lelieveld, J., Butler, T. M., Crowley, J. N., Dillon, T. J., Fischer, H., Ganzeveld, L., Harder, H., Lawrence, M. G., Martinez, M., Taraborrelli, D., and Williams, J.: Atmospheric oxidation capacity sustained by a tropical forest, *Nature*, 452, 737-740, 2008.
- Lelieveld, J., Gromov, S., Pozzer, A., and Taraborrelli, D.: Global tropospheric hydroxyl distribution, budget and reactivity, *Atmos. Chem. Phys.*, 16, 12477-12493, 10.5194/acp-16-12477-2016, 2016.
- 30 Liebmann, J., Karu, E., Sobanski, N., Schuladen, J., Ehn, M., Schallhart, S., Quéléver, L., Hellen, H., Hakola, H., Hoffmann, T., Williams, J., Fischer, H., Lelieveld, J., and Crowley, J. N.: Direct measurement of NO₃ radical reactivity in a boreal forest, *Atmos. Chem. Phys.*, 2018, 3799-3815, 10.5194/acp-18-3799-2018, 2018.
- Liebmann, J. M., Schuster, G., Schuladen, J. B., Sobanski, N., Lelieveld, J., and Crowley, J. N.: Measurement of ambient NO₃ reactivity: Design, characterization and first deployment of a new instrument, *Atmos. Meas. Tech.*, 2017, 1241-1258, 10.5194/amt-2016-381, 2017.
- 35 Long, B., Tan, X. F., Long, Z. W., Ren, D. S., and Zhang, W. J.: Theoretical Study on Decomposition of CF₃OH Catalyzed by Water Dimer and Ammonia, *Chinese Journal of Chemical Physics*, 24, 16-21, 2011.
- Mannschreck, K., Gilge, S., Plass-Duelmer, C., Fricke, W., and Berresheim, H.: Assessment of the applicability of NO-NO₂-O₃ photostationary state to long-term measurements at the Hohenpeissenberg GAW Station, Germany, *Atmos. Chem. Phys.*, 4, 1265-1277, 10.5194/acp-4-1265-2004, 2004.
- 40 Martinez, M., Perner, D., Hackenthal, E. M., Kulzer, S., and Schutz, L.: NO₃ at Helgoland during the NORDEX campaign in October 1996, *J. Geophys. Res. -Atmos.*, 105, 22685-22695, 2000.



- Ng, N. L., Brown, S. S., Archibald, A. T., Atlas, E., Cohen, R. C., Crowley, J. N., Day, D. A., Donahue, N. M., Fry, J. L., Fuchs, H., Griffin, R. J., Guzman, M. I., Herrmann, H., Hodzic, A., Iinuma, Y., Jimenez, J. L., Kiendler-Scharr, A., Lee, B. H., Luecken, D. J., Mao, J., McLaren, R., Mutzel, A., Osthoff, H. D., Ouyang, B., Picquet-Varrau, B., Platt, U., Pye, H. O. T., Rudich, Y., Schwantes, R. H., Shiraiwa, M., Stutz, J., Thornton, J. A., Tilgner, A., Williams, B. J., and Zaveri, R. A.: Nitrate radicals and biogenic volatile organic compounds: oxidation, mechanisms, and organic aerosol, *Atmos. Chem. Phys.*, 17, 2103-2162, 10.5194/acp-17-2103-2017, 2017.
- Nölscher, A. C., Williams, J., Sinha, V., Custer, T., Song, W., Johnson, A. M., Axinte, R., Bozem, H., Fischer, H., Pouvesle, N., Phillips, G., Crowley, J. N., Rantala, P., Rinne, J., Kulmala, M., Gonzales, D., Valverde-Canossa, J., Vogel, A., Hoffmann, T., Ouwersloot, H. G., Vilà-Guerau de Arellano, J., and Lelieveld, J.: Summertime total OH reactivity measurements from boreal forest during HUMPPA-COPEC 2010, *Atmos. Chem. Phys.*, 12, 8257-8270, 10.5194/acp-12-8257-2012, 2012.
- Nölscher, A. C., Bourtsoukidis, E., Bonn, B., Kesselmeier, J., Lelieveld, J., and Williams, J.: Seasonal measurements of total OH reactivity emission rates from Norway spruce in 2011, *Biogeosciences*, 10, 4241-4257, 10.5194/bg-10-4241-2013, 2013.
- Novelli, A., Hens, K., Ernest, C., Martinez, M., Nölscher, A., Sinha, V., Paasonen, P., Petäjä, T., Sipilä, M., Elste, T., Plass-Dülmer, C., J. Phillips, G., Kubistin, D., Williams, J., Vereecken, L., Lelieveld, J., and Harder, H.: Estimating the atmospheric concentration of Criegee intermediates and their possible interference in a FAGE-LIF instrument, 7807-7826 pp., 2017.
- Noxon, J. F., Norton, R. B., and Henderson, W. R.: Observation of Atmospheric NO_3 , *Geophys. Res. Lett.*, 5, 675-678, 1978.
- Osthoff, H. D., Roberts, J. M., Ravishankara, A. R., Williams, E. J., Lerner, B. M., Sommariva, R., Bates, T. S., Coffman, D., Quinn, P. K., Dibb, J. E., Stark, H., Burkholder, J. B., Talukdar, R. K., Meagher, J., Fehsenfeld, F. C., and Brown, S. S.: High levels of nitryl chloride in the polluted subtropical marine boundary layer, *Nature Geoscience*, 1, 324-328, 2008.
- Phillips, G. J., Tang, M. J., Thieser, J., Brickwedde, B., Schuster, G., Bohn, B., Lelieveld, J., and Crowley, J. N.: Significant concentrations of nitryl chloride observed in rural continental Europe associated with the influence of sea salt chloride and anthropogenic emissions, *Geophys. Res. Lett.*, 39, L10811, doi:10.1029/2012GL051912, 2012.
- Phillips, G. J., Thieser, J., Tang, M. J., Sobanski, N., Schuster, G., Fachinger, J., Drewnick, F., Borrmann, S., Bingemer, H., Lelieveld, J., and Crowley, J. N.: Estimating N_2O_5 uptake coefficients using ambient measurements of NO_3 , N_2O_5 , ClNO_2 and particle-phase nitrate, *Atmos. Chem. Phys.*, 16, 13231-13249, 10.5194/acp-16-13231-2016, 2016.
- Plass-Dulmer, C., Michl, K., Ruf, R., and Berresheim, H.: C-2-C-8 hydrocarbon measurement and quality control procedures at the Global Atmosphere Watch Observatory Hohenpeissenberg, *J. Chromatogr. A*, 953, 175-197, 2002.
- Schlosser, E., Brauers, T., Dorn, H. P., Fuchs, H., Häsel, R., Hofzumahaus, A., Holland, F., Wahner, A., Kanaya, Y., Kajii, Y., Miyamoto, K., Nishida, S., Watanabe, K., Yoshino, A., Kubistin, D., Martinez, M., Rudolf, M., Harder, H., Berresheim, H., Elste, T., Plass-Dülmer, C., Stange, G., and Schurath, U.: Technical Note: Formal blind intercomparison of OH measurements: results from the international campaign HOxComp, *Atmos. Chem. Phys.*, 9, 7923-7948, 10.5194/acp-9-7923-2009, 2009.
- Schuster, G., Labazan, I., and Crowley, J. N.: A cavity ring down / cavity enhanced absorption device for measurement of ambient NO_3 and N_2O_5 , *Atmos. Meas. Tech.*, 2, 1-13, 2009.
- Shorees, B., Atkinson, R., and Arey, J.: Kinetics of the gas-phase reactions of β -phellandrene with OH and NO_3 radicals and O_3 at 297 ± 2 K, *Int. J. Chem. Kinet.*, 23, 897-906, 10.1002/kin.550231005, 1991.
- Sobanski, N., Tang, M. J., Thieser, J., Schuster, G., Pöhler, D., Fischer, H., Song, W., Sauvage, C., Williams, J., Fachinger, J., Berkes, F., Hoor, P., Platt, U., Lelieveld, J., and Crowley, J. N.: Chemical and meteorological influences on the lifetime of NO_3 at a semi-rural mountain site during PARADE, *Atmos. Chem. Phys.*, 16, 4867-4883, 10.5194/acp-16-4867-2016, 2016.
- Stutz, J., Alicke, B., Ackermann, R., Geyer, A., White, A., and Williams, E.: Vertical profiles of NO_3 , N_2O_5 , O_3 , and NO_x in the nocturnal boundary layer: 1. Observations during the Texas Air Quality Study 2000 *J. Geophys. Res. -Atmos.*, 109, art. D12306, 10.1029/2003JD004209, 2004.



- Thieser, J., Schuster, G., Phillips, G. J., Reiffs, A., Parchatka, U., Pöhler, D., Lelieveld, J., and Crowley, J. N.: A two-channel, thermal dissociation cavity-ringdown spectrometer for the detection of ambient NO_2 , RO_2NO_2 and RONO_2 , Atmos. Meas. Tech., 9, 553-576, 2016.
- Voigt, S., Orphal, J., and Burrows, J. P.: The temperature and pressure dependence of the absorption cross-sections of NO_2 in the 250-800 nm region measured by Fourier-transform spectroscopy, J. Photochem. Photobiol. A-Chem., 149, 1-7, doi:10.1016/s1010-6030(01)00650-5, 2002.
- von Friedeburg, C., Wagner, T., Geyer, A., Kaiser, N., Vogel, B., Vogel, H., and Platt, U.: Derivation of tropospheric NO_3 profiles using off-axis differential optical absorption spectroscopy measurements during sunrise and comparison with simulations, J. Geophys. Res. -Atmos., 107, 10.1029/2001JD000481, 2002.
- 10 Wayne, R. P., Barnes, I., Biggs, P., Burrows, J. P., Canosamas, C. E., Hjorth, J., Lebras, G., Moortgat, G. K., Perner, D., Poulet, G., Restelli, G., and Sidebottom, H.: The Nitrate Radical - Physics, Chemistry, and the Atmosphere, Atmos. Env., 25, 1-203, 1991.




Figure 1: Overview of measurements during the 2017-intensive. The grey shaded area represents nighttime. GR = global radiation, RF = rainfall, RH = relative humidity, T = temperature, WS = wind speed, WD = wind direction.




Figure 2: Upper panel: $k_{\text{OTG}}^{\text{NO}_3}$ (black) and O_3 mixing ratios (red) from the 29th to the 30th of July. From 23:50 UTC until sunrise the measurement site is located in the residual layer / free troposphere. Lower panel: temperature (T), relative humidity (RH) and wind direction (WD) during the same period.




Figure 3: Median, diel profile of the NO_3 -reactivity. Type 1 nights (black line) show values around the detection limit during night, type 2 nights (red line) are above the detection limit.




Figure 4: Upper panel: measured values of $k_{\text{OTG}}^{\text{NO}_3}$ (black) in comparison with the loss rate constant assigned to individual VOCs. The term “other” includes terpinolene, β -phellandrene, α -terpinene, γ -terpinene, α -thujene and camphene. Myrcene and α -phellandrene were also measured but below the detection limit during the whole campaign. The lower panel indicates the campaign averaged contribution of each measured VOC to the NO_3 loss rate as well as reactivity that was not accounted for by measured VOCs (“missing reactivity”).




Figure 5: Median diel profiles of $k_{\text{VOC}}^{\text{NO}_3}$ and $k_{\text{OTG}}^{\text{NO}_3}$. The error bars on the $k_{\text{OTG}}^{\text{NO}_3}$ measurements are total uncertainty, including systematic error and variability. The uncertainty in $k_{\text{VOC}}^{\text{NO}_3}$ (shaded red area) is dominated by uncertainty in the mixing ratios of the VOCs.




Figure 6: NO_3 mixing ratios measured in the night from the 2th – 3rd of August as well as NO_2 and O_3 (which define the NO_3 production rate). T = temperature, RH = relative humidity, WD = wind directions, WS = wind speed.




Figure 7: Comparison of measured NO_3 mixing ratio (black) with calculated stationary-state mixing ratios using $k_{\text{VOC}}^{\text{NO}_3}$ (red), $k_{\text{VOC}}^{\text{NO}_3} + k_{\text{Het}}$ (blue), and $k_{\text{VOC}}^{\text{NO}_3} + k_{\text{Het}} + k_{\text{NO}}$ (green).




Figure 8: Upper panel: Stationary-state NO_3 mixing ratios calculated using $k_{\text{OTG}}^{\text{NO}_3}$, $[\text{NO}]k_3$, $K_5[\text{NO}_2]f_{\text{het}}$ and J_{NO_3} for the entire campaign and comparison with the measured NO_3 mixing ratios (03.08). The lower panel plots the time series of production and loss rates used for calculation of $[\text{NO}_3]_{\text{ss}}$.




Figure 9: The fraction, f , of the total NO_3 loss with organic trace gases as a campaign mean, diel cycle where $f = k_{\text{OTG}}^{\text{NO}_3} / (k_{\text{OTG}}^{\text{NO}_3} + J_{\text{NO}_3} + [\text{NO}]k_3 + K_5[\text{NO}_2]f_{\text{het}})$. The error bars reflect variability only and do not consider systematic uncertainty.




Figure 10: Upper panel: Time series of $k_{\text{Total}}^{\text{OH}}$ (shaded region is 1σ uncertainty). Lower panel: Time series of $k_{\text{OTG}}^{\text{NO}_3}$ and $k_{\text{OTG}}^{\text{OH}}$. The data is plotted so that the curves overlay at the peak reactivity (01/08).




Figure 11: Correlation between OH-reactivity and NO_3 -reactivity. The coloured lines are relative NO_3 and OH reactivity for single VOCs. The measured NO_3 and OH reactivities are depicted as black (nighttime) and orange datapoints (daytime).