

Brussels, 13 September 2018

Dear William,

I have now prepared the replies and changed the text according to the reviewer's comments.

During the review process we found that it is more consistent for a model intercomparison to calculate the mean age of air using as reference the tropical tropopause. The figures did not change much from the discussion paper, indicating that this is a methodological issue which does not have a large impact on our findings. In the reply to the second reviewer I explain all the details about this change and its impacts on the manuscript.

Sincerely,

Simon Chabrillat (on behalf of all co-authors).

Response to Reviewer #1 for discussion paper

Comparison of mean age of air in five reanalyses using the BASCOE transport model

Chabrilat et al., ACPD, 2018

We thank the reviewer for his/her positive and useful comments. In our replies below the bold type is used to highlight text in the revised manuscript.

- *It is stated that the model input is wind and surface pressure. Is the latter used to convert from model levels to pressure levels? Perhaps this could be said directly at some point.*

Yes: the output AoA datasets are indeed interpolated from model levels to constant pressure levels using the instantaneous and two-dimensional input surface pressures, i.e. prior to any averaging in the longitudinal or time dimension. This is now stated explicitly at the end of section 2.1.

- *Is the sentence in L15-16 (P1) of the abstract needed? It seems redundant.*

This sentence has been removed from the abstract.

- *L4 P2: “considerably depending on the considered period”: consider changing one of the two to avoid repetition.*

"Considerably" has been replaced by "substantially"

- *L30 P3: explain how these...*

Typo corrected

- *L32 P3: each tailored for a different reanalysis dataset.*

This sentence was outdated because the version published in ACPD compared with only one other transport model (TOMCAT) which is tailored for ERA-I. We have updated the sentence accordingly.

- *L14-15 P4: It could be mentioned why it is chosen not to use the vertical wind component directly.*

The revised manuscript states:

These models are usually set on a different grid than their input reanalysis dataset. Since this prevents the direct usage of the vertical wind component in the reanalysis, they rely on mass continuity to derive the vertical mass fluxes corresponding to their own grid.

- L31 P4: “idealized tracer which increases linearly at the surface”: throughout the surface or just in the tropics?

The choice of the surface as source region introduced confusing inconsistencies in the discussion paper (see the general comment by the second referee and also the next comment here). Hence we decided to re-run our calculations and re-plot all figures (except for figure 8, see below) using as source region the tropical tropopause region (defined as the 100 hPa isobar between latitudes 10°S and 10°N), and computing the AoA at each gridpoint as the time elapsed since the mixing ratio of the ideal tracer reached the same value in that source region. The figures did not change significantly from the discussion paper, indicating that this is a methodological issue which does not have any impact on our findings. The last paragraph of section 2.1 has been re-written to fully explain the updated procedure for computing AoA.

For figure 8 (and figure 8 only) we have kept the original calculations where the tracer was set to increase linearly *throughout* the surface, because this figure includes a comparison with observational values of AoA which used the surface as reference. We have moved to the discussion of figure 8 the description of this *surface* boundary condition and its propagation through the troposphere, because it is now irrelevant for all other figures. This description now states explicitly that it uses

... a synthetic tracer which is set to a global constant increasing linearly with time at the surface.

- L31-32 P6: “the AoA at the equatorial tropopause has been subtracted from the fields...”: did you use the climatological or time-dependent tropopause altitude?

See previous question: the revised manuscript shows AoA computes AoA directly from the tropical tropopause region and has dropped all *a posteriori* corrections by subtraction of AoA values at the equatorial tropopause. We have removed the sentences describing this procedure from the revised manuscript.

- L9 P8: GCCM: this has not been introduced before, do you mean CCM?

Yes. We have replaced all occurrences of "GCCM" by "CCM".

- L14-15 P8: “different latitude gradients between the tropics and ...”

Done.

- L13 P9: remove “globally”, it is not global but midlatitude average.

Done.

- L30 and 31 P9: “different with” should be “different from”

Done.

- L11 P10: “not significant”: in ERA-Interim or in observations?

Replaced "sign of observational trend not significant" by clearer

...sign of trends not significant in the observations.

- *L12 P10: “ERA-I does not show any overall trend after 2000...”. Why do you point out these trends after 2000 in ERA-I specifically? Same thing on P16 L22-23.*

This focus on ERA-I is due to the exclusive use of ERA-I in previous studies modeling the latitudinal structure of AoA for the post-2000 period (see first paragraph of section 4.3, P12 of ACPD manuscript). But this context had not been introduced yet for the discussion of figure 8 (i.e. L12 P10 and P16 L22-23). Since this question is specifically investigated through figures 11 and 12, we have simply removed the premature sentence from the discussion of figure 8.

- *L23 P12: standard error for which confidence level?*

This important information has been added in section 4.1 which describes our methodology for multi-linear regressions:

The uncertainties arising from the fit are calculated for the 95% confidence interval and corrected for auto-correlation in the residuals (Eqs. 3,4 and 6 in Santer et al., 2000).

and in the section 4.3 (discussing linear trends):

It is expressed in years per decade (yr dec^{-1}) and is deemed significant at a given grid point if its absolute value is larger than its uncertainty (as defined in section 4.1).

- *L1 P13: remove “unexpected”.*

Done.

- *L2 P13: remove “much”*

Done.

- *L7 P13: “Diallo et al. (2012)...” Ploeger et al. (2015a) show AoA trends for the total and later periods.*

Thanks for pointing this out. In the revised manuscript we now also compare our results with those by Ploeger et al. (2015a), both for the discussion of the latter period (figure 11):

Our results also agree well with those obtained by a diabatic model driven by ERA-I over the same period (Ploeger et al., 2015a).

and for the discussion of the total period (first paragraph discussing figure 12):

Our ERA-I results for the overall period partly contradict those obtained by diabatic models which use not only the wind fields from ERA-I but also its heating rates (Diallo et al., 2012; Ploeger et al., 2015a). Looking at slightly shorter periods of two decades (1989–2010 for the former and 1990–2013 for the latter), these papers reported negative AoA trends for both hemispheres below 28km altitude.

- *L10 P13: “using only wind fields”: do you mean not using heating rates? Perhaps this should be explicit.*

Indeed we meant that our results did not use the heating rates. During our revision we found that the whole sentence was confusing and removed it from the manuscript. The additional use of ERA-I heating rates by the diabatic models (Diallo et al., 2012; Ploeger et al., 2015a) is now explicitly

stated as soon as they are cited (see previous comment).

- L26-27 P14: *“While this may be a coincidence...”*: but having more wave drag would imply a faster BDC, so I do not see the point of this sentence.

Agreed. This sentence has been removed from the revised manuscript.

- L34 P14: *“Miyazaki et al. (2016) Fig. 11”*: The trends in annual mean tropical upwelling for these reanalyses are shown in Abalos et al. (2015) Fig. 11.

The following sentence has been added to the discussion:

Similar disagreements have also been reported between the trends of the annual mean tropical upwelling in three reanalyses over the period 1979–2012, with vertical residual velocities (w^*) increasing in MERRA and JRA-55 and decreasing in ERA-I (Abalos et al., 2015, Fig. 11).

- L18 P15: *Another difference with CLaMS is that it includes a mixing parameterization.*

The revised version states also this difference, citing Konopka et al. (JGR, doi:10.1029/2003JD003792, 2004).

- *Figure 7 caption: “No impact...”* This sentence does not belong to the caption but to the main text.

This sentence has been removed from the caption of Fig.~7.

Added references

Konopka, P., Steinhorst, H.-M., Grooß, J.-U., Günther, G., Müller, R., Elkins, J. W., Jost, H.-J., Richard, E., Schmidt, U., Toon, G., and McKenna, D. S.: Mixing and ozone loss in the 1999–2000 Arctic vortex: Simulations with the three-dimensional Chemical Lagrangian Model of the Stratosphere (CLaMS), *Journal of Geophysical Research: Atmospheres*, 109, doi:10.1029/2003JD003792, 2004.

Response to Reviewer #2 for discussion paper

Comparison of mean age of air in five reanalyses using the BASCOE transport model

Chabrillat et al., ACPD, 2018

We thank the reviewer for his/her insightful comments. It appears that the version of the manuscript which was reviewed by this referee is the version first submitted to ACPD (on 4 April 2018) rather than the version finally published in ACPD (on 7 May 2018). Fortunately all comments apply equally to both versions. In our replies below the bold type is used to highlight text in the revised manuscript.

Replies to general comments

- *The principal conclusion is that the simulations of AoA obtained when BASCOE is constrained by different reanalyzed datasets differ substantially from one another. This is not at all unexpected given the differences among the reanalysis models.*

The reanalysis systems are based on different models but they assimilate very similar satellite datasets. Many users of reanalyses are neither modellers of stratospheric dynamics nor aware of the lack of observational information to constrain the BDC in the reanalyses. From feedback obtained at the 5th International Conference on Reanalysis (ICR5, Rome, November 2017), such users do not expect to see a spread between the reanalyses which is as large as between unconstrained GCMs (Fig. 4). On Fig. 8 they easily understand that the uncertainties in the observational timeseries are large (due to sparse and irregular sampling) but they do not expect to see that the spread between the reanalyses is as large as these observational uncertainties.

A third highlight of this paper is the intercomparison of AoA trends between the reanalyses. Several reanalysis intercomparisons of diagnostics related to stratospheric dynamics have already been published and showed significant differences with respect to their trends (e.g. Abalos et al., 2015; Miyazaki et al., 2016). Yet for the AoA diagnostic, most recent studies rely on ERA-I with much interest in the latitudinal structure of its trends. We found that over the post-2002 period ERA-I is the *only* reanalysis to deliver opposite trends of AoA in the two hemispheres (Fig. 12, middle column). This is also an unexpected result.

- *The paper is well organized and clearly written, with some exceptions, the main one being that the procedure for computing AoA is not well explained. In particular, it is not clear whether AoA is calculated with respect to a reference level at the tropical tropopause or in the troposphere, and this introduces some ambiguity in the interpretation of the results.*

We agree with the referee that the handling of the reference level was problematic in the submitted manuscript. All our calculations used the surface both as the source region and to compute the time lag defining the mean Age of Air. But in order to better highlight the different transit times from the equatorial tropopause, Fig.1 and 3 were corrected a posteriori by subtraction of the time-averaged AoA at 100hPa, 10°S-10°N. All other figures used the surface as reference, hence including the transit time from the surface to the tropical tropopause. This distinction was not clearly made and led to inconsistent figures, as shown by several specific comments made by both reviewers.

Hence we decided to re-run our calculations and re-plot all figures (except for figure 8, see below) using as source region the tropical tropopause region (still defined as the 100 hPa isobar between latitudes 10°S and 10°N), and computing the AoA at each grid point as the time elapsed since the mixing ratio of the ideal tracer reached the same value in that source region. The last paragraph of section 2.1 has been re-written to fully explain the updated procedure for computing AoA.


The figures did not change much from the discussion paper, indicating that this is a methodological issue which does not have a large impact on our findings. Besides figures 3 and 9 which are discussed below for specific comments, there is one other case where the figure changed sufficiently to warrant a minor update in the text: on figure 12 the positive AoA trends for ERA-I in 2002-2015 (top row, middle column) have become significant at all latitudes (in the discussion paper they were significant only in the polar latitudes). On figure 12 the signs and patterns of AoA trends did not change for any other reanalysis or period but the range of these trends increased by up to 50% (see min/max values above the plots); this led us to extend the scale of the color bar, from [-0.6,0.6] to [-0.9,0.9].

For figure 8 (and figure 8 only) we have kept the original calculations where the tracer was set to increase linearly throughout the surface, because this figure includes a comparison with observational values of AoA which used the surface as reference. We have moved to the discussion of figure 8 the description of this surface boundary condition and its propagation through the troposphere, because it is now irrelevant for all other figures. We have also added in this figure a plot showing tropical AoA computed both from the surface and from the tropical tropopause to show that the difference does not vary significantly with the simulated year (see next comment).

Replies to specific comments

- (4, 20) “There is no other representation of convection”: *It is not clear that, in the Tropics, where deep convection can reach the 14-15 km level, this artificial diffusion can simulate vertical transport realistically. But perhaps this does not matter for assessments of AoA in the stratosphere if the base point for AoA calculations is taken to be at or near the tropical tropopause? Please comment (especially since it is not clear how the reference level for computing AoA is chosen). See also comment at (7, 24).*

As explained above, nearly all figures now use (100hPa, 10°S-10°N) as reference region hence the absence of deep convection in the CTM is not an issue for the updated figures. Since for figure 8 we keep using the surface as reference, we added tropical timeseries in the tropics which show both the surface-based evaluation (solid lines) and the tropopause-based evaluation (dashed lines):



Middle panel of revised Fig. 8. Mean AoA in the mid-stratosphere (5-30hPa) for the tropical latitudes (30°S–30°N). Solid lines show AoA using the surface as reference, dashed lines show AoA using the tropical tropopause as reference (i.e. as in all other figures of the revised manuscript).

This comparison between the two evaluations also allows a discussion on the impact of the omission of deep convection in the model. The discussion of Fig. 8 in the revised manuscript includes the following paragraph:

The differences between the two calculations represent the transit times from the surface to the tropical tropopause, are nearly independent of the simulated year and range between 3 months (with ERA-I or JRA-55) and 6 months (with MERRA). These values are close to the longest transit times reported in a recent intercomparison of global models (Krol et al., 2018), indicating a rather slow transport from the surface to the tropical tropopause which we attribute to the omission of deep convective transport in our model. While the surface-based model AoA (solid lines in Fig. 8) may be slightly overestimated, these biases have no significant inter-annual variations and do not hinder the intercomparison between reanalyses.

- (6, 5) “at the wavelength number 47”: “at wavenumber 47” might be better.

Text corrected.

- (6, 8) “Figure 1 compares the results”: I do not believe you have stated how AoA is calculated. [...]

Section 2.1 now describes explicitly the revised procedure to calculate AoA from the tropical tropopause region:

The age of air is defined as the spectrum of transit times from a source region to a given location, with the tropical tropopause usually defining the source region for studies of the stratosphere. In the case of ideal tracers which increase linearly in the source region and have no photochemical production or losses, the mean of this spectrum (denoted here AoA) is simply the time elapsed since the mixing ratio of this ideal tracer reached the same value in the source region (see e.g. Waugh and Hall, 2002). We follow here this classical approach, using for most simulations the 100 hPa isobar between latitudes 10°S and 10°N as source region.

Section 3.2 describes explicitly the original procedure which has been kept only for figure 8:

For consistency the modeled AoA in this figure are evaluated as the time elapsed since the mixing ratio of an ideal tracer reached the same value at the surface, using as boundary condition a global constant increasing linearly with time at the surface.

- (7, 18) Figure 2: This figure, as well as Figure 3, would benefit from a color bar to indicate the values of the AoA isolines not explicitly labeled.

A color bar has been added to figures 2 and 3.

- (7, 24) Figure 3: I am confused by this comparison [...] So, where is the reference point in these simulations? If it is at the surface, then AoA will reflect the effects of transport not just in the stratosphere, but also in the troposphere, including the artificial diffusive transport between the surface and the middle troposphere. Unless I am misunderstanding what you have done here, it seems to me that, if AoA is intended to highlight transport in the stratosphere (e.g., Waugh and Hall, 2002, Sec. 3.1) then the choice of a base point in the troposphere confuses the issue, especially given the use of artificial diffusive transport in the lower troposphere.

The reviewer was rightly confused and his interpretation is correct. We have followed this advice, choosing the tropical tropopause as reference point in the revised manuscript (see above). The relative differences between ERA-I and the four other reanalyses vanish at the reference point and the difference is not plotted at grid points where ERA-I AoA is smaller than 5 days.

- (8, 5) *“The intercomparison at 50 hPa”: You should state explicitly in the text that in these comparisons AoA is “normalized” to be zero at the tropical tropopause (this is only stated in the caption of Figure 4). Otherwise, the reader will wonder, as I did, why the AoA shown in Figures 2-3 are different from the AoA in Figure 4. By the way, a problem with the “normalization” of AoA to zero at 100 hPa is that it gives the impression that AoA above that level is determined only by the stratospheric circulation, when in fact the AoA also contains the effect of transport in the troposphere.*

thanks to the direct calculation of AoA using the tropical tropopause as reference point, no "normalization" is performed any more for the figures 1 and 4 of the revised manuscript.

- (8, 12) *“overall, the spread . . . is larger than the 1-sigma. . .”: One wonders how this result would change if AoA were computed with respect to a reference point at 100 hPa.*

In the revised manuscript the AoA are computed with respect to a reference point at 100 hPa. The differences in Figure 4 between the submitted and revised manuscripts are nearly indistinguishable. Hence the spread between the five simulations at 50 hPa is still larger than the 1- σ observational uncertainties in the tropics, and still nearly as large in the extratropics. We have not modified this sentence in the revised manuscript.

- (8, 26) *“The spread between the four reanalyses reaches a maximum of 0.2 years at 30 hPa”: Are you referring here to the gradient comparison, Figure 4d? How is this “gradient” calculated? The figure legend refers to “MLNH-Tropics” and shows values in months, not per unit distance, so this is really a difference between the Tropics and midlatitudes of the NH. How are Tropics and NH midlatitudes defined?*

The words "(mean age) gradient profiles" or "latitudinal gradients (of mean age)" were meant with the same meaning as Neu et al. (2010) and Chipperfield et al. (2014) i.e. as the difference between AoA in NH midlatitudes and AoA in the Tropics. The vertical profiles on figure 4d simply show the differences between the corresponding profiles on figures 4c and 4b which are mean values for latitude bands 35°N-45°N and 10°S-10°N respectively (as stated in the figure of caption 4).

In the revised manuscript we have added the definition of the latitude bands in the discussion of figures 4b and 4c and we have added the following sentences in the discussion of figure 4d:

These "latitudinal gradients of AoA" were used in several CCM intercomparisons (Neu et al., 2010; Chipperfield et al., 2014). Figure 4d shows this diagnostic for the five reanalyses, i.e. the differences between the AoA profiles on Fig. 4c and Fig. 4b.

We have replaced the words "latitudinal gradients" by "AoA differences" in the remainder of this discussion and in the caption of the figure.

- (8, 30) “MERRA-2 yields an outlying vertical profile of AoA at northern midlatitudes”: True with respect to the other reanalyses except for MERRA (Fig. 4c), and in fact, MERRA and MERRA-2 midlatitude profiles of AoA agree best with the observations. You keep referring to MERRA-2 as an “outlier”, which carries negative connotations, but in fact being an outlier in this comparison is a good thing if one considers the data to be the “truth”..

Thanks for pointing this out. We have corrected the discussion according to your comment:

MERRA and MERRA-2 yield larger AoA at northern midlatitudes than the three other reanalyses. In the case of MERRA-2 this results in a profile of AoA differences which are significantly larger than the profiles obtained with the four other reanalyses but agrees much better with the profile derived from the observations. Hence MERRA-2 apparently underestimates the tropical upwelling in the lowermost stratosphere (100-60~hPa), agrees better with the observations at 50~hPa than any other reanalysis, and joins the results of the four other reanalyses at higher levels.


- (9, 14) “MERRA-2 starts with much older values”: This behavior does appear to be anomalous. Any idea what might be causing it?

This issue is discussed in detail in section 5 (see paragraph starting with "MERRA-2"). We have inserted the following sentence in section 3.2:

The possible causes for this apparently anomalous behavior of MERRA-2 are discussed in section 5.

- (9, 18) “The Pinatubo eruption does not appear to have any impact of the simulated AoA at 50 hPa”: Insofar as one might expect that the largest impact of Pinatubo would be in the Tropics, it might be worthwhile to examine the AoA time series averaged over, say, 30N-30S.

This was done and no impact was found for the tropical latitude band, as shown by the corresponding plot:



Note that any impact in the 30°S-30°N latitude band would have been seen in figure 7 which shows the 72°S-72°N band. The revised manuscript mentions the absence of volcanic impact in the tropical latitude band as well.

- (10, 8) “observational trend is not significant”: One would not expect any trend calculated from the smoothed, model time series shown on the right pane of Figure 8 to be significant either. By the way, you keep referring to “trends” in connection with the model results, but you have not calculated any trends...

Not yet. As outlined in the introduction, modelled trends are evaluated extensively in section 4. The sentence now refers to this later finding:

ERA-I delivers a weakly positive trend over the period 1989-2015 and we will assess in section 4.3 that this trend in the model results is significant.

And this finding has been added in the discussion of figure 12 (section 4.3):

The same plot (i.e. Fig. 12, top right) also shows that the positive trend which had been inferred visually for the northern mid-latitudes of the middle stratosphere (Fig. 8, left) is significant.

- ...Note also that Garcia et al. (2011) have argued that, even using model output for an ideal AoA tracer, trends over periods as long as 30 years are often not significant when the ideal tracer is sampled like the available observations of stratospheric tracers....

Great caution should indeed be exercised in comparisons of trends between model output and observational datasets which are sparsely and irregularly sampled. This is what we meant in the original manuscript with "..., but Engel et al. (2009, 2017) showed that the sign of this observational trend is not significant". Referring to Garcia et al. (2011) allows us to reinforce and clarify this call to caution. The revised manuscript states:

While the overall trend simulated with ERA-I is apparently in agreement with the balloon observations, this comparison should be considered with great caution because the sign of the AoA trend is not significant in the observations (Engel et al., 2009, 2017) and modelled trends over periods as long as 30 years are often not significant when the ideal tracer is sampled like the available observations of stratospheric tracers (Garcia et al., 2011).

- ... Furthermore, trends derived from observation are also confounded by the fact that no real atmospheric tracer has a constant, linear growth rate.


The non-linearity of the growth rate of CO₂ and SF₆ has of course been taken into account by Engel et al. (2009, 2017) for their derivation of AoA and also for their error analysis. The procedure is described in the supplementary information of Engel et al. (2009) and in the first paragraph of section 4 in Engel et al. (2017).

- (11,16) “in the polar regions and midlatitudes”: Why limit this to extratropical behavior only? It would be interesting to show the seasonal amplitude in the Tropics as well, say a $\pm 30^\circ$ average.

Thanks for the suggestion. The seasonal amplitudes with ERA-I and MERRA-2 have a different vertical structure in the Tropics. We have added this plot to figure 9 in the revised manuscript.

- (11, 24) “MERRA and MERRA-2 . . . different amplitudes depending on the period used or the analysis”: Does this have to do with the development (1989-2001) and stabilization/decline (2002-2015) of the Antarctic ozone hole? To explore this issue, one would have to examine the actual seasonal climatology at high SH latitudes, not just the annual amplitude.

The left plot below is extracted from fig. 9 in the ACPD manuscript while the right plot shows results with the new AoA calculation (i.e. using the tropical tropopause as reference):



This shows that the new AoA calculation delivers amplitudes of AoA seasonal variations which are much closer for periods 1989-2001 and 2002-2015. This is the case also for the other latitude bands, so for the revised manuscript we have removed from figure 9 the results for the early period 1989-2001 and we have dropped the corresponding part of the discussion.

- (12, 9) “could not be expected from inspection of the native dynamic variables”: I do not understand what you are trying to say here. Please elaborate. And note also that the discrepancies you mention (“up to 50% dependencies on the considered time period”) are not even illustrated, so it is very difficult to even guess what the intent of your statement is.

The dependencies of seasonal amplitudes on the considered time period reached 50% in fig. 9a and 9c of the ACPD manuscript but, as explained in the previous comment, they have disappeared from the revised manuscript thanks to the corrected AoA calculation. We removed the whole paragraph (i.e. last paragraph of section 4.2) from the revised manuscript because it is redundant with the second paragraph of section 5.

- (12, 15) “unexpectedly increasing”: Given the very short period covered by the SF₆ observations, it is not clear that one should “expect” any particular sign for the trends. Determination of AoA trends from observations of stratospheric tracers is fraught with many uncertainties; even in models where an ideal, linearly increasing artificial tracer is used, one has to rely on zonal-mean results over long periods to obtain trends that are clearly statistically significant. Arguably, examination of AoA trends determined from observations of stratospheric tracers is not the best tool for documenting changes in the BD circulation. See Garcia et al. (2011).

We acknowledge that it is important to provide proper context about this topic. Section 4.3 now starts with a new paragraph explaining some caveats of interpreting AoA trends as changes in BDC. The next paragraph has been expanded and corrected to justify the interest of the section:

It is delicate to infer changes in the BDC on the basis of AoA trends over periods shorter than several decades. Even in models where an ideal, linearly increasing artificial tracer is used, one has to rely on zonal-mean results over long periods to obtain trends that are clearly statistically significant (Garcia et al., 2011). The statement that is often made that climate models simulate a decreasing AoA throughout the stratosphere only applies over long time periods and is not necessarily the case for the past 25 years, when most tracer measurements were taken (Garfinkel et al., 2017). For example, the analysis of a 1700 year simulation showed that it takes around 30 years for a modeled BDC trend to emerge from the noise of natural climate variability (assuming a 2%/decade trend in the BDC; Hardiman et al., 2017).

While linear trends of AoA over shorter periods may represent transient changes due to climate variability, such changes over timescales which are intermediate between the QBO and the multidecadal scales are still relevant to the study of stratospheric dynamics. Current research on AoA trends has largely focused on a dipole-like latitudinal structure for the period 2002-2012, which was first derived from satellite observation of SF₆ by the MIPAS instrument (Stiller et al., 2012). This structure of trends shows AoA decreasing in the Southern Hemisphere but ~~unexpectedly~~ increasing in the Northern Hemisphere **which was used to explain a recent increase of stratospheric HCl in the Northern Hemisphere (Mahieu et al., 2014) and interpreted as the consequence of a southward shift of the subtropical transport barriers (Stiller et al., 2017).**

- (13, 1) “unexpected growth”: Again, there are no clear expectations about trends for short periods.

We have removed the word "unexpected" from this sentence.

- (13, 23) “the reversal is found for all five reanalyses”: This is true, but the reversals are in the opposite sense in ERAi and CFS vs. JRA-55, MERRA and MERRA2, so it is hard to know what to make of this.

Yes but we still want to highlight this striking feature in case some reader does find what to make of this. We have changed this sentence to:

This reversal is found for all five reanalyses and in all regions of the stratosphere **but it is difficult to interpret because it goes in opposite directions in ERA-I and CFSR versus JRA-55, MERRA and MERRA-2.**

- (13, 25) “unexpectedly large disagreements”: I am not sure why you think the disagreements are “unexpected”. While presumably all reanalyses use more or less the same observational data, the manner in which the data are assimilated and the physical parameterizations included in the different models (in particular, those for convection and mesoscale gravity waves) are different. Note that at (14, 8) you suggest that “the disagreements found here may lie in the differences between the underlying models”; I agree that this is the most plausible working hypothesis.

See above the reply to the first general comment. The word "unexpectedly" had already been removed before publication in ACPD.

Grammar, typos, etc.

All errors have been corrected.

Added references

- Garcia, R. R., Randel, W. J., and Kinnison, D. E.: On the Determination of Age of Air Trends from Atmospheric Trace Species, *J. Atmos. Sci.*, 68, 139–154, doi:10.1175/2010JAS3527.1, 2011.
- Garfinkel, C. I., Aquila, V., Waugh, D. W., and Oman, L. D.: Time-varying changes in the simulated structure of the Brewer–Dobson Circulation, *Atmospheric Chemistry and Physics*, 17, 1313–1327, doi:10.5194/acp-17-1313-2017, 2017.
- Hardiman, S. C., Lin, P., Scaife, A. A., Dunstone, N. J., and Ren, H.-L.: The influence of dynamical variability on the observed Brewer–Dobson Circulation trend, *Geophys. Res. Lett.*, doi:10.1002/2017GL072706, 2017.
- Krol, M., de Bruine, M., Killaars, L., Ouwersloot, H., Pozzer, A., Yin, Y., Chevallier, F., Bousquet, P., Patra, P., Belikov, D., Maksyutov, S., Dhomse, S., Feng, W., and Chipperfield, M. P.: Age of air as a diagnostic for transport timescales in global models, *Geoscientific Model Development*, 11, 3109–3130, doi:10.5194/gmd-11-3109-2018, 2018.

Comparison of mean age of air in five reanalyses using the BASCOE transport model

Simon Chabrillat¹, Corinne Vigouroux¹, Yves Christophe¹, Andreas Engel², Quentin Errera¹, Daniele Minganti¹, Beatriz M. Monge-Sanz³, Arjo Segers⁴, and Emmanuel Mahieu⁵

¹Royal Belgian Institute for Space Aeronomy, BIRA-IASB, Brussels, 1180 Belgium

²Institute for Atmospheric and Environmental Science, Goethe University Frankfurt, Frankfurt, Germany

³European Centre for Medium-Range Weather Forecasts, Shinfield Park, Reading, RG29AX, UK

⁴TNO, Department of Climate, Air and Sustainability, P.O. box 80015, 3508 TA Utrecht, the Netherlands

⁵Institute of Astrophysics and Geophysics, University of Liège, 4000 Liège, Belgium

Correspondence to: Simon Chabrillat

(Simon.Chabrillat@aeronomie.be)

Abstract.

We present a consistent intercomparison of the mean Age of Air (AoA) according to five modern reanalyses: the European Centre for Medium-Range Weather Forecasts Interim Reanalysis (ERA-Interim), the Japanese Meteorological Agency's Japanese 55-year Reanalysis (JRA-55), the National Centers for Environmental Prediction Climate Forecast System Reanalysis (CFSR) and the National Aeronautics and Space Administration's Modern Era Retrospective-analysis for Research Applications version 1 (MERRA) and version 2 (MERRA-2). The modeling tool is a kinematic transport model driven only by the surface pressure and wind fields. It is validated for ERA-I through a comparison with the AoA computed by another transport model.

The five reanalyses deliver AoA which differ in the worst case by one year in the tropical lower stratosphere and more than two years in the upper stratosphere. At all latitudes and altitudes, MERRA-2 and MERRA provide the oldest values (~5-6 years in mid-stratosphere at mid-latitudes) while JRA-55 and CFSR provide the youngest values (~4 years) and ERA-I delivers intermediate results. The spread of AoA at 50 hPa is as large as the spread obtained in a comparison of Chemistry-Climate Models. The differences between tropical and mid-latitudes AoA are in better agreement except for MERRA-2. Compared with in-situ observations, they indicate that the upwelling is too fast in the tropical lower stratosphere. The ~~general hierarchy of reanalyses delivering older AoA (MERRA, MERRA-2) and younger AoA (JRA-55, CFSR) holds during the whole 1989–2015 period, with AoA derived from ERA-I keeping intermediate values. The~~ spread between the five simulations in the northern mid-latitudes is as large as the observational uncertainties in a multidecadal time series of balloon observations, i.e., approximately two years. No global impact of the Pinatubo eruption can be found in our simulations of AoA, ~~contrarily~~ contrary to a recent study which used a diabatic transport model driven by ERA-I and JRA-55 winds and heating rates.

The time variations are also analyzed through multiple linear regression analyses taking into account the seasonal cycles, the Quasi-Biennial Oscillation and the linear trends over four time periods. The amplitudes of AoA seasonal variations in the lower stratosphere are significantly larger using MERRA and MERRA-2 than with the other reanalyses (~~up to twice as large at the~~

50 hPa pressure level). The linear trends of AoA using ERA-I confirm those found by earlier model studies, especially for the period 2002–2012 where the dipole structure of the latitude–height distribution (positive in the northern mid-stratosphere and negative in the southern mid-stratosphere) also matches trends derived from satellite observations of SF₆. Yet the linear trends vary considerably-substantially depending on the considered period. Over 2002–2015 the ERA-I results still show a dipole structure but it is much less pronounced, with positive trends in the northern hemisphere remaining significant only in the polar lower stratosphere (where they reach 0.2 Northern Hemisphere reaching up to 0.3 years per decade). No reanalysis other than ERA-I finds any dipole structure of AoA trends. The signs of the trends depend strongly on the input reanalysis and on the considered period, with values above 10 hPa varying between approximately -0.4 and 0.4 years per decade. Using ERA-I and CFSR, the 2002–2015 trends are negative above 10 hPa but using the three other reanalyses these trends are positive. Over the whole period 1989–2015 each reanalysis delivers opposite trends, i.e., AoA is mostly increasing with CFSR and ERA-I but mostly decreasing with MERRA, JRA-55 and MERRA-2.

In view of these large disagreements, we urge great caution for studies aiming to assess AoA trends derived only from reanalysis winds. We briefly discuss some possible causes for the dependency of AoA on the input reanalysis and highlight the need for complementary intercomparisons using diabatic transport models.

15 1 Introduction

The mean age of air (hereafter AoA) is an evaluation of the time necessary for variations of long-lived (e.g., greenhouse or ozone-depleting) species to propagate from the troposphere to various regions in the stratosphere. This classical diagnostic provides insights on the strength and structure of the Brewer-Dobson Circulation (BDC), the polar vortex, and irreversible mixing in the mid-latitudes (Waugh and Hall, 2002). Due to increased greenhouse gas forcing, Chemistry-Climate Model (CCM) simulations of the 1990–2090 period predict an acceleration of the BDC and a decrease of AoA at all latitudes in the lower part of the stratosphere (Austin and Li, 2006; Butchart, 2014). The observational detection of trends in the BDC strength turns out to be quite difficult. They can be indirectly derived from multidecadal records of stratospheric temperatures but these derivations are indirect and do not yet allow a clear confirmation of the acceleration predicted by CCM, mainly due to an insufficient-quality-insufficiently constrained accuracy of the temperature observations (Fu et al., 2015; Ossó et al., 2015).

25 Observation-based AoA is derived from concentration measurements of very long-lived tracers which increase (nearly) monotonously-monotonically at the surface, such as CO₂ or SF₆. Multidecadal datasets were compiled from balloon soundings or aircraft flights (e.g., Andrews et al., 2001; Ray et al., 2014, and references therein). The corresponding time series are precise but sparse in time and space, leading to large sampling uncertainties. Global coverage time series have been derived from satellite observations, but the precision is lower. The SF₆ retrievals from the Michelson Interferometer for Passive Atmospheric Sounding (MIPAS) satellite instrument delivered a continuously updated dataset with global coverage for the period 2002–2012, leading to breakthrough studies about observed AoA and its time variations during this comparatively short period (Stiller et al., 2008; Haenel et al., 2015). The magnitude, distribution and detectability of the AoA trends observed over the past years and

decades are currently a topic of intense research (e.g., Engel et al., 2009; Stiller et al., 2012; Mahieu et al., 2014; Engel et al., 2017).

Reanalysis systems combine a global weather forecast model, observations, and an assimilation scheme to provide the best estimates (analyses) of past atmospheric states including surface pressure, temperature, and wind over a long (usually multi-decadal) period. While they are derived from assimilation systems used operationally to deliver weather forecasts, they aim to achieve more consistent variations over long timescales, e.g., avoiding spurious discontinuities and trends (Trenberth and Olson, 1988; Bengtsson and Shukla, 1988). Hence the same model version and assimilation scheme are used for the whole period and special care is given to the time-varying biases between the assimilated observations (see, e.g., Simmons et al., 2014). The resulting reanalysis datasets provide a multivariate, spatially complete, and coherent record of the global atmospheric circulation.

The Stratosphere–troposphere Processes And their Role in Climate (SPARC) Reanalysis Intercomparison Project (S-RIP) is a coordinated intercomparison of all major global atmospheric reanalyses. Its introductory paper (Fujiwara et al., 2017) provides an overview of the past and current reanalysis systems and datasets. The present study deals with the five modern reanalyses of surface and satellite data retained in S-RIP: the European Centre for Medium-Range Weather Forecasts Interim Reanalysis (ERA-Interim), the Japanese Meteorological Agency’s Japanese 55-year Reanalysis (JRA-55), the National Centers for Environmental Prediction Climate Forecast System Reanalysis (NCEP-CFSR) and the National Aeronautics and Space Administration’s Modern Era Retrospective-analysis for Research Applications version 1 (MERRA) and version 2 (MERRA-2).

The absolute value of AoA and its evolution over the past decades can be derived from the surface pressure and wind fields available in such reanalyses, using either an offline transport model (see, e.g., Chipperfield, 2006) or a chemistry-climate model nudged to the input reanalysis (Kunz et al., 2011; Kovács et al., 2017) to model the transport of inert tracers propagating from the troposphere to the stratosphere. This approach helped to identify shortcomings in the Brewer-Dobson circulation described by early reanalyses (Meijer et al., 2004; Pawson et al., 2007) and to assess the improvements in the next generation of reanalyses, e.g., from ERA-40 to ERA-Interim (Monge-Sanz et al., 2007; Dee et al., 2011; Monge-Sanz et al., 2012).

Few AoA comparisons have been performed between reanalyses originating from different reanalysis centers. This is mainly due to technical difficulties ~~which~~that are not limited to file formatting issues. While all modern systems use hybrid $\sigma - p$ vertical coordinates (Simmons and Burridge, 1981), each reanalysis comes with a wind field computed on a different grid with different horizontal and vertical resolutions. Some reanalysis forecast models use spectral dynamical cores (Krishnamurti et al., 2006) while others use finite-volume dynamics (Lin, 2004, see next section for details). A common offline transport model may have difficulties dealing with such different grids because it is usually tailored for a specific family of reanalyses, e.g., using an advection algorithm similar to the dynamical core of the driving reanalysis system or climate model (Strahan and Polansky, 2006).

Section 2 describes the input reanalyses and our modeling tools to ~~explains~~explain how these difficulties were circumvented. It also validates our approach with a classical set of observations and with the results of ~~two other transport models, each tailored for its own reanalysis dataset~~another transport model which is tailored for ERA-I.

The main purpose of this paper is to provide a comparison of the AoA obtained from five modern reanalyses included in the S-RIP project in order to assess their level of agreement or to identify outliers. Its focus is not on detailed comparisons with observations (which are deferred to a follow-on study) but rather on a consistent intercomparison between the reanalyses through the use of a common transport model.

Section 3 compares the distribution of the AoA obtained from each reanalysis for a reference period and its time evolution in the middle latitudes. Section 4 uses a multiple linear regression model to characterize the time variations of AoA, including an intercomparison of their linear trends for several periods. Section 5 proposes a brief overview of the possible causes for the disagreements between the reanalyses and states the further work required to elucidate these disagreements. Section 6 concludes the paper with a summary of our findings and their implications.

2 Methodology

2.1 Description and set-up of the offline transport model

Depending on their vertical coordinate system and the reanalysis data used as input, one may distinguish between kinematic and diabatic transport models (Mahowald et al., 2002; Chipperfield, 2006). Diabatic models use isentropic (θ) or hybrid $\sigma - \theta$ vertical coordinates and calculate the vertical transport from diabatic heating rates which may be read from the input reanalysis or re-computed using a separate radiation scheme. Kinematic transport models on the other hand need on input only the surface pressure and horizontal wind fields, ~~relying~~. These models are usually set on a different grid than their input reanalysis dataset. Since this prevents the direct usage of the vertical wind component in the reanalysis, they rely on mass continuity to derive the vertical mass fluxes corresponding to their own grid. The present study uses the kinematic transport model developed for the Belgian Assimilation System for Chemical Observations (BASCOE: see Errera et al., 2008; Skachko et al., 2014; Lefever et al., 2015). Its advection module is the Flux-Form Semi-Lagrangian (FFSL) scheme (Lin and Rood, 1996) configured to follow closely the recommendations of Rotman et al. (2001). We briefly summarize here this configuration because it has an important impact on the simulated distribution of AoA in the stratosphere.

The FFSL advection scheme is run on a evenly-spaced latitude-longitude grid with $2^\circ \times 2.5^\circ$ increments. This grid spacing is typical for current simulations of stratospheric chemistry and transport over several decades (Morgenstern et al., 2017). Using the FFSL algorithm, Strahan and Polansky (2006) showed that this is the minimum resolution allowing a realistic representation of the tropical and high latitude mixing barriers. The FFSL algorithm does not require satisfaction of the Courant-Friedrichs-Lewy (CFL) condition in the longitudinal direction, which is a big computational advantage for regular longitude-latitude grids. The time step is set to 30 minutes by default and automatically split into integer fractions in order to satisfy the CFL condition in the meridional direction. The algorithmic structure of the FFSL scheme allows multiple choices for monotonicity constraints that have implications on the subgrid tracer distribution used to calculate fluxes across cell edges. These choices are made separately in the longitudinal, meridional and vertical directions. Rotman et al. (2001) showed that AoA calculations are very sensitive to the choice of constraint in the vertical direction: realistic results require a positive-definite Piece-wise Parabolic Method, where the constraint on the subgrid distribution is only strong enough to prevent generation of negative

values but overshoots and undershoots are allowed. There is no representation of convection in the model nor any explicit mechanism for horizontal diffusion.

~~In all our simulations the AoA is computed from the distribution of an idealized~~ The age of air is defined as the spectrum of transit times from a source region to a given location, with the tropical tropopause usually defining the source region for studies of the stratosphere. In the case of an ideal tracer which increases linearly at the surface, in the source region and has no photochemical productions or losses, one can obtain the mean of this spectrum (denoted here AoA) at any time and location from the corresponding mixing ratio of the tracer: in such a case the AoA is simply the time elapsed since the ideal tracer had the same mixing ratio in the source region (Vaugh and Hall, 2002) . We follow here this classical approach, using for most simulations the 100 hPa isobar between latitudes 10°S and 10°N as source region. In one case we have used the surface as source region in order to enable a comparison with a long time series of balloon observations (see section 3.2). The output AoA datasets are interpolated from model levels to constant pressure levels using the instantaneous and two-dimensional input surface pressures, i.e. prior to any averaging in the longitudinal or time dimension.

~~In order to allow quick propagation of this boundary condition to the free troposphere, eddy vertical diffusion is modeled in the lower half of the troposphere with a vertical diffusion coefficient K_{zz} decreasing from an arbitrary value of $10 \text{ m}^2 \text{ s}^{-1}$ at the surface to zero at the pressure level halfway between the surface and the tropopause. There is no other representation of convection in the model nor any explicit mechanism for horizontal diffusion.~~

2.2 Description of the input reanalyses

We compute and compare the AoA in five recent reanalyses which are described in detail by Fujiwara et al. (2017): ERA-Interim (~~Dee et al., 2011~~) (European Centre for Medium-Range Weather Forecasts Interim Reanalysis; Dee et al., 2011) , JRA-55 (~~Kobayashi et al., 2015~~) (Japanese 55-year Reanalysis; Kobayashi et al., 2015) , MERRA (Modern Era Retrospective-Analysis for Research; Rienecker et al., 2011) , MERRA-2 (Gelaro et al., 2017) and NCEP-CFSR (~~Saha et al., 2010~~) (National Centers for Environmental Prediction–Climate Forecast System Reanalysis; Saha et al., 2010) . These data-sets were used over the period January 1980 to December 2015, except for NCEP-CFSR which originally ended in December 2010 and is extended here with the CFSv2 data-set (Climate Forecast System version 2 Saha et al., 2014) from January 2011 to December 2014. Hereafter we use “ERA-I” to refer to ERA-Interim and “CFSR” to refer to the combined NCEP-CFSR reanalyses.

Each reanalysis is available on two vertical grids: the native grid of the underlying atmospheric model (product on “model levels”) and an output grid of constant pressures (product interpolated to “pressure levels”). Our simulations are run on the native model levels in order to account for the different vertical resolution of each reanalysis system and also to avoid any interference from the interpolation methods used to deliver the products on constant pressure levels. All reanalysis systems use the hybrid sigma-pressure vertical coordinate with levels extending from the surface up to ~0.266 hPa (~57 km height) in CFSR, 0.1 hPa (~64 km) in ERA-I and JRA-55, or 0.01 hPa (~78 km) in MERRA and MERRA-2. The reader is referred to Fujiwara et al. (2017) for a comparison of the vertical resolutions of the reanalysis systems.

The forecast models use two different frameworks to discretize their primitive variables on the horizontal plane: MERRA and MERRA-2 solve for mass fluxes on a regular latitude-longitude grid (Lin, 2004) while ERA-I, JRA-55 and CFSR use spectral dynamical cores, i.e., they solve for vorticity and divergence expressed on a spherical harmonics basis (e.g., Krishnamurti et al., 2006). Users of the reanalyses often download velocity fields which are derived from the primitive variables and evaluated on varying regular grids: these may be reduced Gaussian grids (ERA-I and JRA-55), regular Gaussian grids (CFSR) or regular latitude-longitude grids (MERRA and MERRA-2). This pre-processing is described in detail in the next subsection.

We use in all cases the analyses valid at 00 h, 06 h, 12 h and 18 h, i.e., datasets with a 6-h time resolution. The assimilation procedure for MERRA and MERRA-2 uses an iterative predictor–corrector approach, generating two separate sets of reanalysis products designated “ANA” for analysis state and “ASM” for assimilated state (Rienecker et al., 2011). The latter products use a 6h “corrector” forecast centered on the analysis time and an incremental analysis update to apply the previously calculated assimilation increment gradually rather than abruptly at the analysis time (Bloom et al., 1996). Thanks to this procedure, the ASM products have smaller wind imbalances than the ANA products (Fujiwara et al., 2017) hence they are preferable for tracer transport simulations. We used the ASM products in MERRA-2 but could not do so with MERRA where the ASM products are only available on constant pressure levels. Since we aim to evaluate each reanalysis on its native vertical grid, we had to fall back on the ANA product in the case of MERRA.

2.3 Pre-processing of the reanalyses

~~Our offline transport model~~ The BASCOE Transport Model (hereafter BASCOE TM) is used as a tool to perform a fair comparison of advective transport in each reanalysis data-set, using their native vertical grids but a common, low-resolution latitude-longitude grid. It requires on input the surface pressure and horizontal velocity on a so-called Arakawa C-grid, i.e., the zonal wind u must be staggered in longitude and the meridional wind v must be staggered in latitude. As indicated by its name, the FFSL algorithm evaluates internally the corresponding mass fluxes and derives the vertical winds (w) from mass conservation. Hence the reanalysis datasets must be carefully pre-processed from spectral or high-resolution gridded fields to the low-resolution C-grid. We have paid special attention to this pre-processing of the reanalyses to make sure that the different types of wind fields are expressed in a consistent manner for our transport algorithm.

Due to its assimilation procedure, the early ERA-40 reanalysis contained large dynamical imbalances which deteriorated the Brewer-Dobson circulation through excessive upward motion in the tropics and excessive transport from the tropics to the mid-latitudes (Meijer et al., 2004; Monge-Sanz et al., 2007). Pawson et al. (2007) described a similar issue with MERRA and proposed to use time-averaged input wind fields in order to remove these imbalances, but this approach is available only for MERRA and MERRA-2. ~~Here we use~~ To filter out such dynamical imbalances, BASCOE uses a pre-processor ~~; originally developed by Segers et al. (2002) for analyses made at ECMWF, to filter out such dynamical imbalances which was originally developed only for the analyses computed by the European Centre for Medium-Range Weather Forecast (ECMWF) including ERA-I (Segers et al., 2002; Bregman et al., 2003)~~. Using the primitive variables of spectral dynamical cores, i.e., the vorticity and divergence expressed on a spherical harmonics basis, this pre-processor evaluates the zonal and meridional winds on a regular latitude-longitude grid while correcting for the small inconsistencies in the pressure tendency compared with the

divergence fields. This correction ensures consistent mass fields even in the presence of spurious surface pressure increments which may be caused by data assimilation.

Our pre-processing for the five reanalysis systems is based on this algorithm, with a preliminary derivation of the spherical harmonics coefficients of vorticity, divergence and surface pressure for the reanalyses other than ERA-I. In all cases these spectral coefficients are truncated at ~~the wavelength-number~~ wavenumber 47 to avoid aliasing on the 2°x2.5° target grid (Krishnamurti et al., 2006, section 7.4).

2.4 Comparison of Age of Air output with another model

Figure 1 compares the results of the BASCOE ~~CTM~~ TM driven by ERA-I with those by a reference Eulerian model, using the standard layout of zonal means at 20 km height and at equatorial, middle and polar latitudes (e.g., Waugh and Hall, 2002). Both models transport idealized tracers which increase linearly at the ~~surface-tropical tropopause~~ and are driven during 20 years by repeating reanalyses of the year 2000. The reference model is TOMCAT, driven by ERA-I analyses with 6-hourly updates. At 20 km height we use the results published by Dee et al. (2011, Fig. 28) while the vertical profiles are those published by Monge-Sanz et al. (2012, Fig. 1). Some observational context is provided with in-situ observations of SF₆ and CO₂ (Hall et al., 1999). ~~As done in recent intercomparisons of climate models (Neu et al., 2010; Chipperfield et al., 2014), the AoA value at the equatorial tropopause has been subtracted from the fields in order to exclude the transit time from the surface to the tropopause.~~

Very good agreement is obtained between TOMCAT and BASCOE ~~CTM~~ TM. At 20 km height the results are nearly identical except in the ~~southern hemisphere~~ Southern Hemisphere where TOMCAT delivers a slightly weaker latitudinal gradient, resulting in a difference of around 0.5 years above the South Pole between both models. All three vertical profiles show that TOMCAT delivers slightly weaker vertical gradients in the lower stratosphere than the BASCOE ~~CTM~~ TM. This results in younger mid-stratospheric AoA by TOMCAT, but here also the largest difference does not exceed 0.5 years (latitude 5°S, height 45 km).

3 Intercomparison of AoA values

Time-varying distributions of AoA were derived from each reanalysis for the whole period 1980-2015. The initial conditions were obtained from twenty-year spin-up runs simulating the 1960-1980 period with repeating reanalyses of the year 1980. The importance of the initialization procedure was evaluated with an alternative set of transport experiments starting in 1981 from forty-year spin-up runs driven by repeating reanalyses of the year 1981. While the initial AoA could be significantly different depending on the initialization procedure (up to 15% difference in 1981 in the case of CFSR), by 1989 these differences were smaller than 1% at all latitudes and pressure levels for each reanalysis (not shown). Hence the five AoA datasets are studied only over the period 1989-2015.

For the sake of convenience the results of each simulation will be designated by its driving reanalysis but the reader is reminded that all results presented here are obtained indirectly through an offline and kinematic transport model. The outcome of the intercomparison could have been different if the AoA had been computed directly in each reanalysis system.

3.1 Mean distribution in 2002–2007

5 The AoA distributions are first averaged over the period 2002–2007 in order to remove seasonal and quasi-biennial oscillations and also to allow comparisons with the distribution most recently derived from MIPAS observations of SF₆ (Kovács et al., 2017).

The global distribution of AoA is first compared with latitude-pressure cross-sections. The ERA-I reanalysis is taken as reference because it delivers intermediate values and has been used in AoA studies with several other CTMs transport models (see, e.g., Diallo et al., 2012; Monge-Sanz et al., 2012; Konopka et al., 2015). Figure 2 shows the latitude-height cross-sections of AoA for the period 2002–2007, with a noticeable hemispheric asymmetry: as expected, the latitudinal gradient is significantly stronger in the southern mid-latitudes and polar regions than in the ~~northern hemisphere~~ Northern Hemisphere, and old air masses reach much lower altitudes above the Antarctic than above the Arctic (e.g., the 5-year isoline starts at 50 hPa above the South Pole and ends at 20 hPa above the North Pole). This is qualitatively confirmed by AoA derived from MIPAS
15 observations of SF₆ for the same period (Kovács et al., 2017, Fig. 7d).

The four other reanalyses deliver quite noticeably different distributions of AoA (Fig. 3). One can distinguish JRA-55 and CFSR as the “younger reanalyses” with AoA not exceeding 5 years in the polar upper stratosphere; MERRA as the “older reanalysis” with maximum AoA values as large as 6.5 years; and ERA-I with intermediate results (5.8 years in the same regions). MERRA-2 is a special case, with upper stratospheric values similar to those reached by ERA-I but quite different
20 latitudinal gradients. The hemispheric asymmetry is more pronounced with ERA-I than with any other reanalysis, e.g., the 3 and 4-year isolines (JRA-55 and CFSR, respectively) or the 5-year isoline (MERRA-2 and MERRA) reach nearly the same level above the North Pole than above the South Pole. MERRA-2 stands out in the middle stratosphere with nearly vertical isolines, i.e., very small vertical gradients which are not supported by MIPAS observations (Haenel et al., 2015; Kovács et al., 2017).

25 While this qualitative comparison of the AoA distributions points to different gradients in the mid-latitudes and polar regions, the relative differences with respect to ERA-I are largest in the tropical lower stratosphere (bottom row of Fig. 3). Hence we focus on this region and its differences with the midlatitudes. Figure 4, inspired by the AoA intercomparisons in CCMs (Neu et al., 2010; Chipperfield et al., 2014), shows the intercomparison of ~~AOA~~ AoA zonal means at 50 hPa, at tropical and northern midlatitudes, and the latitude gradient AoA difference between these two latitude bands. ~~This layout is inspired by the~~
30 ~~AoA intercomparisons in GCCMs (Neu et al., 2010; Chipperfield et al., 2014), including the subtraction of the AoA values at the equatorial tropopause.~~

The intercomparison at 50 hPa (Fig. 4a) shows again the important disagreements between the five model simulations. JRA-55 yields the youngest AoA at all latitudes with values ranging from 0.8 years at the equator to 3.6 years at the South Pole, while MERRA and MERRA-2 yield the oldest AoA with 1.6 years at the equator and around 5 years at the South Pole.

CFSR and ERA-I yield intermediate results with nearly identical values in the northern extratropics but different latitude gradients ~~in-between~~ the tropics and ~~southern hemisphere~~ Southern Hemisphere. The sole simulation to deliver a minimum AoA in the southern tropics is driven by CFSR (~~which yields the minimum AoA~~ at 6°S). In the other simulations this minimum is either exactly at the equator (JRA-55, MERRA) or slightly north of the equator (ERA-I, MERRA-2). In the ~~southern~~ 5 Southern Hemisphere CFSR results in AoA nearly as young as JRA-55 while ERA-I reaches larger values which are very close to the observations. Overall, the spread between the five simulations at 50 hPa is larger than the 1σ observational uncertainties in the tropics, and nearly as large in the extratropics. ~~This spread~~ Since the reanalyses are constrained by very similar satellite datasets, they could have been expected to deliver more similar AoA than an intercomparison of unconstrained climate models. Yet we note that the spread shown on Fig. 4a is as large as in an intercomparison of ~~7-GCCMs~~ 7-GCCMs (Chipperfield et al., 2014, Fig. 2), ~~which is a remarkable result considering the use of reanalyses rather than unconstrained climate models~~ seven CCMs (Chipperfield et al., 2014, Fig. 2).

The vertical profiles of AoA (Fig. 4b and 4c) ~~confirms~~ confirm that this large spread and general hierarchy of AoA (youngest with JRA-55, oldest with MERRA and MERRA-2) are found at all stratospheric levels. In the northern midlatitudes ~~;(35°N–45°N, Fig. 4c)~~ MERRA-2 stands out with vertical gradients which are ~~significantly~~ larger in the lower stratosphere but ~~significantly~~ smaller in the upper stratosphere than in all other reanalyses. While the intermediate values by ERA-I and CFSR agree well with observations in the tropics ~~;(10°S–10°N, Fig. 4b)~~, this is not the case in the northern midlatitudes where only MERRA and MERRA-2 deliver AoA as old as the observations.

The AoA ~~difference~~ differences between the tropics and mid-latitudes (~~Fig. 4d) is~~ are directly related to the inverse of the tropical upwelling velocity and ~~is~~ independent of quasi-horizontal mixing: a smaller AoA latitudinal gradient indicates faster tropical ascent (Neu and Plumb, 1999; Linz et al., 2016). ~~These “latitudinal gradients of AoA” were used in several CCM intercomparisons (Neu et al., 2010; Chipperfield et al., 2014).~~ Figure 4d shows this diagnostic for the five reanalyses, i.e. the ~~differences between the AoA profiles on Fig. 4c and Fig. 4b~~. Except for MERRA-2, the ~~AoA latitudinal gradients~~ profiles of AoA differences delivered by the four other reanalyses agree much more closely than the AoA profiles themselves, at least during the 2002–2007 period. The spread of AoA differences between the four reanalyses reaches a maximum of 0.2 years at 30 hPa, much tighter than the spread of 0.8 years ~~(after removal of one outlier)~~ in the corresponding intercomparison of ~~6-GCCMs~~ six CCMs (Chipperfield et al., 2014, Fig. 3c). While there is good agreement with the observation-derived ~~latitudinal gradient~~ AoA differences below 60 hPa and at 10 hPa, these four reanalyses significantly underestimate it at intermediate pressure levels. This indicates an overestimation of the tropical upwelling obtained with ERA-I, CFSR, JRA-55 and MERRA in the lower stratosphere. MERRA and MERRA-2 yields an outlying vertical profile of yield larger AoA at northern midlatitudes ~~;(35°N–45°N, Fig. 4c)~~ resulting in a latitudinal gradient which is an outlier as well. than the three other reanalyses. In the case of MERRA-2 this results in a profile of AoA differences which are significantly larger than the profiles obtained with the four other reanalyses but agrees much better with the profile derived from the observations. Hence MERRA-2 apparently underestimates the tropical upwelling in the lowermost stratosphere (100–60 hPa), agrees well with better with the observations at 50 hPa ~~and joins the results of the~~ than the four other reanalyses, and is in accordance with them at higher levels.

3.2 Time evolution and absence of volcanic impact

The Pinatubo eruption, which started on 15 June 1991, is expected to have had a significant impact on AoA (Muthers et al., 2016; Diallo et al., 2017). The assimilation of satellite radiance measurements by the Advanced Microwave Sounding Unit (AMSU) started in 1998 (on 1 August in ERA-I and JRA-55 and 1 November in CFSR, MERRA and MERRA-2) and was repeatedly shown to have a important influence on their description of the stratospheric dynamics (e.g., Simmons et al., 2014; Kawatani et al., 2016; Long et al., 2017). Hence we repeat the latitudinal gradient diagnostic but for the period 1992–1997, i.e., after the Pinatubo eruption and before the ingestion of AMSU radiances (Fig. 5). The general outcome is the same as during the later period: the tropical ascent is too fast with all reanalyses except with MERRA-2. Yet MERRA-2 provides a better match with the observations during this earlier period, and the four other reanalyses do not agree as closely.

Figure 6 shows the globally-averaged time evolution of simulated AoA according to the five reanalyses, from 1989 until 2015 at 50 hPa in the midlatitudes. The results are smoothed with a one-year running mean in order to highlight the long-term trends. The overall hierarchy of ages shown on previous figures for year 2002–2007 holds for the whole 1989–2015 period: MERRA and MERRA-2 deliver the oldest AoA, JRA-55 and CFSR the youngest. While MERRA and MERRA-2 agree well in the southern-hemisphere Southern Hemisphere, this is not the case in the northern-hemisphere Northern Hemisphere where MERRA-2 starts with much older values. A rapid decrease of MERRA-2 values during the 1990's 1990s allows these two datasets to reach better agreement after 1998, i.e., the beginning of AMSU assimilation. The possible causes for this apparently anomalous behavior of MERRA-2 will be discussed in section 5. The MERRA output in the northern-hemisphere Northern Hemisphere delivers seasonal cycles with much larger amplitudes than those obtained from all other reanalyses. This will be investigated in the next section.

The Pinatubo eruption does not appear to have any impact of on the simulated AoA at 50 hPa except with MERRA-2 which shows an increase in the southern midlatitudes. The same time series for the tropical latitude band (30°S–30°N) does not show any impact of the Pinatubo eruption either (not shown). This absence of volcanic impact in the other reanalyses is even more evident in a deseasonalized time series of the extra-polar lower stratosphere (Fig. 7). This diagnostic is inspired by Diallo et al. (2017) who showed a significant impact of the Pinatubo eruption on AoA using ERA-I and JRA-55 but with another offline transport model. Since our results apparently contradict this finding, this issue will also be further discussed in section 5.

Figure 8 displays time series of AoA in the middle stratosphere (mean values between 30 hPa and 5 hPa). The intercomparison in the southern-hemisphere left plot compares the model results in the Northern Hemisphere with balloon observations collected since the 1970s (Engel et al., 2017) where the derivation of AoA uses the surface as reference and the outer error bars denote the overall uncertainty of the mean-age value including an assessment of the representativeness of a single profile (Engel et al., 2009). To allow a consistent comparison the solid lines in Fig. 8 show modeled AoA using the surface as reference, i.e. AoA evaluated from a tracer which uses as boundary condition a global constant increasing linearly with time at the surface. This boundary condition is propagated to the free troposphere through vertical diffusion with a coefficient K_{zz} decreasing from an arbitrary value of $10 \text{ m}^2 \text{ s}^{-1}$ at the surface to zero at the pressure level halfway between the surface and the tropopause. Figure 8b compares the resulting time series in the Tropics with the usual calculation of AoA using the

tropical tropopause as reference (dashed lines). The differences between the two calculations represent the transit times from the surface to the tropical tropopause, are nearly independent of the simulated year and range between three months (with ERA-I or JRA-55) and six months (with MERRA). These values are close to the longest transit times reported in a recent intercomparison of global models (Krol et al., 2018) indicating a slow transport from the surface to the tropical tropopause which we attribute to the omission of deep convective transport in our model. While the surface-based model AoA (solid lines in Fig. 8) may be slightly overestimated, these biases have no significant inter-annual variations and do not hinder the intercomparison between reanalyses.

The spread between the five simulations is as large as the observational uncertainties, highlighting again the importance of the disagreements between the five reanalyses. In the northern mid-latitudes (Fig. 8a) no reanalysis delivers any change larger than half a year over the whole period 1989-2015 except for MERRA-2 which indicates a large decrease of 0.8 years, but this decrease starts from values much larger than the observations and happens mostly before 2000. ERA-I delivers a weakly positive trend over the period 1989-2015 and we will assess in section 4.3 that this trend in the model results is significant. While the overall trend simulated with ERA-I is in agreement with the balloon observations, this comparison should be considered with great caution because the sign of the AoA trend is not significant in the observations (Engel et al., 2009, 2017) and modeled trends over periods as long as 30 years are often not significant when the ideal tracer is sampled like the available observations of stratospheric tracers (Garcia et al., 2011).

The intercomparison in the Southern Hemisphere (Fig. 8c) also shows large disagreements between the long-term trends among the five reanalyses. MERRA and MERRA-2 values decrease quickly until 1995 and increase after 2007 while ERA-I values follow an opposite pattern. The long-term evolution of AoA in this region is completely different with from JRA-55 (gradual decrease during until 2002 followed by a stabilization) and differs yet again with from CFSR (no apparent trend before 1997 and rapid increase during 1997-2003).

The thin lines allow a qualitative comparison of faster variations. The seasonal signal dominates in all cases, with similar phases: AoA is oldest in fall and youngest in spring. The seasonal amplitudes are very dependent on the input reanalysis and on the considered year, so their detailed analysis is deferred to the next section. Yet we note already that some reanalyses exhibit a strong modulation of the seasonal cycle by the Quasi-Biennial Oscillation (QBO; for a general review see Baldwin et al., 2001) while others do not. This can be seen very clearly during the period 2005-2009 when the seasonal amplitudes of AoA by ERA-I and MERRA are approximately twice smaller as small during the easterly phase of the QBO (i.e., in 2006 and 2008) than during the westerly phase (i.e., in 2005, 2007, 2009). This modulation of the seasonal variations is weaker in the MERRA-2 and JRA-55 datasets and absent from the CFSR dataset.

~~The right plot in Fig. 8 compares the model results in the northern hemisphere with balloon observations collected since the 1970's (Engel et al., 2017), where the outer error bars denote the overall uncertainty of the mean age value including an assessment of the representativeness of a single profile (Engel et al., 2009). The spread between the five simulations is as large as the observational uncertainties, highlighting again the importance of the disagreements between the five reanalyses. ERA-I delivers a weakly positive trend over the period 1989-2015, apparently in agreement with the balloon observations, but Engel et al. (2009, 2017) showed that the sign of this observational trend is not significant. More importantly, the AoA from the~~

BASCOE simulation with ERA-I does not show any overall trend after 2000, unless one arbitrarily ends the period in 2010. No reanalysis delivers any change larger than half a year over the whole period 1989-2015 except for MERRA-2 which indicates a large decrease of 0.8 years, but this decrease starts from values much larger than the observations and happens mostly before 2000.

5 4 Analysis of temporal variations

We now perform a quantitative investigation of the temporal variations in order to derive the amplitudes of periodic variations and the linear trends of AoA at all latitudes and pressure levels, including their uncertainties.

4.1 Methodology

Vigouroux et al. (2015) used a multiple linear regression model to study the trends of ozone total columns and vertical distribution at several ground-based stations. Here we apply the same tool to $A(t)$, the monthly zonal means of AoA as a function of time, latitude and pressure (after interpolation to a constant log-pressure grid with 2km increments). The multiple linear regression model is expressed as:

$$A(t) = A_0 + A_1 \cdot t + S(t) + Q(t) + \epsilon(t) \quad (1)$$

where t is time, A_0 is the baseline value, A_1 is the annual trend of AoA and $\epsilon(t)$ represents the residuals. The term $S(t)$ describes the seasonal variations in $A(t)$:

$$S(t) = S_1 \cdot \cos(2\pi t/12) + S_2 \cdot \sin(2\pi t/12) + S_3 \cdot \cos(4\pi t/12) + S_4 \cdot \sin(4\pi t/12) \quad (2)$$

where the coefficients S_1 to S_4 describe the seasonal cycle. The term $Q(t)$ describes the variations due to the QBO and its seasonal modulations:

$$Q(t) = Q_{10}(t) \cdot [Q_1 + Q_2 \cdot \cos(2\pi t/12) + Q_3 \cdot \sin(2\pi t/12) + Q_4 \cdot \cos(4\pi t/12) + Q_5 \cdot \sin(4\pi t/12)] + Q_{30}(t) \cdot [Q_6 + Q_7 \cdot \cos(2\pi t/12) + Q_8 \cdot \sin(2\pi t/12) + Q_9 \cdot \cos(4\pi t/12) + Q_{10} \cdot \sin(4\pi t/12)] \quad (3)$$

where the explanatory variables $Q_{10}(t)$ and $Q_{30}(t)$ are the zonal winds observed above Singapore at 10hPa and 30hPa (data from the FU Berlin: <http://www.geo.fu-berlin.de/en/met/ag/strat/produkte/qbo/index.html>) and Q_1 to Q_{10} are the coefficients associated to these two proxies, including their seasonal dependence.

The uncertainties arising from the fit are [calculated for the 95% confidence interval and](#) corrected for auto-correlation in the residuals (~~Eqs. 3,4 and 6 in Santer et al., 2000~~) ([Eqs. 3, 4 and 6 in Santer et al., 2000](#)). Preliminary tests also included

additional terms to account for the El Niño-Southern Oscillation (ENSO), the 11-year solar cycle and volcanic forcings but it was found that these terms do not impact significantly the linear trends nor the amplitudes of seasonal and quasi-biennial oscillations. Hence they were removed from the regression model in order to avoid any over-fitting of the data and to ease the interpretation of the results.

5 An important goal of this analysis is the determination of linear trends. As seen in Fig. 6 and 8, such trends depend closely on the considered time period. Hence the regression model was applied not only to the whole simulation period (~~1989–2015~~1989–2015) but also to an ~~"early period"~~ (~~1989–2001~~"early period" (1989–2001)) and a ~~"recent period"~~ (~~2002–2015~~"recent period" (2002–2015)) which starts after the assimilation of AMSU and on the same year as the MIPAS mission (Stiller et al., 2008, 2012).

10 4.2 Amplitudes of the seasonal cycle and Quasi-Biennial Oscillation

The amplitude of the seasonal variations is approximated by the difference between the maximum and minimum values reached by the term $S(t)$ in the linear regression model. Figure 9 shows the dependence of this approximated amplitude with respect to pressure in ~~the polar regions and midlatitudes. The five latitude bands for the period 2002–2015 (the results are similar for the period 1989–2001). The results with ERA-I are in agreement with an earlier modeling study (Diallo et al., 2012, Fig. 9). The~~
 15 vertical structure agrees broadly across all five reanalyses ~~and in all four regions in the extratropics~~ with maximum amplitudes in the lower stratosphere (around 100 hPa), except above the South Pole where the amplitudes are maximum in the middle stratosphere (10–30 hPa). ~~The ERA-I results agree with an earlier modeling study which also found that seasonal amplitudes are maximum in the lower stratospheric levels of the midlatitudes and North Pole (Diallo et al., 2012, Fig. 9).~~

MERRA and MERRA-2 stand out with larger amplitudes in the lower stratosphere, resulting above the South Pole in a
 20 ~~second~~secondary maximum which is not found by the three other reanalyses. One may argue that ~~these the~~ larger seasonal amplitudes of MERRA and MERRA-2 are a direct consequence of ~~the their~~ larger annual means (see Fig. 3) but this is not supported by the agreement of JRA-55 and ~~(after 2001)~~ CFSR with ERA-I despite their significantly younger annual means. ~~MERRA and MERRA-2 also deliver very different amplitudes depending on the period used for the analysis, with seasonal amplitudes significantly larger during 1985–2001 than for the recent period. An opposite dependence is noted with CFSR~~
 25 ~~above the South Pole, where after 2001 the seasonal amplitudes became much larger and closer to those delivered by the other reanalyses. The considered period has a comparatively much smaller impact on the seasonal amplitudes in JRA-55 and~~ In the Tropics ERA-I stands out with larger amplitudes in the upper stratosphere (around 5 hPa) and MERRA-2 with larger amplitudes in the lower stratosphere (around 50 hPa) while the three other reanalyses are in good agreement.

We now investigate the differences in the QBO among all reanalyses. Kawatani et al. (2016) have compared the monthly-
 30 mean zonal wind in the equatorial stratosphere ~~between~~among reanalyses and found that their degree of disagreement depends on latitude, longitude, height, and the phase of the QBO. They also noted a tendency for the agreement to be best near the longitude of Singapore, suggesting that the Singapore observations act as a strong constraint on all the reanalyses.

Here we perform an intercomparison of the amplitude of the QBO signal (in years) in each reanalysis. We approximate it again as the difference between the maximum and minimum values reached by the term $Q(t)$ in the linear regression model. Our

results for ERA-I show that the QBO amplitude is largest in the subtropics around 30 hPa (not shown), which confirms again the results of Diallo et al. (2012). ~~Fig. Figure~~ 10 compares the results at this pressure level. Except for CFSR, the latitudinal dependence is similar in all reanalyses: the approximated QBO amplitude reaches maximum values around 15 degrees latitude in both hemispheres and presents a marked minimum around the equator. Outside of the equatorial region, the QBO amplitudes by JRA-55 are significantly smaller than by ERA-I, MERRA and MERRA-2. The amplitudes computed from CFSR show no clear structure in the ~~southern hemisphere and reach unexpectedly~~ Southern Hemisphere and reach large values at the North Pole.

~~Overall, the reanalyses have large (up to a factor of 2) disagreements with respect to the seasonal amplitudes in the polar regions and QBO amplitudes in the Tropics. The results for MERRA, MERRA-2 and CFSR have strong (up to 50%) dependencies on the considered time period. Such disagreements and dependencies could not be expected from inspection of the native dynamic variables contained in the reanalyses, neither for the seasonal amplitudes of polar stratospheric temperatures nor for the QBO zonal wind anomalies at 10 hPa (see, e.g., Long et al., 2017, Fig. 11 and 10, respectively).~~

4.3 Linear trends

It is difficult to infer changes in the BDC on the basis of AoA trends over periods shorter than several decades. Even in models where an ideal, linearly increasing artificial tracer is used, one has to rely on zonal-mean results over long periods to obtain trends that are clearly statistically significant (Garcia et al., 2011) . The statement that is often made that climate models simulate a decreasing age throughout the stratosphere only applies over long time periods and is not necessarily the case for the past 25 years, when most tracer measurements were taken (Garfinkel et al., 2017) . For example, the analysis of a 1700 year simulation showed that it takes around 30 years for a modeled BDC trend to emerge from the noise of natural climate variability (assuming a 2%/decade trend in the BDC; Hardiman et al., 2017) .

While linear trends of AoA over shorter periods may represent transient changes due to climate variability, such changes over timescales which are intermediate between the QBO and the multidecadal scales are still relevant to the study of stratospheric dynamics. Current research on AoA trends has largely focused on a dipole-like latitudinal structure for the period 2002-2012, which was first derived from satellite observation of SF₆ by the MIPAS instrument (Stiller et al., 2012). This structure of trends shows AoA decreasing in the Southern Hemisphere but ~~unexpectedly~~ increasing in the Northern Hemisphere ~~and could be due to a~~ which was used to explain a recent increase of stratospheric HCl in the Northern Hemisphere (Mahieu et al., 2014) and interpreted as the consequence of a southward shift of the subtropical transport barriers (Stiller et al., 2017).

The ERA-I reanalysis supports a dipole-like latitudinal structure of AoA trends, at least since 2002. Haenel et al. (2015, hereafter H2015) derived AoA trends from the distribution of SF₆ over the period 2002-2012, using MIPAS observations and a CCM nudged towards ERA-I below 1 hPa, and found a good agreement for the signs, range and latitudinal structure of AoA trends (see ~~FigFigs~~ 6 and 10 in H2015). Here we aim to verify our methodology through a comparison of our results with H2015, to check the consistency of AoA trends derived from the four other reanalyses, and to explore the latitudinal structure of AoA trends for periods starting earlier than 2002.

The linear trend is represented by A_1 in the multiple regression linear model (Eq. 1). It is expressed in years per decade (yr dec^{-1}) and is deemed significant at a given grid point if its absolute value is larger than its ~~standard-error~~uncertainty (as defined in section 4.1). Figure 11 presents the ERA-I trends during the period 2002-2012 in order to compare with H2015. In the polar regions, H2015 showed large and positive trends while they are insignificant according to our model (Fig. 11).

5 This disagreement can be attributed to different approaches: here we study the true age of air using a theoretical tracer with no losses, while H2015 evaluated the apparent mean age of air taking into account the mesospheric sink of SF_6 which has the largest impact in the polar regions (Reddmann et al., 2001). Outside of the polar regions, Figure 11 shows good agreement with both observational and modeling results in H2015, including with respect to the significance of the trends: in the 30-60 hPa (approx. 25-20 km) layer the trends are significant at all extra-tropical latitudes, negative in the ~~southern-hemisphere~~Southern Hemisphere and positive in the ~~northern-hemisphere~~Northern Hemisphere. They reach -0.6 years per decade in the Southern Tropics and close to 0.5 years per decade in the Northern Tropics. Our results also agree well with those obtained by a diabatic model driven by ERA-I over the same period (Ploeger et al., 2015a).

Figure 12 compares the latitude-pressure distributions of AoA trends across all five reanalyses and for the early (1989-2001), recent (2002-2015) and overall periods (1989-2015). It is important to note that the trends over the early and overall periods
15 should be considered with caution since there was little data to constrain the stratospheric winds until 1998 (see the discussion in the next section). The AoA trends derived from ERA-I wind fields during the early period (Fig. 12, upper left) ~~show-unexpected growth-grow~~ in both hemispheres ~~,except-in~~except for the northern lowermost stratosphere. During the recent period, the dipole structure derived from ERA-I (Fig. 12, upper middle) is similar ~~but-much-less-clear~~ than over the slightly shorter period 2002-2012 (Fig. 11),~~with-weaker-~~. The increases in the ~~northern-hemisphere-which-remain-significant-only-in-the-polar-lower stratosphere~~.The-Northern Hemisphere become weaker but remain significant at all latitudes, although at fewer grid points. The maximum trend is located at 24°N and 25 hPa where it slightly exceeds 0.3 ± 0.2 years per decade. The extension of this trend analysis for the overall period (Fig. 12, upper right) shows a dipole structure with negative but mostly insignificant trends in the ~~southern-hemisphere~~Southern Hemisphere; positive trends in the northern middle stratosphere which mostly corresponds to the region with positive trends during the 1989-2001 period; and significantly negative trends in the lowermost stratosphere at all
25 extra-polar latitudes. ~~Diallo et al. (2012) used adiabatic transport model driven by-~~The same plot also shows that the positive trend which had been inferred visually for the northern mid-latitudes of the middle stratosphere (Fig. 8a) is significant. Our ERA-I and found for the period 1989-2010 results for the overall period partly contradict those obtained by diabatic models which use not only the wind fields from ERA-I but also its heating rates (Diallo et al., 2012; Ploeger et al., 2015a) . Looking at slightly shorter periods of two decades (1989–2010 for the former and 1990–2013 for the latter), these papers reported
30 negative AoA trends ~~in-the-lower-stratosphere-and-positive-trends-in-the-mid-stratosphere~~for both hemispheres below 28km altitude. Diallo et al. (2012) also looked at the middle stratosphere where positive trends were found at all latitudes, suggesting that the shallow and deep Brewer-Dobson circulations may evolve in opposite directions.

~~Our transport model, using only wind fields and surface pressure from ERA-I, confirms this finding.~~

Comparing the results obtained with ERA-I with those from other reanalyses, one notes immediately general agreement
35 between ERA-I and CFSR on one hand (Fig. 12, first and second row) and opposite trends in JRA-55, MERRA and MERRA-2

(third to fifth row). The agreement between multidecadal trends in ERA-I and CFSR may be related to their closeness in AoA distribution and spatial gradients (section 3.1). For all reanalyses except ERA-I, the trends for the overall period (1989–2015: Fig. 12, right column) appear dominated by the results from the early period which are subject to caution.

To summarize, the signs of the trends depend strongly on the input reanalysis and on the considered period with values above 10 hPa varying between approximately -0.4 and 0.4 years per decade. JRA-55, MERRA and MERRA-2 indicate an AoA increasing globally over ~~2002–2015~~2002–2015, except in the lowermost stratosphere; while ERA-I and CFSR indicate ~~exactly~~ the opposite (Fig. 12, middle column). These trends are significant only in specific regions of the stratosphere, and the regions of significance vary depending on the considered reanalysis. ERA-I stands out as the only reanalysis yielding a dipole structure of AoA trends for the period ~~2002–2015~~2002–2015, although one may note that in the lower stratosphere, the AoA growth derived for this period from MERRA and MERRA-2 (Fig. 12, middle column, fourth and fifth row) is faster in the ~~northern hemisphere~~Northern Hemisphere than in the ~~southern hemisphere~~Southern Hemisphere. ~~The most striking result, in this intercomparison of AoA trends for different periods, is the reversal of~~ One notes also a reversal of the trends between the early (~~1989–2001~~1989–2001) and recent (~~2002–2015~~2002–2015) periods. This reversal is found for all five reanalyses and in all regions of the stratosphere but it is difficult to interpret because it goes in opposite directions in ERA-I and CFSR versus JRA-55, MERRA and MERRA-2.

5 Discussion and outlook

~~This~~The present intercomparison reveals large disagreements between the AoA derived from the five reanalyses, both with respect to their values and their linear trends. The spread of AoA at 50 hPa (Fig. 4a) is as large as in an intercomparison of ~~CCM~~CCMs (Chipperfield et al., 2014). An intercomparison of AoA trends during the twenty-first century among ~~6-CCM~~five CCMs shows negative trends in the whole middle atmosphere (about -0.05 years per decade) with no large hemispheric asymmetry (Butchart et al., 2010) while our results for 1989–2015 show faster changes (-0.4 to 0.4 years per decade) with different signs depending on the reanalysis and the stratospheric region. Since these results call for further research, we propose here a summary overview of the possible causes for these disagreements and some venues to attempt their identification.

Many intercomparisons of reanalyses have focused on the instantaneous values or long-term evolution of direct output fields such as temperature or zonal winds (Simmons et al., 2014; Lawrence et al., 2015; Long et al., 2017; Kozubek et al., 2017). These intercomparisons do not find large discrepancies, especially after the introduction of new satellite instruments around year 2000. The large disagreements obtained here may ~~look surprising unless one considers~~ be explained by the lack of wind observations available for assimilation in the tropics, high latitudes and stratosphere (Baker et al., 2014). This deficiency of wind information ~~may explain~~explains the divergences between trajectories obtained with different reanalyses in the lower stratosphere, e.g., in the equatorial region during some phases of the QBO (Podglajen et al., 2014) or above the Antarctic during the vortex break-up season (Hoffmann et al., 2017). Such divergent trajectories could have a significant cumulative impact on the mean Age of Air because it is a time-integrated diagnostic spanning several years.

Since the wind fields are weakly constrained, the causes for the disagreements found here may lie in the differences between the underlying models which were summarized recently in the context of S-RIP (Fujiwara et al., 2017). Let us first look at vertical resolution, which has an important impact on the modeling of lower stratospheric dynamics (Richter et al., 2014). In the lower stratosphere, the vertical resolution of CFSR is finest while the resolution of and ERA-I and JRA-55 is the coarsest, with the resolution of MERRA and MERRA-2 in between (Fujiwara et al., 2017). This has no clear impact on AoA since CFSR and JRA-55 deliver the youngest AoA while the MERRA and MERRA-2 deliver the oldest, with ERA-I results in between. Hence one cannot establish a simple link between vertical resolution and AoA in this intercomparison.

The present intercomparison cannot establish the impact of different horizontal resolutions because it uses a common horizontal grid with a coarse resolution of $2^\circ \times 2.5^\circ$ (see sections 2.1 and 2.3). For example, the intercomparison of AoA distributions (section 3.1) showed that JRA-55 and CFSR yield the weakest latitudinal gradients despite their horizontal grid spacing which is finest among the five reanalyses studied here (see Fujiwara et al., 2017, table 2). Another intercomparison could yield different results if it uses the wind fields in each reanalysis at its original resolution – but this could lead to difficulties in the handling of horizontal diffusion (Jablonowski and Williamson, 2011).

Different parametrizations of gravity wave drag are another possible modeling cause for the disagreements in AoA. ERA-I, JRA-55 and CFSR all neglect non-orographic gravity wave drag (except for CFSv2, i.e., CFSR after 2010) and each uses its own parametrization of orographic gravity wave drag. MERRA and MERRA-2 on the other hand use the same parametrization for orographic gravity wave drag (McFarlane, 1987) and both take non-orographic gravity wave drag into account. ~~While this may be a coincidence, MERRA and MERRA-2 happen to provide much older AoA than the three other reanalyses.~~

Miyazaki et al. (2016) compared the mean-meridional circulations and also the mixing strengths in six reanalyses – including ERA-I and JRA-55 – and also found significant disagreements. Their diagnostics are closely related to AoA since a faster mean-meridional circulation evidently leads to younger AoA and increased mixing corresponds mostly to additional aging of air due to recirculation from the extra-tropics to the Tropics (Garny et al., 2014). For example, the disagreements of linear trends for 1989-2015 (right column in Fig. 12) confirm the finding that ERA-I and JRA-55 have opposite linear trends of tropical upward mass flux for the period 1979-2012, with fluxes increasing at all levels in JRA while in ERA-I they increase only in a shallow layer of the lower troposphere but decrease in the middle stratosphere (Miyazaki et al., 2016, Fig. 11). Similar disagreements have also been reported between the trends of the annual mean tropical upwelling in three reanalyses over the period 1979–2012, with vertical residual velocities (\overline{w}^*) increasing in MERRA and JRA-55 and decreasing in ERA-I (Abalos et al., 2015, Fig. 11) .

MERRA-2 stands out with outlying AoA values during the ~~1990's~~ 1990s. A connection is plausible with its difficulties to represent correctly the QBO before 1995 (Kawatani et al., 2016; Coy et al., 2016). ~~Interestingly,~~ Gelaro et al. (2017) noted on that same year a marked decrease in temperature near 1hPa and associated it with a change in assimilated radiance data. Gelaro et al. (2017) describe three features which are absent from the other reanalysis systems and could also play a role in the description of middle atmosphere dynamics in MERRA-2, contributing to its outlying AoA. With respect to assimilated observations, MERRA-2 is the only reanalysis to assimilate Aura-MLS temperatures, from 2004 onwards and above 5 hPa. While this has an important impact on temperatures in the upper stratosphere and lower mesosphere, it does not seem to have

an impact on the AoA time series in the middle stratosphere (Fig. 8) and cannot explain the large values obtained during the ~~1990's~~1990s. With respect to forward model forcings, MERRA-2 is the only reanalysis which includes a large source of non-orographic gravity wave drag in the tropics (Molod et al., 2015) and realistic aerosol optical depths. This last feature most ~~probably-likely~~ explains the sensitivity of the MERRA-2 AoA at 50 hPa to the Pinatubo eruption, which cannot be seen with any other reanalysis (Fig. 6).

Yet the impact of the Pinatubo eruption on MERRA-2 AoA at 50 hPa cannot be seen in the northern midlatitudes, and in the southern midlatitudes it is not larger than the amplitude of seasonal variations. In section 3.2 we could not find any influence of volcanic aerosols at the global scale (Fig. 7), ~~contrarily-contrary~~ to recent results obtained by Diallo et al. (2017) using the Chemical Lagrangian Model of the Stratosphere (CLaMS) driven by ERA-I and JRA-55. While CLaMS is a Lagrangian transport model ~~and BASCOE-CTM including a mixing parametrization (Konopka et al., 2004) and BASCOE TM~~ a Eulerian transport model, we ~~believe-suggest~~ that these conflicting results are better explained by the different approaches with respect to vertical transport: BASCOE ~~CTM-TM~~ is a kinematic model (see section 2.1) while CLaMS is a diabatic transport model, hence also driven by the heating rates from the reanalysis forecast models (Ploeger et al., 2010, 2015b).

Wright and Fueglistaler (2013) have shown that the heat budgets differ significantly in the ~~Tropical-Tropopause-tropical~~ tropopause layer among the reanalyses, with substantial implications for representations of transport and mixing in this region. Abalos et al. (2015) evaluated the vertical component of the advective BDC in ERA-I, MERRA and JRA-55 and found substantial differences between direct (i.e., kinematic) estimates and indirect estimates derived from the thermodynamic balance (i.e., using diabatic heating rates). These intercomparisons of dynamical diagnostics highlight the need for another intercomparison of AoA using a diabatic transport model, because this approach would also reflect the differences between the diabatic heat budgets of each reanalysis - including the temperature increments from the assimilation of temperature radiances (Diallo et al., 2017).

Future work will also involve the disentangling of the contributions to AoA of the residual circulation, mixing on resolved scales and mixing on unresolved scales (i.e., diffusion) as recently performed with ERA-I (Ploeger et al., 2015a; Dietmüller et al., 2017) and quantitative comparisons with observational data-sets, using both MIPAS observations of SF₆ (Stiller et al., 2012; Haenel et al., 2015) and balloon observations of SF₆ and CO₂ (Ray et al., 2014). Comparisons with long-term records of other long-lived tracers will provide further insight at multidecadal scales. A recent study by Douglass et al. (2017) explained that the relationship between AoA and the fractional release of such tracers is a stronger test of the realism of simulated transport than the simple comparisons of mean age distributions. This approach seems very promising not only in the context of S-RIP but also for observation-based evaluations of stratospheric transport in global circulation-chemistry models.

6 Summary and conclusions

We have developed a pre-processor to feed a Eulerian and kinematic transport model with any of the available global reanalysis datasets. This has allowed us to compute the mean Age of Air (AoA) in the stratosphere and its evolution from 1985 to 2015,

according to five modern reanalyses: ERA-Interim, JRA-55, MERRA, MERRA-2 and CFSR. Our results compare well with those published previously using other transport models driven by ERA-Interim and MERRA-2.

The five reanalyses deliver very different and diverse results. In the middle and upper stratosphere, MERRA yields the oldest AoA (~5-6 years at mid-latitudes) and JRA-55 the youngest one (~3.5 years). MERRA-2 provides a different distribution of latitudinal and vertical AoA gradients than any other reanalysis, with near-zero vertical gradients in the middle stratosphere which ~~do not seem are not~~ supported by observations. CFSR and ERA-I give the most similar AoA distributions, with the latter providing stronger gradients vertically in the middle stratosphere and latitudinally in the ~~southern hemisphere~~ Southern Hemisphere. The relative differences between ERA-I and the four other reanalyses are largest in the lower tropical stratosphere. Tropical ascent rates have been compared through the difference between AoA in the northern mid-latitudes and in the tropics, showing good agreements between all reanalyses except for MERRA-2 and an overestimation of the upwelling in the tropical lower stratosphere.

The time variations of AoA were studied first through a qualitative analysis of raw time series in the mid-latitudes, then through a fit with a multiple linear regression model. While the linear trends vary considerably depending on the considered period (2002-2012, 2002-2015 or 1985-2015), the general hierarchy of “older” (MERRA, MERRA-2) and “younger” (JRA-55, CFSR) reanalyses holds during the whole 1985-2015 period, with ERA-I keeping intermediate AoA values. The MERRA-2 results stand out again, with an exceptionally large initial AoA in the ~~northern hemisphere~~ Northern Hemisphere which quickly decreases during the ~~1990's~~ 1990s to reach values similar to those in MERRA. A comparison was performed with a time series of balloon observations realized since the ~~1970's~~ 1970s in the northern mid-latitudes where the uncertainties include an evaluation of the sampling error (Engel et al., 2017). The spread between the five simulations is as large as the observational uncertainties, highlighting again the importance of the disagreements between the five reanalyses. ~~The AoA using ERA-I does not show any overall trend after 2000, unless one arbitrarily ends the period in 2010.~~

The amplitudes of seasonal variations agree broadly across all five reanalyses but in the lower stratosphere they are larger in MERRA and MERRA-2 than in the three other reanalyses. The ~~seasonal amplitudes of MERRA-2 decrease significantly during the 2002-2015 period but at the North Pole they remain 50% larger than those of the three other reanalyses. The~~ latitudinal dependence of QBO amplitudes is similar in all five reanalyses except for CFSR which ~~show~~ shows no clear structure in the ~~southern hemisphere~~ Southern Hemisphere.

The linear trends of ERA-I AoA confirm again the dipole structure of the latitude-height distribution of AoA trends as derived from MIPAS observations of SF₆ for the 2002-2012 period (Haenel et al., 2015), with a decrease in the ~~southern hemisphere~~ Southern Hemisphere reaching about -0.6 years per decade, and an increase in the northern lower stratosphere ~~which is significant and not expected from reaching about 0.5 years per decade. The increase in the Northern Hemisphere is significant (at the 95% confidence level) and it is not obtained in multidecadal~~ climate model simulations. Yet the trends derived from ERA-I are shown to closely depend on the considered period. When it is extended to 2002-2015, the positive trends in the ~~northern hemisphere become mostly insignificant and the dipole structure becomes much less pronounced~~ Northern Hemisphere become weaker (about 0.3 years per decade) and they are significant at fewer grid points. A further extension to 1989-2015 shows ~~positive trends which become significant again in the northern middle stratosphere, but the~~ that the negative trends in

the southern middle stratosphere become insignificant. For all five reanalyses the trends over the early period (1989-2001) have opposite signs than over the recent period (2002-2015). Looking only at the recent period which is better constrained by observations, the main outcome is again large disagreements between the reanalyses: JRA-55, MERRA and MERRA-2 provide increasing AoA in the middle stratosphere while CFSR provides a decreasing but mostly insignificant trend. To summarize, the signs of the trends depend strongly on the input reanalysis and on the considered period with values above 10 hPa varying between approximately -0.4 and 0.4 years per decade. Independently of the considered period, no reanalysis other than ERA-I finds any dipole structure in the latitude-height distribution of AoA trends.

~~No~~ Since the wind fields are weakly constrained, the causes for the disagreements found here may lie in the differences between the underlying models. While no obvious cause could be found ~~for these disagreements. The~~, we suggest that the

parametrization of non-orographic gravity wave drag ~~in the underlying dynamical model~~ deserves further investigation, especially in the case of MERRA-2 which has difficulties to represent correctly the QBO before 1995. No global impact of the Pinatubo eruption can be found in our simulations of AoA, ~~contrarily~~ contrary to a recent study which used ERA-I and JRA-55 to drive a diabatic transport model. This highlights the need to repeat the present intercomparison with diabatic transport models because they would reflect directly the significant differences between the heating rates in the reanalyses (Wright and Fueglistaler, 2013). Future work will also focus on quantitative comparisons with AoA derived from MIPAS observations of SF₆; comparisons with the long-term records of other long-lived tracers to provide further insight at multidecadal scales; and disentangling the contributions to AoA of residual circulation, mixing on resolved scales and mixing on unresolved scales.

The main conclusion of this study is the significant diversity in the distribution of mean AoA which we obtain with our transport model, depending on the input reanalysis. This casts doubt on our ability to model accurately the time necessary for variations of greenhouse or ozone-depleting species to propagate from the troposphere to the stratosphere. We have also found large disagreements between the five reanalyses with respect to the long-term trends of age of air. This suggests that with our type of offline transport model, the wind fields in modern reanalyses are not sufficiently constrained by observations to evaluate the actual changes of stratospheric circulation. Yet this conclusion should not be hastily extended to other types of transport models which also use the reanalyses of temperature and heating rates.

Code and data availability. The monthly zonal averages of AoA, as delivered by the BASCOE TM experiments driven by the five input reanalyses, are distributed as an online supplement to this article. The source code of the BASCOE TM, including its tools to pre-process the reanalyses, is available by email request to the corresponding author. The ERA-Interim reanalysis (Dee et al., 2011) is provided by the ECMWF, see <http://www.ecmwf.int/en/forecasts/datasets>. MERRA data (Rienecker et al., 2011) and MERRA-2 data (Gelaro et al., 2017) are provided by the Global Modeling and Assimilation Office at NASA Goddard Space Flight Center through the NASA GES DISC online archive; see <https://disc.gsfc.nasa.gov/information/glossary?keywords=merra>. The CFSR (Saha et al., 2010) and CFSv2 (Saha et al., 2014) reanalyses data were obtained from NOAA NCEP; see <http://cfs.ncep.noaa.gov/>. The JRA-55 reanalysis (Kobayashi et al., 2015) was obtained from the NCAR Research Data Archive; see <https://rda.ucar.edu/datasets/ds628.0/>.

Acknowledgements. We thank the reanalysis centers (ECMWF, NASA GSFC, NOAA NCEP and JMA) for providing their support and data products. We thank Dr. Gabriele Stiller, Dr. Paul Konopka and Dr. Bernard Legras for fruitful discussions during the preliminary steps leading to this study, and Dr. Masatomo Fujiwara for his coordination of the S-RIP. [We would also like to thank the editor and two anonymous reviewers for their valuable comments.](#) Yves Christophe's contribution was partly supported by the European Commission project MACC-II under the EU Seventh Research Framework Programme (contract number 283576). Daniele Minganti's contribution was financially supported by the Fonds de la Recherche Fondamentale Collective through research project ACCROSS (convention PDRT.0040.16). Emmanuel Mahieu is Research Associate with the F.R.S.-FNRS.

References

- Abalos, M., Legras, B., Ploeger, F., and Randel, W. J.: Evaluating the advective Brewer-Dobson circulation in three reanalyses for the period 1979–2012, *Journal of Geophysical Research: Atmospheres*, 120, 7534–7554, doi:10.1002/2015JD023182, <http://dx.doi.org/10.1002/2015JD023182>, 2015JD023182, 2015.
- 5 Andrews, A. E., Boering, K. A., Daube, B. C., Wofsy, S. C., Loewenstein, M., Jost, H., Podolske, J. R., Webster, C. R., Herman, R. L., Scott, D. C., Flesch, G. J., Moyer, E. J., Elkins, J. W., Dutton, G. S., Hurst, D. F., Moore, F. L., Ray, E. A., Romashkin, P. A., and Strahan, S. E.: Mean ages of stratospheric air derived from in situ observations of CO₂, CH₄, and N₂O, *J. Geophys. Res.*, 106, 32 295–32 314, doi:10.1029/2001JD000465, <http://dx.doi.org/10.1029/2001JD000465>, 2001.
- Austin, J. and Li, F.: On the relationship between the strength of the Brewer-Dobson circulation and the age of stratospheric air, *Geophys. Res. Lett.*, 33, n/a–n/a, doi:10.1029/2006GL026867, <http://dx.doi.org/10.1029/2006GL026867>, 117807, 2006.
- 10 Baker, W. E., Atlas, R., Cardinali, C., Clement, A., Emmitt, G. D., Gentry, B. M., Hardesty, R. M., Källén, E., Kavaya, M. J., Langland, R., Ma, Z., Masutani, M., McCarty, W., Pierce, R. B., Pu, Z., Riishojgaard, L. P., Ryan, J., Tucker, S., Weissmann, M., and Yoe, J. G.: Lidar-Measured Wind Profiles: The Missing Link in the Global Observing System, *Bull. Am. Meteorol. Soc.*, 95, 543–564, doi:10.1175/BAMS-D-12-00164.1, <https://doi.org/10.1175/BAMS-D-12-00164.1>, 2014.
- 15 Baldwin, M. P., Gray, L. J., Dunkerton, T. J., Hamilton, K., Haynes, P. H., Randel, W. J., Holton, J. R., Alexander, M. J., Hirota, I., Horinouchi, T., Jones, D. B. A., Kinnersley, J. S., Marquardt, C., Sato, K., and Takahashi, M.: The quasi-biennial oscillation, *Rev. Geophys.*, 39, 179–229, doi:10.1029/1999RG000073, <http://dx.doi.org/10.1029/1999RG000073>, 2001.
- Bengtsson, L. and Shukla, J.: Integration of Space and In Situ Observations to Study Global Climate Change, *Bulletin of the American Meteorological Society*, 69, 1130–1143, doi:10.1175/1520-0477(1988)069<1130:IOSAIS>2.0.CO;2, [https://doi.org/10.1175/1520-0477\(1988\)](https://doi.org/10.1175/1520-0477(1988)069<1130:IOSAIS>2.0.CO;2)
- 20 069<1130:IOSAIS>2.0.CO;2, 1988.
- Bloom, S., Takacs, L., DaSilva, A., and Ledvina, D.: Data assimilation using incremental analysis updates, *Mon. Weather Rev.*, pp. 1256–1271, 1996.
- Bregman, B., Segers, A., Krol, M., Meijer, E., and van Velthoven, P.: On the use of mass-conserving wind fields in chemistry-transport models, *Atmos. Chem. Phys.*, 3, 447–457, doi:10.5194/acp-3-447-2003, <https://www.atmos-chem-phys.net/3/447/2003/>, 2003.
- 25 Butchart, N.: The Brewer-Dobson circulation, *Rev. Geophys.*, 52, 157–184, doi:10.1002/2013RG000448, <http://dx.doi.org/10.1002/2013RG000448>, 2014.
- Butchart, N., Cionni, I., Eyring, V., Shepherd, T. G., Waugh, D. W., Akiyoshi, H., Austin, J., Brühl, C., Chipperfield, M. P., Cordero, E., Dameris, M., Deckert, R., Dhomse, S., Frith, S. M., Garcia, R. R., Gettelman, A., Giorgetta, M. A., Kinnison, D. E., Li, F., Mancini, E., McLandress, C., Pawson, S., Pitari, G., Plummer, D. A., Rozanov, E., Sassi, F., Scinocca, J. F., Shibata, K., Steil, B., and Tian, W.:
- 30 Chemistry–Climate Model Simulations of Twenty-First Century Stratospheric Climate and Circulation Changes, *J. Climate*, 23, 5349–5374, doi:10.1175/2010JCLI3404.1, <https://doi.org/10.1175/2010JCLI3404.1>, 2010.
- Chipperfield, M. P.: New version of the TOMCAT/SLIMCAT off-line chemical transport model: Intercomparison of stratospheric tracer experiments, *Quart. J. R. Meteorol. Soc.*, 132, 1179–1203, doi:10.1256/qj.05.51, <http://dx.doi.org/10.1256/qj.05.51>, 2006.
- Chipperfield, M. P., Liang, Q., Strahan, S. E., Morgenstern, O., Dhomse, S. S., Abraham, N. L., Archibald, A. T., Bekki, S., Braesicke, P., Di Genova, G., Fleming, E. L., Hardiman, S. C., Iachetti, D., Jackman, C. H., Kinnison, D. E., Marchand, M., Pitari, G., Pyle, J. A., Rozanov, E., Stenke, A., and Tummon, F.: Multimodel estimates of atmospheric lifetimes of long-lived ozone-depleting substances: Present and future, *J. Geophys. Res.*, 119, 2555–2573, doi:10.1002/2013JD021097, <http://dx.doi.org/10.1002/2013JD021097>, 2014.

- Coy, L., Wargan, K., Molod, A. M., McCarty, W. R., and Pawson, S.: Structure and Dynamics of the Quasi-Biennial Oscillation in MERRA-2, *Journal of Climate*, 29, 5339–5354, doi:10.1175/JCLI-D-15-0809.1, <https://doi.org/10.1175/JCLI-D-15-0809.1>, 2016.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haim-
5 berger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quart. J. R. Meteorol. Soc.*, 137, 553–597, doi:10.1002/qj.828, <http://dx.doi.org/10.1002/qj.828>, 2011.
- Diallo, M., Legras, B., and Chédin, A.: Age of stratospheric air in the ERA-Interim, *Atmos. Chem. Phys.*, 12, 12 133–12 154, doi:10.5194/acp-12-12133-2012, <http://www.atmos-chem-phys.net/12/12133/2012/>, 2012.
10
- Diallo, M., Ploeger, F., Konopka, P., Birner, T., Müller, R., Riese, M., Garny, H., Legras, B., Ray, E., Berthet, G., and Jegou, F.: Significant Contributions of Volcanic Aerosols to Decadal Changes in the Stratospheric Circulation, *Geophys. Res. Lett.*, pp. n/a–n/a, doi:10.1002/2017GL074662, <http://dx.doi.org/10.1002/2017GL074662>, 2017GL074662, 2017.
- Dietmüller, S., Garny, H., Plöger, F., Jöckel, P., and Cai, D.: Effects of mixing on resolved and unresolved scales on stratospheric age of air, *Atmos. Chem. Phys.*, 17, 7703–7719, doi:10.5194/acp-17-7703-2017, <https://www.atmos-chem-phys.net/17/7703/2017/>, 2017.
15
- Douglass, A. R., Strahan, S. E., Oman, L. D., and Stolarski, R. S.: Multi-decadal records of stratospheric composition and their relationship to stratospheric circulation change, *Atmos. Chem. Phys.*, 17, 12 081–12 096, doi:10.5194/acp-17-12081-2017, <https://www.atmos-chem-phys.net/17/12081/2017/>, 2017.
- Engel, A., Mobius, T., Bonisch, H., , Schmidt, U., Heinz, R., Levin, I., Atlas, E., Aoki, S., Nakazawa, T., Sugawara, S., Moore, F., Hurst, D.,
20 Elkins, J., Schauffler, S., Andrews, A., and Boering, K.: Age of stratospheric air unchanged within uncertainties over the past 30 years, *Nature Geosci.*, 2, 28–31, doi:10.1038/ngeo388, <http://dx.doi.org/10.1038/ngeo388>, 2009.
- Engel, A., Bönisch, H., Ullrich, M., Sitals, R., Membrive, O., Danis, F., and Crevoisier, C.: Mean age of stratospheric air derived from AirCore observations, *Atmos. Chem. Phys.*, 17, 6825–6838, doi:10.5194/acp-17-6825-2017, <http://www.atmos-chem-phys.net/17/6825/2017/>, 2017.
- Errera, Q., Daerden, F., Chabrillat, S., Lambert, J. C., Lahoz, W. A., Viscardy, S., Bonjean, S., and Fonteyn, D.: 4D-Var assimilation of MIPAS chemical observations: ozone and nitrogen dioxide analyses, *Atmos. Chem. Phys.*, 8, 6169–6187, <http://www.atmos-chem-phys.net/8/6169/2008/>, 2008.
25
- Fu, Q., Lin, P., Solomon, S., and Hartmann, D. L.: Observational evidence of strengthening of the Brewer-Dobson circulation since 1980, *J. Geophys. Res.*, 120, 10,214–10,228, doi:10.1002/2015JD023657, <http://dx.doi.org/10.1002/2015JD023657>, 2015.
- Fujiwara, M., Wright, J. S., Manney, G. L., Gray, L. J., Anstey, J., Birner, T., Davis, S., Gerber, E. P., Harvey, V. L., Hegglin, M. I., Homeyer, C. R., Knox, J. A., Krüger, K., Lambert, A., Long, C. S., Martineau, P., Molod, A., Monge-Sanz, B. M., Santee, M. L., Tegtmeier, S., Chabrillat, S., Tan, D. G. H., Jackson, D. R., Polavarapu, S., Compo, G. P., Dragani, R., Ebisuzaki, W., Harada, Y., Kobayashi, C., McCarty, W., Onogi, K., Pawson, S., Simmons, A., Wargan, K., Whitaker, J. S., and Zou, C.-Z.: Introduction to the SPARC Reanalysis Intercomparison Project (S-RIP) and overview of the reanalysis systems, *Atmos. Chem. Phys.*, 17, 1417–1452, doi:10.5194/acp-17-1417-
30 2017, <http://www.atmos-chem-phys.net/17/1417/2017/>, 2017.
- Garcia, R. R., Randel, W. J., and Kinnison, D. E.: On the Determination of Age of Air Trends from Atmospheric Trace Species, *J. Atmos. Sci.*, 68, 139–154, doi:10.1175/2010JAS3527.1, <http://dx.doi.org/10.1175/2010JAS3527.1>, 2011.

- Garfinkel, C. I., Aquila, V., Waugh, D. W., and Oman, L. D.: Time-varying changes in the simulated structure of the Brewer–Dobson Circulation, *Atmospheric Chemistry and Physics*, 17, 1313–1327, doi:10.5194/acp-17-1313-2017, <https://www.atmos-chem-phys.net/17/1313/2017/>, 2017.
- Garny, H., Birner, T., Bönisch, H., and Bunzel, F.: The effects of mixing on age of air, *J. Geophys. Res.*, 119, 7015–7034, doi:10.1002/2013JD021417, <http://dx.doi.org/10.1002/2013JD021417>, 2014.
- 5 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *Journal of Climate*, 30, 5419–5454, doi:10.1175/JCLI-D-16-0758.1, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- 10 Haenel, F. J., Stiller, G. P., von Clarmann, T., Funke, B., Eckert, E., Glatthor, N., Grabowski, U., Kellmann, S., Kiefer, M., Linden, A., and Reddmann, T.: Reassessment of MIPAS age of air trends and variability, *Atmos. Chem. Phys.*, 15, 13 161–13 176, doi:10.5194/acp-15-13161-2015, <http://www.atmos-chem-phys.net/15/13161/2015/>, 2015.
- Hall, T. M., Waugh, D. W., Boering, K. A., and Plumb, R. A.: Evaluation of transport in stratospheric models, *J. Geophys. Res.*, 104, 18 815–18 839, doi:10.1029/1999JD900226, <http://dx.doi.org/10.1029/1999JD900226>, 1999.
- 15 Hardiman, S. C., Lin, P., Scaife, A. A., Dunstone, N. J., and Ren, H.-L.: The influence of dynamical variability on the observed Brewer–Dobson Circulation trend, *Geophys. Res. Lett.*, 44, 2885–2892, doi:10.1002/2017GL072706, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017GL072706>, 2017.
- Hoffmann, L., Hertzog, A., Röbber, T., Stein, O., and Wu, X.: Intercomparison of meteorological analyses and trajectories in the Antarctic lower stratosphere with Concordiasi superpressure balloon observations, *Atmos. Chem. Phys.*, 17, 8045–8061, doi:10.5194/acp-17-8045-2017, <https://www.atmos-chem-phys.net/17/8045/2017/>, 2017.
- 20 Jablonowski, C. and Williamson, D. L.: The Pros and Cons of Diffusion, Filters and Fixers in Atmospheric General Circulation Models, in: *Numerical Techniques for Global Atmospheric Models*, edited by Lauritzen, P. H., Jablonowski, C., Taylor, M. A., and Nair, R. D., vol. 80 of *Lecture Notes in Computational Science and Engineering*, pp. 381–493, Springer, 2011.
- 25 Kawatani, Y., Hamilton, K., Miyazaki, K., Fujiwara, M., and Anstey, J. A.: Representation of the tropical stratospheric zonal wind in global atmospheric reanalyses, *Atmos. Chem. Phys.*, 16, 6681–6699, doi:10.5194/acp-16-6681-2016, <https://www.atmos-chem-phys.net/16/6681/2016/>, 2016.
- Kobayashi, S., Ota, Y., Harada, y., Ebata, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H., Kobayashi, C., Endo, H., Miyaoka, K., and Takahashi, K.: The JRA-55 Reanalysis: General Specifications and Basic Characteristics, *J. Meteorol. Soc. Jpn.*, 93, 5–48, doi:10.2151/jmsj.2015-001, 2015.
- 30 Konopka, P., Steinhorst, H.-M., Grob, J.-U., Günther, G., Müller, R., Elkins, J. W., Jost, H.-J., Richard, E., Schmidt, U., Toon, G., and McKenna, D. S.: Mixing and ozone loss in the 1999–2000 Arctic vortex: Simulations with the three-dimensional Chemical Lagrangian Model of the Stratosphere (CLaMS), *Journal of Geophysical Research: Atmospheres*, 109, doi:10.1029/2003JD003792, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2003JD003792>, 2004.
- 35 Konopka, P., Ploeger, F., Tao, M., Birner, T., and Riese, M.: Hemispheric asymmetries and seasonality of mean age of air in the lower stratosphere: Deep versus shallow branch of the Brewer–Dobson circulation, *J. Geophys. Res.*, 120, 2053–2066, doi:10.1002/2014JD022429, <http://dx.doi.org/10.1002/2014JD022429>, 2014JD022429, 2015.

- Kovács, T., Feng, W., Totterdill, A., Plane, J. M. C., Dhomse, S., Gómez-Martín, J. C., Stiller, G. P., Haenel, F. J., Smith, C., Forster, P. M., García, R. R., Marsh, D. R., and Chipperfield, M. P.: Determination of the atmospheric lifetime and global warming potential of sulfur hexafluoride using a three-dimensional model, *Atmos. Chem. Phys.*, 17, 883–898, doi:10.5194/acp-17-883-2017, 2017.
- Kozubek, M., Krizan, P., and Lastovicka, J.: Comparison of the long-term trends in stratospheric dynamics of four reanalyses, *Annales Geophysicae*, 35, 279–294, doi:10.5194/angeo-35-279-2017, <http://www.ann-geophys.net/35/279/2017/>, 2017.
- Krishnamurti, T. N., Bedi, H. S., Hardiker, V. M., and Ramaswamy, L., eds.: *An Introduction to Global Spectral Modeling*, Springer Science, 2 edn., 2006.
- Krol, M., de Bruine, M., Killaars, L., Ouwersloot, H., Pozzer, A., Yin, Y., Chevallier, F., Bousquet, P., Patra, P., Belikov, D., Maksyutov, S., Dhomse, S., Feng, W., and Chipperfield, M. P.: Age of air as a diagnostic for transport timescales in global models, *Geoscientific Model Development*, 11, 3109–3130, doi:10.5194/gmd-11-3109-2018, <https://www.geosci-model-dev.net/11/3109/2018/>, 2018.
- Kunz, A., Pan, L. L., Konopka, P., Kinnison, D. E., and Tilmes, S.: Chemical and dynamical discontinuity at the extratropical tropopause based on START08 and WACCM analyses, *J. Geophys. Res.*, 116, n/a–n/a, doi:10.1029/2011JD016686, <http://dx.doi.org/10.1029/2011JD016686>, d24302, 2011.
- Lawrence, Z. D., Manney, G. L., Minschwaner, K., Santee, M. L., and Lambert, A.: Comparisons of polar processing diagnostics from 34 years of the ERA-Interim and MERRA reanalyses, *Atmos. Chem. Phys.*, 15, 3873–3892, doi:10.5194/acp-15-3873-2015, <http://www.atmos-chem-phys.net/15/3873/2015/>, 2015.
- Lefever, K., van der A, R., Baier, F., Christophe, Y., Errera, Q., Eskes, H., Flemming, J., Inness, A., Jones, L., Lambert, J.-C., Langerock, B., Schultz, M. G., Stein, O., Wagner, A., and Chabrillat, S.: Copernicus stratospheric ozone service, 2009–2012: validation, system intercomparison and roles of input data sets, *Atmos. Chem. Phys.*, 15, 2269–2293, doi:10.5194/acp-15-2269-2015, <http://www.atmos-chem-phys.net/15/2269/2015/>, 2015.
- Lin, S.-J.: A “Vertically Lagrangian” Finite-Volume Dynamical Core for Global Models, *Mon. Weather Rev.*, 132, 2293–2307, doi:10.1175/1520-0493(2004)132<2293:AVLFDC>2.0.CO;2, [http://dx.doi.org/10.1175/1520-0493\(2004\)132<2293:AVLFDC>2.0.CO;2](http://dx.doi.org/10.1175/1520-0493(2004)132<2293:AVLFDC>2.0.CO;2), 2004.
- Lin, S.-J. and Rood, R. B.: Multidimensional Flux-Form Semi-Lagrangian Transport Schemes, *Mon. Weather Rev.*, 124, 2046–2070, doi:10.1175/1520-0493(1996)124<2046:mffslt>2.0.co;2, 1996.
- Linz, M., Plumb, R. A., Gerber, E. P., and Sheshadri, A.: The Relationship between Age of Air and the Diabatic Circulation of the Stratosphere, *Journal of the Atmospheric Sciences*, 73, 4507–4518, doi:10.1175/JAS-D-16-0125.1, <https://doi.org/10.1175/JAS-D-16-0125.1>, 2016.
- Long, C. S., Fujiwara, M., Davis, S., Mitchell, D. M., and Wright, C. J.: Climatology and interannual variability of dynamic variables in multiple reanalyses evaluated by the SPARC Reanalysis Intercomparison Project (S-RIP), *Atmos. Chem. Phys.*, 17, 14 593–14 629, doi:10.5194/acp-17-14593-2017, <https://www.atmos-chem-phys.net/17/14593/2017/>, 2017.
- Mahieu, E., Chipperfield, M. P., Notholt, J., Reddmann, T., Anderson, J., Bernath, P. F., Blumenstock, T., Coffey, M. T., Dhomse, S. S., Feng, W., Franco, B., Froidevaux, L., Griffith, D. W. T., Hannigan, J. W., Hase, F., Hossaini, R., Jones, N. B., Morino, I., Murata, I., Nakajima, H., Palm, M., Paton-Walsh, C., III, J. M. R., Schneider, M., Servais, C., Smale, D., and Walker, K. A.: Recent Northern Hemisphere stratospheric HCl increase due to atmospheric circulation changes, *Nature*, 515, 104–107, doi:10.1038/nature13857, <http://dx.doi.org/10.1038/nature13857>, 2014.

- Mahowald, N. M., Plumb, R. A., Rasch, P. J., del Corral, J., Sassi, F., and Heres, W.: Stratospheric transport in a three-dimensional isentropic coordinate model, *Journal of Geophysical Research: Atmospheres*, 107, doi:10.1029/2001JD001313, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2001JD001313>, 2002.
- McFarlane, N. A.: The Effect of Orographically Excited Gravity Wave Drag on the General Circulation of the Lower Stratosphere and Troposphere., *J. Atmos. Sci.*, 44, 1775–1800, doi:10.1175/1520-0469(1987)044<1775:TEOOEG>2.0.CO;2, 1987.
- Meijer, E. W., Bregman, B., Segers, A., and van Velthoven, P. F. J.: The influence of data assimilation on the age of air calculated with a global chemistry-transport model using ECMWF wind fields, *Geophys. Res. Lett.*, 31, doi:10.1029/2004GL021158, 2004.
- Miyazaki, K., Iwasaki, T., Kawatani, Y., Kobayashi, C., Sugawara, S., and Hegglin, M. I.: Inter-comparison of stratospheric mean-meridional circulation and eddy mixing among six reanalysis data sets, *Atmos. Chem. Phys.*, 16, 6131–6152, doi:10.5194/acp-16-6131-2016, <http://www.atmos-chem-phys.net/16/6131/2016/>, 2016.
- Molod, A., Takacs, L., Suarez, M., and Bacmeister, J.: Development of the GEOS-5 atmospheric general circulation model: evolution from MERRA to MERRA2, *Geosci. Model Dev.*, 8, 1339–1356, doi:10.5194/gmd-8-1339-2015, <https://www.geosci-model-dev.net/8/1339/2015/>, 2015.
- Monge-Sanz, B. M., Chipperfield, M. P., Simmons, A. J., and Uppala, S. M.: Mean age of air and transport in a CTM: Comparison of different ECMWF analyses, *Geophys. Res. Lett.*, 34, doi:10.1029/2006gl028515, 2007.
- Monge-Sanz, B. M., Chipperfield, M. P., Dee, D. P., Simmons, A. J., and Uppala, S. M.: Improvements in the stratospheric transport achieved by a chemistry transport model with ECMWF (re)analyses: identifying effects and remaining challenges, *Quart. J. R. Meteorol. Soc.*, doi:10.1002/qj.1996, <http://dx.doi.org/10.1002/qj.1996>, 2012.
- Morgenstern, O., Hegglin, M. I., Rozanov, E., O'Connor, F. M., Abraham, N. L., Akiyoshi, H., Archibald, A. T., Bekki, S., Butchart, N., Chipperfield, M. P., Deushi, M., Dhomse, S. S., Garcia, R. R., Hardiman, S. C., Horowitz, L. W., Jöckel, P., Josse, B., Kinnison, D., Lin, M., Mancini, E., Manyin, M. E., Marchand, M., Marécal, V., Michou, M., Oman, L. D., Pitari, G., Plummer, D. A., Revell, L. E., Saint-Martin, D., Schofield, R., Stenke, A., Stone, K., Sudo, K., Tanaka, T. Y., Tilmes, S., Yamashita, Y., Yoshida, K., and Zeng, G.: Review of the global models used within phase 1 of the Chemistry–Climate Model Initiative (CCMI), *Geosci. Model Dev.*, 10, 639–671, doi:10.5194/gmd-10-639-2017, <http://www.geosci-model-dev.net/10/639/2017/>, 2017.
- Muthers, S., Kuchar, A., Stenke, A., Schmitt, J., Anet, J. G., Raible, C. C., and Stocker, T. F.: Stratospheric age of air variations between 1600 and 2100, *Geophys. Res. Lett.*, 43, 5409–5418, doi:10.1002/2016GL068734, <http://dx.doi.org/10.1002/2016GL068734>, 2016GL068734, 2016.
- Neu, J. L. and Plumb, R. A.: Age of air in a “leaky pipe” model of stratospheric transport, *J. Geophys. Res.*, 104, 19 243–19 255, doi:10.1029/1999JD900251, <http://dx.doi.org/10.1029/1999JD900251>, 1999.
- Neu, J. L., Douglass, A., Huck, P., Oman, L., Pendlebury, D., and Tegtmeier, S.: SPARC CCMVal Report on the Evaluation of Chemistry–Climate Models, chap. 5. Transport, WCRP, <http://www.sparc-climate.org/publications/sparc-reports/sparc-report-no-5/>, 2010.
- Ossó, A., Sola, Y., Rosenlof, K., Hassler, B., Bech, J., and Lorente, J.: How Robust Are Trends in the Brewer–Dobson Circulation Derived from Observed Stratospheric Temperatures?, *J. Climate*, 28, 3024–3040, doi:10.1175/JCLI-D-14-00295.1, <https://doi.org/10.1175/JCLI-D-14-00295.1>, 2015.
- Pawson, S., Stajner, I., Kawa, S. R., Hayashi, H., Tan, W.-W., Nielsen, J. E., Zhu, Z., Chang, L.-P., and Livesey, N. J.: Stratospheric transport using 6-h-averaged winds from a data assimilation system, *J. Geophys. Res.*, 112, n/a–n/a, doi:10.1029/2006JD007673, <http://dx.doi.org/10.1029/2006JD007673>, d23103, 2007.

- Ploeger, F., Konopka, P., Günther, G., Grooß, J., and Müller, R.: Impact of the vertical velocity scheme on modeling transport in the tropical tropopause layer, *Journal of Geophysical Research: Atmospheres*, 115, doi:10.1029/2009JD012023, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2009JD012023>, 2010.
- Ploeger, F., Abalos, M., Birner, T., Konopka, P., Legras, B., Müller, R., and Riese, M.: Quantifying the effects of mixing and residual circulation on trends of stratospheric mean age of air, *Geophys. Res. Lett.*, 42, 2047–2054, doi:10.1002/2014GL062927, <http://dx.doi.org/10.1002/2014GL062927>, 2015a.
- Ploeger, F., Riese, M., Haenel, F., Konopka, P., Müller, R., and Stiller, G.: Variability of stratospheric mean age of air and of the local effects of residual circulation and eddy mixing, *J. Geophys. Res.*, 120, 716–733, doi:10.1002/2014JD022468, <http://dx.doi.org/10.1002/2014JD022468>, 2015b.
- 10 Podglajen, A., Hertzog, A., Plougonven, R., and Zagar, N.: Assessment of the accuracy of (re)analyses in the equatorial lower stratosphere, *J. Geophys. Res.*, 119, 11,166–11,188, doi:10.1002/2014JD021849, <http://dx.doi.org/10.1002/2014JD021849>, 2014.
- Ray, E. A., Moore, F. L., Rosenlof, K. H., Davis, S. M., Sweeney, C., Tans, P., Wang, T., Elkins, J. W., Bönnisch, H., Engel, A., Sugawara, S., Nakazawa, T., and Aoki, S.: Improving stratospheric transport trend analysis based on SF₆ and CO₂ measurements, *J. Geophys. Res.*, 119, 14,110–14,128, doi:10.1002/2014JD021802, <http://dx.doi.org/10.1002/2014JD021802>, 2014JD021802, 2014.
- 15 Reddmann, T., Ruhnke, R., and Kouker, W.: Three-dimensional model simulations of SF₆ with mesospheric chemistry, *J. Geophys. Res.*, 106, 14 525–14 537, doi:10.1029/2000JD900700, <http://dx.doi.org/10.1029/2000JD900700>, 2001.
- Richter, J. H., Solomon, A., and Bacmeister, J. T.: Effects of vertical resolution and nonorographic gravity wave drag on the simulated climate in the Community Atmosphere Model, version 5, *J. Adv. Model. Earth Syst.*, 6, 357–383, doi:10.1002/2013MS000303, <http://dx.doi.org/10.1002/2013MS000303>, 2014.
- 20 Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G.-K., Bloom, S., Chen, J., Collins, D., Conaty, A., da Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick, A. G., Sienkiewicz, M., and Woollen, J.: MERRA: NASA's Modern-Era Retrospective Analysis for Research and Applications, *J. Clim.*, 24, 3624–3648, doi:10.1175/JCLI-D-11-00015.1, <http://dx.doi.org/10.1175/JCLI-D-11-00015.1>, 2011.
- 25 Rotman, D. A., Tannahill, J. R., Kinnison, D. E., Connell, P. S., Bergmann, D., Proctor, D., Rodriguez, J. M., Lin, S. J., Rood, R. B., Prather, M. J., Rasch, P. J., Considine, D. B., Ramarosan, R., and Kawa, S. R.: Global Modeling Initiative assessment model: Model description, integration, and testing of the transport shell, *J. Geophys. Res.*, 106, 1669–1691, 2001.
- Saha, S., Moorthi, S., Pan, H.-L., Wu, X., Wang, J., Nadiga, S., Tripp, P., Kistler, R., Woollen, J., Behringer, D., Liu, H., Stokes, D., Grumbine, R., Gayno, G., Wang, J., Hou, Y.-T., Chuang, H.-Y., Juang, H.-M. H., Sela, J., Iredell, M., Treadon, R., Kleist, D., Van Delst, P., Keyser, D., Derber, J., Ek, M., Meng, J., Wei, H., Yang, R., Lord, S., Van Den Dool, H., Kumar, A., Wang, W., Long, C., Chelliah, M., Xue, Y., Huang, B., Schemm, J.-K., Ebisuzaki, W., Lin, R., Xie, P., Chen, M., Zhou, S., Higgins, W., Zou, C.-Z., Liu, Q., Chen, Y., Han, Y., Cucurull, L., Reynolds, R. W., Rutledge, G., and Goldberg, M.: The NCEP Climate Forecast System Reanalysis, *Bull. Am. Meteorol. Soc.*, 91, 1015–1057, doi:10.1175/2010BAMS3001.1, <http://dx.doi.org/10.1175/2010BAMS3001.1>, 2010.
- Saha, S., Moorthi, S., Wu, X., Wang, J., Nadiga, S., Tripp, P., Behringer, D., Hou, Y.-T., ya Chuang, H., Iredell, M., Ek, M., Meng, J., Yang, R., Mendez, M. P., van den Dool, H., Zhang, Q., Wang, W., Chen, M., and Becker, E.: The NCEP Climate Forecast System Version 2, *J. Climate*, 27, 2185–2208, doi:10.1175/JCLI-D-12-00823.1, <http://dx.doi.org/10.1175/JCLI-D-12-00823.1>, 2014.
- 35

- Santer, B. D., Wigley, T. M. L., Boyle, J. S., Gaffen, D. J., Hnilo, J. J., Nychka, D., Parker, D. E., and Taylor, K. E.: Statistical significance of trends and trend differences in layer-average atmospheric temperature time series, *J. Geophys. Res.*, 105, 7337–7356, doi:10.1029/1999JD901105, <http://dx.doi.org/10.1029/1999JD901105>, 2000.
- Segers, A., van Velthoven, P., Bregman, B., and Krol, M.: On the Computation of Mass Fluxes for Eulerian Transport Models from Spectral Meteorological Fields, pp. 767–776, Springer Berlin Heidelberg, Berlin, Heidelberg, doi:10.1007/3-540-46080-2_81, http://dx.doi.org/10.1007/3-540-46080-2_81, 2002.
- Simmons, A. J. and Burridge, D. M.: An energy and angular-momentum conserving finite-difference scheme and hybrid vertical coordinates, *Mon. Wea. Rev.*, 109, 758–766, doi:10.1175/1520-0493(2002)130<2459:ANTFVC>2.0.CO;2, [http://dx.doi.org/10.1175/1520-0493\(2002\)130<2459:ANTFVC>2.0.CO;2](http://dx.doi.org/10.1175/1520-0493(2002)130<2459:ANTFVC>2.0.CO;2), 1981.
- 10 Simmons, A. J., Poli, P., Dee, D. P., Berrisford, P., Hersbach, H., Kobayashi, S., and Peubey, C.: Estimating low-frequency variability and trends in atmospheric temperature using ERA-Interim, *Quart. J. R. Meteorol. Soc.*, 140, 329–353, doi:10.1002/qj.2317, <http://dx.doi.org/10.1002/qj.2317>, 2014.
- Skachko, S., Errera, Q., Ménard, R., Christophe, Y., and Chabrillat, S.: Comparison of the ensemble Kalman filter and 4D-Var assimilation methods using a stratospheric tracer transport model, *Geosci. Model Dev.*, 7, 1451–1465, doi:10.5194/gmd-7-1451-2014, <http://www.geosci-model-dev.net/7/1451/2014/>, 2014.
- 15 Stiller, G. P., von Clarmann, T., Höpfner, M., Glatthor, N., Grabowski, U., Kellmann, S., Kleinert, A., Linden, A., Milz, M., Reddmann, T., Steck, T., Fischer, H., Funke, B., López-Puertas, M., and Engel, A.: Global distribution of mean age of stratospheric air from MIPAS SF₆ measurements, *Atmos. Chem. Phys.*, 8, 677–695, doi:10.5194/acp-8-677-2008, <http://www.atmos-chem-phys.net/8/677/2008/>, 2008.
- Stiller, G. P., von Clarmann, T., Haenel, F., Funke, B., Glatthor, N., Grabowski, U., Kellmann, S., Kiefer, M., Linden, A., Lossow, S., and López-Puertas, M.: Observed temporal evolution of global mean age of stratospheric air for the 2002 to 2010 period, *Atmos. Chem. Phys.*, 12, 3311–3331, doi:10.5194/acp-12-3311-2012, <http://www.atmos-chem-phys.net/12/3311/2012/>, 2012.
- 20 Stiller, G. P., Fierli, F., Ploeger, F., Cagnazzo, C., Funke, B., Haenel, F. J., Reddmann, T., Riese, M., and von Clarmann, T.: Shift of subtropical transport barriers explains observed hemispheric asymmetry of decadal trends of age of air, *Atmos. Chem. Phys.*, 17, 11 177–11 192, doi:10.5194/acp-17-11177-2017, <https://www.atmos-chem-phys.net/17/11177/2017/>, 2017.
- Strahan, S. E. and Polansky, B. C.: Meteorological implementation issues in chemistry and transport models, *Atmos. Chem. Phys.*, 6, 2895–2910, doi:10.5194/acp-6-2895-2006, <http://www.atmos-chem-phys.net/6/2895/2006/>, 2006.
- Trenberth, K. E. and Olson, J. G.: An Evaluation and Intercomparison of Global Analyses from the National Meteorological Center and the European Centre for Medium Range Weather Forecasts, *Bulletin of the American Meteorological Society*, 69, 1047–1057, doi:10.1175/1520-0477(1988)069<1047:AEAIOG>2.0.CO;2, [https://doi.org/10.1175/1520-0477\(1988\)069<1047:AEAIOG>2.0.CO;2](https://doi.org/10.1175/1520-0477(1988)069<1047:AEAIOG>2.0.CO;2), 1988.
- 30 Vigouroux, C., Blumenstock, T., Coffey, M., Errera, Q., García, O., Jones, N. B., Hannigan, J. W., Hase, F., Liley, B., Mahieu, E., Mellqvist, J., Notholt, J., Palm, M., Persson, G., Schneider, M., Servais, C., Smale, D., Thölix, L., and De Mazière, M.: Trends of ozone total columns and vertical distribution from FTIR observations at eight NDACC stations around the globe, *Atmos. Chem. Phys.*, 15, 2915–2933, doi:10.5194/acp-15-2915-2015, <http://www.atmos-chem-phys.net/15/2915/2015/>, 2015.
- 35 Waugh, D. and Hall, T.: Age of Stratospheric Air: Theory, Observations, and Models, *Rev. Geophys.*, 40, 1–1–26, doi:10.1029/2000RG000101, <http://dx.doi.org/10.1029/2000RG000101>, 2002.
- Wright, J. S. and Fueglistaler, S.: Large differences in reanalyses of diabatic heating in the tropical upper troposphere and lower stratosphere, *Atmos. Chem. Phys.*, 13, 9565–9576, doi:10.5194/acp-13-9565-2013, <http://www.atmos-chem-phys.net/13/9565/2013/>, 2013.




Figure 1. Mean Age of Air (AoA, in years) from two model simulations using idealized tracers advected by ERA-I for fixed year 2000. Models shown are BASCOE ~~CTM~~ CTM (blue solid lines) and TOMCAT (blue dotted lines). The modeled AoA fields are ~~corrected so that mean age = 0 at Equator~~ calculated using as reference the tropical tropopause region (10°S-10°N, 15-km 100hPa). Upper left: values at 20km height; upper right: vertical profiles at 5°S; lower left: vertical profiles at 40°N; lower right: vertical profiles at 65°N. The symbols represent in situ observations collected during the ~~1990's~~ 1990s (see Hall et al., 1999; Waugh and Hall, 2002, for details). The legend in the upper left panel applies to all four panels.




Figure 2. Latitude-pressure distribution of ~~mean-AoA~~ in 2002–2007 from the BASCOE simulation driven by ERA-I. ~~Blue colors indicate relatively small values, orange and red colors indicate relatively large values.~~




Figure 3. Latitude-pressure distribution of AoA (years) in 2002–2007 from BASCOE simulations driven by all reanalyses but ERA-I (top row; same color scale as previous figure) and relative difference with respect to the mean AoA by the ERA-I-driven simulation for the same period (bottom row; darker blues indicate more negative differences and darker reds more positive differences difference is not plotted at grid points where ERA-I AoA is smaller than 5 days). These reanalyses are, from left to right: JRA-55, CFSR, MERRA-2, MERRA.




Figure 4. AoA (years) in 2002–2007 by the BASCOE TM driven by five reanalyses (solid lines) versus in-situ observations (symbols) with their 1 σ uncertainties (grey shading). The five reanalyses are ERA-I (blue), MERRA-2 (red), MERRA (pink), JRA-55 (purple) and CFSR (green). ~~The modeled AoA fields are corrected so that mean age = 0 at the tropical tropopause (100 hPa).~~ (a) AoA at 50 hPa with aircraft observations of CO₂ (Andrews et al., 2001; Neu et al., 2010). (b) AoA in the tropics (10°N–10°S) with aircraft observations (Andrews et al., 2001; Chipperfield et al., 2014). (c) AoA in the northern mid-latitudes (35°N–45°N) with balloon observations (Engel et al., 2009; Chipperfield et al., 2014). (d) AoA gradient differences between the northern mid-latitudes and tropics (Neu et al., 2010; Chipperfield et al., 2014). The legend in panel (d) applies to panels (b) and (c) as well.




Figure 5. Same as figure 4(d) but for the period 1992–1997.




Figure 6. Time evolution of AoA (years) interpolated to a pressure of 50 hPa in the northern mid-latitudes (40°N – 50°N mean, top) and in the southern mid-latitudes (50°S – 40°S mean, bottom). Thin lines show instantaneous model output every 5 days using the five reanalyses with color codes according to the legend shown in the lower panel. Thick lines are smoothed with a one-year running mean. The black vertical lines highlight the start of the Pinatubo eruption and the first assimilation of AMSU (see text).




Figure 7. Time evolution of the globally averaged (72°S – 72°N) anomalies of AoA (years) with respect to their mean (1989–2015) annual cycles, between 16 km and 28 km, using the five reanalyses with same color codes as in previous figure. ~~No impact of the Pinatubo eruption can be seen, contrarily to the results of Diallo et al. (2017, Fig. 1b).~~




Figure 8. Time evolution of AoA (years) averaged from 30 hPa to 5 hPa (approximately 24 to 36 km). Thick lines show model output smoothed with a one-year running mean and line color codes as in the southern previous figure. (50S–40S, left) and Mean for the northern mid-latitudes (40°N–50°N, right) : Solid lines show model output with color codes according to where the legend shown in the left panel. Thin lines (left panel only; omitted from right panel for clarity) show instantaneous model output every 5 days while thick lines are smoothed with a one-year running mean. Northern mid-latitude symbols (right panel) represent values derived from balloon observations of SF₆ (circles) and CO₂ (triangles) with color code showing the latitude of the measurements (according to the inset color bar) and outer error bars including sampling uncertainties (Engel et al., 2017). (b) Mean for the tropical latitudes (30°S–30°N) where the dashed lines show AoA using the tropical tropopause as reference. (c) Mean for the southern mid-latitudes (50°S–40°S) where the thin lines show instantaneous model output every 5 days. Except for the dashed lines in panel (b), all AoA in this figure use the surface as reference.




Figure 9. Amplitude (in years) of the seasonal variation in the 2002–2015 linear regression fit of AoA, as a function of pressure and averaged in four latitude bands, from left to right: a) North Pole, (70°N–90°N); b) Mid-latitudes North, (40°N–50°N); c) South Pole, 90°S–70°S; d) Tropics (30°S–30°N); e) Mid-latitudes South, (50°S–40°S). The regression model was applied separately to AoA results for the periods 1989–2001 (dashed thin lines) and 2002–2015 (thick solid lines). Same color codes as in previous figures: the legends in panels (a) and (c) apply to all four panels.




Figure 10. Amplitude (in years) of the QBO variation in the 2002-2015-2002-2015 linear regression fit of AoA, as a function of latitude at pressure 30 hPa. Same color codes as in previous figures.




Figure 11. Latitude-pressure distribution of AoA trends (in years /per decade) using the ERA-Interim reanalysis over 2002-2012. White crosses indicate grid points where the sign of the trend is not significant, i.e., its absolute value is smaller than the ~~standard error~~ uncertainty delivered by the regression analysis at the 95% confidence level. The color scale is the same as in Haenel et al. (2015, Fig. 6 and 10) with darker blues indicating more negative trends and darker reds more positive trends.




Figure 12. Latitude-pressure distributions of AoA trends (in years ^{per} decade) over 1989-2001 (left column), 2002-2015 (middle column) and 1989-2015 (right column) using the five reanalyses (from top to bottom: ERA-I, CFSR, JRA-55, MERRA, MERRA-2). White crosses and colors have the same meaning than in the previous figure, but note the different scale (top of figure).