

Response to Anonymous Referee #1

The authors would like to thank Anonymous Referee #1 for his comments. Below, please find our response to the referee's comments (RC to denote Reviewer's comment, AR to denote Authors' reply).

RC-1) Main objective: It is unclear if the goal of the study is to understand the seasonal/diurnal cloud evolution of the Namibia-Angola stratiform cloud regime (as suggested by the title) or to evaluate SEVIRI cloud retrievals with other datasets (most of the analysis revolves around the differences between SEVIRI and other datasets, and potential bias in SEVIRI retrievals). If the focus is to characterize the diurnal cycle, then please provide more detailed information about the amplitude of the daytime cycle and explain spatial/temporal changes in the context of the atmospheric circulation and thermodynamical structure. For instance, in Painemal et al. (2012, JGR), we attempted to understand the dynamical factors behind the cloud diurnal cycle in the SE Pacific, and showed hourly composites (maps) of cloud retrievals. In Painemal et al. (2013, J. Atmos Sc.), we further endeavored to understand variations in liquid water path and cloud fraction, in the context of the boundary layer depth evolution and subsidence variability. Similarly, we utilized a super-parameterized climate model and NASA-Langley SEVIRI retrievals for describing the diurnal evolution of cloud fraction and height over the Namibia-Angola stratocumulus cloud deck (Painemal et al., 2015 J. Climate). If the focus is mostly evaluating the ability of SEVIRI to reproduce the diurnal cycle, please modify the title and the introduction accordingly.

AR-1) We do agree with the reviewer's comment that the original title could mislead the reader about the objective of the study. Therefore, we modified the title to "Evaluating the diurnal cycle of South Atlantic stratocumulus clouds as observed by MSG-SEVIRI". The primary objective of our manuscript is to *evaluate* the diurnal cycle of South Atlantic stratocumulus clouds based on cloud property retrievals from the SEVIRI CLAAS-2 algorithm. In order to make this clearer we modified the text in introduction section in the original manuscript **in line# 118**.

RC-2) Inhomogeneity and cloud mask: I was surprised that cloud fraction (mask) differences between SEVIRI CLAAS and MODIS collection 6 were not analyzed. I would speculate that the spatial pattern of the SEVIRI-MODIS difference near the equator in Figure S5d is most likely due to cloud fraction differences between both sensors/algorithms. Also, the use of

some sort of inhomogeneity index would provide support to the hypothesis that pixel resolution is in part responsible for the COT difference between SEVIRI and MODIS. Although the plane parallel bias is likely playing a role, it is puzzling that both SEVIRI COT and reff are generally smaller than their MODIS counterparts (1.6 μ m), as one would expect that reff (COT) is overestimated (underestimated) as the pixel resolution becomes coarser (e.g. table 1a and 1b in Painemal et al., 2012 JGR). This points to other issues associated with differences in the retrieving algorithms, since SEVIRI visible channels were calibrated against MODIS. The question could be answered if the CLAAS algorithm were applied to MODIS (I do not know if this is even possible). At the very least, the authors should speculate about the causes for the inconsistencies between MODIS and SEVIRI that cannot be explained by the plane parallel bias or absorbing aerosols. Lastly, if the COT threshold was applied to SEVIRI (and MODIS), then a comparable threshold should be applied to TMI. If not, the comparison between TMI and SEVIRI.

AR-2) Thank you for this suggestion. In the revised manuscript we included the diurnal cycle of cloud fraction for 2-yr mean in Fig. 10 and for seasonal means in Fig. S8. The cloud fraction maps for SEVIRI, MODIS and their differences are shown in Figure 7 for all-sky condition, and described in the revised manuscript. Figure S5d in the original manuscript (Fig. 6d in the revised manuscript) is SEVIRI-MODIS 1.6-micron LWP differences for the overcast case. Therefore, SEVIRI and MODIS LCF are >95 % and also both LCFs agree within ± 1 %.

We performed a preliminary analysis of factors that might explain SEVIRI COT/CER low biases. Firstly, spatial heterogeneity was considered. Similar to Painemal et al. (2013b), we found both for SEVIRI and for MODIS a decrease in COT and increase in CER with increasing scene heterogeneity under constant TMI LWP. However, the negative SEVIRI-MODIS CER difference remained a robust feature, independent of the magnitude of the scene heterogeneity. Secondly, the different view geometries of SEVIRI (fixed) and MODIS (varying from orbit to orbit) were analysed. A dependence of the SEVIRI-MODIS CER bias on the MODIS view zenith angle (VZA) was observed, with the bias varying between -0.5 μ m to -2 μ m and generally being lowest for the MODIS oblique backscatter view direction. Further analysis in this direction, including also the relative azimuth angle, is promising but deferred to a future study, in which also spectral and algorithmic differences between the MODIS and SEVIRI instruments and cloud property retrievals should be considered. The description is embedded **in original manuscript line#586** in the revised manuscript.

SEVIRI CER being smaller than MODIS is indeed unexpected based on the plane parallel bias effect alone. Of course, there are algorithmic differences too. For example, SEVIRI uses the 0.6-micron channel instead of the 0.8-micron as MODIS does over the ocean. Redoing the full CLAAS retrievals with the SEVIRI 0.8 micron channel or applying the CLAAS algorithms to MODIS could indeed shed light on the observed differences, but this would imply a huge effort and we feel this is beyond the scope of this study (which is an evaluation of the current official CLAAS-2 cloud properties).

Regarding the COT threshold applied to SEVIRI data, we confirm that a comparable threshold was effectively applied to both MODIS and TMI data. In the comparison we only included $0.25^\circ \times 0.25^\circ$ gridboxes (and the corresponding SEVIRI, TMI, and MODIS retrievals) with a gridbox-mean SEVIRI COT > 3 . All gridboxes (and the corresponding SEVIRI, TMI, and MODIS retrievals) with a mean SEVIRI COT < 3 were excluded from the analysis. **We clarified this in the first paragraph of section 3 in line# 245** in the revised manuscript.

RC-3) Inconsistent results: The curve “SEV” in Fig. 2a is SEVIRI collocated with MODIS, correct? Does it mean that not always MODIS and SEVIRI are collocated? All the retrievals should be spatially and temporally collocated. I also noticed that in Fig S9, SEVIRI reff for overcast and all-sky samples is almost the same (Fig. S9a and b). In contrast, the all-sky and overcast averages are quite different for MODIS, why?

AR-3) ‘SEVIRI’ is SEVIRI collocated with TMI. ‘SEV’ is SEVIRI collocated with MODIS. Thus, yes SEVIRI observations are always collocated in time and space with either TMI or MODIS. Unfortunately collocations in time and space of SEVIRI, TMI, and MODIS, so called triple collocations, are very rare, and would only be available when TMI and MODIS line-up. Therefore we did not use triple collocations. The fact that SEVIRI CER between overcast and all-sky are very similar, while MODIS shows larger differences, could be explained by the weighting applied in CLAAS. For thin clouds (which are the ones added in the all-sky averages) CLAAS-2 will tend to retrieve lower CER because of the weighting with a climatological a priori of 8-microns. MODIS will probably tend to retrieve higher CER for thin clouds because these are often also broken clouds for which CER is overestimated (esp. with 1.6-micron). Indeed, the differences between all-sky and overcast

MODIS CER are much smaller for the 3.7 micron channel (see Fig. S7), which is known to be less affected by cloud heterogeneity.

RC-4) What is the purpose of showing the three MODIS reff's as the apples-to-apples comparison is between MODIS-SEVIRI at 1.6 um? I understand that this is useful for understanding the reff bias due to absorbing aerosols (which is not a novel result), but it is extremely confusing to understand figures 6-7 with so many symbols and retrievals, and the overall objective of including MODIS 2.1 and 3.7 reff is unclear. Similarly, just report one MODIS COT as the three MODIS COT are essentially the same. The authors mention that differences in MODIS reff at 1.6, 2.1, and 3.7 um might be providing information about the cloud vertical structure and precipitation; however, numerous papers (e.g. Zhang and Platnick, 2011, Painemal et al. 2013 ACP: "The impact of horizontal heterogeneities, cloud fraction, and liquid water path on warm cloud effective radii from CERES-like Aqua MODIS retrievals) have shown that the difference between the 3.7 um and 2.1 um reff (or 1.6 um) mostly reflect the effect of spatial inhomogeneity and clear-sky contamination in the retrievals (for stratiform clouds).

AR-4) We agree that the different MODIS channel retrievals are useful for understanding the absorbing aerosol effect. Therefore, we propose to keep them in the new Figs. 2 and 3. We agree that the different MODIS channels don't add much in the scatter density plots and may cause confusion (new Fig. 8). Therefore, we removed the 2.1 and 3.7 micron plots from Fig. 8. In the overcast maps (Fig. 6) and diurnal cycle plots we do think the different channel MODIS CER retrievals (Fig. S7) do add important information regarding heterogeneity effects (as correctly argued by the referee). So we propose to keep these in the new version, while for LWP and COT the different MODIS retrievals are not shown anymore.

RC-5) Figs. S2 and S5 deserve to be included in the manuscript:

AR-5) We have included Figures S2 and S5 as Figs. 3 and 6 in the revised manuscript.

RC-6) Remove 'Discussion' from the title of Section 4:

AR-6) 'Discussion' has been removed from the respective section title in the revised version.

RC-7) Page 12, line 349, I disagree, Figure S1 does not show any difference for COT:

AR-7) We believe it does. Fig. S1c does show a median SEVIRI-MODIS COT bias of -1.

RC-8) Page 5, first paragraph: This is non-raining pixels according to the RSS algorithm. Depending on the threshold used to define a rainy pixel, drizzle or light precipitation is still possible:

AR-8) This is indeed possible. We have replaced the word ‘avoid’ by ‘minimize’ in the revised manuscript to indicate that not all rainy pixels will be filtered out in practice.

RC-9) Page 15, line 435: I cannot find the figure that shows the overestimation of SEVIRI (relative to MODIS) for broken scenes.

AR-9) This remark in particular concerns the 3.7-micron MODIS retrievals. Fig. S51 in the original manuscript (Fig. 6l in revised version) does show SEVIRI being 2-4 micron lower than MODIS, but for the other channels, the difference is less. However, this refers to ‘the area outside the Sc region’ not to ‘broken clouds’, since all plots in Fig. 6l are for overcast clouds only. **The discussion is elaborated in the revised version in line# 435 of the original manuscript.**

RC-10) Page 16, first paragraph. This explanation is unlikely, since you removed samples with high liquid water path (precipitating samples according to the RSS algorithms). Moreover, the spectral difference in reff is mostly indicative of the effect of spatial inhomogeneity and 3D radiative effects in the retrievals rather than information about the vertical structure or precipitation.

AR-10) Agree: the reason that MODIS 3.7 Reff is higher than that of 2.1 and 1.6 is probably mainly related to inhomogeneity. Horizontal inhomogeneity has a larger impact in the 1.6-micron than in the 3.7-micron channel. A deeper discussion is included in the revised manuscript **in line# 435 of the original manuscript.**

RC-11) Page 17, line 487, what do you mean by “smaller ...”.

AR-11) The relative variation in effective radius is smaller than the relative variation in COT during the day. The text in the revised version is now modified for clarity.

RC-12) Page 19 lines 547-549: I suspect this is mostly due to cloud thinning. If cloud fraction (mask) played a role, then cloud effective radius would be biased high due to clear-sky contamination and 3D radiative effects.

AR-12) We now included the diurnal cycle of SEVIRI liquid cloud fraction for our Sc study domain, both 2-yr (Fig. 10) and seasonal (Fig. S8) means. The diurnal variation in cloud fraction is very similar to the variation in all-sky LWP. COT and CER are in-cloud means, whereas LWP is in-cloud water content multiplied by liquid cloud fraction in order to compare with the gridbox mean TMI at a $0.25^\circ \times 0.25^\circ$ resolution aggregated from the original SEVIRI (and MODIS) resolution. Therefore, the diurnal variation in TMI and SEVIRI LWP follow the diurnal variation of cloud fraction in all-sky scene. For the overcast case, most of the diurnal variation in LWP is likely because of cloud thinning. In the all-sky case, MODIS 1.6 and 2.1 channel CER are indeed biased high due to clear-sky contamination and cloud heterogeneity impacts. On the other hand, 3.7 micron CER agrees well with SEVIRI. SEVIRI values are lower because of the climatological weighting applied to CER corresponding to the lower COT pixels.

RC-13) Page 20 line 558: Use austral winter instead:

AR-13) Modified accordingly in the revised version.

RC-14) Page 4, line 102, replace “evaluated” with “analyzed”.

AR-14) Modified accordingly in the revised manuscript.

RC-15) Page 4, line 110, Add “In contrast,” before Painemal et al. (2012). Painemal et al. (2012) also utilized in-situ cloud probe to assess the bias in satellite cloud properties.

AR-15) Have been added to the revised manuscript version.

RC-16) Page 5 line 143: Define VIS/NIR.

AR-16) In the revised version this has been defined.

RC-17) Page 6, line 157, If the physical retrievals are derived for $COT > 4$, it seems logical to use the same threshold for comparing SEVIRI with MODIS.

AR-17) For $COT = 4$ there is still a 13% influence of the a priori CER according to Eq. (2), so to completely get rid of this influence, one would even have to apply a larger threshold (say

COT>5). For COT=3 the weight of the a priori CER is 35%. This seemed a reasonable compromise between a modest weighting with the a priori on the one hand and keeping enough samples for robust statistics on the other hand.

RC-18) Maps: It is very difficult to extract quantitative information from the maps due to the use of a continuous color palette (too many tones). Instead, it would be better to define only 10 or 12 discrete colors.

AR-18) In the revised version, all the maps are plotted with discrete color palettes. Indeed, the figures are more readable now.

Response to Anonymous Referee #2

The authors would like to thank Anonymous Referee #2 for his/her comments. Below, please find our response to the referee's comments (RC to denote Reviewer's comment, AR to denote Authors' reply).

RC-1) Figure 2 indicates that the SEVIRI retrieval is substantially more sensitive to the presence of smoke above the cloud. The difference in the response is robust and indicates something meaningful about the differences in how the retrievals are performed, but is only addressed very briefly on line 375 as "partially explained by the spectral difference that for SEVIRI retrievals the 0.6um channel is used as a non-absorbing channel in contrast to the 0.8um channel for MODIS." I feel that this needs some deeper discussion. In what way is the SEVIRI channel more sensitive? Perhaps there is a citation that documents that spectral absorption features that explain this. Could the MODIS bands be chosen for SEVIRI in light of this additional bias due to absorbing aerosol? If the is only partly explained by the differing spectral absorption of smoke between 0.6 um and 0.8 um, then what are the other contributing factors?

AR-1) Indeed, a study by Haywood et al. (2004) investigated the spectral dependence of aerosol optical thickness for different non-absorbing and water absorbing channels and found that the effect of aerosol on the 0.63-micron radiance is significantly larger than that on the 0.86 micron radiance. The presence of aerosols above clouds reduces the 0.63-micron radiance substantially more than the 0.86-micron radiance. This could potentially introduce a low bias of (20% to >30%) 2 to >6 in retrieved cloud optical thickness for clouds with true COT 10 and 20, depending upon how large the true COT is, as the bias is highest for clouds with the largest COT. This low bias is larger for the retrieval at 0.63-micron SEVIRI channel than at 0.86-micron MODIS non-absorbing visible channel. Depending upon the paired water-absorbing NIR channel, the CER retrieval is also biased. Combining the 3.7-micron channel with the 0.63 or 0.86-micron visible channel, there is a relatively modest CER high bias of <1 micron, as the constant CER lines are more or less parallel to the 0.63 or 0.86-micron axis. However, the radiance pair of 0.86/1.63 introduced a significant low bias in CER of about 3 micron for a cloud with actual CER of 10 micron due to the apparent indirect effect induced by the decreased 0.86-micron radiance on non-parallel constant CER lines. This low bias will be even larger for the 0.6/1.63 radiance pair as used for the CLAAS-2 SEVIRI retrievals, and hence both SEVIRI COT and CER are expected to be lower than the

corresponding MODIS values retrieved from the 0.86/1.63 micron radiance pair. **This description is embedded in line# 347 of the original manuscript.**

RC-2) In line 319 it is noted that the SEVIRI retrieval exhibits a strong decrease in effective radius with increasing smoke above the cloud, but only a very weak decrease in cloud optical thickness. Is this consistent the cases presented in the Haywood et al. (2004) paper? Many of the cases in that paper exhibited a strong decrease in the optical thickness and only a weak decrease in the retrieved effective radius, although the details depend on the spectral bands chosen for the retrieval. Also, is this consistent with the explanation offered above for the stronger sensitivity of SEVIRI to smoke? I would expect that if the so-called “non-absorbing” band chosen is substantially more sensitive to smoke absorption, that this would cause a more substantial impact on the retrieved optical thickness than the effective radius. This needs to be clarified.

AR-2) The sentence in line 319 in original manuscript, about the “only a very weak decrease in cloud optical thickness with increasing AI”, is further described (in the original manuscript itself) in the paragraph immediately following it **in line# 330**. As such, cloud optical thickness has shown a weak decrease, but the expectation is that the true COT has to increase with AI, as strongly suggested by the sharp TMI LWP increase with AI. Thus, the overall low bias in COT due to smoke is indeed more substantial, in agreement with Haywood et al. (2004), because there is a simultaneous increase in true COT with AI (i.e. getting closer to the coast). The low bias in SEVIRI CER will also be larger than in MODIS CER due to the use of the 0.63/1.6-micron spectral pair, as the 0.63-micron reflectance would be substantially more affected by smoke than the 0.86-micron reflectance used in MODIS and also because the 1.6-micron based constant CER lines are less parallel to the 0.63-micron reflectance axis than to the 0.86-micron reflectance axis of the MODIS LUTs.

RC-3) In the paragraph beginning in line 435 comparing SEVIRI and MODIS in broken cloud scenes, it is noted that SEVIRI is biased high relative to MODIS primarily because of a high bias in the effective radius retrieved. The authors argue that this could be caused by the SEVIRI algorithm’s artificial use of a climatological effective radius for optically thin clouds. However, I wonder if it might also be contributed by the differences in resolution between SEVIRI and MODIS. Could it be that SEVIRI with a larger footprint than MODIS is simply more likely in broken cloud scenes to report a valid retrieval in a pixel that in reality

is contaminated by some inhomogeneity or clear-sky regions? That would presumably lead to a high bias in effective radius that is more substantial for SEVIRI than MODIS.

AR-3) The results presented in line 435 (in original manuscript) are for overcast gridboxes, where overcast is selected with the criteria of LCF>95% and COT>3. Outside the identified Sc regime, near the equator it is possible that the positive plane-parallel CER bias in heterogeneous cloud scenes with low COTs is the dominant one for the larger SEVIRI pixel size, whereas towards south it seems that the effect of climatological weighting applied to SEVIRI CER is dominant. The MODIS 1.6 and 2.1 micron CERs could have been impacted by plane-parallel bias and overestimated the retrieved CER. **This description is embedded in line# 428 in the original manuscript.**

Evaluating the diurnal cycle of South Atlantic stratocumulus clouds as observed by MSG-SEVIRI

Authors 8/16/2018 9:23 AM

Deleted: Characterizing

Authors 8/16/2018 9:23 AM

Deleted: cloud properties from satellite retrievals

Chellappan Seethala¹, Jan Fokke Meirink², Ákos Horváth³, Ralf Bennartz⁴, Rob Roebeling⁵

5 ¹Finnish Meteorological Institute, Kuopio, Finland

²Royal Netherlands Meteorological Institute, De Bilt, Netherlands

³University of Hamburg, Hamburg, Germany

⁴Vanderbilt University, Nashville, TN & University of Wisconsin – Madison, Madison, WI.

⁵EUMETSAT, Darmstadt, Germany

10 *Correspondence to:* Chellappan Seethala (seethala.chellappan@fmi.fi)

15

20

25

30

Abstract

Marine stratocumulus (Sc) clouds play an essential role in the earth radiation budget. Here, we compare liquid water path (LWP), [cloud](#) optical thickness (COT), and [cloud](#) effective radius (CER) retrievals from two years of collocated
40 Spinning Enhanced Visible and InfraRed Imager (SEVIRI), MODerate resolution Imaging Spectroradiometer (MODIS), and Tropical Rainfall Measuring Mission Microwave Imager (TMI) observations, estimate the effect of biomass burning smoke on passive imager retrievals, as well as evaluate the diurnal cycle of South Atlantic marine Sc clouds.

The effect of absorbing aerosols from biomass burning on the retrievals was investigated using aerosol index (AI)
45 obtained from the Ozone Monitoring Instrument (OMI). SEVIRI and MODIS LWPs were found to decrease with increasing AI relative to TMI LWP, consistent with well-known negative visible/near-infrared retrieval biases in COT and CER. In the aerosol-affected months of July-August-September, SEVIRI LWP – based on the 1.6- μm CER – was biased low by 14 g m^{-2} (~16 %) compared to TMI in overcast scenes, while MODIS LWP showed a smaller low bias of 4 g m^{-2} (~5 %) for the 1.6- μm channel and a high bias of 8 g m^{-2} (~10 %) for the 3.7- μm channel
50 compared to TMI. Neglecting aerosol-affected pixels reduced the mean SEVIRI-TMI LWP bias considerably. On a two-year data base, SEVIRI LWP had a correlation with TMI and MODIS LWP of about 0.86 and 0.94, respectively, and biases of only 4–8 g m^{-2} (5–10 %) for overcast cases.

The SEVIRI LWP diurnal cycle was in good overall agreement with TMI except in the aerosol-affected months.

Both TMI and SEVIRI LWP decreased from morning to late afternoon, after which a [slow](#) increase was observed.

55 Terra and Aqua MODIS mean LWPs also suggested a similar diurnal variation. The relative amplitude of the two-year mean and seasonal mean LWP diurnal cycle varied between 35–40 % from morning to late afternoon for overcast cases. The diurnal variation in SEVIRI LWP was mainly due to changes in COT, while CER showed only little diurnal variability.

60 1. Introduction

Changes in marine boundary layer (MBL) clouds over eastern subtropical oceans and associated differences in cloud radiative forcing are thought to be the main source of uncertainty in climate feedback simulations (Bony and Dufresne, 2005; Meehl et al., 2007, Zelinka et al., 2017). Climate models do not yet adequately parameterize the physical and dynamical processes affecting the formation of these clouds and fail to represent their variability on
65 different time scales. Thus, understanding MBL cloud variability and its driving mechanisms remains crucial.

Authors 8/16/2018 9:23 AM

Deleted: slight

70 Marine stratocumulus (Sc), [the dominant cloud type](#) prevalent over eastern subtropical oceans, [plays a vital role in](#) radiation budget calculations because [it reflects](#) most of the incoming solar radiation back to space while having little effect on terrestrial radiation. Marine Sc clouds tend to form [over](#) relatively cold sea surface temperatures (SSTs), within a shallow, well-mixed MBL capped by strong subsidence and a strong temperature inversion (e.g., Albrecht et al., 1995; Norris, 1998; Wood and Hartmann, 2006; Sandu et al., 2010). Several studies investigated the synoptic to inter-annual variability and the driving mechanisms of these clouds from both an observational and a modeling perspective (e.g., Klein and Hartmann, 1993; Klein et al., 1995; Bretherton and Wyant, 1997; Wood and Bretherton, 2006; Eastman et al., 2011; Wood, 2012; [Painemal et al., 2012, 2013a, 2015](#); [Adebiyi et al., 2015](#); [Adebiyi](#) and Zuidema, 2016; Horowitz et al., 2017; Kar et al., 2018; Lu et al., 2018).

80 Marine Sc clouds are prevalent throughout the year and exhibit an explicit diurnal cycle (Minnis and Harrison, 1984; Wood et al., 2002, 2012). The daily maximum in marine Sc clouds tends to occur during the early morning hours before sunrise, while the minimum usually occurs in the afternoon (Minnis et al., 1992; Rozendaal et al., 1995; Bretherton et al., 1995; Wood et al., 2002). During daytime, shortwave absorption by clouds effectively reduces or even cuts off the transport of heat and moisture from the surface into the cloud layer, resulting in a decoupled MBL (Nicholls, 1984; Betts, 1990); simultaneously, enhanced cloud-top entrainment of dry air from above promotes a weaker inversion (Duynderke et al., 2004), which leads to thinner or even disappearing clouds. During the night, on the other hand, strong longwave radiative cooling near cloud top produces negative buoyancy and, hence, a vertically well-mixed stable MBL (James, 1957; Moeng et al., 1992; Bretherton and Wyant, 1997), which increases cloud amount. Previous studies documented that subtropical Sc plays a significant role in the entire tropical response to climate perturbations (Miller, 1997), and underestimating [the amount of](#) these clouds in global climate models can lead to a positive SST bias as large as ~5 K (Ma et al., 1996). GCMs often fail to capture the diurnal variation of important processes in the cloud-topped MBL, such as the reduction of cloud fraction and the likelihood of decoupling in the afternoon (Abel et al., 2010; Medeiros et al., 2012). Wilson and Mitchell (1986) and Rozendaal et al. (1995) also demonstrated that introducing, or simply altering the resolution of, the diurnal cycle of these clouds in a GCM could trigger cloud radiative forcing both at the surface and at the top-of-atmosphere. Moreover, in the Fourth Assessment Report of the Intergovernmental Panel on Climate Change Forster et al. (2007) highlighted the diurnal cycle of stratiform clouds as one of the major uncertainties in current estimates of cloud radiative forcing. Comparisons of observations with models also revealed large and potentially systematic errors in the modeled diurnal cycle (O'Dell et al., 2008; Roebeling and van Meijgaard, 2009; Greuell et al., 2011).

Authors 8/16/2018 9:23 AM
Deleted:) clouds,

Authors 8/16/2018 9:23 AM
Deleted: are

Authors 8/16/2018 9:23 AM
Deleted: for

Authors 8/16/2018 9:23 AM
Deleted: they reflect

Authors 8/16/2018 9:23 AM
Deleted: under

Authors 8/16/2018 9:23 AM
Deleted: Adebiyi and Zuidema

To fully evaluate the diurnal cycle of Sc clouds, reliable observations with high spatial and temporal resolution are needed; the paucity of such data is one of the main reasons for the current level of uncertainty. A few studies took advantage of measurements available from intensive field campaigns, satellites, and model simulations to investigate the diurnal variations of these clouds. Notably, Blaskovic et al. (1990) evaluated the diurnal cycle of northeast Pacific Sc off the California coast using observations during the First International Satellite Cloud Climatology Project Regional Experiment (FIRE). [Painemal et al. \(2013a\)](#) reported that cloud top height and cloud fraction over the southeast Pacific were increased in the early morning hours and reached minimum in the afternoon. Most recently, Painemal et al. (2017) evaluated the diurnal cycle of cloud entrainment rate over northeast Pacific marine boundary layer clouds based on geostationary satellite retrievals and a mixed-layer model, and reported that the cloud top height tendency term dominates the entrainment. Ciesielski et al. (2001) evaluated the diurnal variation of northeast Atlantic Sc from the Atlantic Stratocumulus Transition Experiment (ASTEX). Zuidema and Hartmann (1995) and Wood et al. (2002) studied the diurnal variation in liquid water path (LWP) based on observations from microwave imagers. Rahn and Garreaud (2010) and Burleyson et al. (2013) analyzed the diurnal cycle of southeast Pacific Sc using the Variability of the American Monsoon Systems' Ocean-Cloud-Atmosphere-Land Study Regional Experiment (VOCALS-REx) datasets. Kniffka et al. (2014) studied the temporal and spatial characteristics of LWP of different types of clouds from SEVIRI data, for most of Europe and Africa. In general, all of these studies revealed an early morning maximum and afternoon minimum in cloud amount and LWP, linked to solar insolation/absorption. Rozendaal et al. (1995) and Wood et al. (2002) showed that the amplitude of diurnal variations in cloud amount and LWP could exceed 20 % of the mean value. These studies, however, did not consider diurnal variations in cloud optical thickness (COT) or cloud droplet effective radius (CER) and were usually based on limited measurements from a single instrument, the uncertainties of which were not well characterized. In contrast, Painemal et al. (2012) evaluated the diurnal cycle of LWP, COT, and CER for southeast Pacific Sc based on GOES-10 (Geostationary Operational Environmental Satellite-10) visible/near-infrared and microwave satellite observations, as well as in-situ cloud probe data but only for a two-month period. They noted that variations in COT drive the diurnal cycle of LWP mostly.

In this study, we investigate the diurnal variations of southeast Atlantic Sc clouds. This geographic domain is notable for its unique feature that part of the year a smoke layer transported from the African continent resides above the Sc deck, which poses a challenge to the retrieval of aerosol and cloud properties from space. In recent years several field campaigns have been initiated, to investigate aerosol-cloud interactions and their role in climate.

Authors 8/16/2018 9:23 AM
Deleted: the

Authors 8/16/2018 9:23 AM
Deleted: evaluated

Authors 8/16/2018 9:23 AM
Deleted: of two months

Authors 8/16/2018 9:23 AM
Deleted: The southeast Atlantic

Authors 8/16/2018 9:23 AM
Deleted: clouds

Authors 8/16/2018 9:23 AM
Deleted: as described in Zuidema et al. (2016),

Authors 8/16/2018 9:23 AM
Deleted: . The purpose

140 | [some of them in our region of interest \(Zuidema et al., 2016\). Satellite observations of cloud properties in this region](#)
| [are provided by the geostationary](#) Meteosat Second Generation (MSG) Spinning Enhanced Visible and InfraRed
Imager (SEVIRI) Cloud property dataset using SEVIRI - Edition 2 (CLAAS-2) from the Satellite Application
145 | Facility on Climate Monitoring (CM SAF) (Benas et al., 2017). [The purpose of our study is to evaluate the CLAAS-](#)
| [2 cloud properties using Version 7.1](#) TMI (Tropical Rainfall Measuring Mission (TRMM) Microwave Imager)
(Wentz 2018) and Collection 6 MODIS (Moderate Resolution Imaging Spectro-radiometer) retrievals (Platnick et
150 | al., 2017). [In this process,](#) the effect of above-cloud aerosols on LWP retrievals from the SEVIRI and MODIS
passive imagers [is quantified, and particular attention is paid to](#) the diurnal cycle of the Sc clouds in the South
Atlantic, which is a somewhat neglected region as most previous studies focused on the North or South Pacific (west
of California and Chile). The main strength of our study is the use of an extensive two-year dataset, which allows us
to investigate the seasonal variation of the diurnal cycle. SEVIRI's higher temporal resolution of 15 minutes allows
examining the diurnal cycle with greater detail than offered by earlier GOES instruments. We only consider non-
155 | raining warm liquid clouds to [minimize](#) significant retrieval uncertainties associated with the presence of rain and
ice clouds at higher altitudes. Retrieval artifacts related to absorbing aerosols (e.g., Haywood et al., 2004) have been
evaluated and aerosol-affected grid boxes have subsequently been removed from the analysis.

The paper is structured as follows. A description of our datasets including retrieval artifacts and
uncertainties is provided in Section 2. The comparison methodology is described in Section 3. Section 4 discusses
160 | retrieval biases related to the presence of smoke from continental biomass burning over clouds and analyzes spatial
distributions, comparison statistics, and diurnal variations of Sc properties from SEVIRI, TMI, and Terra and Aqua
MODIS on seasonal and two-year timescales. Finally, a summary is offered in Section 5.

2. Satellite datasets

160 2.1 Visible/Near-infrared (VIS/NIR) retrievals

2.1.1 Spinning Enhanced Visible and InfraRed Imager (SEVIRI)

SEVIRI is an optical radiometer onboard the MSG geostationary satellite series operated by the European
Organization for the Exploitation of Meteorological Satellites (EUMETSAT). SEVIRI measures radiances in 12
165 | spectral bands including 4 [visible/near-infrared \(VIS/NIR\)](#) channels (0.6–1.6 μm plus a broadband high-resolution
VIS channel) and 8 IR channels (3.9–13.4 μm). It has a spatial resolution of $3 \times 3 \text{ km}^2$ at nadir and a repeat frequency
of 15 minutes for full-disk images covering Europe, Africa, and the Atlantic Ocean.

Authors 8/16/2018 9:23 AM

Deleted: study is three-fold. One, to compare

Authors 8/16/2018 9:23 AM

Deleted:) against Version 7

Authors 8/16/2018 9:23 AM

Deleted: Two, to quantify

Authors 8/16/2018 9:23 AM

Deleted: . Three, to study

Authors 8/16/2018 9:23 AM

Deleted: avoid

The CM SAF CLAAS-2 climate data record is described in Benas et al. (2017). Part of the cloud processing software is the CPP (cloud physical properties) algorithm, which retrieves cloud optical thickness and cloud [droplet](#) effective radius based on measured reflectances in the 0.63- μm and 1.6- μm channels. The retrieval scheme is based on earlier bispectral methods (hereafter also referred to as visible/near-infrared or VIS/NIR technique) that retrieve cloud optical thickness and cloud [droplet effective radius](#) from satellite radiances at wavelengths in the (for clouds) non-absorbing visible and the moderately absorbing solar infrared part of the spectrum (Nakajima and King 1990; Han et al. 1994; Nakajima and Nakajima, 1995; Watts et al., 1998; Roebeling et al., 2006). The liquid water path is computed from the retrieved [cloud](#) optical thickness (τ or COT) and [cloud](#) droplet effective radius (r_e or CER) as

$$\text{LWP} = \frac{2}{3} \tau r_{e(1.6\mu\text{m})} \rho_l, \text{ where } \rho_l \text{ is the density of liquid water (Stephens 1978).} \quad (1)$$

The SEVIRI retrievals are available only during daytime and are performed assuming plane parallel clouds. Because r_e is not well constrained by the measured 1.6- μm channel reflectance for thin clouds, it is weighted towards a climatological a priori value of 8 μm for pixels with $\tau \leq 4$ -similar to the handling of small optical thicknesses in optimal estimation methods. The relationship used to weight the CER retrieval is,

$$r_{e,\text{assign}} = r_{e,\text{clim}} (1 - w) + r_{e,\text{ret}} w \quad (2)$$

where, $w = 1 / (1 + e^{(-1.25(\tau_{\text{ret}} - \tau_{w,\text{clim}})})$; $r_{e,\text{clim}} = 8 \mu\text{m}$; $\tau_{w,\text{clim}} = 2.5$

In part of our analysis, a $\tau > 3$ threshold is applied to minimize the impact of strongly weighted effective radii for thin clouds on the results. The SEVIRI shortwave channels were calibrated with Aqua-MODIS as described in Meirink et al. (2013). More details on the CPP retrieval algorithm are provided in CM SAF (2016).

2.1.2 Moderate Resolution Imaging Spectroradiometer (MODIS)

MODIS is the flagship instrument aboard the Terra and Aqua polar orbiter satellites. Terra has a 10:30am descending node sun-synchronous orbit, while Aqua has a 13:30pm ascending node sun-synchronous orbit. Terra and Aqua MODIS image the entire Earth's surface every 1 to 2 days, acquiring data in 36 spectral bands. The MODIS Collection 6 (C6) cloud property datasets (Platnick et al., 2017) with 1x1 km^2 spatial resolution from both Terra (MOD06) and Aqua (MYD06) have been used in this study.

Similar to CLAAS-2 SEVIRI, the MODIS C6 algorithm uses the VIS/NIR technique to retrieve cloud properties. Over ocean, the 0.86- μm band is used for optical thickness information in conjunction with one of three water-absorbing near-infrared bands located at 1.6, 2.2, and 3.7- μm , which are particularly sensitive to droplet

Authors 8/16/2018 9:23 AM

Deleted: particle

Authors 8/16/2018 9:23 AM

Deleted: 6

Authors 8/16/2018 9:23 AM

Deleted: particle size

Authors 8/16/2018 9:23 AM

Deleted: .

Authors 8/16/2018 9:23 AM

Deleted: .

Authors 8/16/2018 9:23 AM

Deleted: micron

effective radius. Although all three near-infrared channels generally observe the upper portion of clouds, the vertical sampling of droplets becomes progressively deeper from 3.7 to 1.6- μm due to decreasing absorption (Platnick, 2000).

210 The C6 algorithm is a revamped version of the Collection 5 (C5) algorithm that has gone through several updates to improve performance. Modifications include improved radiative transfer and lookup tables with finer COT and CER bins, redesigned cloud thermodynamic phase detection based on a variety of independent tests, and separate spectral retrievals of COT, CER, and derived LWP for channel combinations using the 1.6, 2.2, and 3.7- μm bands. Differences in CER between C5 and C6 are evaluated in Rausch et al. (2017). Depending on a subpixel
215 heterogeneity index, the properties of partly cloudy pixels are listed separately and the algorithm also provides retrieval failure metrics for pixels where the observed reflectances fall outside the LUT solution space.

2.1.3 Known uncertainties in VIS/NIR retrievals

While these datasets offer excellent resources for investigating warm, overcast single-layer clouds, they are subject
220 to certain retrieval artifacts due to algorithm assumptions and complexities in the retrieval technique. The VIS/NIR cloud property retrievals rely on 1-D radiative transfer model-generated LUTs, which do not account for subpixel cloud heterogeneity and 3-D cloud structure, and that could lead to significant biases in retrieved cloud properties for inhomogeneous and partially cloudy scenes. Cloud vertical stratification is essential to consider when computing LWP. Although MODIS retrieves effective radius at three separate water-absorbing channels, 1.6, 2.2, and 3.7- μm ,
225 all three are most sensitive to near cloud-top properties (Platnick 2000; Zhang and Platnick 2011). Hence, the LWP derived by combining retrieved τ and retrieved r_e from any one of the near IR channels could potentially under- or overestimate the true value depending upon the actual cloud stratification. For stratocumulus that typically follows a sub-adiabatic r_e profile, bigger droplets will be located near cloud top, and thus the derived LWP could be an overestimate. As a first-order correction, an adiabatic model is proposed by Wood and Hartmann (2006), which
230 results in a $\sim 17\%$ reduction from the standard vertically homogeneous LWP in eq. 1 (Bennartz 2007; Bennartz and Rausch, 2017). More details about the retrieval uncertainties of the VIS/NIR technique can be found in Horváth and Davies (2007), Seethala and Horváth (2010), Horváth et al. (2014), Zhang et al., (2012), Grosvenor et al., (2018) and references therein.

235 2.2 TRMM Microwave Imager (TMI)

Authors 8/16/2018 9:23 AM

Deleted: retrieval artifacts

TMI was a 5-channel, dual-polarized, passive microwave imager onboard the Tropical Rainfall Measuring Mission (TRMM) satellite that was operational between December 1997 – April 2015, continuously monitoring the tropics between 40° S and 40° N. Unlike the sun-synchronous polar orbiters hosting the similar SSM/I (Special Sensor Microwave/Imager) instruments, the TRMM satellite precessed west to east in a semi-equatorial orbit, producing data at different local times. The radiometer measured microwave radiation at 10.7, 19.4, 21.3, 37, and 85.5 GHz. The Wentz absorption-emission based algorithm (Wentz, 1997; Wentz and Spencer, 2000; Hilburn and Wentz, 2008) is used to retrieve meteorological parameters such as sea surface temperature (SST), surface wind speed (W), water vapor path (V), liquid water path (LWP), and rain rate (R) over the ocean. Our primary interest, LWP, is mainly derived from 37-GHz observations at a native resolution of 13 km, although here we used the 0.25° gridded product available from Remote Sensing Systems (RSS). The error characteristics of TMI data are similar to those of the RSS SSM/I and AMSR-E (Advanced Microwave Scanning Radiometer for Earth Observing System) products, as all microwave retrievals are produced by the same unified algorithm. Various sources of potential errors are documented in Horváth and Gentemann (2007), O'Dell et al. (2008), Seethala and Horváth (2010), Elsaesser et al. (2017), and Greenwald et al. (2018). Because the diurnal cycle is targeted here, the non-sun-synchronous TMI observations are particularly useful. The precessing orbit of TRMM allows for a comparison of observations at different local times, which cover the entire diurnal cycle over the course of a month.

TMI data were recently reprocessed using the significantly improved version 7.1 (V7.1) of the radiometer data processing algorithm (Wentz, 2015). The following major modifications were introduced: TMI brightness temperatures were recalibrated using the same procedures applied to all other RSS microwave products, the previously removed small negative LWP values are now reported, some large geolocation errors were corrected, the roll of the satellite was recalculated, the radiation contribution from the emissive antenna itself was removed, and Radio Frequency Interference (RFI) from the cold mirror was minimized. This improved V7.1 TMI product available at www.remss.com was utilized in this study.

2.3 Ozone Monitoring Instrument (OMI)

Areas affected by biomass burning smoke or desert dust were identified using OMI ultraviolet Aerosol Index (AI). OMI AI represents the deviation of measured 354-nm radiance from model estimates calculated for a purely molecular atmosphere bounded by a Lambertian surface, with positive values indicating the presence of absorbing aerosols (Torres et al. 2007). A distinguishing feature of OMI AI is its ability to detect absorbing aerosols above

Authors 8/16/2018 9:23 AM

Deleted: TMI

(and even mixed with) clouds. Specifically, we used the 0.25° resolution daily Level-2 gridded product (OMAERUVG).

270 3. Comparison methodology

For our study we used two years of data (December 2010 – November 2012) from SEVIRI, TMI, and Terra- and Aqua-MODIS. We consider JJA (June-July-August), SON (September-October-November), DJF (December-January-February), and MAM (March-April-May) respectively to represent austral winter, spring, summer, and autumn; henceforth, ‘seasonal’ refers to an average over a given season in 2 consecutive years. SEVIRI pixel-level

275 data were averaged down to TMI’s 0.25° x 0.25° resolution, only using SEVIRI retrievals within ±7.5 minutes of the

TMI observation time. Note that SEVIRI [and MODIS LWPs are](#) representative of the in-cloud LWP. For compatibility with the TMI gridbox-mean LWP, we multiplied SEVIRI LWP with the successful cloud retrieval fraction (henceforth referred to as “liquid cloud fraction” or LCF) within the TMI gridbox. Similarly, when matching COT, CER, and LWP from SEVIRI and MODIS, both datasets were averaged down to 0.25° x 0.25°

280 resolution, using the same temporal collocation criterion of ±7.5 minutes, [and the MODIS LWP was also scaled by the corresponding LCF. In order to minimize the impact of SEVIRI CER values that were strongly weighted towards a climatological a priori value in thin clouds, only those 0.25° x 0.25° gridboxes \(and the corresponding TMI, SEVIRI, and MODIS retrievals\) were included in the analysis of overcast scenes, which had a gridbox-mean SEVIRI COT >3.](#)

285 Our study domain is a 70° x 40° (50° W-20° E, 35° S-5° N) area in the Southeast Atlantic. Over the relatively cold SSTs near the Namibian coast extensive sheets of marine Sc clouds form, which transition into scattered trade Cu as they are advected towards [warmer ocean near the equator.](#) Decks of subtropical marine Sc, scattered Cu, and occasionally deep convective clouds cover the study domain. We however restricted our analysis to marine Sc clouds.

290 Because microwave and optical techniques represent fully independent approaches, each having their own shortcomings, the analysis of retrieval discrepancies does not necessarily establish absolute accuracies. A considerable number of studies have investigated the differences between LWP retrievals based on passive microwave and VIS/NIR satellite observations (Bennartz, 2007; Borg and Bennartz, 2007; Horváth and Davies, 2007; Horváth and Gentemann, 2007; Wilcox et al., 2009; Greenwald, 2009; Seethala and Horváth, 2010; Horváth et al., 2014, Cho et al., 2015; Greenwald et al., 2018). The major shortcomings of microwave [retrievals](#) were found

Authors 8/16/2018 9:23 AM

Deleted: LWP is

Authors 8/16/2018 9:23 AM

Deleted: .

Authors 8/16/2018 9:23 AM

Deleted: the

Authors 8/16/2018 9:23 AM

Deleted: measurements

300 to be the uncertain retrieval of LWP in the presence of rain and a wet (positive) bias of 10-15 g m⁻² in broken cloud fields. However, V7.1 TMI data now includes the small negative LWP values that were previously discarded [in V4 data, leading to a significantly reduced microwave wet bias](#) (Greenwald et al., 2018).

Authors 8/16/2018 9:23 AM
Deleted: , and thus the microwave wet-bias has been

305 The major issues affecting VIS/NIR measurements are the dependence of retrievals on cloud fraction, variations with sun-view geometry, horizontal and vertical [subpixel inhomogeneity](#), 3D radiative effects, and the presence of aerosols/cirrus above the liquid cloud layer. Agreement between the VIS/NIR and microwave techniques is generally better for more stratiform clouds, where a near-adiabatic cloud liquid water profile can be assumed. To minimize these retrieval problems, we examine the diurnal characteristics of only low-level non-raining warm (liquid) clouds, which typically dominate the South Atlantic marine Sc domain. Additional criteria are applied to reduce as much as possible the influence of rain and ice clouds; [gridboxes](#) are included only if flagged as confident liquid clouds with valid LWP retrieval, [cloud top temperature \(CTT\) > 275 K](#), [ice fraction = 0](#) in SEVIRI and MODIS retrievals, and [rain rate = 0](#) in TMI retrievals. Our study domain is also affected by continental biomass burning during austral winter and spring, which in turn affects VIS/NIR cloud retrievals; therefore, special attention is paid to the analysis of retrieval artifacts related to the presence of smoke over the Sc deck.

Authors 8/16/2018 9:23 AM
Deleted: causing

310 We noticed that the extent and location of South Atlantic Sc clouds vary from month to month; hence we opted to define the Sc domain dynamically, rather than selecting a fixed rectangular area [as](#) study domain. Thresholding the spatial mean map of liquid cloud fraction (LCF) and the heterogeneity parameter ($H = \frac{\text{SEVIRI } 0.63 \mu\text{m reflectance standard deviation}}{\text{mean reflectance}}$) was found to delineate Sc regions in good agreement with visual observations. To precisely define the Sc domain, we used a region-growing algorithm to find adjacent, connected [grid-boxes](#) with $\text{LCF} > 80\%$. The identified Sc regions were typically within 20° W – 20° E and 5° S – 35° S. Cloud properties were separately evaluated for two cases:

Authors 8/16/2018 9:23 AM
Deleted: -grid boxes

Authors 8/16/2018 9:23 AM
Deleted: and

Authors 8/16/2018 9:23 AM
Deleted: with ice fraction ≤ 0

Authors 8/16/2018 9:23 AM
Deleted: ,

Authors 8/16/2018 9:23 AM
Deleted: \leq

Authors 8/16/2018 9:23 AM
Deleted: ,

Authors 8/16/2018 9:23 AM
Deleted: to specify the

- 320
1. 'all-sky': including all grid boxes [from](#) the identified Sc domain.
 2. 'overcast': only including grid boxes [from](#) the identified Sc domain, [which had an LCF \$\geq 95\%\$ and a mean SEVIRI COT \$> 3\$](#) . These criteria were imposed to minimize retrieval artifacts related to broken clouds as well as thin clouds for which the [SEVIRI CER](#) retrieval in particular is relatively uncertain.

Authors 8/16/2018 9:23 AM
Deleted: ..

Authors 8/16/2018 9:23 AM
Deleted: in

Authors 8/16/2018 9:23 AM
Deleted: with $\text{LCF} \geq 95\%$ and $\text{COT} > 3$ in

325

4. Results

Authors 8/16/2018 9:23 AM
Deleted: and Discussion

4.1 Effect of biomass burning smoke on SEVIRI and MODIS retrievals

This section presents the analysis of the effect of smoke and/or aerosols above marine Sc on passive VIS/NIR imager retrievals of cloud properties. Our study domain, especially the Sc region located off the Namibia coast, is severely influenced by biomass burning on the African continent, as it produces episodic plumes of dark smoke that drift over the southeast Atlantic Ocean during the dry season JJASO (June-through-October). Beneath the elevated smoke layer, there is a persistent deck of bright marine Sc clouds. Previous research (Hobbs, 2002; McGill et al., 2003; Wilcox, 2010) has shown that the smoke is typically located in layers (at 2 to 4 km altitude) that are vertically separated from the Sc clouds below (at ~1.5 km altitude) and, hence, direct microphysical interaction between the aerosols and the Sc is often inhibited by the strong temperature inversion above the cloud layer. However, more recent studies e.g., Rajapakshe et al. (2017) reported that smoke layers are closer to the cloud layer, and significantly enhance the brightness of stratocumulus over there (Lu et al. 2018). Recently, several studies evaluated the dynamical and climatological impacts of the presence of smoke above Sc clouds from both modeling as well as satellite and/or field campaign measurements (Adebiyi et al., 2015; Adebiyi and Zuidema, 2016; Zuidema et al., 2016; Das et al., 2017; Horowitz et al., 2017; Chang and Christopher, 2017; Lu et al., 2018; Kar et al., 2018). When smoke resides above low-level clouds, the observed visible channel (0.6- or 0.8- μm) reflectance is reduced due to absorption by smoke, which is not taken into account in the LUTs and can introduce a negative bias in the retrieved COT as well as CER, and hence in LWP. According to Haywood et al. (2004), this negative bias in the 1.6- μm CER is significantly larger than that in the 2.1- μm CER (which is estimated to be less than 1 μm), while the bias in retrieved COT can be up to 30 %. Previous studies also noticed a domain-mean underestimation of ~3 to 6 g m^{-2} in MODIS LWP over the South Atlantic Sc region in the presence of absorbing aerosols (Bennartz and Harshvardhan, 2007; Wilcox et al., 2009; Seethala and Horváth, 2010). Therefore, we need to quantify the impact of absorbing aerosols on SEVIRI and MODIS VIS/NIR retrievals in our Sc domain for our study period. The presence of absorbing aerosols can be diagnosed using the OMI Aerosol Index (AI), because large positive AIs correspond to absorbing aerosols, such as dust and smoke, and small positive or negative AIs correspond to non-absorbing aerosols and clouds.

Figure 1a depicts the spatial distribution of average OMI aerosol index during JAS for 2011 and 2012, with the black contour representing the Sc region. It is clear that absorption by smoke is highest near the Namibian coast and decreases away from shore. The locations of greater cloud amount partly coincide with the locations of larger AIs. The spatial distribution of SEVIRI and TMI LWP and their bias for overcast conditions are shown in Figs. 1b-d. Near the coast where the smoke absorption is stronger, SEVIRI LWPs increasingly underestimated the TMI

LWPs (SEVIRI values were approximately half of the corresponding TMI values). Over the smoke-free areas of the stratocumulus region, on the other hand, SEVIRI-retrieved LWPs were slightly higher than TMI LWPs. The domain-mean TMI LWP is 85 g m^{-2} , whereas the mean SEVIRI LWP is only 71 g m^{-2} , indicating an LWP low bias of 14 g m^{-2} or $\sim 16 \%$ in SEVIRI retrievals.

375 In Fig. 2, cloud properties from TMI, SEVIRI, and MODIS retrievals are binned into AI bins of 0.5 for the overcast Sc conditions. In the SEVIRI $1.6\text{-}\mu\text{m}$ CER retrievals, a steady and strong decrease from 11 to $6 \mu\text{m}$ is observed, while the COT decrease is weaker from 10.8 to 9 with AI increasing from 0 to 3.5. As a result, SEVIRI LWP sharply decreases from 86 to 45 g m^{-2} over the same AI range. TMI LWP, in contrast, increases from 84 to 101 g m^{-2} between clean and increasingly polluted regions. For overcast grid boxes with little to no smoke absorption
380 (AI < 0.5), SEVIRI LWP agrees well with TMI LWP, having only a 2 g m^{-2} high bias. However, SEVIRI has a low bias of $6\text{--}25 \text{ g m}^{-2}$ for moderate AI between 1 and 2, and a large negative bias $\leq -40 \text{ g m}^{-2}$ for grid-boxes with AI > 2.5; the bias increases linearly with AI.

Considering that cloud amount happens to be spatially correlated with AI and that microwave retrievals are unaffected by absorbing aerosols, the increase in TMI LWP with increasing AI, that is closer to shore, seems
385 plausible. Because the variability of LWP is mostly controlled by COT rather than CER in the absence of smoke-induced retrieval biases (Seethala and Horváth, 2010; Painemal et al., 2012), the microwave retrievals suggest that the true COT should also increase with AI. Taken together, the microwave and VIS/NIR retrievals imply that SEVIRI COT is increasingly underestimated as AI increases, [in line with Haywood et al. \(2004\)](#). The low bias in SEVIRI LWP in smoke-affected areas arises from the combination of the negative COT and CER retrieval biases. A
390 similar underestimation is reported in aircraft retrievals of COT and CER for a stratus deck residing below an absorbing aerosol layer (Coddington et al., 2010).

Interestingly, a systematic overall increase in LWP with AI as indicated by TMI LWP in Fig. 2a was also noticed in previous observational and modeling studies, e.g., Johnson et al. (2004), Wilcox (2010), Randles and Ramaswamy (2010), Adebisi et al. (2015), Adebisi and Zuidema (2016). While this could partly be explained by the
395 fortuitous spatial correlation between higher aerosol loads and thicker clouds in this Sc region, these studies argue that strong atmospheric absorption by the smoke warms the 700 hPa air temperature and increases upward motion. This increased buoyancy inhibits cloud-top entrainment and promotes a stronger inversion, thereby helping to preserve humidity and cloud cover in the MBL, resulting in increased cloud amount and LWP compared to a smoke-free environment. Similar to our SEVIRI results, Bennartz and Harshvardhan (2007), Wilcox et al. (2009), and

Authors 8/16/2018 9:23 AM

Deleted: >

Authors 8/16/2018 9:23 AM

Deleted: .

Seethala and Horváth (2010) also noted a systematic MODIS LWP underestimation in Sc off southern Africa during the biomass burning seasons. Painemal et al. (2014) also noted a CER decrease in MODIS data despite increased LWP north of 5° S during the biomass burning season.

405 Retrieval discrepancies due to the presence of absorbing aerosols above Sc clouds were also evaluated
between SEVIRI and MODIS. [Unlike the TMI microwave technique, SEVIRI and MODIS rely on VIS/NIR
channels for cloud property retrieval and hence are heavily impacted by above-cloud aerosols. SEVIRI uses 0.63-
410 0.86- \$\mu\text{m}\$ reflectances, whereas MODIS uses 0.86- \$\mu\text{m}\$ reflectances over ocean, primarily to acquire COT. According to
the radiative transfer calculations of Haywood et al. \(2004\), aerosols over bright Sc reduce both the 0.63- and the
0.86- \$\mu\text{m}\$ radiance, but the reduction is more pronounced in the former due to the wavelength dependence of aerosol
optical thickness. Their calculations indicated a low bias of 2 to 6 in COT retrieved from 0.86- \$\mu\text{m}\$ reflectances for a
true COT of 10 and 20 respectively, and this low bias would be larger in COT retrieved from 0.63- \$\mu\text{m}\$ reflectances.
The water-absorbing channels used primarily to retrieve CER are 1.6-, 2.1-, and 3.7- \$\mu\text{m}\$ for MODIS, whereas for
415 CLAAS-2 only the SEVIRI 1.6- \$\mu\text{m}\$ channel is used. The 3.7- \$\mu\text{m}\$ and to a smaller extent the 2.1- \$\mu\text{m}\$ retrieved CERs
are less affected by aerosols above clouds, because the constant CER lines are nearly parallel to the visible
reflectance axis in the bispectral LUT. In contrast, the 1.6- \$\mu\text{m}\$ based constant CER lines are less parallel to the 0.63-
or 0.86- \$\mu\text{m}\$ reflectance axis in the LUT and hence there is a stronger underestimation of 1.6- \$\mu\text{m}\$ CER. For example,
according to Haywood et al. \(2004\), the 0.86/1.63- \$\mu\text{m}\$ radiance pair produced a significant low bias in CER of about
420 3 \$\mu\text{m}\$ for a cloud with actual CER of 10 \$\mu\text{m}\$, due to the apparent indirect effect induced by the decreased 0.86- \$\mu\text{m}\$
radiance on non-parallel constant CER lines. This low bias will be even larger for the 0.63/1.63 radiance pair as used
for the SEVIRI retrievals. As a result, SEVIRI COT and CER in smoke-affected regions are both expected to be
smaller than their MODIS counterparts retrieved from the 0.86/1.63- \$\mu\text{m}\$ radiance pair. In general, such absorbing-
aerosol biases are more pronounced for bright optically thick clouds.](#)

425 [Figure 3 depicts the spatial distribution of SEVIRI-MODIS LWP, COT, and CER differences for all three
water absorbing MODIS channels, averaged for JAS 2011 and JAS 2012 overcast conditions. Within the Sc regime,
SEVIRI COT is biased low by 1 compared to MODIS where the smoke absorption is highest. As expected, little
variation is observed in the SEVIRI-MODIS COT bias as a function of the MODIS water-absorbing channel,
because COT is mostly determined by the VIS channel reflectance. However, CER from the three MODIS channels
and thus the corresponding SEVIRI-MODIS CER bias, dramatically differs over the largest smoke absorption areas.
430 As discussed above, MODIS CER retrieved from the 3.7- \$\mu\text{m}\$ channel is expected to be the least affected by](#)

absorption effects. As shown, MODIS 3.7- μm CER values are 2 to 5 μm larger than the SEVIRI 1.6- μm CER, with the largest differences occurring in grid boxes with the strongest smoke absorption effects (in the 1.6- μm channel). The low bias between SEVIRI CER and MODIS 2.1- μm CER is 1 to 3 μm , whereas it is only ~ 1 μm compared to MODIS 1.6- μm CER, consistently over the Sc regime. As a result of the SEVIRI COT and CER low biases, the SEVIRI-MODIS LWP bias also increases from the MODIS 1.6- μm to the 3.7- μm channel. Not surprisingly, SEVIRI LWP agrees best with MODIS 1.6- μm LWP, with a typical bias of ± 5 g m^{-2} and a maximum bias of ~ 10 g m^{-2} over areas with the strongest smoke absorption. Compared to the MODIS 2.1- and 3.7- μm retrievals, the SEVIRI LWP bias ranges from 10-20 and 10-30 g m^{-2} , respectively, again showing the maximum over the strongest smoke absorption regions. These results are fully consistent with the differential absorption effects found by Haywood et al. (2004) and confirm that the 3.7- μm channel is the least affected by biomass smoke and generally performs best in aerosol-above-cloud situations.

Frequency histograms of SEVIRI minus MODIS LWP, COT, and CER biases, as well as the biases relative to MODIS CPP for overcast conditions aggregated for JAS 2011 and JAS 2012 are shown in supplemental Fig. S1. SEVIRI COT is biased low by ~ 1 compared to MODIS. Compared to the 1.6- μm MODIS CERs, ~ 70 % of SEVIRI CERs have a mean bias of -1.5 μm . Although SEVIRI CERs are biased low compared to all three MODIS CERs, the ~ 1 μm additional negative bias relative to the 2.1- and 3.7- μm CERs indicates much smaller smoke-induced retrieval artifacts in these two channels. In general, the CER retrievals from SEVIRI tend to be lower than corresponding retrievals from the three MODIS channels, with SEVIRI having about 1.5 μm to 2.5 μm lower CER values. The SEVIRI minus MODIS LWP distributions peak at about -10 g m^{-2} irrespective of the MODIS channel used for the retrieval.

The mean MODIS LWPs are 80 g m^{-2} , 87 g m^{-2} , and 90 g m^{-2} respectively for 1.6-, 2.1-, and 3.7- μm channel retrievals, while the corresponding mean SEVIRI LWP is 71 g m^{-2} . As shown in Fig. 2a, MODIS 1.6- μm retrieved LWP undergoes the largest decrease from 92 to 72 g m^{-2} with AI. In clean cases, MODIS 1.6- μm LWP is 10 % higher than the SEVIRI 1.6- μm LWP. The difference between MODIS and SEVIRI LWP is even larger for the 2.1- and 3.7- μm channel retrievals due to the wavelength-dependent absorption effects.

A decrease from 12 to 9 μm is observed in MODIS 1.6- μm CER, whereas both the MODIS 2.1- and 3.7- μm CERs show a smaller decrease of ~ 1.5 μm with increasing AI (Fig. 2c). This, again, indicates the reduced effect of absorbing aerosols on 2.1- and 3.7- μm reflectances. Surprisingly, SEVIRI 1.6- μm CERs are about 1.5 μm (2.5

Authors 8/16/2018 9:23 AM
Deleted: ,

Authors 8/16/2018 9:23 AM
Deleted: appeared to be

Authors 8/16/2018 9:23 AM
Deleted: high

Authors 8/16/2018 9:23 AM
Deleted: likely

Authors 8/16/2018 9:23 AM
Deleted: shows

Authors 8/16/2018 9:23 AM
Deleted: and the

Authors 8/16/2018 9:23 AM
Deleted: (Fig. 2c)

Authors 8/16/2018 9:23 AM
Deleted: .

μm and 3 μm) lower than MODIS 1.6-μm (2.1- and 3.7-μm channels) CERs, even in [less polluted \(AI < 0.5\)](#) overcast conditions.

Authors 8/16/2018 9:23 AM
Deleted: unpolluted

470 MODIS COT decreased slightly until AI < 1.5, increased steeply until AI = 2.5, and then leveled-off in all three channels. However, SEVIRI and MODIS COTs differ by 1 with MODIS being higher even in grid-boxes [least affected](#) by smoke. Taken together both COT and CER variations, MODIS LWPs show a decreasing trend with AI in all three channels, with the largest decrease of ~20 g m⁻² seen in the 1.6-μm retrieval. The 2.1- and 3.7-μm MODIS LWPs show a reduction of only ~10 g m⁻². The SEVIRI minus MODIS differences in LWP, COT, and CER increased with AI even in the common 1.6-μm channel; although differences were the smallest in this channel, 475 especially for CER. This is somewhat surprising, considering that the CLAAS-2 SEVIRI and MODIS C6 COT-CER retrieval algorithms are rather similar, the SEVIRI 1.6-μm channel has been calibrated with the corresponding MODIS channel, and the comparison is done for the most favorable overcast condition. The finding that AI has a stronger impact on SEVIRI 1.6-μm LWP than on MODIS 1.6-μm LWP [could be explained, as discussed above, by spectral differences in the visible channel used:](#) for SEVIRI retrievals the [0.63-μm](#) channel is used as a non- 480 absorbing channel in contrast to the [0.86-μm](#) channel for MODIS, the latter of which is less affected by aerosol absorption.

Authors 8/16/2018 9:23 AM
Deleted: unaffected

Authors 8/16/2018 9:23 AM
Deleted: may partially

Authors 8/16/2018 9:23 AM
Deleted: the

Authors 8/16/2018 9:23 AM
Deleted: difference that

Authors 8/16/2018 9:23 AM
Deleted: 6

Authors 8/16/2018 9:23 AM
Deleted: 8

Because the presence of absorbing aerosols above Sc clouds introduces a large negative bias in both SEVIRI and MODIS COT and CER retrievals, in the remainder of this work we will exclude grid-boxes with AI>0.1.

485 **4.2 Spatial distribution and mean statistics of SEVIRI, MODIS, and TMI cloud properties**

Authors 8/16/2018 9:23 AM
Formatted: Font:Bold

Authors 8/16/2018 9:23 AM
Formatted: Indent: First line: 0"

Authors 8/16/2018 9:23 AM
Deleted: -

This section presents the results of the comparison of SEVIRI, MODIS, and TMI LWP retrievals, as well as the comparison of SEVIRI and MODIS COT and CER retrievals. Significant variation in the distribution and amount of clouds is observed over the Sc region from month to month. During SON, we observe frequent Sc clouds with large 490 spatial extent. During JJA there are relatively fewer clouds that are shifted slightly to the north. The lowest cloud fractions are seen during DJF and MAM. From a surface-based cloud climatology, Klein and Hartmann (1993) also showed that there is strong seasonal variability in the amount of Sc clouds, which is closely tied to the seasonal cycle of static stability. Over the South Atlantic Sc region, SON had the largest lower tropospheric stability (LTS), and DJF had the smallest. The strongest net cloud radiative effect also occurred during August through November, 495 which further motivates us to examine the seasonal variability of these clouds.

The spatial distributions of two-year-mean SEVIRI cloud properties and TMI LWP for the overcast condition are shown in Fig. 4, whereas the results for the all-sky case are shown in Fig. S2. In the all-sky case, the spatial distribution of LWP indicates that over the marine Sc region the measurement techniques show good agreement, but SEVIRI overestimates TMI by $\sim 15 \text{ g m}^{-2}$ in smooth coastal fog. In contrast, the two-year mean SEVIRI LWP is much lower than the corresponding TMI mean LWP in regions with generally lower cloud fractions, and clouds with structured tops. This could be either due to a high bias in TMI LWP in broken scenes (Seethala and Horváth, 2010; Greenwald et. al., 2018) or an enhanced plane-parallel bias in broken more heterogeneous scenes underestimating COT and overestimating CER in SEVIRI 3km retrievals. In the Sc region, SEVIRI COT varies from 6 to 11 and CER ranges between 8 and 14 μm . The two-year-mean liquid cloud fraction varies between 75 % and 100 %. The mean statistics also show robust skill in LWP retrieval for both SEVIRI and TMI with a high correlation of 0.89 for the Sc regime. Both TMI and SEVIRI show a mean LWP of $\sim 53 \text{ g m}^{-2}$ with negligible bias and a standard deviation of 24 g m^{-2} for the study period.

In the overcast case over the Sc regime, the two-year mean LWP increases to 84 g m^{-2} and 80 g m^{-2} respectively for SEVIRI and TMI, i.e., the mean SEVIRI LWP is about 5 % larger than the mean TMI LWP. In this case, applying an adiabatic correction to SEVIRI LWP would lead to a larger bias of -10 g m^{-2} (-12 %) and standard deviation of 28 g m^{-2} . The unbiased LWPs observed in the all-sky Sc case could be associated with the cancellation of errors between fully overcast and lower LCF grid-boxes within the domain. A higher mean COT of ~ 11 characterizes the overcast Sc case, whereas the mean COT is only about 7 in the all-sky case, suggesting the presence of optically thin clouds which are more prone to retrieval biases. Figure 5 shows a density scatterplot of TMI and SEVIRI LWPs in the overcast Sc region. Most data points are close to the one-to-one line, although at the lower end TMI LWP is slightly higher, while the reverse is true at the higher end -the same feature is also found in monthly and seasonal results.

The daytime-averaged two-year and seasonal statistics of SEVIRI and TMI LWP are listed in Table 1. Seasonally, in the overcast Sc domain, the average LWP varies from 73 to 92 g m^{-2} in standard SEVIRI, 61 to 76 g m^{-2} in adiabatic SEVIRI, and 73 to 82 g m^{-2} in TMI retrievals. In the aerosol-free seasons of DJF and MAM, standard SEVIRI overestimates TMI LWP; applying an adiabatic correction to SEVIRI in these months brings the LWP bias within 5 %, similar to previous studies. The standard SEVIRI likely overestimates the actual LWP in the overcast Sc regime due to the overestimation of CER, as the observed CER in the 1.6- μm channel corresponds to the top layer and is higher than the cloud layer-mean in sub-adiabatic stratocumulus. However, for JJA, when all three

- Authors 8/16/2018 9:23 AM
Deleted: 3
- Authors 8/16/2018 9:23 AM
Deleted: S3
- Authors 8/16/2018 9:23 AM
Deleted: with negligible bias, while
- Authors 8/16/2018 9:23 AM
Deleted: .
- Authors 8/16/2018 9:23 AM
Deleted: is mostly
- Authors 8/16/2018 9:23 AM
Deleted:).

Authors 8/16/2018 9:23 AM
Deleted: LWP

Authors 8/16/2018 9:23 AM
Deleted: 4

months were heavily affected by smoke aerosol, the standard SEVIRI already shows ~10 % lower LWP than TMI; therefore, applying the adiabatic correction would only enhance this negative bias. For SON, only September was heavily affected by aerosol for the analysis years we considered. As a result, the mean standard SEVIRI LWP was ~5 % larger than TMI LWP and applying adiabatic correction would lead to a ~14 % underestimation in SEVIRI LWP. We found that SEVIRI underestimates LWP more during the aerosol-affected months, even after excluding grid-boxes with AI > 0.1 Applying a stricter criteria by excluding grid-boxes with AI > 0 did not improve the results, hinting at [residual](#) OMI AI retrieval biases.

The spatial distribution of SEVIRI and MODIS cloud properties averaged for the study period for the overcast condition is shown in Fig. 6. In general, over the overcast Sc regime, MODIS retrieves higher LWPs in all three channels compared to SEVIRI, but [outside the identified Sc regime, MODIS values are lower than SEVIRI LWP with the exception that 1.6- \$\mu\text{m}\$ \(and to a certain extent 2.1- \$\mu\text{m}\$ \) MODIS LWPs are lower in the north, i.e., closer to the equator and higher in the south. This LWP bias pattern can be explained by the respective CER spatial variation shown in Fig. 6\(f and i\) and that in cloud heterogeneity, as the fractional cloud cover is greater than 95 % in this case. SEVIRI CER values are 1 – 2 \$\mu\text{m}\$ higher in the north closer to the equator, probably indicating the plane-parallel bias in the larger SEVIRI pixel. However, SEVIRI CER is 1 – 2 \$\mu\text{m}\$ lower in the south of the domain due to the increased frequency of the climatological weighting of SEVIRI CER for lower COTs, while MODIS provides an actual retrieved CER. MODIS 3.7- \$\mu\text{m}\$ CER is consistently lower than SEVIRI likely because this channel is least affected by horizontal cloud heterogeneity and 3D cloud structure. The observed two-year-mean COT in the overcast Sc regime is 10.2 for SEVIRI, whereas it is 11.2 for MODIS, indicating SEVIRI mean COT is about 9 % lower than MODIS mean COT. Similarly, the observed two-year-mean CER in the overcast Sc domain is 10.1 \$\mu\text{m}\$ for SEVIRI, but for MODIS it varies between 11.3 – 11.7 \$\mu\text{m}\$ depending on the absorption channel, indicating that SEVIRI mean CER is 11–12 % lower than MODIS CER. The lower SEVIRI LWP value over the Sc regime is due to the combination of lower COT and lower CER values compared to MODIS. \[As expected, the COT bias varied little with MODIS absorption channel, whereas the bias in CER significantly depended on the MODIS NIR channel –MODIS 1.6- \\$\mu\text{m}\\$ CER is consistently ~1 \\$\mu\text{m}\\$ higher than the corresponding SEVIRI, however, MODIS 2.1- and 3.7- \\$\mu\text{m}\\$ CERs are 2 to 3 \\$\mu\text{m}\\$ larger closer to the Namibia coast, indicating the potentially still existing effect of smoke absorption in austral winter months in two-year mean regional distribution, even after discarding the pixels with OMI AI > 0.1.\]\(#\)](#)

The higher SEVIRI LWP values [closer to the equator outside the identified Sc cloud regime](#) are exclusively

Authors 8/16/2018 9:23 AM

Deleted: S5

Authors 8/16/2018 9:23 AM

Deleted: for more broken scenes MODIS values are lower than SEVIRI LWP.

Authors 8/16/2018 9:23 AM

Deleted: in broken scenes

575 due to a 1–2- μm overestimation in CER compared to MODIS, as SEVIRI COTs remain underestimated in almost all
 grid boxes in the study domain. The lower SEVIRI LWP values in the south of the domain outside the identified Sc
 cloud regime are due to a combination of low biases in both SEVIRI COT (~ 1) and CER ($\sim 1 \mu\text{m}$). This geographic
 variation in SEVIRI LWP could be caused by the combined contribution of two factors: (i) in thin clouds with COT
 < 4, the SEVIRI CPP algorithm weighting CER with an a priori (climatological) value of 8 μm , but MODIS
 providing the retrieved values, and (ii) the plane-parallel bias in heterogeneous scenes causing underestimated COT
 580 and overestimated CER values due to the large SEVIRI pixel size. The MODIS 3.7- μm CER values are consistently
 lower outside the Sc cloud regime than the 1.6- and 2.1- μm band retrieved CERs, likely because the latter channels
 are more strongly influenced by cloud heterogeneity and associated 3D radiative effects in broken clouds.

585 Figure 7 depicts the regional distribution of two-year mean SEVIRI and MODIS liquid fractional cloud
 cover and their differences for all-sky case. The LCF varied between 60 – 100 % within the identified Sc regime in
 both datasets; the difference in cloud fraction is within $\pm 5 \%$, with SEVIRI being smaller near the coast and larger
 further offshore. SEVIRI retrieves 5-10 % larger LCF in the mid-Atlantic, which also coincides with the largest H,
 indicating the occurrence of the most heterogeneous clouds. The typical liquid clouds here are small broken Cu
 leading to larger LCF estimates at the larger SEVIRI pixel size. Around the equator, in contrast, SEVIRI
 considerably underestimates MODIS LCF by about 10-30 %. We speculate that the frequent occurrence of ice phase
 590 clouds (deep convection, Ci) here results in overestimating the ice cloud fraction and thus underestimating the liquid
 cloud fraction, at the large SEVIRI pixel size. As expected, in the overcast only scenes (LCF $\geq 95 \%$ and COT > 3)
 SEVIRI and MODIS LCF agree within $\pm 1 \%$ (not shown).

The daytime-averaged two-year and seasonal statistics of SEVIRI and MODIS LWP are listed in Table 2,
 whereas the respective mean COTs and CERs are listed in Table S1. For overcast marine Sc clouds the two-year
 595 mean LWP is 80 g m^{-2} for SEVIRI, 84 g m^{-2} for MODIS 1.6- μm , 88 g m^{-2} for MODIS 2.1- μm , and 87 g m^{-2} for
 MODIS 3.7- μm channels. The differences in retrieved LWP values vary from 4 to 8 g m^{-2} (5–10 %), whereas the
 differences in root mean square deviation (RMSD) values vary between 16–20 g m^{-2} . The SEVIRI and MODIS
 LWP retrievals are highly correlated, with correlations > 0.9. In the aerosol-unaaffected seasons of DJF and MAM,
 the difference between SEVIRI and MODIS LWPs is within 0–5 %. In the heavily polluted months of JJA, LWP
 600 retrievals from SEVIRI are about 10 % lower than those from the MODIS 1.6- μm and 20 % lower than those from
 the MODIS 3.7- μm band. This again suggests that SEVIRI retrievals are more strongly affected by the presence of
 absorbing aerosols in the Sc regime than the corresponding MODIS 1.6- μm retrievals and that these polluted scenes

Authors 8/16/2018 9:23 AM
 Deleted: -4

Authors 8/16/2018 9:23 AM
 Deleted: these clouds similar to overcast clouds. This overestimation

Authors 8/16/2018 9:23 AM
 Deleted: .

Authors 8/16/2018 9:23 AM
 Deleted: weighing

Authors 8/16/2018 9:23 AM
 Deleted: smaller

Authors 8/16/2018 9:23 AM
 Deleted: . Also note that the SEVIRI CER overestimation in broken clouds systematically increases as the sampling height of the comparison MODIS channel gets closer to cloud top, i.e. moving from the 1.6 μm

Authors 8/16/2018 9:23 AM
 Deleted:

Authors 8/16/2018 9:23 AM
 Deleted: band. This could indicate a vertically decreasing CER profile, sometimes seen in raining/drizzling small Cu clouds, in which case the cloud-top

Authors 8/16/2018 9:23 AM
 Deleted: value, especially from the 3.7

Authors 8/16/2018 9:23 AM
 Deleted: , would underestimate the cloud layer-

Authors 8/16/2018 9:23 AM
 Deleted: .

Authors 8/16/2018 9:23 AM
 Deleted: a corresponding underestimation in LWP as well.

Authors 8/16/2018 9:23 AM
 Deleted: re

625 are not sufficiently filtered out by the OMI AI threshold. Indeed, unlike in other seasons, in JJA MODIS LWP_{1.6-μm} < MODIS LWP_{2.1-μm} < MODIS LWP_{3.7-μm}, hinting at the influence of absorbing aerosols on MODIS LWP retrievals as the 3.7-μm channel is known to be the least affected by smoke. In SON, since September is the only month strongly affected by aerosols, the comparison of SEVIRI and MODIS LWPs is better, with SEVIRI low biases of 6–12 %.

630 Figure 8 shows the density scatterplots of SEVIRI versus MODIS 1.6-μm LWPs, COTs, and CERs in the overcast Sc region for the study period. Most data points are close to the one-to-one line, but with a SEVIRI low bias; the same feature is also found in monthly and seasonal results. A low COT bias of 1 compared to all three MODIS channels, and a low CER bias of 1 μm compared to MODIS 1.6-μm and a low CER bias of 1.5 μm compared to MODIS 2.1-μm and 3.7-μm bands are observed in SEVIRI overcast retrievals. The frequency histograms of SEVIRI minus MODIS CPP difference, as well as the differences with respect to different MODIS
635 channels are shown for the all-sky case in Fig. S3 and for the overcast case in Fig. S4. The peak of the LWP absolute difference as well as the relative difference distribution is centred on zero, although the distribution is negatively skewed. Interestingly, in the all-sky case ~40 % of the data have shown negligible difference (zero LWP bias bin), whereas only about 30 % of the data have shown a negligible difference in the overcast case. Histograms of both COT and CER differences reveal that the distribution is off centered. Histograms of COT differences reveal
640 a narrow distribution, which peaks at -1 in the overcast case; however, in the all-sky case there is a broader peak between -1 and 0. Histograms of CER differences reveal wider distributions, especially when compared against the 2.1- and 3.7-μm channels, which peak at -1 μm in the overcast case; however, in the all-sky case the peak is again broader between -2 and -1 μm.

645 4.3 Diurnal cycle of SEVIRI, TMI, and MODIS cloud properties

Figures 9-11 and S5-S8 show the two-year mean and seasonal diurnal cycle of Sc cloud properties. The diurnal cycles shown here are limited to cases with AI values lower than 0.1, in order to minimize VIS/NIR retrieval biases due to biomass burning smoke (see section 4.1). Because SEVIRI retrievals (black standard and green adiabatic) are only available during daytime, TMI LWP is shown separately for day (red) and night (gray) observations, which
650 combined depict the entire 24-hour diurnal cycle. As before, the analysis is done separately for the all-sky case (solid lines with open circles) and the overcast case (dash-dotted lines with plus signs). MODIS Terra and Aqua values at 10h LST and 14h LST, respectively, are plotted as light blue color symbols (open circles or plus signs) in

Authors 8/16/2018 9:23 AM
Deleted: 5

Authors 8/16/2018 9:23 AM
Deleted: S4

Authors 8/16/2018 9:23 AM
Deleted: S6

Authors 8/16/2018 9:23 AM
Deleted: 6-7

Authors 8/16/2018 9:23 AM
Deleted: S7-S9

both cases, [for the common 1.6- \$\mu\text{m}\$ water-absorbing channel, but we report on MODIS CER for all three absorption channels in the diurnal cycle as they hint at distinct features of cloud heterogeneity and retrieval artifacts.](#)

660

For the two-year means (Fig. 9), both TMI and SEVIRI indicate a maximum LWP at 06h LST in the morning before sunrise, followed by a decrease until about 16h LST and an increase afterwards. During the night LWP continues to increase until sunrise, as indicated by the TMI night retrievals. At around 06h LST the two-year-mean all-sky LWP values are $\sim 75 \text{ g m}^{-2}$ for both TMI and SEVIRI, but they decrease to $\sim 40 \text{ g m}^{-2}$ by $\sim 14\text{h LST}$.

665

This decrease in LWP is linked to a sharp decline in COT from 11.5 to 5.5, [as well as a 20 % decrease in mean fractional cloud cover.](#) The relative variation in CER is much smaller [than that in COT during the day](#), in agreement with Zuidema and Hartmann (1995). CER increased by $2 \mu\text{m}$ in the early hours between 06h and 10h LST, stayed around 11.0-11.5 μm most of the day, and decreased by $\sim 1 \mu\text{m}$ in the late afternoon by 18h LST. As a result, the diurnal cycle of LWP was mainly driven by COT. Note that the all-sky two-year-mean TMI (red solid line circles), SEVIRI (black solid line circles), and MODIS ([light blue circles](#)) LWPs exhibit excellent agreement not only in their relative diurnal variations but also in their absolute values –the curves almost completely overlap.

670

For the overcast case, a $\sim 30 \%$ increase in COT and a slight $< 1 \mu\text{m}$ decrease in CER lead to an overall increase of $25\text{--}30 \text{ g m}^{-2}$ ($\sim 40 \%$) in mean LWP compared to the all-sky case. Apart from that, the diurnal cycles of LWP, COT, and CER are very similar between the overcast and all-sky cases. The standard SEVIRI (black dash-dot line, plus signs) and TMI day (red dash-dot line, plus signs) overcast LWPs also show very good quantitative agreement, with SEVIRI being biased high only about 5 g m^{-2} . Note that for the two-year means, adiabatic SEVIRI LWPs (green dashed line, triangles) had larger and negative biases than standard SEVIRI retrievals. As shown later, this was the consequence of the significant smoke-induced negative biases in SEVIRI retrievals in the aerosol-affected seasons of JJA and SON. In the smoke-free seasons of DJF and MAM, adiabatic SEVIRI LWPs were in better agreement with TMI microwave LWPs than were standard SEVIRI LWPs, echoing the findings of Bennartz (2007) and Seethala and Horváth (2010) for MODIS –[AMSRE](#)-E LWP comparison.

675

680

Comparing MODIS Terra (10h LST) and Aqua (14h LST) LWPs, a similar decreasing diurnal trend can be observed, except that MODIS LWPs are $5\text{--}10 \text{ g m}^{-2}$ larger than SEVIRI LWPs for the overcast case, [due to a more pronounced smoke effect \(i.e. larger negative COT biases\) in the SEVIRI 0.63- \$\mu\text{m}\$ channel than in the MODIS 0.86- \$\mu\text{m}\$ channel and also due to the larger SEVIRI pixel size \(3 km vs. 1 km\).](#) The CM SAF (2016) validation report also suggests that the coarser resolution of SEVIRI retrievals results in somewhat lower COT and LWP values compared to MODIS, [because of non-linear averaging effects \(plane-parallel albedo bias\).](#)

685

Authors 8/16/2018 9:23 AM
Deleted: with color (orange, light blue, yellow) indicating
Authors 8/16/2018 9:23 AM
Deleted: used
Authors 8/16/2018 9:23 AM
Deleted: 6

Authors 8/16/2018 9:23 AM
Deleted: .
Authors 8/16/2018 9:23 AM
Deleted: over most of this time period

Authors 8/16/2018 9:23 AM
Deleted: colored

Authors 8/16/2018 9:23 AM
Deleted: AMSRE
Authors 8/16/2018 9:23 AM
Deleted: probably
Authors 8/16/2018 9:23 AM
Deleted: difference in
Authors 8/16/2018 9:23 AM
Deleted: 1
Authors 8/16/2018 9:23 AM
Deleted: 3
Authors 8/16/2018 9:23 AM
Deleted: due to

700 Our results are consistent with Wood et al. (2002) and Painemal et al. (2012), who studied the diurnal
variation of LWP over the southeast Atlantic and southeast Pacific Sc, based on microwave and near-infrared
satellite data. Similar to our results, Painemal et al. (2012) also noted that COT rather than CER explains most of the
LWP variation. Blaskovic et al. (1990) associated the daytime decrease of LWP with the decrease of cloud
705 [geometric](#) thickness observed in their ground-based measurements, as the cloud base height increased from sunrise
till mid-afternoon, while cloud top height decreased in the late afternoon. Duyenkerke et al. (2004) found that the
diurnal variation of Sc LWP is related to the transition from a decoupled MBL during daytime to a vertically well-
mixed MBL during the night. The observed diurnal cycle of Sc is characterized by a cloud layer that gradually
thickens during the night but gets thinner during the day due to absorption of shortwave radiation and decoupling.
The latter state exhibits slightly negative buoyancy fluxes and a minimum vertical velocity variance near cloud base.
710 This implies that surface-driven, moist thermals cannot penetrate the cloud layer, while entrainment maintains a
steady supply of relatively warm and dry air from just above the inversion into the cloud layer, resulting in a distinct
LWP diurnal cycle with minimum values during the day. The diurnal cycle of LWP also consistently follows the
variation of cloud fraction in our data, [as shown in Figs. 10 and S8](#). This is in agreement with Fairall et al. (1990)
and Ciesielski et al. (2001), who observed that fractional cloudiness is maximum in the predawn hours and
715 minimum in the mid-afternoon, which is accompanied by an opposite trend in the MBL moisture with a predawn
drying and an afternoon moistening.

The seasonal mean diurnal cycles of Sc clouds are qualitatively similar to the two-year mean, except for the
aerosol affected months of JJA (Figs. [J1](#) and [S5-S8](#)). The maximum LWP tends to occur between 06h and 10h LST.
The largest diurnal variation is seen during SON, which is also the season with the greatest cloud cover. We found
720 that the relative amplitude of the two-year- and seasonal-mean LWP diurnal cycle is typically 35-40 %. Wood et al.
(2002) reported diurnal amplitudes of 15-35 % in MBL clouds using TMI data and Zuidema and Hartmann (1995)
obtained a 25 % variation in LWP over the North/South Pacific as well as South Atlantic stratus clouds using SSM/I
data for the summer months. However, Fairall et al. (1990) found larger amplitudes of 60-70 % for Californian Sc
clouds using a 17-day period of near-continuous ground-based microwave radiometer data.

725 In the all-sky case, the diurnal variation of TMI and SEVIRI LWP is in good absolute agreement within ± 5
 g m^{-2} , for all seasons and the two-year mean, except JJA. In JJA, however, a $\pm 10 \text{ g m}^{-2}$ or even slightly larger mean
difference is found between the techniques, despite the exclusion of aerosol-affected pixels with $\text{AI} > 0.1$. MODIS

Authors 8/16/2018 9:23 AM

Deleted: .

Authors 8/16/2018 9:23 AM

Deleted: 7

Authors 8/16/2018 9:23 AM

Deleted: S7-S9

Terra and Aqua mean LWPs also show excellent agreement with the corresponding SEVIRI LWPs within $\pm 5 \text{ g m}^{-2}$, for all seasons and the two-year mean.

In the overcast case, SEVIRI LWPs are $10\text{--}20 \text{ g m}^{-2}$ larger than TMI LWPs especially for the aerosol-free seasons of DJF and MAM. After applying the adiabatic correction, the biases become negligible between the datasets. For the aerosol-affected seasons of JJA and SON, the mean SEVIRI LWPs likely underestimate the actual values and hence applying adiabatic correction (reduction) worsens the comparison with TMI LWPs. In the overcast case, MODIS Terra and Aqua LWPs deviate by $5\text{--}10 \text{ g m}^{-2}$ from SEVIRI LWPs for the aerosol-free seasons, but by a larger amount of $5\text{--}20 \text{ g m}^{-2}$ for the aerosol-affected seasons due to smoke-induced biases being larger in SEVIRI than MODIS retrievals (see section 4.2).

Seasonally, COT varies between 4 and 16, typically showing a relative decrease of 40–50 % from early morning to late afternoon. Not surprisingly, the diurnal amplitude of COT is similar to that of LWP. Although the absolute value of CER varies from 7 to 12 μm between different seasons, the relative diurnal variation is negligible. The diurnal reduction in COT is likely due to the reduction in cloud fraction and cloud geometric thickness, while the variation in cloud-top CER is probably indicative of enhanced cloud-top entrainment of dry air and associated droplet evaporation. Although MODIS COTs are slightly higher in both Aqua and Terra data, the difference with SEVIRI is only about 1, while MODIS CER values are within 2 μm of the SEVIRI CER. Interestingly in the all-sky case, MODIS 1.6- and 2.1- μm CERs are 1–2 μm larger than the SEVIRI CER, and the MODIS 3.7- μm CER agrees best with SEVIRI. This suggests that in lower cloud fraction, more heterogeneous scenes, the MODIS 1.6- and 2.1- μm channels retrieve larger values (plane-parallel CER bias), whereas in SEVIRI the applied climatological weighting lowers the actual CER values and makes them more comparable to the MODIS 3.7- μm CERs, which are least affected by cloud heterogeneity and 3D effects.

5. Summary

The objective of this work was to compare LWP, COT, and CER retrievals from SEVIRI, and MODIS, and LWP from TMI, in order to quantify the effect of biomass burning smoke on passive VIS/NIR imager retrievals as well as to evaluate the diurnal cycle of South Atlantic marine stratocumulus clouds. In general, SEVIRI and TMI showed good agreement for instantaneous and domain-mean LWPs in the extensive Sc region, while the agreement in broken clouds was worse. Spatial distributions showed a correlation higher than 0.85 between the all-sky retrievals, with negligible bias on a two-year and seasonal basis for all smoke-free months. Austral winter months were heavily

Authors 8/16/2018 9:23 AM

Deleted: physical

Authors 8/16/2018 9:23 AM

Deleted:

Authors 8/16/2018 9:23 AM

Deleted: ,

Authors 8/16/2018 9:23 AM

Deleted: maritime

Authors 8/16/2018 9:23 AM

Deleted: the

Authors 8/16/2018 9:23 AM

Deleted: cloud region

Authors 8/16/2018 9:23 AM

Deleted: Boreal summer

770 smoke-affected and hence a larger bias was observed between the VIS/NIR and microwave techniques, due to an underestimation in the former. For overcast cases, the mean LWPs were ~60 % greater than the all-sky LWPs in both SEVIRI and TMI. In biomass smoke-free months, the overcast SEVIRI LWPs were higher than the corresponding TMI LWPs; however, an adiabatic correction could reduce this high bias to the 5 % level. In smoke-affected months, in contrast, the adiabatic correction, which amounts to a ~17 % reduction in VIS/NIR LWP, further increased the (negative) bias between SEVIRI and TMI. This was so because SEVIRI retrievals were already biased low by the presence of absorbing aerosols over clouds, even though aerosol index-based filtering was applied [in an attempt](#) to exclude the most polluted pixels.

775 SEVIRI and MODIS LWPs showed excellent correlation of > 0.9 in the Sc region on a two-year and seasonal basis. However, mean MODIS COTs and CERs were [both](#) 5–10 % higher in smoke-free months and 10–20 % higher in smoke-affected months than corresponding SEVIRI mean values. Interestingly, in overcast cases the relative magnitudes of MODIS CER retrievals from the 1.6-, 2.1-, and 3.7- μ m channels were different in smoke-free and smoke-affected months. Especially in JJA, the 1.6- μ m MODIS CER was significantly lower than the other two values, indicating a strong low bias in this channel due to smoke absorption. Overall, the difference between SEVIRI and MODIS LWPs was within 5 % in smoke-free seasons, but the retrieved SEVIRI LWPs were 10–25 % lower than MODIS LWPs in smoke-affected months.

785 Prompted by the above, we separately investigated the influence of absorbing aerosols over the Sc domain using aerosol index obtained from OMI. While TMI LWP showed a steep increase with AI, SEVIRI LWP showed a systematic decrease. This indicates that absorbing aerosols above liquid clouds introduce substantial negative retrieval biases in VIS/NIR cloud optical thickness and droplet effective radius, and, hence, in the deduced LWP. This bias in SEVIRI LWP increased with AI and could be as large as 40 g m⁻² in instantaneous retrievals. Neglecting aerosol-affected pixels with AI>0.1, the domain-mean TMI minus SEVIRI LWP bias could be reduced but not completely removed. Similar to SEVIRI, all three MODIS channels showed a decrease in LWP with AI, with the largest decrease occurring in the 1.6- μ m channel. The overall reduction in LWP with AI was 10–20 % in MODIS retrievals, whereas it was ~50 % in SEVIRI retrievals. [The larger sensitivity of SEVIRI retrievals to smoke can](#) partially be explained by the [difference in the](#) non-absorbing VIS channel [employed in the bispectral methods](#): the 0.86- μ m channel used for MODIS [oceanic](#) retrievals is less affected by aerosol absorption than the 0.63- μ m channel used for SEVIRI.

- Authors 8/16/2018 9:23 AM
Deleted: ; the difference
- Authors 8/16/2018 9:23 AM
Deleted: different
- Authors 8/16/2018 9:23 AM
Deleted: used
- Authors 8/16/2018 9:23 AM
Deleted: 8
- Authors 8/16/2018 9:23 AM
Deleted: 6

800 [Our finding that SEVIRI COT and CER were both lower than their \(1.6- \$\mu\text{m}\$ \) MODIS counterparts, however, is puzzling. In the absence of considerable net 3D effects, subpixel heterogeneity within the larger SEVIRI footprint would alone lead to a simultaneous underestimate of COT \(plane-parallel albedo bias\) and overestimate of CER \(plane-parallel effective radius bias; Zhang et al., 2012\) compared to the higher resolution MODIS retrievals. Indeed, such opposite-sign biases were found by Painemal et al. \(2012\) in GOES-10 COT and CER compared to](#)
805 [MODIS data in the southeast Pacific marine Sc region. Therefore, the SEVIRI COT underestimation is consistent with plane-parallel biases, while the SEVIRI CER underestimation is not.](#)

[We performed a preliminary analysis of factors that might explain these biases. Firstly, spatial heterogeneity was considered. Similar to Painemal et al. \(2013b\), we found both for SEVIRI and for MODIS a decrease in COT and increase in CER with increasing scene heterogeneity under constant TMI LWP. However, the](#)
810 [negative SEVIRI-MODIS CER difference remained a robust feature, independent of the magnitude of the scene heterogeneity. Secondly, the different view geometries of SEVIRI \(fixed\) and MODIS \(varying from orbit to orbit\) were analysed. A dependence of the SEVIRI-MODIS CER bias on the MODIS view zenith angle \(VZA\) was observed, with the bias varying between -0.5 \$\mu\text{m}\$ to -2 \$\mu\text{m}\$ and generally being lowest for the MODIS oblique backscatter view direction. Further analysis in this direction, including also the relative azimuth angle, is promising but deferred to a future study, in which also spectral and algorithmic differences between the MODIS and SEVIRI instruments and cloud property retrievals should be considered.](#)
815

[In the all-sky case, the diurnal variations of TMI and SEVIRI LWP were in good absolute agreement, being within \$\pm 5 \text{ g m}^{-2}\$ for all seasons and the two-year mean, except JJA. In JJA, the season most affected by biomass smoke, a larger mean difference was found between the techniques, although we](#)
820 [made an attempt to eliminate aerosol-affected pixels with AI > 0.1. MODIS Terra and Aqua mean LWPs also showed excellent agreement with corresponding SEVIRI LWPs in the all-sky case, differences being within \$\pm 5 \text{ g m}^{-2}\$ for all seasons and two-year means.](#)

[In the overcast case, SEVIRI LWPs were 10–20 \$\text{g m}^{-2}\$ larger than TMI LWPs especially in the smoke-free seasons of DJF and MAM. After applying an adiabatic correction to SEVIRI retrievals, however, the biases between](#)
825 [the datasets became negligible. In the smoke-affected seasons of JJA and SON, the mean SEVIRI LWPs already underestimated the TMI values due to smoke-induced retrieval biases and hence applying the adiabatic correction \(i.e. further reduction\) worsened the comparison with TMI LWPs. In the overcast case, MODIS Terra and Aqua](#)

830 LWPs differed by 5–10 g m⁻² from SEVIRI LWPs in smoke-free seasons and by a larger amount of 5–20 g m⁻² in smoke-affected seasons, due to the different magnitudes of smoke-induced biases in SEVIRI and MODIS retrievals.

Irrespective of season, both TMI and SEVIRI LWP decreased from morning to mid-afternoon, and after that a [slow](#) increase was observed. [Clouds](#) are the thickest [prior to sunrise](#) and as the day progresses the cloud layer thins due to the absorption of solar radiation and associated decoupling of the sub-cloud layer. We found that the relative amplitude of the LWP diurnal cycle is typically 30-50%, which is close to but slightly larger than the diurnal amplitude reported in most previous studies. The [diurnal](#) variation in SEVIRI LWP was mainly due to that in cloud optical thickness, while droplet effective radius showed relatively small diurnal variability. MODIS Terra (morning) and Aqua (afternoon) LWPs indicated a similar diurnal trend, but MODIS LWPs were 5–10 g m⁻² larger than SEVIRI/TMI values in the overcast case. This [was](#) maybe partly due to the plane-parallel albedo bias affecting the larger SEVIRI pixels.

840 While the discrepancies between microwave and VIS/NIR LWP retrievals in areas of broken clouds with low cloud fraction require further research to be fully resolved, our study has shown that there is a reasonable consensus between the techniques about the seasonal and diurnal cycles of LWP in nearly overcast stratocumulus fields. This lends some credibility to the VIS/NIR retrievals of the underlying cloud microphysical properties. In our opinion, SEVIRI-derived CLAAS-2 cloud property observations provide a useful resource for the evaluation of stratocumulus [diurnal](#) cycles in climate models.

Authors 8/16/2018 9:23 AM

Deleted: slight

Authors 8/16/2018 9:23 AM

Deleted: Prior to sunrise clouds

Authors 8/16/2018 9:23 AM

Deleted: temporal

Authors 8/16/2018 9:23 AM

Deleted: cloud

Acknowledgements

855 The work of the first author was funded by EUMETSAT through the CM SAF Visiting Scientist scheme and was
completed at the Royal Netherlands Meteorological Institute, De Bilt, Netherlands. SEVIRI CLAAS-2 CPP
retrievals were obtained from https://doi.org/10.5676/EUM_SAF_CM/CLAAS/V002. TMI data were produced by
Remote Sensing Systems and sponsored by the NASA Earth Sciences Program. Data are available at
www.remss.com/missions/tmi. OMI data were downloaded from <http://mirador.gsfc.nasa.gov/> and Aqua/Terra
860 MODIS Level-2 data were obtained from <http://ladsweb.nascom.nasa.gov/>. The first author acknowledges Dan
Cayan at Scripps Institution of Oceanography, La Jolla, California, for providing access to his computational facility
to finish a part of this work. We also thank two anonymous reviewers for their insightful comments and suggestions,
which greatly improved our manuscript.

865

870

875

Authors 8/16/2018 9:23 AM
Deleted: done

Authors 8/16/2018 9:23 AM
Deleted:

Authors 8/16/2018 9:23 AM
Deleted: - ... [1]

References

- 885 Abel, S. J., Walters, D. N., and Allen, G.: Evaluation of stratocumulus cloud prediction in the Met Office forecast model during VOCALS-REx, *Atmos. Chem. Phys.*, 10, 10541-10559, doi:10.5194/acp-10-10541-2010, 2010.
- Adebiyi, A. A. and Zuidema, P.: The role of the southern African easterly jet in modifying the southeast Atlantic aerosol and cloud environments, *Quart. J. Roy. Met. Soc.*, 142:697, 1574-1589, <https://doi.org/10.1002/qj.2765>, 2016.
- 890 Adebiyi, A. A., Zuidema, P., and Abel, S. J.: The Convolution of Dynamics and Moisture with the Presence of Shortwave Absorbing Aerosols over the Southeast Atlantic, *J. Climate*, Volume 28, pg.1997-2024, doi: 10.1175/JCLI-D-14-00352.1, 2015.
- Albrecht, B.A., Jensen, M. P., and Syrett, W. J.: Marine boundary layer structure and fractional cloudiness, *J. Geophys. Res.*, 100 (D7), 14209-14222, <https://doi.org/10.1029/95JD00827>, 1995.
- 895 Benas, N., Finkensieper, S., Stengel, M., van Zadelhoff, G.J., Hanschmann, T., Hollmann, R., and Meirink, J. F.: The MSG-SEVIRI-based cloud property data record CLAAS-2, *Earth Syst. Sci. Data*, 9, 415-434, <https://doi.org/10.5194/essd-9-415-2017>, 2017.
- Bennartz, R.: Global assessment of marine boundary layer cloud droplet number concentration from satellite, *J. Geophys. Res.*, 112, D02201, doi:10.1029/2006JD007547, 2007.
- 900 Bennartz, R. and Harshvardhan: Correction to “Global assessment of marine boundary layer cloud droplet number concentration from satellite”, *J. Geophys. Res.*, 112, D16302, doi:10.1029/2007JD008841, 2007.
- Bennartz, R., and Rausch, J.: Global and regional estimates of warm cloud droplet number concentration based on 13 years of AQUA-MODIS observations. *Atmos. Chem. Phys.*, doi:10.5194/acp-17-9815-2017, 2017
- Betts, A. K.: The diurnal variation of California coastal stratocumulus from two days of boundary layer soundings, *Tellus*, 42A, 302–304, <https://doi.org/10.3402/tellusa.v42i2.11879>, 1990.
- 905 Blaskovic, M., Davies, R., and Snider, J. B.: Diurnal variation of marine stratocumulus over San Nicolas island during July, 1987, C2GGR Report No. 90-6, <https://doi.org/10.1175/1520-0493>, 1990.
- Bony, S., and Dufresne, J. L.: Marine boundary layer clouds at the heart of tropical cloud feedback uncertainties in climate models, *Geophys. Res. Lett.*, 32, L20806, doi:10.1029/2005GL023851, 2005.
- 910 Borg, L. A., and Bennartz, R.: Vertical structure of stratiform marine boundary layer clouds and its impact on cloud albedo, *Geophys. Res. Lett.*, 34, L05807, doi:10.1029/2006GL028713, 2007.

- Bretherton, C. S., and Wyant, W. C.: Moisture transport, lower-tropospheric stability, and decoupling of cloud-topped boundary layers, *J. Atmos. Sci.*, 54, 148–167, <https://doi.org/10.1175/1520-0469>, 1997.
- 915 Bretherton, C. S., Austin, P. H., and Siems, S. T.: Cloudiness and Marine Boundary Layer Dynamics in the ASTEX Lagrangian Experiments. Part II: Cloudiness, Drizzle, Surface Fluxes and Entrainment, *J. Atmos. Sci.*, 52, 2724–2735, <https://doi.org/10.1175/1520-0469>, 1995.
- Burleyson, C. D., De Szoek, S. P., Yuter, S. E., Wilbanks, M., and Brewer, W. A.: Ship-Based Observations of the Diurnal Cycle of Southeast Pacific Marine Stratocumulus Clouds and Precipitation, *J. Atmos. Sci.*, 70, 3876–3894, <https://doi.org/10.1175/JAS-D-13-01.1>, 2013.
- 920 Chang, I. and Christopher, S. A.: The impact of seasonalities on direct radiative effects and radiative heating rates of absorbing aerosols above clouds, *Quart. J. Roy. Met. Soc.*, 143:704, 1395-1405, <https://doi.org/10.1002/qj.3012>, 2017.
- Cho, H. M., Zhang, Z., Meyer, K., Lebsock, M., Platnick, S., Ackerman, A. S., Di Girolamo, L., Labonnote, L., Cornet, C., Riedi, J., and Holz, R.: Frequency and causes of failed MODIS cloud property retrievals for liquid phase clouds over global oceans: Failed MODIS Cloud Retrievals. *J. Geophys. Res.: Atmospheres*, 120, Doi:10.1002/2015JD023161, 2015.
- 925 Ciesielski, P. E., Schubert, W. H., and Johnson, R. H.: Diurnal Variability of the Marine Boundary Layer during ASTEX, *J. Atmos. Sci.*, 58, 2355-2376, <https://doi.org/10.1175/1520-0469>, 2001.
- CM SAF: Algorithm Theoretical Basis Document, SEVIRI Cloud Physical Products, CLAAS Edition 2, EUMETSAT Satellite Application Facility on Climate Monitoring, SAF/CM/KNMI/ATBD/SEVIRI/PPP, Issue 2, Rev. 2, https://doi.org/10.5676/EUM_SAF_CM/CLAAS/V002, 10 June 2016b.
- 930 Coddington, O. M., Pilewskie, P., Redemann, J., Platnick, S., Russell, P. B., Schmidt, K. S., Gore, W. J., Livingston, J., Wind, G., and Vukicevic, T.: Examining the impact of overlying aerosols on the retrieval of cloud optical properties from passive remote sensing, *J. Geophys. Res.*, 115, D10211, doi:10.1029/2009JD012829, 2010.
- 935 Duykerke, P. G., De Roode, S. R., Van Zanthen, M. C., Calvo, J., Cuxart, J., Cheinet, S., Chlond, A., Grenier, H., Jonker, P. J., Köhler, M., Lenderink, G., Lewellen, D., Lappen, C., Lock, A. P., Moeng, C. Müller, F., Olmed, D., Piriou, J., Sanchez, E., and Sednev, I.: Observations and numerical simulations of the diurnal cycle of the EUROCS stratocumulus case, *Quart. J. Roy. Met. Soc.*, 130, 3269-3296, doi: 10.1256/qj.03.139, 2004.
- 940 Eastman, R., Warren, S. G., and Hahn, C. J.: Variations in cloud cover and cloud types over the ocean from surface observations, 1954–2008, *J. Climate*, 24, 5914–5934, <https://doi.org/10.1175/2011JCLI3972.1>, 2011.

- Elsaesser, G.S., O'Dell, C. W., Lebsock, M. D., Bennartz, R., Greenwald, T. J., and Wentz, F. J.: The Multisensor Advanced Climatology of Liquid Water Path (MAC-LWP), *J. Climate*, **30**, 10193–10210, <https://doi.org/10.1175/JCLI-D-16-0902.1>, 2017.
- 945 Fairall, C.W., Hare, J. E., and Snider, J. B.: An eight-month of marine stratocumulus cloud fraction, albedo, and integrated liquid water, *J. Climate*, **3**, 847-864, <https://doi.org/10.1175/1520-0442>, 1990.
- Forster, P., et al.: Changes in atmospheric constituents and in radiative forcing, in *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by S. Solomon et al., pp. 130–234, Cambridge Univ. Press, New York, 2007.
- 950 Greenwald T. J.: A 2 year comparison of AMSR-E and MODIS cloud liquid water path observations, *Geophys. Res. Lett.*, **36**, L20805, <https://doi.org/10.1029/2009GL040394>, 2009.
- Greenwald, T. J., L'Ecuyer, T. S., and Christopher, S. A.: Evaluating specific error characteristics of microwave-derived cloud liquid water products, *Geophys. Res. Lett.*, **34**, L22807, doi:10.1029/2007GL031180, 2007.
- Greenwald, T. J., Bennartz, R., Lebsock, M., & Teixeira, J.: An Uncertainty Data Set for Passive Microwave 955 Satellite Observations of Warm Cloud Liquid Water Path. *J. Geophys. Res. Atmos.*, **123**, 3668–3687, <https://doi.org/10.1002/2017JD027638>, 2018.
- Greuell, W., Van Meijgaard, E., Meirink, J. F., and Clerbaux, N.: Evaluation of model predicted top-of-atmosphere radiation and cloud parameters over Africa with observations from GERB and SEVIRI, *J. Climate*, **24**, 4015-4036, doi:10.1175/2011JCLI3856.1, 2011.
- 960 Grosvenor, D. P., Sourdeval, O., Zuidema, P., et al.: Remote sensing of droplet number concentration in warm clouds: A review of the current state of knowledge and perspectives, *Reviews of Geophysics*, Doi:10.1029/2017rg000593, 2018.
- Han Q, Rossow, W. B., and Lasis, A. A.: Near-Global Survey of Effective Droplet Radii in Liquid Water Clouds Using ISCCP Data, *J. Climate* **7**, 465-497, <https://doi.org/10.1175/1520-0442>, 1994.
- 965 Haywood, J. M., Osborne, S. R., and Abel, S. J.: The effect of overlying absorbing aerosol layers on remote sensing retrievals of cloud effective radius and cloud optical depth, *Quart. J. Roy. Met. Soc.*, **130**, 779-800, doi: 10.1256/qj.03.100, 2004.

Authors 8/16/2018 9:23 AM

Deleted: Journal of Geophysical Research: Atmospheres,

- 970 Hilburn, K., and Wentz, F. J.: Intercalibrated passive microwave rain products from the unified microwave ocean retrieval algorithm (UMORA), *J. Appl. Met. Clim.*, 47, 778–794, <https://doi.org/10.1175/2007JAMC1635.1>, 2008.
- Hobbs, P. V.: Clean air slots amid atmospheric pollution, *Nature*, 415, 861, doi:10.1038/415861a, 2002.
- Horowitz, H. M., Garland, R. M., Thatcher, M., Landman, W. A., Dedekind, Z., Van der Merwe, J.,
975 and Engelbrecht, F. A.: Evaluation of climate model aerosol seasonal and spatial variability over Africa using AERONET, *Atmos. Chem. Phys.*, 17:22, 13999-14023, <https://doi.org/10.5194/acp-17-13999-2017>, 2017.
- Horváth, Á., and Gentemann, C.: Cloud-fraction-dependent bias in satellite liquid water path retrievals of shallow, non-precipitating marine clouds, *Geophys. Res. Lett.*, 34, L22806, doi:10.1029/2007GL030625, 2007.
- Horváth, Á., and Davies, R.: Comparison of microwave and optical cloud water path estimates from TMI, MODIS,
980 and MISR, *J. Geophys. Res.*, 112, D01202, doi:10.1029/2006JD007101, 2007.
- Horváth, Á., Seethala, C., and Deneke, H.: View angle dependence of AMSR-E and MODIS cloud liquid water path retrievals in warm oceanic clouds, *J. Geophys. Res.*, 119, 8304–8328, doi:10.1002/2013JD021355, 2014.
- James, D. G.: Observations from aircraft of temperatures and humidities near stratocumulus clouds, *Quart. J. Roy. Met. Soc.*, 85, 120-130, <https://doi.org/10.1002/qj.49708536405>, 1959.
- 985 Johnson, B. T., Shine, K. P., Forster, P. M.: The semi-direct aerosol effect: Impact of absorbing aerosols on marine stratocumulus, *Quart. J. Roy. Met. Soc.*, 130, 1407-1422, <https://doi.org/10.1256/qj.03.61>, 2004.
- Kar, J., Vaughan, M., Tackett, J., Liu, Z., Omar, A., Rodier, S., Trepte, C., and Lucker, P.: Swelling of transported smoke from savanna fires over the Southeast Atlantic Ocean, *Rem. Sen. Env.*, 211, 105-111, <https://doi.org/10.1016/j.rse.2018.03.043>, 2018.
- 990 Klein S. A., and Hartmann, D. L.: The seasonal cycle of low stratiform clouds, *J. Climate*, 6, 1587-1606, <https://doi.org/10.1175/1520-0442-1993,1993>.
- Klein, S. A., Hartmann, D. L., and Norris, J.: On the Relationships among Low-Cloud Structure, Sea Surface Temperature, and Atmospheric circulation in the Summer time Northeast Pacific, *J. Climate*, 8, 1140–1155, <https://doi.org/10.1175/1520-0442-1995,1995>.
- 995 Kniffka, A., Stengel, M., Lockhoff, M., Bennartz, R., and Hollmann, R.: Characteristics of cloud liquid water path from SEVIRI onboard the Meteosat Second Generation 2 satellite for several cloud types, *Atmos. Meas. Tech.*, 7, 887-905, doi:10.5194/amt-7-887-2014, 2014.

Authors 8/16/2018 9:23 AM

Formatted: Font:Not Italic

Authors 8/16/2018 9:23 AM

Formatted: Font:Not Italic

- 1000 | Liu, Z., Liu, X., Zhang, Z., Zhao, C., Meyer, K., Rajapakshe, C., Wu, C., Yang, Z., and Penner, J. E.: (2018)
Biomass smoke from southern Africa can significantly enhance the brightness of stratocumulus over the
southeastern Atlantic Ocean, *Pro. Nat. Acad. Sci.*, 115:12, 2924-2929,
<https://doi.org/10.1073/pnas.1713703115>, 2018.
- McGill, M. J., Hlavka, D. L., Hart, W. D., Welton, E. J., and Campbell, J. R.: Airborne lidar measurements of
aerosol optical properties during SAFARI-2000, *J. Geophys. Res.*, 108, 8493, doi:10.1029/2002JD002370,
2003.
- 1005 Medeiros, B., and Stevens, B.: Revealing differences in GCM representations of low clouds, *Clim. Dyn.*, 36, 385-
399, doi:10.1007/s00382-009-0694-5, 2011.
- Medeiros, B., Williamson, D. L., Hannay, C., and Olson, J. G.: Southeast Pacific stratocumulus in the Community
Atmosphere Model, *J. Climate*, 25, 6175-6192, <https://doi.org/10.1175/JCLI-D-11-00503.1>, 2012.
- Meehl, G. A., Covey, C., Delworth, T., Latif, M., McAvaney, B., Mitchell, J. F. B., Stouffer, R. J., and Taylor, K.
1010 E.: The WCRP CMP3 multi-model dataset: A new era in climate change research, *Bull. Amer. Met. Soc.*, 88,
1383-1394, <https://doi.org/10.1175/BAMS-88-9-1383>, 2007.
- Meirink, J. F., Roebeling, R. A., and Stammes, P.: Inter-calibration of polar imager solar channels using SEVIRI,
Atm. Meas. Tech., 6, 2495-2508, doi:10.5194/amt-6-2495-2013, 2013.
- Miller, R. L.: Tropical thermostats and low cloud cover. *J. Climate*, 10, 409-440, [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0442(1997)10)
1015 0442(1997), 1997.
- Minnis, P., and Harrison, E. F.: Diurnal variability of regional cloud and clear-sky radiative parameters derived from
GOES data. Part I: Analysis method, *J. Climate*, 23, 993-1011, [https://doi.org/10.1175/1520-0450\(1984\)](https://doi.org/10.1175/1520-0450(1984)23),
1984.
- Minnis, P., Heck, P., Young, D., Fairall, C., and Snider, J.: Stratocumulus cloud properties derived from
1020 simultaneous satellite and island-based instrumentation during fire, *J. Appl. Met.*, 31, 317-339,
[https://doi.org/10.1175/1520-0450\(1992\)](https://doi.org/10.1175/1520-0450(1992)31), 1992.
- Moeng, C. H., Shen, S., and Randall, D. A.: Physical processes within the nocturnal stratus-topped boundary layer,
J. Atmos. Sci., 49, 2384-2401, [https://doi.org/10.1175/1520-0469\(1992\)](https://doi.org/10.1175/1520-0469(1992)49), 1992.
- Nakajima T. Y., Nakajima, T.: Determination of Cloud Microphysical Properties from NOAA-17 Measurements for
1025 FIRE and ASTEX regions, *J. Atmos. Sci.*, 52, 4043-4059, [https://doi.org/10.1175/1520-0469\(1995\)](https://doi.org/10.1175/1520-0469(1995)52), 1995.

- Nakajima, T., and King, M. D.: Determination of the optical thickness and effective particle radius of clouds from reflected solar radiation measurements, I, Theory, *J. Atmos. Sci.*, 47, 1878-1893, [https://doi.org/10.1175/1520-0469\(1990\)](https://doi.org/10.1175/1520-0469(1990)), 1990.
- 1030 Nicholls, S.: The dynamics of stratocumulus: Aircraft observations and comparisons with a mixed layer model, *Quart. J. Roy. Meteor. Soc.*, 110, 783–820, <https://doi.org/10.1002/qj.49711046603>, 1984.
- Norris, J. R.: Low cloud type over the ocean from surface observations. Part I: Relationship to surface meteorology and the vertical distribution of temperature and moisture, *J. Climate*, 11, 369-382, [https://doi.org/10.1175/1520-0442\(1998\)](https://doi.org/10.1175/1520-0442(1998)), 1998.
- 1035 O'Dell, C. W., Wentz, F. J., and Bennartz, R.: Cloud liquid water path from satellite-based passive microwave observations: A new climatology over the global oceans, *J. Climate*, 21, 1721-1739, <https://doi.org/10.1175/2007JCLI1958.1>, 2008.
- Painemal, D.: Entrainment rate diurnal cycle in marine stratiform clouds estimated from geostationary satellite retrievals and a meteorological forecast model, *Geophys. Res. Lett.*, Doi: 10.1002/2017GL074481, 2017.
- 1040 [Painemal, D., K. M. Xu, A. Cheng, P. Minnis, and R. Palikonda: Mean structure and diurnal cycle of southeast Atlantic boundary layer clouds: Insights from satellite observations and multiscale modeling framework simulations, *J. Climate*, 28 \(1\), 324-341, doi:10.1175/JCLI-D-14-00368.1, 2015.](#)
- Painemal, D., Kato, S, and Minnis, P.: Boundary layer regulation in the southeast Atlantic cloud microphysics during the biomass burning season as seen by the A-train satellite constellation, *J. Geophys. Res. Atmos.*, 119, 11,288–11,302, doi:10.1002/2014JD022182, 2014.
- 1045 Painemal, D., Minnis, P., Ayers, J. K., and O'Neill, L.: GOES-10 microphysical retrievals in marine warm clouds: Multi-instrument validation and daytime cycle over the southeast Pacific, *J. Geophys. Res.*, 117, D19212, doi:10.1029/2012JD017822, 2012.
- 1050 [Painemal, D., Minnis, P., and O'Neill, L.: The diurnal cycle of cloud top height and cloud cover over the Southeastern Pacific as observed by GOES-10, *J. Atmos. Sci.*, Volume 70, pages: 2393-2408, Doi:10.1175/JAS-D-12-0325.1, 2013a.](#)
- [Painemal, D., Minnis, P., and Sun-Mack, S.: The impact of horizontal heterogeneities, cloud fraction, and liquid water path on warm cloud effective radii from CERES-like Aqua MODIS retrievals, *Atmos. Chem. Phys.*, 13, 9997–10003, doi:10.5194/acp-13-9997-2013, 2013b.](#)

- 1055 Platnick, S.: Vertical photon transport in cloud remote sensing problems, *J. Geophys. Res.*, 105, 22,919-22,935,
<https://doi.org/10.1029/2000JD900333>, 2000.
- Platnick, S., Meyer, K. G., King, M. D., Wind, G., Amarasinghe, N., Marchant, B., Arnold, G. T., Zhang, Z.,
Hubanks, P. A., Holz, R. E., Yang, P., Ridgway, W. L., and Riedi, J.: The MODIS Cloud Optical and
Microphysical Products: Collection 6 Updates and Examples From Terra and Aqua, *IEEE T. Geosci. Remote*,
55, 502–525, <https://doi.org/10.1109/TGRS.2016.2610522>, 2017.
- 1060 Rahn, D., and Garreaud, R.: Marine boundary layer over the subtropical southeast Pacific during VOCALS-REx -
Part 1: Mean structure and diurnal cycle, *Atmos. Chem. Phys.*, 10, 4491-4506, doi:10.5194/acp-10-4491-
2010, 2010.
- Rajapakshe, C., Zhang, Z., Yorks, J., Yu, H., Tan, Q., Meyer, K., Platnick, S., Winker, D. M.: Seasonally
Transported Aerosol Layers over Southeast Atlantic are Closer to Underlying Clouds than Previously
1065 Reported, *Geophys. Res. Lett.*, 44, Doi:10.1002/2017GL073559, 2017.
- Rausch, J., Meyer, K., Bennartz, R., and Platnick, S.: Differences in liquid cloud droplet effective radius and number
concentration estimates between MODIS collections 5.1 and 6 over global oceans, *Atmos. Meas. Tech.*, 10,
2105–2116, <https://doi.org/10.5194/amt-10-2105-2017>, 2017.
- 1070 Randles, C. A. and Ramaswamy, V.: Direct and semi-direct impacts of absorbing biomass burning aerosol on the
climate of southern Africa: a Geophysical Fluid Dynamics Laboratory GCM sensitivity study, *Atmos. Chem.*
Phys., 10, 9819-9831, doi:10.5194/acp-10-9819-2010, 2010.
- Roebeling, R. A., Feijt, A. J., and Stammes, P.: Cloud property retrievals for climate monitoring: Implications of
differences between Spinning Enhanced Visible and Infrared Imager (SEVIRI) on METEOSAT-8 and
Advanced Very High Resolution Radiometer (AVHRR) on NOAA-17, *J. Geophys. Res.*, 111D20210
10.1029/2005jd006990, 2006.
- 1075 Roebeling, R. A., and E. Van Meijgaard, E.: Evaluation of the Daylight Cycle of Model-Predicted Cloud Amount
and Condensed Water Path over Europe with Observations from MSG SEVIRI, *J. Climate*, 22, 1749-
1766.10.1175/2008jcli2391.1, 2009.
- Rozendaal M. A., Leovy, C. B., and Klein, S. A.: An observational study of diurnal variations of marine stratiform
1080 cloud, *J. Climate*, 8, 1795-1809, [https://doi.org/10.1175/1520-0442\(1995\)](https://doi.org/10.1175/1520-0442(1995)), 1995.
- Sandu, I., Stevens, B., and Pincus, R.: On the transitions in marine boundary layer cloudiness, *Atmos. Chem. Phys.*,
10, 2377-2391, <https://doi.org/10.5194/acp-10-2377-2010>, 2010.

Authors 8/16/2018 9:23 AM

Deleted: .

- Schulz, J., Albert, P., Behr, H. D., Caprion, D., Deneke, H., Dewitte, S., Durr, B., Fuchs, P., Gratzki, A., Hechler, P.,
1085 Hollmann, R., Johnston, S., Karlsson, K. G., Manninen, T., Muller, R., Reuter, M., Riihela, A., Roebeling, R.,
Selbach, N., Tetzlaff, A., Thomas, W., Werscheck, M., Wolters, E., and Zelenka, A.: Operational climate
monitoring from space: the EUMETSAT Satellite Application Facility on Climate Monitoring (CM SAF),
Atmos. Chem. Phys., 9, 1687-1709, <https://doi.org/10.5194/acp-9-1687-2009>, 2009.
- Seethala, C., and Horváth, Á.: Global assessment of AMSR-E and MODIS cloud liquid water path retrievals in
1090 warm oceanic clouds, J. Geophys. Res., 115, D13202, doi:10.1029/2009JD012662, 2010.
- Stephens G. L., Paltridge, G. W., and Platt, C. M. R.: Radiation profiles in extended water clouds. III. Observations.
J. Atmos. Sci., 35, 2133-2141, [https://doi.org/10.1175/1520-0469\(1978\)](https://doi.org/10.1175/1520-0469(1978)), 1978.
- Torres, O., Tanskanen, A., Veihelmann, B., Ahn, C., Braak, R., Bhartia, P. K., Veefkind, P., and Levelt, P.: Aerosols
and surface UV products from Ozone Monitoring Instrument observations: An overview, J. Geophys. Res.,
1095 112, D24S47, doi:10.1029/2007JD008809, 2007.
- Watts P. D., Mutlow C. T., Baran A. J., and Zavody A. M.: "Study on Cloud Properties derived from Meteosat
Second Generation Observations," Final Report, EUMETSAT ITT no. 97/181, 1998.
- Wentz, F. J.: A 17-Yr Climate Record of Environmental Parameters Derived from the Tropical Rainfall Measuring
Mission (TRMM) Microwave Imager, J. Climate, 28, 6882-6902, <https://doi.org/10.1175/JCLI-D-15-0155.1>,
1100 2015.
- Wentz, F. J., and Spencer, R.: SSM/I rain retrievals within a unified all-weather ocean algorithm, J. Atmos. Sci., 55,
1613-1627, [https://doi.org/10.1175/1520-0469\(1998\)](https://doi.org/10.1175/1520-0469(1998)), 1998.
- Wentz, F. J.: A well-calibrated ocean algorithm for special sensor microwave/imager, J. Geophys. Res., 102, 8703-
8718, <https://doi.org/10.1029/96JC01751>, 1997.
- 1105 Wilcox, E. M., Harshvardhan, and Platnick, S.: Estimate of the impact of absorbing aerosol over cloud on the
MODIS retrievals of cloud optical thickness and effective radius using two independent retrievals of liquid
water path, J. Geophys. Res., 114, D05210, doi:10.1029/2008JD010589, 2009.
- Wilcox, E. M.: Stratocumulus cloud thickening beneath layers of absorbing smoke, Aerosol, Atmos. Chem. Phys.,
10, 11769-11777, <https://doi.org/10.5194/acp-10-11769-2010>, 2010.
- 1110 Wilson, C. A., and Mitchell, J. F. B.: Diurnal variation and cloud in a general circulation model, Quart. J. Roy. Met.
Soc., 112, 347-369, <https://doi.org/10.1002/qj.49711247205>, 1986.

- Wood, R.: Stratocumulus Clouds, *Monthly Weather Review*, 140(8), 2373–2423, doi:10.1175/MWR-D-11-00121.1, 2012.
- 1115 Wood, R., and Bretherton, C. S.: On the relationship between stratiform low cloud cover and lower-tropospheric stability, *J. Climate*, 19, 6425–6432, <https://doi.org/10.1175/JCLI3988.1>, 2006.
- Wood, R., Bretherton, C. S., and Hartmann, D. L.: Diurnal cycle of liquid water path over the subtropical and tropical oceans, *Geophys. Res. Lett.*, 29 (23), 2092, doi:10.1029/2002GL015371, 2002.
- Wood, R., and Hartmann, D. L.: Spatial Variability of Liquid Water Path in Marine Low Cloud: The Importance of Mesoscale Cellular Convection, *J. Climate*, 19, 1748-1764, <https://doi.org/10.1175/JCLI3702.1>, 2006.
- 1120 Zelinka, M. D., Randall, D. A., Webb, M. J., and Klein, S. A.: Clearing clouds of uncertainty, *Nature Climate Change* 7, 674-678, 10.1038/nclimate3402, 2017.
- Zhang, Z., Ackerman, A. S., Feingold, G., Platnick, S., Pincus, R., and Xue, H.: Effects of cloud horizontal inhomogeneity and drizzle on remote sensing of cloud droplet effective radius: Case studies based on large-eddy simulations, *J. Geophys. Res. Atmos.*, 117, D19208, <https://doi.org/10.1029/2012jd017655>, 2012.
- 1125 | [Zhang, Z. and Platnick, S.: An assessment of differences between cloud effective particle radius retrievals for marine water clouds from three MODIS spectral bands, *J. Geophys. Res.*, 116, D20215, <https://doi.org/10.1029/2011jd016216>, 2011.](https://doi.org/10.1029/2011jd016216)
- Zuidema P., and Hartmann, D. L.: Satellite determination of stratus cloud microphysical properties, *J. Climate*, 8, 1638-1657, [https://doi.org/10.1175/1520-0442\(1995\)](https://doi.org/10.1175/1520-0442(1995)), 1995.
- 1130 Zuidema, P., Redemann, J., Haywood, J., Wood, R., Piketh, S., Hipondoka, M., and Formenti, P.: Smoke and clouds above the southeast Atlantic: Upcoming field campaigns probe absorbing aerosol's impact on climate, *Bull. Am. Met. Soc.*, 97: 1131–1135, <https://doi.org/10.1175/BAMS-D15-00082.1>, 2016.

1135

Authors 8/16/2018 9:23 AM

Deleted: -

TABLES

1140 **Table 1.** Two-year mean and seasonal statistics of collocated SEVIRI and TMI LWP retrievals for rain-free, ice-free, smoke-free
 (AI_v < 0.1), COT > 3, and overcast (LCF ≥ 95%) grid cells over the marine stratocumulus region. LWP means, biases (SEVIRI-
 TMI), and Root Mean Square Differences (RMSD) are given in g m⁻². COT means and CER means (in micron) are also
 1145 tabulated. The values in brackets are statistics without filtering for LCF ≥ 95% and COT > 3, i.e., for the all-sky case. “adb”
 refers to the overcast LWP calculation assuming adiabatic clouds.

Authors 8/16/2018 9:23 AM
 Deleted: <

	JJA	SON	DJF	MAM	Two-year
Stratocumulus (SEVIRI vs. TMI)					
SEVIRI LWP	73 (48)	87 (63)	92 (52)	83 (41)	84 (53)
SEVIRI LWP adb	61	72	76	69	70
TMI LWP	82 (57)	82 (62)	76 (45)	73 (39)	80 (53)
SEVIRI-TMI LWP	-9 (-9)	5 (1)	16 (7)	10 (2)	4 (0)
SEV adb -TMI LWP	-21	-10	0	-4	-10
RMSD	31 (26)	28 (26)	22 (21)	21 (20)	28 (24)
#Samples	1.6E+5 (3.2E+5)	3.3E+5 (5.2E+5)	1.4E+5 (3.2E+5)	9.1E+4 (2.6E+5)	7.3E+5 (1.4E+6)
Correlation	0.81 (0.87)	0.86 (0.89)	0.92 (0.93)	0.93 (0.92)	0.86 (0.89)
SEVIRI COT	10.2 (6.8)	11.0 (8.2)	10.7 (6.6)	10.3 (5.7)	10.7 (7.0)
SEVIRI CER	9.4 (10.8)	10.6 (11.2)	11.6 (11.7)	10.8 (11.0)	10.6 (11.2)

1150

1155

1160

1165

Table 2. Two-year mean and seasonal statistics of collocated SEVIRI and MODIS retrievals in rain-free, ice-free, smoke-free ($AL \leq 0.1$), $COT > 3$, and overcast ($LCF \geq 95\%$) grid cells over the marine stratocumulus region. LWP means, biases (MODIS-SEVIRI), and Root Mean Square Differences (RMSD) are given in $g\ m^{-2}$. Corresponding COT means and CER means are tabulated in Table S1. The values in brackets are statistics without filtering for $LCF \geq 95\%$ and $COT > 3$, i.e., for the all-sky case. “S” and “M” refer SEVIRI and MODIS.


	JJA	SON	DJF	MAM	Two-year
Stratocumulus (SEVIRI vs. MODIS)					
SEVIRI LWP	71 (44)	81 (58)	88 (45)	81 (39)	80 (48)
MODIS 1.6 LWP	79 (46)	86 (59)	88 (45)	84 (41)	84 (49)
MODIS 2.1 LWP	85 (48)	90 (60)	89 (44)	85 (40)	88 (50)
MODIS 3.7 LWP	88 (51)	89 (59)	84 (42)	82 (38)	87 (49)
M 1.6 – S LWP	8 (2)	5 (1)	0 (0)	3 (2)	4 (1)
M 2.1 – S LWP	14 (4)	9 (2)	1 (-1)	4 (1)	8 (2)
M 3.7 – S LWP	17 (7)	8 (1)	-4 (-3)	1 (-1)	7 (1)
RMSD 1.6	17 (21)	16 (20)	17 (18)	16 (18)	17 (19)
RMSD 2.1	16 (21)	15 (19)	17 (18)	16 (19)	16 (19)
RMSD 3.7	19 (20)	19 (20)	18 (19)	18 (17)	20 (19)
#Samples	3.7E+5 (9.6E+5)	6.4E+5 (1.4E+6)	2.4E+5 (8.2E+5)	2.0E+5 (7.8E+5)	1.5E+6 (4.0E+6)
Corr. LWP 1.6	0.95 (0.91)	0.95 (0.93)	0.95 (0.94)	0.96 (0.94)	0.95 (0.93)
Corr. LWP 2.1	0.95 (0.91)	0.95 (0.93)	0.95 (0.94)	0.96 (0.93)	0.95 (0.93)
Corr. LWP 3.7	0.94 (0.92)	0.93 (0.92)	0.94 (0.93)	0.95 (0.95)	0.93 (0.93)

1170

1175

FIGURES


1180




1185

Figure 1. Spatial distribution of (a) OMI ultraviolet aerosol index, (b) SEVIRI minus TMI liquid water path bias, (c) SEVIRI liquid water path, and (d) TMI liquid water path, averaged for JAS in 2011 and 2012 for overcast ($LCF \geq 95\%$ and $COT > 3$) rain- and ice-free conditions. The black contour denotes the identified stratocumulus region.

Authors 8/16/2018 9:23 AM




Deleted:




1195 **Figure 2.** OMI aerosol index versus (a) SEVIRI, TMI and MODIS LWPs, (b) SEVIRI LWP biases compared to TMI and
 1200 MODIS, (c) SEVIRI and MODIS COTs, (d) SEVIRI – MODIS COT biases, (e) SEVIRI and MODIS CER, and (f) SEVIRI –
 MODIS CER biases, over the overcast Sc region for JAS 2011 and JAS 2012 for rain- and ice-free conditions. Solid lines
 correspond to SEVIRI vs. TMI comparison, whereas dash-dotted lines correspond to SEVIRI vs. MODIS comparison. The label
 “SEV” refers SEVIRI values at MODIS collocations.

Authors 8/16/2018 9:23 AM



Deleted:

... [2]




1205

1210


1215

1220

Figure 3. Spatial distribution of SEVIRI liquid water path biases (a, d, g), cloud optical thickness biases (b, e, h), and droplet effective radius biases (c, f, i), compared to MODIS 1.6-, 2.1-, and 3.7- μm channel retrievals, respectively, averaged for JAS 2011 and JAS 2012 for overcast ($\text{LCF} \geq 95\%$ and $\text{COT} > 3$) rain- and ice-free conditions.




1225 | **Figure 4.** Two-year mean map of (a) SEVIRI minus TMI LWP bias, (b) SEVIRI LWP, (c) TMI LWP, (d) SEVIRI COT, (e) SEVIRI 1.6- μm CER, for the overcast case ($\text{LCF} \geq 95\%$ and $\text{COT} > 3$). The solid black contour denotes the identified Sc region. Rain-, ice-, and smoke-free conditions were applied.





1230

1235




1240 | **Figure 5.** Scatter density plot of SEVIRI versus TMI liquid water path for the overcast case ($\text{LCF} \geq 95\%$ and $\text{COT} > 3$) in two years of data. Rain-, ice-, and smoke-free conditions were applied.






1245

Figure 6. Two-year mean map of (a) SEVIRI LWP, (b) SEVIRI COT, (c) SEVIRI CER, (d, g, j) SEVIRI minus MODIS liquid water path biases, (e, h, k) SEVIRI minus MODIS cloud optical thickness biases, (f, i, l) SEVIRI minus MODIS droplet effective radius biases for the overcast case (LCF \geq 95% and COT $>$ 3) in ice- and smoke-free conditions.




1255

Figure 7. Two-year mean map of liquid fraction of cloud amount of (a) SEVIRI, (b) MODIS, and (c) SEVIRI-MODIS bias for all-sky case. Contours represent the heterogeneity measure H , computed from 3-km resolution SEVIRI 0.63- μm reflectances.




1260

Figure 8. Scatter density plot of SEVIRI versus MODIS 1.6-μm liquid water path, cloud optical thickness, and effective radius in the overcast case (LCF ≥ 95% and COT > 3) in two years of data. Rain-, ice-, and smoke-free conditions were applied.



Authors 8/16/2018 9:23 AM





Deleted: 6

1265

1270

Figure 9. Two-year mean diurnal cycle of cloud properties over the Sc region, both for all-sky and overcast ($LCF \geq 95\%$ and $COT > 3$) cases: (a) SEVIRI and TMI LWPs, (b) SEVIRI minus TMI LWP bias, (c) SEVIRI COT, and (d) SEVIRI 1.6- μm CER. MODIS Terra and Aqua values are also plotted. Rain-, ice-, and smoke-free conditions were applied.

1275



1280

1285

1290

1295

[Figure 10. Two-year mean diurnal cycle of SEVIRI LCF over the Sc region for the all-sky case. MODIS Terra and Aqua values are also plotted. Rain-, ice-, and smoke-free conditions were applied.](#)




Figure 11. Seasonal mean diurnal cycle of SEVIRI and TMI LWPs over the Sc region, both for all-sky and overcast ($\text{LCF} \geq 95\%$ and $\text{COT} > 3$) cases: (a) DJF, (b) MAM, (c) JJA, and (d) SON for December 2010 to November 2012. MODIS Terra and Aqua values are also plotted. Rain-, ice-, and smoke-free conditions were applied.

Authors 8/16/2018 9:23 AM
Deleted: -

Supplemental Materials

Table S1. Two-year mean and seasonal statistics of collocated SEVIRI and MODIS retrievals in rain-free, ice-free, smoke-free

($AI_{\lambda} < 0.1$), $COT > 3$, and overcast ($LCF \geq 95\%$) grid cells over the marine stratocumulus region. COT means and CER means (in micron) are listed. The values in brackets are statistics without filtering for $LCF \geq 95\%$ and $COT > 3$, i.e., for the all-sky case.

	JJA	SON	DJF	MAM	Two-year
Stratocumulus (SEVIRI vs. MODIS)					
SEVIRI COT	10.2 (5.9)	10.3 (7.1)	9.9 (5.2)	9.9 (4.8)	10.2 (6.0)
MODIS 1.6 COT	11.3 (7.3)	11.3 (8.3)	10.8 (6.3)	10.8 (5.9)	11.1 (7.2)
SEVIRI CER	8.8 (10.2)	10.2 (11.1)	11.6 (11.4)	10.5 (10.7)	10.1 (10.9)
MODIS 1.6 CER	10.3 (12.6)	11.5 (12.9)	12.2 (13.4)	11.4 (13.2)	11.3 (13.0)
MODIS 2.1 CER	11.1 (13.8)	11.9 (13.3)	12.3 (13.4)	11.6 (13.6)	11.7 (13.5)
MODIS 3.7 CER	11.5 (12.6)	11.7 (12.5)	11.6 (11.9)	11.2 (11.8)	11.6 (12.2)
Correl. COT 1.6	0.96 (0.93)	0.97 (0.96)	0.96 (0.95)	0.96 (0.94)	0.96 (0.95)
Correl. CER 1.6	0.93 (0.73)	0.90 (0.76)	0.92 (0.60)	0.92 (0.62)	0.92 (0.70)
Correl. CER 2.1	0.90 (0.77)	0.87 (0.79)	0.91 (0.62)	0.92 (0.67)	0.89 (0.72)
Correl. CER 3.7	0.86 (0.81)	0.78 (0.76)	0.87 (0.64)	0.87 (0.74)	0.80 (0.74)

Discussion: Frequency histograms of SEVIRI – MODIS LWP, COT, and CER difference, as well as, the differences relative to MODIS LWP, COT, and CER for the overcast condition aggregated during JAS 2011 and JAS 2012 are shown in Fig. S1. The histogram of SEVIRI – MODIS COT differences revealed that the peak of the distribution is off zero with ~35 % of the data falling into the -1 bin. Only ~17 % of data showed mean zero difference, while ~23 % of data showed a difference of -2. The SEVIRI COT relative to MODIS COT was within 10 % for 36 % of the data, within 20 % for 80 % of the data, and within 30 % for 95 % of the retrievals. Overall, SEVIRI COT appeared to be low by ~1 compared to MODIS COT.

SEVIRI CER retrieved in the 1.6- μm channel was compared with MODIS CERs retrieved in three water absorbing channels at 1.6-, 2.1-, and 3.7- μm . Compared to the 1.6- μm MODIS CERs, ~70 % of SEVIRI CERs have a mean difference of -1.5 μm . Compared to the 2.1- and 3.7- μm MODIS CERs, the difference histograms indicate larger differences: ~55 % and ~50 % of SEVIRI CERs have a difference of -2.5 μm , respectively. Although SEVIRI CERs are biased low compared to all three MODIS CERs, the ~1 μm additional low bias relative to the 2.1- and 3.7- μm CERs likely indicates much smaller smoke-induced retrieval artifacts in these two channels. In general, the

Authors 8/16/2018 9:32 AM

Deleted: <

Authors 8/16/2018 9:32 AM

Deleted: MODIS 2.1 COT .

11.4 (7.2) -

11.3 (8.2) -

10.8 (6.2) -

10.8 (5.8) -

11.2 (7.1) -

... [1]

Authors 8/16/2018 9:32 AM

Formatted: Line spacing: 1.5 lines

Authors 8/16/2018 9:32 AM

Formatted: Line spacing: 1.5 lines

Authors 8/16/2018 9:32 AM

Formatted: Line spacing: 1.5 lines

Authors 8/16/2018 9:32 AM

Formatted: Line spacing: 1.5 lines

Authors 8/16/2018 9:32 AM

Formatted: Line spacing: 1.5 lines

Authors 8/16/2018 9:32 AM

Formatted: Line spacing: 1.5 lines

Authors 8/16/2018 9:32 AM

Deleted: -

... [2]

Authors 8/16/2018 9:32 AM

Deleted: -

... [3]

Authors 8/16/2018 9:32 AM

Deleted: -

0.96 (0.95) -

... [4]

Authors 8/16/2018 9:32 AM

Deleted: 0.96 (0.95) -

... [5]

Authors 8/16/2018 9:32 AM

Deleted: -

... [6]

Authors 8/16/2018 9:32 AM

Deleted: -

... [7]

Authors 8/16/2018 9:32 AM

Deleted: -


Authors 8/16/2018 9:32 AM


Deleted: .

CER retrievals from SEVIRI tend to be lower than corresponding retrievals from the three MODIS channels, with SEVIRI having about 1.5 μm to 2.5 μm lower CER values.

45 The SEVIRI minus MODIS LWP distributions peak at about -10 g m^{-2} irrespective of the MODIS channel used for the retrieval. The differences between MODIS 1.6- μm and SEVIRI retrievals are within 10 % for about 30 % of SEVIRI pixels, within 20 % for about 60 % of the SEVIRI pixels, and within 30 % for about 80 % of the SEVIRI pixels. However, differences between SEVIRI and MODIS 2.1- μm and 3.7- μm channel retrievals are larger, with relative differences being smaller than 10 % for about 22 % of the SEVIRI pixels against MODIS 2.1-
50 μm and for about 16 % of the SEVIRI pixels against MODIS 3.7- μm values.

The frequency histograms of SEVIRI – MODIS LWP, COT, and CER differences, as well as the difference with respect to different MODIS channels [for the 2-year aggregate](#) are shown in Fig. [S3](#) (all-sky case) and Fig. [S4](#) (overcast case). The peak of the LWP absolute/relative difference distribution is centred on zero, although the distribution is negatively skewed. Interestingly, in the all-sky case ~ 40 % of the data have shown negligible difference (zero LWP bias bin), whereas, only about 30 % of the data have shown a negligible difference in the overcast case. About 20–30 % of the data have fallen into the LWP difference bin of -10 g m^{-2} in either cases. In the overcast case, ~ 40 % of the data have shown a relative LWP difference < 10 % and ~ 90 % of the data have shown a relative LWP difference < 30 %; however, for the all-sky case, only about 25 % and 60 % of the data have shown relative LWP differences < 10 % and < 30 %. Respectively, about 48 %, 84 %, 95 % of the observations show relative COT differences within 10 %, 20 %, and 30 % in the overcast case. Similarly, about 90 % of the observations show relative CER differences within 30 % in the overcast case. Histograms of both COT and CER differences reveal that the distribution is off centered. Histograms of COT differences reveal a narrow distribution which peaks at -1 especially in the overcast case; however in the all-sky case a broader peak is noticed between -1 and 0. Histograms of CER differences reveal wider distributions (especially when compared against the 2.1- and
60 3.7- μm channels), which peak at -1 μm in the overcast case; however, in the all-sky case a broader peak is noticed between -2 μm and -1 μm .





75

Figure S1. Histogram of SEVIRI – MODIS liquid water path differences (a), cloud optical thickness differences (c), and droplet effective radius differences (e), as well as, histogram of SEVIRI – MODIS LWP, COT, CER differences relative to MODIS LWP (b), COT (d), and CER (f) for JAS 2011 and JAS 2012 for overcast ($LCF \geq 95\%$ and $COT > 3$) rain- and ice-free conditions.

80

Authors 8/16/2018 9:32 AM
Formatted: Font:9 pt, Bold
 Authors 8/16/2018 9:32 AM
Formatted: Justified

Authors 8/16/2018 9:32 AM
Formatted: Left




Figure S2. Two-year mean map of (a) SEVIRI minus TMI LWP difference, (b) SEVIRI LWP, (c) TMI LWP, (d) SEVIRI COT, (e) SEVIRI 1.6- μm CER, for the all-sky case. The solid black contour denotes the identified Sc region. Rain-, ice-, and aerosol-free conditions were applied.






Figure S3. Histogram of SEVIRI – MODIS liquid water path differences (a), cloud optical thickness differences (c), and droplet effective radius differences (e), as well as, histogram of SEVIRI – MODIS LWP, COT, CER differences relative to MODIS LWP (b), COT (d), and CER (f) for December 2010 to November 2012 for the all-sky case with rain- and ice-free conditions.



100

105

Figure S4. Histogram of SEVIRI – MODIS liquid water path differences (a), cloud optical thickness differences (c), and droplet effective radius differences (e), as well as, histogram of SEVIRI – MODIS LWP, COT, CER differences relative to MODIS LWP (b), COT (d), and CER (f) for December 2010 to November 2012 for the overcast case ($LCF \geq 95\%$ and $COT > 3$) in rain- and ice-free conditions.


Authors 8/16/2018 9:32 AM
Formatted: Font:Not Bold

Authors 8/16/2018 9:32 AM
Formatted: Left, Line spacing: single

Authors 8/16/2018 9:32 AM
Formatted: Font:12 pt, Not Bold

Authors 8/16/2018 9:32 AM
Deleted: all-sky case with rain- and ice-free conditions. -

... [9]



115

120

125

Figure S5. Seasonal mean diurnal cycle of SEVIRI LWP bias compared to TMI as well as Terra and Aqua MODIS, over the Sc region, both for all-sky and overcast-cases ($LCF \geq 95\%$ and $COT > 3$): (a) DJF, (b) MAM, (c) JJA, and (d) SON of the study period. Rain-, ice-, and smoke-free conditions were applied.





Figure S6. Seasonal mean diurnal cycle of SEVIRI and Terra and Aqua MODIS cloud optical thicknesses over the Sc region, both for all-sky and overcast-cases ($LCF \geq 95\%$ and $COT > 3$): (a) DJF, (b) MAM, (c) JJA, and (d) SON of the study period.


Authors 8/16/2018 9:32 AM




Deleted:

Authors 8/16/2018 9:32 AM

Deleted: S8




145



150

Figure S7. Seasonal mean diurnal cycle of SEVIRI and Terra and Aqua MODIS cloud droplet effective radius over the Sc region, both for all-sky and overcast-cases ($LCF \geq 95\%$ and $COT > 3$): (a) DJF, (b) MAM, (c) JJA, and (d) SON of the study period.

Authors 8/16/2018 9:32 AM
Deleted: S9



155

160

165

[Figure S8. Seasonal mean diurnal cycle of SEVIRI and Terra and Aqua MODIS liquid cloud fraction over the Sc region, for all-sky case: \(a\) DJF, \(b\) MAM, \(c\) JJA, and \(d\) SON of the study period.](#)