

Estimation of black carbon emissions from Siberian fires using satellite observations of absorption and extinction optical depths

Igor B. Kononov et al.

Correspondence to: Igor B. Kononov (konov@appl.sci-nnov.ru)

Here, we report several supplementary tests that were designed to evaluate the sensitivity of the empirical AAOD parameterization (see Eq. 4) discussed in Sect. 2.2.3 with respect to possible biases in the simulated AAOD and AOD data for the "bgr" scenario (see Eq. 5). The different test cases involved different $AAOD_b$ and AOD_b values that were modified by applying to them the constant scaling factors, s_1 and s_2 , respectively.

First, we considered the case where the both s_1 and s_2 were equal zero. That is, an impact of the "background" aerosol on the columnar SSA values was entirely disregarded. This case is analogous to one addressed in Kononov et al. (2017a), except that the data selection criteria were different (see Sect. 2.2.3). The estimate of κ_I obtained in this case (see Fig. S1) is 15 % lower than that for the base case (see Fig. 4). The difference can be explained by the uncertainty at the 90th percentile confidence level. Using the parameterization presented in Fig. S1 for estimation of BB BC emission estimation procedure would accordingly result in about 15 % larger top-down estimates of BC emissions than those reported in Sect. 3.5.




Figure S1: The AAOD/AOD ratio as a function of the EC/(EC+OC) ratio. Both ratios shown by red crosses have been derived from the observations at the AERONET sites as explained in Sect. 2.2.3, except that the $AAOD_b$ and AOD_b values involved in Eq. (5) were zeroed. A linear regression fitted with the ODR method and its 68.3 % (1-sigma) confidence intervals are shown by solid and dashed blue lines, respectively.

In the second test, s_1 was equal to 0.5 or 1.5 ($AAOD_b$ was increased or decreased by 50 %), while s_2 was equal to 1.0. The results indicate (see Fig. S2) that the parameterization is quite insensitive to the large changes in the background AAOD values. Although we cannot estimate possible biases in $AAOD_b$, the fact that the decrease of the $AAOD_b$ values decreases the uncertainty of both κ_I and κ_2 may be regarded as evidence that $AAOD_b$ is biased positively.

Finally, in the third test, the $AAOD_b$ values were kept constant (s_1 was equal 1.0) but the AOD_b values were increased or decreased by 50 % (s_2 was set to be 0.5 or 1.5). The decrease of AOD_b results in a rather small decrease of κ_I (by 8 %), but the increase of AOD_b leads to a substantial increase (by 20 %) of the same coefficient. Therefore, if AOD_b is strongly underestimated, the AAOD/AOD ratio may also be considerably underestimated in our simulations, while the BB BC emissions obtained in our analysis may be accordingly overestimated. However, although we do not have any information about probable biases in AOD_b at the AERONET sites considered in this study, there is evidence (see Sect. 3.6) that AOD_b is

actually overestimated by about 40 % on average for the whole study region. Therefore, the underestimation of AAOD/AOD ratio by the parameterization presented used in our analysis (see Fig. 4) is possible but not likely.




Figure S2: The same as in Fig. S1, but with the original AOD_b values and with the AAOD_b values (see Eq. (5)) that were (a) decreased by a factor 0.5 and (b) increased by a factor of 1.5.




Figure S3: The same as in Figs. S1 and S2, but with the original AAOD_b values and with the AOD_b values that were (a) decreased by a factor 0.5 and (b) increased by a factor of 1.5.