

Supplemental Information

Is there an aerosol signature of cloud processing?

Barbara Ervens^{1,2}, Armin Sorooshian^{3,4}, Abdulmonam M. Aldhaif³, Taylor Shingler^{5,6}, Ewan Crosbie^{5,6}, Luke Ziembra⁶, Pedro Campuzano-Jost^{2,7}, Jose L. Jimenez^{2,7}, Armin Wisthaler^{8,9}

¹ NOAA/ESRL/Chemical Sciences Division, Boulder, CO, USA

² CIRES, University of Colorado, Boulder, CO, USA

³ Department of Chemical and Environmental Engineering, University of Arizona, Tucson, AZ, USA

⁴ Department of Hydrology and Atmospheric Sciences, University of Arizona, Tucson, AZ, USA

⁵ Science Systems and Applications, Inc., Hampton, VA, USA

⁶ NASA Langley Research Center, Hampton, VA, USA

⁷ Department of Chemistry and Biochemistry, University of Colorado, Boulder, Colorado, USA

⁸ Department of Chemistry, University of Oslo, Oslo, Norway

⁹ Institute for Ion Physics and Applied Physics, University of Innsbruck, Innsbruck, Austria

Correspondence to: barbara.ervens@noaa.gov and armin@email.arizona.edu

This supplemental information includes:

Table S1: Initial gas phase mixing ratios [ppb] used as input for the box and parcel models. Data are from SEACR⁴S for six different air masses.

Figure S1: Predicted relative mass concentration increase due to cloud processing in six air masses as identified during SEAC⁴RS. Black symbols show the measured, initial size distributions; colored symbols are model results, color-coded by log ([absolute mass increase / $\mu\text{g m}^{-3}$]).

Figure S2: Predicted relative mass concentration increase due to cloud processing in six air masses as identified during SEAC⁴RS. Black symbols show the measured, initial size distributions; colored symbols are model results, color-coded by O/C ratio.

Table S1: Initial gas phase mixing ratios [ppb]

	Marine	Urban	Biomass Burning	Agric biomass burning	Back-ground	Biogenic
SO ₂	0.43	0.56	0.252	1.44	0.64	0.582
NO	0.028	0.2	0.137	0.352	0.112	0.062
NO ₂	0.066	1.196	0.311	2.37	0.521	0.283
HCHO	0.883	2.64	4.04	14.4	2.8	4.2
H ₂ O ₂	1.172	2.58	3.72	4.3	2.6	2.8
O ₃	27	57	68	63	57	50
Isoprene	0.003	0.14	0.18	1.13	0.147	1.61
Monoterpenes	0.007	0.03	0.098	0.328	0.039	0.123
Methylvinyl ketone ¹⁾	0.008	0.124	0.113	0.33	0.18	1.1
Methacrolein ¹⁾	0.008	0.124	0.113	0.33	0.18	1.1
Toluene	0.009	0.07	0.125	0.243	0.041	0.056
Xylene ²⁾	0.005	0.035	0.067	0.12	0.02	0.028
Ethylene	0.0486	0.215	0.842	2.68	0.212	0.191

¹⁾ The total of methyl vinyl ketone and methacrolein was measured by PTRMS. For model purposes, the total was equally split between the two compounds.

²⁾ Xylene measurements were not available. The mixing ratio was assumed to be 50% toluene.




Figure S 1: Predicted relative mass concentration increase due to cloud processing in six air masses as identified during SEAC⁴RS. Black symbols show the measured, initial size distributions; colored symbols are model results, color-coded by the absolute increased in mass




Figure S 2: Predicted relative mass concentration increase due to cloud processing in six air masses as identified during SEAC⁴RS. Black symbols show the measured, initial size distributions; colored symbols are model results, color-coded by O/C ratio.