



A comprehensive organic nitrate chemistry: insights into the lifetime of atmospheric organic nitrates

Azimeh Zare¹, Paul S. Romer¹, Tran Nguyen², Frank N. Keutsch^{3,a}, Kate Skog^{3,b}, Ronald C. Cohen^{1,4}

¹Department of Chemistry, University of California Berkeley, Berkeley, CA, USA

5 ²College of Agricultural and Environmental Sciences, University of California, Davis, CA, USA

³Department of Chemistry, University of Wisconsin-Madison, Madison, WI, USA

^anow at: School of Engineering and Applied Sciences and Department of Chemistry & Chemical Biology, Harvard University, Cambridge, MA, USA

^bnow at: Department of Chemical & Environmental Engineering, Yale University, New Haven, CT, USA

10 ⁴Department of Earth and Planetary Sciences, University of California Berkeley, Berkeley, CA, USA

Correspondence to: Ronald C. Cohen (rc Cohen@berkeley.edu)

Abstract. Organic nitrate chemistry is the primary control over the lifetime of nitrogen oxides (NO_x) in rural and remote continental locations. As NO_x emissions decrease, organic nitrate chemistry becomes increasingly important to urban air quality. However, the lifetime of individual organic nitrates and the reactions that lead to their production and removal remain relatively poorly constrained, causing organic nitrates to be poorly represented by models. Guided by recent laboratory and field studies, we developed a detailed gas phase chemical mechanism representing most of the important individual organic nitrates. We use this mechanism within the WRF-Chem model to describe the role of organic nitrates in nitrogen oxide chemistry and in comparisons to observations. We find the daytime lifetime of total organic nitrates with respect to all loss mechanisms to be 2.6 h in the model. This is consistent with analyses of observations at a rural site in central Alabama during the Southern Oxidant and Aerosol Study (SOAS) in summer 2013. The lifetime of the first-generation organic nitrates is ~2 h versus the 3.2 h lifetime of secondary nitrates produced by oxidation of the first-generation nitrates. The different generations are subject to different losses, with dry deposition to the surface dominant loss process for the second-generation organic nitrates, and chemical loss dominant for the first-generation organic nitrates. Removal by hydrolysis is found to be responsible for the loss of ~30% of the total organic nitrate pool.

25 1 Introduction

In remote continental regions, biogenic volatile organic compounds (BVOCs), including isoprene and terpenes, are the most reactive organic compounds in the atmosphere (Guenther, 2013). The oxidative chemistry of BVOCs affects the distribution of oxidants (OH, O₃, NO₃) and the lifetime of NO_x (=NO+NO₂), creating a feedback loop that affects oxidant concentrations, the lifetime of BVOCs and secondary organic aerosol formation. Along the pathway to complete oxidation of BVOCs, reactions with the nitrogen oxide family radicals (NO, NO₂ and NO₃) to form organic nitrate products (e.g. Perring et al., 2013) are an important branch point that sets the importance of this feedback.



During the day, BVOCs react with the hydroxyl radical (HO) and peroxy radicals (RO_2) are formed. At modest concentrations of NO_x , the peroxy radicals react primarily with NO. The major products of that reaction are NO_2 and an alkoxy radical (RO). There is also a minor channel (with a branching fraction α) that results in addition of the NO to the peroxy radical resulting in an organic nitrate (RONO_2) product. During the night, nitrate radicals (NO_3), the product of the oxidation of NO_2 by O_3 , are also a major source of RONO_2 . BVOCs react with NO_3 , resulting in the formation of nitrooxy peroxy radicals in high yields. The radicals subsequently react to form closed shell RONO_2 , with branching ratio β .

In the last decade, there have been major updates to our understanding of the chemical reactions that occur during isoprene oxidation (Paulot et al., 2009a, 2009b; Crounse et al., 2011; Liu et al., 2013; Peeters et al., 2014; Nguyen et al., 2014; Wolfe et al., 2016; Mills et al., 2016; Teng et al., 2017). This understanding includes recognition that the yield of RONO_2 from reaction of isoprene peroxy radicals with NO is 11–15%, at the high end of the range reported in earlier laboratory experiments (Wennberg et al., 2018). The yield of nitrates from monoterpene oxidation is less clear as laboratory data indicate a very wide range (e.g. from greater than 1% (Aschmann et al., 2002) to 26% (Rindelaub et al., 2015)). For NO_3 oxidation of isoprene experimental data show that the yield, β , is high and varies in the range of 65%–80% (Perring et al., 2009a; Rollins et al., 2009; Kwan et al., 2012).

Once formed, RONO_2 can be photolyzed or oxidized to produce NO_x or HNO_3 along with an organic partner, or they can serve as reservoirs of NO_x that can be transported or deposited to the surface. An additional pathway for gas-phase RONO_2 loss is partitioning into aerosol in either an organic phase where vapor pressure would describe partitioning or an aqueous phase where a Henry's law constant would describe solubility. In the aerosol, the RONO_2 can undergo liquid phase reaction. Some RONO_2 are rapidly hydrolyzed with time scales on the order of hours to minutes under environmentally-relevant pH conditions (Jacobs et al., 2014; Boyd et al., 2015; Rindelaub et al., 2016), while other nitrates are thought to be relatively stable against hydrolysis in neutral conditions (Hu et al., 2011). The main nitrogen-containing product of organic nitrate hydrolysis is nitric acid (Darer et al., 2011). Using measurements of organic nitrates and nitric acid over the Canadian boreal forest and southeast US, Browne et al. (2013) and Romer et al. (2016) provide evidence that hydrolysis of monoterpene and isoprene nitrates is likely a significant loss process and contributes to HNO_3 production. The short lifetime of HNO_3 to deposition in the boundary layer means that organic nitrate loss through hydrolysis in the boundary layer is a permanent sink of NO_x .

For any organic nitrate, its structure determines the rate of its oxidation and photolysis as well as the rate of hydrolysis and deposition. Multifunctional nitrates containing hydroxyl or peroxide groups are likely to have deposition rates much faster than the rates for monofunctional nitrates (Shepson et al., 1996). The dry deposition of organic nitrates has been discussed in the studies by Farmer and Cohen (2008) and Nguyen et al. (2015). Nguyen et al. (2015) directly measured deposition rates of organic nitrates from BVOCs and the first-generation isoprene nitrates were observed to have daytime dry deposition velocity of $\sim 2 \text{ cm s}^{-1}$, which is higher than the values currently used in most models (Ito et al., 2009; Mao et al., 2013; Browne et al., 2014).



Unlike hydrolysis of organic nitrates in aerosol and deposition of organic nitrates to the surface, which is considered a sink of nitrogen oxides in the atmosphere, oxidation and photolysis of RONO_2 may recycle NO_x . Different assumptions regarding NO_x recycling during organic nitrate oxidation result in large variations in simulation of NO_x and O_3 (von Kuhlmann et al., 2004; Fiore et al., 2005; Wu et al., 2007; Horowitz et al., 2007; Paulot et al., 2012). For example, Xie et al. (2013) showed that the uncertainty in the fraction of NO_x returned to the atmosphere during isoprene nitrate oxidation had a larger impact than uncertainty in isoprene nitrate yield on O_3 production. This affirms the need for characterization of the fate and lifetime of RONO_2 in the atmosphere. New clarity is available for the chemical fate of the first-generation isoprene nitrates (e.g. (Lee et al., 2014; Xiong et al., 2015, 2016), while much less is known about the fate of organic nitrates formed from monoterpenes. Because few of these loss processes have been measured, especially for highly oxidized or monoterpene nitrates, there is large uncertainty associated with any description of the lifetime of organic nitrates. Several modeling studies (Paulot et al., 2012; Xie et al., 2013; Mao et al., 2013) have focused specifically on the fate of isoprene nitrates, and have found that how their chemistry is represented has major consequences for NO_x and O_3 . Recently, Browne et al., (2014) extended the representation of organic nitrate chemistry by including in detail the gas-phase chemistry of monoterpenes and discussed different scenarios for uncertain loss processes of monoterpene nitrates. Their improved mechanism for BVOC chemistry has been used as a skeleton for several subsequent modeling studies (e.g. Fisher et al., 2016 and this work). However, none of these models has yet combined detailed molecular representations of individual RONO_2 derived from anthropogenic, isoprene, and monoterpene VOC precursors. Here we describe the development of a gas phase mechanism along those lines. In a forthcoming paper we couple the mechanism described here to aerosol and cloud properties. Here we approximate the effects of aerosols and clouds with simpler parameters representing the effects of the condensed phase chemistry. The model calculations are compared to observations from the SOAS (the Southern Oxidant and Aerosol Study) campaign in the rural Southeastern United States in summer 2013. We explore the relative contributions of OH and NO_3 chemistry to the production of organic nitrates from BVOCs and investigate the importance of different organic nitrate loss processes. Then we explore the lifetime of organic nitrates and consequences of organic nitrate chemistry for atmospheric NO_x to understand the role of RONO_2 in the NO_x and ozone budgets in the moderate NO_x , BVOC-dominated terrestrial environments that represent the most common chemical regime on the continents during summer.

2 Model Description; WRF-Chem model

We use WRF-Chem version 3.5.1 (Grell et al., 2005) with a horizontal resolution of 12 km and 30 vertical layers over the eastern United States. Our simulation domain is defined on the Lambert projection, which is centered at 35°N, 87°W and has 290 and 200 grid points in the west–east and south–north directions, respectively (see Fig. 3 for the horizontal domain). Meteorological data for initial and boundary conditions are driven by the North American Regional Reanalysis (NARR) and data for the chemical initial and boundary conditions are taken from MOZART (Emmons et al., 2010). The model simulation



period is from 27 May to 30 June 2013, with the first 5 days as spin-up, similar to Browne et al. (2014), to remove the impact of initial condition.

Anthropogenic emissions are based on the US EPA 2011 National Emission Inventory (NEI) and scaled to 2013 based on the changes in the annual average emissions from 2011 to 2013. The appropriate scale factors have been derived from the NEI Air Pollutant Emissions Trend Data. We also adjust NO_x emissions (uniformly reduced by 50%) following Travis et al. (2016), who suggest that reduced NO_x emissions can better reproduce the SEAC4RS aircraft measurements for the Southeastern United States. Lightning emissions of NO_x are not included in the model. Lightning NO_x is mainly released at the top of convective updrafts (Ott et al., 2010) and does not strongly impact the distribution of NO_2 in the boundary layer (e.g. Laughner and Cohen, 2017). Biogenic emissions of isoprene, monoterpenes, other BVOCs, oxygenated VOCs (OVOCs), and nitrogen gas emissions from the soil are parameterized using the Model of Emissions of Gases and Aerosol from Nature (MEGAN) (Guenther et al., 2006).

Gas phase reactions are simulated using the second generation Regional Atmospheric Chemistry Mechanism (RACM2) (Goliff et al., 2013), as updated by Browne et al. (2014) and with additions to the mechanisms as described below, which is implemented with the kinetic preprocessor (KPP) software (Damian et al., 2002). The Modal Aerosol Dynamics Model for Europe/Secondary Organic Aerosol Model (MADE/SORGAM) aerosol module (Ackermann et al., 1998; Schell et al., 2001) is used to treat organic and inorganic aerosols. This two-product aerosol scheme does not treat the partitioning of individual chemical species such as organic nitrates. Therefore, we focus here only on investigating the impacts of the gas-phase representation of the chemistry and a full consideration of the gas and aerosol in a coupled framework is a subject of continuing research.

2.1 Chemical Mechanism

We base our core chemistry on a modified version of RACM2 (Goliff et al., 2013). The base we begin with is described by Browne and Cohen (2012) and Browne et al. (2014), and is referred to here as the RACM2_Berkeley scheme. Full details of the RACM2_Berkeley mechanism and a complete list of the compounds can be found in Browne et al. (2014). RACM2_Berkeley includes updates to the isoprene oxidation mechanism (Paulot et al., 2009a, 2009b; Peeters and Müller, 2010; Lockwood et al., 2010; Stavrou et al., 2010; Crouse et al., 2011), an extended mechanism for anthropogenic-originated organic nitrates (Carter and Atkinson, 1989; Middleton et al., 1990; Arey et al., 2001) and updates for monoterpene chemistry. Browne et al. (2014) evaluated the RACM2_Berkeley mechanism using aircraft observations over the Canadian boreal forest.

In this study, the RACM2_Berkeley scheme is further updated with recent advances in the representation of OH- and NO_3 -initiated BVOC oxidation under both low- and high- NO_x conditions, as well as with improved deposition rates and is denoted RACM2_Berkeley2 (see Table S1-S3 in the Supplement). We begin with a more complete description of recent advances in our understanding of isoprene chemistry (Fig. 1). The hydroxy peroxy radical (ISOPO2) that is the product of



isoprene oxidation by OH has multiple potential fates. ISOPO₂ can undergo unimolecular isomerization, leading to the production of hydroperoxy aldehydes (HPALD), among other products. It can react with HO₂ to produce isoprene hydroxy hydroperoxide (ISHP), methyl vinyl ketone (MVK), methacrolein (MACR) and CH₂O. The latter three species can also be formed from the reactions of ISOPO₂ with other (acetyl or methyl) peroxy radicals. ISHP react with OH to form isoprene epoxydiols (IEPOX) and regenerate OH. St. Clair et al. (2015) found that the reaction rate of ISHP + OH is approximately 10% faster than the rate given by Paulot et al. (2009b) and indicate the relative role of the different isomers of ISHP. Here we use kinetics and products of the reactions of three different isomers of ISHP with OH based on St. Clair et al. (2015). We also increase the molar yield of total ISHP from the ISOPO₂ + HO₂ reaction to 93.7% (Liu et al., 2013), with a decrease in the yields of MVK, MACR and HO_x to maintain mass balance. We use rates from Bates et al. (2016) for reactions of three different isomers of IEPOX with OH.

We maintain the overall branching ratio of isoprene nitrates at 11.7% as in Browne et al. (2014) while changing the mix of isomers. Browne et al. (2014) implemented a scheme for the reaction of ISOPO₂ with NO, based on experiments conducted by Paulot et al. (2009b), including β and δ-hydroxy isoprene nitrates (ISOPNB and ISOPND) with yields 4.7% and 7.0%, respectively. Here we update the yield of β versus δ isomers to 10.5% and 1.2% respectively. A theoretical study by Peeters et al. (2014) showed that the peroxy radical redissociations are fast and peroxy isomers may interconvert, so that β-isomers comprise ~95% of the radical pool. The experimental findings of Teng et al. (2017) are also consistent with that idea. Simulations by Fisher et al. (2016) showed that an isoprene hydroxy radical distribution leading to 90% ISOPNB and 10% ISOPND is consistent with SOAS observations.

ISOPND and ISOPNB then photolyze or react with O₃ or OH to yield products that are either NO_x or second-generation organic nitrates. We update reaction rates of isoprene hydroxy nitrate oxidation based on Lee et al. (2014). Compared to the RACM2_Berkeley mechanism (based on Paulot et al. (2009b) and Lockwood et al. (2010)), the reaction rates for ISOPNs+OH are increased and the rate coefficients of ISOPNs+O₃ are decreased. The model represents the products of the reactions of ISOPND and ISOPNB with OH as ISOPNDO₂ and ISOPNBO₂. We update the reaction of ISOPNB+OH to include a small yield of IEPOX and NO₂ (12%) as found by Jacobs et al. (2014). We also update the rate constants for reaction of ISOPNDO₂ and ISOPNBO₂ with NO producing second-generation isoprene nitrates following Lee et al. (2014). Second-generation isoprene nitrates from the OH-initiated pathway include ethanal nitrate (ETHLN), propanone nitrate (PROPNN), multifunctional isoprene nitrate (IMONIT), methacrolein nitrate (MACRN) and methylvinylketone nitrate (MVKN). MACRN and MVKN can also be formed directly from photooxidation of MVK and MACR under high-NO_x conditions. Here, we follow Praske et al. (2015) to update MVK chemistry under both low- and high-NO_x conditions, resulting in greater recycling of OH and NO₂ and decreased formation of organic nitrates from reactions with NO or HO₂.

In addition to OH chemistry, isoprene is oxidized by NO₃. In RACM2_Berkeley, the isoprene + NO₃ chemistry was parameterized with one generic organic nitrate as the product. Recently, Schwantes et al. (2015) developed a kinetic



mechanism for NO₃-initiated oxidation of isoprene in which products, branching ratios, and rate constants are estimated based on recent experimental results. Their suggested products from NO₃ oxidation of isoprene are consistent with organic nitrates detected in the ambient atmosphere during SOAS. RACM2_Berkeley2 treats the NO₃ initiated oxidation of isoprene in some detail with formation and subsequent oxidation of isoprene nitrates largely based on the work of Schwantes et al. (2015) and Rollins et al. (2009) (Fig. 2). In the first step, NO₃ addition to isoprene forms a nitrooxy peroxy radical (INO₂) and then, depending on the radical the INO₂ reacts with, first-generation isoprene nitrates are formed, namely C5 carbonyl nitrate (ICN), β and δ-hydroxy nitrate (IHNB and IHND) and β and δ-nitrooxyhydroperoxide (INPB and INPD). We set the ICN yield at 54% and 72% for the INO₂+NO₃ and INO₂+INO₂ reactions, respectively. ICN is a first-generation isoprene nitrate that is reactive towards NO₃ (Rollins et al., 2009). Subsequent oxidation of ICN by NO₃ forms second-generation isoprene nitrates as well as nitric acid at rates and yields based on Master Chemical Mechanism (MCM v3.2) (Jenkin et al., 1997; Saunders et al., 2003). MCM v3.2 uses a reaction rate coefficient of 1.22 x 10⁻¹⁴ cm³ molec⁻¹ s⁻¹ at 298 K, which is 5 times slower than the rate given by Rollins et al. (2009) and an order of magnitude faster than the rate given by Schwantes et al. (2015). Given the differences in the experimental data, splitting the difference by using the MCM rate seems a reasonable choice.

IHNB and IHND are also identified in the chamber experiments as products of the INO₂+INO₂ reaction. We follow Schwantes et al. (2015) and use rate constants and products, respectively from Lee et al. (2014) and Jacob et al. (2014), for the subsequent fate of IHNB and IHND upon reaction with OH. We note that these rate constants and products correspond to the rate constants and products for the reactions of ISOPND and ISOPNB (OH-initiated isoprene hydroxy nitrates) with OH. In Fig. 2, we only show reactions for one of the isomers.

Nitrooxy hydroperoxides (INPD and INPB) are the dominant products of the INO₂+HO₂ reaction with a combined 77% yield, most of which is δ-isomers (INPD). Schwantes et al. (2015) found that the total molar yield of INPD and INPB per reacted isoprene is higher than the yield found in previous studies (Ng et al., 2008; Kwan et al., 2012). In these previous studies, the nitrooxycarbonyl (ICN) was the main contributor to the family of isoprene nitrates produced by NO₃-initiated chemistry and nitrooxyhydroperoxides (INPD and INPB) were a minor fraction. The difference is likely caused by variation in the fate of the nitrooxy peroxy radical (INO₂+HO₂ vs. INO₂+NO₃) under different experimental conditions. The fate of this radical (INO₂) in the nighttime atmosphere is still highly uncertain (Brown and Stutz, 2012; Boyd et al., 2015). Chamber experiments by Schwantes et al. (2015) suggested that OH reacts with INPD and INPB to form second-generation isoprene nitrates and nitrooxy hydroxy epoxides (INHE), a newly identified product that undergoes a similar heterogeneous chemistry as IEPOX and can impact secondary organic aerosol (SOA) formation. We set the yield of the INHE δ-isomer from INPD at 37% and yield of the INHE β-isomer from INPB at 78%. Those are lower than the IEPOX yield formed from ISHP. INHE isomers will be further oxidized by OH, leading to recycling of NO_x or forming a later generation of organic nitrates. We use the same rates that we use for OH oxidation of IEPOX isomers (Schwantes et al., 2015; Bates et al., 2016). In this scheme, we also assume that INPD and INPB undergo hydrogen abstraction from the hydroperoxide group with the



same rate constants as those we use for hydrogen abstraction from the hydroperoxide group of ISHP as suggested by St. Clair et al. (2015).

C4 carbonyl hydroperoxy nitrates (R4NO) and C4 carbonyl hydroxynitrates (R4N) form from oxidation of the first-generation isoprene nitrates. In reaction with OH, they produce propanone nitrate, PROPNN. The subsequent fate of PROPNN is dominated by oxidation by OH and photolysis resulting in the return of NO_x to the atmosphere. Here, we include a photolysis rate for PROPNN and other carbonyl nitrates (ICN, ETHLN), MVKN and MACRN, that is faster than that used in some other recent mechanisms following the recommendations in Müller et al. (2014). These rates are 10 times larger than the rates given by Paulot et al. (2009a). We use the enhanced photolysis rate of PROPNN for carbonyl nitrate ICN formed from NO₃-initiated isoprene oxidation, and also include photolysis reactions for other new isoprene nitrate species, following Schwantes et al. (2015) and MCM v3.2. The fast photolysis reactions of organic nitrates implemented in this work leads to an increase in NO_x concentrations.

2.2 Organic nitrate deposition to surfaces

We use the resistance-based approach based on the original formulation of Wesely (1989) to calculate dry deposition velocities. Dry deposition rates of RONO₂ in RACM2_Berkeley follow Ito et al. (2007). In RACM2_Berkeley2, using the same formalism, we update the dry deposition parameters (effective Henry's Law coefficients (H*) and reactivity factors (f₀)) for isoprene nitrates as recommended by Nguyen et al. (2015), which results in more rapid removal than in RACM2_Berkeley (Table S3 in the Supplement). We also update the dry deposition parameters for monoterpene-derived nitrates that were previously assumed to deposit at a rate similar to the deposition rate of isoprene hydroxy nitrates (Browne et al., 2014; Ito et al., 2007). Dry deposition velocities of the nitrooxy hydroxy epoxides (INHE) are assumed to be similar to the updated depositional loss rate of IEPOX, given by Nguyen et al. (2015).

2.3 Hydrolysis of tertiary nitrates

In addition to oxidation, photolysis and deposition to the surface, another possible fate of organic nitrates is uptake to the aerosol phase followed by hydrolysis. A rapid hydrolysis (Hu et al., 2011; Jacobs et al., 2014) is recognized for tertiary nitrates, while non-tertiary nitrates under atmospheric conditions are considered unreactive (Darer et al., 2011; Boyd et al., 2015). Due to the limitations of the model representation of organic nitrate aerosol from either aqueous (Marais et al., 2016) or vapor-pressure dependent pathways (Pye et al., 2015), we represent this process for gas-phase organic nitrates by applying a time scale of 3h for tertiary nitrates based on the laboratory chamber study by Boyd et al. (2015). The fraction of tertiary (F_{tertiary}) vs. non-tertiary nitrates is estimated, depending upon the molecular structure of the nitrate, from MCM v3.2.

We apply F_{tertiary} at 41% for β isomers of isoprene nitrates from OH oxidation while we assume that all δ isomers are non-tertiary. Most of the nitrates formed by NO₃-initiated chemistry of isoprene are not tertiary nitrates. The fraction of tertiary nitrates for monoterpene-derived nitrates is also different for species formed by OH oxidation than from NO₃ oxidation. In



RACM2_Berkeley2, we introduce TONIH (C10 nitrooxy hydroperoxide), TONIN (saturated) and UTONIN (unsaturated) monoterpene-derived nitrates from NO₃ oxidation, which differ from the unsaturated (UTONIT) and saturated (TONIT) monoterpene-derived nitrates from OH oxidation. Since contributions of tertiary limonene, α -pinene and β -pinene nitrates from NO₃ reaction are 35%, 15% and 50% (MCM v3.2), respectively, we define F_{tertiary} at 35% as an average for TONIH, UTONIN and TONIN. F_{tertiary} is defined at 77% for UTONIT and TONIT. The value is average of the 62% for α -pinene nitrates and 92% for β -pinene nitrates and is equal to the 77% for limonene nitrates from OH chemistry.

Further changes in the RACM2_Berkeley2 mechanism for monoterpene nitrate chemistry consist of a revised reaction rate for API+NO₃. The rate constant is calculated as an average of the rates given in MCM v3.2 for α -pinene and β -pinene, as API in the mechanism indicates a 50-50 mixture of α -pinene and β -pinene. In our mechanism, following Browne et al. (2014), first-generation monoterpene nitrates react with O₃ and OH and form second-generation nitrates. Here, we also add reaction of first-generation monoterpene nitrates with NO₃ with the rate constant $K=3.15 \times 10^{-13} \exp(-448.0/\text{Temp})$, following Fisher et al. (2016). We assume the second-generation monoterpene nitrate can oxidize, photolyze and deposit identically to nitric acid (Browne et al., 2014). In summary, we have described a detailed chemical mechanism tracking individual organic nitrates in some detail through second-generation products of isoprene and monoterpene oxidation.

3 Results and Discussion

We evaluate our mechanism by comparison to SOAS observations in Bibb County, Alabama (32.90° N latitude, 87.25° W longitude) in summer 2013. These observations together with field campaign data from the long-term monitoring site in the SouthEastern Aerosol Research and CHaracterization (SEARCH) Network (Hansen et al., 2003) (at the same location) provide unique resources for evaluation of our model of organic nitrate chemistry. The measurements include total and speciated organic nitrates, gas phase and aerosol organic nitrates, HO_x radicals, a wide range of VOCs, and ozone (Pye et al., 2015; Romer et al., 2016; Lee et al., 2016).

3.1 Organic nitrate concentrations

Figure 3a shows the spatial distribution of total organic nitrates for the 24-h average of the model simulation period at the surface. The location of SOAS ground site (at the Centreville) is circled in the figure. The campaign area is in a location with among the highest modeled organic nitrate concentrations in the region, up to 350 ppt. Figure 3b highlights that the modeled RONO₂ originating from biogenic VOCs dominate over the organic nitrates with anthropogenic VOC precursors over most of the domain. In the southeast, up to 80% of organic nitrates are biogenic. Biogenic nitrates are 40-50% in the northern portion.

Figure 4 compares median diurnal cycle of observed total organic nitrates from the SOAS campaign to the model simulation during the simulation period. Total organic nitrates in both the gas and particle phase were measured by TD-LIF (Thermal Dissociation Laser-Induced Fluorescence, (Day et al., 2002)). Temporal variability in the total organic nitrates is reproduced



with little bias ($r=0.8$ and normalized mean bias (NMB) =32%). Although the mean of the simulated organic nitrates (0.26 ± 0.19) slightly overestimates the observation mean (0.20 ± 0.1), the medians are found within variability of the observations. The highest bias in median values and variability are observed after sunset to sunrise, which is likely caused by mismatch in vertical turbulent mixing in the simulated and actual boundary layers. Inclusion of hydrolysis as a possible fate for tertiary organic nitrates results in significant improvement of the simulations compared to the observations (not shown here). Tertiary nitrates have shorter lifetime against hydrolysis under atmospheric conditions, compared to the lifetime against deposition (Fig. S1 in the Supplement) making them the most important sink of nitrates.

Diurnal cycles of measured and simulated RONO_2 have a sharp peak at 320-370 ppt around 10:00 and a slow decline through the rest of the day. Throughout the night, the mixing ratios were observed and modeled to remain nearly constant. Around 10:00, when the highest total organic nitrates are observed during SOAS, the simulated OH-initiated and second-generation organic nitrate concentrations both reach their maximum (Fig. S2 in the Supplement). Second-generation nitrates do not show sharp variability over day and night because of their longer lifetime but they do slightly increase after sunrise (around 7:00). OH-initiated organic nitrates that can remain in the residual layer overnight contribute to the total organic nitrate during the morning. At sunrise, when OH and NO began to increase (Fig. S3 in the Supplement), OH-initiated organic nitrates increase until they reach their maximum at around 10:00. In contrast, NO_3 -initiated organic nitrates reach their peak mixing ratio before sunrise and immediately after sunrise, they decline sharply to a minimum concentration during the day (Fig. S2 in the Supplement). As the sun sets, NO drops to near zero and NO_3 production initiates the formation of organic nitrates. OH and NO_3 -initiated reactions occur out of phase in the diurnal cycle resulting in the relatively flat diurnal profile for total organic nitrate throughout the night. Observations of individual molecules are predicted to have more strongly varying diurnal cycles, consistent with observations (Xiong et al., 2015).

3.2 Organic nitrate composition

The composition of the simulated organic nitrates by our model during SOAS at Centreville site (CTR) is shown in Fig. 5a. Monoterpene nitrates are calculated to be one third of total organic nitrates, which is comparable to the Browne et al. (2014) calculation for boreal regions of North America. We define total isoprene-derived nitrates in WRF-Chem as the sum of isoprene hydroxy nitrates, isoprene carbonyl nitrates, MVK and MACR nitrates, isoprene nitrooxyhydroperoxides, ethanal nitrate, propanone nitrate and multifunctional isoprene nitrates. We find the contribution of the total isoprene-derived nitrates to total organic nitrates to be 44%. This is consistent with the range of 25-50% observed from SEAC4RS airborne measurements taken onboard the NASA DC-8 in August–September 2013 over the southeastern US (Fisher et al., 2016). However, it is in contrast to other recent modeling studies over the Southeastern US by Mao et al. (2013) and Xie et al. (2013) that suggested, respectively, more than 90% and 60% of total organic nitrates are from isoprene oxidation. This discrepancy is likely due to the simulated longer lifetime of these nitrates as well as omission of organic nitrates produced from monoterpenes and anthropogenic VOCs in those models.



The observed RONO_2 composition during SOAS is shown in Fig. 5b. The sum of the individual isomers of isoprene nitrates, terpene hydroxynitrates and terpene nitrooxyhydroperoxides were measured in the gas phase by Chemical Ionization Time of Flight Mass Spectrometry using CF_3O^- reagent ion (Crouse et al., 2006; Schwantes et al., 2015; Teng et al., 2015) and ethyl and isopropyl nitrates were measured by gas chromatography–mass spectrometry (de Gouw et al., 2003). Similar to the model results, the largest contributions to the total organic nitrates in the observations are isoprene oxidation products, which represent 22% against 44% in the model. Carbonyl isoprene nitrates including ICN, ETHLN and PROPNN as a fraction of total RONO_2 (8% in the model and <7% in the observations) and their concentrations (Fig. S4a and b in the Supplement) are reproduced well by the model. However, the model overestimates the fraction of RONO_2 that is isoprene hydroxynitrates and MVKN+MACRN (21% modeled vs. 13% observed). Isoprene hydroxynitrates from NO_3 -initiated chemistry are a small portion of the total simulated isoprene hydroxynitrates (~15%). The difference between the modeled and observed contribution of isoprene hydroxynitrates to total organic nitrates is thus more likely a result of differences between the modeled and observed nitrates that are products of OH-initiated chemistry. Insights from very recent studies by Teng et al. (2017) and Wennberg et al. (2018) suggests a larger F_{tertiary} for OH-initiated isoprene hydroxynitrates than the value we calculated from MCM. That reflects a larger fraction of these nitrates to be subject of hydrolysis and thus it perhaps explains part of the discrepancy between the model simulations and observations.

The largest difference between the modeled and observed contribution of isoprene nitrates to total organic nitrates is due to the modeled gas-phase multifunctional isoprene nitrates and isoprene nitrooxy hydroperoxides. Aerosol- and gas-phase second-generation multifunctional isoprene nitrates and aerosol-phase isoprene nitrooxy hydroperoxides were not individually measured during SOAS. Instead, total aerosol-phase organic nitrates were measured by TD-LIF, using an activated charcoal denuder to remove gas-phase organic nitrates, and found to contribute around 40% of total organic nitrates at the SOAS CTR site (Fig. 5b). Ng et al. (2008) and Rollins et al. (2009) found isoprene oxidation can form 4–23% nitrate aerosol yields and showed multifunctional nitrates to be a dominant nitrate aerosol. If the isoprene nitrooxy hydroperoxides are favored to partition to aerosol this would explain the model-measurement discrepancy for the calculated contribution of multifunctional isoprene nitrates and isoprene nitrooxy hydroperoxide. They are simulated in the gas phase using RACM2_Berkeley2 but we might interpret them as contributing to particle phase organic nitrate. That is consistent with the Lee et al. (2016) finding from observations of speciated particle organic nitrates during the SOAS campaign. They showed total particle organic nitrates have a dominant contribution from highly functionalized isoprene nitrates containing between six and eight oxygen atoms.

Nitrate aerosol yields for monoterpene oxidation reactions from different laboratory chamber experiments, field measurements and modeling studies have been reported to be very high (up to 100%) (Russell and Allen, 2005; Fry et al., 2009, 2011, 2013; Pye et al., 2015; Boyd et al., 2015). Among monoterpene nitrates, NO_3 -initiated nitrates (Ayres et al., 2015) and functionalized nitrates (Lee et al., 2016) have been shown to be an especially significant fraction of the total particle organic nitrate source at SOAS site. These findings imply that the remainder of the measured particle organic



nitrates can be attributed to mono- or sesquiterpene derived RONO_2 including NO_3 -initiated terpene hydroxynitrates, terpene nitrooxyhydroperoxides and multifunctional terpene nitrates, which are simulated and present in the gas phase in our mechanism. If we interpret the aerosol nitrates to be these compounds, then we find a rough correspondence between the model and observations (see Fig. 5a and b).

- 5 In RACM2_Berkeley2, contribution from organic nitrates of anthropogenic origin is simulated to be 21% of total RONO_2 (referred as Other at Fig. 5a). That is higher than the 10% inferred from the observations for the measured anthropogenic organic nitrates at SOAS. Some caution should be taken in the interpretation of such a comparison, as the observations at SOAS do not represent the same species as the modeled ones and include only ethyl and propyl nitrates (Fig. 5b). In RACM2_Berkeley2, a wide range of the organic nitrates of anthropogenic origin (C_1 - C_5 nitrates) are categorized into four
- 10 groups including monofunctional saturated, multifunctional unsaturated, multifunctional saturated, and aromatic-derived nitrates that are partitioned from the lumped precursors including alkanes, aromatics, alcohols and alkenes. However, the remaining unspciated measured RONO_2 contribute 27% of total organic nitrates observed by TD-LIF. One hypothesis to explain this difference is that the rest of simulated organic nitrates of anthropogenic origin might be a portion of the unspecified measured RONO_2 .

15 3.3 Relationships between RONO_2 , O_x and CH_2O

As ozone and total organic nitrates are produced in a common reaction with branches that yield one or the other, their observed and modeled correlation provides an additional constraint on our understanding of organic nitrates. The sum of O_3 and NO_2 is conserved on longer time scales than O_3 alone, accordingly we use $\text{O}_x = \text{O}_3 + \text{NO}_2$ in this analysis. As shown in Fig. 6, during the daytime (from 8:00 to 18:00 local time) the modeled and observed correlations between O_x and RONO_2 are

20 nearly identical. A linear fit to the observations yields a line with slope of $129 \pm 4 \text{ ppbv}(\text{O}_x) \text{ ppbv}(\text{RONO}_2)^{-1}$ and a fit to the model output yields $125 \pm 4 \text{ ppbv}(\text{O}_x) \text{ ppbv}(\text{RONO}_2)^{-1}$.

This slope has typically been used to estimate the approximate branching ratio of the entire VOC mixture (α_{eff}) with an assumption that photochemical production is rapid as compared to removal processes; α_{eff} is inversely proportional to the O_x vs. RONO_2 slope (Perring et al., 2013). The quantified α in the laboratory for BVOCs are much higher than typical α 's for

25 anthropogenic VOCs (Perring et al. (2013) and references therein). Therefore, for regions like Southeastern U.S. where BVOCs dominate the VOC mixture, a much lower slope than our calculated value is expected. We conclude that the observed slope is reflecting the short lifetime of organic nitrates at SOAS.

Formaldehyde (CH_2O) is another co-product to RONO_2 and as Perring et al. (2009b) discussed, the slope of the $\text{RONO}_2/\text{CH}_2\text{O}$ correlation is related to the ratio of the production of both species, as both have similar lifetimes. We would

30 expect the slope could provide a constraint on the yield of isoprene nitrates, especially since in much of the domain isoprene is the dominant source of both RONO_2 and CH_2O . Figure 7 shows the correlation between observed CH_2O and RONO_2 during SOAS. The slope of the best fit line, with an intercept allowed to differ from zero to consider the possibility of a



background CH₂O that mixes in from the free troposphere, is found to be 0.116, consistent with previous estimates by Perring et al. (2009b) who observed a slope of 0.119 during INTEX-NA in 2004. The slope would imply an OH-initiated isoprene nitrate yield of 12% (Perring et al., 2009b) if we use a lifetime of 1.7 h at SOAS for RONO₂ as reported by Romer et al. (2016). This is nearly identical to the yield used in the mechanism described in this manuscript. The correlation of modeled CH₂O and total RONO₂ has a smaller slope of 0.085. The discrepancy between the slopes from the simulated and observed data can be attributed to model overestimation of CH₂O (Fig. S5 in the Supplement).

3.4 Organic nitrate formation

Figure 8a shows the diurnal cycle of fractional RONO₂ production simulated using RACM2_Berkeley2 averaged over the boundary layer at the CTR site during SOAS. During the day, production of organic nitrates is dominated by reaction of isoprene with OH. This is consistent with the OH reactivity (OHR) of individually measured compounds at SOAS which was dominated by reaction with isoprene (Kaiser et al., 2016; Romer et al., 2016). In contrast, the vast majority of RONO₂ production at night is monoterpene nitrates, which are formed as a result of NO₃ oxidation of nighttime monoterpene emissions. NO₃ chemistry of isoprene leading to isoprene nitrate formation is also found to be significant during the nighttime. This fraction from isoprene that accumulates in the boundary layer in late afternoons can reach 35% of the total organic nitrates formed at night. In addition to investigating the relative importance of instantaneous organic nitrate production based on the VOC precursors, we calculate the fraction of isoprene nitrates produced from NO₃ chemistry to the total isoprene nitrate production (~44%, Fig. S6 in the Supplement) over day and night of the entire modeling period confirming the relative importance of this pathway for producing isoprene nitrates versus OH oxidation of isoprene. This finding is consistent with the modeling result (~40%) from Xie et al. (2013).

The fraction of anthropogenic VOC-derived organic nitrates to total simulated production of organic nitrates is estimated to be negligible; however their contribution to the simulated concentrations of organic nitrates is much higher and reached up to 0.25 (Fig. 8b). This is due to their relatively long lifetime (> 100 h lifetime to oxidation by 1×10^6 molecules cm⁻³ of OH at 298 K and long lifetime to deposition (Henry's law constant of $\sim 1 \text{ Matm}^{-1}$), Browne et al. (2014) and references therein) that causes their concentrations to increase with time in the boundary layer in comparison to short-lived first-generation biogenic organic nitrates. Similarly, the fraction of second-generation nitrates formed from oxidation of first-generation isoprene and monoterpene nitrates (Fig. 8a) is predicted to be (~ 0.04), much less than the calculated contribution of their concentrations to total organic nitrate concentrations (~ 0.3) (Fig. 8b). In the next subsection we will discuss more about the loss processes and lifetime of these organic nitrates.

3.5 Organic nitrate lifetime

We define the lifetime of organic nitrates as the concentration of RONO₂ divided by the combined loss rate via all proposed loss mechanisms. The loss mechanisms include chemical loss processes (oxidation, photolysis, and hydrolysis of RONO₂)



and deposition. The nighttime lifetime of organic nitrates might be longer than the daytime value (and might be similar to or longer than the length of a single night). Because of uncertainties associated with simulation of the boundary layer height and organic nitrate concentrations at nighttime, we focus on the daytime lifetime as a guide for thinking about the organic nitrate fate. Figure 9 shows the estimated daytime lifetime of 2 h for first-generation biogenic organic nitrates and a longer lifetime for the second-generation organic nitrates (3.2 h). Including organic nitrates from anthropogenic sources we estimate a fairly short overall lifetime of 2.6 h for total RONO_2 . This short lifetime results in the less efficient transport of organic nitrates to the free troposphere and over large distances from sources. Using the SOAS field observations, Romer et al. (2016) suggested ~ 1.7 h for the atmospheric lifetime of RONO_2 . They calculated the lifetime by the assumption that RONO_2 are near steady state in the afternoon. If we constrain our calculation to 12:00-16:00 and give an intercept of 40 ppt as Romer et al. (2016) did, the overall estimated lifetime in the model is estimated to be 2.9 h, but using the production rates of organic nitrates instead of the loss rates (by assumption of the atmospheric steady state condition applied in Romer et al. (2016)) our result remarkably shows very good agreement with their finding (identical value, Fig. S7 in the Supplement). GEOS-Chem simulations by Fisher et al. (2016) reported a similar short lifetime by assuming a hydrolysis lifetime of 1 h lifetime for all tertiary and non-tertiary nitrates and not including the longer-lived small alkyl nitrates.

Accurate determination of the lifetime of organic nitrates is a major challenge for assessing the influence of organic nitrates on atmospheric chemistry. However, The estimated lifetime of ~ 3 h for organic nitrates found here as well as other studies over the Southeastern United States (Perring et al., 2009b; Romer et al., 2016; Fisher et al., 2016) is less than the range of NO_x lifetimes (5.5–11 h) calculated by observational studies (e.g. Valin et al., 2013; Romer et al., 2016). Organic nitrates should therefore generally be categorized as short-lived NO_x reservoirs, which remove NO_x in a plume, but act as a source of NO_x in remote regions.

3.6 NO_x recycling efficiency

To determine the fraction of NO_x converted to RONO_2 and then released back to the gas phase as NO_x , the relative importance of different loss pathways of organic nitrates must be known. Oxidation and photolysis of organic nitrates recycle NO_x but hydrolysis and deposition cause permanent removal of NO_x from the atmosphere. Recent field studies suggested that isoprene nitrates are removed quickly by dry deposition (Nguyen et al., 2015) and some have concluded that deposition is the primary sink of nitrates (Rosen et al., 2004; Horii et al., 2006; Horowitz et al., 2007), while others estimated that oxidation or ozonolysis is the dominant loss mechanism of isoprene nitrates (Shepson et al., 1996; Ito et al., 2007; Perring et al., 2009a; Browne et al., 2013). Similar uncertainty for the fate and dominant loss processes of monoterpene nitrates was found by Browne et al. (2014). Here, we update possible fates of organic nitrates in WRF-Chem from recent findings including photolysis (Müller et al., 2014), oxidation and ozonolysis (Lee et al., 2014), deposition (Nguyen et al., 2015) and hydrolysis (Boyd et al., 2015) and then estimated the contribution of different fates to first- and second-generation isoprene and monoterpene nitrates (Fig. 10a and b). We note that our calculation represents the loss of the nitrate



functionality and does not include the fraction of loss processes of first-generation organic nitrates by oxidation and ozonolysis that retain the nitrate functionality by forming second-generation organic nitrates.

Figure 10b shows that the loss of second-generation organic nitrates is dominated by deposition (60%), causing permanent loss of NO_x . That is due to the assumed rapid depositional loss of second-generation monoterpene nitrates (deposited as fast as HNO_3) in this study following Browne et al. (2014). Fractional contributions of photolysis (~25%) and oxidation (~15%) are not negligible and are much larger than those estimated by Browne et al. (2014), which is a consequence of using the rapid photolysis rates of PROPNN and ETHLN as second-generation isoprene nitrates (Müller et al., 2014). In contrast, the loss of first-generation nitrates occurs largely by the sum of chemical mechanisms that recycle NO_x to the atmosphere: reaction with ozone (24%) and OH (11%) and photolysis (20%), with additional loss by deposition (15%) and hydrolysis (~29%). Fisher et al. (2016) predicted much larger losses of RONO_2 by aerosol hydrolysis, (~60% of total nitrate losses), reflecting the short lifetime of nitrates with respect to hydrolysis applied in their study for all nitrates. However, with our representation of hydrolysis of nitrates in aqueous solutions (slower rate and only tertiary nitrates), hydrolysis is still important (the calculated loss rate for nitrates is ~ 3 to 11 pptv h^{-1}), accounting for one-third of the organic nitrate loss and leading to a large increase in HNO_3 production in the atmosphere. Assessing the impact of hydrolysis of nitrates on the budget of nitric acid is beyond the scope of this work. We note that the relative contribution of nitrate hydrolysis in aqueous solutions differs widely for individual RONO_2 species from each other due to their different structures.

Including loss of RONO_2 from anthropogenic sources, we find that loss of overall RONO_2 via hydrolysis with an additional contribution from deposition become comparable to loss via other processes that return NO_x to the atmosphere. Figure 11 shows the NO_x recycling efficiency, defined as the ratio between the instantaneous production of NO_x from loss of organic nitrates and the instantaneous loss of NO_x to production of organic nitrates. Since chemical degradation of nitric acid is much slower than deposition, the slope of 0.52 is interpreted as the fraction of the sequestered NO_x that can be exported away from its emission region and released downwind through organic nitrate chemistry. Our finding is consistent with a recycling of about half of isoprene nitrates to NO_x calculated by Paulot et al. (2009a) and Horowitz et al. (2007). Of this total, we calculated ~38% of NO_x cycled back relatively quickly while first-generation nitrates are oxidizing and producing second-generation nitrates.

4 Conclusions

The lifetimes of organic nitrates with respect to hydrolysis, oxidation, and deposition play an important role in the NO_x budget and formation of O_3 and secondary organic aerosols. Analyses from recent field studies in the Southeastern United States found a lifetime of ~2-3 h for organic nitrates. By incorporating new findings from recent laboratory, field and modeling studies into a gas-phase mechanism we provide a state-of-the-science representation of expanded organic nitrates in the WRF-Chem model. Using the updated model we are able to reproduce a short organic nitrate lifetime (2.6 h), similar to that observed during SOAS.



After adding hydrolysis as a possible fate of tertiary gas-phase biogenic organic nitrates in our mechanism and in combination with all other loss mechanism, we find that the lifetime of second-generation organic nitrates is longer than the lifetime of first-generation nitrates. We find dry deposition is the dominant loss process for second-generation organic nitrates and chemical mechanisms of ozonolysis, photolysis and oxidation that can recycle NO_x to the atmosphere have a more important role in loss of first-generation organic nitrates from the atmosphere. The contribution of tertiary nitrate hydrolysis to total organic nitrate removal from the atmosphere is found to be 30%. We find, therefore, that 52% of the NO_x sequestered by production of organic nitrates can be cycled back to the atmosphere.

To accurately estimate organic nitrate lifetime, the production, loss and fate of these compounds must be well constrained. Evaluation of our updated mechanism using SOAS observations in summer 2013 indicates the model represents much of the important chemistry governing organic nitrates. We show that the simulated concentrations of total organic nitrates, correlations with CH_2O and ozone and contribution of individual RONO_2 to total organic nitrates are in fairly good agreement with observations at the SOAS CTR ground site. We find the largest difference between the modeled and observed contributions of individual organic nitrate compounds to total RONO_2 is for highly functionalized isoprene nitrates and monoterpene nitrates. We attribute this difference to possible high aerosol yields of these organic nitrate species, which are represented in the gas phase in our mechanism. Future analysis for developing a complete representation of organic nitrate chemistry including an organic nitrate aerosol formation mechanism, either from aqueous-phase uptake or vapor-pressure partitioning onto pre-existing organic aerosol, in addition to the detailed gas-phase mechanism described here will benefit the model approximation.

Data availability

Measurements from the SOAS campaign are available at <https://esrl.noaa.gov/csd/groups/csd7/measurements/2013senex/Ground/DataDownload> (SOAS Science Team, 2013).

Acknowledgements

We gratefully acknowledge support from NSF grants AGS-1352972, AGS-1247421 and AGS-1628530, NOAA Office of Global Programs grant NA13OAR4310067 and NASA grant NNX 15AE37G. We thank Allen Goldstein and Pawel Konrad Misztal for PTRTOFMS VOC data and William Brune for OH data. We also thank the SOAS field campaign team including Ann Marie Carlton.



References

- Ackermann, I. J., Hass, H., Memmesheimer, M., Ebel, A., Binkowski, F. S. and Shankar, U.: Modal Aerosol Dynamics model for Europe: Development and first applications, *Atmos. Environ.*, 32(17), 2981–2999, doi:10.1016/S1352-2310(98)00006-5, 1998.
- 5 Arey, J., Aschmann, S. M., Kwok, E. S. C. and Atkinson, R.: Alkyl Nitrate, Hydroxyalkyl Nitrate, and Hydroxycarbonyl Formation from the NO_x-Air Photooxidations of C₅-C₈ n-Alkanes, *J. Phys. Chem. A*, 105(6), 1020–1027, doi:10.1021/jp003292z, 2001.
- Aschmann, S. M., Atkinson, R. and Arey, J.: Products of reaction of OH radicals with α -pinene, *J. Geophys. Res. Atmospheres*, 107(D14), ACH 6-1, doi:10.1029/2001JD001098, 2002.
- 10 Ayres, B. R., Allen, H. M., Draper, D. C., Brown, S. S., Wild, R. J., Jimenez, J. L., Day, D. A., Campuzano-Jost, P., Hu, W., de Gouw, J., Koss, A., Cohen, R. C., Duffey, K. C., Romer, P., Baumann, K., Edgerton, E., Takahama, S., Thornton, J. A., Lee, B. H., Lopez-Hilfiker, F. D., Mohr, C., Wennberg, P. O., Nguyen, T. B., Teng, A., Goldstein, A. H., Olson, K. and Fry, J. L.: Organic nitrate aerosol formation via NO₃ + biogenic volatile organic compounds in the southeastern United States, *Atmos Chem Phys*, 15(23), 13377–13392, doi:10.5194/acp-15-13377-2015, 2015.
- 15 Bates, K. H., Nguyen, T. B., Teng, A. P., Crounse, J. D., Kjaergaard, H. G., Stoltz, B. M., Seinfeld, J. H. and Wennberg, P. O.: Production and Fate of C₄ Dihydroxycarbonyl Compounds from Isoprene Oxidation, *J. Phys. Chem. A*, 120(1), 106–117, doi:10.1021/acs.jpca.5b10335, 2016.
- Boyd, C. M., Sanchez, J., Xu, L., Eugene, A. J., Nah, T., Tuet, W. Y., Guzman, M. I. and Ng, N. L.: Secondary organic aerosol formation from the β -pinene+NO₃ system: effect of humidity and peroxy radical fate, *Atmos Chem Phys*, 15(13), 7497–7522, doi:10.5194/acp-15-7497-2015, 2015.
- 20 Brown, S. S. and Stutz, J.: Nighttime radical observations and chemistry, *Chem. Soc. Rev.*, 41(19), 6405–6447, doi:10.1039/C2CS35181A, 2012.
- Browne, E. C. and Cohen, R. C.: Effects of biogenic nitrate chemistry on the NO_x lifetime in remote continental regions, *Atmos Chem Phys*, 12(24), 11917–11932, doi:10.5194/acp-12-11917-2012, 2012.
- 25 Browne, E. C., Min, K.-E., Wooldridge, P. J., Apel, E., Blake, D. R., Brune, W. H., Cantrell, C. A., Cubison, M. J., Diskin, G. S., Jimenez, J. L., Weinheimer, A. J., Wennberg, P. O., Wisthaler, A. and Cohen, R. C.: Observations of total RONO₂ over the boreal forest: NO_x sinks and HNO₃ sources, *Atmos Chem Phys*, 13(9), 4543–4562, doi:10.5194/acp-13-4543-2013, 2013.
- Browne, E. C., Wooldridge, P. J., Min, K.-E. and Cohen, R. C.: On the role of monoterpene chemistry in the remote
30 continental boundary layer, *Atmos Chem Phys*, 14(3), 1225–1238, doi:10.5194/acp-14-1225-2014, 2014.
- Carter, W. P. L. and Atkinson, R.: Alkyl nitrate formation from the atmospheric photooxidation of alkanes; a revised estimation method, *J. Atmospheric Chem.*, 8(2), 165–173, doi:10.1007/BF00053721, 1989.



- Crouse, J. D., McKinney, K. A., Kwan, A. J. and Wennberg, P. O.: Measurement of Gas-Phase Hydroperoxides by Chemical Ionization Mass Spectrometry, *Anal. Chem.*, 78(19), 6726–6732, doi:10.1021/ac0604235, 2006.
- Crouse, J. D., Paulot, F., Kjaergaard, H. G. and Wennberg, P. O.: Peroxy radical isomerization in the oxidation of isoprene, *Phys. Chem. Chem. Phys.*, 13(30), 13607–13613, doi:10.1039/C1CP21330J, 2011.
- 5 Damian, V., Sandu, A., Damian, M., Potra, F. and Carmichael, G. R.: The kinetic preprocessor KPP—a software environment for solving chemical kinetics, *Comput. Chem. Eng.*, 26(11), 1567–1579, doi:10.1016/S0098-1354(02)00128-X, 2002.
- Darer, A. I., Cole-Filipiak, N. C., O'Connor, A. E. and Elrod, M. J.: Formation and Stability of Atmospherically Relevant Isoprene-Derived Organosulfates and Organonitrates, *Environ. Sci. Technol.*, 45(5), 1895–1902, doi:10.1021/es103797z, 2011.
- 10 Day, D. A., Wooldridge, P. J., Dillon, M. B., Thornton, J. A. and Cohen, R. C.: A thermal dissociation laser-induced fluorescence instrument for in situ detection of NO₂, peroxy nitrates, alkyl nitrates, and HNO₃, *J. Geophys. Res. Atmospheres*, 107(D6), ACH 4-1, doi:10.1029/2001JD000779, 2002.
- Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L. and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci Model Dev*, 3(1), 43–67, doi:10.5194/gmd-3-43-2010, 2010.
- Farmer, D. K. and Cohen, R. C.: Observations of HNO₃, ΣAN, ΣPN and NO₂ fluxes: evidence for rapid HO_x chemistry within a pine forest canopy, *Atmos Chem Phys*, 8(14), 3899–3917, doi:10.5194/acp-8-3899-2008, 2008.
- Fiore, A. M., Horowitz, L. W., Purves, D. W., Levy, H., Evans, M. J., Wang, Y., Li, Q. and Yantosca, R. M.: Evaluating the contribution of changes in isoprene emissions to surface ozone trends over the eastern United States, *J. Geophys. Res. Atmospheres*, 110(D12), D12303, doi:10.1029/2004JD005485, 2005.
- Fisher, J. A., Jacob, D. J., Travis, K. R., Kim, P. S., Marais, E. A., Chan Miller, C., Yu, K., Zhu, L., Yantosca, R. M., Sulprizio, M. P., Mao, J., Wennberg, P. O., Crouse, J. D., Teng, A. P., Nguyen, T. B., St. Clair, J. M., Cohen, R. C., Romer, P., Nault, B. A., Wooldridge, P. J., Jimenez, J. L., Campuzano-Jost, P., Day, D. A., Hu, W., Shepson, P. B., Xiong, F., Blake, D. R., Goldstein, A. H., Misztal, P. K., Hanisco, T. F., Wolfe, G. M., Ryerson, T. B., Wisthaler, A. and Mikoviny, T.: Organic nitrate chemistry and its implications for nitrogen budgets in an isoprene- and monoterpene-rich atmosphere: constraints from aircraft (SEAC4RS) and ground-based (SOAS) observations in the Southeast US, *Atmos Chem Phys*, 16(9), 5969–5991, doi:10.5194/acp-16-5969-2016, 2016.
- Fry, J. L., Kiendler-Scharr, A., Rollins, A. W., Wooldridge, P. J., Brown, S. S., Fuchs, H., Dubé, W., Mensah, A., dal Maso, M., Tillmann, R., Dorn, H.-P., Brauers, T. and Cohen, R. C.: Organic nitrate and secondary organic aerosol yield from NO₃ oxidation of β-pinene evaluated using a gas-phase kinetics/aerosol partitioning model, *Atmos Chem Phys*, 9(4), 1431–1449, doi:10.5194/acp-9-1431-2009, 2009.
- 30



- Fry, J. L., Kiendler-Scharr, A., Rollins, A. W., Brauers, T., Brown, S. S., Dorn, H.-P., Dubé, W. P., Fuchs, H., Mensah, A., Rohrer, F., Tillmann, R., Wahner, A., Wooldridge, P. J. and Cohen, R. C.: SOA from limonene: role of NO₃ in its generation and degradation, *Atmos Chem Phys*, 11(8), 3879–3894, doi:10.5194/acp-11-3879-2011, 2011.
- Fry, J. L., Draper, D. C., Zarzana, K. J., Campuzano-Jost, P., Day, D. A., Jimenez, J. L., Brown, S. S., Cohen, R. C., Kaser, L., Hansel, A., Cappellin, L., Karl, T., Hodzic Roux, A., Turnipseed, A., Cantrell, C., Lefer, B. L. and Grossberg, N.: Observations of gas- and aerosol-phase organic nitrates at BEACHON-RoMBAS 2011, *Atmos Chem Phys*, 13(17), 8585–8605, doi:10.5194/acp-13-8585-2013, 2013.
- Goliff, W. S., Stockwell, W. R. and Lawson, C. V.: The regional atmospheric chemistry mechanism, version 2, *Atmos. Environ.*, 68, 174–185, doi:10.1016/j.atmosenv.2012.11.038, 2013.
- 10 de Gouw, J. A., Goldan, P. D., Warneke, C., Kuster, W. C., Roberts, J. M., Marchewka, M., Bertman, S. B., Pszenny, A. a. P. and Keene, W. C.: Validation of proton transfer reaction-mass spectrometry (PTR-MS) measurements of gas-phase organic compounds in the atmosphere during the New England Air Quality Study (NEAQS) in 2002, *J. Geophys. Res. Atmospheres*, 108(D21), 4682, doi:10.1029/2003JD003863, 2003.
- Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C. and Eder, B.: Fully coupled “online” chemistry within the WRF model, *Atmos. Environ.*, 39(37), 6957–6975, doi:10.1016/j.atmosenv.2005.04.027, 2005.
- 15 Guenther, A.: Biological and Chemical Diversity of Biogenic Volatile Organic Emissions into the Atmosphere, *Int. Sch. Res. Not.*, 2013, e786290, doi:10.1155/2013/786290, 2013.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I. and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos Chem Phys*, 6(11), 3181–3210, doi:10.5194/acp-6-3181-2006, 2006.
- 20 Hansen, D. A., Edgerton, E. S., Hartsell, B. E., Jansen, J. J., Kandasamy, N., Hidy, G. M. and Blanchard, C. L.: The Southeastern Aerosol Research and Characterization Study: Part 1—Overview, *J. Air Waste Manag. Assoc.*, 53(12), 1460–1471, doi:10.1080/10473289.2003.10466318, 2003.
- Horii, C. V., William Munger, J., Wofsy, S. C., Zahniser, M., Nelson, D. and Barry McManus, J.: Atmospheric reactive nitrogen concentration and flux budgets at a Northeastern U.S. forest site, *Agric. For. Meteorol.*, 136(3), 159–174, doi:10.1016/j.agrformet.2006.03.005, 2006.
- 25 Horowitz, L. W., Fiore, A. M., Milly, G. P., Cohen, R. C., Perring, A., Wooldridge, P. J., Hess, P. G., Emmons, L. K. and Lamarque, J.-F.: Observational constraints on the chemistry of isoprene nitrates over the eastern United States, *J. Geophys. Res. Atmospheres*, 112(D12), D12S08, doi:10.1029/2006JD007747, 2007.
- 30 Hu, K. S., Darer, A. I. and Elrod, M. J.: Thermodynamics and kinetics of the hydrolysis of atmospherically relevant organonitrates and organosulfates, *Atmos Chem Phys*, 11(16), 8307–8320, doi:10.5194/acp-11-8307-2011, 2011.
- Ito, A., Sillman, S. and Penner, J. E.: Effects of additional nonmethane volatile organic compounds, organic nitrates, and direct emissions of oxygenated organic species on global tropospheric chemistry, *J. Geophys. Res. Atmospheres*, 112(D6), D06309, doi:10.1029/2005JD006556, 2007.



- Ito, A., Sillman, S. and Penner, J. E.: Global chemical transport model study of ozone response to changes in chemical kinetics and biogenic volatile organic compounds emissions due to increasing temperatures: Sensitivities to isoprene nitrate chemistry and grid resolution, *J. Geophys. Res. Atmospheres*, 114(D9), D09301, doi:10.1029/2008JD011254, 2009.
- Jacobs, M. I., Burke, W. J. and Elrod, M. J.: Kinetics of the reactions of isoprene-derived hydroxynitrates: gas phase epoxide formation and solution phase hydrolysis, *Atmos Chem Phys*, 14(17), 8933–8946, doi:10.5194/acp-14-8933-2014, 2014.
- Jenkin, M. E., Saunders, S. M. and Pilling, M. J.: The tropospheric degradation of volatile organic compounds: a protocol for mechanism development, *Atmos. Environ.*, 31(1), 81–104, doi:10.1016/S1352-2310(96)00105-7, 1997.
- Kaiser, J., Skog, K. M., Baumann, K., Bertman, S. B., Brown, S. B., Brune, W. H., Crounse, J. D., de Gouw, J. A., Edgerton, E. S., Feiner, P. A., Goldstein, A. H., Koss, A., Misztal, P. K., Nguyen, T. B., Olson, K. F., St. Clair, J. M., Teng, A. P., Toma, S., Wennberg, P. O., Wild, R. J., Zhang, L. and Keutsch, F. N.: Speciation of OH reactivity above the canopy of an isoprene-dominated forest, *Atmos Chem Phys*, 16(14), 9349–9359, doi:10.5194/acp-16-9349-2016, 2016.
- von Kuhlmann, R., Lawrence, M. G., Pöschl, U. and Crutzen, P. J.: Sensitivities in global scale modeling of isoprene, *Atmos Chem Phys*, 4(1), 1–17, doi:10.5194/acp-4-1-2004, 2004.
- Kwan, A. J., Chan, A. W. H., Ng, N. L., Kjaergaard, H. G., Seinfeld, J. H. and Wennberg, P. O.: Peroxy radical chemistry and OH radical production during the NO₃-initiated oxidation of isoprene, *Atmos Chem Phys*, 12(16), 7499–7515, doi:10.5194/acp-12-7499-2012, 2012.
- Laughner, J. L. and Cohen, R. C.: Quantification of the effect of modeled lightning NO₂ on UV–visible air mass factors, *Atmos Meas Tech*, 10(11), 4403–4419, doi:10.5194/amt-10-4403-2017, 2017.
- Lee, B. H., Mohr, C., Lopez-Hilfiker, F. D., Lutz, A., Hallquist, M., Lee, L., Romer, P., Cohen, R. C., Iyer, S., Kurtén, T., Hu, W., Day, D. A., Campuzano-Jost, P., Jimenez, J. L., Xu, L., Ng, N. L., Guo, H., Weber, R. J., Wild, R. J., Brown, S. S., Koss, A., Gouw, J. de, Olson, K., Goldstein, A. H., Seco, R., Kim, S., McAvey, K., Shepson, P. B., Starn, T., Baumann, K., Edgerton, E. S., Liu, J., Shilling, J. E., Miller, D. O., Brune, W., Schobesberger, S., D’Ambro, E. L. and Thornton, J. A.: Highly functionalized organic nitrates in the southeast United States: Contribution to secondary organic aerosol and reactive nitrogen budgets, *Proc. Natl. Acad. Sci.*, 113(6), 1516–1521, doi:10.1073/pnas.1508108113, 2016.
- Lee, L., Teng, A. P., Wennberg, P. O., Crounse, J. D. and Cohen, R. C.: On Rates and Mechanisms of OH and O₃ Reactions with Isoprene-Derived Hydroxy Nitrates, *J. Phys. Chem. A*, 118(9), 1622–1637, doi:10.1021/jp4107603, 2014.
- Liu, Y. J., Herdinger-Blatt, I., McKinney, K. A. and Martin, S. T.: Production of methyl vinyl ketone and methacrolein via the hydroperoxyl pathway of isoprene oxidation, *Atmos Chem Phys*, 13(11), 5715–5730, doi:10.5194/acp-13-5715-2013, 2013.
- Lockwood, A. L., Shepson, P. B., Fiddler, M. N. and Alaghmand, M.: Isoprene nitrates: preparation, separation, identification, yields, and atmospheric chemistry, *Atmos Chem Phys*, 10(13), 6169–6178, doi:10.5194/acp-10-6169-2010, 2010.



- Mao, J., Paulot, F., Jacob, D. J., Cohen, R. C., Crounse, J. D., Wennberg, P. O., Keller, C. A., Hudman, R. C., Barkley, M. P. and Horowitz, L. W.: Ozone and organic nitrates over the eastern United States: Sensitivity to isoprene chemistry, *J. Geophys. Res. Atmospheres*, 118(19), 2013JD020231, doi:10.1002/jgrd.50817, 2013.
- Marais, E. A., Jacob, D. J., Jimenez, J. L., Campuzano-Jost, P., Day, D. A., Hu, W., Krechmer, J., Zhu, L., Kim, P. S.,
5 Miller, C. C., Fisher, J. A., Travis, K., Yu, K., Hanisco, T. F., Wolfe, G. M., Arkinson, H. L., Pye, H. O. T., Froyd, K. D., Liao, J. and McNeill, V. F.: Aqueous-phase mechanism for secondary organic aerosol formation from isoprene: application to the southeast United States and co-benefit of SO₂ emission controls, *Atmos Chem Phys*, 16(3), 1603–1618, doi:10.5194/acp-16-1603-2016, 2016.
- Middleton, P., Stockwell, W. R. and Carter, W. P. L.: Aggregation and analysis of volatile organic compound emissions for
10 regional modeling, *Atmospheric Environ. Part Gen. Top.*, 24(5), 1107–1133, doi:10.1016/0960-1686(90)90077-Z, 1990.
- Mills, G. P., Hiatt-Gipson, G. D., Bew, S. P. and Reeves, C. E.: Measurement of isoprene nitrates by GCMS, *Atmos Meas Tech*, 9(9), 4533–4545, doi:10.5194/amt-9-4533-2016, 2016.
- Müller, J.-F., Peeters, J. and Stavrou, T.: Fast photolysis of carbonyl nitrates from isoprene, *Atmos Chem Phys*, 14(5), 2497–2508, doi:10.5194/acp-14-2497-2014, 2014.
- 15 Ng, N. L., Kwan, A. J., Surratt, J. D., Chan, A. W. H., Chhabra, P. S., Sorooshian, A., Pye, H. O. T., Crounse, J. D., Wennberg, P. O., Flagan, R. C. and Seinfeld, J. H.: Secondary organic aerosol (SOA) formation from reaction of isoprene with nitrate radicals (NO₃), *Atmos Chem Phys*, 8(14), 4117–4140, doi:10.5194/acp-8-4117-2008, 2008.
- Nguyen, T. B., Crounse, J. D., Schwantes, R. H., Teng, A. P., Bates, K. H., Zhang, X., St. Clair, J. M., Brune, W. H., Tyndall, G. S., Keutsch, F. N., Seinfeld, J. H. and Wennberg, P. O.: Overview of the Focused Isoprene eXperiment at the
20 California Institute of Technology (FIXCIT): mechanistic chamber studies on the oxidation of biogenic compounds, *Atmos Chem Phys*, 14(24), 13531–13549, doi:10.5194/acp-14-13531-2014, 2014.
- Nguyen, T. B., Crounse, J. D., Teng, A. P., Clair, J. M. S., Paulot, F., Wolfe, G. M. and Wennberg, P. O.: Rapid deposition of oxidized biogenic compounds to a temperate forest, *Proc. Natl. Acad. Sci.*, 112(5), E392–E401, doi:10.1073/pnas.1418702112, 2015.
- 25 Ott, L. E., Pickering, K. E., Stenchikov, G. L., Allen, D. J., DeCaria, A. J., Ridley, B., Lin, R.-F., Lang, S. and Tao, W.-K.: Production of lightning NO_x and its vertical distribution calculated from three-dimensional cloud-scale chemical transport model simulations, *J. Geophys. Res. Atmospheres*, 115(D4), D04301, doi:10.1029/2009JD011880, 2010.
- Paulot, F., Crounse, J. D., Kjaergaard, H. G., Kroll, J. H., Seinfeld, J. H. and Wennberg, P. O.: Isoprene photooxidation: new insights into the production of acids and organic nitrates, *Atmos Chem Phys*, 9(4), 1479–1501, doi:10.5194/acp-9-1479-
30 2009, 2009a.
- Paulot, F., Crounse, J. D., Kjaergaard, H. G., Kürten, A., Clair, J. M. S., Seinfeld, J. H. and Wennberg, P. O.: Unexpected Epoxide Formation in the Gas-Phase Photooxidation of Isoprene, *Science*, 325(5941), 730–733, doi:10.1126/science.1172910, 2009b.



- Paulot, F., Henze, D. K. and Wennberg, P. O.: Impact of the isoprene photochemical cascade on tropical ozone, *Atmos Chem Phys*, 12(3), 1307–1325, doi:10.5194/acp-12-1307-2012, 2012.
- Peeters, J. and Müller, J.-F.: HOx radical regeneration in isoprene oxidation via peroxy radical isomerisations. II: experimental evidence and global impact, *Phys. Chem. Chem. Phys.*, 12(42), 14227–14235, doi:10.1039/C0CP00811G, 5 2010.
- Peeters, J., Müller, J.-F., Stavrou, T. and Nguyen, V. S.: Hydroxyl Radical Recycling in Isoprene Oxidation Driven by Hydrogen Bonding and Hydrogen Tunneling: The Upgraded LIM1 Mechanism, *J. Phys. Chem. A*, 118(38), 8625–8643, doi:10.1021/jp5033146, 2014.
- Perring, A. E., Wisthaler, A., Graus, M., Wooldridge, P. J., Lockwood, A. L., Mielke, L. H., Shepson, P. B., Hansel, A. and 10 Cohen, R. C.: A product study of the isoprene+NO₃ reaction, *Atmos Chem Phys*, 9(14), 4945–4956, doi:10.5194/acp-9-4945-2009, 2009a.
- Perring, A. E., Bertram, T. H., Wooldridge, P. J., Fried, A., Heikes, B. G., Dibb, J., Crouse, J. D., Wennberg, P. O., Blake, N. J., Blake, D. R., Brune, W. H., Singh, H. B. and Cohen, R. C.: Airborne observations of total RONO₂: new constraints on the yield and lifetime of isoprene nitrates, *Atmos Chem Phys*, 9(4), 1451–1463, doi:10.5194/acp-9-1451-2009, 2009b.
- 15 Perring, A. E., Pusede, S. E. and Cohen, R. C.: An Observational Perspective on the Atmospheric Impacts of Alkyl and Multifunctional Nitrates on Ozone and Secondary Organic Aerosol, *Chem. Rev.*, 113(8), 5848–5870, doi:10.1021/cr300520x, 2013.
- Praske, E., Crouse, J. D., Bates, K. H., Kurtén, T., Kjaergaard, H. G. and Wennberg, P. O.: Atmospheric Fate of Methyl Vinyl Ketone: Peroxy Radical Reactions with NO and HO₂, *J. Phys. Chem. A*, 119(19), 4562–4572, doi:10.1021/jp5107058, 20 2015.
- Pye, H. O. T., Luecken, D. J., Xu, L., Boyd, C. M., Ng, N. L., Baker, K. R., Ayres, B. R., Bash, J. O., Baumann, K., Carter, W. P. L., Edgerton, E., Fry, J. L., Hutzell, W. T., Schwede, D. B. and Shepson, P. B.: Modeling the Current and Future Roles of Particulate Organic Nitrates in the Southeastern United States, *Environ. Sci. Technol.*, 49(24), 14195–14203, doi:10.1021/acs.est.5b03738, 2015.
- 25 Rindelaub, J. D., McAvey, K. M. and Shepson, P. B.: The photochemical production of organic nitrates from α -pinene and loss via acid-dependent particle phase hydrolysis, *Atmos. Environ.*, 100, 193–201, doi:10.1016/j.atmosenv.2014.11.010, 2015.
- Rindelaub, J. D., Borca, C. H., Hostetler, M. A., Slade, J. H., Lipton, M. A., Slipchenko, L. V. and Shepson, P. B.: The acid-catalyzed hydrolysis of an α -pinene-derived organic nitrate: kinetics, products, reaction mechanisms, and atmospheric 30 impact, *Atmos Chem Phys*, 16(23), 15425–15432, doi:10.5194/acp-16-15425-2016, 2016.
- Rollins, A. W., Kiendler-Scharr, A., Fry, J. L., Brauers, T., Brown, S. S., Dorn, H.-P., Dubé, W. P., Fuchs, H., Mensah, A., Mentel, T. F., Rohrer, F., Tillmann, R., Wegener, R., Wooldridge, P. J. and Cohen, R. C.: Isoprene oxidation by nitrate radical: alkyl nitrate and secondary organic aerosol yields, *Atmos Chem Phys*, 9(18), 6685–6703, doi:10.5194/acp-9-6685-2009, 2009.



- Romer, P. S., Duffey, K. C., Wooldridge, P. J., Allen, H. M., Ayres, B. R., Brown, S. S., Brune, W. H., Crouse, J. D., de Gouw, J., Draper, D. C., Feiner, P. A., Fry, J. L., Goldstein, A. H., Koss, A., Misztal, P. K., Nguyen, T. B., Olson, K., Teng, A. P., Wennberg, P. O., Wild, R. J., Zhang, L. and Cohen, R. C.: The lifetime of nitrogen oxides in an isoprene-dominated forest, *Atmos Chem Phys*, 16(12), 7623–7637, doi:10.5194/acp-16-7623-2016, 2016.
- 5 Rosen, R. S., Wood, E. C., Wooldridge, P. J., Thornton, J. A., Day, D. A., Kuster, W., Williams, E. J., Jobson, B. T. and Cohen, R. C.: Observations of total alkyl nitrates during Texas Air Quality Study 2000: Implications for O₃ and alkyl nitrate photochemistry, *J. Geophys. Res. Atmospheres*, 109(D7), doi:10.1029/2003JD004227, 2004.
- Russell, M. and Allen, D. T.: Predicting secondary organic aerosol formation rates in southeast Texas, *J. Geophys. Res. Atmospheres*, 110(D7), D07S17, doi:10.1029/2004JD004722, 2005.
- 10 Saunders, S. M., Jenkin, M. E., Derwent, R. G. and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, *Atmos Chem Phys*, 3(1), 161–180, doi:10.5194/acp-3-161-2003, 2003.
- Schell, B., Ackermann, I. J., Hass, H., Binkowski, F. S. and Ebel, A.: Modeling the formation of secondary organic aerosol within a comprehensive air quality model system, *J. Geophys. Res. Atmospheres*, 106(D22), 28275–28293, doi:10.1029/2001JD000384, 2001.
- 15 Schwantes, R. H., Teng, A. P., Nguyen, T. B., Coggon, M. M., Crouse, J. D., St. Clair, J. M., Zhang, X., Schilling, K. A., Seinfeld, J. H. and Wennberg, P. O.: Isoprene NO₃ Oxidation Products from the RO₂ + HO₂ Pathway, *J. Phys. Chem. A*, doi:10.1021/acs.jpca.5b06355, 2015.
- Shepson, P. B., Mackay, E. and Muthuramu, K.: Henry's Law Constants and Removal Processes for Several Atmospheric β-Hydroxy Alkyl Nitrates, *Environ. Sci. Technol.*, 30(12), 3618–3623, doi:10.1021/es960538y, 1996.
- 20 St. Clair, J. M., Rivera-Rios, J. C., Crouse, J. D., Knap, H. C., Bates, K. H., Teng, A. P., Jørgensen, S., Kjaergaard, H. G., Keutsch, F. N. and Wennberg, P. O.: Kinetics and Products of the Reaction of the First-Generation Isoprene Hydroxy Hydroperoxide (ISOPOOH) with OH, *J. Phys. Chem. A*, doi:10.1021/acs.jpca.5b06532, 2015.
- Stavrakou, T., Peeters, J. and Müller, J.-F.: Improved global modelling of HO_x recycling in isoprene oxidation: evaluation against the GABRIEL and INTEX-A aircraft campaign measurements, *Atmos Chem Phys Discuss*, 10(7), 16551–16588, doi:10.5194/acpd-10-16551-2010, 2010.
- 25 Teng, A. P., Crouse, J. D., Lee, L., St. Clair, J. M., Cohen, R. C. and Wennberg, P. O.: Hydroxy nitrate production in the OH-initiated oxidation of alkenes, *Atmos Chem Phys*, 15(8), 4297–4316, doi:10.5194/acp-15-4297-2015, 2015.
- Teng, A. P., Crouse, J. D. and Wennberg, P. O.: Isoprene Peroxy Radical Dynamics, *J. Am. Chem. Soc.*, 139(15), 5367–5377, doi:10.1021/jacs.6b12838, 2017.
- 30 Travis, K. R., Jacob, D. J., Fisher, J. A., Kim, P. S., Marais, E. A., Zhu, L., Yu, K., Miller, C. C., Yantosca, R. M., Sulprizio, M. P., Thompson, A. M., Wennberg, P. O., Crouse, J. D., St. Clair, J. M., Cohen, R. C., Laughner, J. L., Dibb, J. E., Hall, S. R., Ullmann, K., Wolfe, G. M., Pollack, I. B., Peischl, J., Neuman, J. A. and Zhou, X.: Why do models overestimate surface ozone in the Southeast United States?, *Atmos Chem Phys*, 16(21), 13561–13577, doi:10.5194/acp-16-13561-2016, 2016.



- Valin, L. C., Russell, A. R. and Cohen, R. C.: Variations of OH radical in an urban plume inferred from NO₂ column measurements, *Geophys. Res. Lett.*, 40(9), 1856–1860, doi:10.1002/grl.50267, 2013.
- Wennberg, P. O., Bates, K. H., Crouse, J. D., Dodson, L. G., McVay, R. C., Mertens, L. A., Nguyen, T. B., Praske, E., Schwantes, R. H., Smarte, M. D., St Clair, J. M., Teng, A. P., Zhang, X. and Seinfeld, J. H.: Gas-Phase Reactions of
5 Isoprene and Its Major Oxidation Products, *Chem. Rev.*, 118(7), 3337–3390, doi:10.1021/acs.chemrev.7b00439, 2018.
- Wesely, M. L.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models, *Atmospheric Environ.* 1967, 23(6), 1293–1304, doi:10.1016/0004-6981(89)90153-4, 1989.
- Wolfe, G. M., Kaiser, J., Hanisco, T. F., Keutsch, F. N., de Gouw, J. A., Gilman, J. B., Graus, M., Hatch, C. D., Holloway, J., Horowitz, L. W., Lee, B. H., Lerner, B. M., Lopez-Hilfiker, F., Mao, J., Marvin, M. R., Peischl, J., Pollack, I. B.,
10 Roberts, J. M., Ryerson, T. B., Thornton, J. A., Veres, P. R. and Warneke, C.: Formaldehyde production from isoprene oxidation across NO_x regimes, *Atmos Chem Phys*, 16(4), 2597–2610, doi:10.5194/acp-16-2597-2016, 2016.
- Wu, S., Mickley, L. J., Jacob, D. J., Logan, J. A., Yantosca, R. M. and Rind, D.: Why are there large differences between models in global budgets of tropospheric ozone?, *J. Geophys. Res. Atmospheres*, 112(D5), D05302, doi:10.1029/2006JD007801, 2007.
- 15 Xie, Y., Paulot, F., Carter, W. P. L., Nolte, C. G., Luecken, D. J., Hutzell, W. T., Wennberg, P. O., Cohen, R. C. and Pinder, R. W.: Understanding the impact of recent advances in isoprene photooxidation on simulations of regional air quality, *Atmos Chem Phys*, 13(16), 8439–8455, doi:10.5194/acp-13-8439-2013, 2013.
- Xiong, F., McAvey, K. M., Pratt, K. A., Groff, C. J., Hostetler, M. A., Lipton, M. A., Starn, T. K., Seeley, J. V., Bertman, S. B., Teng, A. P., Crouse, J. D., Nguyen, T. B., Wennberg, P. O., Misztal, P. K., Goldstein, A. H., Guenther, A. B., Koss, A.
20 R., Olson, K. F., de Gouw, J. A., Baumann, K., Edgerton, E. S., Feiner, P. A., Zhang, L., Miller, D. O., Brune, W. H. and Shepson, P. B.: Observation of isoprene hydroxynitrates in the southeastern United States and implications for the fate of NO_x, *Atmos Chem Phys*, 15(19), 11257–11272, doi:10.5194/acp-15-11257-2015, 2015.
- Xiong, F., Borca, C. H., Slipchenko, L. V. and Shepson, P. B.: Photochemical degradation of isoprene-derived 4,1-nitrooxy enal, *Atmos Chem Phys*, 16(9), 5595–5610, doi:10.5194/acp-16-5595-2016, 2016.




Figure 1: Schematic representation of the formation of isoprene nitrates (in green) initiated by OH oxidation. Re-released of the consumed NO_x to the atmosphere by chemical loss processes of oxidation, ozonolysis and photolysis of organic nitrate is shown in red. See Table S2 in the Supplement for species descriptions.

5




Figure 2: Schematic representation of the formation of isoprene nitrates (in green) initiated by NO_3 oxidation. For simplification, fates of only one isomer of hydroxy nitrates (IHNB and IHND) and nitrooxyhydroperoxide (INPB and INPD) are shown. See Table S2 in the Supplement for species descriptions.




Figure 3: (a) Concentrations of total organic nitrates for average of the model simulation period (June 2013). (b) Fractional contribution of BVOC-derived organic nitrates to total organic nitrates. The location of SOAS CTR ground site is circled in the figure.

5




Figure 4: Median diurnal cycles of observed and simulated total organic nitrates at Centreville during the 2013 SOAS campaign. The vertical bars show the interquartile range of the hourly data. The panel includes mean of the simulated and observed organic nitrates and the normalized mean bias (NMB) in percent.




Figure 5: The composition of the (a) simulated organic nitrates by WRF-Chem using RACM2_Berkeley2 and (b) observed organic nitrates during SOAS at CTR.




Figure 6: The modeled and observed correlations between O_x (=O₃+NO₂) and organic nitrates concentrations at daytime during SOAS. The lines indicate linear regression (intercept fixed at 0) and confidence intervals. The panel includes slopes of the lines and root mean square errors (RMSE).




Figure 7: The modeled and observed correlations between CH_2O and organic nitrates at daytime during SOAS. The slope shows the best fit line, with an intercept allowed to differ from zero to consider the possibility of a background CH_2O . The panel includes slopes of the lines and root mean square errors (RMSE).




Figure 8: Diurnal cycle of fractional organic nitrate (a) productions and (b) concentrations simulated by WRF-Chem averaged at boundary layer at the CTR site during SOAS.




Figure 9: The simulated concentration of 1st- (blue) and 2nd- (red) generation organic nitrates versus their loss rates at daytime during SOAS. Slopes of the linear fit give their lifetimes. The Concentrations and loss rates of 1st-generation nitrates are divided by 2.




Figure 10: Contribution of different fates to (a) the first and (b) second generation of isoprene and monoterpene nitrates loss.




Figure 11: The simulated instantaneous production of NO_x from loss of organic nitrates versus the instantaneous loss of NO_x to production of organic nitrates. The slope shows the NO_x recycle efficiency.

5