

Nitrogen oxides in the global upper troposphere: interpreting cloud-sliced NO₂ observations from the OMI satellite instrument

Eloise A. Marais^{1,2,*}, Daniel J. Jacob^{2,3}, Sungyeon Choi⁴, Joanna Joiner^{4,5}, Maria Belmonte-Rivas⁶, Ronald C. Cohen^{7,8}, Steffen Beirle⁹, Lee T. Murray¹⁰, Luke D. Schiferl^{11,**}, Viral Shah¹², Lyatt Jaeglé¹²

5 ¹School of Geography, Earth, and Environmental Sciences, University of Birmingham, Birmingham, UK.

²John A Paulson School of Engineering and Applied Sciences, Harvard University, Cambridge, MA, USA.

³Earth and Planetary Sciences, Harvard University, Cambridge, MA, USA.

⁴Science Systems and Applications Inc., Lanham, MD.

⁵NASA Goddard Space Flight Center, Greenbelt, MD.

10 ⁶Royal Netherlands Meteorology Institute, De Bilt, the Netherlands.

⁷Department of Chemistry, University of California at Berkeley, Berkeley, CA.

⁸Department of Earth and Planetary Science, University of California at Berkeley, Berkeley, CA.

⁹Max-Planck-Institut für Chemie, Mainz, Germany.

¹⁰Department of Earth and Environmental Sciences, University of Rochester, Rochester, New York, USA.

15 ¹¹Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, USA.

¹²Department of Atmospheric Sciences, University of Washington, Seattle, WA, USA.

* Now at: Department of Physics and Astronomy, University of Leicester, Leicester, UK.

** Now at: Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY, USA.

Correspondence to: Eloise A. Marais (eloise.marais@le.ac.uk)

20 **Abstract.** Nitrogen oxides (NO_x ≡ NO + NO₂) in the upper troposphere (UT) have a large impact on global tropospheric ozone and OH (the main atmospheric oxidant). New cloud-sliced observations of UT NO₂ at 450-280 hPa (~6-9 km) from the OMI satellite instrument produced by NASA and KNMI provide global coverage to test our understanding of the factors controlling UT NO_x. We find that these products offer useful information when averaged over coarse scales (20° × 32°, seasonal), and that the NASA product is more consistent with aircraft observations of UT NO₂. Correlation with LIS/OTD satellite observations of lightning flash frequencies suggests that lightning is the dominant source of NO_x to the upper troposphere except for extratropical latitudes in winter. The NO₂ background in the absence of lightning is 10-20 pptv. We infer a global mean NO_x yield of 280 ± 80 moles per lightning flash, with no significant difference between the tropics and mid-latitudes, and a global lightning NO_x source of 5.9 ± 1.7 Tg N a⁻¹. There is indication that the NO_x yield per flash increases with lightning flash footprint and with flash energy.

1. Introduction

30 Nitrogen oxides (NO_x ≡ NO + NO₂) in the upper troposphere (UT) have profound effects on the oxidizing capacity of the atmosphere and on climate, but the factors controlling their concentrations are poorly understood. NO_x in the UT impacts climate by efficiently producing ozone where it is a potent greenhouse gas (Dahlmann et al., 2011; Worden et al., 2011; Rap et al., 2015) and by increasing the concentration of OH (the main tropospheric oxidant) (Murray et al., 2012; Murray et al., 2014). Primary NO_x sources in the UT include lightning, aircraft, convective injection, and downwelling from the stratosphere (Ehhalt et al., 1992; Jaeglé et al., 1998; Bertram et al., 2007). NO_x cycles chemically with reservoir species including nitric acid (HNO₃), pernitric acid (HNO₄), dinitrogen pentoxide (N₂O₅), peroxyacetyl nitrate (PAN), and other organic nitrates, thus defining the NO_y chemical family (NO_y ≡ NO_x + reservoirs). Effective loss of NO_x from the UT is through subsidence of NO_y to lower altitudes where deposition of HNO₃ provides the ultimate sink. The residence time of NO_y in the UT is 10-20 days (Prather and Jacob, 1997). The lifetime of

NO_x against conversion to short-lived reservoirs varies from ~3 hours in the convective outflow of thunderstorms to 0.5-1.5 days
40 in background air (Nault et al., 2016). Chemical recycling from these reservoirs maintains relatively high UT NO_x background
concentrations (Bradshaw et al., 2000; Baehr et al., 2003; Nault et al., 2016).

Representation of lightning NO_x in chemical transport models (CTMs) is particularly uncertain. Physically-based parameterizations
relating lightning frequency to deep convective cloud tops, convective mass flux, convective precipitation, or high-cloud ice
45 content have poor predictive capability (Tost et al., 2007; Allen et al., 2010; Murray et al., 2012; Finney et al., 2014), limiting our
ability to estimate the response of lightning NO_x to future climate (Finney et al., 2016; 2018). An alternative is to prescribe flash
densities from space-based observations and static NO_x production rates per flash (Sauvage et al., 2007; Allen et al., 2010; Murray
et al., 2012). NO_x production efficiencies per flash in the literature vary from <10 to 5000 moles nitrogen per flash (mol N fl⁻¹)
(Schumann and Huntrieser, 2007; Murray, 2016). Global chemical transport models (CTMs) typically use 100-500 mol N fl⁻¹,
50 sometimes assuming higher production rates at mid-latitudes than in the tropics (Hudman et al., 2007; Ott et al., 2010), and a global
lightning NO_x source of 3-7 Tg N a⁻¹ to match observations of tropospheric ozone and NO_y species (Sauvage et al., 2007).

Our understanding of UT NO_x has so far been evaluated with observations from aircraft campaigns (Drummond et al., 1988; Jacob
et al., 1996; Crawford et al., 1997; Jaeglé et al., 1998; Bradshaw et al., 2000; Hudman et al., 2007; Stratmann et al., 2016). There
55 are also long-term NO_x measurements from instruments onboard commercial aircraft dating back to the 1990s, but these are mostly
over the north Atlantic and the NO₂ measurements have low precision and interference from thermally unstable NO_x reservoir
compounds (Brunner et al., 2001). A number of studies have used satellite observations of tropospheric NO₂ columns from solar
backscatter to infer lightning NO_x emissions (Beirle et al., 2010; Pickering et al., 2016; Bucselá et al., 2010), including in
combination with global models (Boersma et al., 2005; Martin et al., 2007; Miyazaki et al., 2014). These studies estimate global
60 lightning NO_x emission of 1 to 8 Tg N a⁻¹.

New cloud-sliced satellite products of tropospheric NO₂ mixing ratios at 280-450 hPa (~6-9 km) offer additional vertical resolution
by retrieving partial NO₂ columns above clouds and exploiting differences in heights of neighboring clouds to calculate NO₂
mixing ratios (Choi et al., 2014; Belmonte-Rivas et al., 2015). There are two new products of seasonal mean UT NO₂ mixing ratios
65 retrieved from Ozone Monitoring Instrument (OMI) partial NO₂ columns by research groups at KNMI and NASA. The KNMI
product has been evaluated against UT NO₂ from the Tracer Model version 4 (TM4) CTM. Large regional differences between
OMI and TM4 are attributed to model deficiencies in lightning NO_x and uplift of anthropogenic pollution (Belmonte-Rivas et al.,
2015). The NASA UT product is new to this work and follows a similar retrieval approach to the mid-tropospheric (900-650 hPa)
product of Choi et al. (2014). That product was evaluated with aircraft observations of NO₂ and interpreted with the Global
70 Modeling Initiative (GMI) CTM (Choi et al., 2014). Choi et al. (2014) identified large discrepancies between modeled and observed
NO₂ seasonality over regions influenced by pollution and lightning.

Here we compare the two UT NO₂ products, obtained with distinct retrieval methods, and use aircraft observations of NO₂ from
multiple NASA DC8 aircraft campaigns to arbitrate and evaluate the information that can be derived from the satellite datasets.
75 We go on to test current understanding of UT NO_x and the implications for lightning emissions using the GEOS-Chem CTM.

2. OMI observations of upper troposphere NO₂


OMI is onboard the NASA Aura satellite launched into sun-synchronous orbit in July 2004. It has an overpass time of about 13h30 local time (LT), a swath width of 2600 km, and a horizontal resolution of $13 \text{ km} \times 24 \text{ km}$ at nadir (Levelt et al., 2006). Columns of NO_2 along the instrument viewing path (slant columns) are obtained by spectral fitting of solar backscattered radiation in the 405-465 nm window (Boersma et al., 2011). Standard products of total and tropospheric column NO_2 are screened for cloudy scenes using a cloud radiance fraction threshold of 0.5. Partial columns of NO_2 above cloudy scenes can be used to estimate vertically resolved NO_2 mixing ratios, as was first demonstrated with satellite observations of ozone (Ziemke et al., 2001). This approach, so-called cloud slicing, assumes a uniform trace gas concentration between two horizontally nearby clouds at different altitudes, so that the UT NO_2 mixing ratio is proportional to the slope of the partial columns versus the corresponding cloud pressures at the optical centre of the cloud. Two products of seasonal mean UT NO_2 have been retrieved from OMI following distinct retrieval steps detailed below: a product from KNMI at 330-450 hPa for 2006 (Belmonte-Rivas et al., 2015) and from NASA at 280-450 hPa for 2005-2007 following an approach similar to that used to retrieve mid-tropospheric NO_2 (Choi et al., 2014). In what follows we distinguish the two OMI NO_2 products as KNMI and NASA.

The KNMI product uses DOMINO v2.0 slant columns (Boersma et al., 2011) and OMCLDO2 cloud fractions and altitudes (Acarreta et al., 2004) over partially to very cloudy scenes (cloud radiance fraction > 0.5). Contamination due to NO_2 from below (up to 66% over polluted land masses) is estimated using the TM4 model and removed. Stratospheric NO_2 from an assimilated product (Belmonte-Rivas et al., 2014) is also removed. An air mass factor (AMF) (detailed in Boersma et al. (2004)) that accounts for viewing geometry, surface albedo, light attenuation by gases along the viewing path, and sensitivity to NO_2 vertical distribution is applied to the resultant partial slant columns to convert to vertical columns. Additional data filtering removes scenes with solar zenith angle (SZA) $\geq 70^\circ$ and surface albedo $\geq 30\%$. Resultant daily vertical partial columns are aggregated on consistent pressure and horizontal ($1^\circ \times 1^\circ$) grids and used to determine seasonal mean UT NO_2 mixing ratios for gridsquares with at least 30 measurements. UT NO_2 centred at 380 hPa (range 330-450 hPa) is estimated as the difference between partial tropospheric columns retrieved above two neighboring clouds with cloud pressures in the ranges 330-450 hPa and 380-500 hPa, respectively. Biases from sampling cloudy scenes, such as the effect of clouds on photochemistry, are corrected using TM4. These are small (typically $< 20\%$) in the UT (Belmonte-Rivas et al., 2015).

The NASA UT NO_2 product for 2005-2007, centred at 350 hPa (~ 280 -450 hPa), uses updated version 3 slant columns (OMNO2 v3.0) (Krotkov et al., 2017) that correct for a positive bias in the DOMINO v2.0 product with improved spectral fitting (Marchenko et al., 2015; van Geffen et al., 2015). Partial columns from the cloud height to the top of the atmosphere are retrieved for individual OMI pixels above very cloudy scenes (cloud radiance fraction > 0.7) to minimize contamination from below. Cloud fraction and height is from the OMCLDO2 product (Acarreta et al., 2004). The AMF accounts for viewing path geometry and light scattering by clouds with uniform scatter that are optically thick and geometrically thin (near-Lambertian clouds). Data filtering is applied to remove scenes with SZA $> 80^\circ$, snow/ice cover, and severe aerosol pollution that could be misclassified as clouds. Daily UT NO_2 is estimated for neighboring partial columns with sufficient cloud variability (cloud pressure distance > 160 hPa) and well-mixed NO_2 (NO_2 vertical gradient < 0.33 pptv hPa $^{-1}$ diagnosed with the GMI CTM). The stratospheric column is assumed uniform above neighboring clouds and so is removed when differencing two nearby partial columns. Daily values of UT NO_2 are gridded to obtain seasonal means at $5^\circ \times 8^\circ$ (latitude \times longitude) for scenes with at least 50 measurements. Gaussian weighting is applied to assign higher weighting to UT NO_2 closest to 350 hPa. Choi et al. (2014) used a similar approach to retrieve mid-tropospheric NO_2 except that cloud fraction and height were from the OMCLDRR product, and successful retrieval required a stricter cloud radiance fraction of 0.9, a minimum of 30 measurements, and a wider minimum cloud pressure distance of 200 hPa. A shift in cloud radiance fraction

threshold from 0.9 (Choi et al., 2014) to 0.7 (this work) only introduces a small (<5%) difference in the retrieved partial columns due to contamination from below, as estimated by Pickering et al. (2016) for OMI scenes over the Gulf of Mexico.

120 Figure 1 compares seasonal mean UT NO₂ from the two satellite products in December-February and June-August. KNMI NO₂ is gridded to the NASA coarse grid. Data for March-May and September-November are in the Supplement (Figure S1). KNMI NO₂ has greater coverage than the NASA product, due to a lower cloud fraction threshold in the retrieval. The two products exhibit very different spatial features. Spatial correlation between the two products (Pearson's correlation coefficient between coincident gridsquares) is R = 0.41 in December-February and R = 0.38 in June-August. There is marginal improvement in the correlation with further spatial averaging. At 20° × 32° we find R = 0.50 in December-February and R = 0.45 in June-August. The correlation only increases substantially in September-November from R = 0.49 at 5° × 8° (Figure S1) to R = 0.66 at 20° × 32°. KNMI is systematically lower than NASA in all seasons for coincident gridsquares, varying from 16% lower in June-August to 48% lower in December-February at 20° × 32°.



130 **Figure 1. Upper troposphere (UT) NO₂ from the OMI satellite instrument. Seasonal mean UT NO₂ from KNMI in 2006 at 330-450 hPa (top) is compared to NASA in 2005-2007 at 280-450 hPa (bottom). Data are at 5° × 8° horizontal resolution for December-February (left) and June-August (right). Grey areas indicate no data and, for NASA, scenes with fewer than 50 measurements.**


135 Contamination of UT NO₂ from below the cloud may still be present in the datasets despite attempts to correct for this using the TM4 model in the case of KNMI and by only considering very cloudy scenes in the case of NASA. These include a large enhancement in KNMI NO₂ (> 90 pptv) over southern Africa in June-August when there is intense biomass burning, and the NO₂ hotspot over northeast China in all seasons in both products (Figures 1, S1). Belmonte-Rivas et al. (2015) caution that the contamination correction in the KNMI product relies on accurate simulation of NO₂ vertical distribution.

140 3. Evaluation of OMI upper troposphere NO₂ with aircraft observations

We evaluate the OMI UT NO₂ products with observations from NASA DC8 aircraft campaigns over North America and Greenland in spring-summer, for which dense coverage is available (Figure 2). These include the INTEX-A, INTEX-B, ARCTAS, DC3, and SEAC⁴RS campaigns. Only INTEX-B is in the same year as the OMI products but we consider interannual variability to be only a small source of error. All NO₂ measurements are from thermal-dissociation laser-induced fluorescence (TD-LIF) instruments
145 (Day et al., 2002). These are susceptible to interference from decomposition of thermally unstable reservoir compounds including methyl peroxy nitrate (CH₃O₂NO₂) and HNO₄ (Browne et al., 2011). Publicly available DC3 and SEAC⁴RS TD-LIF NO₂ are already corrected for this interference. We apply a correction for the other campaigns using the relationship between temperature and percentage interference from Browne et al. (2011). Observed mean ambient air temperature in the UT during INTEX-A was
150 246 K, corresponding to 20% interference. That for INTEX-B was 241 K (30% interference) and 236 K for ARCTAS (38% interference).

There are also NO₂ observations from the recent NASA ATom campaign, and from the In-service Aircraft for a Global Observing System (IAGOS) commercial aircraft campaign (Berkes et al., 2017). These use chemiluminescence instruments that are also susceptible to interference. Chemiluminescence and TD-LIF NO₂ are consistent during the SEAC⁴RS campaign for the altitude
155 range considered in this work (6-9 km) (Travis et al., 2016), but the interference from chemiluminescence is challenging to quantify, due to dependence on the operator and instrument design that varies across campaigns (Reed et al., 2016).

Figure 2 shows the sampling extent of TD-LIF UT NO₂ over North America and Greenland in spring-summer at 450-280 hPa around the satellite overpass (11h00-16h00 LT) for scenes not influenced by the stratosphere (diagnosed with collocated ozone/CO
160 > 1.25 mol mol⁻¹ (Hudman et al., 2007)). Concentrations of UT NO₂ exceed 80 pptv over the eastern US due to lightning NO_x emissions and convective transport of boundary layer pollution, and are < 30 pptv over the rest of the domain.



165 **Figure 2. NASA DC8 upper troposphere NO₂ over North America in spring-summer (March-August). Observations are from the TD-LIF instrument at 450-280 hPa, 11h00-16h00 local time, and without stratospheric influence. Campaigns include INTEX-A in June-August 2004 (Singh et al., 2006), INTEX-B in March-May 2006 (Singh et al., 2009), ARCTAS in March-April and June-July 2008 (Jacob et al., 2010), DC3 in May-June 2012 (Barth et al., 2015), and SEAC⁴RS in August 2013 (Toon et al., 2016).**

Figure 3 shows the spatial correlation between the March-August mean gridded aircraft data and the OMI UT NO₂ from the KNMI and NASA products as a function of horizontal resolution. There is no significant spatial correlation between the OMI products and aircraft NO₂ at 5° × 8° (R < 0.1) and 10° × 16° (R < 0.2). The correlation improves with further spatial averaging, peaking at 20° × 32° (R = 0.56 for KNMI, R = 0.64 for NASA). The satellite products are also spatially consistent at this resolution (R = 0.89), but KNMI is 43% lower than NASA.




Figure 3. Evaluation of OMI and GEOS-Chem upper troposphere NO₂ with aircraft observations. Individual points are Pearson's correlation coefficients between gridded March-August mean UT NO₂ measured from the aircraft and OMI KNMI in 2006 (blue), OMI NASA in 2005-2007 (red), and GEOS-Chem in 2006 (green), for grid averaging domains of 5° × 8° (latitude × longitude), 10° × 16°, 15° × 24°, 20° × 32°, and 25° × 40°. Values inset are the number of points at each resolution. The domain sampled is shown in Figure 2.

Figure 4 compares the spatial distribution of OMI and aircraft UT NO₂ at 20° × 32° over North America. Domain mean KNMI UT NO₂ is 38% lower than the aircraft observations, compared to 2.2% higher for NASA UT NO₂. Both products exhibit less variability (reduced major axis, RMA, regression slopes < 1) and high bias in background NO₂ compared to the aircraft observations (positive RMA intercepts of 5.9 ± 1.4 pptv for KNMI and 9.2 ± 2.7 pptv for NASA). We proceed with the NASA UT NO₂ product at 20° × 32°, as correlation peaks at this resolution and the NASA product is more consistent with domain mean aircraft UT NO₂ than the KNMI product.

4. Constraints on upper tropospheric NO_x

The NASA product provides near-global coverage of UT NO₂ to assess current understanding of regional UT NO_x sources and dynamics by comparing to UT NO₂ from the GEOS-Chem CTM (version 10-01; http://wiki.seas.harvard.edu/geos-chem/index.php/GEOS-Chem_v10-01) driven with NASA MERRA-2 reanalysis meteorology. The model horizontal resolution is 2° × 2.5° and the output is regridded to 20° × 32° for comparison with OMI. GEOS-Chem is sampled under all-sky conditions in the satellite overpass window (12h00-15h00 local time). We find that the effect on NO₂ of sampling the model under cloudy conditions is small. Isolating NO₂ under very cloudy conditions using MERRA-2 cloud fractions decreases modeled UT NO₂ by no more than 5 pptv in the tropics/subtropics and less at higher latitudes. We use output from the model for 2006 following a one-

195 year spin-up for chemical initialization. Interannual variability in UT NO₂, determined as the difference between modeled 2006 and multi-year mean (2005-2007) UT NO₂, is small (< 4 pptv) everywhere except central Africa year-round (4-12 pptv), the Arctic north of 60°N (up to 25 pptv), and the Middle East in June-August and northern India in March-May (both 10-20 pptv). Recent evaluation of model NO₂ with observed vertical profiles from the SEAC⁴RS aircraft campaign show no significant bias in the 6-9 km range of the OMI product (Travis et al., 2016; Silvern et al., 2018).

200




Figure 4. March-August upper troposphere NO₂ over North America. All data are at 20° × 32°. Background colors in the different panels show concentrations from KNMI, NASA, and GEOS-Chem (GC). Circles show the aircraft observations (same in all panels). Aircraft observations are for 11h00-16h00 local time (LT). The model is sampled in the satellite overpass time window (12h00-15h00 LT). Model and aircraft data are at 280-450 hPa and screened for stratospheric influence using ozone/CO > 1.25 mol mol⁻¹. Inset boxes show reduced major axis (RMA) regression statistics and mean NO₂ for coincident gridsquares. Grey gridsquares indicate no observations.

205


Local GEOS-Chem emissions of NO_x in the UT include aircraft and lightning. Aircraft emissions from the AEIC inventory (Stettler et al., 2011) total 0.82 Tg N in 2006; much less than lightning in the same year (6.5 Tg N). Transport from the stratosphere is simulated using a climatology of NO_y species concentrations from the GMI model above the tropopause (Murray et al., 2012) and is very small (0.4 Tg N a⁻¹ as total NO_y). Lightning in the model is estimated using the parameterization implemented by Murray et al. (2012). This includes an initial estimate of lightning flashes using the Price and Rind (1992, 1993, 1994) relationship between cloud-top height and lightning flashes. These are then scaled to the same annual global flash frequency (46 fl s⁻¹) and regional

210

distribution as the climatology from the combined Lightning Imaging Sensor (LIS) and Optical Transient Detector (OTD) high-
215 resolution monthly climatology (LIS/OTD HRMC) (Cecil et al., 2014). The standard GEOS-Chem model has higher NO_x yields
per flash at northern mid-latitudes (north of 35°N) than in the tropics ($500 \text{ mol N fl}^{-1}$ versus $260 \text{ mol N fl}^{-1}$), but we find that this
overestimates observed OMI UT NO_2 by 10-20 pptv (20-40%) at northern mid-latitudes in summer when the lightning source is
dominant. Here we address this overestimate by assuming a NO_x yield of $260 \text{ mol N fl}^{-1}$ everywhere. This decreases global lightning
220 NO_x emissions by 15% from 6.5 to 5.5 Tg N a^{-1} . The lightning parameterization in GEOS-Chem does not distinguish lightning
 NO_x production from flashes within or between clouds (intra- or inter-cloud) or from the cloud to the Earth's surface (cloud-to-
ground).

Figure 3 shows the spatial correlation between the model and aircraft observations. The model is more consistent with the aircraft
observations than OMI at fine spatial resolution. Like OMI, GEOS-Chem correlation with the aircraft observations improves with
225 spatial averaging, peaking at $20^\circ \times 32^\circ$ ($R = 0.75$). Figure 4 also shows comparison of March-August GEOS-Chem UT NO_2 with
the aircraft observations at $20^\circ \times 32^\circ$. The model is sampled over the same pressure range as NASA (280-450 hPa) around the
OMI overpass (12h00-15h00 LT) and is filtered for stratospheric influence using model ozone/ $\text{CO} > 1.25 \text{ mol mol}^{-1}$. Domain
average UT NO_2 from the model is 19% lower than the aircraft measurements and the model also overestimates background UT
 NO_2 (intercept = $7.5 \pm 1.0 \text{ pptv}$) and underestimates the variability (slope = 0.45 ± 0.09).


230 Figure 5 compares seasonal mean OMI and GEOS-Chem UT NO_2 in December-February and June-August. The other seasons are
shown in the Supplement (Figure S2). Formation of PAN, HNO_4 and $\text{CH}_3\text{O}_2\text{NO}_2$ accounts for over 75% of NO_x loss in the model
in all seasons. Lower concentrations of UT NO_2 in the northern hemisphere winter compared to summer in the model is mostly
because lightning activity is at a minimum. The model underestimates UT NO_2 in the northern mid-latitudes in winter by 20-40
235 pptv, suggesting misrepresentation of another process in the model, such as excessive NO_x loss by N_2O_5 hydrolysis in aerosols
(Kenagy et al., 2018). The particularly large bias over polluted regions in winter could also be due to contamination of the UT NO_2
retrievals by enhanced boundary layer NO_2 .




240 **Figure 5. Observed and modelled upper troposphere NO₂. The figure shows NASA OMI seasonal mean UT NO₂ for 2005-2007 (top) and corresponding GEOS-Chem model values (bottom). The model is sampled at 280-450 hPa during the satellite overpass (12h00-15h00 LT), and filtered for stratospheric influence. Data are at 20° × 32° horizontal resolution for December-February (left) and June-August (right). Grey gridsquares in the top panel indicate no OMI data.**

Figure 6 shows the log-log relationship between seasonal mean LIS/OTD lightning flash climatology and seasonal mean UT NO₂ from OMI and GEOS-Chem, and the corresponding reduced major axis linear regression fits. Data are divided into northern mid-latitudes and tropics. We exclude the contaminated observations over northeast China and the wintertime northern mid-latitude gridsquares that show no correlation with lightning flashes ($R < 0.1$). Results from multi-model sensitivity studies indicate that UT NO_x in winter is predominantly from surface sources, with a smaller contribution from extra-tropical lightning (Grewe et al., 2001). Background concentration of UT NO₂ (intercepts in Figure 6) from non-lightning sources is 10-20 pptv and is 3-5 pptv higher in the northern mid-latitudes than the tropics. The slopes for the linear fits to lightning flash frequency are consistent between the OMI observations and GEOS-Chem, and show similar slopes for northern mid-latitudes and the tropics. Fitting the ratio between OMI observations and GEOS-Chem on the 20° × 32° grid implies a NO_x yield per flash of 280 ± 80 mol N fl⁻¹ with no significant difference between mid-latitudes and the tropics, and no significant difference with the GEOS-Chem prior estimate of 260 mol N fl⁻¹. Our prior estimate of global lightning source was 5.5 Tg N a⁻¹, and the improved estimate is 5.9 ± 1.7 Tg N a⁻¹.

255



260 **Figure 6. Log-log relationship between upper troposphere NO₂ and lightning flash frequencies, and linear regression fits between the two. Individual points are coincident seasonal mean UT NO₂ from OMI (left) and GEOS-Chem (right) versus seasonal mean LIS/OTD lightning flash climatologies for coincident 20° × 32° gridsquares in the northern mid-latitudes (> 30°N; blue) and tropics (< 30°N; orange). Northern mid-latitude points exclude December-February that show poor correlation with lightning flashes (see text for details). Lines and legends show reduced major axis linear regression fits to the data with corresponding Pearson's correlation coefficients. The regression lines plot as curves on the log-log scale, highlighting the NO₂ background at low lightning flash rates and the correlation of NO₂ with lightning at high flash rates.**



270 **Figure 7. Relationship between OMI and GEOS-Chem derived lightning NO_x production rates and LIS lightning properties: energy (radiance), duration, and footprint area. Individual points are seasonal mean 20° × 32° gridsquares at 40°N-40°S.**

Properties of lightning flashes including energy, duration, and footprint area, have been retrieved from the OTD and LIS sensors (Beirle et al., 2014). The flash footprint area is the spatial extent of lightning detection events contributing to the flash (collection of local events) diagnosed by the satellite data. Figure 7 shows the relationship between OMI and GEOS-Chem derived lightning NO_x production rates and LIS lightning properties from Beirle et al. (2014). The strongest correlation is with lightning extent (R = 0.50), followed by energy (R = 0.40). The correlation with flash duration is weak (R = 0.25). The relationships in Figure 7 suggest a dependence of lightning NO_x production rates on lightning flash energy of $510 \pm 80 \text{ mol N (J m}^{-2} \text{ sr}^{-1} \mu\text{m}^{-1})^{-1}$ and on flash footprint

area of $2.0 \pm 0.3 \text{ mol N km}^{-2}$, possibly offering guidance for relating NO_x yields to physical properties in global models rather than the current approach of assigning static values.

5. Conclusions

280 Measurements of NO_x in the upper troposphere (UT) have mainly been from aircraft campaigns that are limited in space and time. Two new cloud-slicing UT NO_2 products from the Ozone Monitoring Instrument (OMI) produced by KNMI and NASA offer the potential to address uncertainties in our understanding of UT NO_x sources. Here we intercompared these products, evaluated them with aircraft observations, and used them together with GEOS-Chem model simulations to demonstrate a dominance of lightning as a source of UT NO_x .

285

The KNMI and NASA UT NO_2 products use very different retrieval methods. Seasonal mean concentrations from the two products show weak global correlation at the $5^\circ \times 8^\circ$ (latitude \times longitude) resolution of the NASA retrieval, with some improvement when the data are further averaged to $20^\circ \times 32^\circ$ ($R = 0.5\text{-}0.7$). At that resolution they show correlation with in situ aircraft observations of UT NO_2 over North America for different years ($R = 0.56\text{-}0.64$). The KNMI product is biased low by 38% relative to the aircraft observations while the NASA product has no significant bias.

290

We find from the relationship of OMI UT NO_2 with LIS/OTD flash rates that most NO_x in the upper troposphere is from lightning, except in the mid-latitudes in winter. The background NO_2 concentration in the absence of lightning is 10-20 pptv. The relationship suggests no difference in NO_x yields per flash between the mid-latitudes and the tropics, in contrast to the higher yields at mid-latitudes often assumed in models. We derive a global mean lightning NO_x production rate per flash of $280 \pm 80 \text{ mol N fl}^{-1}$, from which we infer a best estimate for the global lightning NO_x emission of $5.9 \pm 1.7 \text{ Tg N a}^{-1}$.

295

Data Availability

Data from this work can be made available upon request: E. A. Marais for GEOS-Chem output, M. Belmonte-Rivas for KNMI OMI UT NO_2 , S. Choi and J. Joiner for NASA OMI UT NO_2 , and S. Beirle for LIS lightning properties.

300 Competing Interests

The authors declare that they have no conflicts of interest.

Author Contributions

EAM conducted model simulations, analysed and interpreted satellite, model, and aircraft data, and prepared the manuscript, DJJ provided supervisory guidance and assisted in the writing. SC, JJ, and MR-B retrieved the OMI UT NO_2 products, RCC aided in interpreting aircraft observations., LTM contributed LIS/OTD lightning flash observations, SB contributed lightning flash properties, LS, VS, and LJ contributed updated GEOS-Chem code.

305

Acknowledgements

This work was funded by the NASA Tropospheric Chemistry Program and a University of Birmingham Research Fellowship and NERC/EPSRC grant (EP/R513465/1) awarded to E. A. Marais. Model simulations were performed on the University of Birmingham's BlueBEAR High Performance Cluster. The authors would like to thank the BlueBEAR support team for IT and HPC support.

References

- Acarreta, J. R., De Haan, J. F., and Stammes, P.: Cloud pressure retrieval using the O₂-O₂ absorption band at 477 nm, *J. Geophys. Res.*, 109, doi:10.1029/2003jd003915, 2004.
- 315 Allen, D., Pickering, K., Duncan, B., and Damon, M.: Impact of lightning NO emissions on North American photochemistry as determined using the Global Modeling Initiative (GMI) model, *J. Geophys. Res.*, 115, doi:10.1029/2010jd014062, 2010.
- Baehr, J., Schlager, H., Ziereis, H., Stock, P., van Velthoven, P., Busen, R., Strom, J., and Schumann, U.: Aircraft observations of NO, NO_y, CO, and O₃ in the upper troposphere from 60°N to 60°S - Interhemispheric differences at midlatitudes, *Geophys. Res. Lett.*, 30, doi:10.1029/2003gl016935, 2003.
- 320 Barth, M. C., Cantrell, C. A., Brune, W. H., Rutledge, S. A., Crawford, J. H., Huntrieser, H., Carey, L. D., MacGorman, D., Weisman, M., Pickering, K. E., Bruning, E., Anderson, B., Apel, E., Biggerstaff, M., Campos, T., Campuzano-Jost, P., Cohen, R., Crouse, J., Day, D. A., Diskin, G., Flocke, F., Fried, A., Garland, C., Heikes, B., Honomichl, S., Hornbrook, R., Huey, L. G., Jimenez, J. L., Lang, T., Lichtenstern, M., Mikoviny, T., Nault, B., O'Sullivan, D., Pan, L. L., Peischl, J., Pollack, I., Richter, D., Riemer, D., Ryerson, T., Schlager, H., St Clair, J., Walega, J., Weibring, P., Weinheimer, A., Wennberg, P., Wisthaler, A., Wooldridge, P. J., and Ziegler, C.: The Deep Convective Clouds and Chemistry (DC3) Field Campaign, *B. Am. Meteorol. Soc.*, 96, 1281-1309, doi:10.1175/Bams-D-13-00290.1, 2015.
- 325 Beirle, S., Huntrieser, H., and Wagner, T.: Direct satellite observation of lightning-produced NO_x, *Atmos. Chem. Phys.*, 10, 10965-10986, doi:10.5194/acp-10-10965-2010, 2010.
- 330 Beirle, S., Koshak, W., Blakeslee, R., and Wagner, T.: Global patterns of lightning properties derived by OTD and LIS, *Nat Hazard Earth Sys*, 14, 2715-2726, doi:10.5194/nhess-14-2715-2014, 2014.
- Belmonte-Rivas, M. B., Veefkind, P., Boersma, F., Levelt, P., Eskes, H., and Gille, J.: Intercomparison of daytime stratospheric NO₂ satellite retrievals and model simulations, *Atmos. Meas. Tech.*, 7, 2203-2225, doi:10.5194/amt-7-2203-2014, 2014.
- Belmonte-Rivas, M. B., Veefkind, P., Eskes, H., and Levelt, P.: OMI tropospheric NO₂ profiles from cloud slicing: constraints on surface emissions, convective transport and lightning NO_x, *Atmos. Chem. Phys.*, 15, 13519-13553, doi:10.5194/acp-15-13519-2015, 2015.
- 335 Berkes, F., Houben, N., Bundke, U., Franke, H., Pätz, H.-W., Rohrer, F., Wahner, A., and Petzold, A.: The IAGOS NO_x Instrument – Design, operation and first results from deployment aboard passenger aircraft, *Atmos. Meas. Tech.*, doi:10.5194/amt-2017-435, 2017.
- 340 Bertram, T. H., Perring, A. E., Wooldridge, P. J., Crouse, J. D., Kwan, A. J., Wennberg, P. O., Scheuer, E., Dibb, J., Avery, M., Sachse, G., Vay, S. A., Crawford, J. H., McNaughton, C. S., Clarke, A., Pickering, K. E., Fuelberg, H., Huey, G., Blake, D. R., Singh, H. B., Hall, S. R., Shetter, R. E., Fried, A., Heikes, B. G., and Cohen, R. C.: Direct measurements of the convective recycling of the upper troposphere, *Science*, 315, 816-820, doi:10.1126/science.1134548, 2007.
- Boersma, K. F., Eskes, H. J., and Brinksma, E. J.: Error analysis for tropospheric NO₂ retrieval from space, *J. Geophys. Res.*, 109, doi:10.1029/2003jd003962, 2004.
- 345 Boersma, K. F., Eskes, H. J., Meijer, E. W., and Kelder, H. M.: Estimates of lightning NO_x production from GOME satellite observations, *Atmos. Chem. Phys.*, 5, 2311-2331, doi:10.5194/acp-5-2311-2005, 2005.
- Boersma, K. F., Eskes, H. J., Dirksen, R. J., van der A, R. J., Veefkind, J. P., Stammes, P., Huijnen, V., Kleipool, Q. L., Sneep, M., Claas, J., Leitao, J., Richter, A., Zhou, Y., and Brunner, D.: An improved tropospheric NO₂ column retrieval algorithm for the Ozone Monitoring Instrument, *Atmos. Meas. Tech.*, 4, 1905-1928, doi:10.5194/amt-4-1905-2011, 2011.
- 350 Bradshaw, J., Davis, D., Grodzinsky, G., Smyth, S., Newell, R., Sandholm, S., and Liu, S.: Observed distributions of nitrogen oxides in the remote free troposphere from the NASA global tropospheric experiment programs, *Rev. Geophys.*, 38, 61-116, doi:10.1029/1999rg900015, 2000.
- Browne, E. C., Perring, A. E., Wooldridge, P. J., Apel, E., Hall, S. R., Huey, L. G., Mao, J., Spencer, K. M., St Clair, J. M., Weinheimer, A. J., Wisthaler, A., and Cohen, R. C.: Global and regional effects of the photochemistry of CH₃O₂NO₂: evidence from ARCTAS, *Atmos. Chem. Phys.*, 11, 4209-4219, doi:10.5194/acp-11-4209-2011, 2011.
- 355 Brunner, D., Staehelin, J., Jeker, D., Wernli, H., and Schumann, U.: Nitrogen oxides and ozone in the tropopause region of the Northern Hemisphere: Measurements from commercial aircraft in 1995/1996 and 1997, *J. Geophys. Res.*, 106, 27673-27699, doi:10.1029/2001jd900239, 2001.

- Bucsela, E. J., Pickering, K. E., Huntemann, T. L., Cohen, R. C., Perring, A., Gleason, J. F., Blakeslee, R. J., Albrecht, R. I., Holzworth, R., Cipriani, J. P., Vargas-Navarro, D., Mora-Segura, I., Pacheco-Hernandez, A., and Laporte-Molina, S.: Lightning-generated NO_x seen by the Ozone Monitoring Instrument during NASA's Tropical Composition, Cloud and Climate Coupling Experiment (TC4), *J. Geophys. Res.*, 115, doi:10.1029/2009jd013118, 2010.
- Cecil, D. J., Buechler, D. E., and Blakeslee, R. J.: Gridded lightning climatology from TRMM-LIS and OTD: Dataset description, *Atmos. Res.*, 135, 404-414, doi:10.1016/j.atmosres.2012.06.028, 2014.
- 365 Choi, S., Joiner, J., Choi, Y., Duncan, B. N., Vasilkov, A., Krotkov, N., and Bucsela, E.: First estimates of global free-tropospheric NO₂ abundances derived using a cloud-slicing technique applied to satellite observations from the Aura Ozone Monitoring Instrument (OMI), *Atmos. Chem. Phys.*, 14, 10565-10588, doi:10.5194/acp-14-10565-2014, 2014.
- Crawford, J. H., Davis, D. D., Chen, G., Bradshaw, J., Sandholm, S., Kondo, Y., Merrill, J., Liu, S., Browell, E., Gregory, G., Anderson, B., Sachse, G., Barrick, J., Blake, D., Talbot, R., and Pueschel, R.: Implications of large scale shifts in tropospheric NO_x levels in the remote tropical Pacific, *J. Geophys. Res.*, 102, 28447-28468, doi:10.1029/97jd00011, 1997.
- 370 Dahlmann, K., Grewe, V., Ponater, M., and Matthes, S.: Quantifying the contributions of individual NO_x sources to the trend in ozone radiative forcing, *Atmos. Environ.*, 45, 2860-2868, doi:10.1016/j.atmosenv.2011.02.071, 2011.
- Day, D. A., Wooldridge, P. J., Dillon, M. B., Thornton, J. A., and Cohen, R. C.: A thermal dissociation laser-induced fluorescence instrument for in situ detection of NO₂, peroxy nitrates, alkyl nitrates, and HNO₃, *J. Geophys. Res.*, 107, doi:10.1029/2001jd000779, 2002.
- 375 Drummond, J. W., Ehhalt, D. H., and Volz, A.: Measurements of nitric oxide between 0-12 km altitude and 67°N to 60°S latitude obtained during STRATOZ III, *J. Geophys. Res.*, 93, 15831-15849, doi:10.1029/JD093iD12p15831, 1988.
- Ehhalt, D. H., Rohrer, F., and Wahner, A.: Sources and distribution of NO_x in the upper troposphere at northern midlatitudes, *J. Geophys. Res.*, 97, 3725-3738, doi:10.1029/91JD03081, 1992.
- 380 Finney, D. L., Doherty, R. M., Wild, O., Huntrieser, H., Pumphrey, H. C., and Blyth, A. M.: Using cloud ice flux to parametrise large-scale lightning, *Atmos. Chem. Phys.*, 14, 12665-12682, doi:10.5194/acp-14-12665-2014, 2014.
- Finney, D. L., Doherty, R. M., Wild, O., Young, P. J., and Butler, A.: Response of lightning NO_x emissions and ozone production to climate change: Insights from the Atmospheric Chemistry and Climate Model Intercomparison Project, *Geophys. Res. Lett.*, 43, 5492-5500, doi:10.1002/2016gl068825, 2016.
- 385 Finney, D. L., Doherty, R. M., Wild, O., Stevenson, D. S., MacKenzie, I. A., and Blyth, A. M.: A projected decrease in lightning under climate change, *Nat. Clim. Change*, 8, 210-213, doi:10.1038/s41558-018-0072-6, 2018.
- Grewe, V., Brunner, D., Dameris, M., Grenfell, J. L., Hein, R., Shindell, D., and Staehelin, J.: Origin and variability of upper tropospheric nitrogen oxides and ozone at northern mid-latitudes, *Atmos. Environ.*, 35, 3421-3433, doi:10.1016/S1352-2310(01)00134-0, 2001.
- 390 Hudman, R. C., Jacob, D. J., Turquety, S., Leibensperger, E. M., Murray, L. T., Wu, S., Gilliland, A. B., Avery, M., Bertram, T. H., Brune, W., Cohen, R. C., Dibb, J. E., Flocke, F. M., Fried, A., Holloway, J., Neuman, J. A., Orville, R., Perring, A., Ren, X., Sachse, G. W., Singh, H. B., Swanson, A., and Wooldridge, P. J.: Surface and lightning sources of nitrogen oxides over the United States: Magnitudes, chemical evolution, and outflow, *J. Geophys. Res.*, 112, doi:10.1029/2006jd007912, 2007.
- 395 Jacob, D. J., Heikes, B. G., Fan, S. M., Logan, J. A., Mauzerall, D. L., Bradshaw, J. D., Singh, H. B., Gregory, G. L., Talbot, R. W., Blake, D. R., and Sachse, G. W.: Origin of ozone and NO_x in the tropical troposphere: A photochemical analysis of aircraft observations over the South Atlantic basin, *J. Geophys. Res.*, 101, 24235-24250, doi:10.1029/96jd00336, 1996.
- Jacob, D. J., Crawford, J. H., Maring, H., Clarke, A. D., Dibb, J. E., Emmons, L. K., Ferrare, R. A., Hostetler, C. A., Russell, P. B., Singh, H. B., Thompson, A. M., Shaw, G. E., McCauley, E., Pederson, J. R., and Fisher, J. A.: The Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS) mission: design, execution, and first results, *Atmos. Chem. Phys.*, 10, 5191-5212, doi:10.5194/acp-10-5191-2010, 2010.
- 400 Jaeglé, L., Jacob, D. J., Wang, Y., Weinheimer, A. J., Ridley, B. A., Campos, T. L., Sachse, G. W., and Hagen, D. E.: Sources and chemistry of NO_x in the upper troposphere over the United States, *Geophys. Res. Lett.*, 25, 1705-1708, doi:10.1029/97gl03591, 1998.
- 405 Kenagy, H. S., Sparks, T. L., Ebben, C. J., Wooldridge, P. J., Lopez-Hilfiker, F., Lee, B. H., Thornton, J. A., McDuffe, E. E., Fibiger, D. L., Brown, S. S., Montzka, D. D., Weinheimer, A. J., Schroder, J. C., Campuzano-Jost, P., Day, D. A., Jimenez, J. L., Dibb, J. E., Campos, T., Shah, V., Jaegle, L., and Cohen, R. C.: NO_x Lifetime and NO_y Partitioning During WINTER, *J. Geophys. Res.*, 123, doi:10.1029/2018JD028736, 2018.
- 410 Krotkov, N. A., Lamsal, L. N., Celarier, E. A., Swartz, W. H., Marchenko, S. V., Bucsela, E. J., Chan, K. L., Wenig, M., and Zara, M.: The version 3 OMI NO₂ standard product, *Atmos. Meas. Tech.*, 10, 3133-3149, doi:10.5194/amt-10-3133-2017, 2017.
- Levelt, P. F., Van den Oord, G. H. J., Dobber, M. R., Malkki, A., Visser, H., de Vries, J., Stammes, P., Lundell, J. O. V., and Saari, H.: The Ozone Monitoring Instrument, *IEEE T. Geosci. Remote*, 44, 1093-1101, doi:10.1109/Tgrs.2006.872333, 2006.

- 415 Marchenko, S., Krotkov, N. A., Lamsal, L. N., Celarier, E. A., Swartz, W. H., and Bucsela, E. J.: Revising the slant column density retrieval of nitrogen dioxide observed by the Ozone Monitoring Instrument, *J. Geophys. Res.*, 120, 5670-5692, doi:10.1002/2014jd022913, 2015.
- Martin, R. V., Sauvage, B., Folkins, I., Sioris, C. E., Boone, C., Bernath, P., and Ziemke, J.: Space-based constraints on the production of nitric oxide by lightning, *J. Geophys. Res.*, 112, doi:10.1029/2006jd007831, 2007.
- 420 Miyazaki, K., Eskes, H. J., Sudo, K., and Zhang, C.: Global lightning NO_x production estimated by an assimilation of multiple satellite data sets, *Atmos. Chem. Phys.*, 14, 3277-3305, doi:10.5194/acp-14-3277-2014, 2014.
- Murray, L. T., Jacob, D. J., Logan, J. A., Hudman, R. C., and Koshak, W. J.: Optimized regional and interannual variability of lightning in a global chemical transport model constrained by LIS/OTD satellite data, *J. Geophys. Res.*, 117, doi:10.1029/2012jd017934, 2012.
- 425 Murray, L. T., Mickley, L. J., Kaplan, J. O., Sofen, E. D., Pfeiffer, M., and Alexander, B.: Factors controlling variability in the oxidative capacity of the troposphere since the Last Glacial Maximum, *Atmos. Chem. Phys.*, 14, 3589-3622, doi:10.5194/acp-14-3589-2014, 2014.
- Murray, L. T.: Lightning NO_x and Impacts on Air Quality, *Current Pollution Reports*, 2, 115-133, doi:10.1007/s40726-016-0031-7, 2016.
- 430 Nault, B. A., Garland, C., Wooldridge, P. J., Brune, W. H., Campuzano-Jost, P., Crouse, J. D., Day, D. A., Dibb, J., Hall, S. R., Huey, L. G., Jimenez, J. L., Liu, X. X., Mao, J. Q., Mikoviny, T., Peischl, J., Pollack, I. B., Ren, X. R., Ryerson, T. B., Scheuer, E., Ullmann, K., Wennberg, P. O., Wisthaler, A., Zhang, L., and Cohen, R. C.: Observational Constraints on the Oxidation of NO_x in the Upper Troposphere, *J. Phys. Chem. A*, 120, 1468-1478, doi:10.1021/acs.jpca.5b07824, 2016.
- Ott, L. E., Pickering, K. E., Stenchikov, G. L., Allen, D. J., DeCaria, A. J., Ridley, B., Lin, R. F., Lang, S., and Tao, W. K.: Production of lightning NO_x and its vertical distribution calculated from three-dimensional cloud-scale chemical transport model simulations, *J. Geophys. Res.*, 115, doi:10.1029/2009jd011880, 2010.
- 435 Pickering, K. E., Bucsela, E., Allen, D., Ring, A., Holzworth, R., and Krotkov, N.: Estimates of lightning NO_x production based on OMI NO₂ observations over the Gulf of Mexico, *J. Geophys. Res.*, 121, 8668-8691, doi:10.1002/2015jd024179, 2016.
- Prather, M. J., and Jacob, D. J.: A persistent imbalance in HO_x and NO_x photochemistry of the upper troposphere driven by deep tropical convection, *Geophys. Res. Lett.*, 24, 3189-3192, doi:10.1029/97gl03027, 1997.
- 440 Price, C., and Rind, D.: A simple lightning parameterization for calculating global lightning distributions, *J. Geophys. Res.*, 97, 9919-9933, doi:10.1029/92JD00719, 1992.
- Price, C., and Rind, D.: What determines the cloud-to-ground lightning fraction in thunderstorms, *Geophys. Res. Lett.*, 20, 463-466, doi:10.1029/93gl00226, 1993.
- 445 Price, C., and Rind, D.: Modeling global lightning distributions in a General-Circulation Model, *Mon. Weather Rev.*, 122, 1930-1939, doi:10.1175/1520-0493(1994)122<1930:Mgldia>2.0.Co;2, 1994.
- Rap, A., Richards, N. A. D., Forster, P. M., Monks, S. A., Arnold, S. R., and Chipperfield, M. P.: Satellite constraint on the tropospheric ozone radiative effect, *Geophys. Res. Lett.*, 42, 5074-5081, doi:10.1002/2015gl064037, 2015.
- Reed, C., Evans, M. J., Di Carlo, P., Lee, J. D., and Carpenter, L. J.: Interferences in photolytic NO₂ measurements: explanation for an apparent missing oxidant?, *Atmos. Chem. Phys.*, 16, 4707-4724, doi:10.5194/acp-16-4707-2016, 2016.
- 450 Sauvage, B., Martin, R. V., van Donkelaar, A., Liu, X., Chance, K., Jaeglé, L., Palmer, P. I., Wu, S., and Fu, T. M.: Remote sensed and in situ constraints on processes affecting tropical tropospheric ozone, *Atmos. Chem. Phys.*, 7, 815-838, doi:10.5194/acp-7-815-2007, 2007.
- Schumann, U., and Huntrieser, H.: The global lightning-induced nitrogen oxides source, *Atmos. Chem. Phys.*, 7, 3823-3907, doi:10.5194/acp-7-3823-2007, 2007.
- 455 Silvern, R. F., Jacob, D. J., Travis, K. R., Sherwen, T., Evans, M. J., Cohen, R. C., Laughner, J. L., Hall, S. R., Ullmann, K., Crouse, J. D., Wennberg, P. O., Peischl, J., and Pollack, I. B.: Observed NO/NO₂ ratios in the upper troposphere imply errors in NO-NO₂-O₃ cycling kinetics or an unaccounted NO_x reservoir." *Geophysical Research Letters* 45: 4466-4474.
- Singh, H. B., Brune, W. H., Crawford, J. H., Jacob, D. J., and Russell, P. B.: Overview of the summer 2004 intercontinental chemical transport experiment - North America (INTEX-A), *J. Geophys. Res.*, 111, doi:10.1029/2006jd007905, 2006.
- 460 Singh, H. B., Brune, W. H., Crawford, J. H., Flocke, F., and Jacob, D. J.: Chemistry and transport of pollution over the Gulf of Mexico and the Pacific: spring 2006 INTEX-B campaign overview and first results, *Atmos. Chem. Phys.*, 9, 2301-2318, doi:10.5194/acp-9-2301-2009, 2009.
- Stettler, M. E. J., Eastham, S., and Barrett, S. R. H.: Air quality and public health impacts of UK airports. Part I: Emissions, *Atmos. Environ.*, 45, 5415-5424, doi:10.1016/j.atmosenv.2011.07.012, 2011.
- 465 Stratmann, G., Ziereis, H., Stock, P., Brenninkmeijer, C. A. M., Zahn, A., Rauthe-Schoch, A., Velthoven, P. V., Schlager, H., and Volz-Thomas, A.: NO and NO_y in the upper troposphere: Nine years of CARIBIC measurements onboard a passenger aircraft, *Atmos. Environ.*, 133, 93-111, doi:10.1016/j.atmosenv.2016.02.035, 2016.
- Toon, O. B., Maring, H., Dibb, J., Ferrare, R., Jacob, D. J., Jensen, E. J., Luo, Z. J., Mace, G. G., Pan, L. L., Pfister, L., Rosenlof, K. H., Redemann, J., Reid, J. S., Singh, H. B., Thompson, A. M., Yokelson, R., Minnis, P., Chen, G., Jucks, K. W., and
- 470

- Pszenny, A.: Planning, implementation, and scientific goals of the Studies of Emissions and Atmospheric Composition, Clouds and Climate Coupling by Regional Surveys (SEAC⁴RS) field mission, *J. Geophys. Res.*, 121, 4967-5009, doi:10.1002/2015jd024297, 2016.
- 475 Tost, H., Jockel, P. J., and Lelieveld, J.: Lightning and convection parameterisations - uncertainties in global modelling, *Atmos. Chem. Phys.*, 7, 4553-4568, doi:10.5194/acp-7-4553-2007, 2007.
- Travis, K. R., Jacob, D. J., Fisher, J. A., Kim, P. S., Marais, E. A., Zhu, L., Yu, K., Miller, C. C., Yantosca, R. M., Sulprizio, M. P., Thompson, A. M., Wennberg, P. O., Crouse, J. D., St Clair, J. M., Cohen, R. C., Laughner, J. L., Dibb, J. E., Hall, S. R., Ullmann, K., Wolfe, G. M., Pollack, I. B., Peischl, J., Neuman, J. A., and Zhou, X. L.: Why do models overestimate surface ozone in the Southeast United States?, *Atmos. Chem. Phys.*, 16, 13561-13577, doi:10.5194/acp-16-13561-2016, 480 2016.
- van Geffen, J. H. G. M., Boersma, K. F., Van Roozendaal, M., Hendrick, F., Mahieu, E., De Smedt, I., Snee, M., and Veefkind, J. P.: Improved spectral fitting of nitrogen dioxide from OMI in the 405-465 nm window, *Atmos. Meas. Tech.*, 8, 1685-1699, doi:10.5194/amt-8-1685-2015, 2015.
- 485 Worden, H. M., Bowman, K. W., Kulawik, S. S., and Aghedo, A. M.: Sensitivity of outgoing longwave radiative flux to the global vertical distribution of ozone characterized by instantaneous radiative kernels from Aura-TES, *J. Geophys. Res.*, 116, doi:10.1029/2010jd015101, 2011.
- Ziemke, J. R., Chandra, S., and Bhartia, P. K.: "Cloud slicing": A new technique to derive upper tropospheric ozone from satellite measurements, *J. Geophys. Res.*, 106, 9853-9867, doi:10.1029/2000jd900768, 2001.