



Mediterranean Precipitation Response to Greenhouse Gases and Aerosols

Tao Tang¹, Drew Shindell¹, Bjørn H. Samset², Olivier Boucher³, Piers M. Forster⁴, Øivind Hodnebrog², Gunnar Myhre², Jana Sillmann², Apostolos Voulgarakis⁵, Timothy Andrews⁶,
5 Gregory Faluvegi⁷, Dagmar Fläschner⁸, Trond Iversen⁹, Matthew Kasoar⁵, Viatcheslav Kharin¹⁰,
Alf Kirkevåg⁹, Jean-Francois Lamarque¹¹, Dirk Olivie⁹, Thomas Richardson⁴, Camilla W. Stjern²,
Toshihiko Takemura¹²

¹Nicholas School of the Environment, Duke University, Durham, USA

10 ²CICERO Center for International Climate and Environmental Research – Oslo, Norway

³Institute Pierre-Simon Laplace, Université Pierre et Marie Curie / CNRS, Paris, France

⁴University of Leeds, Leeds, United Kingdom

⁵Imperial College London, London, United Kingdom

⁶Met Office Hadley Centre, Exeter, UK

15 ⁷Columbia University, New York, USA and NASA Goddard Institute for Space Studies, New York, USA

⁸Max-Planck-Institut für Meteorologie, Hamburg, Germany

⁹Norwegian Meteorological Institute, Oslo, Norway

¹⁰Canadian Centre for Climate Modelling and Analysis, Victoria, BC, Canada

¹¹National Center for Atmospheric Research, Boulder, USA

20 ¹²Kyushu University, Fukuoka, Japan

Correspondence to: Tao Tang (tao.tang@duke.edu)

Abstract. Atmospheric aerosols and greenhouse gases affect cloud properties, radiative balance and thus, the hydrological cycle. Observations show that precipitation has decreased in the Mediterranean since the 20th century,
25 and many studies have investigated possible mechanisms. So far, however, the effects of aerosol forcing on
Mediterranean precipitation remain largely unknown. Here we compare Mediterranean precipitation responses to
individual forcing agents in a set of state-of-the-art global climate models (GCMs). Our analyses show that both
greenhouse gases and aerosols can cause drying in the Mediterranean, and that precipitation is more sensitive to black
carbon (BC) forcing than to well-mixed greenhouse gases (WMGHGs) or sulfate aerosol. In addition to local heating,
30 BC appears to reduce precipitation by causing an enhanced positive North Atlantic Oscillation (NAO)/Arctic
Oscillation (AO)-like sea level pressure (SLP) pattern, characterized by higher SLP at mid-latitudes and lower SLP at
high-latitudes. WMGHGs cause a similar SLP change, and both are associated with a northward diversion of the jet
stream and storm tracks, reducing precipitation in the Mediterranean while increasing precipitation in Northern
Europe. Though the applied forcings were much larger, if forcings are scaled to those of the historical period of 1901-
35 2010, roughly one-third ($31 \pm 17\%$) of the precipitation decrease would be attributable to global BC forcing with the
remainder largely attributable to WMGHGs whereas global scattering sulfate aerosols have negligible impacts. The
results from this study suggest that future BC emissions may significantly affect regional water resources, agricultural
practices, ecosystems, and the economy in the Mediterranean region.



1 Introduction

Aerosols, fine particles in the atmosphere produced by both natural processes and anthropogenic activities, impact the Earth's climate by scattering and absorbing solar radiation (direct effect), or by modifying the properties of clouds (indirect effects) through a variety of mechanisms including atmospheric heating and changes in ice nuclei and cloud condensation nuclei (CCN), including their size, location and concentration. These changes may significantly affect solar radiation and precipitation (*Ramanathan et al.*, 2001; *Kaufman et al.*, 2002; *Shindell et al.*, 2012; *Bond et al.*, 2013; *Boucher et al.*, 2013). The effects of aerosols on climate have been widely studied both on global and regional scales (*Ramanathan and Carmichael*, 2008; *Shindell and Faluvegi*, 2009). For example, *Menon et al.* (2002) reported slight cooling and drying trends in the northern part of China in the 2nd half of the 20th century, and attributed such trends to the emissions of BC aerosols based on model simulations. Similarly, *Hodnebrog et al.* (2016) reported a precipitation decrease in southern Africa due to local biomass burning aerosols based on analyses of model simulations and local energy budget. On the other hand, *Koren et al.* (2012) argued that aerosols could intensify rainfall events in the lower and mid-latitudes by analyzing satellite observations. However, *Stevens and Feingold* (2009) contended that the effects of aerosols on clouds and precipitation are very limited due to the buffering effects of the climate system itself. In addition to their influence on temperature and precipitation, aerosols may also affect large-scale atmospheric circulation. *Takahashi and Watanabe* (2016), based on a series of model simulations, suggested that the Pacific trade winds were accelerated partially by sulfate aerosols during the past two decades. *Dunstone et al.* (2013) reported that aerosols could modulate Atlantic tropical storm frequency due to aerosol-induced shifts in the Hadley circulation. These differing results suggest that aerosol effects on regional climate may depend on the aerosol types, seasons, and regions of interest.

A decreasing precipitation trend in the Mediterranean area since the 20th century has been reported and its possible causes have been investigated in many studies (*Piervitali et al.*, 1998; *Buffoni et al.*, 1999; *Mariotti et al.*, 2002; *Düneloh and Jacobeit*, 2003; *Xoplaki et al.*, 2004). For instance, *Quadrelli et al.* (2001) observed a strong correlation between winter Mediterranean precipitation and the NAO (*Hurrell et al.*, 2001). *Krichak and Alpert* (2005) suggested that the East Atlantic-West Russia (EA-WR) pattern may also play an important role in modulating the precipitation in the Mediterranean. Hence, the responses of Mediterranean precipitation to these large-scale variability patterns (e.g., NAO, EA-WR), and to some extent how these patterns might be responding to external drivers, are fairly well-understood (*Black et al.*, 2010). However, prior studies included all the drivers at once, so cannot discern the relative roles of WMGHGs and other agents. Anthropogenic aerosols have been reported to greatly influence the temperature in the Mediterranean (*Nabat et al.*, 2014), but the effects of aerosols on Mediterranean precipitation have not been carefully examined. Since precipitation impacts water availability for both ecosystems and human societies, it is crucial to understand the different impacts of the climate drivers that are responsible for the Mediterranean precipitation trend. To bridge this knowledge gap, here we analyze Mediterranean precipitation changes based on a group of state-of-the-art GCMs that examined the precipitation response to individual climate drivers, which could help inform management of water resources, regional societal activities such as agriculture, and even emissions mitigation.



2 Data and Method

2.1 Data

This study employed output from nine models participating in the Precipitation Driver and Response Model Intercomparison Project (PDRMIP), utilizing simulations examining the individual responses to CO₂, sulfate and BC aerosols. In these simulations, perturbations were performed with each model at global scale: a doubling of CO₂ concentration (CO₂×2), 10 times present-day BC concentration (BC×10), and 5 times present-day SO₄ concentration (SO₄×5). All perturbations were abrupt. CO₂×2 perturbations were applied relative to the models' own baseline values. For aerosol perturbations, monthly present-day concentrations were derived from the AeroCom Phase II initiative (Myhre *et al.*, 2013a). The concentrations were multiplied by the stated factors. A few models instead perturbed aerosol emissions, in most cases again using AeroCom Phase II data. Each perturbation was run in two configurations, a 15-yr fixed sea surface temperature (SST) simulation and a 100-yr coupled simulation. Each fixed-SST simulation is compared with its fixed-SST control simulation to diagnose the effective radiative forcing (ERF) due to each perturbation (Myhre *et al.*, 2013b), whereas each coupled run is compared with its coupled control run to examine climate response. The nine models used in this study are listed in Table 1. A more detailed description of PDRMIP and its initial findings are given in Samset *et al.* (2016) and Myhre *et al.* (2017).

2.2 Method

In addition to direct analysis of meteorological fields (e.g. precipitation, sea-level pressure) in the models, we also analyse the energy budget associated with the hydrological cycle. Following Hodnebrog *et al.* (2016) and Muller and O'Gorman (2011), the precipitation change is related to diabatic cooling and the horizontal transport of dry static energy as follows:

$$L_c \Delta P = \Delta Q + \Delta H \quad (1)$$

Here L_c is the latent heat of condensation of water vapor, which is 29 W m⁻² mm⁻¹ day. ΔP is the precipitation change. ΔH is the column-integrated dry static energy flux divergence and ΔQ is the column-integrated diabatic cooling, which is calculated as:

$$\Delta Q = \Delta LW + \Delta SW - \Delta SH \quad (2)$$

where ΔLW is the change of longwave radiation in the atmospheric column and ΔSW is the change of shortwave radiation in the atmospheric column. ΔSH is the change of upward sensible heat flux.

Since most of the precipitation events occur in the wet season (Oct-Mar) in the Mediterranean, roughly 70% of total annual precipitation (Mariotti *et al.*, 2002; Kottek *et al.*, 2006), the analysis was restricted to the wet season in the current study unless noted otherwise. All of the data used in this study were re-gridded into 2.5°×2.5° horizontal resolution for analyses.



3 Results

Fig. 1 shows the multi-model mean (MMM) of normalized ΔP for each forcing. Both CO₂ and BC caused a substantial drying over Mediterranean, with a larger magnitude from BC (Fig. 1a-b) whereas SO₄ contributed very little in the Mediterranean region (Fig. 1c). Moreover, in stark contrast to the drying of the Mediterranean, Northern Europe shows increasing precipitation trends for CO₂ and BC (Fig. 1a-b), which will be discussed in more details later. To compare the precipitation response quantitatively, the domain-averaged (purple rectangle in Fig. 1, 30°N-45°N, 10°W-40°E) trends are shown in Fig. 2. For CO₂, all of the nine models show drying trends (Fig. 2a). The MMM is -0.03 ± 0.03 mm/day per W/m², with individual model values ranging from -0.01 to -0.06 mm/day per W/m². For BC (Fig. 2b), all nine models again show drying trends, with the MMM value -0.12 ± 0.07 mm/day per W/m², which is four times as large as that of CO₂. When it comes to SO₄ (Fig. 2c), the model results even differ in the sign of change and the MMM value is small (-0.01 ± 0.04 mm/day per W/m²). These analyses show that the precipitation response is more sensitive to BC forcing than to CO₂ and SO₄ in this region.

In order to investigate the mechanisms governing the precipitation response, we performed an energy budget analysis for this region (Fig. 3). For CO₂, the drying is dominated by horizontal energy transport (gray box in the CO₂ panel), albeit some offset by diabatic cooling (pink box in the CO₂ panel). For BC, the net radiation change, which is primarily SW (red box in the BC panel), has a larger impact than the horizontal energy change (gray box in the BC panel), but the latter is nonetheless a substantial fraction of the net change. When it comes to SO₄, the small precipitation response results from the offsetting of net radiation change (pink box in the SO₄ panel) and horizontal energy transport (gray box in the SO₄ panel). The energy budget analysis implies that the dynamical responses to CO₂ and BC played a crucial role in modulating the precipitation in this region.

We then analyzed the ΔSLP from the model output (Fig. 4). Specifically, it is seen that CO₂ induced strong SLP changes. The SLP increased at mid-latitudes, with increases extending from the North Atlantic to Southern Europe, and decreased at high latitudes (Fig. 4a). BC led to a similar pattern of SLP change, but with increased magnitude (Fig. 4b), characterized by two increases centered in Europe and the Western North Atlantic. Compared with CO₂ and BC, SO₄ caused an opposite change (Fig. 4c). The CO₂ and BC forcings appear to induce a pattern similar to the positive phase of the NAO /AO (Lorenz, 1951), in which the jet streams and storm tracks are displaced northward, leading to a drier Mediterranean and precipitation increases in Northern Europe (Fig. 1a-b). Such a shift in response to forcings is more clearly seen in the changes of zonal winds (Fig. 5). The CO₂ caused a strengthening of zonal winds in the whole upper atmosphere and a strengthening around 60°N from the near-surface to the top of the atmosphere, as well as weakening around 30°N from the near-surface to the mid-troposphere (Fig. 5a), as in prior studies (Shindell *et al.*, 2001) The strengthening around 60°N is more apparent for BC (Fig. 5b). Similar results were seen in response to aerosol forcing in a prior study (Allen and Sherwood, 2011). This shift is possibly due to the enhancement of the tropospheric temperature gradient between mid-latitudes and high-latitudes, as suggested by Allen *et al.* (2012).



Our analyses illustrate that BC aerosols may modulate regional precipitation in part via modifying large-scale circulation patterns. Many previous studies suggest that BC could impact regional precipitation by changing the local vertical temperature profile, in which BC aerosols absorb solar radiation and heat the atmosphere, thus suppressing convection and cloud formation (Kaufman *et al.*, 2002; Meehl *et al.*, 2008; Ramanathan and Carmichael, 2008; 5 Hodnebrog *et al.*, 2016). Our results (analyses of the energy budget, SLP and zonal winds) suggest that a portion of the drying is also associated with large-scale circulation responses. In addition, our pattern of jet stream/storm track changes (Fig. 4 & 5) is also in agreement with the projections from the latest IPCC report (Collins *et al.*, 2013) based on a set of CMIP5 models, with increasing storm activities in Northern Europe and decreasing storms in the Mediterranean. Such a shift of storm tracks may further reduce the precipitation in the Mediterranean, though 10 reductions in WMGHG or BC emissions may help to mitigate the projected drying.

4 Case Study – Historical Observations and Scaled Model Results

The above analyses demonstrated how the precipitation and circulation responded to each forcing both qualitatively and quantitatively. In order to explore their potential relative contributions to the total precipitation change, here we apply linear scaling to the model output. Since PDRMIP utilized large aerosol and greenhouse gas changes in order 15 to achieve strong signals that could be statistically significant with a relatively modest amount of computational time, the precipitation change from those model outputs needs to be scaled in order to compare with observations. Uncertainties related to this approach are discussed further in section 5.

In this study, we focus on the period from 1901 to 2010. The scaled precipitation change for each individual forcing 20 is defined as:

$$\Delta P_{scaled} = \Delta P \times (ERF_{1901-2010} / ERF_{model}) \quad (3)$$

In equation (3), ΔP is the precipitation change over Mediterranean in the model during the last 50 years in the coupled 25 run, since the model has reached near-equilibrium state after 30 years. $ERF_{1901-2010}$ is the historical global ERF for the period of 1901-2010. The values were obtained from the latest Intergovernmental Panel on Climate Change (IPCC) assessment report (Myhre *et al.*, 2013b). The $ERF_{1901-2010}$ value used for CO₂ is 2.33 W/m², which is larger than the CO₂ value from the IPCC report as CO₂ was used to represent all WMGHGs in this case study. $ERF_{1901-2010}$ values for BC and SO₄ are 0.28 W/m² and -0.33 W/m², respectively. ERF_{model} is the global ERF in the PDRMIP models, which 30 was obtained by calculating the energy flux change at the top of the atmosphere from years 6 to 15 of the fixed-SST simulations, since present models largely equilibrate within 5 years of fixed-SST running (Kvalevåg *et al.*, 2013). In addition to the direct effects of the aerosols, the indirect effects of aerosols were also included in most of the models and thus, in the ERF_{model} . The value of $(ERF_{1901-2010} / ERF_{model})$ is the scaling factor applied to model precipitation output to match historical forcing levels. They are 0.64 [0.57, 0.69], 0.33 [0.10, 0.68] and 0.11 [0.04, 0.16] for CO₂, 35 BC and SO₄, respectively (where the values indicate the mean [min, max] across the nine models). An important assumption here is that the ΔP changes linearly with ERF_{model} .



ΔP_{scaled} is calculated for CO₂, BC, and SO₄ separately. The total ΔP_{scaled} is the combination of the three, assuming their responses to those forcings can be added linearly. It should be noted that in this analysis, we use the combined responses to WMGHGs, BC and SO₄ to approximate the total historical response over 1901-2010. Several additional factors may have also played a role, including natural forcing (solar and volcanic activities), land use/land cover change, contrails, ozone (O₃) (both tropospheric and stratospheric) and stratospheric water vapor, which have forcings of -0.03, -0.09, 0.05, 0.26 and 0.06 W/m², respectively (Myhre *et al.*, 2013b). As all these forcings are fairly small, simulations to isolate their impacts would be extremely computationally expensive and hence were not performed but to first order we expect their exclusion is unlikely to greatly affect our results. Characterization of the influence of these other drivers merits future study, particularly as some operate via different physical processes (e.g. tropospheric ozone is both a greenhouse gas and an absorber of incoming solar radiation). Similar analyses were also performed to obtain scaled SLP change (ΔSLP_{scaled}), zonal wind change and energy budget change in the atmospheric column.

Several observational and reanalysis datasets were also employed in this part of our study. For precipitation, Global Precipitation Climatology Center (GPCC) monthly precipitation data (Schneider *et al.*, 2011), provided by NOAA/OAR/ESRL from their website (<https://www.esrl.noaa.gov/psd/data/gridded/data.gpc.html#detail>), is employed. It is a high quality gridded dataset that is mainly terrestrial station-derived. For SLP, we use HadSLP2 data (Allan and Ansell, 2006), which is created by combining marine observations from ICOADS data (Worley *et al.*, 2005) and land (terrestrial and island) observations (available at <https://www.esrl.noaa.gov/psd/data/gridded/data.hadslp2.html>). We also use NCEP/NCAR reanalysis data (Kalnay *et al.*, 1996), downloaded from <https://www.esrl.noaa.gov/psd/data/gridded/data.ncep.reanalysis.derived.surface.html>, for the comparisons of zonal wind. All these datasets have undergone rigorous quality control and have been widely used in the climate community, including the IPCC 2013 assessment report (Hartmann *et al.*, 2013). The trends of the observed and reanalysis data were estimated by a simple linear regression applied to the same period of the datasets.

The combination of WMGHGs, BC and SO₄ exerted a strong drying trend in the Mediterranean (Fig. 6a). The drying trends shown here are statistically significant and consistent with the observations (Fig. 6b), as well as previous studies (Buffoni *et al.*, 1999; Mariotti *et al.*, 2002; Dünkeloh and Jacobeit, 2003). When averaged over the whole domain, the scaled drying trends caused by WMGHGs, BC and SO₄ are -1.28 ± 1.21 mm/decade, -0.58 ± 0.34 mm/decade and -0.03 ± 0.21 mm/decade, respectively (not shown here). When combined (Fig. 6c), all nine models show decreased precipitation, with MMM value of -1.89 ± 1.39 mm/decade, which is roughly a 5% decrease relative to the climatology of the control simulations. Such a decreasing trend is indistinguishable from the observations (-2.78 ± 1.10 mm/decade, a 10% decrease compared with its 110-yr climatology). In spite of the dominant role of WMGHGs in the drying of the Mediterranean, BC contributed roughly one-third ($31 \pm 17\%$) of the total forced precipitation decrease in this region whereas the contribution of the scattering aerosol-SO₄ is negligible ($\sim 1.6\%$). We also examined the trend of precipitation in the control simulations and found only very weak responses (Fig. 6c), with a mean value of 0.004 ± 0.03 mm/decade and maximum value of 0.03 mm/decade in any individual model. Since current GCMs are able to capture



the broad spatial and temporal features of internal variability (Flato *et al.*, 2013), and the forced drying signal is almost equal to the total signal (Fig. 6a-c), the consistent drying trend in the models is very unlikely to be attributable to unforced variability and appears realistic. The energy budget change (Fig. 6d) clearly shows that the net precipitation decrease is mainly due to horizontal energy transport (gray box) rather than diabatic cooling (pink box), because the absorption of SW radiation (red box) and LW radiative cooling (green box) offset one another in total.

Fig. 7a shows the overall response of SLP to these forcings, with strong SLP increases at mid-latitudes and strong decreases at higher latitudes. Such patterns of SLP changes are also found in the observed datasets (Fig. 7b), albeit with a larger magnitude. The combined pattern of zonal wind responses shows winds intensified at the northern edge of the jet stream and weakened at the southern edge (Fig. 7c). The NCEP dataset depicts a similar pattern of changes, with winds intensifying at 60°N and weakening at 30°N, but as with SLP, with a stronger magnitude (Fig. 7d). Some previous studies have pointed out that current GCMs significantly underestimate the tropical expansion and jet stream shift, which could be related to the short observational record, large internal variability or model deficiencies (Johanson and Fu, 2009; Allen *et al.*, 2012). Despite the underestimations, our analyses clearly demonstrate the shift of the jet stream in response to these forcings that appears qualitatively consistent with observations.

Based on the model simulations in the current study, the pattern of climate response to BC forcing over the past ~110 years is similar to the response to WMGHGs over Europe and the North Atlantic, including precipitation, SLP and zonal winds. At the same time, our results suggest that SO₄ played a very limited role in modulating Mediterranean precipitation trends and North Atlantic storm tracks. In other words, the precipitation trends during the past 110 years in the Mediterranean are likely to be only weakly sensitive to scattering aerosols that were not modeled (e.g., organic carbon, nitrate) or the uncertainties in aerosol negative forcing (probably not even for indirect forcing, as they were included in sulfate simulations for most models). The small sensitivity of SO₄ is likely due to compensation between local and remote effects (Liu *et al.*, 2018). Combined with its small *ERF*, the role of SO₄ appears to be negligible during this period. However, the simulations examined here were not designed to determine whether the aerosol effects are due to local or remote emissions from the models. Initial analysis from PDRMIP regional experiments (in which BC over Asia only is multiplied by 10, with everything else being held at present-day levels) indicate that BC from Asia contributes as much as 60% to the drying signal in the Mediterranean, and in fact a larger average rainfall change in the Mediterranean than averaged over Asia itself. This suggests that the remote effects of BC may have dominated the Mediterranean precipitation changes. Hence the response to global BC increases may be a reasonable proxy for the 20th century changes, although it would be useful to explore the effects of local reductions from Europe itself in the late 20th century. The relative impacts of local versus remote forcing will be further explored in forthcoming PDRMIP analyses.

5 Discussion and Conclusion

Since PDRMIP experiments are equilibrium simulations while the real-world is transient, and we scaled PDRMIP forcing to match historical levels, we examined related experiments to test both these aspects of the methodology used



in our comparison with historical observations. Historical GHG-only simulations using the same CMIP5 models (Taylor *et al.*, 2012) that participated in the PDRMIP project were collected and analyzed (data available at <http://strega.ldeo.columbia.edu:81/CMIP5/monthly/byModel/>). Six models are available and each model has 1-5 ensemble members. All of the six models show drying trends (Fig. 8), with a MMM value of -1.32 ± 1.65 mm/decade
5 (-1.29 mm/decade when weighted by ensemble size) which is quite close to the WMGHGs result of our scaled equilibrium PDRMIP output (-1.28 ± 1.21 mm/decade). In fact, the overlap of their probability density functions is 0.85, assuming a normal distribution. This comparison indicates that our methodology does not appear to be a large source of uncertainty in the current study, though response to other agents may not be as linear as those to WMGHGs (unfortunately, simulations are not currently available to evaluate other forcings, and, given the enormous expense of running enough ensemble members to isolate the relatively small signals for individual aerosols, are unlikely to be
10 anytime soon). Similar analyses were also performed for SLP and zonal winds, and again there is no appreciable difference between the historical transients and the scaled equilibrium responses. The consistent results suggest that the methodology works surprisingly well.

15 In addition to the wet season, precipitation during the dry season (Apr-Sep) for the PDRMIP model was also analyzed. The modelled dry season precipitation trends, however, do not match the observations well (not shown). The modelled results also show a statistically strong drying trend while the observations do not show significant changes. Two possible reasons may be responsible for the apparent discrepancies. One is that only 30% of the total precipitation occurs during the dry season (boreal summer months) and it is difficult to simulate the uneven distribution of
20 infrequent rainfall events. The other is that there are large uncertainties in the observational data itself. Unlike the wet season, in which nearly half of the grid boxes show statistically significant trends (Fig. 6b), almost none of the grid boxes show statistically significant trends in the dry season, undermining the robustness of the observational results.

The drying influence of WMGHGs will be more prominent in the future due to their projected continued growth. In
25 contrast, many studies suggest that aerosol concentrations may decrease rapidly in the future due to air quality and climate policies along with their relatively short lifetime compared with WMGHGs (Andreae *et al.*, 2005; Myhre *et al.*, 2013b; Shindell *et al.*, 2013). Reductions of BC could, to some extent, slow down the drying trend in the Mediterranean. Overall, a drier Mediterranean region is expected owing to increasing WMGHGs, but the pace of change in global BC emissions may substantially modify the drying rate in the near term.

30 Some limitations and uncertainties still exist in our current study. First, it is important to keep in mind that the case study in Section 4 is not a formal attribution analysis, despite the estimation of BC contribution. Our aim is to give a first grasp of the effects of aerosol on regional precipitation in the Mediterranean. Second, although our comparison of scaled equilibrium and unscaled transient simulations indicates that our methodology works well at least for
35 WMGHGs, there is no systematic study so far exploring the linearity (or non-linearity) of the precipitation responses to BC or the linearity of responses to multiple versus individual forcings. Third is that the $ERF_{1901-2010}$ of BC represents direct effects only (Myhre *et al.*, 2013b). Semidirect and indirect effects, however, are included in many of our



PDRMIP models, and thus in ERF_{model} . We did not include these effects in the scaling in this study for two reasons: first, the indirect effects of BC in the PDRMIP models do not include ice particles and are difficult to evaluate as BC concentrations were prescribed so that they cannot interact fully with clouds, indicating that they are not fully resolved, and second, the net $ERF_{1901-2010}$ of semidirect plus indirect effects is likely small (-0.1 to $+0.2$ W/m^2) with a very large overall uncertainty range (-0.4 to $+0.9$ W/m^2) (Bond *et al.*, 2013). If the semi-direct and indirect effects of BC (-0.1 to $+0.2$ W/m^2) are considered in the scaling, the ΔP_{scaled} of BC aerosol would be -0.44 to -0.87 mm/decade and still contribute a substantial part (25 to 40 %) to the drying. The situation is similar for sulfate aerosol, for which indirect effects are included in ERF_{model} , but not in $ERF_{1901-2010}$. We did not include indirect effects in our scaling as these were not attributed to individual aerosol species in the IPCC AR5 (Boucher *et al.*, 2013). If the indirect effects are considered, the negative $ERF_{1901-2010}$ could increase roughly by a factor of two (assuming indirect effects are largely associated with sulfate). However, the ΔP_{scaled} of sulfate aerosol would still be very small compared with WMGHGs or BC, which would not impact our conclusions. The final issue is related to the design of the model simulations. The perturbations are $5\times$ or $10\times$ present-day aerosol concentrations, which are time-invariant. The aerosols, however, have significant spatial and temporal variations. For instance, aerosol concentrations have been increasing in Asia continuously since 1950, but decreasing in Europe since the 1970s (Allen *et al.*, 2013). As noted, further work is needed to determine how much of the Mediterranean trends result from local relative to remote forcing. To the extent that the trends are driven by remote forcing the potential influence of such spatio-temporal variations will be small. This will be explored in future PDRMIP simulations.

Our analyses show that both WMGHGs and BC influence wet season Mediterranean rainfall by causing an enhanced positive NAO/AO-like SLP pattern as well as by some local heating due to SW absorption. The SLP pattern is characterized by higher SLP in the North Atlantic and Mediterranean and lower SLP in the Northern part of Europe, which diverts the jet stream and storm tracks further northward, reducing the precipitation in the Mediterranean and increasing precipitation in Northern Europe. In contrast, global perturbations of the scattering aerosol SO_4 have a negligible impact. The results from this study may have important implications to the management of regional water resources, agricultural practice, ecosystems, environment, and economics as well as social development and behavior in a warming climate. They also stress the importance of accounting for the aerosols (and generally short-lived forcers) for short-term (e.g., decadal) regional climate prediction.

Acknowledgement

All model results used for this study are available to the public through the Norwegian NORSCORE data storage facility. We thank the three reviewers for their insightful comments. We also acknowledge the NASA High-End Computing Program through the NASA Center for Climate Simulation at Goddard Space Flight Center for computational resources to run the GISS-E2R model and support from NASA GISS. PDRMIP is partly funded through the Norwegian Research Council project NAPEX (project number 229778). O. Boucher acknowledges HPC resources from CCRT under the genmip6 allocation provided by GENCI (Grand Equipement National de Calcul Intensif). P. Forster and T. Richardson were supported by NERC grants NE/K007483/1 and NE/N006038/1. Ø. Hodnebrog was



partly funded through the Norwegian Research Council project HYPRE (project no. 243942). A.Voulgarakis and M.Kasoar are supported by NERC under grant NE/K500872/1. Simulations with HadGEM3-GA4 were performed using the MONSoon system supplied under the Joint Weather and Climate Research Programme of the Met Office and NERC. D. Olivić, A. Kirkevåg and T. Iversen were supported by the Norwegian Research Council through the projects EVA (grant 229771), EarthClim (207711/E10), NOTUR (nn2345k), and NorStore (ns2345k). T. Takemura was supported by the supercomputer system of the National Institute for Environmental Studies, Japan, the Environment Research and Technology Development Fund (S-12-3) of the Ministry of the Environment, Japan and JSPS KAKENHI Grant Numbers 15H01728 and 15K12190. Computing resources for CESM1-CAM5 (ark:/85065/d7wd3xhc) simulations were provided by the Climate Simulation Laboratory at NCAR Computational and Information System Laboratory, sponsored by the National Science Foundation and other agencies.

References

- Allan, R., and Ansell, T. (2006). A new globally complete monthly historical gridded mean sea level pressure dataset (hadslp2): 1850–2004. *Journal of climate*, 19(22), 5816–5842. doi: 10.1175/JCLI3937.1
- Allen, R., Norris, J., and Wild, M. (2013). Evaluation of multidecadal variability in cimp5 surface solar radiation and inferred underestimation of aerosol direct effects over europe, china, japan, and india. *Journal of Geophysical Research: Atmospheres*, 118(12), 6311–6336. doi: 10.1002/jgrd.50426
- Allen, R. J., and Sherwood, S. C. (2011). The impact of natural versus anthropogenic aerosols on atmospheric circulation in the community atmosphere model. *Climate Dynamics*, 36(9–10), 1959–1978. doi: 10.1007/s00382-010-0898-8
- Allen, R. J., Sherwood, S. C., Norris, J. R., and Zender, C. S. (2012). Recent northern hemisphere tropical expansion primarily driven by black carbon and tropospheric ozone. *Nature*, 485(7398), 350–354. doi: 10.1038/nature11097
- Andreae, M. O., Jones, C. D., and Cox, P. M. (2005). Strong present-day aerosol cooling implies a hot future. *Nature*, 435(7046), 1187. doi: 10.1038/nature03671
- Arora, V., Scinocca, J., Boer, G., Christian, J., Denman, K., Flato, G., Kharin, V., Lee, W., and Merryfield, W. (2011). Carbon emission limits required to satisfy future representative concentration pathways of greenhouse gases. *Geophysical Research Letters*, 38(5). doi: 10.1029/2010GL046270
- Bellouin, N., Rae, J., Jones, A., Johnson, C., Haywood, J., and Boucher, O. (2011). Aerosol forcing in the climate model intercomparison project (cmip5) simulations by hadgem2-es and the role of ammonium nitrate. *Journal of Geophysical Research: Atmospheres*, 116(D20). doi: 10.1029/2011JD016074
- Bentsen, M., Bethke, I., Debernard, J., Iversen, T., Kirkevåg, A., Seland, Ø., Drange, H., Roelandt, C., Seierstad, I., and Kristjánsson, J. (2013). The norwegian earth system model, noresm1-m—part 1: Description and basic evaluation of the physical climate. *Geosci. Model Dev*, 6(3), 687–720. doi: 10.5194/gmd-6-687-2013
- Black, E., Brayshaw, D. J., and Rambeau, C. M. (2010). Past, present and future precipitation in the middle east: Insights from models and observations. *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 368(1931), 5173–5184. doi: 10.1098/rsta.2010.0199
- Bond, T. C., Doherty, S. J., Fahey, D., Forster, P., Berntsen, T., DeAngelo, B., Flanner, M., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S. (2013). Bounding the role of black carbon in the climate system: A scientific assessment. *Journal of Geophysical Research: Atmospheres*, 118(11), 5380–5552.
- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S. K., Sherwood, S., Stevens, B., and Zhang, X. Y., 2013: Clouds and aerosols. In: *Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [T. F. Stoker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex & P. M. Midgley (eds.)]. Cambridge University Press, Cambridge, UK and New York, USA.
- Buffoni, L., Maugeri, M., and Nanni, T. (1999). Precipitation in italy from 1833 to 1996. *Theoretical and Applied Climatology*, 63(1–2), 33–40. doi: 10.1007/s007040050089
- Collins, M., Knutti, R., Arblaster, J., Dufresne, J.-L., Fichetef, T., Friedlingstein, P., Gao, X., Gutowski, W., Johns, T., Krinner, G., Shongwe, M., Tebaldi, C., Weaver, A., and Wehner, M., 2013: Long-term climate change: Projections, commitments and irreversibility. In: *Climate change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [T. F. Stoker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex & P. M. Midgley (eds.)]. Cambridge University Press, Cambridge, UK and New York, USA.



- Collins, W., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Halloran, P., Hinton, T., Hughes, J., Jones, C., Joshi, M., Liddicoat, S., Martin, G., O'Connor, F., Rae, J., Senior, C., Storch, S., Totterdell, I., Wiltshire, A., and Woodward, S. (2011). Development and evaluation of an earth-system model-hadgem2. *Geoscientific Model Development*, 4(4), 1051. doi: 10.5194/gmd-4-1051-2011
- 5 Dufresne, J.-L., Foujols, M.-A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki, S., Bellenger, H., Benshila, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., and Cadule, P. (2013). Climate change projections using the ipsl-cm5 earth system model: From cmip3 to cmip5. *Climate Dynamics*, 40(9-10), 2123-2165. doi: 10.1007/s00382-012-1636-1
- Dünkeloh, A., and Jacobeit, J. (2003). Circulation dynamics of mediterranean precipitation variability 1948–98. *International Journal of Climatology*, 23(15), 1843-1866. doi: 10.1002/joc.973
- 10 Dunstone, N., Smith, D., Booth, B., Hermanson, L., and Eade, R. (2013). Anthropogenic aerosol forcing of atlantic tropical storms. *Nature Geoscience*, 6(7), 534-539. doi: 10.1038/ngeo1854
- Flato, G., Marotzke, J., Abiodun, B., Braconnot, P., Chou, S.-C., Collins, W., Cox, P., Driouech, F., Emori, S., Eyring, V., Forest, C., Gleckler, P., Guilyardi, E., Jakob, C., Kattsov, V., Reason, C., and Rummukainen, M., 2013: Evaluations of climate models. In: *Climate change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [T. F. Stoker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex & P. M. Midgley (eds.)]. Cambridge University Press, Cambridge, UK and New York, USA.
- 15 Gent, P. R., Danabasoglu, G., Donner, L. J., Holland, M. M., Hunke, E. C., Jayne, S. R., Lawrence, D. M., Neale, R. B., Rasch, P. J., Vertenstein, M., Worley, P. H., Yang, Z., and Zhang, M. (2011). The community climate system model version 4. *Journal of climate*, 24(19), 4973-4991. doi: 10.1175/2011JCLI4083.1
- 20 Hartmann, D., Klein Tank, A., Rusticucci, M., Alexander, L., Brönnimann, S., Charabi, Y., Dentener, F., Dlugokencky, E., Easterling, D., Kaplan, A., Soden, B. J., Thorne, P. W., Wild, M., and Zhai, P. M., 2013: Observations: Atmosphere and surface. In: *Climate change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [T. F. Stoker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex & P. M. Midgley (eds.)]. Cambridge University Press, Cambridge, UK and New York, USA.
- 25 Hodnebrog, Ø., Myhre, G., Forster, P. M., Sillmann, J., and Samset, B. H. (2016). Local biomass burning is a dominant cause of the observed precipitation reduction in southern africa. *Nature communications*, 7. doi: 10.1038/ncomms11236
- Hurrell, J. W., Holland, M. M., Gent, P. R., Ghan, S., Kay, J. E., Kushner, P. J., Lamarque, J.-F., Large, W. G., Lawrence, D., Lindsay, K., Lipscomb, W., Long, M., Mahowald, N., Marsh, D., Neale, R., Rasch, P., Vavrus, S., Vertenstein, M., Bader, D., Collions, W., Hack, J., Kiehl, J., and Marshall, S. (2013). The community earth system model: A framework for collaborative research. *Bulletin of the American Meteorological Society*, 94(9), 1339-1360. doi: 10.1175/BAMS-D-12-00121.1
- 30 Hurrell, J. W., Kushnir, Y., and Visbeck, M. (2001). The north atlantic oscillation. *science*, 291(5504), 603-605. doi: 10.1126/science.1058761
- Iversen, T., Bentsen, M., Bethke, I., Debernard, J., Kirkevåg, A., Seland, Ø., Drange, H., Kristjansson, J., Medhaug, I., Sand, M., and Seierstad, I. (2013). The norwegian earth system model, noresm1-m-part 2: Climate response and scenario projections. *Geoscientific Model Development*, 6(2), 389. doi: 10.5194/gmd-6-389-2013
- 35 Johanson, C. M., and Fu, Q. (2009). Hadley cell widening: Model simulations versus observations. *Journal of climate*, 22(10), 2713-2725. doi: 10.1175/2008JCLI2620.1
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D. (1996). The ncep/ncar 40-year reanalysis project. *Bulletin of the American Meteorological Society*, 77(3), 437-471. doi: 10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2
- 40 Kaufman, Y. J., Tanré, D., and Boucher, O. (2002). A satellite view of aerosols in the climate system. *Nature*, 419(6903), 215-223. doi: 10.1038/nature01091
- Kay, J., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., Arblaster, J., Bates, S., Danabasoglu, G., Edwards, J., Holland, M., Kushner, P., Lamarque, J., Lawrence, D., Lindsay, K., Middleton, A., Munoz, E., Neale, R., Oleson, K., Polvani, L., and Vertenstein, M. (2015). The community earth system model (cesm) large ensemble project: A community resource for studying climate change in the presence of internal climate variability. *Bulletin of the American Meteorological Society*, 96(8), 1333-1349. doi: 10.1175/BAMS-D-13-00255.1
- 45 Kirkevåg, A., Iversen, T., Seland, Ø., Hoose, C., Kristjansson, J., Struthers, H., Ekman, A. M., Ghan, S., Griesfeller, J., Nilsson, E. D., and Schulz, M. (2013). Aerosol–climate interactions in the norwegian earth system model–noresm1-m. *Geoscientific Model Development*, 6(1), 207-244. doi: 10.5194/gmd-6-207-2013
- Koren, I., Altaratz, O., Remer, L. A., Feingold, G., Martins, J. V., and Heiblum, R. H. (2012). Aerosol-induced intensification of rain from the tropics to the mid-latitudes. *Nature Geoscience*, 5(2), 118-122. doi: 10.1038/ngeo1364
- 50 Kottek, M., Grieser, J., Beck, C., Rudolf, B., and Rubel, F. (2006). World map of the köppen-geiger climate classification updated. *Meteorologische Zeitschrift*, 15(3), 259-263. doi: 10.1127/0941-2948/2006/0130
- 55 Krichak, S. O., and Alpert, P. (2005). Decadal trends in the east atlantic–west russia pattern and mediterranean precipitation. *International Journal of Climatology*, 25(2), 183-192. doi: 10.1002/joc.1124
- Kvalevåg, M. M., Samset, B. H., and Myhre, G. (2013). Hydrological sensitivity to greenhouse gases and aerosols in a global climate model. *Geophysical Research Letters*, 40(7), 1432-1438. doi: 10.1002/grl.50318
- 60 Liu, L., Shawki, D., Voulgarakis, A., Kasoar, M., Samset, B., Myhre, G., Forster, P., Hodnebrog, Ø., Sillmann, J., Aalbergssjø, S., T. A., Boucher, O., Faluvegi, G., Iversen, T., Kirkevåg, A., Lamarque, J.-F., Olivićević, D., Richardson, T., Shindell, D., Shine,



- K., Stjern, C., and Takemura, T. (2018). A pdmip multi-model study on the impacts of regional aerosol forcings on global and regional precipitation. (In preparation).
- Lorenz, E. N. (1951). Seasonal and irregular variations of the northern hemisphere sea-level pressure profile. *Journal of Meteorology*, 8(1), 52-59. doi: 10.1175/1520-0469(1951)008<0052:SAIVOT>2.0.CO;2
- 5 Mariotti, A., Struglia, M. V., Zeng, N., and Lau, K. (2002). The hydrological cycle in the mediterranean region and implications for the water budget of the mediterranean sea. *Journal of climate*, 15(13), 1674-1690. doi: 10.1175/1520-0442(2002)015<1674:THCITM>2.0.CO;2
- Meehl, G. A., Arblaster, J. M., and Collins, W. D. (2008). Effects of black carbon aerosols on the indian monsoon. *Journal of climate*, 21(12), 2869-2882. doi: 10.1175/2007JCLI1777.1
- 10 Menon, S., Hansen, J., Nazarenko, L., and Luo, Y. (2002). Climate effects of black carbon aerosols in china and india. *science*, 297(5590), 2250-2253. doi: 10.1126/science.1075159
- Muller, C., and O’Gorman, P. (2011). An energetic perspective on the regional response of precipitation to climate change. *Nature Climate Change*, 1(5), 266-271. doi: 10.1038/NCLIMATE1169
- Myhre, G., Forster, P., Samset, B., Hodnebrog, Ø., Sillmann, J., Aalbergsjø, S., Andrews, T., Boucher, O., Faluvegi, G., and Fläschner, D. (2017). Pdrmp: A precipitation driver and response model intercomparison project, protocol and preliminary results. *Bulletin of the American Meteorological Society*(2016). doi: 10.1175/BAMS-D-16-0019.1
- 15 Myhre, G., Samset, B., Schulz, M., Balkanski, Y., Bauer, S., Berntsen, T., Bian, H., Bellouin, N., Chin, M., Diehl, T., Easter, R., Feichter, J., Ghan, S., Hauglustaine, D., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Lund, M., Luo, G., Ma, X., Noije, T., Penner, J., Rasch, P., Ruiz, A., Seland, Ø., Skeie, R., Stier, P., Takemura, T., Tsigaridis, K., Wang, P., Wang, Z., Xu, L., Yu, H., Yu, F., Yoon, J., Zhang, K., Zhang, H., and Zhou, C. (2013a). Radiative forcing of the direct aerosol effect from aerocom phase ii simulations. *Atmospheric Chemistry and Physics*, 13(4), 1853. doi: 10.5194/acp-13-1853-2013
- 20 Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglested, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang, H., 2013b: Anthropogenic and natural radiative forcing. In: *Climate change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [T. F. Stoker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex & P. M. Midgley (eds.)]. Cambridge University Press, Cambridge, UK and New York, USA.
- 25 Nabat, P., Somot, S., Mallet, M., Sanchez-Lorenzo, A., and Wild, M. (2014). Contribution of anthropogenic sulfate aerosols to the changing euro-mediterranean climate since 1980. *Geophysical Research Letters*, 41(15), 5605-5611. doi: 10.1002/2014GL060798
- 30 Neale, R. B., Chen, C.-C., Gettelman, A., Lauritzen, P. H., Park, S., Williamson, D. L., Conley, A. J., Garcia, R., Kinnison, D., and Lamarque, J.-F. (2010). *Description of the near community atmosphere model (cam 4.0)*. Retrieved from https://www.cesm.ucar.edu/models/ccsm4.0/cam/docs/description/cam4_desc.pdf
- Otto-Bliessner, B. L., Brady, E. C., Fasullo, J., Jahn, A., Landrum, L., Stevenson, S., Rosenbloom, N., Mai, A., and Strand, G. (2016). Climate variability and change since 850 ce: An ensemble approach with the community earth system model. *Bulletin of the American Meteorological Society*, 97(5), 735-754. doi: 10.1175/BAMS-D-14-00233.1
- 35 Piervitali, E., Colacino, M., and Conte, M. (1998). Rainfall over the central-western mediterranean basin in the period 1951-1995. Part i: Precipitation trends. *Nuovo cimento*, 21C(3), 331-344.
- Quadrelli, R., Pavan, V., and Molteni, F. (2001). Wintertime variability of mediterranean precipitation and its links with large-scale circulation anomalies. *Climate Dynamics*, 17(5-6), 457-466. doi: 10.1007/s003820000121
- 40 Ramanathan, V., and Carmichael, G. (2008). Global and regional climate changes due to black carbon. *Nature Geoscience*, 1(4), 221-227. doi: 10.1038/ngeo156
- Ramanathan, V., Crutzen, P. J., Kiehl, J. T., and Rosenfeld, D. (2001). Aerosols, climate, and the hydrological cycle. *science*, 294(5549), 2119-2124. doi: 10.1126/science.1064034
- Samset, B., Myhre, G., Forster, P., Hodnebrog, Ø., Andrews, T., Faluvegi, G., Flaeschner, D., Kasoar, M., Kharin, V., and Kirkevåg, A. (2016). Fast and slow precipitation responses to individual climate forcings: A pdmip multi-model study. *Geophysical Research Letters*. doi: 10.1002/2016GL068064
- 45 Schmidt, G. A., Kelley, M., Nazarenko, L., Ruedy, R., Russell, G. L., Aleinov, I., Bauer, M., Bauer, S. E., Bhat, M. K., Bleck, R., Canuto, V., Chen, Y.-H., Cheng, Y., Clune, T. L., Genio, A. D., Fainchtein, R. d., Faluvegi, G., Hansen, J. E., Healy, R. J., Kiang, N. Y., Koch, D., Lacis, A. A., Legrande, A. N., Lerner, J., Lo, K. K., Matthews, E. E., Menon, S., Miller, R. L., Oinas, V., Olosa, A. O., Perlwitz, J. P., Puma, M. J., Putman, W. M., Rind, D., Romanou, A., Sato, M., Shindell, D., Sun, S., Syed, R., Tausnev, N., Tsigaridis, K., Unger, N., Voulgarakis, A., Yao, M.-S., and Zhang, J. (2014). Configuration and assessment of the giss modele2 contributions to the cmi5 archive. *Journal of Advances in Modeling Earth Systems*, 6(1), 141-184. doi: 10.1002/2013MS000265
- Schneider, U., Becker, A., Finger, P., Meyer-Christoffer, A., Rudolf, B., and Ziese, M. (2011). *Gpcc full data reanalysis version 6.0 at 2.5°: Monthly land-surface precipitation from rain-gauges built on gts-based and historic data*.
- 55 Shindell, D., and Faluvegi, G. (2009). Climate response to regional radiative forcing during the twentieth century. *Nature Geoscience*, 2(4), 294-300. doi: 10.1038/ngeo473
- Shindell, D. T., Lamarque, J.-F., Schulz, M., Flanner, M., Jiao, C., Chin, M., Young, P., Lee, Y. H., Rotstayn, L., and Mahowald, N. (2013). Radiative forcing in the accmip historical and future climate simulations. *Atmospheric Chemistry and Physics*, 13(6), 2939-2974. doi: 10.5194/acp-13-2939-2013
- 60



- Shindell, D. T., Schmidt, G. A., Miller, R. L., and Rind, D. (2001). Northern hemisphere winter climate response to greenhouse gas, ozone, solar, and volcanic forcing. *Journal of Geophysical Research: Atmospheres*, 106(D7), 7193-7210. doi: 10.1029/2000JD900547
- Shindell, D. T., Voulgarakis, A., Faluvegi, G., and Milly, G. (2012). Precipitation response to regional radiative forcing. *Atmospheric Chemistry and Physics*, 12(15), 6969-6982. doi: 10.5194/acp-12-6969-2012
- 5 Stevens, B., and Feingold, G. (2009). Untangling aerosol effects on clouds and precipitation in a buffered system. *Nature*, 461(7264), 607-613. doi: 10.1038/nature08281
- Takahashi, C., and Watanabe, M. (2016). Pacific trade winds accelerated by aerosol forcing over the past two decades. *Nature Climate Change*. doi: 10.1038/nclimate2996
- 10 Takemura, T., Egashira, M., Matsuzawa, K., Ichijo, H., O'ishi, R., and Abe-Ouchi, A. (2009). A simulation of the global distribution and radiative forcing of soil dust aerosols at the last glacial maximum. *Atmospheric Chemistry & Physics*, 9(9), 3061-3073.
- Takemura, T., Nozawa, T., Emori, S., Nakajima, T. Y., and Nakajima, T. (2005). Simulation of climate response to aerosol direct and indirect effects with aerosol transport-radiation model. *Journal of Geophysical Research: Atmospheres*, 110(D2). doi: 10.1029/2004JD005029
- 15 Taylor, K. E., Stouffer, R. J., and Meehl, G. A. (2012). An overview of cmip5 and the experiment design. *Bulletin of the American Meteorological Society*, 93(4), 485-498. doi: 10.1175/BAMS-D-11-00094.1
- Walters, D., Williams, K., Boutle, I., Bushell, A., Edwards, J., Field, P., Lock, A., Morcrette, C., Stratton, R., Wilkinson, J., Willett, R., Bellouin, N., Bodas-Salcedo, A., Brooks, M., Copsey, D., Earnshaw, P., Hardiman, S., Harris, C., Levine, R., Maclachlan, C., Manners, J., Marin, G., Milton, S., Palmer, M., Roberts, M., Rodríguez, J., Tennant, W., and Vidale, P. (2014). The met office unified model global atmosphere 4.0 and jules global land 4.0 configurations. *Geoscientific Model Development*, 7(1), 20 361-386. doi: 10.5194/gmd-7-361-2014
- Watanabe, M., Suzuki, T., O'ishi, R., Komuro, Y., Watanabe, S., Emori, S., Takemura, T., Chikira, M., Ogura, T., Sekiguchi, M., Takata, K., Yamazaki, D., Yokohata, T., Nozawa, T., Hasumi, H., Tatebe, H., and Kimoto, M. (2010). Improved climate simulation by miroc5: Mean states, variability, and climate sensitivity. *Journal of climate*, 23(23), 6312-6335. doi: 25 10.1175/2010JCLI3679.1
- Worley, S. J., Woodruff, S. D., Reynolds, R. W., Lubker, S. J., and Lott, N. (2005). Icoads release 2.1 data and products. *International Journal of Climatology*, 25(7), 823-842. doi: 10.1002/joc.1166
- Xoplaki, E., Gonzalez-Rouco, J. F., Luterbacher, J., and Wanner, H. (2004). Wet season mediterranean precipitation variability: Influence of large-scale dynamics and trends. *Climate Dynamics*, 23(1), 63-78. doi: 10.1007/s00382-004-0422-0
- 30
- 35
- 40
- 45



Table 1. Descriptions of the nine PDRMIP models (adapted from *Liu et al. (2018)*). GA = Global Atmosphere. HTAP2 = Hemispheric Transport Air Pollution, phase 2.

Model	Version	Resolution	Ocean setup	Aerosol setup	References
Canadian Earth System Model (CanESM2)	2010	2.8×2.8 35 levels	Coupled	Emission	<i>Arora et al. (2011)</i>
Goddard Institute for Space Studies Model E2, coupled with Russell ocean model (GISS-E2-R)	E2-R	2×2.5 40 levels	Coupled	Fixed concentration	<i>Schmidt et al. (2014)</i>
Hadley Center Global Environment Model, version 2-Earth System (HadGEM2-ES)	6.6.3	1.875×1.25 38 levels	Coupled	Emissions	<i>Collins et al. (2011)</i>
HadGEM3	GA 4.0	1.875×1.25 85 levels	Coupled	Fixed concentration	<i>Bellouin et al. (2011)</i> <i>Walters et al. (2014)</i>
Model for Interdisciplinary Research on Climate-Spectral Radiation-Transport Model for Aerosol Species (MIROC-SPRINTARS)	5.9.0	T85 (approx. 1.4×1.4), 40 levels	Coupled	HTAP2 emissions	<i>Takemura et al. (2005)</i> <i>Takemura et al. (2009)</i> <i>Watanabe et al. (2010)</i>
Community Earth System Model, version 1 (Community Atmosphere Model, version 4) [CESM1-CAM4]	1.0.3	2.5×1.9 26 levels	Slab	Fixed concentration	<i>Neale et al. (2010)</i> <i>Gent et al. (2011)</i>
CESM1-CAM5	1.1.2	2.5×1.9 30 levels	Coupled	Emissions	<i>Hurrell et al. (2013)</i> <i>Kay et al. (2015)</i> <i>Otto-Bliesner et al. (2016)</i>
Norwegian Earth System Model (NorESM)	NorESM1-M (intermediate resolution)	2.5×1.9 26 levels	Coupled	Fixed concentration	<i>Bentsen et al. (2013)</i> <i>Iversen et al. (2013)</i> <i>Kirkevåg et al. (2013)</i>
L'institut Pierre-Simon Laplace Coupled Model, version 5A (IPSL-CM5A)	CMIP5	3.75×1.875 39 levels	Coupled	Fixed concentration	<i>Dufresne et al. (2013)</i>

5

10




Figure 1: Normalized ΔP (change per unit forcing) for (a) CO_2 , (b) BC and (c) SO_4 . Black dots indicate the change is significant at 0.95 confidence level. Please note that the sign for SO_4 is flipped due to its negative forcing. Thus, the results shown for SO_4 is the precipitation change per unit negative forcing.

5

10




Figure 2: Domain-averaged ΔP (purple rectangles in Fig. 1) for (a) CO₂, (b) BC and (c) SO₄. Error bars of multi-model mean (MMM) are 90% inter-model spread.

5

10




Figure 3: Domain-averaged (purple rectangles in Fig. 1) energy budget change for each forcing and energy component as shown in Equation (1) and (2). It holds that $L_c \Delta P$ (blue) = ΔQ (pink) + ΔH (gray), where $L_c \Delta P$ is the change in total latent heating, ΔQ (pink) = ΔLW (green) + ΔSW (red) - ΔSH (brown) is the change in diabatic cooling of the atmospheric column due to shortwave and longwave radiation, and sensible heat flux, ΔH is the change in column-integrated dry static energy flux divergence. The error bars indicated 90% inter-model spread.

5

10

15

20




Figure 4: Same as Fig. 1, but for sea level pressure (SLP).

5

10

15

20




Figure 5: Same as Fig. 1, but for zonal wind. The green contours represent the 50-yr climatology of the zonal wind in the control simulations. The contours are at the interval of 5m/s, with positive values indicating eastward winds.

5

10





Figure 6: Scaled change for the combination of WMGHGs, BC and SO_4 for (a) spatial pattern of precipitation, (c) domain-averaged precipitation change, and (d) energy budget change, along with (b) GPCC observational data (for which gray indicate missing or incomplete data). The dots in (a) and (b) indicate changes are significant at 0.95 and 0.9 confidence level, respectively. Error bars in (c) and (d) indicate 90% uncertainty ranges.

5



5 **Figure 7:** SLP (a & b) and zonal wind change (c & d). (a) and (c) are scaled change for the combination of WMGHGs, BC and SO₄, while (b) and (d) are Hadley observational data and NCEP reanalysis data, respectively. Dots indicate the changes are significant at 0.95 confidence level. The green contours in (c) and (d) represent the climatology position of the zonal wind. The contours are at the interval of 5m/s, with positive values indicating eastward winds.




Figure 8: Precipitation trends of CMIP5 HistoricalGHG simulations during 1901-2005, (a) domain-averaged (purple rectangles in Fig. 1) trends for each model. Error bars of each model indicate 90% inter-ensemble spread and error bars of MMM indicate the 90% inter-model spread. The numbers in parentheses indicate the ensembles collected for each model. (b) spatial pattern of MMM value for the trends. Dots indicate that the change is significant at 0.95 confidence level.

5