Long-term trends of instability and associated parameters

2 over the Indian region obtained using radiosonde network

3 Rohit Chakraborty, Madineni Venkat Ratnam* and Shaik Ghouse Basha

4 National Atmospheric Research Laboratory, India.

5 *Correspondence to:* M. Venkat Ratnam (vratnam@narl.gov.in)

6

7 Abstract

8 Long-term trends of the parameters related to convection and instability obtained from 27 radiosonde 9 stations across 6 sub-divisions over the Indian region during the period 1980-2016 is presented. A total of 16 10 parcel and instability parameters along with moisture content, wind shear, and thunderstorm and rainfall 11 frequencies have been utilized for this purpose. Robust fit regression analysis is employed on the regional 12 average time series to calculate the long-term trends on both seasonal and yearly basis. The Level of Free 13 Convection (LFC) and Equilibrium Level (EL) height is found to ascend significantly in all Indian sub-14 divisions. Consequently, the coastal regions (particularly the western coasts) experience increasing in Severe 15 Thunderstorm (TSS) and Severe Rainfall Frequencies (SRF) in the pre-monsoon while the inland regions 16 (especially central India) experience an increase in Ordinary Thunderstorm (TSO) and Weak Rain Frequency 17 (WRF) during the monsoon and post-monsoon. The 16-20 year periodicity is found to dominate the long-term 18 trends significantly compared to other periodicities and the increase in TSS, and Convective Available Potential 19 Energy (CAPE) is found more severe after the year 1999. The enhancement in moisture transport and associated 20 cooling at 100 hPa along with dispersion of boundary layer pollutants is found to be the main cause for the 21 increase in CAPE which leads to more convective severity in the coastal regions. However, in inland regions 22 moisture-laden winds are absent and the presence of strong capping effect of pollutants on instability in the 23 lower troposphere has resulted in more Convective Inhibition Energy (CINE). Hence, TSO and weak rainfall 24 occurrences have increased particularly in these regions.

25 Key words: Instability, Convection, Long-term trends, Radiosonde

26

27 1. Introduction

28 Intense convective phenomena are a common climatic feature in the Indian tropical region which occurs 29 during the pre-monsoon to post-monsoon seasons (April-October) (Ananthakrishnan, 1977) and they are 30 generally accompanied by intense thunderstorms, lightning, wind gusts with heavy rainfall. Hence, they are 31 known to induce immense socio-economic hazards including loss of life and property. Several reports have 32 shown an increase in the climatic extreme occurrence and intensity of these phenomena throughout the world (33 Webster et al., 2005; Emanuel, 2006). In this connection, the traditional surface-based parcel theory has been 34 utilized to understand convective processes using atmospheric soundings as it calculates the atmospheric 35 instabilities and other parameters at various heights (Huntrieser et al., 1997; Santhi et al., 2014; Nelli et al, 36 2018a).

37 Considering the importance of studying the long-term trends in climatic extremes, a series of research 38 attempts have been orchestrated to study and predict intense convection world-wide in the last two decades (Das 39 et al. 2017). Using multiple tropical stations and re-analysis data, Gettleman et al. (2002) and Riemann-Campe 40 et al. (2009) have shown that Convective Available Potential Energy (CAPE) has been increasing very strongly 41 with a growth rate of ~20% per decade during the period 1958-1997 due to increase in surface heating and 42 moisture. Gensini and Mote (2015) projected a 236 % increase in severe thunderstorm frequency from 1980-43 1990 to 2080-2090 over the eastern United States (US). Further, Brooks (2013) used various combinations of 44 CAPE and Vertical Wind Shear (VWSH) products and results hinted towards a probable increase in the severe 45 thunderstorms over the US. It was also observed that the effect of increasing CAPE is more dominant on 46 convective severity than in case of decreasing shear. On the other hand, studies by Prein et al. (2017) showed that a recent increase of temperature has led to a rise of moisture ingress and consequently the frequency and 47 48 severity of extreme precipitation events associated with intense convection have shown a steep rise everywhere 49 in the world. At the same time, an increase in thunderstorm severity and instability has also been reported by 50 many attempts over the Asian region (Wang et al., 2011; Saha et al., 2017).

51 Over the Indian region, Manohar et al. (1999) studied the latitudinal variation and distribution of 52 thunderstorm frequency and CAPE over 78 Indian stations during 1970-1980 and he postulated that the ambient 53 temperature at 100 hPa pressure level has a strong relationship with it. Dhaka et al. (2010) utilized radiosonde 54 observations during 1958-1997 and he obtained very prominent anti-correlations on both yearly and seasonal 55 basis between convection strength (CAPE) and upper troposphere temperatures at 100 hPa (T100). Later, 56 Murugavel et al. (2012) studied the long term trends of CAPE from 32 radiosonde stations during 1984-2008 57 and revealed an alarming growth in monsoon CAPE over India with a slope of 38J/kg/year. However, he 58 additionally stated that the low-level moisture and solar cycle can have additional impact on the increasing 59 CAPE. Recently, Chakraborty et al. (2017a) and Saha et al. (2017) reported a weakening in lower tropospheric 60 instability over few Indian stations due to increasing pollution levels using reanalysis datasets. Apart from that, 61 some studies have also attempted to correlate convective severity with boundary layer phenomena, upper-62 tropospheric moisture, surface fluxes, solar effect and precipitation; (Basha and Ratnam, 2013; Basha et al., 63 2013; Murthy and Sivaramakrishnan, 2006; Allappattu and Kunnikrishnan, 2009; Xie et al., 201;, Nelli et al. 64 2018b; Rakshit at al., 2016).

65 Previous studies over India have shown the distribution of CAPE only whereas other parameters like 66 Convective Inhibition Energy (CINE), Mixed Layer CAPE (MLC), Lifted Index (LI), Total Totals Index (TTI), 67 and Precipitable Water Vapor (PWV) are also important as they explain how the atmospheric instability and 68 moisture changes at various levels of the atmosphere. In addition, the influence of climatic oscillation (Quasi-69 Biennial Oscillation (QBO), El-Nino Southern Oscillation (ENSO) and Solar Cycle) on the seasonal and annual 70 variation of convective parameters was also not studied in detail. Therefore in the present study, long-term 71 variation of parcel parameters (Lifted Condensation Level (LCL), Level of Free Convection (LFC), Equilibrium 72 Level (EL), CAPE and CINE), with instability (LI, Vertical Totals Index (VT)), moisture (PWV, and PWV at 73 low levels (PWL)), thunderstorm and rainfall severity frequencies (Thunderstorm-Severe (TSS), Thunderstorm-74 Ordinary (TSO), Weak Rain Fall (WRF) and Strong Rain Fall (SRF)) followed by Temperature at 100hPa 75 (T100) and Wind Shear (WSH) is investigated using 27 radiosonde stations along with gridded rainfall data over 76 India. This article is structured as follows: Section 2 describes the datasets and methodology adopted for the present study. Section 3 presents the long-term analysis of parcel and instability parameter over Chennai (13.08°N, 80.27°E) and 6 sub-divisions of the Indian subcontinent on both annual and seasonal basis, followed by the periodicity and split trend analysis. Finally, a discussion on the results and conclusions is appended in Section 4 and 5, respectively.

81

82 2. Dataset and Methodology

83 Radiosonde observations from 27 stations over the Indian region from 1980-2016 are obtained from 84 Integrated Global Radiosonde Archives (https://www1.ncdc.noaa.gov/pub/data/igra/derived/derived-por/). 85 These datasets provide daily temperature and humidity profiles from 1538 stations around the world in fixed 86 pressure levels after doing quality checks (Durre et al., 2006; Ferreira et al. 2018). These studies have concluded 87 that the radiosonde data quality from IGRA has faced certain problems from time to time, but such cases are not 88 so prominent over the Indian region, especially after the year 1980. It is mainly because of the higher accuracy 89 and reliability of this in-situ measurement technique that these datasets are widely used worldwide nowadays for 90 calibrating other continuous profiler instruments (Chakraborty and Maitra, 2016). In accordance with data 91 availability and reliability, only 27 stations have been considered out of 37 IGRA Indian radiosonde stations 92 thereby providing descent data availability for carrying out this study. When an in depth investigation is done on 93 the data continuity by plotting the temperature and humidity profiles for all days, a set of gaps in datasets were 94 discovered. Most of the utilized stations have intermittent data gaps of 2-7 days in certain months only but, on 95 the whole, except only a very few cases, the duration of these individual data gaps are mostly limited to less 96 than 1 month. However, these small data gaps are not expected to provide any significant impact on the long-97 term seasonal or annual average variations of 37 years x 12 months span.

98 In addition to data availability, homogeneity also acts as a common concern before using the data. 99 However, such issues should not be considered serious as all three types of homogeneity issues namely: volume, 100 instrument type and quality have been addressed before commencing the study. First, about 5000 radiosonde 101 profiles are available in majority of IGRA stations which are uniformly distributed among all years and 102 seasons(except monsoon); hence it provides a decent data volume for investigation of yearly trends. Secondly, 103 the data of all Indian IGRA stations come from a single type of IM-MK3 radiosondes which has not undergone 104 any change in radiosonde accuracies in the last years and so this addresses the instrument type related issue. 105 Finally, regarding data quality, a set of 7 quality checks are performed by IGRA before accepting the data which 106 should remove any unreliable observations before being used in the study. These 7 quality checks also include 107 repetition check which rejects any possible case of humidity sensor saturation errors during rainy conditions 108 especially in monsoon.. Thus it can be inferred that the obtained climatic trends of instability from IGRA is 109 expected to be far more reliable compared to other data sources.

These 27 stations have been divided into six homogenous regions as defined by the India Meteorological Department (IMD) (Rao, 1976) which are: Central India (CI), East Coast (EC), North East (NE), North West (NW), Peninsula India (PI) and the West Coasts (WC) as shown in Fig. 1. Further, for simplicity, these regions have again been combined into three major categories namely: coastal regions (EC and WC), Inland (CI and PI) and others (NE and NW). After retrieving the profiles some more additional internal quality checks are performed before using the data for every IGRA station. First, the balloon burst height has to be minimum of 15 km to be selected for analysis. Second, any gaps of temperature and humidity in important 117 pressure levels such as 850, 700, 500, 300, 200 and 100 hPa will produce difficulty in calculation of 118 atmospheric instability; hence those profiles were rejected in quality control tests for all stations. Again, out of 119 the available radiosonde profiles, some profiles have displayed absurd variations of temperature and humidity at 120 various heights and hence they are discarded. After completion of these quality checks it was thought that 121 atmospheric instability shows prominent diurnal variation, datasets of only one time slot can be taken for 122 analysis. As datasets at 00Z are more in number than at 12Z, the analysis has been actually progressed with 00Z 123 datasets. A complete detail about the final dataset used for every station is indicated in Table A1. It may be 124 noted here that the volume of observations are found to distributed almost homogenously throughout the 125 measurement period and a detailed year wise breakup of radiosonde launches utilized are not shown to maintain 126 the focus of the work. For calculation of the instability parameters, the temperature and humidity profiles were 127 transformed from the standard pressure levels using cubic spline interpolation at every 100 m height bins. Piece-128 wise linear/ quadratic/ cubic spline interpolation schemes are employed instead of linear interpolation in 129 temperature and humidity retrievals in this study as the former techniques can more faithfully regenerate the 130 nonlinearities in boundary layer variations of meteorological parameters according to recent studies by 131 Chakraborty et al. (2016). After this, a similar surface-based parcel method is utilized for estimating the parcel 132 and instability parameters (LCL, LFC, EL, CAPE, MLC, CINE) as already described by Chakraborty et al. 133 (2018). A small detail about the physical significance of these parameters is now given in the Supplementary 134 Section. For thunderstorm genesis, moisture growth and wind shear are extremely important, therefore we 135 calculated the total amount of water vapor (PWV) and that up to 700 hPa level (PWL) along with the horizontal 136 wind shear between surface and 6 km altitude. In addition to these, we have used temperature at the 100 hPa 137 pressure level as it is found to strongly influence the convective strengths over the Indian region (Manohar et al., 138 1999; Dhaka et al., 2010).

139 Along with these parameters, the long-term impact of instability on the convection has also been 140 studied from thunderstorm and rain frequencies. Daily measurements of surface wind speeds is obtained for all 141 the radiosonde observations at 00Z using the Wyoming Website(weather.uwyo.edu/upperair/sounding.html). 142 The thunderstorm frequencies are calculated on yearly basis based on the criterion given by IMD 143 (http://imd.gov.in/section/nhac/termglossary.pdf). According to this criterion, if the maximum surfaces wind 144 speed is greater than 62 km/h then it is considered as a severe thunderstorm event otherwise if wind speeds are 145 between 31 and 62 km/h then it is considered as an ordinary thunderstorm case (also used by Saha et al. (2014)). 146 Hereafter, the total number of thunderstorm occurrences per year in both severe and ordinary category is 147 counted and represented by thunderstorm frequencies as TSS and TSO. Here it may be noted that, the wind 148 speed measurements are taken from the first measurement of radiosonde balloon flight for all stations. These 149 datasets are always within 10m from the surface and according to WMO criterion, they can assume a maximum 150 error of 1 m/s from surface to 100 hPa level. Since a minimum wind speed of 31 kmph or 8.61 m/s is required 151 for identification as an ordinary thunderstorm, hence this 1 m/s error is not expected to perturb the thunderstorm 152 severity climatology presented in this study. In the same way, IMD provides daily rainfall accumulations in 153 0.25 degree spatial resolution over the Indian region since the year 1900 (Rajeevan et al., 2006, 2008; Pai et al., 154 2014). This daily precipitation data at the closest grid point is used to define the frequency of severe and weak 155 rainfall days hereafter referred to as WRF and SRF respectively. The severe rainfall frequencies constitute those 156 days where the daily accumulation is greater than 124.5 mm/day while for the weak rainfall cases it is less than

157 7.5 mm/day according to IMD glossary as given in http://imd.gov.in/section/nhac/termglossary.pdf.

158 From the previous section it follows that a set of 14 parcel parameters with rainfall and thunderstorm 159 frequencies are essential to understand the convective climatology over India. However, other than this, 8 160 standard instability parameters (LI, KI, TTI, CT, VT, CAPE, CINE and MLC) are also additionally important to 161 quantify the thunderstorm severity, hence must also be considered for analysis. Now, it is known that most of 162 these instability parameters are inter-related; hence PCA analysis is done to identify and use only those 163 instability parameters that can give a complete but independent overview of the atmospheric instability using 164 minimum parameters. In this analysis, introduced by (Hoteling 1936) a set of possibly related parameters are 165 converted into orthogonal independent components after which the primary components are plotted with the 166 initial parameters. Parameter variance scores present at the farthest distance from the primary principal 167 components and also from all the other variables contain the highest variance; hence they are selected for 168 representing the existing group of old inputs. Hence in the present study, daily datasets of all 6 instability 169 parameters are averaged to yearly values for every regions and then the principal component analysis is 170 performed on the datasets are obtained. Daily datasets have not been directly used for PCA as it would have too 171 many fluctuations which would make the redundant parameter identification very difficult in all cases. The 172 variance distribution plot (not shown) for each region showed that only the first two components contribute to 173 more than 70% of the total variance; hence the covariance scores of these two strongest orthogonal components 174 are plotted in Fig. 2 which depict that the LI is completely unrelated to rest of the parameters. Again, since VT 175 is found to lie exactly in the middle of the rest of the parameters, and it also represents the lower tropospheric 176 instability in a much more suitable way hence this parameter is also used with LI to represent the rest of the 177 instability parameters in a convenient way. Consequently, LI and VT are additionally considered along with the 178 previous set of 14 attributes to get the final set of 16 parameters for further analysis.

179 Thus, a set of 16 parameters are finally taken for the analysis: LCL, LFC, EL, LI, VT, CAPE, CINE, 180 MLC, PWV, PWL, WSH, T100, TSO, TSS, WRF and SRF. However, apart from the IGRA radiosonde and the 181 IMD rainfall database, it was believed that some other parameters may also be externally responsible for the 182 changing trends in atmospheric instability and hence they are also included. They comprise the monthly mean 183 aerosol absorption index (AAI) data taken from the Tropospheric Emission Monitoring Internet Service 184 (TEMIS) Air Pollution Archive (De Graaf et al., 2005). In addition, the monthly average gridded data of ozone 185 mixing ratio (OMR) and Specific Humidity (SHUM) along with Downward Long Wave Radiation Flux 186 (DLWRF) are also utilized from **ERA-Interim Re-analysis** datasets 187 (https://apps.ecmwf.int/datasets/data/interim-full-daily/levtype=sfc/).

188 We have estimated all parameters from daily radiosonde data and averaged over a season and annually 189 for obtaining trend at 95% confidence interval using robust regression analysis (Shepard, 1968). Further, the 190 parameters from radiosonde were averaged region wise and then the robust fit algorithm is employed on the 191 normalized time series to get the long-term trends (Andersen, 2008; Raj et al., 2018). These yearly trend values 192 are multiplied by 37 to get the total climatological trend in one parameter over the complete data span of 1980-193 2016. For seasonal trend analysis, the same approach has been utilized for different seasons. The seasonal 194 distribution has been adopted from IMD reports which are as follows: Pre-monsoon (March-May), Monsoon 195 (June-September), Post-monsoon (October-November) and Winter (December-February). Further, for studying the periodicities associated with each of these time series, an Empirical Mode Decomposition (EMD) technique

197 is used (Wu and Huang, 2009). Finally, the robust fit analysis is done on each of these components to compare

198 the trends from each periodicity to determine which of the periodicities dominates in each parameter.

199

200 3. Results

201 3.1. Climatic trends over Chennai

202 In the previous study by Chakraborty et al. (2018), long term trends of instability were investigated over 203 Gadanki(13.5°N,79.2°E) situated on a hilly terrain with an altitude of 370 m above sea level at a distance of 204 ~150 km from the eastern coasts and Bay of Bengal . To see whether, the observed trends of these parameters 205 are behaving similarly in case of IGRA profiles also the climatic trends of instability are now described over 206 Chennai (13.08°N, 80.27°E) which is the closest radiosonde station from Gadanki. The yearly averaged datasets 207 are normalized with respect to their climatic mean and are plotted along with 1 sigma standard errors in Fig 3 208 after which robust fit regression analysis (Andersen 2008) is utilized to obtain the climatological trends in these 209 parameters as shown by red solid lines in the plots. A decreasing trend in VT and increase of magnitude in 210 CINE with LI is noticed which indicates a reduction in the lower atmospheric instability (Fig.3d,e,h,i). 211 However, CAPE (Fig.3f) shows significant increasing trends throughout the period. LFC has a slightly 212 ascending trend (~18 hPa) which leads to increasing CINE and decreasing VT over Chennai, while the EL is 213 found to get lifted up drastically (Fig.3c) resulting in an increase in the total instability and CAPE. The increase 214 in height of EL can be caused by a reduction in temperatures in the upper tropospheric heights (Manohar et al, 215 1999). Hence, it can be inferred that the reduction in temperatures near 100 hPa (Fig.31) plays an important role 216 in modulating the total atmospheric instability and CAPE.

The enhancement in CINE magnitude and reduction in VT leads to the reduction in the frequency of weaker convective systems with medium or lower CAPE values. Again, as CAPE is one of most important parameters that modulate convective severity, hence the frequency of severe thunderstorms and heavy rainfall occurrences is expected to rise (Fig.3n,p). Thus, it is inferred that lower level instability has reduced due to elevated CINE and LFC; while the upper-level atmospheric instability has intensified significantly due to a cooling at 100 hPa and ascension in EL over Chennai. Hence, CAPE value increases drastically leading to more severe thunderstorm and heavy rainfall frequency events during the mentioned period.

Before proceeding to the investigation on the climatological trends of convection and instability over the Indian region, it is necessary to validate whether the obtained hypothetical trends from Chennai are free from any data quality issues. Hence a region wise climatology of the most important parameter CAPE is obtained from all the Indian regions using ERA-Interim Reanalysis data and the trends are shown in Fig. S1. This figure clarifies that all the Indian regions (especially the coastal regions) have experienced a common rise in CAPE especially after 1996-2000. Thus, the stated hypothesis looks clear and hence this can be progressed over a much broader way.

However, it should be noted that Fig 3 provides too much detailed and cumbersome results related to all 16 parameters and the complexity of the analysis is expected to increase further when similar analysis will be presented for all the Indian regions together. But on the other hand, for a complete understanding about the morphology of upper and lower tropospheric instability, all the instability parameters will be required. Hence, to reduce chances of confusion and to make the results more compact, all 16 parameters will be discussed together but only a few of them will be presented in the main study. After a thorough consideration with respect to the
main objective of the present attempt, 8 parameters namely LFC, EL, CAPE, CINE, PWV, T100, TSS and SRF
are retained in the main figures while their complementary aspects such as LCL, LI, VT, MLC, PWL, WSH,
TSO and WRF are shown in the supplementary sections.

240 3.2. Climatological average of parameters

241 The climatological mean values of all instability parameters over six different Indian sub-divisions are 242 shown with boxplot analysis (McGill, 1978) in Fig. 4 and Fig. S3. The utility of using this approach is that, it 243 will reveal which regions of India shows normal expected variation (if it lies within the box limits signifying 25-244 75% percentage of the distribution), while on the other hand it will also identify those regions having the 245 extreme outlier values (lying outside the whiskers signifying the outermost 5% of the distribution). The LCL 246 (Fig.S3a) and LFC (Fig.4a) are found to be at the lowest altitudes in the coastal regions. As these stations 247 receive most of the moisture from sea, the EL (Fig.4b) is also expected to be higher at the coastal areas and 248 lower elsewhere. However, due to low moisture availability, the inland regions experience weaker instability 249 which results in lower CAPE (~900 J/kg) (Fig.4c) with higher CINE (Fig.4d) and WSH (Fig.S3f). During strong 250 convection, the values of LI (Fig.S3b) (which represents that the mid-tropospheric instability) are also expected 251 to be more negative in the coastal regions. Similarly, height integrals of instability such as CAPE (Fig.4c) and 252 MLC (Fig.S3d) are significantly higher (~1500 J/kg) in the coastal regions while the magnitude of MLC 253 (Fig.S3d) is found to be almost half of CAPE. As the trends in total convective strengths below 300 hPa are 254 quite low compared to that over the total atmospheric column, hence it follows that the portion of buoyant 255 column above 300 hPa must have contributed significantly to the total convective developments over the Indian 256 region. Again, being opposite of CAPE, CINE values are minimum in the coastal regions compared to inland 257 and continental regions thereby serving as a potential cause for the reduced instability in these regions.

258 Similar to CAPE and MLC combination, the PWV (Fig.4e) and PWL (Fig.S3e) pair shows the highest 259 averages in the coastal regions due to their closest proximity to the adjoining seas. Also, PWL (moisture integral 260 up to 700 hPa) is found to be almost half of PWV, hence the mid and upper tropospheric humidity is found to 261 play a strong role in modulating the convective systems over India. The instability and moisture are highest in 262 the coastal regions hence the frequency of severe thunderstorms and rainfall occurrences are comparatively 263 higher (Fig.4g,h). The North Western region shows the large values of thunderstorm frequency which is not 264 supported by other parameters. Hence, it may be inferred that this is due to frequent dry storms called "Andhi" 265 which have no relation with convective instability and rainfall (Rajpal and Deka, 1980). Thus, it can be 266 concluded that the effect of convection is large in the coastal regions compared to other regions which resulted 267 in high CAPE with more thunderstorms and intense rain occurrences.

268 **3.3.** Long-term trends in the instability parameters

The long-term trends are calculated for each parameter during the entire study period of 1980-2016 for all regions using the robust regression analysis at 95% confidence interval as depicted in Fig. 5 and Fig. S4. For simplicity, the average trends along with their standard deviation values are depicted in Table 1. Also to investigate about the significance of trend values calculated from these time series datasets, a *t-test* analysis (Gosset, 1908)is done on all parameter and locations. The p values are calculated at 95% confidence limits for ttest analysis on all instability parameters over the Indian sub-divisions and interestingly, all the values are found to be below 0.05. Hence the time series variations to be presented in subsequent sections will always be 276 statistically significant in nature. So, to have a better quantitative measure of the trend significance, the total 277 changes in each of these parameters are presented in percentage form in place of the p values in the table. This 278 process will enable an easy identification of regions experiencing more accelerated convective growth. But on 279 the other hand, while analysing the results of the trend analysis in statistical form, the absolute trend has to be 280 given more importance as the % changes completely depend on the magnitude of the long term mean. The LCL 281 (Fig.S4a) height is found to decrease which may lead to an overall increase in the number of rain occurrences 282 throughout the country (provided that the amount of atmospheric instability is adequate). On contrary, LFC 283 (Fig.5a) is found to ascend in all the regions except NE resulting in the reduction of lower level instability and a 284 an increase of CINE magnitude (Fig.5d). However, the extent of change in LFC (Fig.5a) and LCL (Fig.S4a) is 285 smallest in the coastal regions (~10 hPa). In case of EL (Fig.5b), a very prominent ascent is depicted in all 286 regions (highest in coastal regions) which increase the height of the buoyant column; hence the net effect on 287 total instability and CAPE (Fig.5c) is expected to increase significantly. Similarly, LI (Fig.S4b) values become 288 more negative in all the regions with slightly higher magnitudes in the coastal regions. VT represents the lower 289 level atmospheric instability and hence is expected to be affected by the elevation in LFC. Thus, a reduction in 290 VT (Fig.S4c) is seen with minimum values in the coastal regions (~0.3), medium in the NE and NW regions 291 (~0.5) and highest in deep inland regions such as CI and PI (~0.8). An intensification in CAPE (Fig.5c) is seen 292 in all regions (~1100 J/kg) as expected from EL (Fig.5b) and LI. However, the increase is the highest (~100%) 293 at the coastal regions whereas in MLC (Fig.S4d), which is measured only up to 300 hPa level, the increment is 294 only 20% of that in CAPE. Hence, it follows that the maximum contribution towards the increase in CAPE 295 comes above 300 hPa. In case of CINE, an overall enhancement in values is observed as expected (~60 J/kg). In 296 addition, the trend values suggest a two-fold increase of CINE in inland regions while the values are much lesser 297 (50%) in the coastal regions due to balancing effect from strong convections and CAPE in those regions.

298 The PWV (Fig.5e) and PWL (Fig.S4e) values are increasing similar to CAPE and MLC. The long-term 299 trends in PWV are about 10% of its climatological average with highest in the coastal regions. Further, the 300 lower level moisture content of PWL (up to 700 hPa) showed an increase but the trend values are comparatively 301 smaller (~6%). Hence it follows that it is not the lower tropospheric moisture (below 700 hPa) but the remaining 302 amount which is increasing significantly for all regions. Now, as this growth in upper tropospheric moisture is 303 analogous with a parallel rise in upper level CAPE, hence there should be a possible association between these 304 two factors which needs to be investigated. The WSH (Fig.S4f) parameter increases in all regions of the 305 country, and hence it produces an inhibiting effect on the lower level instability. An upper tropospheric cooling 306 trend is observed in all other regions (Fig. 5f) with minimum values in the inland regions and maximum in the 307 coastal regions. Consequently, the increase in CAPE values is maximum in the coastal regions and lesser 308 elsewhere. The ordinary thunderstorm frequency is also found to increase (Fig. S4g) which may be due to the 309 partial damping effect of an elevated LFC and CINE on lower level instabilities. However, the TSS (Fig.5g) is 310 found to increase at a much higher rate compared to TSO especially in the coastal regions. On the other hand, an 311 increase in CINE and decrease of VT lead to an increase in the number of WRF (Fig.S4h). However, due to rise 312 in CAPE and TSS (Fig.5g), the SRF (Fig.5h) is also found to rise significantly by about 20% particularly in the 313 coastal regions. It may be noted that, as EL has more dominant effect on CAPE hence the rise in SRF is much 314 larger than that WRF (5%). Finally, the long-term trends have been compared between the east and west coastal 315 regions and it is observed that the rate of increase in total instability is the most prominent in the western coasts

while factors related to ascending LFC, CINE and reducing VT are more significant in the central India which is the farthest from both the sea coasts. Thus, the long-term analysis infers that lower atmospheric instability has reduced while the upper tropospheric instability and moisture increased drastically over the Indian region. As a result, convective severity as expressed in terms of CAPE, TSS and SRF is increasing more strongly in the coastal regions while in the continental areas this effect is dampened due to the contribution of increasing CINE and WSH.

322 **3.4.** Seasonal effect on long-term trends in the instability parameters

323 The seasonal variation of the long-term variations of atmospheric instability is shown in Fig. 6 and Fig. 324 S5. LCL shows a uniform descent by 10 hPa in all seasons (Fig.S5a) whereas LFC ascends in most of the 325 regions and seasons (Fig.6a). However, this ascent is more prominent in the monsoon and post-monsoon season. 326 However, the seasonal variation is absent in EL and LI (Fig.6b, Fig S5b) which are mainly associated with an 327 upper layer phenomenon. VT shows the most prominent reduction in monsoon and post-monsoon seasons 328 (Fig.S5c). MLC and CAPE show a lot of regional disparities but with a common increase in its value in all the 329 seasons (Fig.6c, S5d). In monsoon and post-monsoon, the increase in CAPE is slightly lesser due to the effect of 330 decreasing VT and elevated LFC. CINE is closely related to VT and LFC, hence it shows slight increase (of 331 magnitude) in the monsoon and post-monsoon seasons with maximum values in inland regions as expected 332 (Fig.6d).

333 PWV, PWL and WSH represent a prominent increase in the monsoon followed by the post-monsoon 334 (Fig.6e, Fig.S5e-f). T100 is related to an upper atmospheric phenomenon hence no seasonal or spatial variation 335 is displayed, except for a small cooling effect in pre-monsoon (Fig.6f) due to the prevalence of intense 336 convections events which is supported by the strongest increase in CAPE. A decrease in lower atmospheric 337 instability and increase in CINE is observed; hence TSO and WRF are expected to increase. However, this 338 increase is found more dominant only in the monsoon and post-monsoon (Fig.S5g,h). Another interesting result 339 is that TSS and SRF do not behave similarly. TSS increases almost uniformly in all seasons with the highest in 340 the pre-monsoon. However, SRF increases mainly in the monsoon followed by the post-monsoon season 341 (Fig.6g,h). The observed disparity between them is due to the profuse moisture availability during monsoon and 342 post-monsoon compared to the pre-monsoon.

Further, in seasonal trends, east and west coasts show equivalent trends in all instability parameters while the Central India still remains as the region which is most affected by the ascension of LFC and CINE. Thus, the seasonal analysis reveals that the yearly long-term trends are almost uniformly distributed in all the seasons. The ordinary and weak thunderstorm frequencies show the strongest increase during monsoon and post-monsoon while the upper atmospheric instability shows a weak influence in the pre-monsoonal trends on the yearly climatology.

349 3.5. Effect of specific periodicities on long-term trends

In case of both annual and seasonal trend analysis, all Indian sub-divisions are found to follow similar behavior. Hence, to find out the periodicities in the average long-term trends, the time series of all regions are averaged and then subjected to EMD technique which reveals the existence of four main periodicities namely: 1.5 - 2.5 years corresponding to QBO, 4-6 years corresponding to ENSO, 10-12 years corresponding to the solar cycle and the fourth one is of 16-20 years. A similar multi-decadal climatic oscillation was also reported by 355 Dhaka et al. (2010). Hence for simplicity, this periodicity has been renamed as a Multi-decadal Climatic356 Oscillation (MCO).

357 The climatic trends of these periodicities for each parameter are calculated from robust regression 358 analysis. An illustration of the obtained MCO periodicities for CAPE along all the Indian regions is shown in 359 Fig. S2. Further for comparison, the trend values from each periodicity is normalized to percentage with respect 360 to the total trend values for each parameters and the net contribution of these individual periodicities are 361 depicted in Fig. 7 and Fig. S6. The figure suggests that ENSO, QBO and solar cycle have no effect on LCL 362 (Fig.S6a) while MCO is quite strong. LFC shows minimal effects to all periodicities except solar cycle period 363 which may be due to solar-terrestrial heating (Fig.7a). EL and LI are significantly affected by both solar and 364 MCO periodicities (Fig.7b, Fig, S6b). But in LI, the contribution from MCO is much more than solar effect. In 365 case of VT (Fig. S6c) the effect of both ENSO and MCO are found prominent. CAPE is found to be strongly 366 influenced by MCO followed by solar effect (Fig.7c) and this is also discernible from the most strong increasing 367 trends in CAPE especially in the coastal regions after the years 1996-2000 in Fig. S2. However, in case of MLC, 368 contribution of MCO is comparatively lesser (Fig.S6d) hence some separate phenomena above 300 hPa may 369 have prominent influence on increasing CAPE. Apart from CAPE, effect of MCO is also found very strong in 370 case of CINE (Fig.7d).

371 The moisture parameters like PWV and PWL show similar variability as in CAPE and MLC which 372 indicates significant moisture transport changes only above 300 hPa in the past 18 years (Fig.7e, Fig S6e). WSH 373 does not show the dominance of any periodicity (Fig.S6f) while T100 shows the most prominent contribution 374 from the MCO (Fig.7f) thereby showing its connection with the long-term variability in EL and CAPE with 375 associated thunderstorm severity in the recent years. TSO and TSS are both affected by solar and MCO 376 (Fig.S6g, Fig.7g) but TSS shows that the effect of MCO is higher compared to TSO. Finally, the effect of MCO 377 is also found more prominent in case of SRF and WRF (Fig.7h, Fig S6h). In nutshell, the MCO acts as the most 378 dominant periodicity which has influenced the convective severity over India. So, in the coming sections, the 379 MCO trends for both halves of 37 years will be studied, For ease of indication and referencing, these trends of 380 18 years span each will be hereafter mentioned as quasi-bi-decadal trends (since bothe spans are close to 20 381 years in length).

- 382
- 383

3.6. Investigation of quasi-bi-decadal trends between 1980-1997 and 1999-2016

384 In the previous section, the annual averaged time series of many parameters such as EL, LI, VT, CAPE, 385 CINE, T100, TSS, WRF and SRF showed very significant changes with respect to MCO. It has also been 386 indicated from Fig. S2 that the climatic trends before and after the period 1996-2000 are significantly different 387 from each other. Therefore, the trends have been estimated with respect to two time periods before and after the 388 year 1998. The time series for both MCO are produced and their trend values are represented in Fig. 8 and Fig. 389 S7. For simplicity, the MCO are referred as C1 (1980 to 1998) and C2 (1999 to 2016), respectively. Starting 390 with LCL, in C1 there is almost no change, but in C2 there is a strong descent which influences the overall 391 change in the time series (Fig.S7a). In case of LFC, C1 shows an ascending trend, but in C2, a significant 392 increasing pattern of LFC pressure is seen hence an overall descent is obtained (Fig.8a). An ascent in the EL is 393 noticed in both the periods however during C2 the trends show significant enhancement (Fig.8b). LI values are 394 expected to become more negative from 37 years trend, however its absolute magnitude shows a slight reduction in C1 followed by a prominent increase in C2 resulting in a net increase in instability(Fig.S7b). VT shows an overall decreasing pattern in both the periods (Fig.S7c). CAPE (Fig.8c) shows an enhancement in both the cycles but the trends become more prominent in C2 (1500 J/Kg). Similar to CAPE, MLC (Fig.S7d) also shows an increasing trend in both the cycles but the trend values are also much smaller than in CAPE. Hence, the rise in EL height can be considered as a primary factor for increase in CAPE above 300 hPa during C2. CINE shows increasing trend in both C1 and C2 but again the trend values are much stronger (~80 J/kg) during C2 especially in the inland regions (Fig.8d)..

402 The moisture trends in both PWV and PWL have shown a constant increase in both the MCO 403 throughout India (Fig.8e, Fig S7e). The WSH (Fig.8k) also increases uniformly in both MCO with strongest 404 trends in the inland regions. A prominent cooling of ~1.5 degrees is seen in 100 hPa levels everywhere in C1, 405 but in C2 the trend increases to ~-2.5 degrees (Fig.8f) which can be considered responsible for the abrupt 406 elevation in EL and increasing CAPE values during the recent years. TSO increases slightly in C2 compared to 407 C1 (Fig.S7g). But in case of TSS, the positive trend gets doubled in C2 mainly in the coastal regions (Fig.8g). 408 Finally, in case of SRF the trend values in C2 are slightly higher with the maximum magnitudes in the coastal 409 regions as expected (Fig.8h). A further comparison between the six regions reveals that the west coast shows the 410 maximum enhancement in all the instability and convective severity parameters in the past 18 years due to 411 strong growth in moisture content and associated cooling at 100 hPa.

412 On the contrary, during C2 central India suffers from the maximum reduction in lower level instability 413 as seen from the rise in CINE and LFC due to the dearth of moisture content. Similar results are also found in 414 other coastal and inland regions. Hence it follows that mainly during C2, the upper tropospheric instability has 415 enhanced everywhere while the lower tropospheric instability has reduced which has led to the development in 416 both CAPE and CINE. As a result both TSS-TSO and WRF -SRF combination increases.

417

418 4. Discussion

419 From the previous section, it is inferred that a cooling trend at 100 hPa levels has led to the ascent in EL 420 which results in an increase in CAPE, TSS and SRF. To explain the reason behind this, we consider the ozone to 421 be a primary heating agent by absorbing the incoming solar ultraviolet radiation near 100 hPa level 422 (Mohanakumar, 2008). OH hydroxyl radicals are formed by oxidation of water vapor molecules by a reactive 423 oxygen atom at the same height. On the other hand, it has been reported by Forster et al. (2007) that in the recent 424 years there has been a cooling in upper troposphere due to decrease in ozone concentration near 70 hPa. Hence, 425 it is hypothesized that the OH radicals formed from the oxidation of water vapour can take an active role in the 426 breakup of ozone molecules at 100 hPa levels which may lead to this cooling effect. The preceding sections 427 have shown a significant increase in moisture content especially in the coastal areas hinting towards more 428 moisture transport from the adjoining seas. Again, an increase in LI and CAPE values have also been reported in 429 most of the regions which can lift the available moisture to upper atmospheric levels (Das et al., 2016; Guha et 430 al., 2017). To add to this increasing CAPE and LI, many recent researchers' have reported a net increase in the 431 Hadley cell and Brewer-Dobson circulation strength (Liu et al., 2012; Fu et al., 2015; Shepherd and 432 McLandress, 2011) which also assists in the up-liftment of moisture to upper atmospheric levels. Thus, it is 433 inferred that low-level moisture is transported to the upper troposphere and above where it is responsible for 434 ozone depletion and cooling thereby elevating the EL and increasing the thunderstorm severity.

435 To test this hypothesis, yearly averaged time series of specific humidity and ozone mixing ratio data are 436 collected for all stations and the quasi-bi-decadal trend values are depicted in Fig. 9. This figure shows a rise in 437 specific humidity levels by 7% in C2 over entire India (Fig.9a). On the other hand, trends of specific humidity 438 have almost trebled in C2 phase with the maximum values in the coastal regions (Fig.9e). As water vapor 439 concentration increases, ozone concentration is expected to decrease. The ozone trends support this hypothesis 440 by showing a sharp transition from low positive to high negative values during C2 (Fig.9b,f). It may be 441 additionally noted that the specific humidity increase and reduction in ozone content are strongest in the coastal 442 regions leading to a higher increase in CAPE and severe thunderstorms in those regions.

443 In the recent decades, Indian region has experienced a surface warming trend which is mainly caused 444 by an increase in greenhouse gas concentrations as pointed out by Basha et al. (2017). These greenhouse gases 445 are heat absorbing in nature and these particles reside within the lower troposphere (generally below 700 hPa) 446 due to surface heating and boundary layer dynamics as reported by Chakraborty et al. (2017b). Further, these 447 gases has a tendency to absorb and then re-emit the outgoing longwave radiation as emitted by the Earth 448 resulting in more downward longwave radiation flux and atmospheric heating which elevates the LFC. 449 Additionally, this near surface heating reduces the vertical temperature lapse rate leading to a drop in lower 450 instability (VT). To test this hypothesis, yearly averaged Downward Long Wave Radiation Flux (DLWRF) time 451 series is depicted over the Indian region in Fig. 9(c,g) which also suggests that DLWRF values are increasing in 452 C2. To show that the increase in DLWRF is due to the heat absorbing particles only, the trends in Absorbing 453 Aerosol Index (AAI) are shown for all the regions. The figure suggests that the mean of AAI is increasing 454 slightly more in C2 with a positive trend (Fig.9d,h). Due to this heating of lower atmosphere and capping of 455 lapse rates by greenhouse gases and absorptive aerosols, the LFC starts ascending, so WSH and CINE get 456 stronger while VT reduces. As a result the ordinary to weak convective occurrences start increasing.

457 Finally, it has to be explained why the upper air instability and CAPE are increasing mainly in the 458 coastal regions. The coastal regions have high moisture content (Saha et al., 2017). Because of the strong land-459 ocean contrast, low-level winds close to 850 hPa flow into the coastal regions and disperse the pollutants and 460 greenhouse gases to other locations leading to a weaker convective inhibition in those areas. This hypothesis is 461 supported by the lowest AAI values in the coasts despite having high increasing trends in those areas. In 462 addition, the ample moisture supply in the coastal regions is lifted up to the upper troposphere and lower 463 stratosphere (UTLS) where it undergoes prominent cooling due to ozone reduction. Hence, the EL ascends more 464 resulting in higher CAPE which finally led to an abrupt rise in TSS and SRF in the coastal regions. However, in 465 the inland regions the layer of absorptive aerosols and greenhouse gases cannot be dispersed amply due to the 466 dearth of strong lower level winds. As a result, the growth of lower atmospheric instability gets inhibited in the 467 inland regions. Further, due to less moisture availability, UTLS cooling and EL ascent are much lower hence 468 there is a less rise in CAPE which ultimately leads to an increase in TSO and WRF in those sub-divisions. It 469 may be noted that the trend in AAI is not significantly different for the two time periods C1 and C2. Again, it is 470 the EL and not the LFC or LCL which influences CAPE strongly; hence the strong trends of humidity increase 471 and ozone reduction overpowers the weaker inhibitory effect from the atmospheric aerosols and this acts as a 472 major driving force behind the increase in convective severity compared to in most of the cases. 473

474

475 5. Summary and conclusions

476 In recent decades, global warming has become a threat to human life and society in terms of its various 477 implications. Increase in surface temperature leads to stronger atmospheric instabilities which in term may 478 increase the CAPE resulting in more severe thunderstorm and precipitation activity. Hence, the long term 479 variations of instability parameters will help to better understand the changes in the weather extremes with 480 respect to climate change. In light of the above, the main objective of the present study is to analyze whether 481 convective instability is changing over the Indian region during the last 37 years, and then to find its possible 482 effects on thunderstorm and rainfall severity. Radiosonde measurements from Integrated Global Radiosonde 483 Archives (IGRA) pertaining to 27 stations across 6 Indian sub-divisions are utilized to depict the spatial 484 distribution of these long-term trends during the period of 1980-2016. The selection of instability parameters is 485 done based on Principle Component Test (PCT) which showed the importance of taking LI and VT for further 486 investigations. A total of 16 parameters (including parcel and instability data with moisture content, wind shear, 487 and thunderstorm and rainfall frequencies) have been utilized. Robust fit approach is employed on the regional 488 average time series to calculate the long-term trends on both yearly and seasonal basis. The main highlights 489 obtained from the present study are listed below:

- The coastal regions experience the most significant rise in Convective Available Potential Energy (CAPE)
 and Equilibrium Level (EL) leading to more occurrences of Severe Thunderstorms (TSS) and severe
 rainfall events (SRF) while the inland regions undergo a decrease in lower atmospheric instability due to
 elevated Convective Inhibition Energy (CINE) and Level of Free Convection resulting in more Ordinary
 thunderstorm (TSO) and Weak Rainfall occurrences (WRF).
- 495
 2. In the pre-monsoon season, an increasing TSS activity is observed due to higher instability connected to
 496
 496
 497
 497
 498
 498
 498
 498
 499
 499
 499
 490
 490
 490
 490
 490
 490
 491
 491
 492
 493
 494
 494
 495
 495
 496
 496
 497
 498
 498
 498
 498
 499
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
 490
- The Empirical Mode Decomposition (EMD) analysis on the instability parameters reveals that the 16-20
 year multi-decadal oscillation (MCO) as the most dominant component in all six Indian sub-divisions.
- 502 4. The quasi-bi-decadal analysis reveals an increase in magnitude in many parameters like EL, CAPE, CINE,
 503 TSO and TSS along with cooling at 100 hPa level during C2 (1999-2016) which dominates 37-year trend.
- 5. The annual and quasi-bi-decadal trends support that the increase in thunderstorm severity and associated convection is strongest along western coasts due to maximum moisture ingress from the seas while the greatest reduction in lower atmospheric instability is experienced in central India owing to the lack of pollutant dispersal as it is situated very far from the seas.
- 508
 6. In the coastal regions, ample amount of water vapor is advected into the mid-troposphere from the surrounding seas which in presence of strong lifting goes up to upper troposphere and lower stratosphere
 510 (UTLS) where ozone depletion occurs leading to a strong cooling effect. This cooling effect enables the ascent in EL resulting in much stronger LI and CAPE values, hence more TSS and SRF..
- 512 7. In the inland regions, the dispersing effect by sea winds is absent hence the capping effect of lower513 instability is more leading to stronger CINE values. Again, due to the dearth of moisture transport from the

514 seas, the UTLS cooling is lesser; leading to a weaker rise in CAPE consequently the TSO and WRF 515 frequencies increase significantly.

- 516 517
- 518 519

8. However, as the ascent in EL has a stronger contribution over increasing CAPE than the inhibitory effect of LFC, hence the long term trends are expected to be more strongly influenced by the ozone decomposition and cooling at 100 hPa levels than the capping effect of low level inversions from absorptive aerosols; hence the convective severity over the Indian regions is found to increase.

520 Thus, it may be inferred that in the near future also, convective severity will increase strongly in the 521 coastal regions while weak and ordinary thunderstorms will be more common in the inland regions. It may 522 appear at certain sections of this analysis that the trends of CAPE and EL are exorbitantly high; but it is not the 523 actual case because previous studies by Murugavel et al (2012) and Gettlemann et al. (2002) have also shown 524 almost comparable trends in convective severity both in India and abroad. Nevertheless, this study gets an upper 525 hand over the previous approaches as it successfully explains the hypothesis brought forward by early research 526 attempts that UTLS cooling at 100 hPa and greenhouse gases concentration rise can regulate the climatic trends 527 of convective severity and frequency especially over tropical regions in the recent decades.

528 After going through the study, there may be a possibility of thinking that the change in instability 529 trends is due to the change in sensors around 1998. But this is not the actual case because first, there has not 530 been any mention in past literature survey related to any change in radiosonde data quality during late 1990s in 531 IMD or IGRA. Secondly, the yearly variations of all 16 parameters for various IGRA stations as in Chennai do 532 not commonly show any abrupt change in time series during 1996-2004 except for a few cases. Thirdly it has 533 been revealed by IMD reports that the year 2000 was a tipping point for the climate change led warming over 534 India thereby leading to a rise in catastrophic weather events and a cataclysmic fallout will follow by the year 535 2040 if these emission scenarios are not curbed recently (Hindustan Times, 2019). Thus, it follows that the 536 observed changes in the atmospheric instability trends before and after 1996-2004 are due to a synoptic global 537 warming based climate change phenomena and not due to any change in radiosonde sensor type.

538 However, in spite of all this, the present study has certain shortcomings. The most important one 539 among them is that this set of explanations is based on isolated information from selected in-situ observations 540 and hence it needs to be studied in more detail spatially in future using model-based observations. In the recent 541 years, certain studies have utilized multiple GCM outputs over the US to infer the robust increase in 542 thunderstorm frequency (Diffenbaugh et al., 2013; Seeley and Romps, 2015). However, these types of studies 543 have not yet been done over the Indian region. Hence, a combination of multi-station radiosonde data with 544 model data will be utilized to provide a generalized picture about convective severity over the Indian region. On 545 the other hand, this study also introduces the effect of direct aerosol heating on instability and convection; but 546 the probable impact of indirect aerosol loading on modulating the cloud lifetime and convective severity has not 547 been discussed here. This is because, the relationship between indirect aerosol forcing and instability is still 548 unclear and complex (Connoly et al. 2012). A few researches in the recent years have hypothesized that a higher 549 concentration of aerosols may lead to stronger updrafts velocities by altering the latent heat release resulting in 550 growth of CAPE and TSS (Tao et al. 2012; Storer and van den Heever, 2013). However, this is a season and 551 location specific phenomena and hence it is not expected to impact the yearly trend of CAPE and TSS as strong 552 as the upper tropospheric cooling effect projected in this study. But in future, an exhaustive analysis of cloud

- 553 and aerosol components involving both in-situ and modelled data can to be done to investigate its contribution
- 554 on the total CAPE, TSS and SRF trends over the Indian region.
- 555

556 Author contribution

- 557 Rohit Chakraborty performed complete analysis and written the first draft. MVR provided the initial concept,
- 558 guiding, and editing and SGB contributed in analysis, discussion and editing.
- 559

560 Acknowledgments

561 One of the authors (Rohit Chakraborty) thanks, Science and Engineering Research Board, Department 562 of Science and Technology for providing fellowship under National Post-Doctoral Scheme (File 563 No:PDF/2016/001939). He also acknowledges National Atmospheric Research Laboratory, for providing 564 necessary support and data for this work. The authors also thank S.T. Akhil Raj, Sanjeev Dwivedi and N. 565

- Narendra Reddy for their suggestions. Data used in present study can be obtained directly from IGRA website.
- 566

567 References

- 568 Alappattu, D.P. and Kunhikrishnan, P.K.: Premonsoon estimates of convective available potential energy over 569 oceanic region surrounding Indian subcontinent. J. Geophys. Res., Atmos. 114, 2009.
- 570 Ananthakrishnan, R.: Some aspects of the monsoon circulation and monsoon rainfall. Pure Appl. Geophys., 571 115(5-6), 1209-1249, 1977.
- 572 Andersen, R.: Modern methods for robust regression. 152, Quantitative Applications in Social Sciences, Sage, 573 2008.

574 Basha, G., Kishore, P., Ratnam, M. V., Jayaraman, A., Kouchak, A. A., Ouarda, T. B. M. J., and Velicogna, I.: 575 Historical and Projected Surface Temperature over India during 20 th and 21 st century. Sci. Rep., 7(1), 576 2987, 2017.

- 577 Basha, G., and Ratnam, M.V.: Moisture Variability over Indian monsoon regions, Atmos. Res., 132-133, 35-578 45, 2013.
- 579 Basha, G., Ratnam, M.V., and Krishna Murthy, B.V.: Upper tropospheric water vapor variations, J Earth Syst 580 Sci., 122(6), 1583-1591, 2013.
- 581 Brooks, H. E.: Severe thunderstorms and climate change. Atmos. Res., 123, 129-138, 2013.
- 582 Chakraborty, R. and Maitra, A.: Retrieval of atmospheric properties with radiometric measurements using 583 neural network. Atmos. Res., 181:124-132, https://doi.org/10.1016/j.atmosres.2016.05.011, 2016.
- 584 Chakraborty, R., Talukdar, S., Saha, U., Jana, S., and Maitra, A.: Anomalies in relative humidity profile in the 585 boundary layer during convective rain. Atmos. Res., 191, https://doi.org/10.1016/j.atmosres.2017.03.011, 586 2017a.
- 587 Chakraborty, R., Saha, U., Singh, A. K., and Maitra, A.: Association of atmospheric pollution and instability 588 indices: Α detailed investigation over an Indian urban metropolis. Atmos. Res., 589 https://doi.org/10.1016/j.atmosres.2017.04.033, 2017b.
- 590 Chakraborty, R., Basha, G., Ratnam and M.V.: Diurnal and long-term variation of instability indices over a 591 tropical region in India. Atmos. Res., https://doi.org/10.1016/j.atmosres.2018.03.012, 2018.

- 592 Connolly PJ, Vaughan G,May PT, Chemel C, Allen G, Choularton TW, Gallagher MW, Bower KN, Crosier J,
 593 Dearden C. 2013. Can aerosols influence deep tropical convection? Aerosol indirect effects in the Hector
 594 island thunderstorm. Q. J. R. Meteorol. Soc. 139: 2190–2208. DOI:10.1002/qj.2083.
- Das, S., Chakraborty, R. and Maitra, A.: A random forest algorithm for nowcasting of intense precipitation
 events, Adv. Sp. Res., 60(6), 1271-1282, 2017, https://doi.org/10.1016/j.asr.2017.03.026.
- 597 Das, S.S., Ratnam, M.V., Uma, K.N., Patra, A.K., Subrahmanyam, K.V., Girach, I.A., Suneeth, K.V., Kumar
 598 K.K., and Ramkumar, G.: Stratospheric intrusion in to the troposphere during the tropical cyclone Nilam
 599 (2012), Q. J. R. Meteorolog. Soc. doi: 10.1002/qj.2810, 2016.
- bhaka, S. K., Sapra, R., Panwar, V., Goel, A., Bhatnagar, R., Kaur, M.,: Influence of large-scale variations in convective available potential energy (CAPE) and solar cycle over temperature in the tropopause region at Delhi (28.3°N, 77.1°E), Kolkata (22.3° N, 88.2° E), Cochin (10° N, 77° E), and Trivandrum (8.5°N, 77.0°E) using radiosonde during 1980--2005. Earth Planets Space, 62(3), 319–331, 2010.
- 604 De Graaf, M., Stammes, P., Torres, O., and Koelemeijer, R. B. A.: Absorbing Aerosol Index: Sensitivity 605 analysis, application to GOME and comparison with TOMS. J. Geophys. Res. Atmos., 110(D1), 2005.
- Diffenbaugh, N. S., Scherer, M., and Trapp, R. J.: Robust increases in severe thunderstorm environments in response to greenhouse forcing. Proc. Natl. Acad. Sci., 110(41), 16361–16366, 2013.
- Durre, I., Vose, R. S., and Wuertz, D. B.: Overview of integrated global radiosonde archive. J. Clim., 19(1), 53–68, 2006.
- Emanuel, K.: Climate and tropical cyclone activity: A new model downscaling approach. J. Clim., 19(19),
 4797–4802, 2006.
- Ferreira, A.P., Nieto, R., Gimeno, L., Completeness of radiosonde humidity observations based on the IGRA,
 Earth Syst. Sci. Data Discuss., 2018, https://doi.org/10.5194/essd-2018-95.
- Forster, P. M., Bodeker, G., Schofield, R., Solomon, S. and D. Thompson: Effects of ozone cooling in the
 tropical lower stratosphere and upper troposphere, Geophys. Res. Lett., 34, L23813,
 doi:10.1029/2007GL031994, 2007.
- Fu, Q., Lin, P., Solomon, S., and Hartmann, D. L.: Observational evidence of the strengthening of the BrewerDobson circulation since 1980. J. Geophys. Res. Atmos., 120(19), 2015.
- 619 Gosset, W.S., The probable error of a mean, Biometrika, 6(1), 1-25, 1908.
- Gensini, V. A., and Mote, T. L.: Downscaled estimates of late 21st century severe weather from CCSM3. Clim.
 Change, 129 (1-2), 307-321, 2015.
- 622 Gettelman, A., Seidel, D. J., Wheeler, M. C., and Ross, R. J.: Multidecadal trends in tropical convective
 623 available potential energy. J. Geophys. Res. Atmos. 107(D21), 2002.
- 624 Hindustan Times: <u>https://www.hindustantimes.com/environment/freak-weather-to-rise-in-india-</u> 625 over-two-decades/story-T1G8SqfBh8jydT15UnKGuM.html, last access: 24 January, 2019.
- Hotelling, H., Analysis of complex statistical variables into principal components, J. Educational Psychology,
 24, 417-411, 1936.
- Guha, B. K., Chakraborty, R., Saha, U., and Maitra, A.: Tropopause height characteristics associated with ozone
 and stratospheric moistening during intense convective activity over Indian sub-continent. Global Planet.
 Change, 158. https://doi.org/10.1016/j.gloplacha.2017.09.009, 2017.

- Huntrieser, H., Schiesser, H.H., Schmid, W.,and Waldvogel, A.: Comparison of Traditional and Newly
 Developed Thunderstorm Indices for Switzerland, Weather Forecasting, 12, 108-125, 1997.
- Joseph, P. V., Raipal, D. K., and Deka, S. N.: Andhi. The Convective Dust Storms of Northwest India. Mausam,
 31, 431–442, 1980.
- Kharin, V. V, Zwiers, F. W., Zhang, X., and Wehner, M.: Changes in temperature and precipitation extremes in
 the CMIP5 ensemble. Clim. Change, 119(2), 345–357, 2013.
- Liu, J., Song, M., Hu, Y., and Ren, X.: Changes in the strength and width of the Hadley circulation since 1871.
 The climate of the past, 8(4), 1169–1175, 2012.
- Manohar, G. K., Kandalgaonkar, S. S., and Tinmaker, M. I. R: Thunderstorm activity India and Indian
 southwest monsoon. J. Geophys. Res. Atmos. 104(D4), 4169–4188, 1999.
- 641 McGill, R, Turkey, J.W., Larsen, W.A., Variations of Box Plots, The American Staistician. 32(1), 12-16, 1978.
- 642 Mohanakumar, K.:Stratosphere-troposphere interactions: introduction.Springer Science Business Media, 2008.
- Murthy, B.S. and Sivaramakrishnan, S.: Moist convective instability over the Arabian Sea during the Asian
 summer monsoon, 2002, Meteorol. Appl., https://doi.org/10.1017/S135048270500201X, 2006.
- Murugavel, P., Pawar, S. D., and Gopalakrishnan, V.: Trends of convective available potential energy over the
 Indian region and its effect on rainfall. Int. J. Climatol., 32(9), 1362–1372, 2012.
- Nelli, N. R., Ratnam, M.V., Basha, G., and Ravikiran, V.: Cloud vertical structure over a tropical station
 obtained using long-term high resolution radiosonde measurements, Atmos. Chem. Phys. Discuss.,
 https://doi.org/10.5194/acp-2018-194, 2018a.
- Nelli, N. R., and Rao, K.G.: Contrasting variations in the surface layer structure between the convective and
 non-convective periods in the summer monsoon season for Bangalore location during PRWONAM, J.
 Atmos. Sol. Terr. Phys., https://doi.org/10.1016/j.jastp.2017.11.017, 2018b.
- Pai, D. S., Sridhar, L., Rajeevan, M., Sreejith, O. P., Satbhai, N. S., and Mukhopadhyay, B.: Development of a new high spatial resolution (0.25 x 0.25) long period (1901--2010) daily gridded rainfall data set over India and its comparison with existing data sets over the region. Mausam, 65(1), 1–18, 2014.
- Prein, A. F., Rasmussen, R. M., Ikeda, K., Liu, C., Clark, M. P., and Holland, G. J.: The future intensification of
 hourly precipitation extremes. Nat. Clim. Change, 7(1), 48, 2017.
- Raj, S. T. A., Ratnam, M. V., Rao, D. N., and Murthy, B. V. K.: Long-term trends in stratospheric ozone,
 temperature, and water vapor over the Indian region. Ann. Geophys. (Vol. 36, p. 149), 2018.
- Rajeevan, M., Bhate, J., Kale, J. D., and Lal, B.: High resolution daily gridded rainfall data for the Indian region: Analysis of break and active monsoon spells. Current Science, 296–306, 2006.
- Rajeevan, M., Bhate, J., and Jaswal, A. K.: Analysis of variability and trends of extreme rainfall events over
 India using 104 years of gridded daily rainfall data. Geophys. Res. Lett., 35(18), 2008.
- Rakshit, G., Chakraborty, R. and Maitra, A.: Effect of vertical wind on rain drop size distributions in the
 boundary layer, 2016 International Symposium on Antennas and Propagation (ISAP), 846-847, IEEE, 2016,
 URL: http://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=7821528&isnumber=7821043.
- 667 Rao, Y. P.:Southwest monsoon. Synoptic Meteorology, 1976.
- Riemann-Campe, K., Fraedrich, K., and Lunkeit, F.: A global climatology of convective available potential
 energy (CAPE) and convective inhibition (CIN) in ERA-40 reanalysis. Atmos. Res., 93(1), 534–545, 2009.

- Saha, U., Maitra, A., Midya, S. K., and Das, G. K.: Association of thunderstorm frequency with rainfall
 occurrences over an Indian urban metropolis. Atmos. Res., 138, 240–252, 2014.
- Saha, U., Chakraborty, R., Maitra, A., and Singh, A. K.: East-west coastal asymmetry in the summertime nearsurface wind speed and its projected change in future climate over the Indian region. Glob. Planet. Change,
 152., https://doi.org/10.1016/j.gloplacha.2017.03.001, 2017.
- Santhi, Y. D., Ratnam, M. V., Dhaka, S. K., and Rao, S. V.: Global morphology of convection indices observed
 using COSMIC GPS RO satellite measurements. Atmos. Res., 137, 205–215, 2014.
- Seeley, J. T., and Romps, D. M.: The effect of global warming on severe thunderstorms in the United States. J.
 Clim., 28(6), 2443–2458, 2015.
- Shepard, D.: A two-dimensional interpolation function for irregularly-spaced data. In Proceedings of the 1968
 23rd ACM national conference (pp. 517–524), 1968.
- Shepherd, T. G., and McLandress, C.: A robust mechanism for the strengthening of the Brewer-Dobson circulation in response to climate change: Critical-layer control of subtropical wave breaking. J. Atmos. Sci., 68(4), 784–797, 2011.
- Storer, R.L., van den Heever, S.C.: Microphysical processes evident in aerosol forcing of tropical deep
 convective clouds. J. Atmos. Sci., 70, 430–446, 2013 doi.org/10.1175/JAS-D-12-076.1
- Tao, W.K., Chen, J.P., Li, Z., Wang, C., Zhang, C., 2012. Impact of aerosols on convective clouds and
 precipitation. *Review of Geophysics*, 50, RG2001. doi:10.1029/2011RG000369
- Wang, S.-Y., Davies, R. E., Huang, W.-R., and Gillies, R. R.: Pakistan's two-stage monsoon and links with the
 recent climate change. J. Geophys. Res. Atmos., 116(D16), 2011.
- Webster, P. J., Holland, G. J., Curry, J. A., and Chang, H.-R.: Changes in tropical cyclone number, duration, and intensity in a warming environment. Science, 309(5742), 1844–1846, 2005.
- Wu, Z., and Huang, N. E.: Ensemble empirical mode decomposition: a noise-assisted data analysis method.
 Advances in Adaptive Data Analysis, 1(1), 1–41, 2009.
- Kie, B., Zhang, Q., and Ying, Y.: Trends in precipitable water and relative humidity in China: 1979--2005. J.
 Appl. Meteorol. Clim., 50(10), 1985–1994, 2011.
- 696

Name		CI		EC		NE		NW		PI		WC			India							
		μ	o	%	μ	o	%	μ	ď	%	μ	o	%	μ	ď	%	μ	o	%	μ	o	%
LCL	(hPa)	39	2.8	4.43	8.9	0.1	0.93	15.4	0	1.71	24	0.6	2.63	45.5	1.0	5.17	9.1	0.2	0.96	23	6.4	2.49
LFC	(hPa)	-38	1.4	5.57	-11	0.2	1.41	13.4	0.1	1.91	-44	3.8	6.63	-9.2	0.0	1.36	-17	0.6	2.19	-18	8.7	2.46
EL	(hPa)	-188	11.2	49.2	-280	27.1	82.8	-206	8.8	60.5	-230	2.0	67.6	-239	6	63.6	-311	40	91.4	-242	19	68.1
LI	(°C)	-0.8	0.01	16.7	-1.7	0.2	22.2	-1.3	0.1	22.8	-1.1	0.1	17.7	-1.4	0.0	27.2	-1.8	0.1	24.6	-1.3	0.1	20.3
VT	(°C)	-0.7	0.01	2.98	-0.3	0.02	1.50	-0.5	0	2.24	-0.5	0.0	2.32	-0.9	0.1	3.85	-0.4	0.0	1.95	-0.5	0.1	2.34
CAPE	(J/kg)	617	2.9	82.8	1589	90.8	108	1137	30	125	858	53	90.3	1000	39	107	1554	198	98.9	1126	159	97.9
MLC	(J/kg)	55	0.24	12.1	288	9.6	42.8	273	4.8	56.7	134	1.1	29.7	201	16	43.7	323	27	42.0	212	42	36.8
CINE	(J/kg)	-94	7.4	87.8	-36	0.3	46.7	-30	1.2	27.8	-85	6.9	103	-67	2.4	62.6	-44	1.5	55.3	-59	11	73.7
PWV	(mm)	1.4	0.03	5.71	3.2	0.03	10.0	3.7	0.1	13.4	1.3	0.0	4.72	2.2	0.0	8.97	3.9	0.1	11.2	2.6	0.5	8.85
PWL	(mm)	-0.2	0	1.95	0.4	0	2.75	0.7	0.1	6.22	0.0	0	4.65	0.6	0.0	5.71	0.6	0.0	3.98	0.4	0.2	3.32
WSH	(/s)	5.8	0.3	78.4	3.4	0.2	54.4	4.8	0.2	75.0	3.4	0.0	54.4	5.5	0.6	74.8	3.6	0.1	68.6	4.4	0.4	69.8
T100	(°C)	-1.5	0.03	3.00	-2.5	0.1	5.20	-0.4	0	0.83	-0.3	0	0.59	-2.5	0.3	5.13	-2.2	0.2	4.68	-1.6	0.4	3.00
TSO		1.4	0.05	24.3	0.8	0	10.5	2.2	0.0	53.1	3.5	0.2	53.8	2.7	0.2	40.9	0.5	0	4.92	1.8	0.5	27.3
TSS		1.5	0.0	70.5	2.3	0.05	144	2	0.1	250	2.5	0.3	209	1.7	0.1	81.7	2.3	0.1	131	2.1	0.2	147
WRF		2.9	0.1	9.51	3.8	0.1	6.55	4.6	0.4	11.5	2.2	0.0	4.19	2.8	0.1	8.88	6.3	0.2	11.4	3.8	0.6	7.28
SRF		0.4	0.0	32.4	0.8	0.06	22.2	0.2	0.0	8.30	0.2	0	14.8	0.2	0	14.4	1.1	0.1	39.5	0.5	0.2	20.5

Table 1: Statistical information related to the 37-year trend of all instability parameters over the six sub-divisions of India (μ: long-term average, σ: standard
 deviation, %: total percentage trend).

Figures



Figure 1: The locations of the 27 IGRA stations used for the present study. The distribution of the 27 stations over Indian regions is as follows: 4 stations in the CI, 6 stations in EC, 4 stations in NE, 4 stations in the NW, 5 stations in the PI and finally 4 stations in WC.



Figure 2: Principle Component Analysis for selection of instability parameters for the long-term trend study in (a) Central India (CI), (b) East Coast (EC), (c), North East (NE), (d) North West (NW), (e) Peninsular India (PI) and (f) West Coasts (WC) obtained using IGRA observations.



Figure 3: Long-term variation of (a) LCL, (b) LFC, (c) EL, (d) LI, (e) VT, (f) CAPE, (g) MLC, (h) CINE, (i) PWV, (j) PWL, (k) WSH, (l) T100, (m) TSO, (n) TSS, (o) WRF and (p) SRF observed over Chennai.



Figure 4: Climatological mean values of (a) LFC, (b) EL, (c) CAPE, (d) CINE, (e) PWV, (f) T100, (g) TSS and (h) SRF over the six sub-divisions of India. Coastal Regions are represented by red cones, the north eastern and western regions are denoted by black stars and diamonds while the blue cones represent the inland regions. Here the box limits refer to the upper and lower quartiles (25% and 75%) while the whiskers refer to the outlier limit of the data (5% and 95% limit of the population).

1



Figure 5: Long-term variation of (a) LFC, (b) EL, (c) CAPE, (d) CINE, (e) PWV, (f) T100, (g) TSS and (h) SRF over the six sub-divisions of India during the period 1980-2016. Legends are same as in Figure 4.



Figure 6: Seasonal trend of long-term variation for (a) LFC, (b) EL, (c) CAPE, (d) CINE, (e) PWV, (f) T100, (g) TSS, and (h) SRF over India during all seasons. Here 1 refers to pre-monsoon (March-May), 2 refers to Monsoon (June-September), 3 for Post-monsoon (October-November) and 4 for Winter (December-February). Legends are same as in Figure 4.



Figure 7: Percentage contribution of various periodicities on long-term trend of all instability parameters over India namely: 1.5 -2.5 years periodicity denoted as 1, 4 -6 years periodicity denoted as 2, 10-12 years periodicity displayed as 3 and 16-20 years periodicity represented as 4.



Figure 8: Comparison of average values for two time periods indicating the trend of various instability parameter over the six sub-divisions of India in two half periods of 18 years each (the numbers 1 and 2 represent the first and second period, C1 and C2, during 1980-1997 and 1999-2016, respectively) during 1999-2016. Legends are same as in Figure 4.



Figure 9: Average values and climatological trends of specific humidity, ozone mixing ratio at 100 hPa and Downward Longwave Radiation Flux (DLWRF) with Absorptive Aerosol Index (AAI) over the six sub-divisions of India over two half periods of 18 years each (1980-1997 and 1999-2016). Legends are same as in Figure 4.

Appendix

Sl no.	Station	Latitu de	Longit ude	Altitude above MSL	Station Name	Initial No. of profiles available	No. of profiles at 00 & 12Z	No. of profile s at 00Z	Region		
1	42361	26.23	78.25	205	Gwalior	9901	9412	4530			
2	42369	26.75	80.88	122	Lucknow	16869	16387	8963	Central		
3	42379	26.75	26.75 83.37 78 Gorakhpur		12376	11793	6170	India			
4	42667	23.28	77.35	522	Bhopal	14795	13968	4472			
5	42809	22.65	88.45	6	Kolkata	15212	14626	6980			
6	42971	20.25	85.83	45	Bhubaneshwar	18325	17552	6672			
7	43150	17.68	83.3	70	Vishakhapatna m	13225	12856	6355	Eastern Coasts		
8	43185	16.2	81.15	3	Machilipatnam	17108	16374	8014			
9	43279	13	80.18	14	Chennai	14067	13487	8278			
10	43346	10.92	79.83	7	Karaikal	16519	16106	6890			
11	42314	27.48	95.02	110	Dibrugarh	10067	9550	3801			
12	42410	26.1	91.58	54	Guwahati	15280 8934	14803	8681	North		
13	42492	25.6	85.17	51	Patna		8318	4370	Eastern		
14	42724	23.88	91.25	16	Agartala	15234	14732	6340			
15	42101	30.33	76.47	251	Patiala	11572	10129	4663			
16	42182	28.58	77.2	210	New	14077	13982	6581	North		
17	42339	26.3	73.02	217	Jodhpur	13133	12918	5274	Western		
18	42647	23.06	72.63	55	Ahmadabad	11430	11006	5540			
19	42867	21.1	79.05	310	Sonegaon	15626	14971	8532			
20	43014	19.85	75.4	585	Aurangabad	14220	13993	4032	D · 1		
21	43041	19.08	82.03	554	Jagdalpur	10568	10205	5437	Peninsular		
22	43128	17.45	5 78.47 530 Hyderabad		Hyderabad	10234	9723	6195	munu		
23	43295	12.97	77.58	917	Bangalore	10150	9514	4899	9		
24	43003	19.12	72.85	14	Bombay	14102	13808	7030			
25	43192	15.48 73.82		58	Goa	7070	6313	5180	Western		
26	43285	12.95	74.83	31	Mangalore	9866	9406	5020	Coasts		
27	43371	8.48	76.95	60	Trivandrum	11590	11120	8304			

Table A1: Details of the dataset used.

Abbreviation Full Form Slno. LCL (hPa) Lifted Condensation Level 1 2 Level of Free Condensation LFC (hPa) 3 Equilibrium Level EL (hPa) 3 $LI(^{\circ}C)$ Lifted Index 4 $VT(^{o}C)$ Vertical Totals Index 5 Convective Available Potential Energy CAPE (J/kg) 6 MLC (J/kg) Mixed Layer CAPE 7 **Convective Inhibition** CINE (J/kg) 8 PWV (mm) Precipitable Water Vapour 9 PWL (mm) Lower Level PWV 10 WSH (s^{-1}) Wind Shear T100 (°C) 11 Temperature at 100 hPa 12 TSO Ordinary Thunderstorm Frequency TSS 13 Severe Thunderstorm Frequency 14 WRF Weak Rainfall Frequency 15 Severe Rainfall Frequency SRF 16 Specific humidity SHUM (kg/kg) 17 AAI Absorptive Aerosol Index 18 IMD India Meteorological Department 19 IGRA Integrated Global Radiosonde Archive 20 GHG Green House Gas 21 DLWRF (W/m^2) **Downward Long Wave Radiation Flux**

Table A2: List of Abbreviations

23

24

25

EMD

UTLS

QBO

ENSO

Empirical Mode Decomposition

Upper Troposphere Lower Stratosphere

Quasi-biennial oscillation

El Niño-Southern Oscillation