

**Diel variation of mercury stable isotope ratios record photoreduction of PM<sub>2.5</sub>-bound mercury**

Qiang Huang<sup>1,2</sup>, Jiubin Chen<sup>1,3,\*</sup>, Weilin Huang<sup>4</sup>, John R. Reinfelder<sup>4</sup>, Pingqing Fu<sup>5</sup>, Shengliu

5 Yuan<sup>1</sup>, Zhongwei Wang<sup>1</sup>, Wei Yuan<sup>1</sup>, Hongming Cai<sup>1</sup>, Hong Ren<sup>5</sup>, Yele Sun<sup>5</sup> and Li He<sup>6</sup>

<sup>1</sup> SKLEG, Institute of Geochemistry, CAS, Guiyang 550081, China

<sup>2</sup> SKLOG, Guangzhou Institute of Geochemistry, CAS, Guangzhou 510640, China

<sup>3</sup> Institute of Surface-Earth System Science, Tianjin University, 300072, China

<sup>4</sup> Department of Environmental Sciences, Rutgers, The State University of New Jersey, New  
10 Brunswick, NJ 08901, USA

<sup>5</sup> LAPC, Institute of Atmospheric Physics, CAS, Beijing 100029, China

<sup>6</sup> Laboratory of Hebei Institute of Regional Geology and Mineral Resources Survey,  
Shijiazhuang 065000, China

\*Corresponding author.

15 E-mail: jbchen@tju.edu.cn

**Abstract.** Mercury (Hg) bound to fine aerosols (PM<sub>2.5</sub>-Hg) may undergo photochemical reaction that causes isotopic fractionation and obscures the initial isotopic signatures. In this study, we quantified Hg isotopic compositions for 56 PM<sub>2.5</sub> samples collected between Sept. 15<sup>th</sup> and Oct. 16<sup>th</sup>, 2015 from Beijing, China, among which 26 were collected during the daytime (between 8:00 a.m. and 6:30 p.m.) and 30 during night (between 7:00 p.m. and 7:30 a.m.). The results show that diel variation was statistically significant ( $p < 0.05$ ) for Hg content,  $\Delta^{199}\text{Hg}$  and  $\Delta^{200}\text{Hg}$ , with Hg content during the daytime ( $0.32 \pm 0.14 \mu\text{g g}^{-1}$ ) lower than at night ( $0.48 \pm 0.24 \mu\text{g g}^{-1}$ ) and  $\Delta^{199}\text{Hg}$  and  $\Delta^{200}\text{Hg}$  values during the daytime (mean of  $0.26\text{‰} \pm 0.40\text{‰}$  and  $0.09\text{‰} \pm 0.06\text{‰}$ , respectively) higher than during the nighttime ( $0.04\text{‰} \pm 0.22\text{‰}$  and  $0.06\text{‰} \pm 0.05\text{‰}$ , respectively), whereas PM<sub>2.5</sub> concentrations and  $\delta^{202}\text{Hg}$  values showed insignificant ( $p > 0.05$ ) diel variation. Geochemical characteristics of the samples and the air mass backward trajectories (PM<sub>2.5</sub> source related) suggest that diel variation in  $\Delta^{199}\text{Hg}$  values resulted primarily from the photochemical reduction of divalent PM<sub>2.5</sub>-Hg, rather than variations in emission sources. The importance of photoreduction is supported by the strong correlations between  $\Delta^{199}\text{Hg}$  and: (i)  $\Delta^{201}\text{Hg}$  (positive, slope = 1.1), (ii)  $\delta^{202}\text{Hg}$  (positive, slope = 1.15), (iii) content of Hg in PM<sub>2.5</sub> (negative), (iv) sunshine durations (positive), and (v) ozone concentration (positive) observed for consecutive day-night paired samples. Our results provide isotopic evidence that local, daily photochemical reduction of divalent Hg is of critical importance to the fate of PM<sub>2.5</sub>-Hg in urban atmospheres and that, in addition to variation in sources, photochemical reduction appears to be an important process that affects both the particle mass-specific abundance and isotopic composition of PM<sub>2.5</sub>-Hg.

**Keywords:** Mass-independent fractionation; Atmospheric particulate mercury; Transport and transformation; Isotopic evidence of photoreduction

## 1 Introduction

40 Atmospheric mercury (Hg) consists of three operationally-defined forms including particle-bound Hg (PBM), gaseous oxidized Hg (GOM), and gaseous elemental Hg (GEM) (Selin, 2009). GEM is the most abundant (about 90%) and chemically stable form (Selin, 2009), and is transported at regional and global scales. GOM has short residence times as it can readily be dissolved in rain droplets, adsorbed on particulate matter (PM), and it reacts rapidly within both  
45 gaseous and aqueous phases with or without PM. PBM contains mainly reactive Hg species such as  $\text{Hg}^{2+}$  and perhaps trace quantities of  $\text{Hg}^0$ , and is transported at regional or local scales thereby reflecting Hg pollution and cycling within short distances from emission source (Selin, 2009; Subir et al., 2012). PBM has multiple sources and undergoes complex transport and transformation processes in the atmosphere (Subir et al., 2012).

50 This study aimed at characterizing the isotope compositions of  $\text{PM}_{2.5}\text{-Hg}$  (particulate matter with aerodynamic diameter less than 2.5 micrometers) to better understand the complex transformation processes of PBM. Hg has seven stable isotopes and is known to exhibit both mass-dependent (MDF, represented by  $\delta^{202}\text{Hg}$ ) and mass-independent (MIF, including odd-mass-number isotopic MIF (odd-MIF) and even-mass-number isotopic MIF (even-MIF),  
55 represented by  $\Delta^{199}\text{Hg}$ ,  $\Delta^{201}\text{Hg}$ ,  $\Delta^{200}\text{Hg}$  and  $\Delta^{204}\text{Hg}$ ) fractionation during Hg transformations under various environmental conditions (Hintelmann and Lu, 2003; Jackson et al., 2004; Bergquist and Blum, 2007; Jackson et al., 2008; Gratz et al., 2010; Chen et al., 2012; Sherman et al., 2012). Prior studies have shown that MDF can be induced by several Hg transformation and transport processes (Bergquist and Blum, 2007; Kritee et al., 2007; Estrade et al., 2009;  
60 Yang and Sturgeon, 2009; Sherman et al., 2010; Wiederhold et al., 2010; Ghosh et al., 2013; Smith et al., 2015; Janssen et al., 2016), but large extents of odd-MIF mainly occurred during photochemical reactions including photoreduction (Bergquist and Blum, 2007; Zheng and Hintelmann, 2009; Sherman et al., 2010; Zheng and Hintelmann, 2010b; Rose et al., 2015),

photodemethylation (Bergquist and Blum, 2007; Rose et al., 2015) and photooxidation (Sun et al., 2016). Smaller but measurable degrees of odd-MIF were also reported for nonphotochemical abiotic reduction (Zheng and Hintelmann, 2010a) and evaporation of  $\text{Hg}^0$  (Estrade et al., 2009; Ghosh et al., 2013). Interestingly, the results of laboratory and field investigations suggest that specific  $\Delta^{199}\text{Hg}/\Delta^{201}\text{Hg}$  ratios are associated with such transformation processes, with a ratio of about 1.0 for photoreduction of inorganic  $\text{Hg}^{2+}$  (Bergquist and Blum, 2007; Zheng and Hintelmann, 2009; Sherman et al., 2010; Zheng and Hintelmann, 2010b; Rose et al., 2015), about 1.3 for photodemethylation and 1.6 for  $\text{Hg}^0$  evaporation and photooxidation (Bergquist and Blum, 2007; Estrade et al., 2009; Zheng and Hintelmann, 2010a; Ghosh et al., 2013; Rose et al., 2015; Sun et al., 2016). Even-MIF of  $\text{Hg}$  isotope signatures were observed mostly in wet deposition, but the mechanism producing such fractionation remains unknown (Chen et al., 2012; Cai and Chen, 2016).

Prior studies have shown relatively constant  $\text{Hg}$  isotope compositions for GEM and very large variations of  $\text{Hg}$  isotope ratios for dissolved  $\text{Hg}^{2+}$  in wet precipitation (Gratz et al., 2010; Chen et al., 2012; Rolison et al., 2013; Wang et al., 2015; Yuan et al., 2015). A few studies reported that the  $\text{Hg}$  isotope compositions of PBM also show large variations (Rolison et al., 2013; Das et al., 2016; Huang et al., 2016; Yu et al., 2016; Xu et al., 2017). Among these limited studies, Rolison et al. (2013) reported  $\delta^{202}\text{Hg}$  ( $-1.61\text{‰}$  to  $-0.12\text{‰}$ ) and  $\Delta^{199}\text{Hg}$  values ( $0.36\text{‰}$  to  $1.36\text{‰}$ ) for  $\text{Hg}$  bound on total suspended particulates (TSP), with  $\Delta^{199}\text{Hg}/\Delta^{201}\text{Hg}$  ratios of approximately unity, a value typical of photoreduction of inorganic  $\text{Hg}^{2+}$ . Das et al. (2016) found values of  $\Delta^{199}\text{Hg}$  varied between  $-0.31\text{‰}$  and  $0.33\text{‰}$  for  $\text{PM}_{10}$  from Kolkata, eastern India. It was suggested that PBM with longer residence times may have undergone greater photoreduction and hence exhibited more positive MIF. Huang et al. (2016) investigated  $\text{Hg}$  isotope compositions for  $\text{PM}_{2.5}$  samples taken from Beijing, China, and attributed their observed seasonal variations in both MDF ( $\delta^{202}\text{Hg}$  from  $-2.18\text{‰}$  to  $0.51\text{‰}$ ) and MIF ( $\Delta^{199}\text{Hg}$

from  $-0.53\text{‰}$  to  $0.57\text{‰}$ ) to varied contributions from multiple sources of  $\text{PM}_{2.5}\text{-Hg}$ , while the  
90 more positive  $\Delta^{199}\text{Hg}$  values were likely produced by extensive photochemical reduction during  
long-range-transported. These prior results show that the Hg isotope approach can be employed  
for tracking sources and identifying possible transformation processes for airborne PM-Hg, and  
that PBM may undergo photochemical reactions that obscure its initial isotopic signature.

The goal of this study was to quantify short-term (diel) variations in the isotope composition  
95 of  $\text{PM}_{2.5}\text{-Hg}$  in an effort to elucidate if photochemical processes could impact overall contents  
and isotope compositions of PM-bound Hg in an urban environment. Unlike prior studies in  
which PM samples were collected continuously over 24 hrs or longer, we collected two  $\text{PM}_{2.5}$   
samples per 24 hrs with a daytime (D) sample between 8:00 a.m. and 6:30 p.m. and a nighttime  
(N) sample between 7:00 p.m. and 7:30 a.m. The specific objectives of this study were to verify  
100 and quantify whether Hg isotope compositions of  $\text{PM}_{2.5}$  exhibit diel variations, and to elucidate  
whether photochemical transformation is the dominant process for such diel variations.

## 2 Experimental section

### 2.1 Field site, sampling method, and preconcentration of $\text{PM}_{2.5}\text{-Hg}$

105 Beijing was selected as the area of study because of its well-known air pollution issue (Zhang  
et al., 2007). Detailed information on the study site and  $\text{PM}_{2.5}$  sampling procedures were given  
elsewhere (Huang et al., 2016). During the sampling period between Sept. 15<sup>th</sup> and Oct. 16<sup>th</sup>,  
2015, the average outdoor temperatures were  $22.1 \pm 3.0^\circ\text{C}$  and  $18.5 \pm 2.7^\circ\text{C}$ , and the average  
relative humidity were  $45 \pm 20\%$  and  $59 \pm 19\%$ , for D and N, respectively. The  $\text{PM}_{2.5}$  samples  
110 were collected using a Tisch Environmental  $\text{PM}_{2.5}$  high volume air sampler, which collects  
particles at a flow rate of  $1.0 \text{ m}^3 \text{ min}^{-1}$  through a  $\text{PM}_{2.5}$  size selective inlet on a pre-combusted  
( $450^\circ\text{C}$  for 6 hrs) quartz fiber filter (Pallflex 2500 QAT-UP,  $20 \text{ cm} \times 25 \text{ cm}$ , Pallflex Product  
Co., USA). Quartz fiber filters were widely used to collect operationally defined PBM

(Schleicher et al., 2015; Zhang et al., 2015; Xu et al., 2017). A total of 61 samples including 30  
115 D samples and 31 N samples were collected between 8:00 a.m. and 6:30 p.m. and 7:00 p.m. to  
7:30 a.m., respectively, along with 2 field blanks. They were wrapped with aluminum film,  
packed in plastic bags, and stored at  $-20^{\circ}\text{C}$  in the lab prior to analysis. Meteorological data,  
including temperature (T), relative humidity (RH), sunshine duration, daily average wind speed  
(WS), were acquired from China Meteorological Administration (<http://data.cma.cn>), and the  
120 atmospheric ozone content ( $P_{\text{O}_3}$ ) was measured concurrently. These data are summarized in  
Table S1.

## 2.2 Hg content and stable isotope measurements

The mass of each  $\text{PM}_{2.5}$  sample was gravimetrically quantified. Hg bound on each  $\text{PM}_{2.5}$  sample  
was extracted and concentrated for analysis of Hg content and stable Hg isotopes using the  
125 method reported previously (Huang et al., 2015). The details of the procedures are also given  
in Supporting Information (SI).

Among the 61  $\text{PM}_{2.5}$  samples, 56 (including 26 D- and 30 N-samples) had sufficient Hg mass  
( $> 10$  ng) and were further analyzed for Hg isotope compositions using a multicollector  
inductively coupled plasma mass spectrometer (MC-ICP-MS, Nu Instruments Ltd., UK)  
130 equipped with a continuous flow cold vapor generation system. Detailed protocols for the Hg  
isotope analysis can be found in Huang et al. (2015) and also in SI.  $^{196}\text{Hg}$  and  $^{204}\text{Hg}$  were not  
measured due to their very low abundance. Instrumental mass bias was corrected using an  
internal standard (NIST SRM 997 Tl) and strict sample-standard bracketing with NIST SRM  
3133 Hg standard. Delta ( $\delta$ ) notation is used to represent MDF in units of per mil (‰) as defined  
135 by the following equation (Blum and Bergquist, 2007):

$$\delta^x\text{Hg} (\text{‰}) = [({}^x\text{Hg}/{}^{198}\text{Hg})_{\text{sample}}/({}^x\text{Hg}/{}^{198}\text{Hg})_{\text{NIST3133}} - 1] \times 1000 \quad (1)$$

where  $x = 199, 200, 201,$  and  $202$ . MIF is reported as the deviation of a measured delta value  
from the theoretically predicted MDF value according to the equation:

$$\Delta^x\text{Hg} (\text{‰}) = \delta^x\text{Hg} - \beta \times \delta^{202}\text{Hg} \quad (2)$$


140 where the mass-dependent scaling factor  $\beta$  is 0.252, 0.5024, and 0.752 for  $^{199}\text{Hg}$ ,  $^{200}\text{Hg}$  and  $^{201}\text{Hg}$ , respectively (Blum and Bergquist, 2007).

For quality assurance and control, we used NIST SRM 3177 Hg as a secondary standard and analyzed repeatedly during sample analysis session. The collective measurements of the NIST 3177 standard yielded average  $\delta^{202}\text{Hg}$ ,  $\Delta^{199}\text{Hg}$ ,  $\Delta^{200}\text{Hg}$  and  $\Delta^{201}\text{Hg}$  values of  $-0.53\text{‰} \pm 0.09\text{‰}$ ,  $-0.01\text{‰} \pm 0.04\text{‰}$ ,  $0.00\text{‰} \pm 0.04\text{‰}$  and  $-0.01\text{‰} \pm 0.07\text{‰}$  (2SD,  $n = 17$ ). We also analyzed 145 regularly a well-known reference material UM-Almaden and a certified reference material (CRM) GBW07405, and the results showed average  $\delta^{202}\text{Hg}$ ,  $\Delta^{199}\text{Hg}$ ,  $\Delta^{200}\text{Hg}$  and  $\Delta^{201}\text{Hg}$  values of  $-0.60\text{‰} \pm 0.09\text{‰}$ ,  $-0.01\text{‰} \pm 0.04\text{‰}$ ,  $0.01\text{‰} \pm 0.04\text{‰}$  and  $-0.03\text{‰} \pm 0.07\text{‰}$  (2SD,  $n = 17$ ) and of  $-1.77\text{‰} \pm 0.14\text{‰}$ ,  $-0.29\text{‰} \pm 0.06\text{‰}$ ,  $0.00\text{‰} \pm 0.04\text{‰}$  and  $-0.32\text{‰} \pm 0.07\text{‰}$  (2SD,  $n = 155$  6), respectively. These values were consistent with previous results (Blum and Bergquist, 2007; Chen et al., 2010; Huang et al., 2015; Huang et al., 2016). The uncertainties of  $\text{PM}_{2.5}\text{-Hg}$  isotope ratios listed in Table S2 were calculated based on repetitive measurements. However, if uncertainty of the isotopic compositions for a given sample was smaller than the uncertainty of CRM GBW07405, the uncertainty associated with that sample was assigned 2SD uncertainties (0.14‰, 0.06‰, 0.04‰ and 0.07‰ for  $\delta^{202}\text{Hg}$ ,  $\Delta^{199}\text{Hg}$ ,  $\Delta^{200}\text{Hg}$  and  $\Delta^{201}\text{Hg}$ ) obtained for long-term measurement of the CRM GBW07405.

### 2.3 Air mass backward trajectories

To identify possible pathways of  $\text{PM}_{2.5}\text{-Hg}$  transport, backward HYSPLIT trajectories of air masses at a height of 500 m above ground level and arriving at the sampling site were simulated. 160 Backward trajectories for each D or N sample were calculated every 1 hrs using the Internet-Based HYSPLIT Trajectory Model and gridded meteorological data (Global Data Assimilation System, GDAS1) from the U.S. National Oceanic and Atmospheric Administration (NOAA) (Fig. S1). The obtained average directions of arriving air masses for each sample were

summarized in Table S1. The frequencies of backward trajectories were calculated for all the  
165 samples taken during Sept. 15<sup>th</sup> to Oct. 16<sup>th</sup> 2015 using the Internet-Based HYSPLIT Trajectory  
Model and the archived GDAS0p5, with an interval of 3 hrs. Each trajectory had a total run  
time of 72 hrs and a grid resolution of 0.5 × 0.5 degree trajectory frequency. The simulation  
results showed the dominant air mass was arriving from southwest of the sampling site during  
the sampling period (see Fig. 1).



170

**Figure 1.** Geographic location of the PM<sub>2.5</sub> collection site in Beijing, China (Baidu Map image) and average air mass back trajectories during sampling from Sept. 15<sup>th</sup> to Oct. 16<sup>th</sup>, 2015 (left), and characteristics of North-West vs. South-East arriving air masses.


## 2.4 Statistical analysis

175 T-test was performed for uncertainty analysis using IBM SPSS Statistics Version 22. Both paired samples *t*-test and independent samples *t*-test were performed for diel variations of Hg content,  $\delta^{202}\text{Hg}$ ,  $\Delta^{199}\text{Hg}$  and  $\Delta^{200}\text{Hg}$ , and their results were summarized in Table S3.

## 3 Results and discussion

### 180 3.1 Diel variation of PM<sub>2.5</sub>-Hg

The chronological sequence of Hg stable isotope ratios, along with weather conditions for the 56 PM<sub>2.5</sub> samples are presented in Figure 2 (see also Tables S1 and S2 for quantitative atmospheric data and  $\Delta^{201}\text{Hg}$  values). The major features of this dataset include: i) large variations in both MDF and odd-MIF of Hg isotopes, ii) significant diel differences in Hg isotope ratios, iii) correlations of weather conditions and air mass backward trajectories with Hg isotope signals, and iv) detectable even-MIF.



**Figure 2.** Chronological sequence of MIF ( $\Delta^{199}\text{Hg}$  and  $\Delta^{200}\text{Hg}$ ) and MDF ( $\delta^{202}\text{Hg}$ ) of the 56 samples collected during the daytime (D, red) and nighttime (N, blue), along with selected weather data including cumulative hours of sunshine (Solar) and air mass backward-trajectory directions.

The volumetric PM<sub>2.5</sub> concentrations ranged from 4 to 158  $\mu\text{g m}^{-3}$  with an average value of  $52 \pm 40 \mu\text{g m}^{-3}$  (1SD,  $n = 61$ ) and the highest values ( $> 100 \mu\text{g m}^{-3}$ ) detected during a severe

195 haze event of Oct. 4<sup>th</sup>-7<sup>th</sup>. The mass-based Hg contents ranged from 0.08 to 1.22  $\mu\text{g g}^{-1}$  with a mean value of  $0.40 \pm 0.21 \mu\text{g g}^{-1}$  (1SD,  $n = 61$ ).

Hg isotope analysis showed that  $\delta^{202}\text{Hg}$  values varied from  $-1.49\text{‰}$  to  $0.55\text{‰}$  (mean =  $-0.53\text{‰} \pm 0.40\text{‰}$ , 1SD,  $n = 56$ ), with the lowest value found in sample Oct-2-N and the highest in Oct-8-N. Significant odd-MIF of Hg isotopes was found and the  $\Delta^{199}\text{Hg}$  values ranged from  $-0.53\text{‰}$  to  $1.04\text{‰}$  (mean =  $0.14\text{‰} \pm 0.33\text{‰}$ ) with the lowest ( $-0.53\text{‰}$ )  $\Delta^{199}\text{Hg}$  value on Oct-5-D during the severe haze event and the highest ( $1.04\text{‰}$ )  $\Delta^{199}\text{Hg}$  value on Sept-26-D during a sunny day (without cloud) (Fig. 2). All samples also displayed slight even-MIF, with  $\Delta^{200}\text{Hg}$  values ranging from  $-0.02\text{‰}$  to  $0.21\text{‰}$  (average  $0.07\text{‰} \pm 0.06\text{‰}$ , 1SD,  $n = 56$ ), which were significant compared to the detection precision of  $\pm 0.04\text{‰}$ . The overall variations of Hg isotope ratios for these 12-hr D/N  $\text{PM}_{2.5}$  samples are generally consistent with several prior reports for the 24-hr PBM samples (Rolison et al., 2013; Das et al., 2016; Huang et al., 2016).

T-test results (Table S3) showed that diel variation was statistically significant ( $p < 0.05$ ) for Hg contents,  $\Delta^{199}\text{Hg}$ , and  $\Delta^{200}\text{Hg}$  values, as their  $p$  values are 0.005, 0.000 and 0.004 resulting from paired samples  $t$ -test, and are 0.003, 0.017 and 0.019 resulting from independent samples  $t$ -test. For all samples, Hg contents for D-samples ( $0.32 \pm 0.14 \mu\text{g g}^{-1}$ ) were lower than N-samples ( $0.48 \pm 0.24 \mu\text{g g}^{-1}$ ), and  $\Delta^{199}\text{Hg}$  and  $\Delta^{200}\text{Hg}$  values for D-samples (mean of  $0.26\text{‰} \pm 0.40\text{‰}$  and  $0.09\text{‰} \pm 0.06\text{‰}$ , respectively) were higher than N-samples ( $-0.04\text{‰} \pm 0.22\text{‰}$  and  $0.06\text{‰} \pm 0.05\text{‰}$ , respectively). However,  $\text{PM}_{2.5}$  concentrations and  $\delta^{202}\text{Hg}$  had statistically insignificant ( $p > 0.05$ ) diel variation, as their  $p$  values are 0.887 and 0.052 resulting from paired samples  $t$ -test, and are 0.909 and 0.053 resulting from independent samples  $t$ -test.

### 3.2 Diel variation in odd-MIF of $\text{PM}_{2.5}$ -Hg independent of air mass source

Many consecutive D-N sampling intervals had similar air mass back trajectories (Table S1 and Fig. S1), suggesting that the dominant sources of  $\text{PM}_{2.5}$ -Hg did not vary over each such 24 hr

220 sampling period. For example, pairs Sept-16-D and Sept-16-N, Sept-17-D and Sept-17-N, Sept-  
20-D and Sept-20-N, Sept-21-D and Sept-21-N, Oct-1-D and Oct-1-N, Oct-2-D and Oct-2-N,  
Oct-4-D and Oct-4-N have similar air mass trajectories from the southwest, and pairs Oct-8-D  
and Oct-8-N, Oct-9-D and Oct-9-N, Oct-10-D and Oct-10-N, Oct-11-D and Oct-11-N, Oct-12-  
D and Oct-12-N have similar air mass trajectories from the northwest and north (Fig. S1). It is  
225 reasonable to assume, therefore, that each of these D-N PM<sub>2.5</sub> sample pairs had identical  
dominant sources of PM<sub>2.5</sub>-Hg and to expect that they would have very similar Hg isotope  
compositions. Instead, however, the data presented in Table S2 and Figure 2 revealed a unique  
and consistent pattern of diel variation in Hg isotope ratios; specifically, each PM<sub>2.5</sub> D-sample  
had a statistically significantly higher positive  $\Delta^{199}\text{Hg}$  value (up to +1.04‰) than its consecutive  
230 PM<sub>2.5</sub> N-sample.

The more positive  $\Delta^{199}\text{Hg}$  values measured for the PM<sub>2.5</sub> D-samples are highly unlikely  
uncharacteristic of known emission sources of PM<sub>2.5</sub>-Hg. It is possible that PM<sub>2.5</sub>-Hg from  
different emission sources may have different Hg isotope compositions. However, prior studies  
showed that  $\Delta^{199}\text{Hg}$  values of the PBM from dominant anthropogenic emission sources are  
235 generally negative or close to zero. Schleicher et al. (2015) demonstrated that coal combustion  
is likely the major source of PM<sub>2.5</sub>-Hg in Beijing. Huang et al (2016) reported that regional  
anthropogenic activities such as coal combustion ( $\Delta^{199}\text{Hg}$  values from -0.30‰ to 0.05‰),  
metal smelting (-0.20‰ to -0.05‰) and cement production (-0.25‰ to 0.05‰), as well as  
biomass burning (low to -0.53‰) were likely the dominant sources of PM<sub>2.5</sub>-Hg at this study  
240 site. As shown in Figure 2 and Table S1, the PM<sub>2.5</sub> D-samples with high  $\Delta^{199}\text{Hg}$  values (>  
0.60‰) each had very different air mass back trajectories (Fig. S1). For instance, Sept-18-D  
(with  $\Delta^{199}\text{Hg}$  value +0.90‰), Sept-26-D (with  $\Delta^{199}\text{Hg}$  value +1.04‰) and Oct-3-D (with  
 $\Delta^{199}\text{Hg}$  value +0.86‰) were associated with north, southwest, and north-south mixed air  
masses, respectively. A reasonable explanation of these observations is that high positive


245  $\Delta^{199}\text{Hg}$  values measured for D-samples resulted from  $\text{PM}_{2.5}\text{-Hg}$  transformation, specifically photoreduction, during atmospheric transport. Indeed, the diel variation of  $\Delta^{199}\text{Hg}$  for  $\text{PM}_{2.5}$  D-N sample pairs may well reflect strong (D) versus less or no (N) influences of photochemical reactions under time-variant local and regional weather conditions.

### 3.3 Photochemical reduction as a cause of odd-MIF in subset of daytime $\text{PM}_{2.5}$ samples

250 To detail the effects of photochemical reactions on the variation of Hg isotope ratios for  $\text{PM}_{2.5}\text{-Hg}$ , we regrouped our dataset into subsets corresponding to day and night. We further regrouped our results into two source-related subsets, south-east (S-E) and north-west (N-W), according to the air mass backward trajectories during each sampling event (Fig. 1), and two other subsets corresponding to sunny days within the S-E group (sunS-E) and all sunny days (Sun), which  
255 includes sunS-E and N-W as N-W consisted entirely of sunny days. The N-W subset of  $\text{PM}_{2.5}$  was associated with an air mass that tracked from the north, northeast, northwest and west, relatively less polluted areas, and is therefore representative of long-range transport and relatively constant sources of  $\text{PM}_{2.5}$  and Hg (Huang et al., 2016). The S-E subset was associated with an air mass that tracked from the south, southwest, southeast and east, heavily polluted  
260 and highly populated areas, and was characterized by relatively high contents of  $\text{PM}_{2.5}$ , likely from industrial sources in the region (coal fired power plants, coking and steel industries). Unlike the N-W arriving air mass which corresponded to all sunny days during the entire sampling period, the S-E arriving air mass was associated with a range of weather conditions including hazy, cloudy, rainy, and sunny days. According to our results (Table S2 and Fig. 2),  
265  $\text{PM}_{2.5}$  concentrations of the N-W subset ( $23 \pm 19 \mu\text{g m}^{-3}$ ) were significantly ( $p < 0.05$ ) lower than the S-E subset ( $69 \pm 40 \mu\text{g m}^{-3}$ ), which is consistent with the fact that the N-W areas of Beijing were less industrialized, less populated and less polluted than the S-E areas. However, regardless of whether their associated air masses originated from moderately or heavily polluted areas, both N-W and S-E subset samples showed diel variations in their Hg contents and isotope


270 ratios (see Fig. 3 and discussion below). This, as discussed above, indicates that air mass source was not a dominant factor producing the diel variation of Hg isotope ratios in consecutive D-N PM<sub>2.5</sub> samples.

The observed diel difference in  $\Delta^{199}\text{Hg}$  values of PM<sub>2.5</sub>-Hg is even more prominent and statistically robust within subsets of PM<sub>2.5</sub> samples regrouped according to their air mass trajectories (i.e., PM<sub>2.5</sub> source related) and sunny days (with greater extent of photochemical reactions). As shown in Figure 3,  $\Delta^{199}\text{Hg}$  values for N-W subset samples collected during the day had a higher range (0.04‰ to 0.90‰) and mean (0.39‰  $\pm$  0.27‰ SD,  $n = 10$ ) compared to those (-0.07‰ to 0.32‰, mean = 0.09‰  $\pm$  0.13‰ SD,  $n = 9$ ) for N-samples ( $p = 0.02$ ). Similarly, analysis of the sunS-E subset revealed a significant difference in  $\Delta^{199}\text{Hg}$  values ( $p = 0.03$ ) between sunny days and nights, but not when the entire S-E sample set ( $p = 0.22$ ), which includes hazy, raining, and cloudy days, was considered. Since the N-W subset was associated with less polluted areas and the S-E subset was associated with heavily polluted and highly populated areas, the observation of significant diel variation of  $\Delta^{199}\text{Hg}$  in PM<sub>2.5</sub>-Hg within each subset (Fig. 3) is consistent with the above conclusion that such variation of PM<sub>2.5</sub>-Hg isotope ratios was not controlled by variation of Hg emission sources. The highly positive  $\Delta^{199}\text{Hg}$  values observed for daytime samples within the Sun subset (Fig. S2) further supports the conclusion that PM<sub>2.5</sub>-Hg was strongly affected by photochemical reactions on sunny days.



**Figure 3.** Diel variations of  $\Delta^{199}\text{Hg}$  for  $\text{PM}_{2.5}\text{-Hg}$  for the entire dataset, the N-W subset, the S-E subset, and the S-E sunny days only subset. Note that all days included in the N-W subset were sunny. Diel differences within each subset were examined using the Independent Samples T-Test.

Linear correlations of  $\Delta^{199}\text{Hg}$  versus  $\Delta^{201}\text{Hg}$  for all 56  $\text{PM}_{2.5}$  samples (Fig. 4a) and three subsets, N-W (Fig. 4b), S-E (Fig. 4c), and Sun (Fig. 4d), yielded slopes of  $1.06 \pm 0.05$  (1SD,  $r^2 = 0.89$ ),  $1.06 \pm 0.12$  ( $r^2 = 0.81$ ),  $1.13 \pm 0.05$  ( $r^2 = 0.92$ ) and  $1.13 \pm 0.08$  ( $r^2 = 0.84$ ), respectively (Fig. 4). Such slopes are all indicative of photochemical reduction of  $\text{Hg}^{2+}$  according to prior studies (Bergquist and Blum, 2007; Zheng and Hintelmann, 2009). The photoreduction process is further evidenced by a progressive increase in  $\Delta^{199}\text{Hg}$  from zero or slightly negative values to positive values as the content of Hg in  $\text{PM}_{2.5}$  ( $C_{\text{Hg}}$ ) decreased in D-samples (Fig. S1a). This trend is statistically more significant ( $p < 0.05$ ) for D-samples within the N-W and Sun subsets (Figs. S1-b and -d). Similarly, for all sunny day samples, a positive correlation ( $p < 0.05$ ) was also observed between  $\Delta^{199}\text{Hg}$  and  $\delta^{202}\text{Hg}$  (Fig. S3), consistent with prior experimental results (Bergquist and Blum, 2007; Zheng and Hintelmann, 2009). Collectively, the Hg isotope results suggest that photochemical reduction is an important process during the transport of  $\text{PM}_{2.5}\text{-Hg}$  in the atmosphere.



**Figure 4.** Correlations between  $\Delta^{199}\text{Hg}$  and  $\Delta^{201}\text{Hg}$  for different subsets of  $\text{PM}_{2.5}$  samples: a) all data, b) North-West (N-W), c) South-East (S-E) and d) All sunny days (Sun). The slope, intercept and  $r$ -square of the line from simple linear regression. Vertical and horizontal error bars correspond to 2SD analytical precision.

Among all D-samples,  $\Delta^{199}\text{Hg}$  is only weakly correlated with sunshine duration ( $r^2 = 0.20$ ,  $p = 0.02$ ). However,  $\Delta^{199}\text{Hg}$  values for all D-samples collected on days with sunshine durations  $>8$  hrs are positive whereas half of the  $\Delta^{199}\text{Hg}$  values for samples collected on cloudy or hazy days with shorter sunshine durations are negative or near zero (Fig. S4). In addition, a significant positive linear correlation between  $\Delta^{199}\text{Hg}$  values and atmospheric ozone contents ( $P_{\text{O}_3}$ ) ( $r^2 = 0.517$ ,  $p < 0.01$ ) for all but four daytime samples was obtained (Fig. S5). The four outliers (Sept-16-D, Sept-17-D, Oct-5-D and Oct-6-D) were collected on days with high ozone ( $P_{\text{O}_3}$  above 50 ppbv) and severe smog formation. Conversely, no significant correlation ( $p > 0.05$ ) between  $\Delta^{199}\text{Hg}$  and  $P_{\text{O}_3}$  was found for the nighttime samples.

The increase in  $\Delta^{199}\text{Hg}$  of daytime  $\text{PM}_{2.5}\text{-Hg}$  with sunlight duration and ozone concentration indicates that the physical and photochemical conditions of the atmosphere may affect the atmospheric transformation of  $\text{PM}_{2.5}\text{-Hg}$ . A prior experimental study showed that GEM oxidation can produce negative  $\Delta^{199}\text{Hg}$  values in oxidized  $\text{Hg}^{2+}$  with  $\Delta^{199}\text{Hg}/\Delta^{201}\text{Hg}$  ratios of 1.6 and 1.9 for Br and Cl radical initiated oxidations (Sun et al., 2016). We can exclude the possible contribution of  $\text{Hg}^0$  oxidation to  $\text{PM}_{2.5}\text{-Hg}$ , given the fact that  $\Delta^{199}\text{Hg}/\Delta^{201}\text{Hg}$  ratio was about 1.1 and most  $\text{PM}_{2.5}\text{-Hg}$  samples collected during the daytime when Br and Cl radicals could form had positive  $\Delta^{199}\text{Hg}$  values. Thus it is highly unlikely that oxidation would have caused the diel variation in Hg isotopes in  $\text{PM}_{2.5}$ . However, the exception to the observed relationships between  $\Delta^{199}\text{Hg}$  with sunlight duration, and ozone concentration show that in a highly oxidizing atmosphere (higher  $P_{\text{O}_3}$ ) such as occurs during extreme smog events, the odd-MIF of Hg isotopes in  $\text{PM}_{2.5}$  may decrease or reverse. A possible explanation for this effect

may be the increased production of GOM and its collection with PM<sub>2.5</sub>-Hg during such smog  
335 events. While PM<sub>2.5</sub>-Hg samples collected on quartz fiber filters may include some GOM  
(Lynam and Keeler, 2002), this contribution was likely small in most of our D and N samples  
due to the opposing diel trends in the concentrations of PBM and GOM in urban air (Engle et  
al., 2010). GOM would therefore not have had a major effect on the observed diel variations of  
 $\Delta^{199}\text{Hg}$  values for PM<sub>2.5</sub>-Hg and may have in fact masked an even larger MIF signature due to  
340 the photoreduction of PBM during the day.

Interestingly, negative  $\Delta^{199}\text{Hg}$  values in daytime PM<sub>2.5</sub>-Hg were only observed during a rainy  
day and an extreme smog event. Since the Hg emitted from local sources had close to zero and  
negative values of odd-MIF, higher humidity (such as during rainy days) and heavy pollution  
(the extreme smog) may enhance the effect of scavenging of locally produced gaseous or  
345 particulate Hg during rain or smog events, which may therefore have contributed to the reversal  
of the odd-MIF signature of Hg collected as PM<sub>2.5</sub> at these times. In addition, the negative  
 $\Delta^{199}\text{Hg}$  values in PM<sub>2.5</sub> may have resulted from the contribution of biomass burning with limited  
photoreduction effect during periods of less sunshine (Fig. 2 and Table S1) since plant foliage  
has negative  $\Delta^{199}\text{Hg}$  values (Yu et al., 2016) and more negative  $\Delta^{199}\text{Hg}$  values (down to  $-0.53\text{‰}$ )  
350 of PM<sub>2.5</sub>-Hg in Beijing were related to biomass burning, a source of PM<sub>2.5</sub>-Hg south of Beijing  
in autumn (Huang et al., 2016). This could further explain the relatively lower  $\Delta^{199}\text{Hg}$  values  
in the majority of the N-samples (for example, Sept-28-N and Oct-5-N with  $\Delta^{199}\text{Hg}$  of  $-0.46\text{‰}$   
and  $-0.51\text{‰}$ ), even in those collected under clear weathering condition. Indeed, each bulk  
sample collected during night time was a mixture of the leftover PM<sub>2.5</sub> (with positive odd-MIF)  
355 from the previous daytime and the PM<sub>2.5</sub> newly input from various sources including industrial  
emissions (with close to zero  $\Delta^{199}\text{Hg}$ ) and biomass burning (somewhat negative  $\Delta^{199}\text{Hg}$ )  
(Huang et al., 2016) during nighttime.

A possible explanation of the observed effects of diel variation of PM<sub>2.5</sub>-Hg would be the temperature-dependent gas-aerosol partitioning of GOM (Rutter and Schauer, 2007; Amos et al., 2012), which favors more adsorption of GOM on PM during nighttime when atmospheric temperature is relatively lower than daytime. However, the magnitude of such adsorption is also proportional to the GOM concentration in the atmosphere. An inverse calculation exercise (in SI) shows that the higher PM<sub>2.5</sub>-Hg measured for our samples would require higher GOM concentrations during the nighttime, which contradicts with prior findings that GOM concentrations are significantly lower during the nighttime than the daytime as GOM is a product of photo-oxidation processes (Poissant et al., 2005; Liu et al., 2007; Amos et al., 2012). In addition, GOM gas-aerosol partitioning is considered a chemisorption and desorption process (Rutter and Schauer, 2007), which unlikely result in appreciable odd-MIF of Hg isotopes (Jiskra et al., 2012; Smith et al., 2015). Therefore, GOM partitioning would have little or no effect on the observed diel variations of  $\Delta^{199}\text{Hg}$  values for PM<sub>2.5</sub>-Hg.


Variation in atmosphere boundary layer height (ABLH) from 1000 to 1300 m during daytime to less than 200 to 300 m during nighttime may have contributed to the diel variation in Hg isotopic composition of PM<sub>2.5</sub>-Hg (Quan et al., 2013). With a high ABLH during daytime, relatively strong turbulence may help mixing the PM<sub>2.5</sub>-Hg from the surface to the upper free troposphere, where photoreactions may be favored due to higher intensities of ultraviolet radiation on clear days. In contrast, a lower ABLH at night may weaken the vertical transport of PM<sub>2.5</sub>-Hg, but enhance the contribution from newly produced PM<sub>2.5</sub>-Hg, possibly resulting in higher concentrations of PM<sub>2.5</sub>-Hg with negative or close to zero  $\Delta^{199}\text{Hg}$  values from emission sources and/or GOM. However, vertically-resolved, day-night measurements of Hg stable isotope ratios in PBM and GOM are needed to fully evaluate the effects of various physical processes on diel variation of the Hg isotopic compositions for the PM<sub>2.5</sub>.

While our results cannot exclude the effects of other possible processes, such as oxidation, adsorption (and desorption)/gas-aerosol partitioning, and precipitation, based on the limited previous studies (Jiskra et al., 2012; Smith et al., 2015; Sun et al., 2016), these processes are not likely to be important to the diel variation of odd-MIF of Hg isotopes in PM<sub>2.5</sub>-Hg we observed.

### 3.4 Photochemical reduction as a cause of diel variation in odd-MIF in day-night sample pairs of PM<sub>2.5</sub>

To explore the possible causes of diel variation in odd-MIF of Beijing PM<sub>2.5</sub>-Hg further, we examined four subgroups of PM<sub>2.5</sub> samples, each of which included 2 to 4 consecutive pairs of D-N samples that were collected during time periods of relatively constant atmospheric conditions (i.e., not being hazy, rainy or windy or having extremely high ozone ( $P_{O_3}$  above 50 ppbv)). Within each of the four subgroups,  $\Delta^{199}\text{Hg}$  and  $\delta^{202}\text{Hg}$  values were lower at night than during the previous or following day (Figs. 5-a and -b). As shown in Figure 5c, there is a significant positive correlation ( $p < 0.01$ ) between  $\Delta^{199}\text{Hg}$  and  $\delta^{202}\text{Hg}$  values for all samples in these four subgroups, with average values for both day and night falling right on the best fit line. The slope of this line is  $1.15 \pm 0.33$ , which is consistent with the reported value of  $1.15 \pm 0.07$  for photochemical reduction of  $\text{Hg}^{2+}$  in aqueous solution (Bergquist and Blum, 2007). Coincidentally, the contents of Hg in PM<sub>2.5</sub> ( $C_{\text{Hg}}$ ) were higher in N-samples than in immediately preceding or following D-samples (Fig. 5d), indicating a negative linear relationship between  $\Delta^{199}\text{Hg}$  values and  $C_{\text{Hg}}$  (Fig. 5e). Moreover, eight of the total eleven daytime samples among the four subgroups showed a positive linear correlation between  $\Delta^{199}\text{Hg}$  and the total cumulative daily solar radiation on a horizontal surface (SH) (Fig. 5f). These eleven samples also showed a negative correlation between the logarithmic values of  $C_{\text{Hg}}$  and SH (Fig. 5g). These correlations are consistent with the photochemical reduction of divalent Hg observed under laboratory conditions, and thus strongly support the hypothesis that photochemical

reduction is an important process controlling the fate of ambient atmospheric PM<sub>2.5</sub>-Hg. Given its diel trend and relatively large range, the MIF of odd Hg isotopes in Beijing PM<sub>2.5</sub> we observed was most likely due to the magnetic isotope effect (MIE), which has been invoked to explain MIF during the photochemical reduction of aqueous Hg<sup>2+</sup> (Zheng and Hintelmann, 2010b).



**Figure 5.** Hg isotope ratios and contents in four subgroups of consecutive pairs of day-night samples collected during periods of relatively constant atmospheric conditions. Linear correlations between  $\Delta^{199}\text{Hg}$  and  $\delta^{202}\text{Hg}$  (c),  $C_{\text{Hg}}$  (e) and the total cumulative daily solar radiation on a horizontal surface (SH, MJ m<sup>-2</sup>) (f), and between  $C_{\text{Hg}}$  and SH (g) were displayed. The uncertainty for measurement of  $\Delta^{199}\text{Hg}$  and  $\delta^{202}\text{Hg}$  of PM<sub>2.5</sub> samples were 0.06‰ and 0.12‰ in 2SD, respectively.

### 3.5 Even isotope MIF

Diel variation of  $\Delta^{200}\text{Hg}$  signatures was also observed in PM<sub>2.5</sub>-Hg. Prior studies reported mainly positive  $\Delta^{200}\text{Hg}$  values for wet precipitation (up to 1.24‰) (Gratz et al., 2010; Chen et al., 2012; Wang et al., 2015; Yuan et al., 2015) and aerosols (-0.05‰ to 0.28‰) (Rolison et al., 2013; Das et al., 2016; Huang et al., 2016). Even Hg isotope MIF has been attributed to

425 complex redox processes occurring in the upper atmosphere, but the underlying mechanisms remain unclear (Mead et al., 2013; Eiler et al., 2014). Thus, the even-MIF signatures may suggest a small proportion of PM<sub>2.5</sub>-Hg is likely originated from the upper atmosphere, through for example long-term transport and/or ABLH increasing during daytime. Given the fact that all samples displayed only slightly positive  $\Delta^{200}\text{Hg}$  (average  $0.07\text{‰} \pm 0.06\text{‰}$ , 1SD), this  
430 contribution may be very limited. In addition,  $\Delta^{200}\text{Hg}$  values are weakly, but significantly correlated with  $\Delta^{199}\text{Hg}$  ( $r^2 = 0.13$ ,  $p < 0.01$ ) (Fig. S6) and  $\delta^{202}\text{Hg}$  ( $r^2 = 0.27$ ,  $p < 0.01$ ) (Fig. S7). No mechanistic explanation is available yet for such observations, however.

#### 4 Conclusions

435 This study showed significant diel variations of Hg isotopic compositions for ambient PM<sub>2.5</sub>-Hg collected in the city of Beijing. The Hg isotope signatures featured a large range of MDF ( $\delta^{202}\text{Hg}$  value from  $-1.49\text{‰}$  to  $0.55\text{‰}$ , mean of  $-0.53\text{‰} \pm 0.40\text{‰}$ ) and significant ( $p < 0.05$ ) MIF with more positive  $\Delta^{199}\text{Hg}$  values in daytime samples ( $0.26\text{‰} \pm 0.40\text{‰}$ ) than at night ( $0.04\text{‰} \pm 0.22\text{‰}$ ). The results clearly indicated that the Hg isotope compositions of PM<sub>2.5</sub>-Hg are  
440 impacted variously by both weather conditions (such as sunlight duration), which may promote the photochemical reaction, and directions of air mass trajectories, which are related to possible sources of PM<sub>2.5</sub>. D-N paired samples having similar air mass backward trajectories and hence similar sources exhibited strong positive correlations between  $\Delta^{199}\text{Hg}$  and  $\Delta^{201}\text{Hg}$  with a slope of 1.1 and  $\Delta^{199}\text{Hg}$  and  $\delta^{202}\text{Hg}$  with a slope of 1.15, and a decrease in the content of Hg in PM<sub>2.5</sub>  
445 as  $\Delta^{199}\text{Hg}$  increased. These results provide isotopic evidence that local, daytime photochemical reduction of divalent Hg is of critical importance to the fate of PM<sub>2.5</sub>-Hg in urban atmosphere. Although the specific reactions and mechanisms that control Hg isotope fractionation (MDF and MIF) in Beijing PM<sub>2.5</sub> could not be explicitly determined from this field study, our result illustrated that, in addition to variation in sources, photochemical reduction appears to be an

450 important process that affects both the content and isotopic composition of PM<sub>2.5</sub>-Hg. Further systematic study is thus needed to better quantify the photoreduction of PM<sub>2.5</sub>-Hg to estimate the percentage of reduced Hg it produces and its impact on the global biogeochemical cycling of Hg.

455 *Acknowledgments.* We thank the anonymous reviewers for their constructive comments and suggestions. This study was supported financially by the National Key Research and Development Program of China (No. 2017YFC0212702), Natural Science Foundation of China (no. 41701268, 41625012, 41273023), Guizhou Scientific Research Program (No. 20161158) and State Key Laboratory of Organic Geochemistry (OGL-201501).

460

## References

- Amos, H. M., Jacob, D. J., Holmes, C. D., Fisher, J. A., Wang, Q., Yantosca, R. M., Corbitt, E. S., Galarneau, E., Rutter, A. P., Gustin, M. S., Steffen, A., Schauer, J. J., Graydon, J. A., St Louis, V. L., Talbot, R. W., Edgerton, E. S., Zhang, Y., and Sunderland, E. M.: Gas-particle partitioning of atmospheric Hg(II) and its effect on global mercury deposition, *Atmos. Chem. Phys.*, 12, 591-603, 2012.
- 465 Bergquist, B. A., and Blum, J. D.: Mass-dependent and -independent fractionation of Hg isotopes by photoreduction in aquatic systems, *Science*, 318, 417-420, 2007.
- Blum, J. D., and Bergquist, B. A.: Reporting of variations in the natural isotopic composition of mercury, *Anal. Bioanal. Chem.*, 388, 353-359, 2007.
- 470 Cai, H., and Chen, J.: Mass-independent fractionation of even mercury isotopes, *Science Bulletin*, 61, 116-124, 10.1007/s11434-015-0968-8, 2016.
- Chen, J. B., Hintelmann, H., and Dimock, B.: Chromatographic pre-concentration of Hg from dilute aqueous solutions for isotopic measurement by MC-ICP-MS, *J. Anal. At. Spectrom.*, 25, 1402-1409, 2010.

- Chen, J. B., Hintelmann, H., Feng, X. B., and Dimock, B.: Unusual fractionation of both odd and even mercury isotopes in precipitation from Peterborough, ON, Canada, *Geochim. Cosmochim. Acta*, 90, 33-46, 2012.
- Das, R., Wang, X., Khezri, B., Webster, R. D., Sikdar, P. K., and Datta, S.: Mercury isotopes of atmospheric particle bound mercury for source apportionment study in urban Kolkata, India, *Elementa: Science of the Anthropocene*, 4, 000098, 2016.
- Eiler, J. M., Bergquist, B., Bourg, I., Cartigny, P., Farquhar, J., Gagnon, A., Guo, W., Halevy, I., Hofmann, A., Larson, T. E., Levin, N., Schauble, E. A., and Stolper, D.: Frontiers of stable isotope geoscience, *Chem. Geol.*, 372, 119-143, 2014.
- Engle, M. A., Tate, M. T., Krabbenhoft, D. P., Schauer, J. J., Kolker, A., Shanley, J. B., and Bothner, M. H.: Comparison of atmospheric mercury speciation and deposition at nine sites across central and eastern North America, *J. Geophys. Res. Atmos.*, 115, 2010.
- Estrade, N., Carignan, J., Sonke, J. E., and Donard, O. F. X.: Mercury isotope fractionation during liquid-vapor evaporation experiments, *Geochim. Cosmochim. Acta*, 73, 2693-2711, 2009.
- Ghosh, S., Schauble, E. A., Lacrampe Couloume, G., Blum, J. D., and Bergquist, B. A.: Estimation of nuclear volume dependent fractionation of mercury isotopes in equilibrium liquid–vapor evaporation experiments, *Chem. Geol.*, 336, 5-12, 2013.
- Gratz, L. E., Keeler, G. J., Blum, J. D., and Sherman, L. S.: Isotopic composition and fractionation of mercury in Great Lakes precipitation and ambient air, *Environ. Sci. Technol.*, 44, 7764-7770, 2010.
- Hintelmann, H., and Lu, S. Y.: High precision isotope ratio measurements of mercury isotopes in cinnabar ores using multi-collector inductively coupled plasma mass spectrometry, *Analyst*, 128, 635-639, 2003.
- Huang, Q., Liu, Y. L., Chen, J. B., Feng, X. B., Huang, W. L., Yuan, S. L., Cai, H. M., and Fu, X. W.: An improved dual-stage protocol to pre-concentrate mercury from airborne particles for precise isotopic measurement, *J. Anal. At. Spectrom.*, 30, 957-966, 2015.
- Huang, Q., Chen, J., Huang, W., Fu, P., Guinot, B., Feng, X., Shang, L., Wang, Z., Wang, Z., Yuan, S., Cai, H., Wei, L., and Yu, B.: Isotopic composition for source identification of mercury in atmospheric fine particles, *Atmos. Chem. Phys.*, 16, 11773-11786, 2016.

- Jackson, T. A., Muir, D. C. G., and Vincent, W. F.: Historical variations in the stable isotope composition of mercury in Arctic Lake sediments, *Environ. Sci. Technol.*, 38, 2813-2821, 2004.
- Jackson, T. A., Whittle, D. M., Evans, M. S., and Muir, D. C. G.: Evidence for mass-independent and mass-dependent fractionation of the stable isotopes of mercury by natural processes in aquatic ecosystems, *Appl. Geochem.*, 23, 547-571, 2008.
- Janssen, S. E., Schaefer, J. K., Barkay, T., and Reinfelder, J. R.: Fractionation of mercury stable isotopes during microbial methylmercury production by iron- and sulfate-reducing bacteria, *Environ. Sci. Technol.*, 50, 8077-8083, 2016.
- Jiskra, M., Wiederhold, J. G., Bourdon, B., and Kretzschmar, R.: Solution speciation controls mercury isotope fractionation of Hg(II) sorption to goethite, *Environ. Sci. Technol.*, 46, 6654-6662, 2012.
- Kritee, K., Blum, J. D., Johnson, M. W., Bergquist, B. A., and Barkay, T.: Mercury stable isotope fractionation during reduction of Hg(II) to Hg(0) by mercury resistant microorganisms, *Environ. Sci. Technol.*, 41, 1889-1895, 2007.
- Liu, B., Keeler, G. J., Dvonch, J. T., Barres, J. A., Lynam, M. M., Marsik, F. J., and Morgan, J. T.: Temporal variability of mercury speciation in urban air, *Atmos. Environ.*, 41, 1911-1923, 2007.
- Lynam, M. M., and Keeler, G. J.: Comparison of methods for particulate phase mercury analysis: Sampling and analysis, *Anal. Bioanal. Chem.*, 374, 1009-1014, 2002.
- Mead, C., Lyons, J. R., Johnson, T. M., and Anbar, A. D.: Unique Hg stable isotope signatures of compact fluorescent lamp-sourced Hg, *Environ. Sci. Technol.*, 47, 2542-2547, 2013.
- Poissant, L., Pilote, M., Beauvais, C., Constant, P., and Zhang, H. H.: A year of continuous measurements of three atmospheric mercury species (GEM, RGM and Hg-p) in southern Quebec, Canada, *Atmos. Environ.*, 39, 1275-1287, 2005.
- Quan, J. N., Gao, Y., Zhang, Q., Tie, X. X., Cao, J. J., Han, S. Q., Meng, J. W., Chen, P. F., and Zhao, D. L.: Evolution of planetary boundary layer under different weather conditions, and its impact on aerosol concentrations, *Particuology*, 11, 34-40, 2013.

- Rolison, J. M., Landing, W. M., Luke, W., Cohen, M., and Salters, V. J. M.: Isotopic  
535 composition of species-specific atmospheric Hg in a coastal environment, *Chem. Geol.*, 336,  
37-49, 2013.
- Rose, C. H., Ghosh, S., Blum, J. D., and Bergquist, B. A.: Effects of ultraviolet radiation on  
mercury isotope fractionation during photo-reduction for inorganic and organic mercury  
species, *Chem. Geol.*, 405, 102-111, 2015.
- 540 Rutter, A. P., and Schauer, J. J.: The effect of temperature on the gas-particle partitioning of  
reactive mercury in atmospheric aerosols, *Atmos. Environ.*, 41, 8647-8657, 2007.
- Schleicher, N. J., Schäfer, J., Blanc, G., Chen, Y., Chai, F., Cen, K., and Norra, S.: Atmospheric  
particulate mercury in the megacity Beijing: Spatio-temporal variations and source  
apportionment, *Atmos. Environ.*, 109, 251-261, 2015.
- 545 Selin, N. E.: Global biogeochemical cycling of mercury: A review, *Annu. Rev. Env. Resour.*,  
34, 43-63, 2009.
- Sherman, L. S., Blum, J. D., Johnson, K. P., Keeler, G. J., Barres, J. A., and Douglas, T. A.:  
Mass-independent fractionation of mercury isotopes in Arctic snow driven by sunlight,  
*Nature Geosci.*, 3, 173-177, 2010.
- 550 Sherman, L. S., Blum, J. D., Keeler, G. J., Demers, J. D., and Dvonch, J. T.: Investigation of  
local mercury deposition from a coal-fired power plant using mercury isotopes, *Environ. Sci.  
Technol.*, 46, 382-390, 2012.
- Smith, R. S., Wiederhold, J. G., and Kretzschmar, R.: Mercury isotope fractionation during  
precipitation of metacinnabar ( $\beta$ -HgS) and montroydite (HgO), *Environ. Sci. Technol.*, 49,  
555 4325-4334, 2015.
- Subir, M., Ariya, P. A., and Dastoor, A. P.: A review of the sources of uncertainties in  
atmospheric mercury modeling II. Mercury surface and heterogeneous chemistry-A missing  
link, *Atmos. Environ.*, 46, 1-10, 2012.
- Sun, G., Sommar, J., Feng, X., Lin, C.-J., Ge, M., Wang, W., Yin, R., Fu, X., and Shang, L.:  
560 Mass-dependent and -independent fractionation of mercury isotope during gas-phase  
oxidation of elemental mercury vapor by atomic Cl and Br, *Environ. Sci. Technol.*, 50, 9232-  
9241, 2016.

- Wang, Z., Chen, J., Feng, X., Hintelmann, H., Yuan, S., Cai, H., Huang, Q., Wang, S., and Wang, F.: Mass-dependent and mass-independent fractionation of mercury isotopes in precipitation from Guiyang, SW China, *C. R. Geosci.*, 347, 358-367, 2015.
- Wiederhold, J. G., Cramer, C. J., Daniel, K., Infante, I., Bourdon, B., and Kretzschmar, R.: Equilibrium mercury isotope fractionation between dissolved Hg(II) species and thiol-bound Hg, *Environ. Sci. Technol.*, 44, 4191-4197, 2010.
- Xu, H., Sonke, J. E., Guinot, B., Fu, X., Sun, R., Lanzanova, A., Candaudap, F., Shen, Z., and Cao, J.: Seasonal and annual variations in atmospheric Hg and Pb isotopes in Xi'an, China, *Environ. Sci. Technol.*, 51, 3759-3766, 2017.
- Yang, L., and Sturgeon, R.: Isotopic fractionation of mercury induced by reduction and ethylation, *Anal. Bioanal. Chem.*, 393, 377-385, 2009.
- Yu, B., Fu, X., Yin, R., Zhang, H., Wang, X., Lin, C.-J., Wu, C., Zhang, Y., He, N., Fu, P., Wang, Z., Shang, L., Sommar, J., Sonke, J. E., Maurice, L., Guinot, B., and Feng, X.: Isotopic composition of atmospheric mercury in China: New evidence for sources and transformation processes in air and in vegetation, *Environ. Sci. Technol.*, 50, 9262-9269, 2016.
- Yuan, S., Zhang, Y., Chen, J., Kang, S., Zhang, J., Feng, X., Cai, H., Wang, Z., Wang, Z., and Huang, Q.: Large variation of mercury isotope composition during a single precipitation event at Lhasa city, Tibetan Plateau, China, *Procedia Earth and Planetary Science*, 13, 282-286, 2015.
- Zhang, L., Wang, S. X., Wang, L., Wu, Y., Duan, L., Wu, Q. R., Wang, F. Y., Yang, M., Yang, H., Hao, J. M., and Liu, X.: Updated emission inventories for speciated atmospheric mercury from anthropogenic sources in China, *Environ. Sci. Technol.*, 49, 3185-3194, 2015.
- Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimojo, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M., and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophys. Res. Lett.*, 34, 2007.

- Zheng, W., and Hintelmann, H.: Mercury isotope fractionation during photoreduction in natural water is controlled by its Hg/DOC ratio, *Geochim. Cosmochim. Acta*, 73, 6704-6715, 2009.
- 595 Zheng, W., and Hintelmann, H.: Nuclear field shift effect in isotope fractionation of mercury during abiotic reduction in the absence of light, *J. Phys. Chem. A*, 114, 4238-4245, 2010a.
- Zheng, W., and Hintelmann, H.: Isotope fractionation of mercury during its photochemical reduction by low-molecular-weight organic compounds, *J. Phys. Chem. A*, 114, 4246-4253, 2010b.

600