We thank the reviewer for the helpful comments, which certainly improved the manuscript. Especially, by evaluating the different spatial resolutions of the observations and the simulations and associated effects we are sure that the manuscript became more meaningful. The detailed replies on the reviewer's comments are structured as follows. Reviewer comments have bold letters, are labeled, and listed always in the beginning of each answer followed by the author's comments including (if necessary) major revised parts of the manuscript. The revised parts of the manuscript are written in quotation marks and italic letters. Minor revisions of the text can be found in the additionally submitted mark-up file.

1. The real problem of the study is insufficient resolution of the numerical simulations to effectively match the experimental data. The majority of efforts is set to averaging of the experimental data, which allows produce optical thickness fields of the resolution comparable to the model output due to limited model domain and poor resolution of the simulations. This causes that conclusions are weak and too far going.

The reviewer is right that the differences between the original resolution of the AisaEAGLE observations and the COSMO simulations are a big issue. The primary reason for using the selected observations and the applied model resolution was based on a previous study by Loewe et al. (2017), where the COSMO model was compared to observations. For the 100 m grid spacing the comparisons of liquid/ice water content, size distributions of droplets and ice crystals showed good agreements. Therefore, we were confident with this model setup and first did not change the resolution for this study.

However, the reviewer is right that the large differences in the resolutions may not be comparable. Therefore, we added simulations, where we improved the grid spacing to 50 m. We tried to simulate with 25 m grid spacing, but this was not possible due to numerical instabilities. A reduction of the spatial domain could not solve this issue. A modification of the model, e.g. implementation of a different turbulence scheme, is beyond the scope of this work.

Throughout the discussion part of the manuscript, we now discuss the findings for both, 50 m and 100 m grid spacing. The comparison of both simulation runs did show significant differences, which show that the reproduction of small-scale cloud inhomogeneities depends on the model setup. In the revised manuscript this is explicitly analyzed and discussed. Including the new simulations with the new grid spacing of 50 m caused several changes in the manuscript. Please find below the main changes with regard to text parts, graphs, and tables:

Abstract:

"Simulations are performed for spatial resolutions of 50 m (1.6 km x 1.6 km domain) and 100 m (6.4 km x 6.4 km domain)."

"[...] show that COSMO produces more homogeneous clouds by a factor of two (100 m spatial resolution) compared to the measurements. Those differences reduce for the spatial resolution of 50 m."

Introduction:

"For the Arctic Summer Cloud Ocean Study (ASCOS), Loewe et al. (2017) validated COSMO for simulations with a spatial resolution of 100 m with respect to droplet/ice crystal number concentrations, cloud top/bottom boundaries, and surface fluxes. Cloud structures and inhomogeneities were not validated due to the lack of observational data. Here, airborne imaging spectrometer measurements obtained during the VERDI campaign are used to analyze the small–scale cloud inhomogeneities (< 1 km), which are then compared to COSMO simulations using the same model setup as proposed by Loewe et al. (2017) with 64 by 64 grid points and 100 m spatial resolution as well as a finer resolved setup with 32 by 32 grid points and 50 m spatial resolution"

Section 3.1:

"[...] The size of the model domain used by Loewe et al. (2017) was $6.4_{6.4}$ km in horizontal direction with a spatial resolution of 100 m. Here, this setup is applied as well. However, analyzing cloud inhomogeneities requires a fine horizontal spatial resolution of the model simulations. Therefore, for the comparison with the imaging spectrometer measurements analyzed here, the spatial resolution is also increased to 50 m for addition model runs. In those cases, the domain size is reduced to 32 by 32 grid points (1.6 km x 1.6 km) for computational constrains. A further reduction of the spatial resolution was not possible due to numerical instabilities. [...]"

Section 4.3:

"The investigations on the single cases during VERDI are performed for spatial resolutions of 50 m (32 by 32 grid points) and 100m (64 by 64 grid points). All other model parameters are kept constant with respect to the analysis performed by Loewe et al. (2017). "

"In order to average the observed fields of τ_{meas} to the grid spacings of 50 and 100 m, the τ_{meas} -values of distinct numbers of neighboring pixels are averaged. The number depends on the single pixel size of the particular cases, which is a function of the distance between aircraft and cloud. For the four investigated cases this number varies between 13 (26) and 18 (36) pixels, which are needed to generate pixel sizes of τ_{meas} comparable to the 50 m (100 m) spatial resolution of COSMO."

"For the COSMO simulations, which use 50m spatial resolution, the domain size is reduced to 32 by 32 grid points resulting in a total domain of 1.6 km by 1.6 km, which is comparable to the observations. Therefore, the domain of those simulations was not adapted for the comparisons."

"Therefore, in the following analysis, comparing the simulated against observed fields of τ , the simulations with the finer spatial resolution of 50 m are used. The simulations with 100 m spatial resolution are used to discuss the model sensitivity with respect the spatial resolutions."



"Figure 5 (was Fig. 6). Illustrated are sections of one and the same field of τ_{meas} from 14 May 2012 with a spatial resolutions of (a) \approx 3 m (original resolution), (b) 50 m (COSMO resolution), (c) 100 m (COSMO resolution), (d) 150 m, and (e) 300 m. [...]"

Section 5.1:

	Case	$\bar{\tau} \pm \sigma_{\tau}$	$ ho_{ au}$	S_{τ}	$\chi_{ au}$
VERDI (50 m)	14 May	7.8 ± 1.5	0.195	0.086	0.979
	15 May	6.4 ± 0.7	0.121	0.055	0.992
	16 May	6.4 ± 1.0	0.166	0.078	0.983
	17 May	4.2 ± 0.5	0.154	0.071	0.986
VERDI (100 m)	14 May	8.1 ± 1.2	0.209	0.093	0.977
	15 May	6.4 ± 0.5	0.115	0.052	0.993
	16 May	6.6 ± 0.6	0.145	0.065	0.988
	17 May	4.3 ± 0.4	0.132	0.061	0.990
COSMO (50 m)	14 May	7.9 ± 0.6	0.071	0.030	0.997
	15 May	7.1 ± 0.7	0.092	0.040	0.995
	16 May	6.0 ± 0.6	0.094	0.040	0.995
	17 May	5.8 ± 0.5	0.083	0.036	0.996
COSMO (100 m)	14 May	6.9 ± 0.5	0.066	0.028	0.997
	15 May	5.4 ± 0.3	0.053	0.023	0.998
	16 May	5.5 ± 0.5	0.090	0.037	0.996
	17 May	5.6 ± 0.3	0.044	0.019	0.999

Table 2. Mean value of τ , standard deviation σ_{τ} , and the three 1D inhomogeneity parameters ρ_{τ} , S_{τ} , and χ_{τ} calculated for all four cases from the observations and the simulations with the two different grid spacings of 50 and 100 m.

"[...] Table 2 lists the mean value of τ , standard deviation σ_{σ} and the three 1D inhomogeneity parameters ρ_{τ} S_{τ} and χ_{τ} for the observations and the simulations with the two different spatial resolutions of 50 and 100 m.

Both, measurements and simulation show the highest cloud optical thickness on 14 May with $\tau_{meas} = 8.1 + -1.2$ and $\tau_{sim} = 7.9 + -0.6$ at 50 m spatial resolution and $\tau_{sim} = 6.9 + -0.5$ at 100 m spatial resolution, which show an overall agreement. [...] However, compared to the grid spacing of 100 m it is obvious that the finer resolved simulations lead to better agreements between measurements and simulations."

Section 5.2

"The 2D autocorrelation functions are calculated to compare the typical spatial scales and the directional character of the small-scale cloud inhomogeneities (no large-scale inhomogeneities like roll convection) of observations and simulations."

"Furthermore, the results for $P^2_{\tau,meas}$ and $P^2_{\tau,sim}$ show that COSMO simulations using a spatial resolution of 50 m produce similar sizes of the small-scale cloud structures compared to the measurements."

For the simulations with 100 m spatial resolution (graph not shown) the directional features still compare well between observations and simulations."



New Figure 7 (was Fig. 8) shows now the results for 50 m grid spacing and not 100 m grid spacing anymore.

Table 3. Calculated decorrelation lengths $\xi_{\tau,\text{meas}}$ and $\xi_{\tau,\text{sim}}$ for the two different resolutions/grid spacings of 50 and 100 m along (ξ_{τ}^{\ddagger}) and across (ξ_{τ}^{\ddagger}) the observed and simulated predominant directions (compare Fig. 7e-h and Fig. 7m-p for 50 m grid spacing).

	Case	$\xi^{\ddagger}_{ au,50\mathrm{m}}$ [m]	$\xi^{\leftrightarrow}_{ au,50\mathrm{m}}$ [m]	$\xi^{\ddagger}_{ au,100{ m m}}$ [m]	$\xi^{\leftrightarrow}_{ au,100\mathrm{m}}$ [m]
VERDI	14 May	800	330	>1000	400
	15 May	260	180	280	190
	16 May	220	100	350	170
	17 May	250	150	370	260
COSMO	14 May	260	190	530	320
	15 May	250	200	380	260
	16 May	270	180	500	280
	17 May	240	190	430	390

2. p. 5 Fig.2 Why you do not show wind components? Later you discuss directional shear...

It is true that it would be helpful to include the wind components. In the initially submitted version wind was partly included in Fig. 4, at least for COSMO. However, in the resubmitted manuscript we removed Fig. 4. Therefore, we now extended Fig. 2 by two panels (Dropsonde, COSMO) for the wind direction. As the dropsonde does not provide vertical wind, we limited the plot to horizontal winds, e.g. wind direction. Please find below the new Fig. 2 from the resubmitted manuscript.



Figure 2. (*a*, *e*) Potential temperature, (*b*, *f*) relative humidity, (*c*, *g*) wind speed, and (*d*, *h*) wind direction for the four investigated cases. The dropsonde data is shown in the first row (*a*-*d*) and the 2 h domain-averaged profiles after spin-up time of the simulations are shown in the second row (*e*-*h*). Dropsondes were released closest to the imaging spectrometer measurements.

3. 3.1. Simulations Model set-up is not detailed enough. Please describe fluxes, radiation, microphysics in few sentences, referencing is not enough. Subversions on May 14-15 and 16-17 are at very different heights. Was vertical resolution at inversion height comparable?

The two-moment cloud microphysics scheme by Seifert and Beheng (2006) is used in the COSMO model. Within the model the number densities and the masses of six hydrometeor types are predicted. The six hydrometeor types are cloud droplets, cloud ice, raindrops, snow, graupel, and hail. The scheme is based on the partial power moments of the number density size distribution function of cloud droplets and raindrops. The different ice phase hydrometeor growth processes are parameterized, in which the depositional growth of ice particles is dominant in Arctic mixed-phase clouds.

The radiation is a two-stream radiation scheme after Ritter and Geleyn (1992). It is calculated every 2 s and has a direct cloud-radiative feedback.

The vertical resolution at the inversion height on the different days is comparable with a maximum vertical grid spacing of around 15 m up to the inversion height.

We added additional information about the cloud scheme, the radiation scheme and the vertical resolution in section 3.1. Further, the surface fluxes depend on the surface temperature, which is 273.5 K for the sea water surface. We added this information in Sec. 3.1 as well.

"The two-moment cloud microphysics scheme by Seifert and Beheng (2006) predicts the number densities and the masses of six hydrometeor types. The different ice phase hydrometeor growth processes are parameterized in this scheme. In COSMO, the radiative transfer is described by a two-stream radiation scheme after Ritter and Geleyn (1992). It is calculated every 2 s and has a direct cloud-radiative feedback. A three-dimensional prognostic turbulence scheme describes the turbulent fluxes of heat, momentum and mass by a first-order closure after Smagorinsky and Lilly (Herzog et al., 2002; Langhans et al., 2012)."

"The vertical height range of 22 km is divided into 166 vertical levels, which are more dense for the ABL with a typical grid spacing of around 15 m up to the inversion height of the different days of investigation."

"The surface of the model is sea water and the surface fluxes depend on the surface temperature, which is 273.5 K for the sea water surface."

4. p.6 l. 17 "to avoid numerical issues" really? Or data from dropsondes represent actual realization along trajectory, not a good choice for initial profiles?

Our wording "avoid numerical issues" was obviously confusing. We meant, that we are simulating with a very high resolution, which is numerical expensive. The record of a dropsonde is higher in time and in space and thus we have many more vertical levels than in the model. Also because of the horizontal drift of the dropsonde through cloud inhomogeneities the profile is not monotonically. This caused issues in the model initialization and had to be smoothed out.

The dropsonde data are the only information we have from the atmospheric conditions during the campaign and it is a good choice to define the atmospheric BL, because of the high vertical resolution. The aim of the measurement flights during VERDI was to have similar conditions in a specified sector, where e.g. the cloud top height is the same over a certain area.

"The dropsonde are partly affected by horizontal variability, when slowly passing the cloud and drifting horizontally. Therefore parts of the original profiles (Fig.2) are smoothed and brought to a vertical monotonic increasing profile for initialization of the model."

5. p. 7 l. 12 WD in Fig 4 I guess is for wind direction, but generally the figure is hard to interpret. E.g. the same wind shear whether in the middle of the given colour and at the edge of colours can be visible or not. I fill not comfortable with this plot.

The reviewer is right. Figure 4 was hard to interpret. However, in the resubmitted version of the manuscript we removed Fig. 4. For the discussions of the wind direction we now use the extended plot in Fig. 2.

"The simulation of the 16 May shows a wind shear from around 150° to around 100° (Fig. 2) and a decrease of v with height above the cloud top height, which is also seen in the dropsonde profiles (Fig. 2c). The other simulations do not show a turning of the wind directly above the inversion height."

"Figure 2d and Fig. 2h illustrate the temporally averaged wind directions in the simulations."

6. Section 4.2 The section shows nicely discrepancies between the experimental data and the simulation. Why in conclusion do you not call for higher resolution simulations? In the supplementary material of the paper you cite (Pedersen et al., 2016) there are suggestions that basic cloud patterns are reproduced reasonably in smaller domain. Why do you not perform sensitivity analysis due to model resolution?

As replied to the reviewer's first comment, in the resubmitted version of the manuscript, we included a sensitivity study with respect to the model grid spacing. All relevant changes in the manuscript are given in our answer to the reviewer's first comment

7. Section 4.3 Again: your model domain larger than the swath of the measurements. Why not to run model in smaller domain but at higher resolution? In particular when you conclude that the decorrelation length increases with decreasing spatial resolution.

Thanks for this suggestion. As explained above, we added simulations with a higher resolution of 50 m by reducing the grid points. Additionally, we tested a further increase of the resolution. However, due to numerical instabilities, it was not possible to further increase the grid spacing. Please find the changes in the manuscript related to this comment below our answer to the first reviewer comment.

8. Section 5.1. I think that your conclusion that the model captures temporal changes of inhomogeneity is not well justified, there are only 4 points analyzed. Moreover, the maximum modelled inhomogeneity is dated May 16th, while is observed on May 14th. On these days, vertical profiles indicate that clouds and boundary layer properties on these days are substantially different.

Maybe it was misleading that we talked about a trend, which might lead to the impression that we compare the changes of ρ_{τ} on a temporal scale. In making this comparison, our intention was to show that in general COSMO produced larger/lower inhomogeneity, when larger/lower inhomogeneity was observed by the measurements. To avoid a further confusion, we rewrote this part by the following:

"However, the observations show that the cloud field became more homogeneous from 14 to 15 May as indicated by lower values of ρ_{τ} which reduce from 0.209 to 0.115. From 15 to 16 May, ρ_{τ} increases to 0.145, which indicates a cloud field with slightly higher inhomogeneity. Then, on 17 May, ρ_{τ} reduced to 0.132, showing that the cloud field became more homogeneous again. These different cases with high and low ρ_{τ} are reproduced by COSMO independent on the chosen spatial resolution. Larger discrepancy between modeled and observed inhomogeneity parameters only occurred on 14 May, when the observations were influenced by large–scale cloud structures. Nevertheless, the lower/higher inhomogeneity is also imprinted in the inhomogeneity parameters S_{τ} and χ_{τ} , which are smaller/larger in both, measurements and simulations, indicating that COSMO performs well with regard to the 1D inhomogeneity parameters." 9. Section 5.2. Results in this section are more convincing. However, these results could be strengthened discussing dynamical patterns (convective rolls) the boundary layer. Does the maximum optical thickness correlate with location of updraughts and maximum cloud top heights? Analysis of that could help to publish the paper, since conclusions are weak and should be supported with additional investigations; which can increase our understanding of modelled processes. This is particularly important in terms of your sensitivity study in Section 6.

The reviewer is right. A more though rough investigation of the dynamical patterns of the clouds detected in the boundary layer will help to strengthen the results of the manuscript. Unfortunately, larger scale dynamic patterns such as role convection are not fully covered by the narrow view of the AisaEAGLE imaging spectrometer. Therefore, such analysis as suggested by the reviewer would only be possible using the simulations with COSMO. As this would not be very convincing without the observations, we, therefore, focus on the small-scale structure of cloud inhomogeneities.

Anyway, thanks for this suggestion! It is indeed possible using the methods applied in the manuscript to study larger scale dynamic patterns and might be done in a follow up study.

"Thus, also large–scale cloud structures are covered by COSMO (purple stripes in Fig. 4d) but not in the observations. Therefore, the large–scale structures cannot be compared between observations and simulations. With respect to a comparison of the small–scale structures, the spatial sizes (spatial resolution, domain size) of both datasets need to be conformed to make a direct comparison possible."

"Figure 4e and Fig. 4f further illustrate that it is not possible to compare the large–scale structures between observations and simulations. The large–scale structures, which are covered by the COSMO simulations are identified by a second increase of the $P^2_{\tau,sim}$ at distances (≈ 1 km in Fig. 4f) larger than ξ_{τ} . The width of the measured fields is too narrow to cover such a second increase in the $P^2_{\tau,meas}$ (compare Fig. 4e). Therefore, the further comparison of the cloud structures, which are identified in the observations and simulations, is restricted to the small–scale cloud structures with sizes below 1 km only."

"The statistical methods applied in this study can also be applied to characterize the larger-scale dynamic patterns, if the domain is large enough to resolve them."

10. Section 6. Again: this section calls for more thorough analysis as pointed above.

By including simulations with 50 m grid spacing and the following discussion comparing different resolutions, we hope some interesting new results did now improve the manuscript. Comparing the results for the small domain (1.6 x 1.6 km) with the large domain (6.4 x 6.4 km), a major conclusion is that the small-scale structures simulated by COSMO must be influenced by the large-scale structures. This results from the fact that the small-scale structures depend on the wind speed (see graph below), when the domain is large enough so that large-scale structures can evolve. Contrarily, in the small domain, where no large-scale structures evolve, the small-scale structures are independent on changes of wind speed. Therefore, the effect of wind on the small-scale structures (Fig. <u>8</u>s to Fig. <u>8</u>x) acts only indirectly via the change of large-scale cloud structures. This has the consequence that the natural behavior of the small-scale structures (e.g. their size and orientation) might be disturbed, if the simulated domain is too small.

In the resubmitted manuscript, we extensively elaborate those new findings. Please find below our main changes to the manuscript and the revised Fig. 9 (now Fig. 8).



"Figure 8. Exemplary selected fields of τ_{sim} for the 15 May 2012 case simulated for differently scaled initial wind speeds on a grid with 50 m spatial resolution and 1.6 km by 1.6 km domain **(a-f)** and on a grid with 100 m spatial resolution and with 6.4 km by 6.4 km domain **(m-r)**. Calculated 2D autocorrelation coefficients P_{τ}^2 are given for each case in **(g-I)** and **(s-x)**. White lines in (s)--(x) illustrate the orientation used for the calculation of the 1D P_{τ}^2 along (straight white lines) and across (dashed white lines) the dominant directions illustrated in Fig. 10. Red squares in (m)–(r) mark areas of comparable size to the small domains in (a)–(f)."

"To test its influence on the horizontal cloud inhomogeneity, the simulations for 15 May (50 and 100 m grid spacing) are repeated for different initializations, where the wind profile is varied. Here, the case on 15 May is chosen, because it shows the best agreement between observations and simulations (Fig. 7r) to serve as a benchmark case."

"Figure 8a to Fig. 8f show the simulated 2D fields of τ_{sim} for the simulations with the domain size of 1.6 km by 1.6 km and 50 m spatial resolution. Small-scale structures (< 0.5 km) are obvious and rather randomly orientated throughout the simulations for all six different initializing wind profiles. The spatial sizes of the small–scale structures quantified by the decorrelation length depend only little on the wind speed. This is confirmed by the 2D autocorrelation analysis illustrated in Fig. 8g to Fig. 8l. Displayed are only the horizontal scales below 0.8 km, quantified by the 2D autocorrelation coefficients for shifts below +/-0.8 km. A predominant direction of the small–scale structures is only slightly developed and varies independently from cases to case without clear preference. Furthermore, the P^2_{τ} and the decorrelation length, which vary between 150 and 300 m show only slight variations with changing wind speed. This means that the sizes of the small–scale structures is basically independent to the wind speed.

Contrarily, the simulations with a domain size of 6.4 by 6.4 km and 100 m spatial resolution show a clear dependency on the wind speed. [...]''

"Comparing the simulations for the small domain (1.6 x 1.6 km, 50 m spatial resolution) with the large domain (6.4 x 6.4 km, 100 m spatial resolution), indicates that the small–scale structures are most likely influenced by the large–scale structures. Only for the simulations with the large domain, the small-scale structures depend on the wind speed. This indicates that small–scale cloud inhomogeneities are not directly linked to the wind speed but rather are influenced by the large–scale cloud inhomogeneities such as cloud roles. If these large–scale structures are not covered by the simulations (too small domain), the natural behavior of the small–scale structures (e.g. their size and orientation) might be disturbed. With respect to the comparison between observations and simulations, this may explain why only on 14 May larger differences between model and observations were found. All other three cases did not show a significant large–scale cloud structure, while on 14 May cloud roles were observed by the imaging spectrometer. Thus, the simulations of 15, 16, and 17 May are more uncritical with respect to the model domain than for 14 May, when a large domain is required to reproduce the large–scale cloud structures and, therefore, improve the simulation of the small–scale cloud structures."

11. Section 7. After the additional analysis this section (and abstract) should be updated adequately.

Accordingly to the revised manuscript, we updated the abstract and Section 7 – Summary and conclusions. Please find the main changes below.

Abstract:

"[...] Simulations are performed for spatial resolutions of 50 m (1.6 km x 1.6 km domain) and 100 m (6.4 km x 6.4 km domain). Macrophysical cloud properties such as cloud top altitude and vertical extent are well captured by COSMO. Cloud horizontal inhomogeneity quantified by the standard deviation and one-dimensional (1D) inhomogeneity parameters show that COSMO produces more homogeneous clouds by a factor of two (100 m spatial resolution) compared to the measurements. Those differences reduce for the spatial resolution of 50 m. However, for both spatial resolutions the directional structure of the cloud inhomogeneity is well represented by the model. Differences between the individual cases are mainly associated with the wind shear near cloud top and the vertical structure of the atmospheric boundary layer. A sensitivity study changing the wind velocity in COSMO by a vertically constant scaling factor shows that the directional small–scale cloud inhomogeneity structures can range from 250 m to 800 m and depend on the mean wind speed, if the simulated domain is large enough to capture also large–scale structures, which then influence the small–scale structures. For those cases a threshold wind velocity is identified, which determines when the cloud inhomogeneity stops increasing with increasing wind velocity."

Summary and Conclusion:

"[...] For the reason of comparability, the observed fields of cloud optical thickness τ_{meas} were aggregated to pixel sizes of 50 m and 100 m, the applied spatial resolutions of the individual simulations. The general inhomogeneity was compared using 1D inhomogeneity parameters. For 100 m spatial resolution the absolute values of cloud inhomogeneity derived from COSMO are larger by a factor of about two, as compared to the values obtained from the observations. These differences slightly reduce, when the spatial resolution of the simulations is increased by a finer grid of 50 m. However, for both spatial resolutions the cloud inhomogeneity generated by COSMO is too low. [...]

For the COSMO simulations with a spatial resolution of 50 m, the ξ^{\uparrow}_{τ} and $\xi^{\leftrightarrow}_{\tau}$ agree well between observations and simulations, except for the case on 14 May. In contrast, for the simulations with a spatial resolution of 100 m, COSMO produced small–scale cloud structures with characteristic sizes 20 to 30 % larger compared to the observations. However, for both spatial resolutions the best agreement was found for the case on 15 May 2012.

The agreement between COSMO and observations for the case on 15 May 2012 is used as basis for a systematic sensitivity study with respect to the wind speed as a main drivers of the cloud inhomogeneities. Simulations for the case on 15 May with differently scaled initialization wind profiles showed that the degree of horizontal cloud inhomogeneity was not significantly changed for the simulations with a small domain (1.6 km x 1.6 km) and 50 m spatial resolution, but for the simulations using a large domain (6.4 km x 6.4 km) and 100 m spatial resolution. This indicates that the large–scale cloud structures such as cloud roles influence the small–scale cloud inhomogeneity. To correctly simulate the small–scale cloud inhomogeneity, COSMO needs to be run in a large domain, which also covers the large–scale cloud structures. This might have been the reason for the large differences between observations and simulations found for the case of 14 May, when pronounced cloud rolls were observed. All other cases did not show such large–scale cloud structures and were simulated by COSMO closer to reality despite the small domain.

[...]

Altogether, cloud inhomogeneities are a challenge for cloud resolving models. Not only the spatially averaged magnitude of inhomogneity but also the directional structure and the interaction with large–scale cloud structures needs to be reproduced in the simulations. Although COSMO produces more homogeneous clouds, it performed well, because it correctly represented the directional structures and the general degree of cloud inhomogeneity, if no larger–scale cloud structures are present. However, the statistical methods applied in this study can also be applied to characterize the larger–scale dynamic patterns, if the domain is large enough to resolve them."

We thank the reviewer for the helpful comments, which certainly improved the manuscript. Especially, due to more detailed descriptions of the background and discussions of the main topic the manuscript has improved significantly. The detailed replies on the reviewer's comments are structured as follows. Reviewer comments have bold letters, are labeled, and listed always in the beginning of each answer followed by the author's comments including (if necessary) revised parts of the paper. The revised parts of the paper are written in quotation marks and italic letters.

Major Issues:

1. The introduction is too long. Even after reading the second page, I am not sure why we should worry about the inhomogeneity in the cloud radiation properties aka optical depth. Is it because we need better sub-grid characterization of radiative properties in global climate models? It will be better if the authors explicitly state the specific objective of the study.

The reviewer is right. The introduction was written too general and broad on Arctic clouds. We have shortened it and put the focus stronger on the cloud inhomogeneities and their directional structures. Please find below the revised introduction:

"Arctic clouds are expected to be a major contributor to the so-called Arctic Amplification (Serreze and Barry, 2011; Wendisch et al., 2017) and, therefore, need to be represented adequately in model projections of the future Arctic climate (Vavrus, 2004). Especially, low-level Arctic stratus are of importance (Wendisch et al., 2013), because they occur quite frequently (around 40 %, Shupe et al., 2006, 2011), typically persist over several days or even weeks (Shupe et al., 2011), and on annual average, warm the Arctic surface (Shupe and Intrieri, 2004). The numerous physical and microphysical processes that determine the properties of Arctic stratus are complexly linked to each other (e.g., Curry et al., 1996) and still not understood in full detail (Morrison et al., 2012).

Dynamic factors (updrafts), which increase the actual supersaturation in the cloud beyond the equilibrium values for both liquid water and ice, and a steady supply of water vapor from above the cloud act to stabilize the Arctic stratus (Shupe et al., 2008). This facilitates the simultaneous existence of both phases (Korolev, 2007). While in updrafts liquid and ice crystals grow, the cloud top cooling induces downward vertical motion, where Wegener-Bergeron-Findeisen process may dominate. Therefore, small–scale structures can be important to understand the microphysical processes. Additionally, Arctic stratus shows microphysical inhomogeneities, which typically occur on horizontal and vertical scales below a few kilometers and even tens of meters (Chylek and Borel, 2004; Lawson et al., 2010). The small–scale cloud structures, which accompany cloud inhomogeneities, lead to three-dimensional (3D) radiative effects (Varnai and Marshak, 2001), which can be parameterized using inhomogeneity parameters (Iwabuchi and Hayasaka, 2002; Oreopoulos and Cahalan, 2005).

Unfortunately, the understanding of Arctic clouds is impeded by a paucity of comprehensive observations due to a lack of basic research infrastructure and the harsh Arctic environment (Intrieri et al., 2002; Shupe et al., 2011). Therefore, observation of small–scale cloud structures within the Arctic circle are sparse. Satellite observations are typically too coarse to resolve scales below 250 m and space–born passive remote sensing observations suffer from contrast problems over highly reflecting surfaces (snow and ice, Rossow and Schiffer, 1991). Ground–based remote sensing observations with radar and lidar typically point only in zenith direction and are not capable to provide the horizontal 2D–structure of clouds. Only along the wind direction the variability of clouds is resolved (Shiobara et al., 2003; Marchand et al., 2007). For example, using correlation analysis, Hinkelmann (2013) revealed significant differences between along–wind and cross–wind solar irradiance variability on small spatial scales in broken–cloud situations. In comparison, airborne spectral imaging observation of reflected solar radiation provide areal measurements with spatial resolution down to several meters (Schäfer et al., 2015). Bierwirth et al. (2013) used such airborne measurements of reflected solar spectral radiance to retrieve fields of cloud optical thickness τ of Arctic stratus and demonstrated their strong spatial

variability. From similar measurements, Schäfer et al. (2017a) analyzed the directional variability of different cloud types including Arctic stratus. The few analyzed cases revealed that 1D statistics are not sufficient to quantify the variability of horizontal clouds inhomogeneities.

Likewise, treating small–scale inhomogeneities using reanalysis data and atmospheric models is difficult. Global reanalysis products have relatively coarse spatial resolutions (40 km and larger; Lindsay et al., 2014) and, therefore, do not resolve small–scale features. Furthermore, in numerical weather prediction and climate models, the representation of the temporal evolution of mixed-phase clouds is poor (Barrett et al., 2017a, b). Especially, areas of up- and downdrafts in Arctic stratus, which are typically in the range of less than 1 km cannot be resolved but have to be parametrized (Field et al., 2004; Klein et al., 2009). To realistically simulate the spatial structure of these clouds, Large Eddy Simulations (LES) with a spatial resolution of 100 m or less and high vertical resolution (< 20 m within atmospheric boundary layer, ABL) are needed. Those LES can resolve the vertical motion of the turbulent eddies in the ABL and the cores of up- and downdrafts representing the inhomogeneities in the cloud top structure, which can be seen in the amount of liquid water at the cloud top. The size of the up- and downdraft cores may differ depending on the time of the year (Roesler et al., 2016).

Previous LES studies focus for instance on cloud-top entrainment (Mellado, 2017) and emphasize the behavior of changes in the spatial resolution on the liquid water path (Pedersen et al., 2016). Kopec et al. (2016) discussed two main processes, the radiative cooling and wind shear. The radiative cooling sharpened the inversion, while wind shear at the top of the ABL causes the turbulence in the capping inversion and lead to dilution at the cloud top.

In general, LES are helpful to focus on a certain process and to investigate cloud formation, cloud evolution or the small-scale structures in an Arctic stratus under controlled conditions. The further aim is to characterize horizontal small-scale cloud inhomogeneities in the size range of less than 1 km in simulations and measurements to better understand the radiative properties of Arctic mixed-phase clouds. Results from the COSMO (COnsortium for Small-Scale MOdeling) model, which is adjusted to a LES setup with a high horizontal and vertical resolution to resolve the cloud structures of Arctic stratus (Loewe et al., 2017; Stevens et al., 2017) are evaluated. For the Arctic Summer Cloud Ocean Study (ASCOS), Loewe et al. (2017) validated COSMO for simulations with a spatial resolution of 100 m with respect to droplet/ice crystal number concentrations, cloud top/bottom boundaries, and surface fluxes. Cloud structures and inhomogeneities were not validated due to the lack of observational data. Here, airborne imaging spectrometer measurements obtained during the VERDI campaign are used to analyze the small-scale cloud inhomogeneities (< 1 km), which are then compared to COSMO simulations using the same model setup as proposed by Loewe et al. (2017) with 64 by 64 grid points and 100 m spatial resolution as well as a finer resolved setup with 32 by 32 grid points and 50 m spatial resolution. For that, data measured by dropsondes served as input for semi-idealized simulations of clouds using COSMO-LES (Sec. 2.3 and Sec. 3). Airborne measured fields of cloud optical thickness retrieved from imaging spectrometer measurements (Sec. 2.2) are used for a comparison with the resulting COSMO clouds with respect to their overall cloud inhomogeneity and directional features of the cloud inhomogeneities (Sec. 4 and Sec. 5). Observations and modelling are aimed to be combined to quantify the horizontal cloud top structures, which are discussed in Sec. 5 and Sec. 6."

2. It is unclear why you chose wind speed as a tuning parameter. By increasing wind speeds you are simple changing the fluxes in the boundary layer and hence the turbulence. So essentially your results are suggesting that we greater turbulence produces higher inhomogeneity, which makes sense. It will be better if the authors can probe this. One way to tackle this would be to make some simulations where the winds are the same, but you increase the surface fluxes.

Thank you for your comment. Probably we did not describe our aim with changing the wind speed very well.

Our intention for changing the wind speed was to change the wind shear at the inversion and, therefore, turbulent processes at cloud top. As explained by the reviewer, this implicitly also changes

surface fluxes. However, simulations by Loewe (2017) over different surface types (sea ice, open lead, open water), e.g. different surface fluxes, showed only an effect in the LWP, which was increased, but did not change the cloud structure. In these simulations, when the BL was coupled to the surface. The cases of the sensitivity study presented here, except of the 16 May cloud are characterized by a boundary de-coupled to the surface. Therefore, surface fluxes are expected to have a minor impact on the cloud layer. Thus, we chose to influence the cloud top structure by changing the wind shear. In the revised manuscript we added this dicussion in Section 6:

"Influences from the surface fluxes are only expected if the cloud is coupled to the surface and if so, affect only the LWP of the cloud (Loewe, 2017). For de-coupled clouds, it is assumed that the cloud structure depends more strongly on the wind shear, respectively the wind speed."

3. Lastly, the authors should show the comparison between the model reported liquid water paths, and cloud boundaries with those observed during the campaign. I think this will make the article complete. Thanks.

Unfortunately, such a comparison is not reasonably for this study, as the model was initialized with the atmospheric profiles (temperature, humidity, wind) observed during the campaign. Therefore, the cloud boundaries in the simulations are almost identical as those measured by the dropsonde profiles. Similar, the liquid water paths were adapted to the dropsonde profiles.

Minor Issues:

1. Line 58: Need reference to justify that sentence.

It was shown by Schäfer et al. (2017a). However, the tense was incorrect due to a typo. We corrected the sentence for this.

"From similar measurements, Schäfer et al. (2017a) analyzed the directional variability of different cloud types including Arctic stratus. The few analyzed cases revealed that 1D-statistics are not sufficient to quantify the variability of horizontal clouds inhomogeneities."

2. Line 101-102: I would simply say that the cloud fraction decreased. The word "dissolved" seems inappropriate in terms of clouds.

We changed "dissolved" to "decreased".

3. Line 121: By "ten fields" I believe you mean ten snapshots?

The reviewer is right. In general, these are only snapshots, which means only smaller parts of a larger cloud scene. However, referring to Schäfer et al. (2017a), we would like to keep on calling it "fields" of cloud optical thickness, although they do not capture the whole cloud scene.

4. Section 3.1: Please describe the radiation and cloud schemes used in the model. Since you are evaluating optical depth, which is a radiative property, it is important to know this. Also mention how often the two schemes are talking to each other. Thanks.

Thank you for your helpful comment. The other reviewer had a similar comment and we like to apply the same answer her.

The two-moment cloud microphysics scheme by Seifert and Beheng (2006) is used in the COSMO model. Within the model the number densities and the masses of six hydrometeor types are predicted. The six hydrometeor types are cloud droplets, cloud ice, raindrops, snow, graupel, and hail. The scheme is based on the partial power moments of the number density size distribution function of cloud droplets and raindrops. The different ice phase hydrometeor growth processes are parameterized, in which the depositional growth of ice particles is dominant in Arctic mixed-phase clouds.

The radiation is a two-stream radiation scheme after Ritter and Geleyn (1992). It is calculated every 2 s and has a direct cloud-radiative feedback.

The vertical resolution at the inversion height on the different days is comparable with a maximum vertical grid spacing of around 15 m up to the inversion height.

We added additional information about the cloud scheme, the radiation scheme and the vertical resolution in section 3.1. Further, the surface fluxes depend on the surface temperature, which is 273.5 K for the sea-water surface. We added this information in Sec. 3.1 as well.

"The two-moment cloud microphysics scheme by Seifert and Beheng (2006) predicts the number densities and the masses of six hydrometeor types. The different ice phase hydrometeor growth processes are parameterized in this scheme. In COSMO, the radiative transfer is described by a two-stream radiation scheme after Ritter and Geleyn (1992). It is calculated every 2 s and has a direct cloud-radiative feedback. A three-dimensional prognostic turbulence scheme describes the turbulent fluxes of heat, momentum and mass by a first-order closure after Smagorinsky and Lilly (Herzog et al., 2002; Langhans et al., 2012)."

"The vertical height range of 22 km is divided into 166 vertical levels, which are more dense for the ABL with a typical grid spacing of around 15 m up to the inversion height of the different days of investigation."

"The surface of the model is sea water and the surface fluxes depend on the surface temperature, which is 273.5 K for the sea water surface."

5. Line 273-279: Please rephrase these sentences. It is confusing to read "large resolution" etc. thanks.

We revised the relevant sentence by the following:

"Thus, the spatial resolution of AisaEAGLE is relatively high, compared to the grid spacing of 100 m from COSMO."

Figure 5e and 5f: There is no "grey dotted line" in the plot. 6.

We have updated the graph shortly before we initially submitted the manuscript and must have missed to include this line in the new version. Now, it is included in the resubmitted version. Please see the graph below:



7. Figure 7a: the plot is showing mean and standard deviation, however there are two blue dots for each resolution? If you are showing mean+std and mean-std, then I suggest you show vertical error-bars.

We have changed the graph accordingly. Now we are using error bars. Please find the revised Figure below:





Manuscript prepared for Atmos. Chem. Phys. with version 2014/09/16 7.15 Copernicus papers of the LATEX class copernicus.cls. Date: 12 July 2018

Simulated and observed horizontal inhomogeneities of optical thickness of Arctic stratus

Michael Schäfer^{1,*}, Katharina Loewe^{2,*}, André Ehrlich¹, Corinna Hoose², and Manfred Wendisch¹

¹Leipzig Institute for Meteorology, University of Leipzig, Leipzig, Germany ²Institute of Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

^{*}Both authors contributed equally to this work.

Correspondence to: Michael Schäfer (michael.schaefer@uni-leipzig.de), Katharina Loewe (katharina.loewe@kit.edu)

Abstract. Two-dimensional (2D) horizontal fields of cloud optical thickness τ_{c} derived from airborne measurements of solar spectral radiance during the Vertical Distribution of Ice in Arctic Clouds (VERDI) campaign (carried out in Inuvik, Canada in April/May 2012) are compared with semi-idealized Large Eddy Simulations (LES) of Arctic stratus performed with the COnsortium for

- 5 Small-Scale MOdeling (COSMO) atmospheric model. The input for the LES is obtained from collocated airborne dropsonde observations. Four consecutive days of a persistent Arctic stratus observed above the sea-ice free Beaufort Sea are selected for the comparison. Simulations are performed for spatial resolutions of 50 m (1.6 km × 1.6 km domain) and 100 m (6.4 km × 6.4 km domain). Macrophysical cloud properties such as cloud top altitude and vertical extent are well captured by COSMO.
- 10 Cloud horizontal inhomogeneity quantified by the standard deviation and one-dimensional (1D) inhomogeneity parameters show that COSMO produces only half of the measured horizontal cloud inhomogeneities, while more homogeneous clouds by a factor of two (100 m spatial resolution) compared to the measurements. Those differences reduce for the spatial resolution of 50 m. However, for both spatial resolutions the directional structure of the cloud inhomogeneity is well represented
- 15 by the model. Differences between the individual cases are mainly associated with the wind shear near cloud top and the vertical structure of the atmospheric boundary layer. A sensitivity study changing the wind velocity in COSMO by a vertically constant scaling factor shows that the directional small-scale cloud inhomogeneity structures strongly can range from 250 m to 800 m and depend on the mean wind speed. A, if the simulated domain is large enough to capture also large-scale
- 20 structures, which then influence the small-scale structures. For those cases a threshold wind velocity

is identified, which determines when the cloud inhomogeneity stops increasing with increasing wind velocity.

1 Introduction

- Arctic clouds are expected to be a major contributor to the so-called Arctic Amplification (Serreze and Barry, 2011; Wendisch et al., 2017) and, therefore, need to be represented adequately in model projections of the future Arctic climate (Vavrus, 2004). Especially, low-level Arctic stratus are of importance (Wendisch et al., 2013), because they occur quite frequently (around 40%, Shupe et al., 2006, 2011), typically persist over several days or even weeks (Shupe et al., 2011), and on annual average, warm the Arctic surface (Shupe and Intrieri, 2004). The numer-
- 30 ous physical and microphysical processes that determine the properties of Arctic stratus are complexly linked to each other (e.g., Curry et al., 1996) and still not understood in full detail (Morrison et al., 2012). Some are explained in the following: Aretic stratus often appears in mixed phase state (liquid water and solid ice coexist, McFarquhar et al., 2007; Mioche et al., 2015) and, therefore, the cloud should glaciate rapidly due to the Wegener-Bergeron-Findeisen process.
- 35 However, this is not the case. A mixed-phase Arctic stratus can still persist for several days or even weeks.

Dynamic factors (updrafts), which increase the actual supersaturation in the cloud beyond the equilibrium values for both liquid water and ice, and a steady supply of water vapor from above the cloud act to stabilize the Arctic stratus (Shupe et al., 2008). This facilitates the simultaneous existence of

- 40 both phases (Korolev, 2007). Aretic mixed-phase clouds are characterized by such distinct up- and downdraft regions. While in updrafts liquid and ice crystals grow, the cloud top cooling induces downward vertical motion, where Wegener-Bergeron-Findeisen process may dominate. Therefore, these-small-scale structures can be important to understand the microphysical processes. However, mixed-phase Aretic stratus typically shows microphysical properties that are inhomogeneous, often
- 45 Additionally, Arctic stratus shows microphysical inhomogeneities, which typically occur on horizontal and vertical scales of below a few kilometers and even tens of meters (Chylek and Borel, 2004; Lawson et al., 2010). The small–scale cloud structures, which accompany cloud inhomogeneities, lead to three-dimensional (3D) radiative effects (Varnai and Marshak, 2001). Those, which can be parameterized using inhomogeneity parameters (Iwabuchi and Hayasaka, 2002; Oreopoulos and Ca-
- 50 halan, 2005).

Unfortunately, the understanding of Arctic clouds is impeded by a paucity of comprehensive observations due to a lack of basic research infrastructure and the harsh Arctic environment (Intrieri et al., 2002; Shupe et al., 2011). Therefore, also observation of small scale observation of small-scale cloud structures within the Arctic circle are sparse. Satellite observations are typically too coarse to resolve

55 scales below 250 m .- Also, space-born and space-born passive remote sensing observations suffer

from contrast problems over highly reflecting surfaces (snow and ice, Rossow and Schiffer, 1991). Ground-based Ground-based remote sensing observations with Radar and LiDAR (Light Detecting And Ranging) radar and lidar typically point only in zenith direction and are not capable to provide the horizontal 2D-structure 2D-structure of clouds. Only along the wind direction the variability of

- 60 clouds is resolved (Shiobara et al., 2003; Marchand et al., 2007). For example, using correlation analysis, Hinkelmann (2013) revealed significant differences between along-wind and cross-wind solar irradiance variability on small spatial scales in broken-cloud situations. <u>Airborne In comparison</u>, <u>airborne</u> spectral imaging observation of reflected solar radiation provide areal measurements with spatial resolution down to several meters (Schäfer et al., 2015). Bierwirth et al. (2013) used such air-
- borne measurements of reflected solar spectral radiance to retrieve fields of cloud optical thickness τ of Arctic stratus and demonstrated their strong spatial variability. From similar measurements, Schäfer et al. (2017a) analyzed the directional variability of different cloud types including Arctic stratus. The few analyzed cases show, revealed that 1D-statistics are not sufficient to quantify the variability of horizontal clouds inhomogeneities.
- 70 Likewise, treating small-scale inhomogeneities using reanalysis data and atmospheric models is difficult. Global reanalysis products have relatively coarse spatial resolutions (40 km and larger; Lindsay et al., 2014) and therefore, therefore, do not resolve small-scale small-scale features. Furthermore, in numerical weather prediction models and climate models, the representation of the temporal evolution of mixed-phase clouds is poor (Barrett et al., 2017a, b). Especially, areas of up-
- 75 and downdrafts in Arctic stratusthat, which are typically in the range of less than 1 km cannot be resolved but have to be parametrized (Field et al., 2004; Klein et al., 2009). To realistically simulate the spatial structure of these clouds, Large Eddy Simulations (LES) with grid spacings a spatial resolution of 100 m or less and high vertical resolution (≈ 20 m within atmospheric boundary layer, ABL) are needed. Thus, these simulations have a high horizontal and vertical resolution, which are
- 80 important to properly simulate Those LES can resolve the vertical motion of the turbulent eddies in the ABL and the cores of up- and downdrafts representing the inhomogeneities in the cloud top structure, which can be seen in the amount of liquid water at the cloud topstructures. The size of the up- and downdraft cores may differ depending on the time of the year (Roesler et al., 2016). Previous LES studies focus for instance on cloud-top entrainment (Mellado, 2017) and em-
- 85 phasize for instance the behavior of changes in the <u>spatial</u> resolution on the liquid water path (Pedersen et al., 2016). Moreover, a high ratio of horizontal grid spacing to vertical grid spacing, called grid aspect ratio, lead to a better agreement with observations. In Kopec et al. (2016) Kopec et al. (2016) discussed two main processes, the radiative cooling and wind shear, are discussed. In their study, The radiative cooling sharpened the inversionand, while
- 90 wind shear at the top of the atmospheric boundary layer (ABL) is the cause of causes the turbulence in the capping inversion and lead to dilution at the cloud top. Thus in

In general, LES are helpful to focus on a certain process and to investigate cloud formation,

cloud evolution or the small-scale structures in an Arctic stratus under controlled conditions. Mixed-phase clouds are challenging for models because of the metastable coexistence of liquid

- 95 cloud droplets and ice particles. Moreover, the thermal-infrared cooling at cloud top is crucial for the persistence of Arctic stratus (Morrison et al., 2012). Studies have shown that the representation of the temporal evolution of these clouds in numerical weather prediction models and climate models is poor (Barrett et al., 2017a, b). Thus, LES studies are needed to better represent the relevant processes within The further aim is to characterize horizontal small-scale cloud inhomogeneities
- 100 in the size range of less than 1 km in simulations and measurements to better understand the radiative properties of Arctic mixed-phase clouds. Here, results Results from the COSMO (COnsortium for Small-Scale MOdeling) model, which is adjusted to a LES setup with a high horizontal and vertical resolution to resolve the cloud structures of Arctic stratus (Loewe et al., 2017; Stevens et al., 2017) are evaluated. For the Arctic Summer Cloud Ocean Study (ASCOS),
- 105 Loewe et al. (2017) validated COSMO for simulations with a spatial resolution of 100 m with respect to droplet/ice crystal number concentrations, cloud top/bottom boundaries, and surface fluxes. Cloud structures and inhomogeneities were not validated due to the lack of observational data. Here, airborne imaging spectrometer measurements obtained during the VERDI campaign are used to analyze the small–scale cloud inhomogeneities (<1 km), which are then compared to
- 110 COSMO simulations using the same model setup as proposed by Loewe et al. (2017) with 64 by 64 grid points and 100 m spatial resolution as well as a finer resolved setup with 32 by 32 grid points and 50 m spatial resolution. For that, data measured by dropsondes served as input for semi-idealized simulations of clouds using COSMO-LES (Sec. 2.3 and Sec. 3). Airborne measured fields of cloud optical thickness retrieved from imaging spectrometer measurements (Sec. 2.2) are used for a com-
- 115 parison with the resulting COSMO clouds with respect to their overall cloud inhomogeneity and directional features of the cloud inhomogeneities (Sec. 4 and Sec. 5). Observations and modelling are aimed to be combined to quantify the horizontal cloud top structures, which are discussed in Sec. 5 and Sec. 6.

2 Airborne measurements

120 2.1 VERtical Distribution of Ice in Arctic clouds (VERDI) campaign

Cloud remote sensing and atmospheric profiles by dropsondes from the airborne VERDI campaign (Bierwirth et al., 2013; Schäfer et al., 2015, 2017a) conducted in April/May 2012 are exploited in this study. VERDI was based in Inuvik, Canada. All data were observed aboard the Polar 5 research aircraft of the Alfred–Wegener–Institute, Helmholtz Centre for Polar and Marine Research (AWI).

125 The measurement flights were mainly carried out in the region over the Beaufort Sea, which was mostly covered by sea ice but also included sea-ice free areas (Polynias). Mostly stratiform low level liquid and mixed-phase clouds within a temperature range of -19°C to 0°C where investigated

(Costa et al., 2017). Here, the analysis is focused on a persistent cloud layer probed on four consecutive days from 14 to 17 May 2012. The applied measurements were performed in close vicinity

130 (≤ 50 km) over constant surface conditions (open water; Polynias). The persistent cloud layer in the respective area dissolved decreased continuously from day to day with cloud top altitude decreasing from about 880 m on 14 May to around 200 m on 17 May (Klingebiel et al., 2015; Schäfer et al., 2015, 2017a).

The Polar 5 research aircraft was equipped with a set of cloud and aerosol in situ and remote sens-

135 ing instruments (Bierwirth et al., 2013; Schäfer et al., 2015; Klingebiel et al., 2015). Atmospheric profiles of temperature, humidity, wind speed and direction were derived from dropsonde measurements, which were regularly released during all flights.

2.2 Horizontal fields of cloud optical thickness

- The qualitative and quantitative description of the cloud inhomogeneities is performed using fields
 of cloud optical thickness τ. Marshak et al. (1995), Oreopoulos et al. (2000), or Schröder (2004)
 proposed to study horizontal cloud inhomogeneities using cloud-top reflectances. However, Schäfer et al. (2017a) pointed out that radiance measurements include the information of the scattering phase function (e.g., forward–/backward scattering peak, halo features). To avoid artifacts in the inhomogeneity analysis from such features, parameters that are independent of the directional scattering of
 the cloud particles have to be analysed. Therefore, to characterize the observed and simulated cloud
- 145 the cloud particles have to be analysed. Therefore, to characterize the observed and simulated cloud fields regarding their horizontal cloud inhomogeneities the cloud optical thickness is applied, which does not include the fingerprint of the scattering phase function.

The 2D fields of τ used for the comparison with COSMO are retrieved from 2D fields of reflected solar spectral radiance, which were collected with the imaging spectrometer AisaEAGLE (Schäfer

- 150 et al., 2013, 2015). Using those data, Schäfer et al. (2017a) retrieved ten fields of cloud optical thickness τ (data set published on PANGAEA, Schäfer et al., 2017b). From those available ten fields of τ , four cases are selected for the comparison to the LES results obtained from COSMO. Figure 1 exemplary illustrates selected sections (1.2 by 3.0 km) of the four chosen cases. The full widths and lengths of the applied fields of τ range to up to 1.7 km and 26.8 km, respectively. Their spatial reso-
- 155 lution is 2.6 to 3.6 m (depending on the distance between aircraft and cloud). During the time period from 14 to 17 May 2012, τ decreased from 8.1 ± 1.2 to 4.3 ± 0.4 (compare Tab. 2, Schäfer et al., 2017a). The selected sections in Fig. 1 illustrate the influence of the temporal evolution on the cloud features. In particular, from 15 to 17 May 2012 a reduction of the horizontal cloud inhomogeneity occurs, which is confirmed by Schäfer et al. (2017a). They also found a contin-
- 160 uous reduction of cloud inhomogeneity during those four consecutive days. Furthermore, directional features, which are prominent on 14 May, seem to be reduced, which is confirmed by autocorrelation analysis performed by Schäfer et al. (2017a).



Figure 1. Exemplary <u>depicted selected</u> sections (1.2 by 3.0 km) of horizontal fields of τ to illustrate the daily variability of the horizontal cloud inhomogeneities during the VERDI campaign on (a) 14 May 2012, (b) 15 May 2012, (c) 16 May 2012, and (d) 17 May 2012. Data adapted from Schäfer et al. (2017b).

2.3 Atmospheric profiles

- During each measurement flight Vaisala dropsondes (type RD94) were used together with the Vaisala
 AVAPS (Airborne Vertical Atmosphere Profiling System) dropsonde receiving system (Hock and Franklin, 1999; Coleman, 2003). The dropsondes were released to sample profiles of meteorological parameters (air pressure p, air temperature T, relative humidity RH, wind speed v, and wind direction WD) below the aircraft, which then was typically operating at about 3 km altitude and allowed to sample the entire cloud and ABL structure by the dropsondes. The accuracy of the dropsonde
- 170 measurements is given by the manufacturer and specified to ± 0.4 hPa for the air pressure, $\pm 0.2^{\circ}$ C for the air temperature, $\pm 2\%$ for the relative humidity, and ± 0.5 m s⁻¹ for the detected wind speed. For the analysis of the cloud fields, the dropsonde releases closest to the four investigated remote sensing observations had been chosen. The potential temperature (Θ), relative humidity (RH), and RH), wind speed (v), and wind direction (WD) profiles for the four investigated cases are displayed
- 175 in Fig. 2. From 14 May to 15 May the cloud top inversion increased from 810 m to 880 m while for the subsequent two days, the inversion layer decreased to 440 m on 16 May and to 200 m on 17 May 2012. In conjunction with the decrease of the cloud top altitude the cloud base altitude decreased as well until it almost reached the surface on 17 May. The relative humidity, displayed in Fig. 2b confirms the initial increase and consecutive decrease of the cloud top and base altitude. The inversion



Figure 2. (a, d) (a, e) Potential temperature(Θ), (b, d) (b, f) relative humidity(RH), and (c, f) (c, g) wind speed(v), and (d, h) wind direction for the four investigated cases. The dropsonde data is shown in the first row (a-c) (a-d) and the 2 h domain-averaged profiles after spin-up time of the simulations are shown in the second row (d-f)(e-h). Dropsondes were released closest to the imaging spectrometer measurements.

180 strength increased over the time period from ≈ 5 K to ≈ 1 K mainly because the temperature of the surface layer continuously decreased; the ABL became more stable.

Furthermore, Fig. 2c illustrates that the near-surface wind increased during the four days from ≈ 1 to $\approx 10 \text{ m s}^{-1}$, which might be of interest in terms of the generation of cloud inhomogeneities. Except for the case on 14 May, where wind speeds in higher altitudes are larger compared to the other days,

185 the daily increase of the near-surface wind speed is also observed in higher altitudes to up to 1 km. Following Jacobson et al. (2013), this is related to Low-Level-Jets (LLJ) for the days from 15 to 17 May.

3 Simulations

3.1 COSMO: General setup

- 190 COSMO is a non-hydrostatic, limited-area atmospheric forecast model (Schättler et al., 2015). Here it is used in a semi-idealized LES setup, which follows the description by Loewe et al. (2017), based on Ovchinnikov et al. (2014) and Paukert and Hoose (2014). The two-moment cloud microphysics scheme by Seifert and Beheng (2006) predicts the number densities and the masses of six hydrometeor types. The different ice phase hydrometeor growth processes are
- 195 parameterized in this scheme. In COSMO, the radiative transfer is described by a two-stream

radiation scheme after Ritter and Geleyn (1992). It is calculated every 2s and has a direct cloud-radiative feedback. A three-dimensional prognostic turbulence scheme describes the turbulent fluxes of heat, momentum and mass by a first-order closure after Smagorinsky and Lilly (Herzog et al., 2002; Langhans et al., 2012). The size of the model domain is-used by

- 200 Loewe et al. (2017) was 6.4×6.4 km in horizontal direction with a grid spacing spatial resolution of 100 m. Here, this setup is applied as well. However, analyzing cloud inhomogeneities requires a fine horizontal spatial resolution of the model simulations. Therefore, for the comparison with the imaging spectrometer measurements analyzed here, the spatial resolution is also increased to 50 m for addition model runs. In those cases, the domain size is reduced to 32 by 32 grid points
- 205 (1.6 km× 1.6 km) for computational constrains. A further reduction of the spatial resolution was not possible due to numerical instabilities. The vertical height range of 22 km is divided into 166 vertical levels, which are concentrated on more dense for the ABL with a typical grid spacing vertical resolution of around 15 m up to the inversion height of the different days of investigation. The initialization profiles of temperature, humidity, wind speed, and wind direction are based on the dropsonde
- 210 data, whereby. The dropsonde data are partly affected by horizontal variability, when slowly passing the cloud and drifting horizontally. Therefore, parts of the original profiles (Fig. 2) are smoothed to avoid numerical issues and brought to a vertical monotonically increasing profile for initialization of the model. The surface of the model is sea water and the surface fluxes depend on the surface temperature, which is 273.5 K for the sea-water surface. Moreover, ERA (European Reanalysis) -
- 215 Interim reanalysis data (from the European Centre for Medium-Range Weather forecast (ECMWF)) (Dee et al., 2011) have been used to complete the profiles above the altitude where the dropsondes were released. Other model parameters such as the description of the large scale subsidence, which is adjusted to the temperature inversion height, the relaxation to fixed cloud droplet number concentration (CDNC) and ice crystal number concentration (ICNC), and the spin up time of 2 h fol-
- 220 lows Ovchinnikov et al. (2014). The CDNCs are based on measurements of the Small Ice Detector Small Ice Detector mark 3 (SID3) measurements (Vochezer et al., 2016). During the four investigated days, CDNC of 90 to 100 cm⁻³ were observed as summarized in Tab.-1. Unfortunately, the concentration of ice crystals was below or at the detection limit of the SID3. Therefore, the ICNC were assumed to be one particle per liter according to observations of mixed-phase Arctic stratus
- 225 during the Indirect and Semi-Direct Aerosol Campaign Indirect and Semi-Direct Aerosol Campaign (ISDAC) (McFarquhar et al., 2011; Ovchinnikov et al., 2014). The inversion height of the temperature $z(T_{in})$ is necessary for the description of the large scale-large-scale subsidence in the model and is represented by the inversion height of the dropsonde profiles, which are used for initialization of the model simulations (Tab.- 1).

 Table 1. Model setup specifications of the different mixed-phase cloud simulations of four VERDI campaign days.

Case	$z(T_{\rm in})$ [m]	CDNC [cm ⁻³]	ICNC [l ⁻¹]
14 May	870	100	1
15 May	988	100	1
16 May	440	90	1
17 May	350	100	1



Figure 3. Domain averages of LWC (blue color scale) and IWC (red-yellow color scale) of the four simulations during the VERDI campaign. Please note the different color scale for the IWC in (d).

230 3.2 Domain-averaged cloud properties and temporal evolution

Time series of simulated liquid water content (LWC) and ice water content (IWC) for the four selected cases are shown in Fig. 3. During the four flights, which are simulated with COSMO, only few ice crystals were observed. In terms of the model domain average profiles of the LWC and IWC, the simulated clouds consist mostly of liquid water droplets except for the 15 May, in which more IWC

is built from around 4 h on (Fig. 3b). Furthermore, the cloud top is around 1000 m for the 14 May and the 15 May (Fig. 3a, b). However, the cloud top height increases during time in all four simulations because of entrainment of air through the top of the ABL. This is evident in the temporal evolution of LWC, which has a maximum between 0.25 and 0.35 g kg⁻¹ near the cloud top. The Arctic clouds on 16 May and 17 May are the lowest simulated clouds with a cloud top initially around 450 m and

240 350 m, respectively (Fig. 3 c, d).

The four simulations show differences in the temperature, relative humidity and wind speed profiles (Fig. 2d-fe-g), which in general still agree with the initial dropsonde profiles after the spin up time (Fig. 2a-c). The height of the ABLs and the strength of the inversions are lower in the simulations of the 16 May and 17 May. Furthermore, for the simulation on 17 May a second inversion develops in

245 the ABL near the surface around 60 m to 150 m. The ABL structure is well mixed in the simulation of the 16 May although no second temperature inversion is built near the surface. The simulation of the 16 May shows a wind shear from around 150° to around 100° (Fig. ??2g) and a decrease of v with height above the cloud top height, which is also seen in the dropsonde profiles (Fig. 2cand Fig. ??). The other simulations do not show a turning of the wind directly above the inversion height.

- 250 The simulated mixed-phase clouds of the four VERDI flights show a liquid water path (LWP) around 35 to 50 g m⁻². The highest LWP is seen in the simulation of the 14 May, which increases towards 50 g m⁻² at the end of the simulation. The simulation of the 15 May has the lowest LWP values. Furthermore, the LWP remains very stable until the end of the simulation. The ice water path (IWP) and the snow water path (SWP) of all four simulations is small especially for the simulated clouds on the 14, 16, and 17 May, which fits well with observations.
- Domain averaged WD (a, c, c, g) and v (b, d, f, h) of the four VERDIsimulations. The black line marks the averaged cloud top height during the simulation. For the comparison of the simulated and observed horizontal cloud structures (cloud inhomogeneities), fields of simulated cloud optical thickness (τ_{sim}) are compared to retrieved fields of cloud optical thickness from the measurements (τ_{meas}).
- 260 The τ_{sim} is calculated within the COSMO model considering the amount of liquid water and the solar spectrum. However, it cannot be expected that COSMO is capable of reproducing the detailed spatial and temporal cloud evolutions, which are captured by the observed fields of τ , accurately (inhomogeneity features and directional structures). Therefore, besides the comparison of observed and simulated clouds with regard to macrophysical cloud features (cloud vertical extent, cloud optical
- 265 thickness) of the individual cases, instead of point-by-point comparisons of cloud parameters, statistical bulk parameters describing the horizontal cloud inhomogeneities, their directional structures, and the temporal evolution of both will be compared.

4 Quantification of cloud inhomogeneities

4.1 One-dimensional statistical bulk parameters

- 270 For the quantitative description of the cloud inhomogeneities from the simulated fields of cloud optical thickness (τ_{sim}) obtained from COSMO and measurement-based retrieved fields of cloud optical thickness (τ_{meas}) collected during the VERDI campaign, statistical techniques are applied. Following Schäfer et al. (2017a), different statistical quantitative measures of the cloud inhomogeneities are derived using the mean and standard deviation of the particular τ field and three 1D inhomogeneity
- 275 parameters ρ_{τ} (Davis et al., 1999b; Szczap et al., 2000), S_{τ} (Davis et al., 1999b; Szczap et al., 2000), and χ_{τ} (Cahalan , 1994; Oreopoulos and Cahalan, 2005). They are given by:

$$\rho_{\tau} = \frac{\sigma_{\tau}}{\bar{\tau}},\tag{1}$$

$$S_{\tau} = \frac{\sqrt{\ln(\rho_{\tau}^2 + 1)}}{\ln 10},$$
(2)

280

285

$$\chi_{\tau} = \frac{\exp\left(\bar{\ln\tau}\right)}{\bar{\tau}}.$$
(3)

A homogeneous cloud is characterized by $\rho_{\tau} = 0$ and $S_{\tau} = 0$. Higher values of ρ_{τ} and S_{τ} indicate more pronounced cloud inhomogeneity. However, both of them have no predefined upper limit. Therefore, ρ_{τ} and S_{τ} only sustain a quantitatively significance, when their values for different cases are compared to each other. The 1D inhomogeneity parameter χ_{τ} ranges between 0 and 1, with values close to unity indicating horizontal homogeneity and values approaching zero characterizing high horizontal inhomogeneity. Due to the limited range between 0 and 1, χ_{τ} is not only a qualitative

4.2 Two-dimensional autocorrelation analysis

but also quantitative measure.

- 290 Two-dimensional autocorrelation analysis is applied to quantify the typical scales of cloud inhomogeneities and to identify directional patterns of the cloud structure (Schäfer et al., 2017a). To derive the autocorrelation functions, each field of τ is correlated with itself, while it is shifted pixel by pixel (observations) or grid point by grid point (simulations) against itself. The values of the resulting correlation coefficients after each shift are in the range between -1 (perfect negative correlation)
- and 1 (perfect positive correlation). Correlation coefficients with values of 0 identify no correlation. Here, only the degree of correlation matters, not if it has a positive or negative sign. Similar to Schäfer et al. (2017a), squared autocorrelation functions P_{τ}^2 are used to avoid ambiguous interpretations. The P_{τ}^2 reach values between 0 (no correlation) and 1 (perfect correlation).
- The particular correlation coefficients at the derived distances identify the similarity of the horizontal cloud structures. If the cloud is horizontally homogeneous, the correlation coefficients stay constant over large distances. If the cloud is rather inhomogeneous the correlation coefficients already drop at closer distances. Therefore, P_{τ}^2 as a function of distances is a measure of the size of the dominant cloud structures.
- A quantitative value for the distance at which cloud structures are different from each other (namely 305 decorrelated) is the decorrelation length ξ_{τ} (Schäfer et al., 2017a). It is the distance at which P_{τ}^2 drops to:

$$P_{\tau}^{2}(\xi_{\tau}) = \frac{1}{e^{2}}.$$
(4)

In a 2D-autocorrelation function, ξ_{τ} can differ depending on the orientation, if the cloud structures have a predominant orientation. To quantify this directionality, ξ_{τ} is calculated along (ξ_{τ}^{\uparrow}) and across

310 $(\xi_{\tau}^{\leftrightarrow})$ the predominant direction. The larger the differences between ξ_{τ}^{\uparrow} and $\xi_{\tau}^{\leftrightarrow}$, the more cloud structures are orientated.

Figure 4a shows a section of an observed field of τ_{meas} , retrieved from the measurements on 15 May. The <u>depicted selected</u> section has a swath of 1.3 km (oriented in y direction) and a length of 6 km



Figure 4. (a–b) Horizontal fields of normalized τ_{meas} (VERDI) and τ_{sim} (COSMO) for the case on 15 May 2012. (c–d) Two-dimensional autocorrelation coefficients $P_{\tau,\text{meas}}^2$ and $P_{\tau,\text{sim}}^2$, calculated for fields of τ displayed in (a) and (b). (e–f) One-dimensional autocorrelation coefficients along (straight white line marked in (c) and (d)) and across (dashed white line marked in (c) and (d)) predominant directional structure. The grey dotted line illustrates the threshold for the estimation of ξ_{τ}^{\uparrow} and $\xi_{\tau}^{\leftrightarrow}$.

315

(oriented in x direction). Figure 4b shows the corresponding field of τ_{sim} (6 km × 6 km, adapted to the selected length of the measurement case), which is simulated with COSMO two hours after the spin up time for the case on 15 May. For comparability reasons, both fields of τ are normalized by their maximum.

Although the swath (y direction) of the field of au_{meas} is smaller by a factor of almost five compared

to the field of $\tau_{\rm sim}$, larger cloud structures of similar size and shape are obvious in both fields of

- 320 τ_{meas} and τ_{sim} . However, with 488 spatial pixels along the swath (spatial double binning was applied during measurements) and a field of view of 37° AisaEAGLE's spatial resolution is ≈ 1.3 m for a target in a distance of 1 km. Thus, the spatial resolution of AisaEAGLE is relatively largehigh, compared to the grid spacing spatial resolution of 100 m from COSMO. Thereby, the exact pixel size of AisaEAGLE depends on the distance between aircraft and cloud, which leads to pixel sizes
- between 2.6 and 3.6 m for the four investigated cases. Due to the 30 to 40 times higher spatial resolution of AisaEAGLE, compared to COSMO's grid spacing, the measurements shows cloud features, which cannot be resolved by COSMO. Those features on a spatial scale below 100 m may have an effect on the statistical (1D inhomogeneity parameters) and spatial comparison (autocorrelation analysis) of the particular fields of τ .
- To quantify the size and orientation of the represented cloud structures in the observations and simulations, Fig. 4c and Fig. 4d show the calculated squared 2D autocorrelation coefficients P²_τ. To calculate them, different numbers of legs (shifts) have to be applied for P²_{τ,meas} and P²_{τ,sim}. The applied field of τ_{meas} consists of 2700 × 450 spatial pixels. Therefore, restricted to the shorter side, 225 × 225 (half of swath pixel number, calculated into x and y direction) legs are chosen for the calculation of the 2D P²_{τ,meas}. COSMO consists of 64 × 64 grid points. This allows 32 × 32 legs for

the calculation of $P_{\tau,\text{sim}}^2$. The resolved domain and pixel/grid-point sizes spatial resolution displayed in Fig. 4c and Fig. 4d

show significant differences, which reveals that a direct comparison is difficult. Applying the 2D autocorrelation analysis to the observations allows to resolve small-scale cloud structures with high

- spatial resolution ($\approx 2.7 \text{ m}$), but only within a narrow spatial range below 1 km. Contrarily, the same analysis for COSMO delivers $P_{\tau,\text{sim}}^2$ with lower spatial resolution ($\geq 100 \text{ m}$), but over a larger spatial range ($\leq 3.2 \text{ km}$, in Fig. 4d only displayed until 2 km). Thus, also large-scale large-scale cloud structures are covered by COSMO (purple stripes in Fig. 4d) but not in the observations. Therefore, to make a direct comparison possible the large-scale structures cannot be compared between
- 345 observations and simulations. With respect to a comparison of the small-scale structures, the spatial sizes (pixel/grid point sizespatial resolution, domain size) of both datasets need to be conformed to make a direct comparison possible.

Furthermore, both, Fig. 4c and Fig. 4d show predominant directional features of the cloud structures. Their lengths and widths are derived from 1D autocorrelation functions along (straight white line in

- 350 Fig. 4c and Fig. 4d) and across (dashed white line in Fig. 4c and Fig. 4d) those predominant directional structures and a subsequent estimation of ξ[‡]_τ and ξ[↔]_τ. The derived ξ[‡]_τ and ξ[↔]_τ show an overall agreement but still differ from each other. For the observations ξ[‡]_{τ,meas} and ξ[↔]_{τ,meas} reach distances of ≈ 500 m and ≈ 250 m, respectively. Contrarily, for the simulations ξ[‡]_{τ,sim} and ξ[↔]_{τ,sim} reach distances of ≈ 800 m and ≈ 400 m, respectively. This might be is a further indication that it is necessary to
- make the fields of τ_{meas} and τ_{sim} conform with regard respect to their spatial resolution and domain.

In the following this is done by (i) averaging the observed fields of τ_{meas} to the spatial resolution of the simulated fields of τ_{sim} and (ii) improving the spatial resolution of the simulations itself. Figure 4e and Fig. 4f further illustrate that it is not possible to compare the large–scale structures between observations and simulations. The large–scale structures, which are covered by the COSMO

360 simulations are identified by a second increase of the $P_{\tau,\text{sim}}^2$ at distances ($\approx 1 \text{ km}$ in Fig. 4f) larger than ξ_{τ} . The width of the measured fields is too narrow to cover such a second increase in the $P_{\tau,\text{meas}}^2$ (compare Fig. 4e). Therefore, the further comparison of the cloud structures, which are identified in the observations and simulations, is restricted to the small–scale cloud structures with sizes below 1 km only.

365 4.3 Final data preparation - Domain adjustmentAdjustment of spatial resolution and domain

To compare both data sets, the fields of τ_{meas} , which are retrieved from the imaging spectrometer measurements are averaged to the spatial resolution of the COSMO τ_{sim} fields. The investigations on the single cases during VERDI are performed for spatial resolutions of 50 m (32 by 32 grid points) and 100 m (64 by 64 grid points). All other model parameters are kept constant with respect to the

analysis performed by Loewe et al. (2017).

In order to do soaverage the observed fields of τ_{meas} to the spatial resolution of 50 and 100 m, the τ_{meas} -values of distinct numbers of neighbouring neighboring pixels are averaged. The number depends on the single pixel size of the particular cases, which is a function of the distance between aircraft and cloud. For the four investigated cases this number varies between 13 (26and 37-) and

- 375 <u>18 (36)</u> pixels, which are needed to generate pixel sizes of τ_{meas} comparable to the <u>50 m (100 mgrid spacing</u>) spatial resolution of COSMO.
 Furthermore, for the simulations with 100 m spatial resolution, the domain size of the measurements and simulations are different. need to be adapted. The applied COSMO's domain size of 6.4 km by
- 6.4 km is about three to four times larger than the domain size of the measurements. Therefore, to
 compare both data sets, the COSMO domain size is also reduced to the width and length of the corresponding τ_{meas} field from the measurements. Therefore, for the comparison, only a squared domain in the center of COSMO's τ_{sim} field is used, which size corresponds to the size of the particular field from the measurement. For the four investigated cases this results in COSMO domains composed out of 12 × 12 to 16 × 16 grid points (1.2 × 1.2 km to 1.6 × 1.6 km). Longer stripes of τ_{meas}-fields
- and stripes according to their lengths across the COSMO domain are not used, because the investigations are focused on small scale cloud inhomogeneities, which are already covered by the smaller squared domain size given by the swath of the τ_{meas} -fields.

For the COSMO simulations, which use 50 m spatial resolution, the domain size is reduced to 32 by 32 grid points resulting in a total domain of 1.6 km by 1.6 km, which is comparable to the

390 observations. Therefore, the domain of those simulations was not adapted for the comparisons.However, to increase the statistics, which might be otherwise too small because of the finally applied



Figure 5. Illustrated are sections of one and the same field of τ_{meas} from 14 May 2012 with a spatial resolutions of (**a**) \approx 3 m (original resolution), (**b**) 3050 m (COSMO resolution), (**c**) 100 m (COSMO resolution), (**d**) 150 m, and (**e**) 300 m. (**f**-**j**) Squared 2D autocorrelation coefficients P_{τ}^2 calculated for the fields of τ_{meas} displayed in (a) to (e). (**k**-**o**) Squared 1D autocorrelation coefficients P_{τ}^2 calculated along straight red line in (f) to (j). Estimated decorrelation length ξ_{τ} is marked by horizontal and vertical black line and labeled by its value. Red dot marks ξ_{τ} as derived from the case with the original spatial resolution of 3 m.

small domain but large pixel sizes, for COSMO averages of the resulting $P_{\tau,\text{sim}}^2$ over all output time steps after spin up are used. For the measured fields, whose which lengths are much longer than their widths, squared domains (size determined by swath of τ_{meas}) are cut along the measured stripe and the resulting $P_{\tau,\text{meas}}^2$ are averaged accordingly. Increasing the number of available $P_{\tau,\text{meas}}^2$ to average is a further restriction to use squared domains instead of stripes.

395

To test possible effects arising from the change of spatial resolution and to check if the relevant scales of cloud inhomogeneity are lost, when reducing the resolution of the measurements, Fig. 5a to Fig. 5e show sections of one and the same field of τ_{meas} from 14 May, but displayed with

- 400 a different spatial resolution of 3 m (original resolution), 3050 m (COSMO fine resolution), 100 m (COSMO original resolution), 150 m, and 300 m resolution. Figure 5f to Fig. 5j show the corresponding squared 2D autocorrelation coefficients. The red line illustrates the direction, which is used to calculate the squared 1D autocorrelation functions and decorrelation lengths ξ_{τ} displayed in Fig. 5k to Fig. 50. The fields from the 2D autocorrelation analysis show that except for the spatial resolution
- of 300 m the directional structure of the cloud inhomogeneities is still captured, when the spatial resolution is reduced. However, the decorrelation lengths, derived from the 1D autocorrelation analysis increases with decreasing spatial resolution from $\xi_{\tau} = 327$ m at 3 m spatial resolution to $\xi_{\tau} = 600$ m



Figure 6. Comparison of (a) mean and standard deviation and (b) inhomogeneity parameters ρ and S as a function of spatial resolution for the fields of τ_{meas} illustrated in Fig. 5a–e.

at 300 m spatial resolution. Therefore, decreasing spatial resolution leads to larger ξ_{τ} , which indicates larger cloud structures. This means that reduced spatial resolution will generate fields of τ with

410 larger spatial scales.

420

To test the influence of the spatial resolution on the overall inhomogeneity, Fig. 6a shows the results for the mean and standard deviation of the fields of τ , illustrated in Fig. 5. Figure 6b shows the corresponding 1D inhomogeneity parameters ρ_{τ} and S_{τ} . While the mean value of τ stays constant for all spatial resolutions, its standard deviation decreases with increasing pixel size. This indicates

415 that the fields of τ become more homogeneous the larger the pixel size is. Comparably Similarly, the value of both 1D inhomogeneity parameters ρ_{τ} and S_{τ} decrease with increasing pixel size, which further confirms the reduction of the inhomogeneity.

Therefore, in the following analysis, comparing the simulated against observed fields of τ , the simulations with the finer spatial resolution of 50 m are used. The simulations with 100 m spatial resolution are used to discuss the model sensitivity with respect the spatial resolutions.

5 Comparison of modeled against observed cloud structures

5.1 Magnitude of inhomogeneity

The fields of τ obtained from the spectral imaging remote sensing (τ_{meas}) are compared to the fields of τ derived from the COSMO simulations (τ_{sim}). To validate the cloud inhomogeneity in the simulated fields, the statistical techniques from Sect. 4.1 including the averaging of the measured fields to 50 and 100 m pixel size are applied. Tab.Table 2 lists the mean value of τ , standard deviation σ_{τ} , and the three 1D inhomogeneity parameters ρ_{τ} , S_{τ} , and χ_{τ} for the simulations and the simulations with the two different spatial resolutions of 50 and 100 m.

430 Both, measurements and simulation show the highest cloud optical thickness on 14 May with $\bar{\tau}_{\text{meas}} = 8.1 \pm 1.2$ and $\bar{\tau}_{\text{sim}} = 7.9 \pm 0.6$ at 50 m spatial resolution and $\bar{\tau}_{\text{sim}} = 6.9 \pm 0.5$ at 100 m

Table 2. Mean value of τ , standard deviation σ_{τ} , and the three 1D inhomogeneity parameters ρ_{τ} , S_{τ} , and χ_{τ} calculated for all four cases from the observations and the simulations with the two different spatial resolutions of 50 and 100 m.

	Case	$\bar{\tau} \pm \sigma_{\tau}$	$ ho_{ au}$	S_{τ}	$\chi_{ au}$
VERDI (<u>50 m</u>)	<u>14 May</u>	7.8 ± 1.5	0.195	0.086	<u>0.979</u>
	<u>15 May</u>	$\underbrace{6.4\pm0.7}$	0.121	0.055	0.992
	<u>16 May</u>	$\underbrace{6.4 \pm 1.0}_{}$	0.166	0.078	0.983
	<u>17 May</u>	$\underbrace{4.2\pm0.5}_{}$	0.154	0.071	0.986
<u>VERDI (100 m)</u>	14 May	8.1 ± 1.2	0.209	0.093	0.977
	15 May	6.4 ± 0.5	0.115	0.052	0.993
	16 May	6.6 ± 0.6	0.145	0.065	0.988
	17 May	4.3 ± 0.4	0.132	0.061	0.990
COSMO (<u>50 m</u>)	<u>14 May</u>	7.9 ± 0.6	0.071	0.030	0.997
	<u>15 May</u>	7.1 ± 0.7	0.092	0.040	0.995
	<u>16 May</u>	$\underline{6.0\pm0.6}$	0.094	0.040	0.995
	<u>17 May</u>	5.8 ± 0.5	0.083	0.036	0.996
<u>COSMO (100 m)</u>	14 May	6.9 ± 0.5	0.066	0.028	0.997
	15 May	5.4 ± 0.3	0.053	0.023	0.998
	16 May	5.5 ± 0.5	0.090	0.037	0.996
	17 May	5.6 ± 0.3	0.044	0.019	0.999

spatial resolution, which show an overall agreement. During the course of the following days, the large scale subsidence lead to a decrease of the cloud top altitude and cloud geometrical thickness and corresponding lower values of τ and σ_{τ} . For these days, model and observations are still in agreement. However, compared to the spatial resolution of 100 m it is obvious that the finer resolved

simulations lead to better agreements between measurements and simulations. Regarding the cloud inhomogeneity, the absolute values of the 1D inhomogeneity parameters ρ_{τ} , S_{τ} , and χ_{τ} do not compare well for the simulations with 100 m spatial resolution. The results for the COSMO simulations show lower 1D inhomogeneity parameters (more homogeneous)

435

- 440 by a factor of two and higher, compared to the results from the measurements. The agreement between the observations and simulations increase with the finer spatial resolution of 50 m, but still does not match perfectly. The reason might be that the comparably lower inhomogeneity derived from COSMO might be for both spatial resolutions is caused by its effective grid spacingspatial resolution, which is approximately three times 50 m or accordingly three times 100 m (Skamarock
- 445 et al., 2004). Although the pixel size of AisaEAGLE is adapted to the COSMO grid spacing spatial resolution by averaging over neighboring pixels, COSMO's effective grid spacing spatial resolution is larger, which might lead to larger homogeneity of the simulations compared to the observations.

Furthermore, COSMO simulates the cloud at the same location, where it is initialized. Contrarily, the AisaEAGLE measurements took place along a stripe of several kilometers. The simulated clouds

- may not change in between the time steps as much as the measurements of the clouds along the measurement stripe do. Therefore, averaging over COSMO's time steps might further produce more homogeneous results than averaging over AisaEAGLE's squared domains along the flight track. However, the observations show that the cloud field became more homogeneous from 14 to 15 May as indicated by lower values of ρ_τ(ρ_τ reduces, which reduce from 0.209 to 0.115). From 15 to
- 455 16 May, ρ_{τ} increases to 0.145, which indicates a cloud field with slightly higher inhomogeneity. Then, on 17 May, ρ_{τ} reduced to 0.132, showing that the cloud field became more homogeneous again. This trend is reproduced by the These different cases with high and low ρ_{τ} are reproduced by COSMO results, except that ρ_{τ} increases to larger values on 16 May than independent on the chosen spatial resolution. Larger discrepancy between modeled and observed inhomogeneity parameters
- 460 only occurred on 14 May, which is not seen from the observations when the observations were influenced by large-scale cloud structures. This trend of decreasingNevertheless, the lower/increasing higher inhomogeneity is also imprinted in the inhomogeneity parameters S_{τ} and χ_{τ} , which decrease are smaller/increase with time larger in both, measurements and simulations. In this regard, indicating that COSMO performs well with

465 regard to the 1D inhomogeneity parameters.

5.2 Spatial inhomogeneity scale

- The 2D autocorrelation functions are calculated to compare the typical spatial scales and the directional character of the eloud inhomogeneities small-scale cloud inhomogeneities (no large-scale inhomogeneities like roll convection) of observations and simulations. The 2D autocorrelation coefficients (P²_{τ,meas}; P²_{τ,sim}) for each case are shown in Fig. 7e to Fig. 7h for the measurements and in Fig. 7m to Fig. 7p for the simulations (50 m spatial resolution). Additionally, representative fields of normalized τ_{meas} (Fig. 7a–d) and τ_{sim} (Fig. 7i–l) are added. The 2D autocorrelation analysis was applied to the simulated fields of τ_{sim} orientated in a North-South and West-East grid. The orientation
- of the observations is determined by the flight direction. Therefore, the orientation of the fields of τ_{meas} and $P_{\tau,\text{meas}}^2$ are rotated into the direction of the COSMO grid. One-dimensional P_{τ}^2 are calculated manually along the dominant direction (straight red and blue lines in Fig. 7e–h and Fig. 7m–p) and across (dashed red and blue lines in Fig. 7e–h and Fig. 7m–p) it. For $P_{\tau,\text{meas}}^2$ (red) and $P_{\tau,\text{sim}}^2$ (blue) the results are displayed in Fig. 7i to Fig. 7l. The dotted black line illustrates the threshold for

480 the estimation of
$$\xi_{\tau}$$
.

The observations on 14 May are influenced by a large scale cloud structure, which is caused by large scale dynamic forcing and leads to an increase of the autocorrelation coefficients for distances larger than 700800 m. Furthermore, during this day a significant directional structure from



Figure 7. (a-d) Exemplary depicted selected sections of fields of τ_{meas} observed during VERDI from 14 to 17 May 2012. (e-h) Mean 2D autocorrelation coefficients $P_{\tau,\text{meas}}^2$ derived for fields of τ_{meas} from VERDI. (i-l) Exemplary depicted selected fields of τ_{sim} simulated with COSMO (50 m spatial resolution) for the VERDI cases from 14 to 17 May 2012. Dashed red boxes illustrate corresponding domain of observed fields of τ_{meas} . (m-p) Mean 2D autocorrelation coefficients $P_{\tau,\text{sim}}^2$ derived for fields of τ_{sim} . (q-t) Decorrelation length ξ_{τ} along strongest (straight blue and red lines) and weakest (dashed blue and red lines) extend of 2D autocorrelation coefficients derived from $P_{\tau,\text{meas}}^2$ in (m-p), respectively.

North-West to South-East is observed. Along this direction the cloud field stays homogeneous over

a wide range ($\xi_{\tau} = 700\xi_{\tau} = 800$ m). Across this predominant structure the small-scale cloud structures reach a decorrelation length of $\xi_{\tau} = 300$ m. During the following days the orientation of the directional structure turns eastwards in the observations and the difference differences between ξ_{τ}^{\ddagger} and $\xi_{\tau}^{\leftrightarrow}$ decreases decrease. This characterizes a weakening of the directional structure of the cloud field. Only on 16 May a directional orientation can be identified, while on 15 and 17 May the clouds

490 are characterized by a rather indifferent horizontal structure.

Comparing the results for $P_{\tau,\text{sim}}^2$ with $P_{\tau,\text{meas}}^2$ reveals that the large scale cloud structure is not well simulated for the case on 14 May. This results most probably from the small domain size of COSMO, which is fixed over the same location when averaging the $P_{\tau,\text{sim}}^2$ over a set of time steps. Contrarily, the averages of $P_{\tau,\text{meas}}^2$ from the measurements are performed over a set of squared domains

- 495 along the flight track. Thus, the chance to cover also larger structures is higher for the measurements compared to the simulations. However, the influences of the large scale cloud structures on the decorrelation length ξ_{τ} of the small scale cloud inhomogeneities is low and can be neglected overall small-scale directional structures are well simulated. On 14 May, a significant directional structure from North–West to South–East is observed, which then turns eastwards for 15 to 17 May. Except
- 500 on 16 May, the predominant simulated directions of the cloud fields are almost identically to the observations.

Furthermore, the results for $P_{\tau,\text{meas}}^2$ and $P_{\tau,\text{sim}}^2$ show that COSMO produces larger simulations using a spatial resolution of 50 m produce similar sizes of the small-scale cloud structures compared to the measurements. In Fig. 7m to Fig. 7p the covered areas of $P_{\tau,\text{sim}}^2$ are larger of similar sizes compared

- to the areas covered by $P_{\tau,\text{meas}}^2$ in Fig. 7e to Fig. 7h. Table 3 lists the resulting $\xi_{\tau,\text{meas}}$ and $\xi_{\tau,\text{sim}}$ calculated along (ξ_{τ}^{\uparrow}) and across $(\xi_{\tau}^{\leftrightarrow})$ the predominant structures found in Fig. 7e–h and Fig. 7m– p. A comparison reveals <u>only minor</u> differences between $\xi_{\tau,\text{meas}}$ and $\xi_{\tau,\text{sim}}$. The <u>values from the</u> simulations (except for ξ_{τ}^{\uparrow} on 14 May) are larger compared to the values from the observations by 20 to 30 %. For all four cases from the observations and simulations calculated decorrelation lengths
- 510 $\xi_{\tau,\text{meas}}$ and $\xi_{\tau,\text{sim}}$ along (ξ_{τ}^{\uparrow}) and across $(\xi_{\tau}^{\leftrightarrow})$ the predominant directions found in Fig. 7e-h and Fig. 7m-p. Case ξ_{τ}^{\uparrow} m $\xi_{\tau}^{\leftrightarrow}$ mVERDI 14 May 700 290 best agreement is achieved on 15 May 280 190 16 May 350 170 and 17 May370 260 COSMO 14 May 530 320 15 May 380 260, when $\xi_{\tau,\text{meas}}$ and $\xi_{\tau,\text{sim}}$ show almost identically results. On 16 May 500 280 17 May 430 390 However, the $P_{\tau,\text{sim}}^2$ follow the trend, which is observed for $P_{\tau,\text{meas}}^2$, including their predominant directional structure. the
- 515 differences are slightly larger, while on 14 May the differences are significantly larger, which might result from the insufficient simulated large-scale cloud structure. For the simulations with 100 m spatial resolution (graph not shown) the directional features still compare well between observations and simulations. Like for the measurements on 14 May a predominant North–West to South–East direction is simulated, which then turns eastwards. Thereby, the cases on 14 May and 16 May show
- 520 the strongest directional features (largest differences between ξ_{τ}^{\uparrow} and $\xi_{\tau}^{\leftrightarrow}$), compare Tab. 3) with ξ_{τ}^{\downarrow}

	Case	$\xi_{\mathrm{T,5Qm}}^{\uparrow}$ [m]	$\xi_{\tau,50m}^{\leftrightarrow}$ [m]	$\xi^{\ddagger}_{\tau,100m}[m]$	$\xi_{\tau,100m}^{\leftrightarrow}[m]$
VERDI	14 May	800	330	>1000	400
	<u>15 May</u>	260	180	280	190
	<u>16 May</u>	<u>220</u>	100	350	170
	<u>17 May</u>	2 <u>50</u>	150	370	260
COSMO	14 May	<u>260</u>	190	530	320
	<u>15 May</u>	250	200	380	260
	<u>16 May</u>	270	180	500	280
	<u>17 May</u>	240	<u>190</u>	430	390

Table 3. Calculated decorrelation lengths $\xi_{\tau,meas}$ and $\xi_{\tau,sim}$ for the two different spatial resolutions of 50 and 100 m along (ξ_{τ}^{\downarrow}) and across ($\xi_{\tau}^{\leftrightarrow}$) the observed and simulated predominant directions (compare Fig. 7e-h and Fig. 7m-p for 50 m spatial resolution).

on 14 May larger than the width of the observed field of τ_{meas} . Although on 17 May COSMO simulates a more isotropic structure ($\xi_{\tau}^{\uparrow} \approx \xi_{\tau}^{\leftrightarrow} \approx 400 \text{ m}$) of the cloud inhomogeneities compared to the measurements ($\xi_{\tau}^{\uparrow} = 370 \text{ m} \neq \xi_{\tau}^{\leftrightarrow} = 260 \text{ m}$) it captures the reduction of the overall directionality. Therefore, the overall results with regard to the directional structure provided by COSMO are ac-

525 ceptable. However, the covered areas of the 2D autocorrelation functions, where the values of $P_{\tau,\text{sim}}^2$ are higher than e^{-2} are larger compared to the areas covered by the particular $P_{\tau,\text{meas}}^2$. Therefore, the $\xi_{\tau,\text{meas}}$ and $\xi_{\tau,\text{sim}}$ calculated along (ξ_{τ}^{\uparrow}) and across $(\xi_{\tau}^{\leftrightarrow})$ the predominant structures do not compare well (compare Tab. 3). Like expected from Fig. 5, the values from the simulations (except for ξ_{τ}^{\uparrow} on 14 May) are larger compared to the values from the observations by 20 to 30 %.

530

535

6 Sensitivity Study

The reasons for the differences on 16 May are (Fig. 7s) are most probably related to the wind field and the temperature profile. Figure ??e 2d and Fig. ??f illustrate a development of the wind profile 2h illustrate the temporally averaged wind directions in the simulationswith simulation time. While the wind direction does not changed at the cloud top of the 14, 15, and 17 May, the simulation of the 16 May shows a turning of the wind. Together with the well-mixed ABL (Fig. 2) this case shows a typical example of a cold air outbreak roll convection (e.g., Brümmer , 1999). On 16 May the simulated wind speed is significantly higher compared to the other days, resulting from the initial

540 if the cloud is coupled to the surface and if so, affect only the LWP of the cloud (Loewe, 2017). For de-coupled clouds, it is assumed that the cloud structure depends more strongly on the wind shear, respectively the wind speed. However, since the wind speed, wind direction, and temperature profile

conditions in the dropsonde profile (Fig. 2c,d). Influences from the surface fluxes are only expected



Figure 8. (a)–(f) Exemplary depicted selected fields of τ_{sim} for the 15 May 2012 case simulated with for differently scaled initial wind speeds on a grid with 50 m spatial resolution and 1.6 km by 1.6 km domain (a–f) and on a grid with 100 m spatial resolution and with 6.4 km by 6.4 km domain (m–r). (g)–(f) Calculated 2D autocorrelation coefficients P_{τ}^2 are given for each case in (g–l) and (s–x). White lines in (as)–(fx) - Additionally, white lines illustrate the orientation used for the calculation of the 1D P_{τ}^2 along (straight white lines) and across (dashed white lines) the dominant directions illustrated in Fig.9. Red squares in (m)–(r) mark areas of comparable size to the small domains in (a)–(f).

are the only parameters, which have been changed in the model input, the wind speed and wind shear are expected to be main drivers for the degree of horizontal cloud inhomogeneity.

- 545 To test its influence on the horizontal cloud inhomogeneity, the simulation simulations for 15 May is (50 and 100 m spatial resolution) are repeated for different initializations, where the wind profile was varied. is varied. Here, the case on 15 May is chosen, because it shows the best agreement between observations and simulations (Fig. 7r) to serve as a benchmark case. Based on the original wind profile, the wind speeds at each altitude are multiplied by (a) 0.5, (b) 1.0, (c) 1.5, (d) 2.0, (e) 2.5, and (f)
- 550 3.0. This leads to mean wind speeds (vertically averaged over cloudy region) of (a) $\approx 0.7 \text{ m s}^{-1}$, (b) $\approx 1.5 \text{ m s}^{-1}$, c) $\approx 2.2 \text{ m s}^{-1}$, (d) $\approx 3.0 \text{ m s}^{-1}$, (e) $\approx 3.7 \text{ m s}^{-1}$, and (f) $\approx 4.4 \text{ m s}^{-1}$. The wind shear was kept constant throughout all simulations.

Figure 8a to Fig. 8f show the simulated 2D fields of τ_{sim} (6.4for the simulations with the domain size of 1.6 km by 1.6 6.4km and 50 m spatial resolution. Small-scale structures (≤ 0.5 km) are



Figure 9. For the six cases of different wind speed calculated 1D autocorrelation functions (**a**) along and (**b**) across the main structures, identified in Fig. 8(**g**)–8(**l**). The grey dotted line marks the threshold of $P_{\tau}^2(\xi_{\tau}) = e^{-2}$. (**c**) From (**a**) and (**b**) derived discrete values for the decorrelation lengths ξ_{τ}^{\ddagger} and $\xi_{\tau}^{\leftrightarrow}$ as a funktion of wind speed v (symbols). Additionally included are fits derived from Eq. (5) and Eq. (6) (dotted lines).

- 555 obvious and rather randomly orientated throughout the simulations for all six different initializing wind profiles. The spatial sizes of the small-scale structures quantified by the decorrelation length depend only little on the wind speed. This is confirmed by the 2D autocorrelation analysis illustrated in Fig. 8g to Fig. 8l. Displayed are only the horizontal scales below 0.8 km, quantified by the 2D autocorrelation coefficients for shifts below ± 0.8 km, A predominant direction of the small-scale
- 560 structures is only slightly developed and varies independently from cases to case without clear preference. Furthermore, the P_{τ}^2 and the decorrelation length, which vary between 150 and 300 m show only slight variations with changing wind speeds. This means that the sizes of the small-scale structures is basically independent to the wind speed.

Contrarily, the simulations with a domain size of 6.4 km by 6.4 km and 100 m spatial resolution).

- 565 Small-scale show a clear dependency on the wind speed. The corresponding 2D fields of τ_{sim} are illustrated in Fig. 8m to Fig. 8r. The small-scale structures (≤ 0.5 km) are obvious. For still obvious in the simulations with coarse resolution, but for lower wind speeds, these small-scale small-scale structures have a North-West to South-East orientation, which turns into North-East to South-West orientation with increasing wind speeds. Large-scale Additionally, large-scale structures (≥ 2 km),
- orientated perpendicular to the small-scale structures occur at 2.5 × v. The direction of these large-scale structures turns as well and becomes more obvious with increasing wind speeds.
 The fields of τ_{sim} are evaluated using related results for the 2D autocorrelation analysis . The results are given in Fig. 8g to Fig. 8l. Displayed are only the horizontal scales below 1 km, the 2D autocorrelation coefficients for shifts below ± 1 km. With increasing wind speeds the area covered
- 575 by $P_{\tau}^2 \ge P_{\tau}^2(\xi_{\tau})$ increases. This illustrates that with increasing wind speed the size of the smallscale cloud structures increases along the predominant directions. The increased wind speed leads to stretched cloud structures along one direction. Along this predominant direction the stretching of the cloud structures smoothes smooths their variability stronger than across this direction. This leads to more homogeneous cloud structures. The turn of the orientation of the cloud structures to the East
- 580 with increasing wind speed is also represented by the fields of P_{τ}^2 .

Quantitative For the simulations with 100 m resolution, the dependency of the small–scale cloud structures on the wind speed was parameterized. Therefore, quantitative values for the size of the cloud imhomogeneity inhomogeneity structures in terms of the decorrelation length ξ_{τ} and as a function of initialization wind speed are displayed in Fig. 9a (along predominant direction) and in Fig. 9b

(across predominant structure). The threshold of P²_τ(ξ_τ) = e⁻² is marked by a grey dotted line. The derived values for ξ[‡]_τ and ξ[↔]_τ are displayed in Fig. 9c as a function of the vertical mean wind speed within the cloudy region. It shows that along the predominant structure the decorrelation length ξ[‡]_τ increases continuously (slightly quadratic increase) with increasing wind speed. The Therefore, the derived decorrelation length along (ξ[‡]_τ) the predominant structure as a function of wind speed
(vertically averaged over cloudy region) in units of m s⁻¹ can be approximated by:

$$\xi_{\tau}^{\uparrow} = 31 \cdot v^2 - 31 \cdot v + 315. \tag{5}$$

Across the predominant structure (Fig. 9c) it is different, which means that for the lower wind speeds ($\langle 2 \times v \rangle$) no influence on P_{τ}^2 and ξ_{τ} occurs, while it is comparable (slightly quadratic increase) to the values along the predominant structures for the stronger wind speeds ($\geq 2 \times v$). The derived decorrelation length across ($\xi_{\tau}^{\leftrightarrow}$) the predominant structure as a function of wind speed can be approximated by:

 $\xi_{\tau}^{\leftrightarrow} = 60 \cdot v^2 - 183 \cdot v + 365. \tag{6}$

Both, ξ_{τ}^{\uparrow} and $\xi_{\tau}^{\leftrightarrow}$ characterize the small-scale cloud inhomogeneities. Large-scale cloud structures cannot be represented due to the too small domain size. However, comparing ξ_{τ}^{\uparrow} with $\xi_{\tau}^{\leftrightarrow}$ shows that the directionality of the cloud structures first increases (0.5 to 2.0 × v) and afterwards decreases (2.0 to 3.0 × v) again. For the case investigated here, the threshold at 2.0 × v applies to a mean v (vertically averaged over cloudy region) of 3.0 m s⁻¹.

Comparing the simulations for the small domain $(1.6 \times 1.6 \text{ km}, 50 \text{ m} \text{ spatial resolution})$ with the large domain $(6.4 \times 6.4 \text{ km}, 100 \text{ m} \text{ spatial resolution})$, indicates that the small-scale structures

- 605 are most likely influenced by the large–scale structures. Only for the simulations with the large domain, the small–scale structures depend on the wind speed. This indicates that small–scale cloud inhomogeneities are not directly linked to the wind speed but rather are influenced by the large–scale cloud inhomogeneities such as cloud roles. If these large–scale structures are not covered by the simulations (too small domain), the natural behavior of the small–scale structures (e.g. their
- 610 size and orientation) might be disturbed. With respect to the comparison between observations and simulations, this may explain why only on 14 May larger differences between model and observations were found. All other three cases did not show a significant large-scale cloud structure, while on 14 May cloud roles were observed by the imaging spectrometer. Thus, the simulations of 15, 16, and 17 May are more uncritical with respect to the model domain than for 14 May, when
- 615 a large domain is required to reproduce the large–scale cloud structures and, therefore, improve the simulation of the small–scale cloud structures.

7 Summary and Conclusions

Cloud remote sensing results (of cloud optical thickness) and atmospheric profile measurements (and atmospheric dropsonde measurements (profiles of air pressure, temperature, relative humidity,

- 620 wind vector) by dropsondes from the airborne VERDI campaign conducted in April/May 2012 are exploited. The analysis focuses on In particular, a persistent cloud layer probed in close vicinity was analyzed, which was probed on four consecutive days from 14 to 17 May 2012 in almost the same area (≤ 50 km) and over constant surface conditions (open water; Polynia)on four consecutive days from 14 to 17 May 2012... The cloud top altitude of the cloud layer in the respective area shrinked
- 625 <u>shrank</u> from day to day; it decreased from about 880 m on 14 May to around 200 m on 17 May. This case was applied as a test bed for The airborne observations obtained during these days were applied to validate cloud simulations with COSMO by a new approach by comparing, which compares the observed and simulated 2D cloud fields.
- The dropsonde profile measurements from the four consecutive days were used to initialize the cloud simulations with COSMO. It is found that COSMO captures the measured cloud altitude, cloud vertical extent, and retrieved cloud optical thickness. The comparison of the horizontal cloud inhomogeneities small-scale cloud inhomogeneities identified within the observations and simulations was performed for horizontal fields of cloud optical thickness τ using . Those τ were either retrieved from airborne observations of reflected solar radiances and (τ_{meas}) or obtained from
- 635 simulated 3D fields of LWC . The (τ_{sim}) . For the reason of comparability, the observed fields of cloud optical thickness τ_{meas} were aggregated to pixel sizes of 50 m and 100 m. The, the applied spatial resolutions of the individual simulations.

The general inhomogeneity was compared using 1D inhomogeneity parameters. For 100 m spatial resolution the absolute values of cloud inhomogeneity derived from COSMO are larger by a factor

- 640 of about two, as compared to the values obtained from the observations; These differences slightly reduce, when the spatial resolution of the simulations is increased by a finer grid of 50 m. However, for both spatial resolutions the cloud inhomogeneity generated by COSMO is too low. This is mainly related to (i) the larger effective grid spacing ($\approx 3 \times 100$ m, Skamarock et al., 2004) spatial resolution ($\approx 3 \times 50$ m and $\approx 3 \times 100$ m, respectively, Skamarock et al., 2004) of COSMO com-
- 645 pared to the pixel size of the observations and (ii) a mismatch in timing/spacing, meaning that for the fields of τ_{sim} from simulations by COSMO the resulting 2D autocorrelation functions $P_{\tau,sim}^2$ 1D inhomogeneity parameters are averaged over several time steps simulated at always over the same location, while for the observed fields of τ_{meas} the resulting 2D autocorrelation functions $P_{\tau,meas}^2$ are averaged observations the 1D inhomogeneity parameters are averaged over several time steps
- along the flight track. Furthermore, in These results are in agreement with a model intercomparison Ovehinnikov et al. (2014) by Ovchinnikov et al. (2014), who revealed that COSMO underestimates the variance of the vertical wind velocity compared to other LES models and, thus , may cause an underestimation of the standard deviation of τ_{sim} is possible. However, except for the case on 16 May

the trend of the temporal evolution of the overall cloud inhomogeneity different magnitudes of cloud
inhomogeneity of the individual days is well covered by COSMO.

Especially for the cases on 14and 16 May the cloud structures structure showed a distinct directional orientation and on , while from 15 and to 17May a still slightly directional orientation May only a slight directional orientation is observed. Brümmer (1999) points out that such directed cloud structures are typical for Arctic stratus with cloud top altitudes below 1 km, which is the case here,

- 660 while. <u>Contrarily</u>, for Arctic stratus with cloud top altitudes <u>larger than above</u> 1.4 km some cell structures are common. Based on a new method, proposed by Schäfer et al. (2017a), which is applied to COSMO data for the first time, a 2D analysis using autocorrelation functions is used to examine directional features of the cloud structures. The investigations showed that, in general, COSMO captured the observed directional structures of the cloud inhomogeneitiesquite well. The wind di-
- rections of the individual cases showed a significant correlation to the direction of the predominant directional structures. During the four investigated days the orientation of the dominant directional structures within the observations turned eastwards by the same degree the wind direction changed. Similar results were found by (Houze, 1994), who stated that in case of changing wind shear cloud streets will be orientated along the mean wind direction. Here, this is reproduced for the cases with 0.5 to 2.0
 - The autocorrelation analysis was used to derive the characteristic size scale of the small-scale cloud structures by estimating the decorrelation length ξ_{τ} , which is the distance at which the squared autocorrelation coefficients P_{τ}^2 drop below e^{-2} . The decorrelation lengths ξ_{τ} were calculated along (ξ_{τ}^{\uparrow}) and across $(\xi_{\tau}^{\leftrightarrow})$ the strongest extend of the derived $P_{\tau,\text{meas}}^2$ and $P_{\tau,\text{sinn}}^2$. For the COSMO
- 675 simulations with a spatial resolution of $50 \times v$. m, the ξ_{τ}^{\uparrow} and $\xi_{r}^{\leftrightarrow}$ agree well between observations and simulations, except for the case on 14 May. In contrast, for the simulations with a spatial resolution of 100 m, COSMO produced small–scale cloud structures with characteristic sizes 20 to 30 % larger compared to the observations. However, for both spatial resolutions the best agreement was found for the case on 15 May 2012.
- 680 The good agreement between the agreement between COSMO results and the observations justified and observations for the case on 15 May 2012 is used as basis for a systematic sensitivity study regarding the main drivers of the cloud inhomogeneities. The wind speed was expected to be with respect to the wind speed as a main forcing for the degree of cloud inhomogeneity. Repeating the simulations drivers of the cloud inhomogeneities. Simulations for the case on 15 May with
- 685 differently scaled initialization profiles for the wind speed, wind profiles showed that the degree of horizontal cloud inhomogeneity was significantly changed. This not significantly changed for the simulations with a small domain $(1.6 \text{ km} \times 1.6 \text{ km})$ and 50 m spatial resolution, but for the simulations using a large domain $(6.4 \text{ km} \times 6.4 \text{ km})$ and 100 m spatial resolution. This indicates that the large-scale cloud structures such as cloud roles influence the small-scale cloud inhomogeneity.
- 690 To correctly simulate the small-scale cloud inhomogeneity, COSMO needs to be run in a large

domain, which also covers the large-scale cloud structures. This might have been the reason for the large differences between observations and simulations found for the case of 14 May, when pronounced cloud rolls were observed. All other cases did not show such large-scale cloud structures and were simulated by COSMO closer to reality despite the small domain.

- 695 However, the significant impact of the wind on the small-scale cloud structures for simulations with 100 m spatial resolution confirms the importance of the wind speed for cloud inhomogeneities. Furthermore, For this case it was found that increasing wind speeds lead to larger horizontal cloud structures (increased decorrelation lengths). It was concluded that the A directionality of the cloud structures first increases (0.5 to $2.0 \times v$) and afterwards decreases (2.0 to $3.0 \times v$) again. Using
- 700 all averaged wind speeds (in the altitudes of the cloudy region) of the six applied wind profiles parameterizations with wind speed. A parameterization of the decorrelation lengths along and across the strongest autocorrelation were with respect to the average wind speed in cloud altitude was derived, which can be used in future studies to generate cloud structures with specific sizes and shapes. Furthermore, it was is concluded that the wind direction and the atmospheric boundary layer struc-
- 705 ture are the explanation for the differences on 16 May. In contrast to the other three days a change of the wind direction of about 50° is found close to the cloud top. MoreoverAdditionally, the ABL is-was well mixed on 16 May, which increases the turbulent mixing within in the ABL and the cloud layer, and consequently influences the cloud top structure. Local differences in the wind fields at the position where the dropsonde was released and the location where the imaging spectrome-
- 710 ter measured might be the reason that this was not equally well captured by the simulations and measurements. The difficulties of COSMOto represent the small-scale cloud structures is a further reason.

All together Altogether, cloud inhomogeneities are challenging to be captured by models and with this study a new method is applied to combine observed and simulated fields of cloud optical

- 715 thicknessa challenge for cloud resolving models. Not only the spatially averaged magnitude of inhomogneity but also the directional structure and the interaction with large-scale cloud structures needs to be reproduced in the simulations. Although COSMO produces more homogeneous clouds, it performed well, because it correctly represented the directional structures and the tendency of increasing/decreasing general degree of cloud inhomogeneityof most of the cases right, if no
- 720 larger-scale cloud structures are present. However, the statistical methods applied in this study can also be applied to characterize the larger-scale dynamic patterns, if the domain is large enough to resolve them.

8 Data availability

730

The fields of cloud optical thickness retrieved from the AisaEAGLE measurements are published on
PANGAEA (Schäfer et al., 2017b). All other data used in this study are available upon request from the corresponding authors (michael.schaefer@uni-leipzig.de, katharina.loewe@kit.edu).

Acknowledgements. We thank Marco Paukert for introducing the COSMO setup. K. Loewe and C. Hoose acknowledge partial funding through the Helmholtz Programme "Atmosphere and Climate" and this project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under grant agreement No 714062 (ERC Starting Grant "C2Phase"). We

- gratefully acknowledge the support by the SFB/TR 172 "ArctiC Amplification: Climate Relevant Atmospheric and SurfaCe Processes, and Feedback Mechanisms (AC)³" in Project B03 funded by the DFG. We thank Eike Bierwirth for organizing and performing the imaging spectrometer measurements during the VERDI campaign. Furthermore, we thank Paul Vochezer, Martin Schnaiter and Emma Järvinen for providing the SID3 data. We
- 735 are grateful to the Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany for supporting the VERDI campaign with the aircraft and manpower. In addition we like to thank Kenn Borek Air Ltd., Calgary, Canada for the great pilots who made the complicated measurements possible. For excellent ground support with offices and accommodations during the campaign we are grateful to the Aurora Research Institute, Inuvik, Canada.

740 References

- Barrett, A. I., Hogan, R. J. and Forbes, R. M.: Why are mixed-phase altocumulus clouds poorly predicted by large-scale models? Part 1. Physical processes, J. Geophys. Res. Atmos., 122, doi:10.1002/2016JD026321, 2017.
- Barrett, A. I., Hogan, R. J. and Forbes, R. M.: Why are mixed-phase altocumulus clouds poorly predicted by
- 745 large-scale models? Part 2. Vertical resolution sensitivity and parameterization, J. Geophys. Res. Atmos., 122, doi:10.1002/2016JD026322, 2017.
 - Bierwirth, E., Ehrlich, A., Wendisch, M., Gayet, J.-F., Gourbeyre, C., Dupuy, R., Herber, A., Neuber, R., and Lampert, A.: Optical thickness and effective radius of Arctic boundary-layer clouds retrieved from airborne nadir and imaging spectrometry, Atmos. Meas. Tech., 6, 1189–1200,doi:10.5194/amt-6-1189-2013, 2013.
- 750 Brümmer, B.: Roll and Cell Convection in Wintertime Arctic Cold-Air Outbreaks, J. Atmos. Sci., 56, 2613– 2636, doi:10.1175/1520-0469(1999)056<2613:RACCIW>2.0.CO;2, 1999.

Cahalan, R.: Bounded cascade clouds: albedo and effective thickness, Nonlin. Proc. Geophys., 1, 1994.

Chylek, P. and Borel, C.: Mixed phase cloud water/ice structure from satellite data high spatial resolution, Geophys. Res. Lett., 31, L14104, doi:10.1029/2004GL020428, 2004.

- 755 Coleman, D. M.: Evaluation of the Performance of the Dropsonde Humidity Sensor in Clouds, SOARS, 2003. Costa, A., Meyer, J., Afchine, A., Luebke, A., Günther, G., Dorsey, J. R., Gallagher, M. W., Ehrlich, A., Wendisch, M., Baumgardner, D., Wex, H., and Krämer, M.: Classification of Arctic, midlatitude and tropical clouds in the mixed-phase temperature regime, Atmos. Chem. Phys., 17, 12219-12238, doi:10.5194/acp-17-12219-2017, 2017.
- 760 Curry, J. A., Rossow, W. B., Randall, D., and Schramm, J. L.: Overview of Arctic cloud and radiation characteristics, J. Climate, 9, 1.731–1.764, doi:10.1175/1520-0442(1996)009<1731:OOACAR>2.0.CO;2, 1996.

Davis, A., Marshak, A., Gerber, H., and Wiscombe, W.: Horizontal structure of marine boundary layer clouds from centimeter to kilometer scales, J. Geophys. Res., 104, 6123–6144, 1999b.

Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda,

- M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N. and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system ,Q.J.R. Meteorol. Soc., 137, 553-
- 597, doi:10.1002/qj.828, 2011.
 Ehrlich, A., Bierwirth, E., Istomnia, L., and Wendisch, M.: Combined retrieval of Arctic liquid water cloud and
 - surface snow properties using airborne spectral solar remote sensing, Atmos. Chem. Tech., 10, 3215–3230, doi:10.5194/amt-10-3215-2017, 2017.
 - Field, P.R., Hogan, R.J., Brown, P.R.A., Illingworth, A.J., Choularton, T.W., Kaye, P.H., Hirst, E., Greenaway,
- R.: Simultaneous radar and aircraft observations of mixed-phase cloud at the 100 m scale, Q. J. R. Meteorol.
 Soc., 130, 1877–1904, doi:10.1256/qj.03.102, 2004.
 - Herzog, , H.-J., Vogel, G., and Schubert, U.: LLM a nonhydrostatic model applied to high-resolving simulations of turbulent fluxes over heterogeneous terrain, Theor. Appl. Climatol., 73, 67–86, doi:10.1007/s00704-002-0694-4, 2002.

- 780 Hinkelmann, L. M.: Differences between along-wind and cross-wind solar irradiance variability on small spatial scales, Sol. Energy, 88, 192–203, doi:10.1016/j.solener.2012.11.011, 2013.
 - Hock, T. F. and Franklin, J. L.: The NCAR GPS dropwindsonde, Bull. Amer. Meteorol. Soc., 80, 407–420, doi:10.1175/1520-0477(1999)080<0407:TNGD>2.0.CO;2, 1999.

Robert A. Houze, Jr.: Cloud Dynamics, International Geophysics series, 53, p. 166, 1994.

- 785 Intrieri, J. M., Shupe, M. D., Uttal, T., and McCarty, B. J.: An annual cycle of Arctic cloud characteristics observed by radar and lidar at SHEBA, J. Geophys. Res., 107, (C10), doi:10.1029/2000JC000423, 2002.
 Iwabuchi, H. and Hayasaka, T.: Effects of cloud horizontal inhomogeneity on the optical thickness retrieved from moderate-resolution satellite data, J. Atmos. Sci., 59, 2227–2242, doi:10.1175/1520-0469(2002)059<2227:EOCHIO>2.0.CO;2, 2002.
- 790 Jacobson, L., Vihma, T., Jakobson, E., Palo, T., Männik, A., and Jaagus, J.: Low-level jet characteristics over the Arctic Ocean in spring and summer, Atmos. Chem. Phys., 13, 11089–11099, doi:10.5194/acp-13-11089-2013, 2013.
 - Klein, S. A., McCoy, R. B., Morrison, H., Ackerman, A. S., Avramov, A., de Boer, G., Chen, M., Cole, J. N. S., Del Genio, A. D., Falk, M., Foster, M. J., Fridlind, A., Golaz, J.-C., Hashino, T., Harrington, J. Y., Hoose,
- C., Khairoutdinov, M. F., Larson, V. E., Liu, X., Luo, Y., McFarquhar, G. M., Menon, S., Neggers, R. A. J., Park, S., Poellot, M. R., Schmidt, J. M., Sednev, I., Shipway, B. J., Shupe, M. D., Spangenberg, D. A., Sud, Y. C., Turner, D. D., Veron, D. E., Salzen, K. v., Walker, G. K., Wang, Z., Wolf, A. B., Xie, S., Xu, K.-M., Yang, F., and Zhang, G.: Intercomparison of model simulations of mixed-phase clouds observed during the ARM Mixed-Phase Arctic Cloud Experiment. I: single-layer cloud, Q. J. R. Meteorol. Soc., 135, 979–1002, doi:10.1002/qj.416, 2009.
- - Klingebiel, M.; de Lozar, A.; Molleker, S.; Weigel, R.; Roth, A.; Schmidt, L.; Meyer, J.; Ehrlich, A.; Neuber, R.; Wendisch, M. and Borrmann, S.: Arctic low-level boundary layer clouds: in situ measurements and simulations of mono- and bimodal supercooled droplet size distributions at the top layer of liquid phase clouds, Atmos. Chem. Phys., 15, 617-631, doi:10.5194/acp-15-617-2015, 2015.
- 805 Kopec, M. K., Malinowskia, S. P. and Piotrowskib, Z. P.: Effects of wind shear and radiative cooling on the stratocumulus-topped boundary layer, Q. J. R. Meteorol. Soc. 142: 3222–3233, doi:10.1002/qj.2903, 2016. Korolev, A.: Limitations of the Wegener–Bergeron–Findeisen Mechanism in the Evolution of Mixed-Phase Clouds, J. Atmos. Sci., 64, 3372–3375, doi:10.1175/JAS4035.1, 2007.
- Langhans, , W., Schmidli, J., and Szintai, B.: A Smagorinsky-Lilly turbulence closure for COSMO-LES:
 810 Implementation and comparison to ARPS, COSMO newsletter, 12, 20-31, 2012.
- Lawson, R.P., Stamnes, K., Stamnes, J., Zmarzly, P., Koskuliks, J., Roden, C., Mo, Q., Carrithers, M., and Bland, G.L.: Deployment of a Tethered-Balloon System for Microphysics and Radiative Measurements in Mixed-Phase Clouds at Ny–Ålesund and South Pole, J. Atmos. Ocean. Tech., 28, 656–670, doi:10.1175/2010JTECHA1439.1, 2010.
- 815 Lindsay, R., Wensnahan, M., Schweiger, A., and Zhang, J.: Evaluation of seven different atmospheric reanalysis products in the arctic, J. Climate, 27, 2588–2605, doi:10.1175/JCLI-D-13-00014.1, 2014.

Loewe, K.: Arctic mixed-phase clouds Macro- and micropysical insights with a numerical model, KIT Scientific publishing, doi:10.5445/KSP/1000070973, 2017.

Loewe, K., Ekman, A. M. L., Paukert, M., Sedlar, J., Tjernström, M., and Hoose, C.: Modelling micro- and
 macrophysical contributors to the dissipation of an Arctic mixed-phase cloud during the Arctic Summer
 Cloud Ocean Study (ASCOS), Atmos. Chem. Phys., 17, 6693–6704, doi:10.5194/acp-17-6693-2017, 2017.

- Marchand, R. T., Ackermann, T.P., and Moroney, C.: An assessment of Multiangle Imaging Spectroradiometer (MISR) stereo-derived cloud top heights and cloud top winds using ground-based radar, lidar, and microwave radiometers, J. Geophys. Res., 112, D06204, doi:10.1029/2006JD007091, 2007.
- 825 Marshak, A., Davis, A., Wiscombe, W., and Cahalan, R.: Radiative smoothing in fractal clouds, J. Geophys. Res., 100, 26 247–26 261, 1995.
 - McFarquhar, G.M., Zhang, G., Poellot, M.R., Kok, G.L., McCoy, R., Tooman, T., Friedlind, A., and Heymsfield, A.J.: Ice properties of single-layer stratocumulus during the Mixed-Phase Arctic Cloud Experiment: 1. Observations, J. Geophys. Res., 112, D24201, doi:10.1029/2007JD008633, 2007.
- 830 McFarquhar, G.M., S. Ghan, J. Verlinde, A. Korolev, J.W. Strapp, B. Schmid, J.M. Tomlinson, M. Wolde, S.D. Brooks, D. Cziczo, M.K. Dubey, J. Fan, C. Flynn, I. Gultepe, J. Hubbe, M.K. Gilles, A. Laskin, P. Lawson, W.R. Leaitch, P. Liu, X. Liu, D. Lubin, C. Mazzoleni, A. Macdonald, R.C. Moffet, H. Morrison, M. Ovchinnikov, M.D. Shupe, D.D. Turner, S. Xie, A. Zelenyuk, K. Bae, M. Freer, and A. Glen: Indirect and Semi-direct Aerosol Campaign., Bull. Amer. Meteor. Soc., 92, 183–201, doi:10.1175/2010BAMS2935.1,
- 835 2011.

840

855

Mellado, J. P.: Cloud-Top Entrainment in Stratocumulus Clouds, Ann. Rev. of Fluid Mech., 49, 145–69, doi:10.1146/annurev-fluid-010816-060231, 2017.

- Mioche, G., Jourdan, O., Ceccaldi, M., and Delanoë, J.: Variability of mixed-phase clouds in the Arctic with a focus on the Svalbard region: a study based on spaceborne active remote sensing, Atmos. Chem. Phys., 15, 2445—2461, doi:10.5194/acp-15-2445-2015, 2015.
- Morrison, H., de Boer, G., Feingold, G., Harrington, J., Shupe, M. D. and Sulia, K: Resilience of persistent Arctic mixed-phase clouds, Nat. Geosci., 5, 11-17, doi:10.1038/NGEO1332, 2012.

Oreopoulos, L., Cahalan, R., Marshak, A., and Wen, G.: A new normalized difference cloud retrieval technique applied to Landsat radiances over the Oklahoma ARM site, J. Appl. Meteorol., 39, 2305–2321, 2000.

- 845 Oreopoulos, L. and Cahalan, R. F.: Cloud Inhomogeneity from MODIS, J. Climate, 18, 5110–5124, 2005. Ovchinnikov, M., Ackerman, A. S., Avramov, A., Cheng, A., Fan, J., Fridlind, A. M., Ghan, S., Harrington, J., Hoose, C., Korolev, A., McFarquhar, G. M., Morrison, H., Paukert, M., Savre, J., Shipway, B. J., Shupe, M. D., Solomon, A., and Sulia, K.: Intercomparison of large-eddy simulations of Arctic mixedphase clouds: Importance of ice size distribution assumptions, J. Adv. Model. Earth Syst., 6, 223–248,
- doi:10.1002/2013MS000282, 2014.
 - Paukert, M. and Hoose, C.: Modeling immersion freezing with aerosol-dependent prognostic ice nuclei in Arctic mixed-phase clouds, J. Geophys. Res. Atmos., 14, 9073–9092, doi:10.1002/2014JD021917, 2014.

Pedersen, J. G., Malinowski, S. P. and Grabowski, W. W.: Resolution and domain-size sensitivity in implicit large-eddy simulation of the stratocumulus-topped boundary layer, J. Adv. Model. Earth Syst., 8, 885–903,doi:10.1002/2015MS000572, 2016.

Ritter, B. and Geleyn, J.-F.: A Comprehensive Radiation Scheme for Numerical Weather Prediction Models with Potential Applications in Climate Simulations, Mon. Wea. Rev., 120, 303–325, doi:10.1175/1520-0493(1992)120<0303:ACRSFN>2.0.CO;2, 1992. Roesler, E. L., D. J. Posselt, and R. B. Rood: Using large eddy simulations to reveal the size, strength, and phase

- of updraft and downdraft cores of an Arctic mixed-phase stratocumulus cloud, J. Geophys. Res. Atmos., 122, 4378–4400.doi:10.1002/2016JD026055, 2016.
 - Rossow, W. B., and Schiffer, R. A.: ISCCP cloud data products, Bull. Amer. Meteor. Soc., 72, 2–20, 1991.
 - Schäfer, M., Bierwirth, E., Ehrlich, A., Heyner, F., and Wendisch, M.: Retrieval of cirrus optical thickness and assessment of ice crystal shape from ground-based imaging spectrometry, Atmos. Meas. Tech., 6, 1855– 1868, doi:10.5194/amt-6-1855-2013, 2013.
- Schäfer, M., Bierwirth, E., Ehrlich, A., Jäkel, E., and Wendisch, M.: Airborne observations and simulations of three-dimensional radiative interactions between Arctic boundary layer clouds and ice floes, Atmos. Chem. Phys., 15, 8147–8163, doi:10.5194/acp-15-8147-2015, 2015.
 - Schäfer, M., Bierwirth, E., Ehrlich, A., Jäkel, E., Werner, F., and Wendisch, M.: Directional, horizontal inhomo-
- geneities of cloud optical thickness fields retrieved from ground-based and airborne spectral imaging, Atmos.
 Chem. Phys., 17, 2359–2372, doi:10.5194/acp-17-2359-2017, 2017.
 - Schäfer, M., Bierwirth, E., Ehrlich, A., Jäkel, E., Werner, F., and Wendisch, M.: Cloud optical thickness retrieved from horizontal fields of reflected solar spectral radiance measured with AisaEAGLE during VERDI campaign 2012, PANGAEA, doi:10.1594/PANGAEA.874798, 2017.
- 875 Schättler, U., Doms, G., and Schraff, C.: A description of the non-hydrostatic regional COSMO-model, part VII: user's guide, http://www.cosmo-model.org, 2015.
 - Schröder, M.: Multiple scattering and absorption of solar radiation in the presence of three-dimensional cloud fields, Ph.D. thesis, Fachbereich Geowissenschaften der Freien Universität Berlin, 2004.

Seifert, A., and Beheng, K. D.: A two-moment cloud microphysics parameterization for mixed-phase clouds.

880 Part 1: Model description, Meteorol. Atmos. Phys., 92, 45–66, doi:10.1007/s00703-005-0112-4, 2006.

- Serreze, M. C., and Barry, R. G.: Processes and impacts of Arctic amplification: A research synthesis, Global Planet. Change, 77, 85–96, doi:10.1016/j.gloplacha.2011.03.004, 2011.
 - Shiobara, M., Masanori, Y., and Kobayashi, H.: A polar cloud analysis based on Micro-pulse Lidar measurements at Ny-Alesund, Svalbard and Syowa, Antarctica, Phys. Chem. Earth, 28, 1205–1212,
- doi:10.1016/j.pce.2003.08.057, 2003.

865

- Shupe, M. D. and Intrieri, J. M.: Cloud radiative forcing of the Arctic surface: The influence of cloud properties, surface albedo, and solar zenith angle, J. Climate, 17, 616–628, 2004.
- Shupe, M. D., Matrosov, S. Y., and Uttal, T.: Mixed-Phase Cloud Properties Derived from Surface-Based Sensors at SHEBA, J. Atmos. Sci., 63, 697–711, doi:10.1175/JAS3659.1, 2006.
- 890 Shupe, M. D., Kollias, P., Persson, P.O.G., and McFarquhar, G.M.: Vertical Motions in Arctic Mixed-Phase Stratiform Clouds, J. Atmos. Sci., 65, 1304–1321, doi:10.1175/2007JAS2479.1, 2008.
 - Shupe, M. D., Walden, V. P., Eloranta, E., Uttal, T., Campbell, J. R., Starkweather, S. M., and Shiobara, M.: Clouds at Arctic Atmospheric Observatories. Part I: Occurrence and Macrophysical Properties, J. Appl. Meteorol. Clim., 50, 626–644, doi:10.1175/2010JAMC2467.1, 2011.
- 895 Skamarock, W. C.: Evaluating Mesoscale NWP Models Using Kinetic Energy Spectra, Mon. Wea. Rev., 132, 3019–3032, doi:10.1175/MWR2830.1, 2004.
 - Stevens, R. G., Loewe, K., Dearden, C., Dimitrelos, A., Possner, A., Eirund, G. K., Raatikainen, T., Hill, A. A., Shipway, B. J., Wilkinson, J., Romakkaniemi, S., Tonttila, J., Laaksonen, A., Korhonen, H., Connolly,

P., Lohmann, U., Hoose, C., Ekman, A. M. L., Carslaw, K. S. and Field, P. R.: A Model Intercomparison
of CCN-Limited Tenuous Clouds in the High Arctic, Atmos. Chem. Phys. Discuss., doi:10.5194/acp-2017-1128, in reviewaccepted, 2017.

- Szczap, F., Isaka, H., Saute, M., and Guillemet, B.: Effective radiative properties of bounded cascade nonabsorbing clouds: Definition of the equivalent homogeneous cloud approximation, J. Geophys. Res., 105, 20617–20633, 2000.
- 905 Varnai, T. and Marshak, A.: Observations of Three-Dimensional Radiative Effects that Influence MODIS Cloud Optical Thickness Retrievals, J. Atmos. Sci., 59, 1607–1618, 2001.
 - Vochezer, P., Järvinen, E., Wagner, R., Kupiszewski, P., Leisner, T., and Schnaiter, M.: In situ characterization of mixed phase clouds using the Small Ice Detector and the Particle Phase Discriminator, Atmos. Meas. Tech., 9, 159–177, doi:10.5194/amt-9-159-2016, 2016.
- 910 Vavrus, S.: The Impact of Cloud Feedbacks on Arctic Climate under Greenhouse Forcing, J. Climate, 17, 603–615, doi:10.1175/1520-0442(2004)017<0603:TIOCFO>2.0.CO;2, 2004.
 - Wendisch, M., Yang, P., and Ehrlich, A., (Eds.): Amplified climate changes in the Arctic: Role of clouds and atmospheric radiation, 132, 1–34, Sitzungsberichte der S\u00e4chsischen Akademie der Wissenschaften zu Leipzig, Mathematisch-Naturwissenschaftliche Klasse, S. Hirzel Verlag, Stuttgart/Leipzig, 2013.
- 915 Wendisch, M., Brückner, M., Burrows, J. P., Crewell, S., Dethloff, K., Ebell, K., Lüpkes, Ch., Macke, A., Notholt, J., Quaas, J., Rinke, A., and Tegen, I.: Understanding causes and effects of rapid warming in the Arctic, EOS, 98(8), 22–26, doi:10.1029/2017EO064803, 2017.